

WEIGHTED COMPOSITION OPERATORS ACTING FROM THE LIPSCHITZ SPACE TO THE SPACE OF BOUNDED FUNCTIONS ON A TREE

TAKUYA HOSOKAWA

ABSTRACT. We study the weighted composition operators between the Lipschitz space and the space of bounded functions on the set of vertices of an infinite tree. We characterized the boundedness, the compactness, and the boundedness from below of weighted composition operators. We also determine the isometric weighted composition operators.

1. INTRODUCTION

A graph $G = (V, E)$ is a pair of a nonempty set V , which is called the vertex set, and a subset E of $\{[v, w] \in V \times V : v \neq w\}$, which is called the edge set. Two vertices v and w are called neighbors if $[v, w] \in E$, and we denote by $v \sim w$. For any vertex $v \in V$, let $\deg v$ denote the number of neighbors of v . The vertex v is called terminal if $\deg v = 1$. A path is a finite or infinite sequence $[v_0, v_1, \dots]$ of vertices such that $v_k \sim v_{k+1}$ and $v_k \neq v_{k+2}$ for all k .

A tree is a connected, locally finite, undirected graph with no loop, no cycle. Remark that for any two vertices v, w there exists a unique finite path from v to w . For any vertex $v, w \in V$, let the distance between v and w be the numbers of edges in the finite path from v to w , and we denote by $d_T(v, w)$. In this paper, we assume that the tree T is rooted at a vertex o and has no terminal. Hence T is an infinite graph.

A vertex v is called descendant of w if w lies in the unique path from o to v . Then w is called an ancestor of v . The set S_v consisting of v and all descendants of v is called the sector determined by v . For $v \in T$, define that $|v| = d_T(o, v)$. We use the notation $T^* = T \setminus \{o\}$. The parent v^- of $v \in T^*$ is the unique vertex satisfying $v^- \sim v$ and

1991 *Mathematics Subject Classification.* 47B33, 47B38, 05C05.

Key words and phrases. tree, Lipschitz space, weighted composition operator.

The author is partially supported by Grant-in-Aid for Scientific Research, Japan Society for the Promotion of Science (No.16K05190).

$|v^-| = |v| - 1$. Remark that for any neighbors $v \sim w$, we have the alternative of $v = w^-$ or $w = v^-$.

In [3], Cohen and Colonna pointed out the geometric correspondence between the hyperbolic disk and the homogeneous trees, that is, each vertex has the same number of neighbors. The authors described the relation in terms of the Möbius transformations and the hyperbolic tilings. In this line, we can regard the complex-valued functions on the vertices of trees (possibly non-homogeneous) as a discretization of the functions on the unit disk.

The supremum norm of f on T will be denoted by

$$\|f\|_\infty = \sup_{v \in T} |f(v)|,$$

and the space of the bounded functions on T by L^∞ . The discrete derivative of f is defined by

$$Df(v) = \begin{cases} f(v) - f(v^-) & (v \neq o), \\ 0 & (v = o). \end{cases}$$

The set of all functions f on T such that $\|Df\|_\infty < \infty$ is called the Lipschitz space, denote by \mathcal{L} . Then \mathcal{L} is a Banach space under the norm

$$\|f\|_{\mathcal{L}} = |f(o)| + \|Df\|_\infty.$$

Since $\|Df\|_\infty \leq 2\|f\|_\infty$, we have that $L^\infty \subset \mathcal{L}$. It is known that the Lipschitz functions f follows the growth condition:

$$(1) \quad |f(v)| \leq |f(o)| + |v| \cdot \|Df\|_\infty$$

(see also Lemma 3.4 of [4]).

Let the little Lipschitz space \mathcal{L}_o be the subspace of \mathcal{L} consisting of all functions f with $\lim_{|v| \rightarrow \infty} Df(v) = 0$. Colonna and Easley [4] proved that \mathcal{L}_o is the closure in \mathcal{L} of the set

$$\mathcal{P} = \left\{ \sum_{k=1}^n a_k \eta_{v_k} : n \in \mathbb{N}, v_k \in T, a_k \in \mathbb{Q}[i] \right\},$$

where η_v is the characteristic function on the sector S_v determined by v . Thus \mathcal{L}_o is a closed separable subspace of \mathcal{L} .

Let ψ be a function on T . The multiplication operator M_ψ is defined by

$$M_\psi f(v) = \psi(v) f(v).$$

For a self-map φ of T , the composition operator C_φ is defined by

$$C_\varphi f(v) = f(\varphi(v)).$$

Moreover, we define the weighted composition operator ψC_φ by

$$\psi C_\varphi f(v) = M_\psi C_\varphi f(v) = \psi(v)f(\varphi(v)).$$

For the multiplication operators, the boundedness and the compactness has been studied on some function spaces. Allen and Craig [2] characterized the boundedness and the compactness of multiplication operators on the weighted Banach space L_μ^∞ on T . We here present the results for the case of L^∞ .

Theorem A ([2]). *Let ψ be a function on T .*

- (i) M_ψ is bounded on L^∞ if and only if $\psi \in L^\infty$. Moreover, if M_ψ is bounded on L^∞ , then $\|M_\psi\|_{L^\infty} = \|\psi\|_\infty$.
- (ii) M_ψ is compact on L^∞ if and only if $\psi(v) \rightarrow 0$ as $|v| \rightarrow \infty$.

Colonna and Easley [5] characterized the boundedness and the compactness of the multiplication operators acting from the Lipschitz spaces to L^∞ .

Theorem B ([5]). *Let ψ be a function on T .*

- (i) *The following conditions are equivalent.*
 - (a) $M_\psi : \mathcal{L} \rightarrow L^\infty$ is bounded.
 - (b) $M_\psi : \mathcal{L}_o \rightarrow L^\infty$ is bounded.
 - (c) $\sup_{v \in T} |v| |\psi(v)| < \infty$.
- (ii) *Suppose that $M_\psi : \mathcal{L} \rightarrow L^\infty$ is bounded. Then the following conditions are equivalent.*
 - (a) $M_\psi : \mathcal{L} \rightarrow L^\infty$ is compact.
 - (b) $M_\psi : \mathcal{L}_o \rightarrow L^\infty$ is compact.
 - (c) $\lim_{|v| \rightarrow \infty} |v| |\psi(v)| \rightarrow 0$.

In [1], Allen, Colonna, and Easley also studied the composition operators on the Lipschitz space \mathcal{L} .

The following is called the weak convergence lemma. Allen and Craig introduced similar result on more general settings (Lemma 2.5 in [2]).

Lemma 1.1. *Let X and Y be \mathcal{L} , \mathcal{L}_o , or L^∞ . Let A be a bounded operator acting from X to Y . Then A is compact if and only if $\|A f_n\|_Y \rightarrow 0$ for every bounded sequence $\{f_n\}$ in X converging to 0 pointwise.*

Recall that a bounded linear operator $T : X \rightarrow Y$, where X and Y are two Banach spaces, is bounded below if there exists a positive constant M such that

$$\|Tx\|_Y \geq M\|x\|_X$$

for any $x \in X$. In [7], Müller introduced two kinds of minimum moduli of bounded linear operators, one of which is related to the boundedness from below.

Definition 1.2. (p.86, [7]) Let X, Y be two Banach spaces and T be a bounded linear operator from X to Y .

(i) The injectivity modulus $j(T)$ of T is defined by

$$j(T) = \inf\{\|Tx\|_Y : \|x\|_X = 1\}.$$

(ii) The surjectivity modulus $k(T)$ of T is defined by

$$k(T) = \sup\{r \geq 0 : T(U_X) \supset rU_Y\}.$$

It is known that $j(T) > 0$, that is, T is bounded below, if and only if T is injective and the range of T is closed in Y . It is also known that $k(T) > 0$ if and only if T is surjective. See [7] for more properties of those minimum moduli in general setting, and [6] for the estimate on the minimum moduli of weighted composition operators on H^∞ .

Our main purpose is to study the weighted composition operators ψC_φ acting between \mathcal{L} and L^∞ . In section 2, we will study the boundedness, the compactness of ψC_φ on L^∞ . We also characterize the isometric weighted composition operators, and the boundedness below of ψC_φ on L^∞ . Moreover, we determine the operator norm, the essential norm, and the minimum moduli of ψC_φ on L^∞ . In section 3, we will study the boundedness, the compactness of ψC_φ acting from \mathcal{L} and \mathcal{L}_o to L^∞ . We also show that there is no isometric weighted composition operator from \mathcal{L} and \mathcal{L}_o to L^∞ . Those results are generalizations of the results on M_ψ given in [5]. Moreover, we will give the estimate on the minimum moduli of ψC_φ from \mathcal{L} to L^∞ .

2. RESULTS ON $\psi C_\varphi : L^\infty \rightarrow L^\infty$

2.1. Boundedness and compactness.

Theorem 2.1. *Let ψ be a function on T , and φ be a self-map of T .*

(i) *ψC_φ is bounded on L^∞ if and only if $\psi \in L^\infty$. Moreover, if ψC_φ is bounded on L^∞ , then $\|\psi C_\varphi\| = \|\psi\|_\infty$.*

(ii) *Suppose that ψC_φ is bounded on L^∞ . Then ψC_φ is compact on L^∞ if and only if one of the following conditions holds:*

(a) *$\varphi(T)$ is a finite subset of T ,*

(b) *$\psi(v) \rightarrow 0$ if $|\varphi(v)| \rightarrow \infty$.*

Moreover, if $\varphi(T)$ is an infinite subset of T , the following estimate holds:

$$(2) \quad \|\psi C_\varphi\|_e = \limsup_{|\varphi(v)| \rightarrow \infty} |\psi(v)|.$$

Proof. (i) If ψC_φ is bounded on L^∞ , then $\psi = \psi C_\varphi 1 \in L^\infty$. On the other hand, if $\psi \in L^\infty$, then we have for $f \in L^\infty$ with $\|f\|_\infty = 1$,

$$\|\psi C_\varphi f\|_\infty = \|\psi(v)f(\varphi(v))\|_\infty \leq \|\psi(v)\|_\infty.$$

Hence we have ψC_φ is bounded on L^∞ if and only if $\psi \in L^\infty$.

Next we let $\chi_{\varphi(w)}$ be the characteristic function at $\varphi(w) \in T$. Then we get

$$\|\psi C_\varphi\| \geq \|\psi(v)\chi_{\varphi(w)}(\varphi(v))\|_\infty \geq |\psi(w)\chi_{\varphi(w)}(\varphi(w))| = |\psi(w)|.$$

Taking the supremum over all $w \in T$, we obtain

$$\|\psi C_\varphi\| \geq \|\psi\|_\infty.$$

We conclude that $\|\psi C_\varphi\| = \|\psi\|_\infty$.

(ii) We remark that $\|\psi\|_\infty < \infty$ since ψC_φ is bounded on L^∞ . Let $\{f_n\}$ be an arbitrary bounded sequence in L^∞ converging to 0 point-wise. Write $\|f_n\|_\infty < M$ for any n .

Suppose that $\varphi(T)$ is a finite set. We have

$$\|\psi C_\varphi f_n\|_\infty = \sup_{v \in T} |\psi(v)f_n(\varphi(v))| \leq \|\psi\|_\infty \cdot \sup_{v \in \varphi(T)} |f_n(v)| \rightarrow 0.$$

By Lemma 1.1, we have ψC_φ is compact on L^∞ .

We here suppose that $\varphi(T)$ is an infinite subset of T . Then it is enough to prove (2). Let K be a compact operator on L^∞ . For any sequence $\{v_n\}$ in T such that $|\varphi(v_n)| \rightarrow \infty$, we have

$$\|(\psi C_\varphi + K)\| \geq \|(\psi C_\varphi + K)\chi_{\varphi(v_n)}\|_\infty \geq |\psi(v_n)| - \|K\chi_{\varphi(v_n)}\|_\infty.$$

Taking the limit superior over n and the infimum over all compact operators K , we obtain

$$\|\psi C_\varphi\|_e \geq \limsup_{n \rightarrow \infty} |\psi(v_n)|.$$

Next we prove the converse. Let N be a positive integer and define the operator K_N by

$$K_N f(v) = \begin{cases} f(v) & (|v| \leq N) \\ 0 & (|v| > N) \end{cases}.$$

It is easy to see that K_N is compact on L^∞ (see the proof of Theorem 3.7 in [5]). Since $\psi C_\varphi K_N$ is also compact on L^∞ , we have

$$\begin{aligned} \|\psi C_\varphi\|_e &\leq \|(\psi C_\varphi - \psi C_\varphi K_N)\| \leq \sup_{\|f\|_\infty=1} \|(\psi C_\varphi - \psi C_\varphi K_N)f\|_\infty \\ &= \sup_{\|f\|_\infty=1} \sup_{|\varphi(v)| > N} |\psi(v)f(\varphi(v))| \leq \sup_{|\varphi(v)| > N} |\psi(v)|. \end{aligned}$$

Letting $N \rightarrow \infty$, we get

$$\|\psi C_\varphi\|_e \leq \limsup_{n \rightarrow \infty} |\psi(v_n)|.$$

This completes the proof. \square

Corollary 2.2. *Let ψ be a function on T and φ be a self-map of T .*

- (i) $\|M_\psi\| = \|\psi\|_\infty$.
- (ii) $\|C_\varphi\| = 1$.

Recall that a tree T is locally finite, and hence the self-map φ of T has finite range if and only if

$$\sup_{v \in T} |\varphi(v)| < \infty.$$

Corollary 2.3. *Let ψ be a bounded function on T and φ be a self-map of T .*

- (i) M_ψ is compact on L^∞ if and only if

$$\limsup_{|v| \rightarrow \infty} |\psi(v)| = 0.$$

- (ii) C_φ is compact on L^∞ if and only if φ has finite range in T .

Moreover, we have the zero-one law on the essential norm of ψC_φ .

Corollary 2.4. *Let φ be a self-map of T . Then*

$$\|C_\varphi\|_e = \begin{cases} 0 & (\text{if } \varphi \text{ has finite range}) \\ 1 & (\text{if } \varphi \text{ does not have finite range}) \end{cases}.$$

2.2. Isometries.

Theorem 2.5. *Let ψ be a function on T and φ be a self-map of T . Then ψC_φ is an isometry on L^∞ if and only if φ is surjective and*

$$(3) \quad \sup_{v \in \varphi^{-1}(w)} |\psi(v)| = 1$$

for any $w \in T$.

Proof. Suppose ψC_φ is an isometry on L^∞ . Since $\|\chi_w\|_\infty = 1$ for $w \in T$, we have

$$\begin{aligned} 1 &= \|\psi C_\varphi \chi_w\|_\infty = \sup_{v \in T} |\psi(v) \cdot \chi_w(\varphi(v))| \\ &= \begin{cases} 0 & (\text{if } w \notin \varphi(T)) \\ \sup_{v \in \varphi^{-1}(w)} |\psi(v)| & (\text{if } w \in \varphi(T)) \end{cases}. \end{aligned}$$

Thus we get φ is surjective and (3) for any $w \in T$.

To prove the converse, we assume φ is surjective and (3) for any $w \in T$. It follows that $\|\psi\|_\infty = 1$, and hence we have $\|\psi C_\varphi f\|_\infty \leq \|f\|_\infty$ for any $f \in L^\infty$. Let $\{w_n\}$ be a sequence in T such that $|f(w_n)| \rightarrow \|f\|_\infty$. Then we have

$$\begin{aligned} \|\psi C_\varphi f\|_\infty &\geq \lim_{n \rightarrow \infty} \sup_{v \in \varphi^{-1}(w_n)} |\psi(v) \cdot f(\varphi(v))| \\ &= \lim_{n \rightarrow \infty} \left(\sup_{v \in \varphi^{-1}(w_n)} |\psi(v)| \right) \cdot |f(w_n)| \\ &= \lim_{n \rightarrow \infty} |f(w_n)| = \|f\|_\infty. \end{aligned}$$

We get ψC_φ is an isometry on L^∞ . □

Example 2.6. Let \mathbb{Z} be the set of all integers. Define φ be a self-map of \mathbb{Z} by

$$\varphi(n) = \begin{cases} n & (n \geq 0) \\ -n & (n = -1, -3, \dots) \\ \frac{n}{2} & (n = -2, -4, \dots) \end{cases}.$$

Clearly, φ maps \mathbb{Z} onto \mathbb{Z} . Moreover, we define a function ψ on \mathbb{Z} by

$$\psi(n) = \begin{cases} 0 & (n = -1, -3, \dots) \\ 1 & (\text{otherwise}) \end{cases}.$$

Since ψ holds (3) for any $w \in \mathbb{Z}$, we have ψC_φ is an isometry on $L^\infty(\mathbb{Z})$.

Corollary 2.7. *Let ψ be a function on T and φ be a self-map of T .*

- (i) M_ψ is an isometry on L^∞ if and only if $|\psi| \equiv 1$ on T .
- (ii) C_φ is an isometry on L^∞ if and only if φ is surjective.

2.3. Boundedness from below and minimum moduli.

We determine the minimum moduli $j(\psi C_\varphi)$ and $k(\psi C_\varphi)$ for bounded weighted composition operators ψC_φ on L^∞ .

Theorem 2.8. *Let ψ be a bounded function on T and φ be a self-map of T .*

- (i) If φ is not surjective, then $j(\psi C_\varphi) = 0$.
- (ii) If φ is surjective, then

$$(4) \quad j(\psi C_\varphi) = \inf_{w \in T} \left(\sup_{v \in \varphi^{-1}(w)} |\psi(v)| \right).$$

Proof. (i) Suppose φ is not surjective. For $w \notin \varphi(T)$, we have $\psi C_\varphi \chi_w \equiv 0$ on T . This implies that $j(\psi C_\varphi) = 0$.

(ii) Suppose φ is surjective. Let M be the infimum of right side of (4). For any $w \in T$, the preimage $\varphi^{-1}(w)$ is not the empty set and

$$\|\psi C_\varphi \chi_w\|_\infty = \sup_{v \in \varphi^{-1}(w)} |\psi(v)|.$$

Thus we get

$$j(\psi C_\varphi) \leq \inf_{w \in T} \|\psi C_\varphi \chi_w\|_\infty = M.$$

Next we will show $j(\psi C_\varphi) \geq M$. For any $w \in T$ and any $f \in L^\infty$ with $\|f\|_\infty = 1$, we have

$$\begin{aligned} \|\psi C_\varphi f\|_\infty &= \sup_{v \in T} |\psi(v) \cdot f(\varphi(v))| \\ &\geq \sup_{v \in \varphi^{-1}(w)} |\psi(v) \cdot f(\varphi(v))| \\ &= \left(\sup_{v \in \varphi^{-1}(w)} |\psi(v)| \right) \cdot |f(w)| \geq M \cdot |f(w)|. \end{aligned}$$

Since w is arbitrary, we have $\|\psi C_\varphi f\|_\infty \geq M \|f\|_\infty$. Then we get $j(\psi C_\varphi) \geq M$. \square

Corollary 2.9. *Let ψ be a bounded function on T and φ be a self-map of T . Then ψC_φ is bounded below on L^∞ if and only if φ is surjective on T and*

$$\inf_{w \in T} \left(\sup_{v \in \varphi^{-1}(w)} |\psi(v)| \right) > 0.$$

We here consider the surjectivity modulus of ψC_φ on L^∞ . First, we will show that if ψ has zeros on T , then ψC_φ is not surjective on L^∞ .

Proposition 2.10. *Let ψ be a bounded function on T and φ be a self-map of T . If there exists $v_0 \in T$ such that $\psi(v_0) = 0$, then $k(\psi C_\varphi) = 0$.*

Proof. Suppose that $k(\psi C_\varphi) > 0$. For $0 < r < k(\psi C_\varphi)$, there exists $f \in L^\infty$ such that $\|f\|_\infty \leq 1$ and $\psi C_\varphi f = r \cdot \chi_{v_0}$. Then we have

$$r = r \cdot \chi_{v_0}(v_0) = \psi(v_0) \cdot f(\varphi(v_0)) = 0,$$

which is a contradiction. \square

We determine the surjectivity modulus of ψC_φ .

Theorem 2.11. *Let ψ be a bounded function on T and φ be a self-map of T .*

- (i) *If φ is not injective, then $k(\psi C_\varphi) = 0$.*
- (ii) *If φ is injective, then*

$$k(\psi C_\varphi) = \inf_{v \in T} |\psi(v)|.$$

Proof. (i) Let φ be a self-map of T , which is not injective. There exist two distinct vertices $v_1, v_2 \in T$ such that $\varphi(v_1) = \varphi(v_2)$. Suppose that $k(\psi C_\varphi) > 0$ and let $0 < r < k(\psi C_\varphi)$. Then there exists $f \in L^\infty$ such that $\|f\|_\infty \leq 1$ and $\psi C_\varphi f = r \cdot \chi_{v_1}$. Since

$$r = r \cdot \chi_{v_1}(v_1) = \psi(v_1) \cdot f_1(\varphi(v_1)),$$

we have $\psi(v_1) \neq 0$ and $f_1(\varphi(v_1)) \neq 0$. On the other hand, we have

$$0 = r \cdot \chi_{v_1}(v_2) = p(v_2) \cdot f_1(\varphi(v_2)) = p(v_2) \cdot f_1(\varphi(v_1)).$$

Thus we get $p(v_2) = 0$. By Proposition 2.10, we get $k(\psi C_\varphi) = 0$. This is a contradiction.

(ii) Suppose φ is injective. To prove $k(\psi C_\varphi) \leq \inf_{v \in T} |\psi(v)|$, we may assume $k(\psi C_\varphi) > 0$. Let ε be a positive number less than $k(\psi C_\varphi)$. For any $w \in T$, there exists $f \in L^\infty$ such that $\|f\|_\infty \leq 1$ and $\psi C_\varphi f = (k(\psi C_\varphi) - \varepsilon) \cdot \chi_w$. It follows that

$$k(\psi C_\varphi) - \varepsilon = (k(\psi C_\varphi) - \varepsilon) \cdot \chi_w(w) = \psi(w) \cdot f(\varphi(w)) \leq |\psi(w)|.$$

Since w is arbitrary, we get $k(\psi C_\varphi) - \varepsilon \leq \inf_{v \in T} |\psi(v)|$. Letting $\varepsilon \rightarrow 0$, we get $k(\psi C_\varphi) \leq \inf_{v \in T} |\psi(v)|$.

Next we will prove the converse. We may assume $M = \inf_{v \in T} |\psi(v)| > 0$. It is enough to prove that, for any $g \in L^\infty$ with $\|g\|_\infty \leq M$, there exists a function $f \in L^\infty$ such that $\|f\|_\infty \leq 1$ and $\psi C_\varphi f = g$. Remark that φ is injective and $\psi(v) \neq 0$ for any $v \in T$. We here define that

$$f(w) = \begin{cases} \frac{g(\varphi^{-1}(w))}{\psi(\varphi^{-1}(w))} & (w \in \varphi(T)) \\ 0 & (w \notin \varphi(T)) \end{cases}$$

Then we can see $\psi C_\varphi f(v) = g(v)$ and

$$\begin{aligned} \|f\|_\infty &= \sup_{w \in \varphi(T)} \frac{|g(\varphi^{-1}(w))|}{|\psi(\varphi^{-1}(w))|} \leq \frac{\|g\|_\infty}{\inf_{v \in T} |\psi(v)|} \\ &\leq \frac{M}{M} = 1 \end{aligned}$$

Thus we conclude $k(\psi C_\varphi) = \inf_{v \in T} |\psi(v)|$. \square

Corollary 2.12. *Let ψ be a bounded function on T and φ be a self-map of T .*

$$(i) \quad j(M_\psi) = k(M_\psi) = \inf_{v \in T} |\psi(v)|.$$

$$(ii) \quad j(C_\varphi) = \begin{cases} 0 & (\text{if } \varphi \text{ is not surjective}) \\ 1 & (\text{if } \varphi \text{ is surjective}) \end{cases}.$$

$$(iii) \quad k(C_\varphi) = \begin{cases} 0 & (\text{if } \varphi \text{ is not injective}) \\ 1 & (\text{if } \varphi \text{ is injective}) \end{cases}.$$

3. RESULTS ON $\psi C_\varphi : \mathcal{L} \rightarrow L^\infty$

In this section we study the weighted composition operator ψC_φ acting from \mathcal{L} to L^∞ .

3.1. Boundedness and compactness.

Theorem 3.1. *Let ψ be a function on T and φ be a self-map of T . Then the following are equivalent:*

- (i) $\psi C_\varphi : \mathcal{L} \rightarrow L^\infty$ is bounded.
- (ii) $\psi C_\varphi : \mathcal{L}_o \rightarrow L^\infty$ is bounded.
- (iii) $\psi \in L^\infty$ and $\sup_{v \in T} |\psi(v)| |\varphi(v)| < \infty$.

Furthermore, under the above conditions, the following estimate holds:

$$\max\{\|\psi\|_\infty, \|\psi(v) \cdot |\varphi(v)|\|_\infty\} \leq \|\psi C_\varphi\|_{\mathcal{L} \rightarrow L^\infty} \leq \|\psi(v)(1 + |\varphi(v)|)\|_\infty.$$

Proof. The implication (i) \Rightarrow (ii) is trivial. Suppose the condition (iii) holds. Let f be a function in \mathcal{L} with $\|f\|_{\mathcal{L}} \leq 1$. Then the growth condition (1) implies that

$$\begin{aligned} |\psi C_\varphi f(v)| &= |\psi(v) f(\varphi(v))| \leq |\psi(v)| (|f(o)| + |\varphi(v)| \cdot \|Df\|_\infty) \\ &\leq |\psi(v)| (1 + |\varphi(v)|) \|f\|_{\mathcal{L}} \\ &\leq \left\{ \|\psi\|_\infty + \sup_{v \in T} |\psi(v) \cdot \varphi(v)| \right\} \|f\|_{\mathcal{L}} < \infty. \end{aligned}$$

Hence we get (i) and the upper bound of $\|\psi C_\varphi\|_{\mathcal{L} \rightarrow L^\infty}$.

Next, suppose the condition (ii) holds. Then we get $\psi C_\varphi 1 = \psi \in L^\infty$. For any positive integer N , we let

$$(5) \quad F_N(v) = \begin{cases} |v| & (|v| \leq N) \\ N & (|v| > N) \end{cases}.$$

Clearly $F_N \in \mathcal{L}_o$ and $\|F_N\|_{\mathcal{L}} = 1$. For any vertex $w \in T$, take enough large N so that $N > |\varphi(w)|$. Then we have that

$$\|\psi C_\varphi\|_{\mathcal{L}_o \rightarrow L^\infty} \geq \|\psi C_\varphi F_N\|_\infty \geq |\psi(w) F_N(\varphi(w))| = |\psi(w)| |\varphi(w)|.$$

Taking the supremum on w over T , we conclude the condition (iii) and the lower bound of $\|\psi C_\varphi\|_{\mathcal{L} \rightarrow L^\infty}$. \square

Theorem 3.2. *Let ψ be a function on T and φ be a self-map of T . Suppose that $\psi C_\varphi : \mathcal{L} \rightarrow L^\infty$ is bounded. Then the following are equivalent:*

- (i) $\psi C_\varphi : \mathcal{L} \rightarrow L^\infty$ is compact.
- (ii) $\psi C_\varphi : \mathcal{L}_o \rightarrow L^\infty$ is compact.

(iii) $|\psi(v)| \cdot |\varphi(v)| \rightarrow 0$ if $|\varphi(v)| \rightarrow \infty$.

Since the above theorem follows from Theorem 3.4, here we do not give its proof. We here present the example satisfying the conditions of Theorem 3.1 and Theorem 3.2.

Example 3.3. Let φ be a self-map of T with infinite range. Then C_φ is not bounded from \mathcal{L} to L^∞ . Put

$$\psi(v) = \frac{1}{|\varphi(v)| + 1}$$

Then, by Theorem 3.1 and Theorem 3.2, we have that ψC_φ is bounded but is not compact. On the other hand, then $\psi^2 C_\varphi$ is compact.

Theorem 3.4. *Let ψ be a function on T and φ be a self-map of T . Suppose that ψC_φ is a bounded operator from \mathcal{L} (respectively, \mathcal{L}_o) to L^∞ . Then the following holds:*

$$\|\psi C_\varphi\|_{e, \mathcal{L} \rightarrow L^\infty} = \|\psi C_\varphi\|_{e, \mathcal{L}_o \rightarrow L^\infty} = \limsup_{|\varphi(v)| \rightarrow \infty} |\psi(v)| \cdot |\varphi(v)|.$$

Proof. It is trivial that $\|\psi C_\varphi\|_{e, \mathcal{L} \rightarrow L^\infty} \geq \|\psi C_\varphi\|_{e, \mathcal{L}_o \rightarrow L^\infty}$.

For each positive integer n , we define the operator $K_n : \mathcal{L} \rightarrow \mathcal{L}$ by

$$K_n f(v) = \begin{cases} f(v) & (|v| \leq n) \\ f(v_n) & (|v| > n) \end{cases}.$$

where v_n is the unique vertex lying in the path between o and v such that $|v_n| = n$. By Lemma 1.1, we have that K_n is compact. Therefore, we get

$$\begin{aligned} \|\psi C_\varphi\|_{e, \mathcal{L} \rightarrow L^\infty} &\leq \|\psi C_\varphi - \psi C_\varphi K_n\|_{\mathcal{L} \rightarrow L^\infty} \\ &= \sup_{\|f\|_{\mathcal{L}}=1} \|(\psi C_\varphi - \psi C_\varphi K_n)f\|_\infty \end{aligned}$$

For $|\varphi(v)| > n$, put w_n the unique vertex lying in the path between o and $\varphi(v)$ such that $|w_n| = n$. Since $d_T(\varphi(v), w_n) = |\varphi(v)| - n$, we have that

$$\begin{aligned} \|(\psi C_\varphi - \psi C_\varphi K_n)f\|_\infty &= \sup_{v \in T} |\psi C_\varphi f(v) - \psi C_\varphi K_n f(v)| \\ &= \sup_{|\varphi(v)| > n} |\psi(v)| \cdot |f(\varphi(v)) - f(w_n)| \\ &\leq \sup_{|\varphi(v)| > n} |\psi(v)| \cdot (|\varphi(v)| - n) \|f\|_{\mathcal{L}} \\ &\leq \sup_{|\varphi(v)| > n} |\psi(v)| \cdot |\varphi(v)| \cdot \|f\|_{\mathcal{L}}. \end{aligned}$$

Thus we have that

$$\|\psi C_\varphi\|_{e, \mathcal{L} \rightarrow L^\infty} \leq \sup_{|\varphi(v)| > n} |\psi(v)| \cdot |\varphi(v)|.$$

Letting $n \rightarrow \infty$, we get

$$\|\psi C_\varphi\|_{e, \mathcal{L} \rightarrow L^\infty} \leq \limsup_{|\varphi(v)| \rightarrow \infty} |\psi(v)| \cdot |\varphi(v)|.$$

Next we will prove the converse. To do this, let n be a positive integer and $r \in (0, 1)$. We define that

$$g_{n,r}(v) = \begin{cases} 0 & (|v| < \sqrt{n}) \\ \frac{n}{n-\sqrt{n}} \cdot \frac{(|v|-\sqrt{n})^{r+1}}{(n-\sqrt{n})^r} & (\sqrt{n} \leq |v| < n) \\ n & (|v| \geq n) \end{cases}.$$

Then $g_{n,r}$ is in \mathcal{L}_o and $g_{n,r}(v) \rightarrow 0$ pointwise as $n \rightarrow \infty$. By short calculation, we have

$$\|g_{n,r}\|_{\mathcal{L}} = \sup_{v \in T} |Dg_{n,r}(v)| = \frac{n}{n-\sqrt{n}} \cdot \frac{(n-\sqrt{n})^{r+1} - (n-\sqrt{n}-1)^{r+1}}{(n-\sqrt{n})^r}.$$

Since $\{x^{r+1} - (x-1)^{r+1}\}/x^r \rightarrow r+1$ as $x \rightarrow \infty$, we have that $\|g_{n,r}\|_{\mathcal{L}} \rightarrow r+1$ as $n \rightarrow \infty$. By Lemma 1.1, we have $\|Kg_{n,r}\|_{\infty} \rightarrow 0$ as $n \rightarrow \infty$ for any compact operator K . Fix a vertex $w \in T$ and put $n = |\varphi(w)|$. Since $\|\psi C_\varphi g_{n,r}\|_{\infty} \geq |\psi(w) \cdot g_{n,r}(\varphi(w))| = |\psi(w)| \cdot |\varphi(w)|$, we obtain that

$$\begin{aligned} \|\psi C_\varphi - K\|_{\mathcal{L}_o \rightarrow L^\infty} &\geq \limsup_{n \rightarrow \infty} \frac{\|(\psi C_\varphi - K)g_{n,r}\|_{\infty}}{\|g_{n,r}\|_{\mathcal{L}}} \\ &\geq \limsup_{n \rightarrow \infty} \frac{\|\psi C_\varphi g_{n,r}\|_{\infty} - \|Kg_{n,r}\|_{\infty}}{\|g_{n,r}\|_{\mathcal{L}}} \\ &\geq \frac{1}{r+1} \limsup_{|\varphi(w)| \rightarrow \infty} |\psi(w)| \cdot |\varphi(w)|. \end{aligned}$$

Now, letting $r \rightarrow 0$, we get

$$\|\psi C_\varphi\|_{e, \mathcal{L}_o \rightarrow L^\infty} \geq \limsup_{|\varphi(v)| \rightarrow \infty} |\psi(v)| \cdot |\varphi(v)|.$$

□

If we take $\varphi(v) = v$ in Theorem 3.1 and Theorem 3.2, then we get the results on the multiplication operators M_ψ , which are same as Theorem B. For the composition operators, by Theorem 3.1, Theorem 3.2 for $\psi(v) \equiv 1$, and (ii) of Corollary 2.3, and Theorem 4.2 of [1], we get the following result on C_φ .

Corollary 3.5. *Let φ be a self-map of T . Then the following are equivalent:*

- (i) $C_\varphi : \mathcal{L} \rightarrow L^\infty$ is bounded.
- (ii) $C_\varphi : \mathcal{L}_o \rightarrow L^\infty$ is bounded.
- (iii) $C_\varphi : L^\infty \rightarrow L^\infty$ is compact.
- (iv) $C_\varphi : \mathcal{L} \rightarrow L^\infty$ is compact.
- (v) $C_\varphi : \mathcal{L}_o \rightarrow L^\infty$ is compact.
- (vi) $C_\varphi : \mathcal{L} \rightarrow \mathcal{L}$ is compact.
- (vii) φ has finite range.

3.2. Isometries.

Theorem 3.6. *For any function ψ on T and any self-map φ of T , ψC_φ acting from \mathcal{L} (or \mathcal{L}_o) to L^∞ is not an isometry.*

Proof. It is enough to prove the statement for the case $\psi C_\varphi : \mathcal{L}_o \rightarrow L^\infty$. We assume that $\psi C_\varphi : \mathcal{L}_o \rightarrow L^\infty$ is an isometry. If there exists a vertex $w \notin \varphi(T)$, then we have that

$$1 = \|\psi C_\varphi \chi_w\|_\infty = \sup_{v \in T} |\psi(v) \chi_w(\varphi(v))| = 0.$$

This is a contradiction. Hence φ must be surjective and

$$\sup\{|\psi(v)| : v \in \varphi^{-1}(w)\} = 1$$

for any $w \in T$. Fix a vertex $w \in T$ with $|w| > 1$. By Theorem 3.1, we get

$$\begin{aligned} 1 &= \|\psi C_\varphi\| \geq \|\psi(v) |\varphi(v)|\|_\infty \\ &\geq \sup_{\varphi(v)=w} |\psi(v)| |\varphi(v)| \\ &= |w| \cdot \sup_{\varphi(v)=w} |\psi(v)| = |w| > 1. \end{aligned}$$

This is a contradiction. We conclude that ψC_φ is not an isometry. \square

3.3. Boundedness from below and minimum moduli.

We characterize the boundedness from below for the weighted composition operators ψC_φ acting from \mathcal{L} to L^∞ . We denote the injectivity modulus $j(\psi C_\varphi)_{\mathcal{L} \rightarrow L^\infty}$ (the surjectivity modulus $k(\psi C_\varphi)_{\mathcal{L} \rightarrow L^\infty}$, resp.) by $j(\psi C_\varphi)$ ($k(\psi C_\varphi)$, resp.) in short.

Theorem 3.7. *Let ψ be a bounded function on T and φ be a self-map of T . Suppose that ψC_φ is bounded from \mathcal{L} to L^∞ .*

- (i) *If φ is not surjective, then $j(\psi C_\varphi) = 0$.*

(ii) If φ is surjective, then

$$(6) \quad \frac{1}{3} \inf_{w \in T} \left(\sup_{v \in \varphi^{-1}(w)} |\psi(v)| \right) \leq j(\psi C_\varphi) \leq \inf_{w \in T} \left(\sup_{v \in \varphi^{-1}(w)} |\psi(v)| \right).$$

Proof. (i) can be proved exactly in the same way as in the proof of (i) of Theorem 2.8.

(ii) Suppose φ is surjective. Let

$$M = \inf_{w \in T} \left(\sup_{v \in \varphi^{-1}(w)} |\psi(v)| \right).$$

Since $\|\chi_w\|_{\mathcal{L}} = 1$ for any w , we get

$$j(\psi C_\varphi) \leq \inf_{w \in T} \|\psi C_\varphi \chi_w\|_\infty = M.$$

Next we will show $j(\psi C_\varphi) \geq M/3$. Let f be in L^∞ with $\|f\|_{\mathcal{L}} = 1$. We put $r = |f(o)|$, then we have that

$$(7) \quad \begin{aligned} \|\psi C_\varphi f\|_\infty &= \sup_{v \in T} |\psi(v) \cdot f(\varphi(v))| \\ &\geq |f(o)| \sup_{v \in \varphi^{-1}(o)} |\psi(v)| \geq rM. \end{aligned}$$

Since $\sup_{v \in T} |Df(v)| = 1 - r$, there exists a sequence $\{w_n\} \subset T^*$ such that

$$|Df(w_n)| = |f(w_n) - f(w_n^-)| > (1 - r) \left(1 - \frac{1}{n}\right).$$

Then

$$\max\{|f(w_n)|, |f(w_n^-)|\} > \frac{1}{2}(1 - r) \left(1 - \frac{1}{n}\right).$$

We choose one of the vertex from $\{w_n, w_n^-\}$ which attains the maximum above, and put u_n . Since the sequence $\{u_n\}$ in T satisfies that for any n ,

$$|f(u_n)| > \frac{1}{2}(1 - r) \left(1 - \frac{1}{n}\right),$$

we have that

$$\begin{aligned} \|\psi C_\varphi f\|_\infty &\geq |f(u_n)| \sup_{v \in \varphi^{-1}(u_n)} |\psi(v)| \\ &\geq \frac{1}{2}(1 - r) \left(1 - \frac{1}{n}\right) M. \end{aligned}$$

Letting $n \rightarrow \infty$, we get

$$(8) \quad \|\psi C_\varphi f\|_\infty \geq \frac{1}{2}(1 - r)M.$$

By (7) and (8), we have that

$$\|\psi C_\varphi f\|_\infty \geq \inf_{0 \leq r \leq 1} \left(\max \left\{ r, \frac{1}{2}(1-r) \right\} \right) M = \frac{1}{3}M.$$

Therefore, we obtain $j(\psi C_\varphi) \geq M/3$. \square

Corollary 3.8. *Let ψ be a bounded function on T and φ be a self-map of T . Suppose that ψC_φ is bounded from \mathcal{L} to L^∞ . Then ψC_φ is bounded below if and only if φ is surjective on T and*

$$\inf_{w \in T} \left(\sup_{v \in \varphi^{-1}(w)} |\psi(v)| \right) > 0.$$

We give some necessary conditions and a sufficient condition for ψC_φ to be surjective.

Proposition 3.9. *Let ψ be a bounded function on T and φ be a self-map of T . Suppose that ψC_φ is bounded from \mathcal{L} to L^∞ .*

- (i) *If there exists $v_0 \in T$ such that $\psi(v_0) = 0$, then $k(\psi C_\varphi) = 0$.*
- (ii) *If φ is not injective, then $k(\psi C_\varphi) = 0$.*
- (iii) *If φ is injective, then*

$$\frac{1}{3} \inf_{v \in T} |\psi(v)| \leq k(\psi C_\varphi) \leq \inf_{v \in T} |\psi(v)|(1 + |\varphi(v)|).$$

Proof. (i) and (ii) can be proved exactly in the same way as in the proof of Proposition 2.10 and (i) of Theorem 2.11.

(iii) Assume that φ is injective. To prove that $k(\psi C_\varphi) \leq \inf_{v \in T} |\psi(v)|(1 + |\varphi(v)|)$, we may assume $k(\psi C_\varphi) > 0$. Let ε be a positive number less than $k(\psi C_\varphi)$. For any $w \in T$, there exists $f \in \mathcal{L}$ such that $\|f\|_{\mathcal{L}} \leq 1$ and $\psi C_\varphi f = (k(\psi C_\varphi) - \varepsilon) \cdot \chi_w$. By the growth condition, we have that

$$\begin{aligned} k(\psi C_\varphi) - \varepsilon &= (k(\psi C_\varphi) - \varepsilon) \cdot \chi_w(w) \\ &= \psi(w) \cdot f(\varphi(w)) \\ &\leq |\psi(w)|(1 + |\varphi(w)|). \end{aligned}$$

Since w is arbitrary, we get $k(\psi C_\varphi) - \varepsilon \leq \inf_{v \in T} |\psi(v)|(1 + |\varphi(v)|)$. Letting $\varepsilon \rightarrow 0$, we get $k(\psi C_\varphi) \leq \inf_{v \in T} |\psi(v)|(1 + |\varphi(v)|)$.

Next we assume $\inf_{v \in T} |\psi(v)| > 0$. If f is in L^∞ with $\|f\|_\infty$, then $\|f\|_{\mathcal{L}} \leq 3$. Thus, by Theorem 2.11, we have that

$$\psi C_\varphi(U_{\mathcal{L}}) \supset \psi C_\varphi\left(\frac{1}{3} U_{L^\infty}\right) \supset \left(\frac{1}{3} \inf_{v \in T} |\psi(v)|\right) U_{L^\infty}.$$

Thus we conclude $k(\psi C_\varphi) \geq \frac{1}{3} \inf_{v \in T} |\psi(v)|$. \square

The upper estimate of $k(\psi C_\varphi)$ in (iii) of Proposition 3.9 is not best possible. We present the example satisfying $\inf_{v \in T} |\psi(v)|(1+|\varphi(v)|) > 0$ but $k(\psi C_\varphi) = 0$.

Example 3.10. Let $\varphi(n) = 2n$, and define a function ψ on \mathbb{Z} by

$$\psi(n) = \begin{cases} 1 & (n = 0) \\ \frac{1}{n} & (n \neq 0) \end{cases}.$$

Clearly, φ is injective on \mathbb{Z} and

$$\inf_{n \in \mathbb{Z}} |\psi(n)| = 0, \quad \text{and} \quad \inf_{n \in \mathbb{Z}} |\psi(n)|(1 + |\varphi(n)|) = 2 > 0.$$

We will show that ψC_φ is not surjective from \mathcal{L} to L^∞ .

To do this, assume $k(\psi C_\varphi) > M > 0$. Let $g(n) = M(-1)^n$. Since $g \in M \cdot U_{L^\infty}$, there exists a function $f \in \mathcal{L}$ such that $\|f\|_{\mathcal{L}} \leq 1$ and $\psi C_\varphi f = g$. For $n \neq 0$, we have that $f(2n) = ng(n) = Mn(-1)^n$. Choose a positive integer n satisfying $M(2n+1) > 2$. Then we have that

$$\begin{aligned} 2 &\geq |Df(2n+2)| + |Df(2n+1)| \\ &\geq |f(2n+2) - f(2n)| \\ &= M \cdot |(n+1)(-1)^{n+1} - n(-1)^n| \\ &= M(2n+1) > 2. \end{aligned}$$

This is a contradiction, therefore $k(\psi C_\varphi) = 0$, that is, ψC_φ is not surjective from \mathcal{L} to L^∞ .

REFERENCES

- [1] R. F. Allen, F. Colonna, and G. R. Easley, *Composition operators on the Lipschitz space of a tree*, *Mediterr. J. Math.* **11** (2014), 97–108.
- [2] R. F. Allen, I. M. Craig, *Multiplication operators on weighted Banach spaces of a tree*, *Bull. Korean Math. Soc.* **54** (2017), 747–761.
- [3] J. M. Cohen and F. Colonna, *Embeddings of trees in the hyperbolic disk*, *Complex Variables* **94** (1994), 311–335.
- [4] F. Colonna, and G. R. Easley, *Multiplication operators on the Lipschitz space of a tree*, *Integr. Equ. Oper. Theory* **68** (2010), 391–411.
- [5] F. Colonna, and G. R. Easley, *Multiplication operators between the Lipschitz space and the space of bounded functions on a Tree*, *Mediterr. J. Math.* **9** (2012), 423–438.
- [6] T. Hosokawa, *Minimum moduli of weighted composition operators on algebras of analytic functions*, *Kodai Math. J.* **29** (2006), 248–254.
- [7] V. Müller, *Spectral Theory of Linear Operators*, Birkhäuser, Basel, 2003.

WEIGHTED COMPOSITION OPERATORS BETWEEN SPACES ON TREE 17

COLLEGE OF ENGINEERING, IBARAKI UNIVERSITY, 4-12-1, NAKANARUSAWA,
HITACHI, IBARAKI, 316-8511, JAPAN

E-mail address: `takuya.hosokawa.com@vc.ibaraki.ac.jp`