

Extension of the Helleseht-Zinoviev result on the system of equations from Goethals codes and Kloosterman sums

Minglong Qi · Shengwu Xiong

Received: date / Accepted: date

Abstract In the paper of Tor Helleseht and Victor Zinoviev (Designs, Codes and Cryptography, **17**, 269-288(1999)), the number of solutions of the system of equations from Z_4 -linear Goethals codes G_4 was determined and stated in Theorem 4. We found that Theorem 4 is wrong for m even. In this note, we complete Theorem 4, and give new divisibility modulo 12 of the Kloosterman sums deduced from Theorem 4, which is different from the result of the same authors for $K(a^4 + a^3)$ modulo 12.

Keywords Z_4 -linear Goethals codes · nonlinear system of equations · exponential sums · Kloosterman sums

Mathematics Subject Classification (2000) 11L05 · 11T23

1 Introduction

Let m be a positive integer, $q = 2^m$, $F := \mathbb{F}_q$ be the finite field of q elements, $F^* := F \setminus \{0\}$, and $F^{**} := F \setminus \{0, 1\}$. The well known Kloosterman sums [1, 2] are defined by

$$K(a) = \sum_{x \in F^*} (-1)^{\text{Tr}(ax+1/x)} \quad (1)$$

where $a \in F^*$, and $\text{Tr}(\cdot)$ is the trace function of F over \mathbb{F}_2 .

Let $b, c \in F$. The problem of finding the coset weight distribution of Z_4 -linear Goethals codes G_4 , is transformed into solving the following nonlinear system of

Communicated by Pascale Charpin, Alexander Pott, Dieter Jungnickel

Minglong Qi · Shengwu Xiong
School of Computer Science and Technology, Wuhan University of Technology
Mafangshan West Campus, 430070 Wuhan City, China
E-mail: mlqiecully@163.com (Minglong Qi)
E-mail: xiongsw@whut.edu.cn (Shengwu Xiong)

equations over F [3]:

$$\begin{cases} x + y + z + u & = 1 \\ u^2 + xy + xz + xu + yz + yu + zu & = b^2 \\ x^3 + y^3 + z^3 + u^3 & = c \end{cases} \quad (2)$$

where x, y, z and u are pairwise distinct elements of F . The number of solutions of (2) (see p.284 of [3]), denoted by $\mu_2(b, c)$, is given by

$$\mu_2(b, c) = \frac{2}{3} \times \begin{cases} M_2(b, c) & \text{if } \text{Tr}(c) \neq \text{Tr}(1), \\ M_2(b, c) - 1 & \text{if } \text{Tr}(c) = \text{Tr}(1). \end{cases} \quad (3)$$

The $M_2(b, c)$ in (3) is defined by

$$4M_2(b, c) = \sum_{v \in F^{**}} (1 + (-1)^{\text{Tr}(\frac{k_1}{v})+l}) (1 + (-1)^{\text{Tr}(\frac{k_2}{v})+l}) \quad (4)$$

where $k_1 = b^2 + c + 1$, $k_2 = b^2 + b + c + \sqrt{c}$, and $l = \text{Tr}(b)$. The explicit evaluation of $M_2(b, c)$ is given by the following equation (see (33) of [3]):

$$M_2(b, c) = \frac{1}{4} (q - 3 + (-1)^{\text{Tr}(k_1)} K(k_1 k_2) - (-1)^{\text{Tr}(b)} (3 + (-1)^{\text{Tr}(k_1)})). \quad (5)$$

We found that Theorem 4 of [3], which gives the explicit evaluation of $\mu_2(b, c)$, is wrong for m even. It is obvious that the authors of [3] forgot to take account of the fact, that $\text{Tr}(1) = 1$ for m odd and 0 for m even, in the last step of the proof of Theorem 4. The following is the correct version of Theorem 4 of [3]:

Theorem 1 *Let $\mu_2(b, c)$ be the number of different 4-tuples (x, y, z, u) , where x, y, z, u are pairwise distinct elements of F , which are solutions to the system (2) over F , where b, c are arbitrary elements of F .*

(1) *If m is odd and $\text{Tr}(c) = 1$ or m is even and $\text{Tr}(c) = 0$, then*

$$\mu_2(b, c) = \frac{1}{6} (q - 8 + (-1)^{\text{Tr}(b)} (K(k_1 k_2) - 3)).$$

(2) *If m is odd and $\text{Tr}(c) = 0$ or m is even and $\text{Tr}(c) = 1$, then*

$$\mu_2(b, c) = \frac{1}{6} (q - 2 - (-1)^{\text{Tr}(b)} (K(k_1 k_2) + 3)).$$

Where $k_1 = b^2 + c + 1$ and $k_2 = b^2 + b + c + \sqrt{c}$.

From Theorem 4 of [3], the divisibility modulo 12 of the Kloosterman sums $K(a^4 + a^3)$ is obtained and stated in Theorem 5 of [3]. In Section 2, we will establish new divisibility modulo 12 of $K(a^4 + a^3)$.

2 New divisibility modulo 12 of $K(a^4 + a^3)$

Before stating the new results on $K(a^4 + a^3)$ modulo 12, we need several auxiliary results.

Lemma 1 ([1]) *Let $a \in F^*$. Then, $K(a) = K(a^2)$.*

Lemma 2 ([4]) *The quadratic equation $x^2 + ax + b = 0$, where $a \in F^*$ and $b \in F$, has two distinct roots in F , if and only if $\text{Tr}(b/a^2) = 0$.*

Lemma 3 ([4]) *Let $a, b \in F$. Then,*

- (1) $\text{Tr}(a) = \text{Tr}(a^2)$, $\text{Tr}(a + b) = \text{Tr}(a) + \text{Tr}(b)$.
- (2) $\text{Tr}(1) = 1$ if m is odd, and $\text{Tr}(1) = 0$ if m is even.

The following theorem shows that $K(a^4 + a^3)$ modulo 12 may have different values than Theorem 5 of [3] if a is under certain restriction:

Theorem 2 *Let $a = b^2 + b + 1$ where $b \in F$. If $a \neq 0$, then*

$$K(a^4 + a^3) \equiv \begin{cases} 3 & (\text{mod } 12), \text{ if } m \equiv 1 \pmod{2}, \\ 7 & (\text{mod } 12), \text{ if } m \equiv 0 \pmod{2}. \end{cases}$$

Proof Set $k_2 = b^2 + \sqrt{c}$, i.e., $b^2 + b + c + \sqrt{c} = b^2 + \sqrt{c}$. We obtain $b = c$, and $k_1 = b^2 + b + 1$. Further, we have $\text{Tr}(b) = \text{Tr}(c)$, $\text{Tr}(k_1) = \text{Tr}(b^2 + b + 1) = \text{Tr}(1)$, and $k_1^2 k_2^2 = b^8 + b^6 + b^4 + b^5 + b^3 + b$. Let $a = b^2 + b + 1$. It is easy to see that $k_1^2 k_2^2 = b^8 + b^6 + b^4 + b^5 + b^3 + b = a^4 + a^3$ in F . By Lemma 1, $K(k_1 k_2) = K((k_1 k_2)^2) = K(a^4 + a^3)$. Substituting $\text{Tr}(k_1) = \text{Tr}(1)$ and $\text{Tr}(b) = \text{Tr}(c)$ into (5), using (3), and taking account that $\text{Tr}(1) = 1$ if m is odd and 0 if m is even, we obtain that

- (1) Case that m is odd:

$$\mu_2(b, c) = \frac{1}{6}(q - 5 - K(a^4 + a^3)) \text{ for } \text{Tr}(c) \in \{0, 1\}.$$

- (2) Case that m is even:

$$\mu_2(b, c) = \begin{cases} \frac{1}{6}(q - 11 + K(a^4 + a^3)), & \text{if } \text{Tr}(c) = 0, \\ \frac{1}{6}(q + 1 + K(a^4 + a^3)), & \text{if } \text{Tr}(c) = 1. \end{cases}$$

The number $6\mu_2(b, c)$ is a multiple of 12 [3]. It can be verified that $(2^m - 5) \equiv 3 \pmod{12}$ if m is odd, and $-(2^m - 11) \equiv -(2^m + 1) \equiv 7 \pmod{12}$ if m is even. The theorem follows from above argument.

□

Based on computer investigation, we propose the following conjecture:

Conjecture 1 *Let $a = b^2 + b + 1$ where $b \in F$, and $\xi = \sum_{i=0}^n a^{2i} + a$ with $n \geq 0$. If $\xi \neq 0$, then*

$$K(\xi^4 + \xi^3) \equiv \begin{cases} 3 & (\text{mod } 12), \text{ if } m \equiv 1 \pmod{2}, \\ 7 & (\text{mod } 12), \text{ if } m \equiv 0 \pmod{2}. \end{cases}$$

To illustrate Conjecture 1, we present the following theorem:

Theorem 3 Let $\xi = b^4 + b + 1$ where $b \in F$. If $\xi \neq 0$, then

$$K(\xi^4 + \xi^3) \equiv \begin{cases} 3 & (\text{mod } 12), \text{ if } m \equiv 1 \pmod{2}, \\ 7 & (\text{mod } 12), \text{ if } m \equiv 0 \pmod{2}. \end{cases}$$

Proof Let $a = b^2 + b + 1$ where $b \in F$. Remark that $\xi = a^2 + a + 1$. Set $k_2 = b^2 + b + c + \sqrt{c} = c + \sqrt{b}$. Then, $c = b^4 + b^2 + b$ and $k_1 = b^4 + b + 1$. Further, we obtain that $\text{Tr}(c) = \text{Tr}(b^4 + b^2 + b) = \text{Tr}(b)$, $\text{Tr}(k_1) = \text{Tr}(b^4 + b + 1) = \text{Tr}(1)$, and $k_1^2 k_2^2 = (b^8 + b^2 + 1)(b^8 + b^4 + b^2 + b) = \xi^4 + \xi^3$ in F . The rest of the proof is analogous to Theorem 2, we omit the details. \square

If $a \in F$ satisfies a certain condition, the binomial $a^4 + a^3$ of $K(a^4 + a^3)$ from Theorem 5 of [3] may be reduced into a monomial of degree one, which is illustrated by the following theorem:

Theorem 4 Let $a \in F^{**}$ be a third root of unity in F^* . Then,

$$K(a) \equiv \begin{cases} 7 & (\text{mod } 12), \text{ if } \text{Tr}(a) = 0, \\ 11 & (\text{mod } 12), \text{ if } \text{Tr}(a) = 1. \end{cases}$$

Proof Set $k_1 = b^2 + c + 1 = b + \sqrt{c}$, then $k_2 = b^2 + b + c + \sqrt{c} = 1$. The condition $k_1 = b^2 + c + 1 = b + \sqrt{c}$ holds if and only if $\xi^2 + \xi + 1 = 0$ where $\xi = (b + \sqrt{c}) \in F^{**}$, i.e., ξ is a third root of unity in F^* . Further, we have $K(k_1 k_2) = K(\xi)$. The rest of the proof is similar to Theorem 2, we omit the details. \square

3 Conclusion

In this note, we pointed out that Theorem 4 of [3] is wrong for m even, and completed it by taking account of the fact that $\text{Tr}(1) = 1$ for m odd and 0 for m even in the proof of the theorem. We established some new divisibility modulo 12 of the Kloosterman sums $K(a^4 + a^3)$ too.

References

1. Carlitz, L.: Kloosterman sums and finite field extensions, *Acta Arithmetica*, **15**(2), 179-193 (1969).
2. Charpin P., Hellesteth T., Zinoviev V.: The divisibility modulo 24 of Kloosterman sums on $GF(2^m)$, m odd, *Journal of Combinatorial Theory, Series A*, **144**, 322-338 (2007).
3. Hellesteth T., Zinoviev V.: On Z_4 -Linear Goethals Codes and Kloosterman Sums, *Designs, Codes and Cryptography*, **17**, 269-288 (1999).
4. Lidl R., Niederreiter H.: *Finite Fields, Encyclopedia of Mathematics and Its Applications*, 2nd ed., Cambridge University Press, Vol. 20 (1997).