

Long-time asymptotic behavior for the discrete defocusing mKdV equation

Meisen Chen, Engui Fan*

School of Mathematical Sciences, Fudan University,
Shanghai 200433, P.R. China

December 24, 2019

Abstract

In this article, we apply Deift-Zhou nonlinear steepest descent method to analyze the long-time asymptotic behavior of the solution for the discrete defocusing mKdV equation

$$\dot{q}_n = (1 - q_n^2)(q_{n+1} - q_{n-1})$$

with decay initial value

$$q_n(t=0) = q_n(0),$$

where $n = 0, \pm 1, \pm 2, \dots$ is a discrete variable and t is continuous time variable. This equation was proposed by Ablowitz and Ladik.

Key words: discrete defocusing mKdV equation, Lax pair, Riemann-Hilbert problem, Deift-Zhou steepest descent method, long-time asymptotic behavior.

1 Introduction

Since Deift and Zhou developed nonlinear steepest descent method in 1993 [1], it has been successfully applied to analyze the long-time asymptotic behavior of a wide variety of continuous integrable systems, such as the mKdV equation, the

*Corresponding author: faneg@fudan.edu.cn

NLS equation, the sine-Gordon equation, the KdV equation, and the Cammasa-Holm equation [2–6]. However, there still has been little work on the long-time behavior of the discrete integrable systems, except for the Toda lattice and discrete NLS equation [7, 8].

In this article, we consider the following discrete defocusing mKdV equation

$$\dot{q}_n = (1 - q_n^2)(q_{n+1} - q_{n-1}) \quad (1)$$

with initial value

$$q_n(t = 0) = q_n(0), \quad (2)$$

where $\dot{q}_n = dq_n(t)/dt$, $n = 0, \pm 1, \pm 2, \dots$ and t is continuous time variable. The equation (1) was first proposed by Ablowitz and Ladik [9]. They further showed that the equation (1) is the semi-discrete version of the following classical mKdV equation [9]

$$u_\tau(x, \tau) + 6u^2u_x(x, \tau) - u_{xxx}(x, \tau) = 0.$$

Narita found two kinds of Miura transformations

$$q_n = \frac{(u_{n-1} + u_{n+1})u_n - 2u_{n-1}u_{n+1}}{(u_{n-1} - u_{n+1})u_n},$$

$$q_n = \frac{u_{n-1} - 2u_n + u_{n+1}}{u_{n-1} - u_{n+1}}$$

between the equation (1) and Sokolov-Shabat equation [10]

$$\dot{u}_n = 4(u_{n-1} - u_n)(u_n - u_{n+1})/(u_{n-1} - u_{n+1}).$$

The some kinds of exact solutions of equation (1) were obtained by using homotopy analysis method and exp-function method [11, 12]. With Darboux transformation, Wen and Gao obtained the explicit solutions for the equation (1) on the discrete spectral of Lax pair [13]. However, the solutions of the equation (1) with initial-boundary condition on the continuous spectral have been still unknown by using inverse scattering transformation or Riemann-Hilbert approach. So in this paper, we would like to apply Riemann-Hilbert approach/Deift-Zhou nonlinear steepest

descent method to investigate the long-time behavior of the solution for the initial value problem of discrete defocusing mKdV equation (1)-(2).

The organization of this paper is as follows. In section 2, we introduce appropriate eigenfunctions and spectral functions to reformulate initial value problem of discrete mKdV equation (1)-(2) as a Riemann-Hilbert problem(RHP). From section 3 to section 8, we transform the RHP to a model one by using a series of deformations and decompositions. In section 9, we show the existence and boundedness of the Cauchy operators during solving the RHP. At the last section, we provide the asymptotic behavior of the solution of the equation (1).

2 Riemann-Hilbert problem

In this section, we first investigate the solvability of the initial value problem (1)-(2) equation and then transform it into a RHP. Moreover, we further express the potential q_n of the discrete defocusing mKdV equation (1) with a solution of the obtained RHP.

Similar to Proposition 2.1 in [8], we can show the following proposition.

Proposition 2.1. *Let s be a nonnegative integer. If the initial condition (2) satisfies*

$$\|q_n(0)\|_{1,s} = \sum_{n=-\infty}^{\infty} (1 + |n|)^s |q_n(0)| < \infty, \quad (3)$$

$$\|q_n(0)\|_{\infty} = \sup_n |q_n(0)| < 1, \quad (4)$$

then the equation (1) admits an unique solution in the space

$$l^{1,s} = \{\{c_n\}_{n=-\infty}^{\infty} : \sum_{n=-\infty}^{\infty} (1 + |n|)^s |c_n| < \infty, 0 \leq t < \infty\}.$$

Proof. We define

$$c_{-\infty} = \prod_{n=-\infty}^{\infty} (1 - |q_n|^2), \quad \rho_0 = (1 - c_{-\infty})^{\frac{1}{2}},$$

then we have

$$\sup_n |q_n(0)| \leq \rho_0. \quad (5)$$

It is easy to verify that $c_{-\infty}$ and ρ_0 are both conserved quantities. Therefore, we consider equation (1) as an ordinary differential equation with respect to t , whose solution belongs to the Banach space $l^{1,s} \subset l^\infty$ under the condition (3) and (4).

Given the ball in l^∞

$$B := \{\{q_n(t)\} \in l^\infty, |q_n(t) - q_n(0)| < \rho_0\},$$

we show that the equation (1) admits a solution in the space B . Since the right-hand side of (1) is Lipschitz continuous and bounded, there exists a t_1 such that equation (1) admits solution in B for $t \in (0, t_1]$. By the standard argument about ordinary differential equations in [14], we get that t_1 depends on ρ_0 so as we have (5). Because we have known that ρ_0 is conserved, we have

$$\sup_n |q_n(t_1)| \leq \rho_0. \quad (6)$$

Similarly, we could extend the solution of equation (1) to $t \in (t_1, 2t_1]$. Repeating the procedure above, we then can extend the solution to $t \in [0, \infty)$ in the space l^∞ . Therefore, for $0 \leq t < \infty$, we have

$$\sup_n |q_n(t)| \leq \rho_0.$$

Based on the fact that there is a solution for (1)-(2) in l^∞ shown above, we further verify that the solution belongs to the Banach space $l^{1,s}$. From equation (1), we get

$$\|\dot{q}_n(t)\|_{1,s} \leq \text{Const.} \|q_n(t)\|_{1,s}.$$

By integrating $\dot{q}_n(t)$ with respect to t , we get

$$\|q_n(t)\|_{1,s} \leq \|q_n(0)\|_{1,s} + \text{Const.} \int_0^t \|q_n(\tau)\|_{1,s} d\tau.$$

By virtue of the Gronwall inequality, it follows that $\|q_n(t)\|_{1,s}$ grows at most exponentially and does not blow up.

For the uniqueness, if we set

$$q_n(0) = 0,$$

then the problem (1)-(2) only admits a zero solution. \square

In the following, we transform the initial problem (1)-(2) into a RHP. It is known that the discrete mKdV equation admits a Lax pair [9]

$$X_{n+1} = z^{\sigma_3} X_n + Q_n X_n, \quad (7)$$

$$\dot{X}_n = z^{2\sigma_3} X_n + B_n X_n, \quad (8)$$

where X_n is a 2×2 matrix, and

$$\sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad z^{\sigma_3} = \begin{pmatrix} z & 0 \\ 0 & z^{-1} \end{pmatrix}, \quad Q_n = \begin{pmatrix} 0 & q_n \\ q_n & 0 \end{pmatrix},$$

$$B_n = \begin{pmatrix} -q_{n-1}q_n & q_n z + q_{n-1}z^{-1} \\ q_n z^{-1} + q_{n-1}z & -q_{n-1}q_n \end{pmatrix}.$$

The Lax pair (7)-(8) admit the following asymptotic Jost solutions

$$X_n(z, t) \sim z^{n\sigma_3} \begin{pmatrix} e^{z^2 t} & 0 \\ 0 & e^{z^{-2} t} \end{pmatrix}, \quad n \rightarrow \pm\infty.$$

Making transformation

$$Y_n = z^{-n\sigma_3} \begin{pmatrix} e^{-z^2 t} & 0 \\ 0 & e^{-z^{-2} t} \end{pmatrix} X_n,$$

then we have

$$Y_{n+1} - Y_n = z^{-\sigma_3} \tilde{Q}_n Y_n, \quad (9)$$

where

$$\tilde{Q}_n = z^{-n\hat{\sigma}_3} e^{-\frac{t}{2}(z^2 - z^{-2})\hat{\sigma}_3} Q_n$$

$$= \begin{pmatrix} 0 & q_n z^{-2n} e^{-t(z^2 - z^{-2})} \\ q_n z^{2n} e^{t(z^2 - z^{-2})} & 0 \end{pmatrix}, \quad (10)$$

and for a 2×2 matrix A , the symbol $e^{\hat{\sigma}_3}$ is defined by

$$e^{\hat{\sigma}_3} A \equiv e^{\sigma_3} A e^{-\sigma_3}.$$

Denoting $Y_n^{(\pm)}$ as the 2×2 eigenfunctions of (9) such that

$$Y_n^{(-)} \rightarrow I \text{ as } n \rightarrow -\infty, \quad (11)$$

$$Y_n^{(+)} \rightarrow I \text{ as } n \rightarrow +\infty, \quad (12)$$

then we have

$$Y_n^{(-)} = I + \sum_{k=-\infty}^{n-1} z^{-\sigma_3} \tilde{Q}_k Y_k^{(-)}, \quad (13)$$

$$Y_n^{(+)} = I - \sum_{k=n}^{\infty} z^{-\sigma_3} \tilde{Q}_k Y_k^{(+)}. \quad (14)$$

By WKB expansion method, it follows that

$$Y_n^{(-)} = z^{-n\hat{\sigma}_3} e^{\frac{i}{2}(z^2 - z^{-2})\hat{\sigma}_3} \times \begin{pmatrix} 1 + O(z^{-2}, \text{even}) & zq_{n-1} + O(z^3, \text{odd}) \\ z^{-1}q_{n-1} + O(z^{-3}, \text{odd}) & 1 + O(z^2, \text{even}) \end{pmatrix}, \quad (15)$$

$$Y_n^{(+)} = z^{-n\hat{\sigma}_3} e^{\frac{i}{2}(z^2 - z^{-2})\hat{\sigma}_3} \times \begin{pmatrix} c_n^{-1} + O(z^2, \text{even}) & -c_n^{-1}z^{-1}q_n + O(z^{-3}, \text{odd}) \\ -c_n^{-1}zq_n + O(z^3, \text{odd}) & c_n^{-1} + O(z^{-2}, \text{even}) \end{pmatrix}, \quad (16)$$

where $O(z^{\pm 3}, \text{odd})$ ($O(z^{\pm 2}, \text{even})$) means the remaining part containing $z^{\pm 3}, z^{\pm 5}, \dots$ ($z^{\pm 2}, z^{\pm 4}, \dots$, respectively) and

$$c_n = \prod_{k=n}^{\infty} (1 - |q_k|^2).$$

Rewriting

$$Y_n^{(-)} = (Y_{n,1}^{(-)}, Y_{n,2}^{(-)}), \quad Y_n^{(+)} = (Y_{n,1}^{(+)}, Y_{n,2}^{(+)}),$$

where $Y_{n,j}^{(\pm)}$ is the j th column of $Y_n^{(\pm)}$, and letting

$$X_n^{(\pm)} = z^{n\sigma_3} \begin{pmatrix} e^{z^2 t} & 0 \\ 0 & e^{z^{-2} t} \end{pmatrix} Y_n^{(\pm)}, \quad (17)$$

from (11) and (12), we find that (7) admits matrix solutions $X_n^{(-)} = (X_{n,1}^{(-)}, X_{n,2}^{(-)})$ and $X_n^{(+)} = (X_{n,1}^{(+)}, X_{n,2}^{(+)})$ such that

$$X_{n,1}^{(-)} \rightarrow z^n e^{z^2 t} \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad X_{n,2}^{(-)} \rightarrow z^{-n} e^{z^{-2} t} \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \quad \text{as } n \rightarrow -\infty, \quad (18)$$

$$X_{n,1}^{(+)} \rightarrow z^n e^{z^2 t} \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad X_{n,2}^{(+)} \rightarrow z^{-n} e^{z^{-2} t} \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \quad \text{as } n \rightarrow \infty. \quad (19)$$

From spectral problem (7), we know that

$$\det(X_n^{(\pm)}) \neq 0,$$

which implies that $X_n^{(\pm)}$ are invertible. Thus, by the linearity of the eigenfunction, there exist four functions $a(z)$, $b(z)$, $a^*(z)$ and $b^*(z)$, such that

$$X_{n,1}^{(-)}(z, t) = a(z)X_{n,1}^{(+)}(z, t) + b(z)X_{n,2}^{(+)}(z, t), \quad (20)$$

$$X_{n,2}^{(-)}(z, t) = b^*(z)X_{n,1}^{(+)}(z, t) + a^*(z)X_{n,2}^{(+)}(z, t), \quad (21)$$

which combining with (7), we can get the symmetry

$$a^*(z) = \overline{a(\bar{z}^{-1})}, \quad b^*(z) = \overline{b(\bar{z}^{-1})}.$$

By using (17) and (20), we find that

$$a(z) = \frac{\det(X_{n,1}^{(-)}, X_{n,2}^{(+)})}{\det X_n^{(+)}} = \frac{\det(Y_{n,1}^{(-)}, Y_{n,2}^{(+)})}{\det Y_n^{(+)}}$$

$$b(z) = \frac{\det(X_{n,1}^{(+)}, X_{n,1}^{(-)})}{\det X_n^{(+)}} = \frac{\det(Y_{n,1}^{(+)}, Y_{n,1}^{(-)})}{\det Y_n^{(+)}}.$$

From (10), (13) and (14), it is worthy of notice that both $Y_{n,1}^{(-)}$ and $Y_{n,2}^{(+)}$ belong to $C[\{|z| \geq 1\} \cup \{\infty\}]$ and they are both analytic outside the unit circle. While

both $Y_{n,2}^{(-)}$ and $Y_{n,1}^{(+)}$ are continuous on the closed unit disc and analytic inside the unit circle. Thus, using (15)-(16), we know that $a(z)$ is analytic outside the unit circle, moreover $a \rightarrow 1$ as $z \rightarrow \infty$.

Besides, b is continuous on the unit circle, and

$$a^*(0) = 1. \quad (22)$$

By calculating the determinant of $X_n^{(-)}$ and $X_n^{(+)}$, we obtain that

$$|a(z)|^2 - |b(z)|^2 = c_{-\infty} > 0, \quad \text{for } |z| = 1,$$

which implies $a \neq 0$ on the unit circle. Let $r = b/a$, $\bar{r} = b^*/a^*$, we then have

$$\bar{r}(z) = \overline{r(\bar{z}^{-1})}, \quad 0 \leq |r(z)| < 1.$$

We now construct the RHP for equation (1). Defining a 2×2 analytic matrix function on $\mathbb{C} \setminus \{|z| = 1\}$

$$m(z; n, t) = \begin{cases} \begin{pmatrix} 1 & 0 \\ 0 & c_n \end{pmatrix} z^{n\hat{\sigma}_3} e^{-\frac{t}{2}(z^2 - z^{-2})\hat{\sigma}_3} \begin{pmatrix} Y_{n,1}^{(-)} \\ Y_{n,2}^{(+)} \end{pmatrix} & |z| > 1, \\ \begin{pmatrix} 1 & 0 \\ 0 & c_n \end{pmatrix} z^{n\hat{\sigma}_3} e^{-\frac{t}{2}(z^2 - z^{-2})\hat{\sigma}_3} \begin{pmatrix} Y_{n,1}^{(+)} \\ Y_{n,2}^{(-)} \end{pmatrix} & |z| < 1, \end{cases} \quad (23)$$

from (15)-(16), we can get the expansion of m

$$m = I + m_1 z^{-1} + \dots, \quad z \rightarrow \infty. \quad (24)$$

Moreover, with (15), (22) and (23), we find that

$$q_n = \frac{m_{12}}{z} \Big|_{z=0} = \frac{d}{dz} m_{12} \Big|_{z=0}, \quad (25)$$

where m_{12} is the (1,2)-entry of the matrix m .

By using (20) and (21), we can derive that

$$m^+ = m^- v, \quad z \in \Sigma, \quad (26)$$

where $\Sigma = \{z : |z| = 1\}$ is called jump curve (see Figure 1), and

$$v(z; n, t) = z^{n\hat{\sigma}_3} e^{-\frac{t}{2}(z^2 - z^{-2})\hat{\sigma}_3} \begin{pmatrix} 1 - |r(z)|^2 & -\bar{r}(z) \\ r(z) & 1 \end{pmatrix},$$

is called the jump matrix.

In summary, we have got the RHP of m

$$\begin{cases} m \text{ analytic on } \mathbb{C} \setminus \{|z| = 1\}, \\ m^+ = m^- v, & z \in \Sigma, \\ m(z; n, t) \rightarrow I & \text{as } z \rightarrow \infty. \end{cases}$$

Letting

$$\varphi = \frac{t}{2}(z^2 - z^{-2}) - n \log z, \quad (27)$$

we obtain that

$$v(z; n, t) = e^{-\varphi\hat{\sigma}_3} \begin{pmatrix} 1 - |r(z)|^2 & -\bar{r}(z) \\ r(z) & 1 \end{pmatrix}. \quad (28)$$




Figure 1. The jump curve Σ ;
'+' means the orientation of the contour

3 The first RHP deformation

In this section, we change the RHP of m into a new equivalent one, such that its jump matrix admits a helpful lower/upper triangular factorization. We take a

real constant number $0 < V_0 < 2$, we will discuss the asymptotic behavior under assumption $|n| \leq V_0 t$, $t \rightarrow \infty$.

From (27), the function φ has four first order stationary phase points

$$S_1 = A, S_2 = \bar{A}, S_3 = -A, S_4 = -\bar{A},$$

where

$$A = 2^{-1} \left(\sqrt{2 + \frac{n}{t}} - i\sqrt{2 - \frac{n}{t}} \right).$$




Figure 2. *The sign figure of $\text{Re}\varphi$*

Let

$$z = |z|e^{i\theta}.$$

From (27), we have

$$\text{Re}\varphi = \frac{t}{2} \cos(2\theta)(|z|^2 - |z|^{-2}) - n \log |z|,$$

which leads to the sign figure of $\text{Re}\varphi$ as shown in Figure 2.

We denote $\widehat{S_j S_{j+1}}$ as the arc on unit circle from S_j to S_{j+1} such that the central angle of the arc is less than π ($S_5 = S_1$). Define δ as an analytic function

on $\mathbb{C} \setminus \{|z| = 1\}$ satisfying an scalar RHP

$$\begin{cases} \delta_+(z) = \delta_-(z) (1 - |r|^2) & \text{on } \widehat{S_1 S_2} \cup \widehat{S_3 S_4}, \\ \delta_+(z) = \delta_-(z) & \text{on } \widehat{S_2 S_3} \cup \widehat{S_4 S_1}, \\ \delta \rightarrow 1 & \text{as } z \rightarrow \infty, \end{cases}$$

which admits a unique solution

$$\delta(z) = e^{-\frac{1}{2\pi i} \left(\int_{S_1}^{S_2} + \int_{S_3}^{S_4} \right) \log(1 - |r(\tau)|^2) \frac{d\tau}{\tau - z}}, \quad (29)$$

where $\int_{S_j}^{S_{j+1}}$, $j = 1, 3$ denote the integral on $\widehat{S_j S_{j+1}}$. Noticing that

$$\sup_{|z|=1} |r(z)| < 1,$$

it can be shown that both δ and δ^{-1} are bounded.

We introduce a transformation on $\mathbb{C} \setminus \{|z| = 1\}$

$$m^{(1)} = m\delta^{-\sigma_3}, \quad (30)$$

where the jump curve still is $\Sigma^{(1)} = \Sigma$, but its orientation is clockwise on $\widehat{S_1 S_2} \cup \widehat{S_3 S_4}$ and counterclockwise on $\widehat{S_2 S_3} \cup \widehat{S_4 S_1}$ as shown in Figure 3.

On Σ , we find that

$$(m\delta^{-\sigma_3})_+ = m_- v \delta_+^{-\sigma_3} = (m\delta^{-\sigma_3})_- \delta_-^{\sigma_3} v \delta_+^{-\sigma_3}, \quad (31)$$

where

$$\begin{aligned}
\delta_-^{\sigma_3} v \delta_+^{-\sigma_3} &= e^{-\varphi \hat{\sigma}_3} \begin{pmatrix} (1 - |r(z)|^2) \delta_- \delta_+^{-1} & -\bar{r}(z) \delta_+ \delta_- \\ r(z) \delta_-^{-1} \delta_+^{-1} & \delta_-^{-1} \delta_+ \end{pmatrix} \\
&= \begin{cases} e^{-\varphi \hat{\sigma}_3} \begin{pmatrix} 1 - |r(z)|^2 & -\bar{r}(z) \delta_+^2 \\ r(z) \delta_-^{-2} & 1 \end{pmatrix}, \\ e^{-\varphi \hat{\sigma}_3} \begin{pmatrix} 1 & -\bar{r}(z) \delta_+^2 \\ \frac{r(z) \delta_-^{-2}}{1 - |r(z)|^2} & 1 - |r(z)|^2 \end{pmatrix} \end{cases} \\
&= \begin{cases} \begin{pmatrix} 1 & -\bar{r}(z) \delta_+^2 e^{-2\varphi} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ r(z) \delta_-^{-2} e^{2\varphi} & 1 \end{pmatrix} & \text{on } \widehat{S_2 S_3} \cup \widehat{S_4 S_1}, \\ \begin{pmatrix} 1 & 0 \\ \frac{r \delta_-^{-2} e^{2\varphi}}{1 - |r|^2} & 1 \end{pmatrix} \begin{pmatrix} 1 & -\bar{r} \delta_+^2 e^{-2\varphi} \\ 0 & 1 \end{pmatrix} & \text{on } \widehat{S_1 S_2} \cup \widehat{S_3 S_4}. \end{cases}
\end{aligned}$$

Also, we verify that

$$m^{(1)} = m \delta^{-\sigma_3} \rightarrow I \quad \text{as } z \rightarrow \infty.$$

So if setting

$$v^{(1)} = \begin{cases} \delta_-^{\sigma_3} v \delta_+^{-\sigma_3} & \text{on } \widehat{S_1 S_2} \cup \widehat{S_3 S_4}, \\ (\delta_-^{\sigma_3} v \delta_+^{-\sigma_3})^{-1} & \text{on } \widehat{S_2 S_3} \cup \widehat{S_4 S_1}, \end{cases}$$

then we get RHP for $m^{(1)}$ as follows

$$m_+^{(1)} = m_-^{(1)} v^{(1)}, \quad z \in \Sigma^{(1)} \quad (32)$$

To express the jump matrix more concise, we set

$$\rho(z) = \begin{cases} -\frac{\bar{r}}{1 - |r|^2} & \text{on } \widehat{S_1 S_2} \cup \widehat{S_3 S_4}, \\ \bar{r} & \text{on } \widehat{S_2 S_3} \cup \widehat{S_4 S_1}, \end{cases} \quad (33)$$

and

$$\bar{\rho}(z) = \overline{\rho(\bar{z}^{-1})},$$

then we have

$$v^{(1)} = b_-^{-1} b_+, \quad (34)$$

where

$$b_- = \begin{pmatrix} 1 & 0 \\ \bar{\rho}\delta_-^{-2}e^{2\varphi} & 1 \end{pmatrix}, \quad b_+ = \begin{pmatrix} 1 & \rho\delta_+^2e^{-2\varphi} \\ 0 & 1 \end{pmatrix}. \quad (35)$$




Figure 3. The jump curve $\Sigma^{(1)}$

4 The second RHP deformation

In this section, we would decompose ρ and $\bar{\rho}$ into the rational part and two small parts respectively, so that we can make the second RHP transformation with augmented jump contour (see Figure 4).




Figure 4. Jump curve $\Sigma^{(2)}$
L is in blue and \bar{L} is in orange

Denoting

$$d = 2^{-1} \min(\sqrt{2 + n/t}, \sqrt{2 - n/t}),$$

and making the decomposition

$$\rho = R + h_I + h_{II}, \quad (36)$$

$$\bar{\rho} = \bar{R} + \bar{h}_I + \bar{h}_{II}, \quad (37)$$

we then can obtain the following estimates

$$|e^{-2\varphi} h_I| \leq Ct^{\frac{1}{2}-l} \quad \text{on } \Sigma, \quad (38)$$

$$|e^{-2\varphi} h_{II}| \leq Ct^{-q/2} \quad \text{on } L, \quad (39)$$

$$|e^{-2\varphi} R(z)| \leq Ce^{-Ct\epsilon^2} \quad \text{on } L^\epsilon, \quad (40)$$

$$|e^{2\varphi} \bar{h}_I| \leq Ct^{\frac{1}{2}-l} \quad \text{on } \Sigma, \quad (41)$$

$$|e^{2\varphi} \bar{h}_{II}| \leq Ct^{-\frac{q}{2}} \quad \text{on } \bar{L}, \quad (42)$$

$$|e^{2\varphi} \bar{R}(z)| \leq Ce^{-Ct\epsilon^2} \quad \text{on } \bar{L}^\epsilon, \quad (43)$$

where we can see L, \bar{L} in Figure 4 and for a positive real number $\epsilon (< d/2)$

$$L^\epsilon = \{z \in L : |z - S_j| \geq \epsilon \text{ for } j = 1, 2, 3, 4\},$$

$$\bar{L}^\epsilon = \{z \in \bar{L} : |z - S_j| \geq \epsilon \text{ for } j = 1, 2, 3, 4\}.$$

In the following, we first get down to the decomposition of ρ on the one side of $\widehat{S_1S_2}$; then we shall study the decomposition of $\bar{\rho}$ on the other side of $\widehat{S_1S_2}$; At last, in the similar way, we would investigate the decomposition on the other curves, like $\widehat{S_2S_3}$, $\widehat{S_3S_4}$ and $\widehat{S_4S_1}$.

4.1 Decomposition of ρ on $\widehat{S_1S_2}$

We consider ρ as a function of θ . If we set $\theta_0 = -arg(A)$, then the region of ρ is $(-\theta_0, \theta_0)$, which is a symmetric interval about $\theta = 0$. Thus, we could decompose ρ as the sum of an odd function and an even function

$$\rho(\theta) = H_e(\theta^2) + \theta H_o(\theta^2),$$

H_o and H_e can be approximated by Taylor's expansion at the point $\theta^2 = \theta_0^2$

$$H_o(\theta^2) = \mu_0^o + \cdots + \mu_k^o (\theta^2 - \theta_0^2)^k + O(\theta^2 - \theta_0^2)^{k+1},$$

$$H_e(\theta^2) = \mu_0^e + \cdots + \mu_k^e (\theta^2 - \theta_0^2)^k + O(\theta^2 - \theta_0^2)^{k+1}.$$

Define

$$R(\theta) = \sum_{l=0}^k (\mu_l^e (\theta^2 - \theta_0^2)^l + \theta \mu_l^o (\theta^2 - \theta_0^2)^l),$$

$$h(\theta) = \rho(\theta) - R(\theta),$$

$$\alpha(z) = (z - S_1)^q (z - S_2)^q,$$

where k and q are two fixed positive integers and have the relationship

$$k = 4q + 1.$$

Notice that

$$R(\pm\theta_0) = \rho(\pm\theta_0), \quad \theta = -i \log z, \quad \text{on } \widehat{S_1 S_2},$$

we consider R as a function of complex number z , which could be extended analytically to a fairly large neighborhood of $\widehat{S_1 S_2}$.

We consider a function

$$\psi = \varphi/(it),$$

then (27) implies that ψ strictly decrease on $(-\theta_0, \theta_0)$. Thus, we can consider θ as a function of ψ and h/α , which are defined by

$$(h/\alpha)(\psi) = \begin{cases} h(\theta(\psi)) / \alpha(\theta(\psi)), & \psi(\theta_0) \leq \psi \leq \psi(-\theta_0), \\ 0, & \text{otherwise on the real line.} \end{cases}$$

We verify that

$$(h/\alpha)(\theta) = O((\theta \pm \theta_0)^{k+1-q}), \quad \theta \rightarrow \pm\theta_0.$$

Since both S_1 and S_2 are the first order stationary phase points, one gets

$$\frac{d\theta}{d\psi} = \left(\frac{d\varphi}{itd\theta} \right)^{-1} = O((\theta \pm \theta_0)^{-1}), \quad \theta \rightarrow \pm\theta_0.$$

For any integer $1 \leq l \leq \frac{3q+2}{2}$, we deduce that

$$h/\alpha \in H_\psi^l, \tag{44}$$

where H_ψ^l norm is uniformly bounded with respect to (n, t) as long as $|n| \leq V_0 t$.

We apply Fourier transform to $(h/\alpha)(\theta)$ to decompose h into two parts

$$h = h_I + h_{II},$$

where

$$h_I(\theta) = \alpha(\theta) \int_t^\infty e^{is\psi(\theta)} (\widehat{h/\alpha})(s) ds, \tag{45}$$

$$h_{II}(\theta) = \alpha(\theta) \int_{-\infty}^t e^{is\psi(\theta)} (\widehat{h/\alpha})(s) ds, \tag{46}$$

and

$$\widehat{\left(\frac{h}{\alpha}\right)}(s) = \int_{-\infty}^{\infty} e^{-is\psi} \left(\frac{h}{\alpha}\right)(s) ds.$$

Thus, we have

$$\rho = R + h_I + h_{II}.$$

By using (44)-(45) and Schwartz inequality, we have

$$|e^{-2\varphi} h_I| \leq Ct^{\frac{1}{2}-l} \text{ on } \widehat{S_1 S_2}. \quad (47)$$

Let

$$p = \sqrt{d^2 + 2d + \frac{1}{2}} - \sqrt{\frac{1}{2}},$$

and introduce a curve

$$L_{12} = l_{12} \cup l_{21} \cup \tilde{L}_{12}, \quad (48)$$

where

$$\begin{aligned} l_{12} &= \{S_1 + S_1 e^{\frac{i\pi}{4}} u : 0 \leq u \leq p\}, \\ l_{21} &= \{S_2 + S_2 e^{-\frac{i\pi}{4}} (p - u) : 0 \leq u \leq p\}, \\ \tilde{L}_{12} &= \text{arc}(S_1 + S_1 e^{\frac{i\pi}{4}} p, S_2 + S_2 e^{-\frac{i\pi}{4}} p). \end{aligned}$$

From (46), It is shown that h_{II} can be analytically extend to $\{Re\varphi > 0\}$, and

$$|e^{-2\varphi} h_{II}| \leq C e^{-tRe(i\psi)} |\alpha|. \quad (49)$$

Since that both S_1 and S_2 are first order saddle points and \tilde{L}_{12} is a closed curve in the inner of $\{Re\varphi > 0\}$, we have

$$Rei\psi \geq \begin{cases} Cu^2 & \text{on } l_{12}, \\ C(p-u)^2 & \text{on } l_{21}, \\ C & \text{on } \tilde{L}_{12}. \end{cases} \quad (50)$$

It is easy to check that

$$|\alpha| = \begin{cases} O(u^q) & \text{on } l_{12}, \\ O((p-u)^q) & \text{on } l_{21}, \\ O(1) & \text{on } \tilde{L}_{12}. \end{cases} \quad (51)$$

With (50) and (51), we obtain

$$|e^{-2\varphi} h_{II}| \leq Ct^{-q/2} \quad \text{on } L_{12}. \quad (52)$$

For sufficiently small positive number ϵ ($\epsilon < d/2$), we define

$$\begin{aligned} l_{12}^\epsilon &= \{z \in l_{12} : |z - S_1| \geq \epsilon\}, \\ l_{21}^\epsilon &= \{z \in l_{21} : |z - S_2| \geq \epsilon\}, \\ L_{12}^\epsilon &= l_{12}^\epsilon \cup l_{21}^\epsilon \cup \tilde{L}_{12}. \end{aligned}$$

Since $\text{dist}(L_{12}^\epsilon, \{S_1, S_2\}) \geq \epsilon$, it follows that

$$\text{Re}(i\psi) \geq C\epsilon^2,$$

and with the boundedness of $R(z)$ on L_{12}^ϵ , we get

$$|e^{-2\varphi} R(z)| \leq Ce^{-Ct\epsilon^2} \quad \text{on } L_{12}^\epsilon. \quad (53)$$

4.2 Decomposition of $\bar{\rho}$ on $\widehat{S_1 S_2}$

For the one side of $\widehat{S_1 S_2}$, we have decomposed ρ into three parts

$$\rho = R + h_I + h_{II}.$$

Similar result can be obtained for $\bar{\rho}$ on $\widehat{S_1 S_2}$ and set

$$\bar{\rho} = \frac{\sqrt{2(2 - \frac{n}{t})}}{\sqrt{2 + \frac{n}{t}} + \sqrt{2 - \frac{n}{t}}}.$$

Consider the contour $\bar{L}_{12} = \bar{l}_{12} \cup \bar{l}_{21}$, where

$$\begin{aligned}\bar{l}_{12} &= \{S_1 - S_1 e^{-\frac{i\pi}{4}u} : 0 \leq u \leq \bar{p}\}, \\ \bar{l}_{21} &=: \{S_2 - S_2 e^{\frac{i\pi}{4}(\bar{p}-u)} : 0 \leq u \leq \bar{p}\}.\end{aligned}$$

Noticing decomposition

$$\bar{\rho} = \bar{R} + \bar{h}_I + \bar{h}_{II},$$

where

$$\bar{R}(z) = \overline{R(\bar{z}^{-1})},$$

then in a similar way to the derivation of h_I and h_{II} , we can get the estimates

$$|e^{2\varphi} \bar{h}_I| \leq Ct^{\frac{1}{2}-l} \quad \text{on } \widehat{S_1 S_2}, \quad (54)$$

$$|e^{2\varphi} \bar{h}_{II}| \leq Ct^{-\frac{q}{2}} \quad \text{on } \bar{L}_{12}. \quad (55)$$

Let

$$\bar{l}_{12}^\epsilon = \{z \in \bar{l}_{12} : |z - S_1| \geq \epsilon\},$$

$$\bar{l}_{21}^\epsilon = \{z \in \bar{l}_{21} : |z - S_2| \geq \epsilon\},$$

$$\bar{L}_{12}^\epsilon = \bar{l}_{12}^\epsilon \cup \bar{l}_{21}^\epsilon,$$

then for a fixed ϵ , we get

$$|e^{2\varphi} \bar{R}(z)| \leq Ce^{-Ct\epsilon^2} \quad \text{on } \bar{L}_{12}^\epsilon. \quad (56)$$

Remark 4.1. We have decomposed ρ and $\bar{\rho}$ on $\widehat{S_1 S_2}$; and we could get (36)-(37) on $\widehat{S_2 S_3}$, $\widehat{S_3 S_4}$ and $\widehat{S_4 S_1}$: $\rho = R + h_I + h_{II}$, similarly. What's more, we could get (38)-(43) in the same way.

4.3 The second RHP deformation

According to the decomposition for ρ and $\bar{\rho}$, we would make the second deformation to get a new RHP equivalent to the original one. Set

$$b_\pm = b_\pm^e + b_\pm^o,$$

where

$$\begin{aligned} b_+^e &= \begin{pmatrix} 1 & (R + h_{II})\delta^2 e^{-2\varphi} \\ 0 & 1 \end{pmatrix}, & b_+^o &= \begin{pmatrix} 1 & h_I \delta^2 e^{-2\varphi} \\ 0 & 1 \end{pmatrix}, \\ b_-^e &= \begin{pmatrix} 1 & 0 \\ (\bar{R} + \bar{h}_{II})\delta^{-2} e^{2\varphi} & 1 \end{pmatrix}, & b_-^o &= \begin{pmatrix} 1 & 0 \\ \bar{h}_I \delta^{-2} e^{2\varphi} & 1 \end{pmatrix}. \end{aligned}$$

Then, for (34)-(37), we have

$$v^{(1)} = (b_-^o b_-^e)^{-1} b_+^o b_+^e \text{ on } \Sigma^{(1)}.$$

Since both h_{II} and R is analytic on Ω_5 and Ω_6 , b_+^e is analytic on Ω_5 and Ω_6 ; similarly, b_-^e is analytic on Ω_3 and Ω_4 . Thus, we define a new holomorphic function on $\mathbb{C} \setminus \Sigma^{(2)}$ as shown in Figure 4:

$$m^{(2)} = \begin{cases} m^{(1)} & \text{on } \Omega_1 \text{ and } \Omega_2, \\ m^{(1)} (b_-^e)^{-1} & \text{on } \Omega_3 \text{ and } \Omega_4, \\ m^{(1)} (b_+^e)^{-1} & \text{on } \Omega_5 \text{ and } \Omega_6, \end{cases} \quad (57)$$

and it is easy to see that

$$\lim_{z \rightarrow \infty} m^{(2)} = \lim_{z \rightarrow \infty} m^{(1)} = I.$$

From (32) and (57), it follows that

$$m_+^{(2)} = m_-^{(2)} v^{(2)} \text{ on } \Sigma^{(2)}, \quad (58)$$

where

$$v^{(2)} = (b_-^{(2)})^{-1} b_+^{(2)} \quad (59)$$

and

$$b_+^{(2)} = \begin{cases} b_+^e & \text{on } L, \\ I & \text{on } \bar{L}, \\ b_+^o & \text{on } \Sigma, \end{cases} \quad b_-^{(2)} = \begin{cases} I & \text{on } L, \\ b_-^e & \text{on } \bar{L}, \\ b_-^o & \text{on } \Sigma \end{cases}$$

with

$$w_{\pm}^{(2)} = \pm(b_{\pm}^{(2)} - I), \quad w^{(2)} = w_{+}^{(2)} + w_{-}^{(2)}.$$

We have completed the second RHP transformation.

5 Reduction of the RHP

In this section, we would like to reduce the previous RHP to a model RHP. We firstly apply an integral operator introduced in [17] to write $m^{(2)}$ in the integral form. Then, we reduce this integral to the sum of the decaying part and the integral related to the leading order.

5.1 Partition of matrices

Recall that $m^{(2)}$ can be written in the integral form

$$m^{(2)}(z) = I + \frac{1}{2\pi i} \int_{\Sigma^{(2)}} \frac{((I - C_{w^{(2)}})^{-1}I)(\tau)w^{(2)}(\tau)}{\tau - z} d\tau, \quad (60)$$

where $C_{w^{(2)}} = C_{+}(\cdot w_{-}^{(2)}) + C_{-}(\cdot w_{+}^{(2)})$, and C_{\pm} are the Cauchy operators defined by

$$C_{\pm}(f)(z) = \frac{1}{2\pi i} \int_{\Sigma^{(2)}} \frac{f(\tau)d\tau}{\tau - z_{\pm}}, \quad f \in L^2(\Sigma^{(2)}), \quad z \in \Sigma^{(2)}.$$

By (24), (25), (30) and (57), we then get

$$\begin{aligned} q_n &= \frac{d}{dz} \left((m^{(2)})_{12} \delta^{-1} \right) \Big|_{z=0} = \delta(0)^{-1} \frac{d}{dz} (m^{(2)})_{12} \Big|_{z=0} \\ &= \frac{\delta(0)^{-1}}{2\pi i} \int_{\Sigma^{(2)}} z^{-2} \left[((I - C_{w^{(2)}})^{-1}I)(z)w^{(2)}(z) \right]_{12} dz. \end{aligned} \quad (61)$$

Now we would like to decompose $w_{\pm}^{(2)}$ into two parts

$$w_{\pm}^{(2)} = w'_{\pm} + w_{\pm}^c,$$

where

$$\begin{aligned}
w_{\pm}^c &= w_{\pm}^a + w_{\pm}^b, \\
w_{\pm}^a &: \begin{cases} w_{\pm}^a &= w_{\pm}^{(2)} \text{ on } \{|z|=1\}, \\ w_{\pm}^a &\text{is equal to the contribution to } w_{\pm}^{(2)} \text{ from terms of type} \\ &h_{II} \text{ and } \bar{h}_{II}, \end{cases} \\
w_{\pm}^b &: \begin{cases} w_{\pm}^b &= w_{\pm}^{(2)} - w_{\pm}^a \text{ on } L^\epsilon \cup \bar{L}^\epsilon, \\ w_{\pm}^b &= 0 \text{ on } \Sigma^{(2)} \setminus (L^\epsilon \cup \bar{L}^\epsilon). \end{cases}
\end{aligned}$$

So we set a contour consisting of four crosses

$$\Sigma' = \Sigma^{(2)} \setminus (\Sigma \cup L^\epsilon \cup \bar{L}^\epsilon) = \cup_{j=1}^4 \Sigma_j,$$

where Σ_j is the small cross connected to S_j for $j = 1, 2, 3, 4$. Note that the orientation of Σ' and Σ_j are determined by that of $\Sigma^{(2)}$ (see Figure 5).




Figure 5. Σ' consisting of four crosses

Lemma 5.1. *There exists a constant $C > 0$, such that*

$$|w_{\pm}^a|, |w^a| \leq Ct^{-1} \text{ on } \Sigma^{(2)}, \quad (62)$$

$$|w_{\pm}^b|, |w^b| \leq Ce^{-\gamma_\epsilon t} \text{ on } \Sigma^{(2)}. \quad (63)$$

Moreover, we have

$$\|w'_\pm\|_{L^2(\Sigma')}, \|w'\|_{L^2(\Sigma')} \leq Ct^{-1/4}, \quad (64)$$

$$\|w'_\pm\|_{L^1(\Sigma')}, \|w'\|_{L^1(\Sigma')} \leq Ct^{-1/2}. \quad (65)$$

Proof. Since δ and δ^{-1} is bounded, from the estimate (38)-(43), we get (62) and (63), and from (50), we get

$$|\delta^2 Re^{-2\varphi}| \leq Const.e^{-Ct|z-S_1|^2} \quad \text{on } l_{12} \cap \Sigma',$$

which leads to

$$\begin{aligned} \int_{l_{12} \cap \Sigma'} |\delta^2 Re^{-2\varphi}| dz &\leq Const.t^{-1/2}, \\ \int_{l_{12} \cap \Sigma'} |\delta^2 Re^{-2\varphi}|^2 dz &\leq Const.t^{-1/2}. \end{aligned}$$

Since all S_j are the first order stationary phase point of φ , we obtain that

$$\begin{aligned} \int_{L \cap \Sigma'} |\delta^2 Re^{-2\varphi}| dz &\leq Const.t^{-1/2}, \\ \int_{L \cap \Sigma'} |\delta^2 Re^{-2\varphi}|^2 dz &\leq Const.t^{-1/2}. \end{aligned}$$

For $\delta^{-2} \bar{R}e^{2\varphi}$ on $\bar{L} \cap \Sigma'$, we also have

$$\begin{aligned} \int_{\bar{L} \cap \Sigma'} |\delta^{-2} \bar{R}e^{2\varphi}| dz &\leq Const.t^{-1/2}, \\ \int_{\bar{L} \cap \Sigma'} |\delta^{-2} \bar{R}e^{2\varphi}|^2 dz &\leq Const.t^{-1/2}. \end{aligned}$$

Therefore, by simply analyzing the entry of w'_\pm , we deduce (64) and (65). \square

5.2 Some resolvent and estimates

We shall decompose

$$\int_{\Sigma^{(2)}} z^{-2} ((I - C_{w^{(2)}})^{-1} I)(z) w^{(2)}(z) dz,$$

into the principal part and the decaying part by the resolvent identity,

Under the assumption that both $(I - C_{w^{(2)}})^{-1}$ and $(I - C_{w'})^{-1}$ exist and are bounded, by the second resolvent identity, we have

$$\begin{aligned}
& \int_{\Sigma^{(2)}} z^{-2} (I - C_{w^{(2)}})^{-1} I w^{(2)} \\
= & \int z^{-2} (I - C_{w'})^{-1} I w' + \int z^{-2} w^c + \int z^{-2} (I - C_{w'})^{-1} (C_{w^c} I) w^{(2)} \\
& + \int z^{-2} (I - C_{w'})^{-1} (C_{w'} I) w^c + \int z^{-2} (I - C_{w'})^{-1} C_{w^c} (I - C_{w^{(2)}})^{-1} (C_{w^{(2)}} I) w^{(2)} \\
= & \int z^{-2} (I - C_{w'})^{-1} I w' + I + II + III + IV. \tag{66}
\end{aligned}$$

Since the length of $\Sigma^{(2)}$ is finite, by (62) and (63), we have

$$\| w_{\pm}^c \|_{L^s(\Sigma^{(2)})}, \| w^c \|_{L^s(\Sigma^{(2)})} \leq C t^{-1} \quad \text{for } s = 1, 2. \tag{67}$$

For $0 \notin \Sigma^{(2)}$, z^{-2} is bounded on $\Sigma^{(2)}$, so we obtain

$$|I| \leq C t^{-1}.$$

Because the Cauchy integral operator is bounded on $L^2(\Sigma^{(2)})$ and

$$\| C_{\pm} \|_{L^2 \rightarrow L^2} \leq 1,$$

we can get that

$$\begin{aligned}
\| C_{w^c} I \|_{L^2(\Sigma^{(2)})} & \leq \| C_+(w_-^c) \|_{L^2(\Sigma^{(2)})} + \| C_-(w_+^c) \|_{L^2(\Sigma^{(2)})} \\
& \leq \| w_-^c \|_{L^2(\Sigma^{(2)})} + \| w_+^c \|_{L^2(\Sigma^{(2)})} \leq C t^{-1}. \tag{68}
\end{aligned}$$

Similarly, from (64), we have

$$\| C_{w'} I \|_{L^2(\Sigma^{(2)})} = O(t^{-1/4}) \quad \text{as } t \rightarrow \infty. \tag{69}$$

From (62)-(63) and $\| C_{\pm} \|_{L^2 \rightarrow L^2} \leq 1$, for $f \in L^2(\Sigma^{(2)})$, we can show that

$$\begin{aligned}
\| C_{w^c} f \|_{L^2(\Sigma^{(2)})} & \leq \| C_+(f w_-^c) \|_{L^2(\Sigma^{(2)})} + \| C_-(f w_+^c) \|_{L^2(\Sigma^{(2)})} \\
& \leq C \| f \|_{L^2(\Sigma^{(2)})} \left(\sup_{z \in \Sigma^{(2)}} |w_-^c(z)| + \sup_{z \in \Sigma^{(2)}} |w_+^c(z)| \right) \\
& \leq C t^{-1} \| f \|_{L^2(\Sigma^{(2)})}, \tag{70}
\end{aligned}$$

by which, we obtain that

$$\begin{aligned}
|II| &\leq C \|C_{w^c} I\|_{L^2} \|w^{(2)}\|_{L^2} \\
&\leq C \|C_{w^c} I\|_{L^2} (\|w'\|_{L^2} + \|w^c\|_{L^2}) \\
&\leq Ct^{-1}, \\
|III| &\leq C \|C_{w'} I\|_{L^2} \|w^c\|_{L^2} \\
&\leq Ct^{-1}, \\
|IV| &\leq C \|C_{w^c}\|_{L^2 \rightarrow L^2} \|C_{w^{(2)}} I\|_{L^2} \|w^{(2)}\|_{L^2} \\
&\leq C \|C_{w^c}\|_{L^2 \rightarrow L^2} (\|C_{w^c} I\|_{L^2} + \|C_{w'} I\|_{L^2}) (\|w'\|_{L^2} + \|w^c\|_{L^2}) \\
&\leq Ct^{-1}.
\end{aligned}$$

From these estimates, combining (61) and (66), we obtain the following equation

$$q_n = \frac{\delta(0)^{-1}}{2\pi i} \int_{\Sigma^{(2)}} z^{-2} [(I - C_{w'})^{-1} I w']_{12}(z) dz + O(t^{-1}). \quad (71)$$

Let

$$C_{w'}^{\Sigma'} = C_+(\cdot \times w'_-|_{\Sigma'}) + C_-(\cdot \times w'_-|_{\Sigma'}),$$

which is an operator on 2×2 Hilbert space $L^2(\Sigma')$. Since w' vanishes on $\Sigma^{(2)} \setminus \Sigma'$, by (2.58) in [1], we could write (71) as

$$q_n = \frac{\delta(0)^{-1}}{2\pi i} \int_{\Sigma'} z^{-2} [(I - C_{w'}^{\Sigma'})^{-1} I w']_{12}(z) dz + O(t^{-1}), \quad (72)$$

where $C_{w'}$ in (71) is an operator on $L^2(\Sigma^{(2)})$, whose kernel is w' .

Remark 5.2. *We have assumed in this subsection that both $(I - C_{w^{(2)}})^{-1}$ and $(I - C_{w'})^{-1}$ exist and are bounded. In fact, in Section 9, we will prove that both $(I - C_{w^{(2)}})^{-1}$ and $(I - C_{w'})^{-1}$ exist and are bounded uniformly as $t \rightarrow \infty$.*

6 Four crosses

In this section, we may decompose w' into four parts according to the cross.

Moreover, we will make some estimates such that the principal part is more accurate for q_n .

Define matrix functions on Σ'

$$w^j = \begin{cases} w' & \text{on } \Sigma_j, \\ 0 & \text{on } \Sigma' \setminus \Sigma_j, \end{cases} \quad j = 1, 2, 3, 4, \quad (73)$$

then we define integral operators on $L^2(\Sigma')$ with kernels in (73)

$$A_j = C_{w^j}^{\Sigma'}, j = 1, 2, 3, 4. \quad (74)$$

For A_j , we have the following result.

Proposition 6.1. *Given $j, k = 1, 2, 3, 4$ ($j \neq k$), we have*

$$\|A_j A_k\|_{L^2 \rightarrow L^2} \leq C t^{-\frac{1}{2}}, \quad (75)$$

$$\|A_j A_k\|_{L^\infty \rightarrow L^2} \leq C t^{-\frac{3}{4}}. \quad (76)$$

The proof of this proposition is similar to Lemma 3.5 in [1]. Referring to (64) and (65), we only have to replace A' , B' in Lemma 3.5 of [1] with A_j , A_k respectively; then, Proposition 6.1 follows.

Assuming that as $t \rightarrow \infty$, $(1 - A_j)^{-1}$ exists and is uniformly bounded which would be proven in Section 9. because $C_{w'}^{\Sigma'} = \sum_j A_j$, by direct calculation, we get

$$1 - \sum_{j \neq k} A_j A_k (1 - A_k)^{-1} = (1 - C_{w'}^{\Sigma'}) [1 + \sum_j A_j (1 - A_j)^{-1}].$$

We verify that for any $f \in L(\Sigma')$,

$$\begin{aligned} \|A_j(f)\|_{L^2} &\leq \|C_+(f w_-^j)\|_{L^2} + \|C_-(f w_+^j)\|_{L^2} \\ &\leq (\|w_+^j\|_{L^\infty} + \|w_-^j\|_{L^\infty}) \|f\|_{L^2} \\ &\leq 2 \|w'_\pm\|_{L^\infty} \|f\|_{L^2} \leq C t^{-1} \|f\|_{L^2}, \end{aligned}$$

which implies

$$\|A_j\|_{L^2 \rightarrow L^2} = O(t^{-1}). \quad (77)$$

With Proposition 6.1, as $t \rightarrow \infty$, we have

$$(1 - C_{w'}^{\Sigma'})^{-1} = [1 + \sum_j A_j(1 - A_j)^{-1}][1 - \sum_{j \neq k} A_j A_k(1 - A_k)^{-1}]^{-1}, \quad (78)$$

then we obtain that

$$\begin{aligned} & \int z^{-2}[(1 - C_{w'}^{\Sigma'})^{-1}I](z)w'(z) \\ &= \int z^{-2}[I + \sum_j A_j(1 - A_j)^{-1}I](z)w'(z)dz + \int z^{-2}\{[1 + \sum_j A_j(1 - A_j)^{-1}] \\ & \quad \times [1 - \sum_{j \neq k} A_j A_k(1 - A_k)^{-1}]^{-1} \sum_{j \neq k} A_j A_k(1 - A_k)^{-1}I\}(z)w'(z)dz. \end{aligned} \quad (79)$$

and

$$A_j A_k(1 - A_k)^{-1} = A_j A_k + A_j A_k(1 - A_k)^{-1}A_k. \quad (80)$$

By the definition of A_k , we learn that $A_k I \in L^2(\Sigma')$ and

$$\|A_k I\|_{L^2} \leq C \|w^k\|_{L^2} \leq C \|w'\|_{L^2} \leq Ct^{-1/4}.$$

From the uniform boundedness of the operator $(1 - A_k)^{-1}$, we obtain $(1 - A_k)^{-1}A_k I$ belong to $L^2(\Sigma')$ and

$$\|(1 - A_k)^{-1}A_k I\|_{L^2} \leq Ct^{-1/4}. \quad (81)$$

Combining (75), (76), (80) and (81), we get that

$$[\sum_{j \neq k} A_j A_k(1 - A_k)^{-1}]I \in L^2(\Sigma'),$$

and

$$\left\| [\sum_{j \neq k} A_j A_k(1 - A_k)^{-1}]I \right\|_{L^2(\Sigma')} \leq Ct^{-3/4}.$$

If we can verified that

$$\left\| [1 + \sum_j A_j(1 - A_j)^{-1}][1 - \sum_{j \neq k} A_j A_k(1 - A_k)^{-1}]^{-1} \right\|_{L^2 \rightarrow L^2} \quad (82)$$

is bounded uniformly as $t \rightarrow \infty$, by Hölder's inequality, we can write (79) in the form

$$\begin{aligned} & \int z^{-2}[(1 - C_{w'}^{\Sigma'})^{-1}I](z)w'(z) \\ &= \int z^{-2}[I + \sum_j A_j(1 - A_j)^{-1}I](z)w'(z)dz + O(t^{-1}). \end{aligned} \quad (83)$$

With equation (64)-(65), we obtain that

$$[1 - \sum_{j \neq k} A_j A_k (1 - A_k)^{-1}]^{-1}, \quad t \rightarrow \infty$$

exists and is a bounded operator on $L^2(\Sigma')$. Also, by (77), we know that

$$1 + \sum_j A_j(1 - A_j)^{-1}$$

exists and is bounded on $L^2(\Sigma')$. Thus, we have proven the uniform boundedness of (82).

With (79), if we prove that for any pair of different numbers in $\{1, 2, 3, 4\} (j \neq k)$ and some constant C

$$\left| \int_{\Sigma'} z^{-2}[A_j(1 - A_j)^{-1}I](z)w^k(z) \right| \leq Ct^{-1}, \quad (84)$$

and have the following theorem.

Theorem 6.2. *The potential function q_n admits the asymptotic estimate*

$$q_n = \frac{\delta(0)^{-1}}{2\pi i} \sum_j \int_{\Sigma_j} z^{-2} \{[(1 - A_j)^{-1}I](z)w^j(z)\}_{12} dz + O(t^{-1}), \quad t \rightarrow \infty. \quad (85)$$

Proof. By using (72), (83) and (84), we could complete the proof of (85) by consider the following relation

$$A_j(1 - A_j)^{-1} = (1 - A_j)^{-1} - 1.$$

□

To prove (84), considering

$$[A_j(1 - A_j)^{-1}I]w^k = A_j(1 - A_j)^{-1}A_jIw^k + A_jIw^k,$$

we obtain the estimate

$$\begin{aligned} \int_{\Sigma_k} |A_jIw^k| &= \int_{\Sigma_k} \left| \left(\int_{\Sigma_j} \frac{w^j(\eta)d\eta}{\eta - \zeta} \right) w^k(\zeta) \right| d\zeta \\ &\leq C \|w^j\|_{L^1(\Sigma_j)} \|w^k\|_{L^1(\Sigma_k)} \leq Ct^{-1}, \end{aligned}$$

$$\begin{aligned} \int_{\Sigma_k} |A_j(1 - A_j)^{-1}A_jIw^k| &= \int_{\Sigma_k} \left| \left(\int_{\Sigma_j} \frac{(1 - A_j)^{-1}A_jI(\eta)w^j(\eta)d\eta}{\eta - \zeta} \right) w^k(\zeta) \right| d\zeta \\ &\leq C \int_{\Sigma_j} |(1 - A_j)^{-1}A_jI(\eta)w^j(\eta)|d\eta \|w^k\|_{L^1(\Sigma_k)} \\ &\leq C \|(1 - A_j)^{-1}A_jI\|_{L^2(\Sigma_j)} \|w^j\|_{L^1(\Sigma_j)} \|w^k\|_{L^1(\Sigma_k)} \\ &\leq C \|A_jI\|_{L^2(\Sigma_j)} \|w^j\|_{L^2(\Sigma_j)} \|w^k\|_{L^1(\Sigma_k)} \\ &\leq C \|A_jI\|_{L^2(\Sigma_j)} \|w^j\|_{L^2(\Sigma_j)}^2 \leq Ct^{-1}. \end{aligned}$$

Then (84) follows.

With Theorem 6.2, the original RHP can be reduced to four RHPs on four separated crosses and the leading order is only about these four RHPs.

7 Rotation and Scaling

In this section, we would introduce scaling operators concerning each stationary phase point.

Before getting down to the scaling operator, we shall first make some preparations. From (27), we show that for any $j = 1, 2, 3, 4$,

$$\varphi''(S_j) = (-1)^j 2i S_j^{-2} \sqrt{4t^2 - n^2}. \quad (86)$$

Let $T_1 = T_2 = 1$ and $T_3 = T_4 = -1$, we define the following quantities for $j = 1, 2, 3, 4$

$$\delta_j(z) = e^{(-1)^{j-1} \frac{1}{2\pi i} \int_{T_j}^{S_j} \log(1-|r(\tau)|^2) \frac{d\tau}{\tau-z}}, \quad (87)$$

$$\nu_j = -\frac{1}{2\pi} \log(1-|r(S_j)|^2), \quad l_j(z) = \int_{T_j}^{S_j} \frac{d\tau}{\tau-z}, \quad (88)$$

$$\chi_j(z) = \frac{1}{2\pi i} \int_{T_j}^{S_j} \log \frac{1-|r(\tau)|^2}{1-|r(S_j)|^2} \frac{d\tau}{\tau-z}, \quad (89)$$

where the symbol $\int_{T_j}^{S_j}$ stands for the integral on the arc $\widehat{T_j S_j}$ from T_j to S_j . These quantities have the relationship

$$\begin{aligned} \delta_j(z) &= e^{(-1)^{j-1}(\chi_j(z) + i\nu_j l_j(z))} \\ &= \left(\frac{S_j - z}{T_j - z} \right)^{(-1)^{j-1} \nu_j} e^{(-1)^{j-1} \chi_j(z)}. \end{aligned} \quad (90)$$

We verify that $\delta(z) = \prod_{j=1}^4 \delta_j(z)$ by making product directly and define

$$\hat{\delta}_j(z) = \delta(z) / \delta_j(z).$$

Since each S_j is first order stationary phase point, we have

$$\varphi(z) = \varphi(S_j) + \varphi''(S_j)(z - S_j)^2 + \varphi_j(z), \quad (91)$$

where

$$\varphi_j(z) = O(|z - S_j|^3).$$

We extend the four small crosses to four infinite ones

$$\begin{cases} \Sigma(S_j) = (S_j + S_j e^{i\pi/4} \mathbb{R}) \cup (S_j + S_j e^{-i\pi/4} \mathbb{R}), \\ \quad \text{oriented inward like } \Sigma_j \text{ for } j = 1, 3, \\ \Sigma(S_j) = (S_j + S_j e^{i\pi/4} \mathbb{R}) \cup (S_j + S_j e^{-i\pi/4} \mathbb{R}), \\ \quad \text{oriented outward like } \Sigma_j \text{ for } j = 2, 4, \end{cases}$$

and define the contours

$$\Sigma_j(0) = \begin{cases} (e^{i\pi/4} \mathbb{R}) \cup (e^{-i\pi/4} \mathbb{R}), \text{ oriented inward like } \Sigma_j, & j = 1, 3, \\ (e^{i\pi/4} \mathbb{R}) \cup (e^{-i\pi/4} \mathbb{R}), \text{ oriented outward like } \Sigma_j, & j = 2, 4. \end{cases}$$

In fact, we have a mapping from $\Sigma_j(0)$ to $\Sigma(S_j)$

$$\begin{aligned} M_j : \Sigma_j(0) &\rightarrow \Sigma(S_j), \\ z &\mapsto \beta_j z + S_j, \end{aligned}$$

where

$$\beta_j = \frac{1}{2}(4t^2 - n^2)^{-1/4} i S_j (-1)^j. \quad (92)$$

Directly calculating we find that

$$\varphi''(S_j) \beta_j^2 = (-1)^{j+1} \frac{i}{2}.$$

We introduce the scaling operator

$$\begin{aligned} N_j : (C^0 \cup L^2)(\Sigma(S_j)) &\rightarrow (C^0 \cup L^2)(\Sigma_j(0),) \\ f(z) &\mapsto (N_j f)(z) = f((\beta_j z + S_j), \end{aligned}$$

which is the pull-back of M_j .




Figure 6. *the mapping M_1^{-1}*

By equation (90) and (91), we obtain

$$\begin{aligned} N_j(e^{-\varphi})(z) &= e^{-\varphi(S_j) - \frac{\varphi''(S_j)}{2} \beta_j^2 z^2 - N_j \varphi_j(z)} \\ &= S_j^n e^{-\frac{i}{2}(S_j^2 - S_j^{-2}) + (-1)^j \frac{i}{4} z^2 - N_j \varphi_j(z)}, \end{aligned} \quad (93)$$

$$N_j \delta_j(z) = \delta_j(\beta_j z + S_j) + \left(\frac{\beta_j}{\beta_j z + S_j - T_j} \right)^{(-1)^{j-1} i \nu_j} z^{(-1)^{j-1} i \nu_j} \times e^{(-1)^{j-1} \chi_j(\beta_j z + S_j)}. \quad (94)$$

Further from (93) and (94), we get

$$N_j(\delta_j e^{-\varphi})(z) = \delta_j^0 \delta_j^1(z), \quad (95)$$

where

$$\delta_j^0 = S_j^n \left(\frac{\beta_j}{S_j - T_j} \right)^{(-1)^{j-1} i \nu_j} e^{(-1)^{j-1} \chi_j(S_j) - \frac{1}{2}(S_j^2 - S_j^{-2})} \hat{\delta}_j(S_j), \quad (96)$$

$$\delta_j^1(z) = \left(\frac{S_j - T_j}{\beta_j z + S_j - T_j} \right)^{(-1)^{j-1} i \nu_j} z^{(-1)^{j-1} i \nu_j} \frac{N_j \hat{\delta}_j(z)}{\hat{\delta}_j(S_j)} \times e^{(-1)^j \frac{1}{4} z^2 - N_j \varphi_j(z) + (-1)^{j-1} (\chi_j(\beta_j z + S_j) - \chi_j(S_j))}. \quad (97)$$

Note that $\Sigma(S_j)$ is extension of Σ_j . Let \hat{w}_\pm^j be the zero extension of $w_\pm^j|_{\Sigma_j}$ on $\Sigma(S_j)$; then, the related operator on $L^2(\Sigma(S_j))$ is denoted as $\hat{A}_j = C_{\hat{w}^j}$ with kernel

$$\hat{w}^j = \hat{w}_+^j + \hat{w}_-^j, \quad j = 1, 2, 3, 4.$$

Define a 2×2 matrix $\Delta_j^0 = (\delta_j^0)^{\sigma_3}$ and the related operator $\tilde{\Delta}_j^0: \tilde{\Delta}_j^0 \phi = \phi \Delta_j^0$ for 2×2 matrix ϕ ; then $\tilde{\Delta}_j^0$ and its inverse are bounded. Letting

$$\tilde{w}_\pm^j = (\Delta_j^0)^{-1} (N_j \hat{w}_\pm^j) \Delta_j^0, \quad \tilde{w}^j = \tilde{w}_+^j + \tilde{w}_-^j,$$

and $\alpha_j = C_{\tilde{w}^j} : L^2(\Sigma_j(0)) \rightarrow L^2(\Sigma_j(0))$, direct calculation shows that

$$\alpha_j = \tilde{\Delta}_j^0 N_j \hat{A}_j N_j^{-1} (\tilde{\Delta}_j^0)^{-1}, \quad (98)$$

$$\hat{A}_j = N_j^{-1} (\tilde{\Delta}_j^0)^{-1} \alpha_j \tilde{\Delta}_j^0 N_j. \quad (99)$$

Noticing that the support of $N_j \hat{w}^j$ belongs to $M_j^{-1} \Sigma_j$, by using (115), we obtain

$$(\Delta_j^0)^{-1} (N_j \hat{w}^j) \Delta_j^0 = (\Delta_j^0)^{-1} (N_j \hat{w}_+^j) \Delta_j^0 = \begin{pmatrix} 0 & R(\beta_j z + S_j) \delta_j^1(z)^2 \\ 0 & 0 \end{pmatrix}. \quad (100)$$

Similarly, on $M_j^{-1}(\Sigma_j \cap \bar{L}) \setminus \{0\}$, we have

$$(\Delta_j^0)^{-1} (N_j \hat{w}^j) \Delta_j^0 = (\Delta_j^0)^{-1} (N_j \hat{w}_-^j) \Delta_j^0 = \begin{pmatrix} 0 & 0 \\ -\bar{R}(\beta_j z + S_j) \delta_j^1(z)^{-2} & 0 \end{pmatrix}. \quad (101)$$

8 Convergence

Noticing that the principal part of q_n consists of four integrals respectively on four separate crosses: Σ_j ($j = 1, 2, 3, 4$), we consider the convergence of the principal part in this section. That is, we estimate four terms

$$\int_{\Sigma_j} z^{-2}[(1 - A_j)^{-1}I](z)w^j(z)dz, \quad j = 1, 2, 3, 4.$$

By (99), we have the following equalities

$$\begin{aligned} & \int_{\Sigma_j} z^{-2}[(1 - A_j)^{-1}I](z)w^j(z)dz \\ &= \beta_j \int_{\Sigma_j(0)} (\beta_j z + S_j)^{-2}[(1 - \alpha_j)^{-1}(\Delta_j^0)](z)(\Delta_j^0)^{-1}N_j \hat{w}^j(z)dz \\ &= \beta_j (\delta_j^0)^2 \int_{\Sigma_j(0)} [(1 - \alpha_j)^{-1}I](z)(\Delta_j^0)^{-1}N_j(\cdot \times \hat{w}^j)(z)\Delta_j^0 dz. \end{aligned} \quad (102)$$

We estimate the limit of formula (102). For this purpose, we investigate the convergence of the following functions

$$\begin{aligned} & (\beta_j z + S_j)^{-2}N_j R(z)\delta_j^1(z)^2, & N_j R(z)\delta_j^1(z)^2, \\ & (\beta_j z + S_j)^{-2}N_j \bar{R}(z)\delta_j^1(z)^{-2}, & N_j \bar{R}(z)\delta_j^1(z)^{-2}. \end{aligned}$$

Proposition 8.1. *For any arbitrary fixed constant γ ($0 < 2\gamma < 1$), on $M_j^{-1}(\Sigma_j \cap L) \cap \{z : \pm z/e^{i\pi/4} > 0\}$ respectively, we have*

$$\begin{aligned} & \left| (\beta_j z + S_j)^{-2}N_j R(z)\delta_j^1(z)^2 - S_j^{-2}R(S_j \pm)e^{-iz^2/2}z^{2iv_j} \right| \\ & \leq C e^{-\frac{1}{2}\gamma z^2} t^{-\frac{1}{2}} \log t, \end{aligned} \quad (103)$$

$$\begin{aligned} & \left| N_j R(z)\delta_j^1(z)^2 - R(S_j \pm)e^{-iz^2/2}z^{2iv_j} \right| \\ & \leq C e^{-\frac{1}{2}\gamma z^2} t^{-\frac{1}{2}} \log t, \end{aligned} \quad (104)$$

where $R(S_j+) = \bar{r}(S_j)$ on $M_j^{-1}(\Sigma_j \cap L) \cap \{z : z/e^{i\pi/4} > 0\}$ and $R(S_j-) = -\frac{\bar{r}(S_j)}{1-|\bar{r}(S_j)|^2}$ on $M_j^{-1}(\Sigma_j \cap L) \cap \{z : z/e^{i\pi/4} < 0\}$.

This proposition can be proved in a similar way to proposition 10.1 in [8], we omit it here. Similarly, we also have the following proposition.

Proposition 8.2. *For any arbitrary fixed constant γ ($0 < 2\gamma < 1$), on $M_j^{-1}(\Sigma_j \cap \bar{L}) \cap \{z : \pm z e^{-i\pi/4} > 0\}$ respectively, we have*

$$\begin{aligned} & \left| (\beta_j z + S_j)^{-2} N_j \bar{R}(z) \delta_j^1(z)^{-2} - S_j^{-2} \bar{R}(S_j \pm) e^{iz^2/2} z^{-2i\nu_j} \right| \\ & \leq C e^{\frac{i}{2}\gamma z^2} t^{-\frac{1}{2}} \log t, \end{aligned} \quad (105)$$

$$\left| N_j \bar{R}(z) \delta_j^1(z)^{-2} - \bar{R}(S_j \pm) e^{iz^2/2} z^{-2i\nu_j} \right| \leq C e^{\frac{i}{2}\gamma z^2} t^{-\frac{1}{2}} \log t, \quad j = 1, 3, \quad (106)$$

where $\bar{R}(S_j +) = r(S_j)$ on $M_j^{-1}(\Sigma_j \cap \bar{L}) \cap \{z : z e^{i\pi/4} > 0\}$ and $\bar{R}(S_j -) = -\frac{r(S_j)}{1-|r(S_j)|^2}$ on $M_j^{-1}(\Sigma_j \cap \bar{L}) \cap \{z : z e^{i\pi/4} < 0\}$.

Remark 8.3. *For even number j , in the same way, we can get the following result*

$$\begin{aligned} & (\beta_j z + S_j)^{-2} N_j R(z) \delta_j^1(z)^2 = S_j^{-2} R(S_j \pm) e^{iz^2/2} z^{-2i\nu_j} \\ & + O(e^{-\frac{i}{2}\gamma z^2} t^{-\frac{1}{2}} \log t), \quad \text{on } M_j^{-1}(\Sigma_j \cap L) \cap \{z : \pm z e^{i\pi/4} > 0\}, \\ & N_j R(z) \delta_j^1(z)^2 = R(S_j \pm) e^{iz^2/2} z^{-2i\nu_j} \\ & + O(e^{-\frac{i}{2}\gamma z^2} t^{-\frac{1}{2}} \log t), \quad \text{on } M_j^{-1}(\Sigma_j \cap L) \cap \{z : \pm z e^{i\pi/4} > 0\}, \\ & \beta_j z + S_j)^{-2} N_j \bar{R}(z) \delta_j^1(z)^{-2} = S_j^{-2} \bar{R}(S_j \pm) e^{iz^2/2} z^{-2i\nu_j} \\ & + O(e^{\frac{i}{2}\gamma z^2} t^{-\frac{1}{2}} \log t), \quad \text{on } M_j^{-1}(\Sigma_j \cap L) \cap \{z : \pm z / e^{i\pi/4} > 0\}, \\ & N_j \bar{R}(z) \delta_j^1(z)^{-2} = \bar{R}(S_j \pm) e^{iz^2/2} z^{-2i\nu_j} \\ & + O(e^{\frac{i}{2}\gamma z^2} t^{-\frac{1}{2}} \log t), \quad \text{on } M_j^{-1}(\Sigma_j \cap L) \cap \{z : \pm z / e^{i\pi/4} > 0\}. \end{aligned}$$

We consider some new matrices and operators that are limits of those on $\Sigma_j(0)$ which is a union of four parts

$$\Sigma_j(0) = \bigcup_{k=1}^k \Sigma_j^k(0),$$

where

$$\Sigma_j^k(0) = e^{i(2k-1)\pi/4} \mathbb{R}^+.$$

For $j = 1, 2, 3, 4$, we define

$$w_{\pm}^{j,\infty} = \lim_{t \rightarrow \infty} \tilde{w}_{\pm}^j, \quad w^{j,\infty} = w_+^{j,\infty} + w_-^{j,\infty}.$$

Defining operators

$$\alpha_j^\infty = C_{w^{j,\infty}} : L^2(\Sigma_j(0)) \rightarrow L^2(\Sigma_j(0)), \quad (107)$$

then by (104), (106) and Remark 8.3, we have

$$\begin{aligned} \|\tilde{w}_{\pm}^j - w_{\pm}^{j,\infty}\|_{L^2} &\leq Ct^{-1} \log t, \\ \|(C_{\tilde{w}^j} - C_{w^{j,\infty}})I\|_{L^2} &\leq Ct^{-1} \log t. \end{aligned}$$

Direct calculation shows that

$$\|(1 - \alpha_j^\infty)^{-1}I - (1 - \alpha_j)^{-1}I\|_{L^2} \leq Ct^{-1} \log t. \quad (108)$$

In summary, by using (85), (102), (103), (105), (108), Remark 8.3, we get that

$$\begin{aligned} q_n &= \frac{\delta(0)^{-1}}{2\pi i} \sum_{j=1}^4 \beta_j S_j^{-2} (\delta_j^0)^2 \left\{ \int_{\Sigma_j(0)} [(1 - \alpha_j^\infty)^{-1}I](z) w^{j,\infty}(z) dz \right\}_{12} \\ &\quad + O(t^{-1} \log t). \end{aligned} \quad (109)$$

For the branch cut of $z^{i\nu_j}$, we first consider that of δ_j^1 . From (97), it is reasonable to consider $M_j^{-1} \widehat{S_j T_j} \cup (\beta_j^{-1}(T_j - S_j) + \mathbb{R}^-)$ as the branch cut of δ_j^1 . As $t \rightarrow \infty$, we find that this branch cut becomes $\mathbb{R}^- = \{x \in \mathbb{R}\}$. Therefore, we consider \mathbb{R}^- as the branch cut of $z^{i\nu_j}$ in the remaining of this article.

9 The boundedness of operators

In this section, we would investigate the existence and boundedness of the following operators

$$\begin{aligned} &(I - C_{w^{(2)}})^{-1}, \quad (I - C_{w'})^{-1}, \quad (1 - A_j)^{-1}, \quad (1 - \hat{A}_j)^{-1}, \\ &(1 - \alpha_j)^{-1}, \quad (1 - \alpha_j^\infty)^{-1} \quad (j = 1, 2, 3, 4). \end{aligned}$$

Remark 9.1. Assume that operator $(1 - \alpha_j^\infty)^{-1}$ exists and is bounded on $L^2(\Sigma_j(0))$ for all j . With Proposition 8.1, 8.2 and Remark 8.3, we deduce that $(1 - \alpha_j)^{-1}$ exists and is uniformly bounded as $t \rightarrow \infty$. Because M_j is homeomorphic from $\Sigma_j(0)$ to $\Sigma(S_j)$, N_j is invertible and $\|N_j\|_{L^2(\Sigma(S_j)) \rightarrow L^2(\Sigma_j(0))}$ is bounded as well as $\|N_j^{-1}\|_{L^2(\Sigma(S_j)) \rightarrow L^2(\Sigma_j(0))}$. Moreover, noticing that Δ_j^0 and its inverse are invertible, we get the existence and boundedness of $(1 - \hat{A}_j)^{-1}$ from those of $(1 - \alpha_j)^{-1}$. More precisely, by equation (98), we have

$$(1 - \hat{A}_j)^{-1} = N_j^{-1}(\tilde{\Delta}_j^0)^{-1}(1 - \alpha_j)^{-1}\tilde{\Delta}_j^0 N_j.$$

By (2.58) in [1], we derive the existence and boundedness of $(1 - A_j)^{-1}$ from those of $(1 - \hat{A}_j)^{-1}$; thus, $(1 - C_{w'}^{\Sigma'})^{-1}$ exists and is uniformly bounded with equation (78). Moreover, by (2.59) in [1], $(I - C_{w'})^{-1}$ exists and is uniformly bounded.

Because of (62)-(63) and boundedness of $\Sigma^{(2)}$, we get that

$$\|C_{w^{(2)}} - C_{w'}\|_{L^2(\Sigma^{(2)}) \rightarrow L^2(\Sigma^{(2)})} \rightarrow 0.$$

Thus, by the second resolvent identity, we obtain that for sufficiently large t , the existence and boundedness of $(I - C_{w'})^{-1}$ implies those of $(I - C_{w^{(2)}})^{-1}$.

According to Remark 9.1, the remaining work in this section is to prove the existence and boundedness of $(1 - \alpha_j^\infty)^{-1}$. We discuss this problem in two cases when j is odd and j is even.

If j is odd, we change the orientation of $\Sigma_j^1(0)$ and $\Sigma_j^4(0)$ in $\Sigma_j(0)$. In fact, this change doesn't affect α_j^∞ as an operator on $L^2(\Sigma_j(0))$. So we could add the real line with orientation marked in Figure 7, we define matrices

$$w_\pm^{j,e} = \begin{cases} w_\pm^{j,\infty} & \text{on } \Sigma_j^2(0) \cup \Sigma_j^3(0), \\ -w_\mp^{j,\infty} & \text{on } \Sigma_j^1(0) \cup \Sigma_j^4(0), \\ 0 & \text{on } \mathbb{R}, \end{cases}$$

$$w^{j,e} = w_+^{j,e} + w_-^{j,e}, \quad \alpha_j^e = C_{w^{j,e}},$$

where

$$w_+^{j,e} = \begin{cases} \begin{pmatrix} 0 & 0 \\ r(S_j)e^{iz^2/2}z^{-2i\nu_j} & 0 \end{pmatrix} & z \in e^{i\pi/4}\mathbb{R}_+, \\ \begin{pmatrix} 0 & -\frac{\bar{r}(S_j)}{1-|r(S_j)|^2}e^{-iz^2/2}z^{2i\nu_j} \\ 0 & 0 \end{pmatrix} & z \in e^{3i\pi/4}\mathbb{R}_+, \\ 0 & \text{otherwise on } \Sigma^e, \end{cases}$$

$$w_-^{j,e} = \begin{cases} \begin{pmatrix} 0 & -\bar{r}(S_j)e^{-iz^2/2}z^{2i\nu_j} \\ 0 & 0 \end{pmatrix} & z \in e^{-i\pi/4}\mathbb{R}_+, \\ \begin{pmatrix} 0 & 0 \\ \frac{r(S_j)}{1-|r(S_j)|^2}e^{iz^2/2}z^{-2i\nu_j} & 0 \end{pmatrix} & z \in e^{-3i\pi/4}\mathbb{R}_+, \\ 0 & \text{otherwise on } \Sigma^e. \end{cases}$$

Then, by (2.58) in [1], we only need estimate the bound of $(1 - \alpha_j^e)^{-1}$.




Figure 7 Σ_j^e for j is odd

We write

$$v^{j,e} = (b_-^{j,e})^{-1}b_+^{j,e} = (1 - w_-^{j,e})^{-1}(1 + w_+^{j,e}),$$

and define a meromorphic function σ

$$\sigma = \begin{cases} z^{i\nu_j\sigma_3} & z \in \Omega_2^e \cup \Omega_5^e, \\ b_+^{j,e} z^{i\nu_j\sigma_3} & z \in \Omega_1^e \cup \Omega_3^e, \\ b_-^{j,e} z^{i\nu_j\sigma_3} & z \in \Omega_4^e \cup \Omega_6^e. \end{cases} \quad (110)$$

We simply denote $v^e = v^{j,e}$, also $\det \sigma = 1$ implies that σ is invertible. Let

$$v^{e,\sigma} = \sigma_-^{-1} v^e \sigma_+, \quad \text{on } \Sigma_j^e, \quad (111)$$

then we find that

$$v^{e,\sigma} = \begin{cases} v^0 := e^{-\frac{iz^2}{4}\hat{\sigma}_3} \begin{pmatrix} 1 & \bar{r}(S_j) \\ -r(S_j) & 1 - |r(S_j)|^2 \end{pmatrix} & \text{on } \mathbb{R}, \\ I & \text{on } \Sigma^e \setminus \mathbb{R}, \end{cases} \quad (112)$$

where

$$\begin{aligned} v^{e,\sigma} &= (b_-^{e,\sigma})^{-1} b_+^{e,\sigma} = (1 - w_-^{e,\sigma})^{-1} (1 - w_+^{e,\sigma}) \\ &= \begin{pmatrix} 1 & 0 \\ -r(S_j)e^{iz^2/2} & 1 \end{pmatrix} \begin{pmatrix} 1 & \bar{r}(S_j)e^{-iz^2/2} \\ 0 & 1 \end{pmatrix}. \end{aligned} \quad (113)$$

Let

$$w^{e,\sigma} = w_+^{e,\sigma} + w_-^{e,\sigma}.$$

Since the operator $C_{w^{e,\sigma}|_{\mathbb{R}}}: L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})$ satisfies

$$\|C_{w^{e,\sigma}|_{\mathbb{R}}}\| \leq \sup_{z \in \mathbb{R}} |e^{-iz^2/2} \bar{r}(S_j)| \leq \|r\|_{L^\infty(\mathbb{R})} < 1,$$

by Lemma 2.56 in [1], we have $\|C_{w^{e,\sigma}}\| < 1$, where $C_{w^{e,\sigma}}$ is an operator on $L^2(\Sigma^e)$. Thus $(1 - C_{w^{e,\sigma}})^{-1}$ exists and is bounded. Like the step 5 in section 3 of [1], we can deduce the existence and boundedness of $(1 - \alpha_j^e)^{-1}$. So the boundedness of $(1 - \alpha_j^\infty)^{-1}$ follows.

As for j is even, we could go through the process once again as j is odd, but we simply complete it by making conjugate transformation to $w_\pm^{j,e}$. Notice that for j is even, Σ_j^e admits the orientation that is different from the case of j is odd. Exactly, each line of Σ_j^e admits different orientation.

Let $w^{j,e} = w_+^{j,e} + w_-^{j,e}$, where $w_\pm^{j,e}$ are given by

$$w_\pm^{j,e} = \begin{cases} w_\pm^{j,\infty} & \text{on } \Sigma_j^2(0) \cup \Sigma_j^3(0), \\ -w_\mp^{j,\infty} & \text{on } \Sigma_j^1(0) \cup \Sigma_j^4(0), \\ 0 & \text{on } \mathbb{R}. \end{cases}$$

Further,

$$w_+^{j,e} = \begin{cases} \begin{pmatrix} 0 & 0 \\ r(S_j)e^{-iz^2/2}z^{2i\nu_j} & 0 \end{pmatrix} & z \in e^{-i\pi/4}\mathbb{R}_+, \\ \begin{pmatrix} 0 & -\frac{\bar{r}(S_j)}{1-|r(S_j)|^2}e^{iz^2/2}z^{-2i\nu_j} \\ 0 & 0 \end{pmatrix} & z \in e^{-3i\pi/4}\mathbb{R}_+, \\ 0 & \text{otherwise on } \Sigma^e, \end{cases}$$

$$w_-^{j,e} = \begin{cases} \begin{pmatrix} 0 & -\bar{r}(S_j)e^{iz^2/2}z^{-2i\nu_j} \\ 0 & 0 \end{pmatrix} & z \in e^{i\pi/4}\mathbb{R}_+, \\ \begin{pmatrix} 0 & 0 \\ \frac{r(S_j)}{1-|r(S_j)|^2}e^{-iz^2/2}z^{2i\nu_j} & 0 \end{pmatrix} & z \in e^{3i\pi/4}\mathbb{R}_+, \\ 0 & \text{otherwise on } \Sigma^e. \end{cases}$$

Define a conjugate operator T :

$$Tf(z) = \overline{f(\bar{z})}.$$

We could find that $Tw_{\pm}^{j,e}$ is almost the same as $w_{\pm}^{j,e}$ on Σ^e by replacing $\overline{r(S_j)}$ to $r(S_{j-1})$. Noticing that

$$\|\bar{r}\|_{L^\infty} = \|r\|_{L^\infty} < 1,$$

the proof of even number j follows as the proof of odd number j .

10 The asymptotic of discrete mKdV equation

In this section, we give the asymptotic behavior of q_n shown in equation (127).

By (109), we only have to estimate

$$\left\{ \int_{\Sigma_j(0)} [(1 - \alpha_j^\infty)^{-1}I](z)w^{j,\infty}(z)dz \right\}_{12}. \quad (114)$$

We expand the m^j in the form

$$\begin{aligned} m^j(z) &= I + \frac{1}{2\pi i} \int_{\Sigma_j(0)} \frac{[(1 - \alpha_j^\infty)^{-1}I](\tau)w^{j,\infty}(\tau)d\tau}{\tau - z} \\ &= I - z^{-1}m_1^j(z) + \dots, \end{aligned} \quad (115)$$

where

$$m_1^j(z) = \int_{\Sigma_j(0)} [(1 - \alpha_j^\infty)^{-1} I](z) w^{j,\infty}(z) \frac{dz}{2\pi i}. \quad (116)$$

And m^j is analytic on $\mathbb{C} \setminus \Sigma_j(0)$ and satisfy

$$m_+^j = m_-^j v^{j,\infty} \quad \text{on } \Sigma_j(0), \quad (117)$$

where

$$v^{j,\infty} = (1 - w_-^{j,\infty})^{-1} (1 + w_+^{j,\infty}).$$

For odd number j , we set

$$\Sigma_j^\infty = \Sigma_j^e,$$

and the orientation of Σ^∞ is different from Σ^e only on \mathbb{R} (See the Figure 8). Further we denote $\tilde{w}_\pm^{j,\infty}$ as the zero extension of $w_\pm^{j,\infty}$ on Σ_j^∞ .




Figure 8. Σ_j^∞ for j is odd

Define $\Phi(z) = m^j(z)\sigma(z)$ on $\mathbb{C} \setminus \Sigma^\infty$, then we would find that

$$\Phi_+ = \begin{cases} \Phi_- & \text{on } (e^{i\pi/4}\mathbb{R}) \cup (e^{-i\pi/4}\mathbb{R}) \setminus (0), \\ \Phi_-(v^0)^{-1} & \text{on } \mathbb{R}. \end{cases} \quad (118)$$

From the asymptotic behavior of m^j as $z \rightarrow \infty$, we have

$$\Phi(z)z^{-i\nu_j\sigma_3} = I - z^{-1}m_1^j + \dots. \quad (119)$$

which implies that Φ is analytic on $\mathbb{C} \setminus \mathbb{R}$. Letting $\hat{\Phi} = \Phi z^{-\frac{iz^2}{4}\sigma_3}$ on $\mathbb{C} \setminus \mathbb{R}$, we then get a model RHP

$$\begin{cases} \hat{\Phi}_+ = \hat{\Phi}_- \begin{pmatrix} 1 - |r(S_j)|^2 & -\overline{r(S_j)} \\ r(S_j) & 1 \end{pmatrix} \text{ on } \mathbb{R}, \\ \hat{\Phi} z^{-i\nu_j\sigma_3} e^{\frac{iz^2}{4}\sigma_3} \rightarrow I \text{ as } z \rightarrow \infty. \end{cases}$$

By using the result (110) in [16], we could get that

$$(m_1^j)_{12} = \frac{i(2\pi)^{1/2} e^{i\pi/4} e^{-\pi\nu_j/2}}{r(S_j)\Gamma(-i\nu_j)}, \quad j = 1, 3. \quad (120)$$

For the case when j is even, we can write m^j as

$$\begin{aligned} m^j &= I + \int_{\Sigma_j(0)} \frac{[(1 - C_{w^{j,e}})^{-1}I](\tau)w^{j,e}(\tau)d\tau}{\tau - z} \\ &= I - z^{-1}m_1^j(z) + \dots, \end{aligned} \quad (121)$$

where

$$m_1^j = \int_{\Sigma_j(0)} [(1 - C_{w^{j,e}})^{-1}I](z)w^{j,e}(z)dz. \quad (122)$$

Because $C_{w^{j,e}} = TC_{Tw^{j,e}}T$ and $T^2 = 1$, we obtain

$$C_{Tw^{j,e}} = TC_{w^{j,e}}T \quad (123)$$

and

$$\begin{aligned} &\int_{\Sigma_j(0)} [(1 - C_{Tw^{j,e}})^{-1}I](z)Tw^{j,e}(z)dz \\ &= \int_{\Sigma_j(0)} [T(1 - C_{w^{j,e}})^{-1}TI](z)Tw^{j,e}(z)dz \\ &= \int_{\Sigma_j(0)} T[(1 - C_{w^{j,e}})^{-1}I \times w^{j,e}](z)dz \\ &= \int_{\Sigma_j(0)} [(1 - C_{w^{j,e}})^{-1}I](z)w^{j,e}(z)dz. \end{aligned} \quad (124)$$

Therefore, in a similar way to the case when j is odd, we have

$$\left\{ \int_{\Sigma_j(0)} [(1 - C_{Tw^{j,e}})^{-1} I](z) Tw^{j,e}(z) dz \right\}_{12} = \frac{i(2\pi)^{1/2} e^{i\pi/4} e^{-\pi\nu_j/2}}{r(S_j)\Gamma(-i\nu_j)}. \quad (125)$$

Combining (124) and (125) gives

$$(m_1^j)_{12} = -\frac{i(2\pi)^{1/2} e^{-i\pi/4} e^{-\pi\nu_j/2}}{r(S_j)\Gamma(i\nu_j)}, \quad j = 2, 4, \quad (126)$$

where $\nu_j = -\frac{1}{2\pi} \log(1 - |r(S_j)|^2)$.

Finally, combining (109), (115), (116), (120) and (126) gives the asymptotic behavior of the discrete mKdV equation

$$q_n = \delta(0)^{-1} \sum_{j=1}^4 \beta_j S_j^{-2} (\delta_j^0)^2 (m_1^j)_{12} + O(t^{-1} \log t), \quad \text{for } |n| \leq V_0 t, \quad (127)$$

where S_j , $j = 1, 2, 3, 4$ are stationary points; β_j and δ_j^0 are given by (92) and (96) respectively, and

$$(m_1^j)_{12} = (-1)^{j-1} \frac{i(2\pi)^{1/2} e^{-i\pi/4} e^{-\pi\nu_j/2}}{r(S_j)\Gamma[(-1)^j i\nu_j]}, \quad j = 1, 2, 3, 4.$$

The asymptotic formula (127) is composed of a decaying term and a leading term coming from four stationary phase points S_j , $j = 1, 2, 3, 4$. In the leading term δ_j^0 contains three oscillatory factors: S_j^n , $e^{-\frac{i}{2}(S_j^2 - S_j^{-2})}$ and $\beta_j^{(-1)^{j-1} i\nu_j}$. Since n/t is fixed, as t tends to the infinity, n also tends to the infinity. If we set $\theta_j = \arg S_j$ and κ_j the imaginary part of S_j^2 , then we can write the three oscillatory terms in the form

$$S_j^n e^{-\frac{i}{2}(S_j^2 - S_j^{-2})} \beta_j^{(-1)^{j-1} i\nu_j} = e^{\frac{i}{2}(2n\theta_j - 2\kappa_j t + (-1)^j \nu_j \log t)} \psi_j(n/t),$$

where ψ_j is a function about n/t . Then, $\beta_j (\delta_j^0)^2$ behaves like $\text{const.} t^{-1/2} e^{ip_j t + iq_j \log t}$, $p_j \in \mathbb{R}$ and $q_j \in \mathbb{R}$.

In this article, we have got the long time asymptotic formula (127) for the solutions of the initial value problem for the discrete defocusing mKdV equation (1)-(2) by the Deift-Zhou steepest descent method. To our knowledge, with exception

to recent work on Toda lattice and discrete Schrödinger equation [7,8,18–20], there has been little work on asymptotic behavior for discrete integrable systems via the Deift-Zhou steepest descent method. There are almost no work on asymptotic behavior for discrete integrable systems with nonzero boundary conditions.

Acknowledgements Many thanks for referees for helpful suggestions in improving the manuscript. This work was supported by the National Science Foundation of China under Project No.11671095 and No.51879045.

References

- [1] Deift, P.A., Zhou, X.: A steepest descent method for oscillatory Riemann–Hilbert problems. Asymptotics for the MKdV equation. *Annals of Mathematics*. **137**(2). 295-368 (2017)
- [2] Deift, P.A., Its, A.R., Zhou, X.: Long-time asymptotics for integrable nonlinear wave equations. in *Important developments in soliton theory*. Springer. 181-204 (1993)
- [3] Cheng, P., Venakides, S., Zhou, X.: Long-time asymptotics for the pure radiation solution of the Sine-Gordon equation. *History and philosophy of logic*. **24**(7-8). 1195-1262 (1999)
- [4] Vartanian, A.H.: Higher order asymptotics of the modified non-linear Schrödinger equation. *Communications in Partial Differential Equations*. **25**(5-6), 1043-1098 (2000)
- [5] Grunert, K., Teschl, G.: Long-time asymptotics for the Korteweg–de Vries equation via nonlinear steepest descent. *Mathematical Physics, Analysis and Geometry*. **12**(3), 287-324 (2009)

- [6] De Monvel, A.B., Kostenko, A., Shepelsky, D., Teschl, G.: Long-time asymptotics for the Camassa–Holm equation. *SIAM Journal on Mathematical Analysis*. **41**(4), 1559-1588 (2009)
- [7] Krüger, H., Teschl, G.: Long-time asymptotics of the Toda lattice for decaying initial data revisited. *Reviews in Mathematical Physics*. **21**(1), 61-109 (2009)
- [8] Yamane, H.: Long-time asymptotics for the defocusing integrable discrete nonlinear Schrödinger equation. *Journal of the Mathematical Society of Japan*. **66**(3), 765-803 (2014)
- [9] Ablowitz, M.J.: Nonlinear evolution equations continuous and discrete. *Siam Review*. **19**(4), 663-684 (1977)
- [10] Narita, K.: Miura transformations between Sokolov-Shabat’s equation and the discrete MKdV equation. *Journal of the Physical Society of Japan*. **66**(12), 4047-4048 (1997)
- [11] Wang, Z., Zou, L., Zhang, H.Q.: Solitary solution of discrete mKdV equation by homotopy analysis method. *Communications in Theoretical Physics*. **49**(6), 1373 (2008)
- [12] Zhu, S.D.: Exp-function method for the discrete mKdV lattice. *International Journal of Nonlinear Sciences and Numerical Simulation*. **8**(3), 465-468 (2007)
- [13] Wen, X.Y., Gao, Y.T.: Darboux transformation and explicit solutions for discretized modified Korteweg-de Vries lattice equation. *Communications in Theoretical Physics*. **53**(5), 825 (2010)
- [14] Lang, S.: *Differential and Riemannian manifolds*. Springer Science & Business Media. **160** (2012)
- [15] Ablowitz, M.J., Ablowitz, M.A., Prinari, B., Trubatch, A.D., et al: *Discrete and continuous nonlinear Schrödinger systems*. Cambridge University Press. **302** (2004)

- [16] Deift, P.A., Zhou, X.: Long-time behavior of the non-focusing nonlinear Schrödinger equation-a case study. University of Tokyo, (1994).
- [17] Beals, R., Coifman, R.R.: Scattering and inverse scattering for first order systems. *Communications on Pure and Applied Mathematics*. **37**(1), 39-90 (1984)
- [18] Yamane, H.: Long-time asymptotics for the defocusing integrable discrete nonlinear Schrödinger equation II, *Symmetry Integrability and Geometry-Methods and Applications*, **11**, 020(2015)
- [19] Yamane, H.: Riemann-Hilbert factorization of matrices invariant under inversion in a circle, *Proceedings of the American Mathematical Society*, **147**, 2147-2157 (2019)
- [20] Yamane, H.: Long-time asymptotics for the defocusing integrable discrete nonlinear Schrödinger equation. *Funkcialaj Ekvacioj-Serio Inteenacia*, **62**, 227-253 (2019)