

Rationally weighted Hurwitz numbers, Meijer G -functions and matrix integrals

M. Bertola^{1,2,3*} and J. Harnad^{1,2†}

¹*Department of Mathematics and Statistics, Concordia University
1455 de Maisonneuve Blvd. W. Montreal, QC H3G 1M8 Canada*

²*Centre de recherches mathématiques, Université de Montréal,
C. P. 6128, succ. centre ville, Montréal, QC H3C 3J7 Canada*

³*SISSA/ISAS, via Bonomea 265, Trieste, Italy*

Tuesday 24th September, 2019

Abstract

The quantum spectral curve equation associated to KP τ -functions of hypergeometric type serving as generating functions for rationally weighted Hurwitz numbers is solved by generalized hypergeometric series. The basis elements spanning the corresponding Sato Grassmannian element are shown to be Meijer G -functions, or their asymptotic series. Using their Mellin integral representation the τ -function, evaluated at the trace invariants of an externally coupled matrix, is expressed as a matrix integral.

1 Introduction

Several instances of τ -functions of hypergeometric type are known to be generating functions for weighted Hurwitz numbers [21, 22, 24, 29], which enumerate branched covers of the Riemann sphere with specified branching profiles. These are distinguished by the choice of a weight generating function $G(z)$ that selects the weighting attributed to various possible branching configurations.

In particular, such τ -functions serve as generating functions for *simple* Hurwitz numbers [38, 39] (both single and double), *weakly monotonic* (or, equivalently, *signed*) Hurwitz numbers ([20, 22, 29]), *strongly monotonic* Hurwitz numbers [21, 22], weighted Hurwitz numbers for Belyi curves, or *dessins d'enfants* [1, 19, 33, 47], polynomially weighted Hurwitz numbers [1–7, 29]; *quantum* Hurwitz numbers [22, 24, 25] and *multispecies* Hurwitz numbers [26].

*e-mail: Marco.Bertola@concordia.ca, Marco.Bertola@sissa.it

†e-mail: harnad@crm.umontreal.ca

Of these, simple Hurwitz numbers, both weakly and strongly monotonic Hurwitz numbers and polynomially weighted Hurwitz numbers are all known to admit matrix integral representations [1–3, 7, 10, 12, 20, 40]. In this work, we derive a new matrix integral representation, of a different type, for hypergeometric KP τ -functions generating weighted Hurwitz numbers with rational weight generating function, which include several of the above as special cases.

In Section 2, the notion of weighted Hurwitz numbers [21, 22, 24, 29], is recalled and the KP τ -function serving as generating function for these is defined, focussing on the case of rational weight generating functions

$$G_{\mathbf{c},\mathbf{d}}(z) := \frac{\prod_{i=1}^L (1 + c_i z)}{\prod_{j=1}^M (1 - d_j z)}. \quad (1.1)$$

We define a natural basis $\{\phi_k^{(\mathbf{c},\mathbf{d},\beta)}(x)\}_{k \in \mathbf{N}^+}$ for the corresponding element of the infinite Sato Grassmannian, consisting of Laurent series obtained by dividing generalized hypergeometric series of the type

$${}_L F_M \left(\begin{matrix} 1 - k + \frac{1}{\beta c_1}, \dots, 1 - k + \frac{1}{\beta c_L} \\ 1 - k - \frac{1}{\beta d_1}, \dots, 1 - k - \frac{1}{\beta d_M} \end{matrix} \middle| \kappa_{\mathbf{c},\mathbf{d}} x \right), \quad k \in \mathbf{Z} \quad (1.2)$$

by the monomials x^{k-1} . These are either convergent or formal, depending on whether $L \leq M + 1$ or $L > M + 1$. They satisfy the sequence of eigenvalue equations with nonnegative integer eigenvalues,

$$\mathcal{L}_{\mathbf{c},\mathbf{d},\beta} \phi_k^{(\mathbf{c},\mathbf{d},\beta)} := (x G_{\mathbf{c},\mathbf{d}}(\beta \mathcal{D}) - \mathcal{D}) \phi_k^{(\mathbf{c},\mathbf{d},\beta)} = (k - 1) \phi_k^{(\mathbf{c},\mathbf{d},\beta)}, \quad k \in \mathbf{Z}, \quad (1.3)$$

where $\mathcal{D} = x \frac{d}{dx}$ is the Euler operator and $\mathcal{L}_{\mathbf{c},\mathbf{d},\beta}$ the quantum spectral curve operator. The first of these is the quantum curve equation appearing in the topological recursion approach [4–6].

Theorem 3.1, Section 3, relates the basis elements $\{\phi_k^{(\mathbf{c},\mathbf{d},\beta)}(x)\}_{k \in \mathbf{N}^+}$ to Barnes-Mellin type integrals

$$\tilde{\phi}_k^{(\mathbf{c},\mathbf{d},\beta)}(x) \sim \int_{\mathcal{C}_k} \frac{\Gamma(1 - k - s) \prod_{\ell=1}^L \Gamma\left(s + \frac{1}{\beta c_\ell}\right) (-\kappa_{\mathbf{c},\mathbf{d}} x)^s}{\prod_{m=1}^M \Gamma\left(s - \frac{1}{\beta d_m}\right)} ds \quad (1.4)$$

defining Meijer G -functions, either as convergent sums over the residues at the poles at $s = 1 - k, 2 - k, \dots$ (when $L \leq M + 1$) or as asymptotic series (when $L > M + 1$). In Section 4 a finite determinantal formula (4.7) is derived for the τ -function with KP flow parameters evaluated at the trace invariants of an n -dimensional matrix X . This leads to Theorem 5.1, expressing the τ -function as an externally coupled matrix integral of generalized Brézin-Hikami [11] type.

2 Generating functions for weighted Hurwitz numbers

2.1 Pure and weighted Hurwitz numbers

We recall the definition of *pure* Hurwitz numbers [16, 17, 30, 31, 34, 45].

Definition 2.1 (Combinatorial). For a set of k partitions $\{\mu^{(i)}\}_{i=1,\dots,k}$ of $N \in \mathbf{N}^+$, the *pure* Hurwitz number $H(\mu^{(1)}, \dots, \mu^{(k)})$ is $\frac{1}{N!}$ times the number of distinct ways that the identity element $\mathbf{I}_N \in S_N$ in the symmetric group in N elements can be expressed as a product

$$\mathbf{I}_N = h_1 \cdots h_k \quad (2.1)$$

of k elements $\{h_i \in S_N\}_{i=1,\dots,k}$, such that for each i , h_i belongs to the conjugacy class $\text{cyc}(\mu^{(i)})$ whose cycle lengths are equal to the parts of $\mu^{(i)}$:

$$h_i \in \text{cyc}(\mu^{(i)}), \quad i = 1, \dots, k. \quad (2.2)$$

An equivalent definition consists of enumeration of branched coverings of the Riemann sphere.

Definition 2.2 (Geometric). For a set of partitions $\{\mu^{(i)}\}_{i=1,\dots,k}$ of weight $|\mu^{(i)}| = N$, the pure Hurwitz number $H(\mu^{(1)}, \dots, \mu^{(k)})$ is defined geometrically [30, 31] as the number of inequivalent N -fold branched coverings $\mathcal{C} \rightarrow \mathbf{P}^1$ of the Riemann sphere with k branch points $(Q^{(1)}, \dots, Q^{(k)})$, whose ramification profiles are given by the partitions $\{\mu^{(1)}, \dots, \mu^{(k)}\}$, normalized by the inverse $1/|\text{aut}(\mathcal{C})|$ of the order of the automorphism group of the covering.

The equivalence of these two definitions follows [34] from the monodromy homomorphism

$$\mathcal{M} : \pi_1(\mathbf{P}^1 / \{Q^{(1)}, \dots, Q^{(k)}\}) \rightarrow S_N \quad (2.3)$$

from the fundamental group of the Riemann sphere punctured at the branch points into S_N , obtained by lifting closed loops from the base to the covering.

To define *weighted* Hurwitz numbers [21, 22, 24, 29], we introduce a weight generating function $G(z)$, either as an infinite product

$$G(z) = \prod_{i=1}^{\infty} (1 + c_i z) \quad (2.4)$$

or an infinite sum

$$G(z) = 1 + \sum_{i=1}^{\infty} g_i z^i, \quad (2.5)$$

either formally, or under suitable convergence conditions imposed upon the parameters. Alternatively, it may be chosen in the dual form

$$\tilde{G}(z) = \prod_{i=1}^{\infty} (1 - c_i z)^{-1}, \quad (2.6)$$

which may also be developed as an infinite sum,

$$\tilde{G}(z) = 1 + \sum_{i=1}^{\infty} \tilde{g}_i z^i. \quad (2.7)$$

The independent parameters determining the weighting may be viewed as either $\{g_i\}_{i \in \mathbf{N}^+}$, $\{\tilde{g}_i\}_{i \in \mathbf{N}^+}$ or $\{c_i\}_{i \in \mathbf{N}^+}$. They are related by the fact that (2.4) and (2.6) are generating functions for elementary and complete symmetric functions, respectively,

$$g_i = e_i(\mathbf{c}), \quad \tilde{g}_i = h_i(\mathbf{c}), \quad (2.8)$$

in the parameters $\mathbf{c} = (c_1, c_2, \dots)$.

In much of the following, we consider combined rational weight generating functions

$$G_{\mathbf{c}, \mathbf{d}}(z) := \frac{\prod_{i=1}^L (1 + c_i z)}{\prod_{j=1}^M (1 - d_j z)}, \quad (2.9)$$

whose Taylor series expansion coefficients are

$$g_i(\mathbf{c}, \mathbf{d}) = \sum_{j=0}^i e_j(\mathbf{c}) h_{i-j}(\mathbf{d}), \quad (2.10)$$

where $\mathbf{c} = (c_1, \dots, c_L)$, $\mathbf{d} = (d_1, \dots, d_M)$.

Definition 2.3 (Weighted Hurwitz numbers). For the case of a weight generating function expressed in the form (2.4), choosing a nonnegative integer d and a fixed partition μ of weight $|\mu| = N$, the weighted (single) Hurwitz number $H_G^d(\mu)$ is defined as the weighted sum over all k -tuples $(\mu^{(1)}, \dots, \mu^{(k)})$

$$H_G^d(\mu) := \sum_{k=1}^d \sum_{\substack{\mu^{(1)}, \dots, \mu^{(k)}, |\mu^{(i)}| = N \\ \sum_{i=1}^k \ell^*(\mu^{(i)}) = d}} \mathcal{W}_G(\mu^{(1)}, \dots, \mu^{(k)}) H(\mu^{(1)}, \dots, \mu^{(k)}, \mu) \quad (2.11)$$

where

$$\ell^*(\mu^{(i)}) := |\mu^{(i)}| - \ell(\mu^{(i)}) \quad (2.12)$$

is the *colength* of the partition $\mu^{(i)}$, and the weight factor is defined to be

$$\begin{aligned} \mathcal{W}_G(\mu^{(1)}, \dots, \mu^{(k)}) &:= \frac{1}{k!} \sum_{\sigma \in S_k} \sum_{1 \leq b_1 < \dots < b_k \leq M} c_{b_{\sigma(1)}}^{\ell^*(\mu^{(1)})} \dots c_{b_{\sigma(k)}}^{\ell^*(\mu^{(k)})} \\ &= \frac{|\text{aut}(\lambda)|}{k!} m_\lambda(\mathbf{c}). \end{aligned} \quad (2.13)$$

Here $m_\lambda(\mathbf{c})$ is the monomial symmetric function [35] of the parameters $\mathbf{c} := (c_1, c_2, \dots)$

$$m_\lambda(\mathbf{c}) = \frac{1}{|\text{aut}(\lambda)|} \sum_{\sigma \in \mathfrak{S}_k} \sum_{1 \leq b_1 < \dots < b_k} c_{b_{\sigma(1)}}^{\lambda_1} \dots c_{b_{\sigma(k)}}^{\lambda_k}, \quad (2.14)$$

indexed by the partition λ of weight $|\lambda| = d$ and length $\ell(\lambda) = k$, whose parts $\{\lambda_i\}$ are equal to the colengths $\{\ell^*(\mu^{(i)})\}$ (expressed in weakly decreasing order)

$$\{\lambda_i\}_{i=1, \dots, k} \sim \{\ell^*(\mu^{(i)})\}_{i=1, \dots, k}, \quad \lambda_1 \geq \dots \geq \lambda_k > 0 \quad (2.15)$$

and

$$|\text{aut}(\lambda)| := \prod_{i \geq 1} m_i(\lambda)! \quad (2.16)$$

where $m_i(\lambda)$ is the number of parts of λ equal to i . A similar definition applies for weight generating functions $\tilde{G}(z)$ of the dual form (2.6), with the monomial symmetric functions (2.14) appearing in (2.11) replaced by the *forgotten* symmetric functions [35]

$$f_\lambda(\mathbf{c}) := \frac{(-1)^{\ell^*(\lambda)}}{|\text{aut}(\lambda)|} \sum_{\sigma \in \mathfrak{S}_k} \sum_{1 \leq b_1 < \dots < b_k} c_{b_{\sigma(1)}}^{\lambda_1} \dots c_{b_{\sigma(k)}}^{\lambda_k}, \quad (2.17)$$

For the case of rational weight generating functions $G_{\mathbf{c}, \mathbf{d}}(z)$, the weighted (single) Hurwitz numbers are

$$\begin{aligned} H_{G_{\mathbf{c}, \mathbf{d}}}^d(\mu) &:= \sum_{\substack{1 \leq k, l \\ k+l \leq d}} \sum_{\substack{\mu^{(1)}, \dots, \mu^{(k)}, \nu^{(1)}, \dots, \nu^{(l)} \\ \sum_{i=1}^k \ell^*(\mu^{(i)}) + \sum_{j=1}^l \ell^*(\nu^{(j)}) = d \\ |\mu^{(i)}| = |\nu^{(j)}| = N}} \mathcal{W}_{G_{\mathbf{c}, \mathbf{d}}}(\mu^{(1)}, \dots, \mu^{(k)}; \nu^{(1)}, \dots, \nu^{(l)}) \\ &\quad \times H(\mu^{(1)}, \dots, \mu^{(k)}, \nu^{(1)}, \dots, \nu^{(l)}, \mu), \end{aligned} \quad (2.18)$$

where the weight factor is

$$\begin{aligned} &\mathcal{W}_{G_{\mathbf{c}, \mathbf{d}}}(\mu^{(1)}, \dots, \mu^{(k)}; \nu^{(1)}, \dots, \nu^{(l)}) \\ &:= \frac{(-1)^{\sum_{j=1}^l \ell^*(\nu^{(j)}) - l}}{k! l!} \sum_{\substack{\sigma \in S_k \\ \sigma' \in S_l}} \sum_{\substack{1 \leq a_1 < \dots < a_k \leq M \\ 1 \leq b_1 < \dots < b_l \leq L}} c_{a_{\sigma(1)}}^{\ell^*(\mu^{(1)})} \dots c_{a_{\sigma(k)}}^{\ell^*(\mu^{(k)})} d_{b_{\sigma'(1)}}^{\ell^*(\nu^{(1)})} \dots d_{b_{\sigma'(l)}}^{\ell^*(\nu^{(l)})}. \end{aligned} \quad (2.19)$$

The sum

$$d := \sum_{i=1}^k \ell^*(\mu^{(i)}), \text{ or } d := \sum_{i=1}^k \ell^*(\mu^{(i)}) + \sum_{j=1}^l \ell^*(\nu^{(j)}) \quad (2.20)$$

of the colengths of the weighted partitions is related to the Euler characteristic χ or genus g of the covering curve by the Riemann-Hurwitz formula

$$\chi = 2 - 2g = N + \ell(\mu) - d. \quad (2.21)$$

2.2 Hypergeometric τ -functions as generating functions for weighted Hurwitz numbers

We recall the definition of KP τ -functions of hypergeometric type [41, 42] that serve as generating functions for weighted Hurwitz numbers [22, 24, 29]. For a weight generating function $G(z)$ and nonzero parameter β , we define two doubly infinite sequences $\{r_i^{(G,\beta)}, \rho_i\}_{i \in \mathbf{Z}}$, labeled by the integers

$$r_i^{(G,\beta)} := \beta G(i\beta), \quad i \in \mathbf{Z}, \quad \rho_0 = 1, \quad (2.22)$$

$$\rho_i := \prod_{k=1}^i r_k^{(G,\beta)}, \quad \rho_{-i} := \prod_{k=0}^{i-1} (r_{-k}^{(G,\beta)})^{-1}, \quad i \in \mathbf{N}^+ \quad (2.23)$$

related by

$$r_i^{(G,\beta)} = \frac{\rho_i}{\rho_{i-1}}, \quad (2.24)$$

where β is viewed as a small parameter for which $G(i\beta)$ does not vanish for any integer $i \in \mathbf{Z}$.

Remark 2.1. *The particular cases of simple, strongly monotonic, weakly monotonic Hurwitz numbers and Belyi curves correspond, respectively, to weight generating functions $G(z) = e^z$, $\frac{1}{1-z}$, $1+z$ and $(1+c_1z)(1+c_2z)$.*

For general rational weight generating functions (2.9), the parameters $\{\rho_i\}_{i \in \mathbf{Z}}$ become

$$\begin{aligned} \rho_i(\mathbf{c}, \mathbf{d}) &:= \beta^i \prod_{k=1}^i \frac{\prod_{l=1}^L (1 + k\beta c_l)}{\prod_{m=1}^M (1 - k\beta d_m)}, \\ \rho_{-i}(\mathbf{c}, \mathbf{d}) &:= \beta^{-i} \prod_{k=1}^{i-1} \frac{\prod_{m=1}^M (1 + k\beta d_m)}{\prod_{l=1}^L (1 - k\beta c_l)}, \quad i \in \mathbf{N}. \end{aligned} \quad (2.25)$$

For each partition λ of N , we define the associated *content product* coefficient

$$r_\lambda^{(G,\beta)} := \prod_{(i,j) \in \lambda} r_{j-i}^{(G,\beta)}, \quad (2.26)$$

where the product is over the locations of all boxes in the Young diagram for λ . The KP τ -function of hypergeometric type associated to these parameters is defined as the Schur function series [22, 29]

$$\tau^{(G,\beta)}(\mathbf{t}) := \sum_{N=0}^{\infty} \sum_{\lambda, |\lambda|=N} (h(\lambda))^{-1} r_\lambda^{(G,\beta)} s_\lambda(\mathbf{t}), \quad (2.27)$$

where $h(\lambda)$ is the product of the hook lengths of the partition λ and $\mathbf{t} = (t_1, t_2, \dots)$ is the infinite sequence of KP flow parameters, which may be equated to the sequence of normalized power sums $(p_1, \frac{1}{2}p_2, \dots)$,

$$t_i := \frac{1}{i} \sum_a x_a^i = \frac{1}{i} p_i \quad (2.28)$$

in a finite or infinite set of auxiliary variables (x_1, x_2, \dots) .

Using the Schur character formula [18, 35]

$$s_\lambda = \sum_{\mu, |\mu|=|\lambda|=N} \frac{\chi_\lambda(\mu)}{z_\mu} p_\mu, \quad (2.29)$$

where $\chi_\lambda(\mu)$ is the irreducible character of the S_N representation determined by λ evaluated on the conjugacy class $\text{cyc}(\mu)$ consisting of elements with cycle lengths equal to the parts of μ ,

$$z_\mu := \prod_{i=1}^{\ell(\mu)} m_i(\mu)! i^{m_i(\mu)} \quad (2.30)$$

is the order of the stabilizer of the elements of this conjugacy class, and

$$p_\mu := \prod_{i=1}^{\ell(\mu)} p_{\mu_i} \quad (2.31)$$

is the power sum symmetric function corresponding to partition μ , we may re-express the Schur function series (2.27) as an expansion in the basis $\{p_\mu\}$.

Theorem 2.1 ([21, 22, 24, 29]). *The τ -function $\tau^{(G,\beta)}(\mathbf{t})$ may equivalently be expressed as*

$$\tau^{(G,\beta)}(\mathbf{t}) = \sum_{d=0}^{\infty} \beta^d \sum_{\mu} H_G^d(\mu) p_\mu(\mathbf{t}), \quad (2.32)$$

and the particular case of rational weight generating functions $G_{\mathbf{c},\mathbf{d}}(z)$, as

$$\tau^{(G_{\mathbf{c},\mathbf{d}},\beta)}(\mathbf{t}) = \sum_{d=0}^{\infty} \beta^d \sum_{\mu} H_{G_{\mathbf{c},\mathbf{d}}}^d(\mu) p_{\mu}(\mathbf{t}). \quad (2.33)$$

It is thus a generating function for the weighted Hurwitz numbers $H_G^d(\mu)$. (See also [40] for τ -functions of hypergeometric type and, in particular, the case of rational weight generating functions.)

2.3 The (dual) Baker function and adapted basis

By Sato's formula [43, 44, 46], the dual Baker-Akhiezer function $\Psi_{(G,\beta)}^*(z, \mathbf{t})$ corresponding to the KP τ -function (2.27) is

$$\Psi_{(G,\beta)}^*(z, \mathbf{t}) = e^{-\sum_{i=1}^{\infty} t_i z^i} \frac{\tau^{(G,\beta)}(\mathbf{t} + [z^{-1}])}{\tau^{(G,\beta)}(\mathbf{t})}, \quad (2.34)$$

where

$$[z^{-1}] := \left(\frac{1}{z}, \frac{1}{2z^2}, \dots, \frac{1}{nz^n}, \dots \right). \quad (2.35)$$

Evaluating at $\mathbf{t} = \mathbf{0}$ and setting

$$x := \frac{1}{z}, \quad (2.36)$$

we define

$$\phi_1(x) := \Psi_{G,\beta}^*(1/z, \mathbf{0}) = \tau^{(G,\beta)}([x]). \quad (2.37)$$

More generally [4–6], we introduce a doubly infinite sequence of functions $\{\phi_k(x)\}_{k \in \mathbf{Z}}$, defined as contour integrals around a small circle centred at the origin (or formal residues)

$$\phi_k(x) = \frac{\beta}{2\pi i x^{k-1}} \oint_{|\zeta|=\epsilon} \rho_k(\zeta) e^{\beta^{-1}x\zeta} \frac{d\zeta}{\zeta^k}, \quad (2.38)$$

where $\rho_k(\zeta)$ is the semi-infinite Laurent series in $\frac{1}{\zeta}$ centred at the origin

$$\rho_k(\zeta) := \sum_{i=-\infty}^{k-1} \rho_{-i-1} \zeta^i, \quad (2.39)$$

with the ρ_i 's given by eqs. (2.22), (2.23). Then $\{\phi_k(1/z)\}_{k \in \mathbf{N}^+}$ forms a basis for the element $w^{(G,\beta)}$ of the infinite Sato-Segal-Wilson Grassmannian corresponding to the τ -function $\tau^{(G,\beta)}(\mathbf{t})$.

The ϕ_k 's may alternatively be expanded as formal (or convergent) Laurent series by evaluating the integrals as a sum of residues at the origin,

$$\phi_k(x) = \beta x^{1-k} \sum_{j=0}^{\infty} \frac{\rho_{j-k}}{j!} \left(\frac{x}{\beta}\right)^j. \quad (2.40)$$

This has a nonvanishing radius of convergence in x provided ρ_j grows no faster than a power

$$|\rho_j| < K^j, \quad j \rightarrow \infty. \quad (2.41)$$

Following [4–6], we introduce the recursion operator

$$R := \beta x G(\beta \mathcal{D}), \quad (2.42)$$

where \mathcal{D} is the Euler operator

$$\mathcal{D} := x \frac{d}{dx}, \quad (2.43)$$

and verify that the ϕ_k 's satisfy the recursion relations

$$R(\phi_k) = \phi_{k-1}, \quad k \in \mathbf{Z}. \quad (2.44)$$

2.4 The quantum and classical spectral curve

It is easily seen from the series expansions (2.40) that the functions $\{\phi_k(x)\}_{k \in \mathbf{Z}}$ satisfy the sequence of equations

$$\beta(\mathcal{D} + k - 1)\phi_k = \phi_{k-1}, \quad k \in \mathbf{Z}, \quad (2.45)$$

which, combined with (2.44), are equivalent to:

$$\mathcal{L}_{\mathbf{c}, \mathbf{d}, \beta} \phi_k := (xG(\beta \mathcal{D}) - \mathcal{D})\phi_k = (k - 1)\phi_k \quad k \in \mathbf{Z}. \quad (2.46)$$

The case $k = 1$

$$\mathcal{L}\phi_1(x) = 0 \quad (2.47)$$

is known in the *topological recursion* approach [4–6, 13–15], as the *quantum spectral curve* equation. Substituting

$$\beta \frac{d}{dx} \rightarrow y, \quad (2.48)$$

in (2.47) gives the classical spectral curve:

$$y = \beta G(xy). \quad (2.49)$$

(The residual factor β is just an artifact, due to the fact that, for simplicity, we have equated β with another generating function parameter γ , which keeps track of the weights of the partitions appearing in the expansion (2.27); it can here simply be set equal to 1.)

When the weight generating function is a polynomial $G = G_{\mathbf{c},\mathbf{0}}$, eq. (2.47) is an ordinary differential equation of degree L with coefficients that are polynomials of degree $\leq L$. More generally, if G is a rational function $G = G_{\mathbf{c},\mathbf{d}}$, as in (2.9), we denote the ϕ_k 's as

$$\phi_k^{(\mathbf{c},\mathbf{d},\beta)}(x) = \beta x^{1-k} \sum_{j=0}^{\infty} \frac{\rho_{j-k}(\mathbf{c}, \mathbf{d})}{j!} \left(\frac{x}{\beta}\right)^j. \quad (2.50)$$

In particular, for $k = 1$, this is the hypergeometric series

$$\begin{aligned} \phi_1^{(\mathbf{c},\mathbf{d},\beta)}(x) &= \sum_{j=0}^{\infty} \frac{\prod_{l=1}^L \left(\frac{1}{\beta c_l}\right)_j}{\prod_{m=1}^M \left(-\frac{1}{\beta d_m}\right)_j} \frac{(\kappa_{\mathbf{c},\mathbf{d}}x)^j}{j!} \\ &= {}_L F_M \left(\begin{matrix} \frac{1}{\beta c_1}, \dots, \frac{1}{\beta c_L} \\ -\frac{1}{\beta d_1}, \dots, -\frac{1}{\beta d_M} \end{matrix} \middle| \kappa_{\mathbf{c},\mathbf{d}}x \right), \end{aligned} \quad (2.51)$$

where

$$\kappa_{\mathbf{c},\mathbf{d}} := (-1)^M \frac{\prod_{l=1}^L \beta c_l}{\prod_{m=1}^M \beta d_m}. \quad (2.52)$$

When $L \leq M$, this converges for all x ; when $L = M + 1$ it converges within a finite radius of $x = 0$. But if $L > M + 1$, it is a divergent formal series. In general, for any $k \in \mathbf{Z}$, ϕ_k is proportional to x^{1-k} times the hypergeometric series with parameters shifted by $k - 1$,

$$\begin{aligned} \phi_k^{(\mathbf{c},\mathbf{d},\beta)}(x) &= \frac{\beta \rho_{-k}(\mathbf{c}, \mathbf{d})}{(\kappa_{\mathbf{c},\mathbf{d}}x)^{k-1}} \sum_{j=0}^{\infty} \frac{\prod_{l=1}^L \left(1 - k + \frac{1}{\beta c_l}\right)_j}{\prod_{m=1}^M \left(1 - k - \frac{1}{\beta d_m}\right)_j} \frac{(\kappa_{\mathbf{c},\mathbf{d}}x)^j}{j!} \\ &= \frac{\beta \rho_{-k}(\mathbf{c}, \mathbf{d})}{(\kappa_{\mathbf{c},\mathbf{d}}x)^{k-1}} {}_L F_M \left(\begin{matrix} 1 - k + \frac{1}{\beta c_1}, \dots, 1 - k + \frac{1}{\beta c_L} \\ 1 - k - \frac{1}{\beta d_1}, \dots, 1 - k - \frac{1}{\beta d_M} \end{matrix} \middle| \kappa_{\mathbf{c},\mathbf{d}}x \right), \end{aligned} \quad (2.53)$$

and hence is also convergent, or not, depending on whether $L \leq M$, $L = M + 1$ or $L > M + 1$.

Defining an auxiliary function $\Phi^{(\mathbf{c},\mathbf{d},\beta)}(x)$ such that

$$\phi_1^{(\mathbf{c},\mathbf{d},\beta)} = \prod_{j=1}^M (1 - \beta d_j \mathcal{D}) \Phi^{(\mathbf{c},\mathbf{d},\beta)}, \quad (2.54)$$

the quantum spectral curve equation may be expressed as

$$x \prod_{i=1}^L (1 + \beta c_i \mathcal{D}) \Phi^{(\mathbf{c}, \mathbf{d}, \beta)} - \mathcal{D} \prod_{j=1}^M (1 - \beta d_j \mathcal{D}) \Phi^{(\mathbf{c}, \mathbf{d}, \beta)} = 0. \quad (2.55)$$

Remark 2.2. *Note that, although in the space of differentiable functions in a punctured neighbourhood of $x = 0$, with possible branch point at $x = 0$, the operator $\prod_{j=1}^L (1 - \beta d_j \mathcal{D})$ has a kernel of dimension L , if we restrict ourselves to formal (or convergent) Laurent series in x and assume that the βd_j 's are not integers, the kernel vanishes, making $\Phi^{(\mathbf{c}, \mathbf{d}, \beta)}(x)$ unique.*

3 Solutions of the quantum spectral curve equation for rational $G(z) = G_{\mathbf{c}, \mathbf{d}}(z)$

Equation (2.55) is of the type satisfied by Meijer G -functions [9, 37]

$$\left[(-1)^{p-m-n} \zeta \prod_{j=1}^p \left(\zeta \frac{d}{d\zeta} + 1 - a_j \right) - \prod_{j=1}^q \left(\zeta \frac{d}{d\zeta} - b_j \right) \right] G^{m,n}(\zeta) = 0, \quad (3.1)$$

with the identifications

$$\begin{aligned} p &= L, \quad q = M + 1, \quad \zeta = (-1)^{L-m-n} \kappa_{\mathbf{c}, \mathbf{d}} x, \\ a_j &= 1 - \frac{1}{\beta c_j}, \quad b_1 = 0, \quad b_{j+1} = \frac{1}{\beta d_j} \quad (j \geq 1). \end{aligned} \quad (3.2)$$

These all have Mellin-Barnes type integral representations of the form

$$G_{p,q}^{m,n} \left(\begin{matrix} a_1, \dots, a_p \\ b_1, \dots, b_q \end{matrix} \middle| \zeta \right) = \frac{1}{2\pi i} \int_{\mathcal{C}_{\mathbf{ab}}} \frac{\prod_{j=1}^m \Gamma(b_j - s) \prod_{k=1}^n \Gamma(1 - a_k + s) \zeta^s}{\prod_{j=m+1}^q \Gamma(1 - b_j + s) \prod_{k=n+1}^p \Gamma(a_k - s)} ds, \quad (3.3)$$

where $0 \leq m \leq q$, $0 \leq n \leq p$ and the contour $\mathcal{C}_{\mathbf{ab}}$ lies between the poles at $\{s = b_j + i\}_{j=1, \dots, m, i \in \mathbf{N}^+}$ and $\{s = a_k - i - 1\}_{k=1, \dots, n, i \in \mathbf{N}^+}$.

By retracting the contour to the right-hand s -plane and evaluating the residues at the poles of the integrand, we obtain a representation of the Meijer-G function as a sum of powers of ζ within the set $\bigcup (b_j + \mathbf{N})$. If $p \geq q$, this only yields a formal series, which may be interpreted as an asymptotic series in suitable sectors of the complex ζ -plane. However, if $p < q$, it gives a convergent expansion.

For $k \in \mathbf{Z}$, we now define the sequence of functions

$$\begin{aligned} \tilde{\phi}_k^{(\mathbf{c}, \mathbf{d}, \beta)}(x) &:= C_k^{(\mathbf{c}, \mathbf{d}, \beta)} G_{L, M+1}^{1, L} \left(\begin{array}{c} 1 - \frac{1}{\beta c_1}, \dots, 1 - \frac{1}{\beta c_L} \\ 1 - k, 1 + \frac{1}{\beta d_1}, \dots, 1 + \frac{1}{\beta d_M} \end{array} \middle| -\kappa_{\mathbf{c}, \mathbf{d}} x \right) \\ &= \frac{C_k^{(\mathbf{c}, \mathbf{d}, \beta)}}{2\pi i} \int_{\mathcal{C}_k} \frac{\Gamma(1 - k - s) \prod_{\ell=1}^L \Gamma\left(s + \frac{1}{\beta c_\ell}\right) (-\kappa_{\mathbf{c}, \mathbf{d}} x)^s}{\prod_{m=1}^M \Gamma\left(s - \frac{1}{\beta d_m}\right)} ds. \end{aligned} \quad (3.4)$$

where

$$C_k^{(\mathbf{c}, \mathbf{d}, \beta)} := \frac{\prod_{j=1}^M \Gamma\left(-\frac{1}{\beta d_j}\right)}{(-\beta)^{k-1} \prod_{\ell=1}^L \Gamma\left(\frac{1}{\beta c_\ell}\right)}, \quad (3.5)$$

with the contour \mathcal{C}_k chosen so that all the poles of the integrand at the integers $1 - k, 2 - k, \dots$ are to its right and the poles at $\{-i - \frac{1}{\beta c_j}\}_{j=1, \dots, L, i \in \mathbf{N}^+}$ are to its left.

If $L \leq M$ this gives a convergent value for all $x \neq 0$ when the contour \mathcal{C}_k starts at $+\infty$ below the real axis, proceeds in a clockwise sense, passes around all the poles on the real axis and ends at $+\infty$. If $L = M + 1$ it also converges along this contour, within the punctured disc

$$0 < |\kappa_{\mathbf{c}, \mathbf{d}} x| < 1. \quad (3.6)$$

If $L \geq M + 1$ the contour \mathcal{C}_k may be chosen to start and end on the imaginary axis at $-i\infty$ and $+i\infty$. The integral then converges provided that

$$|\arg(-\kappa_{\mathbf{c}, \mathbf{d}} x)| < \left(\frac{L - M + 1}{2}\right) \pi. \quad (3.7)$$

We then have the following theorem.

Theorem 3.1. *The functions $\tilde{\phi}_k^{(\mathbf{c}, \mathbf{d}, \beta)}(x)$ satisfy the sequence of equations (2.45), the recursion relations (2.44) and the eigenfunction equations (2.46).*

If $L \leq M$, the Laurent series (2.50) for $\phi_k^{(\mathbf{c}, \mathbf{d}, \beta)}(x)$ is absolutely convergent for all $x \in \mathbf{C}$, $x \neq 0$, and

$$\tilde{\phi}_k^{(\mathbf{c}, \mathbf{d}, \beta)}(x) = \phi_k^{(\mathbf{c}, \mathbf{d}, \beta)}(x). \quad (3.8)$$

If $L = M + 1$, it converges to $\tilde{\phi}_k^{(\mathbf{c}, \mathbf{d}, \beta)}(x)$ within the punctured disc

$$0 < |\kappa_{\mathbf{c}, \mathbf{d}} x| < 1. \quad (3.9)$$

If $L > M + 1$, eq. (2.50) is the asymptotic series for $\tilde{\phi}_k^{(\mathbf{c}, \mathbf{d}, \beta)}(x)$ within the angular sector

$$0 < \arg(-\kappa_{\mathbf{c}, \mathbf{d}} x) < (L - M + 1) \frac{\pi}{2}. \quad (3.10)$$

Proof. **Case** $L \leq M + 1$. We first prove the result for $k = 1$ by evaluation of the residues of the integrand at the poles $s = 0, 1, \dots$. These all come from the factor $\Gamma(-s)$ in the numerator of the integrand, whose residues are

$$\operatorname{res}_{s=j} \Gamma(-s) = \frac{(-1)^j}{j!}. \quad (3.11)$$

Substituting this, and evaluating all other factors in the integrand at the poles gives, by the Cauchy residue theorem,

$$\begin{aligned} \tilde{\phi}_1^{(\mathbf{c}, \mathbf{d}, \beta)}(x) &= \sum_{j=0}^{\infty} \prod_{l=1}^L \frac{\Gamma(j + \frac{1}{\beta c_l})}{\Gamma(\frac{1}{\beta c_l})} \prod_{m=1}^M \frac{\Gamma(-\frac{1}{\beta d_m})}{\Gamma(j - \frac{1}{\beta d_m})} \frac{(\kappa_{\mathbf{c}, \mathbf{d}} x)^j}{j!} \\ &= \sum_{j=0}^{\infty} \frac{\prod_{l=1}^L \left(\frac{1}{\beta c_l}\right)_j}{\prod_{m=1}^M \left(-\frac{1}{\beta d_m}\right)_j} \frac{(\kappa_{\mathbf{c}, \mathbf{d}} x)^j}{j!} = {}_L F_M \left(\begin{matrix} \frac{1}{\beta c_1}, \dots, \frac{1}{\beta c_L} \\ -\frac{1}{\beta d_1}, \dots, -\frac{1}{\beta d_M} \end{matrix} \middle| \kappa_{\mathbf{c}, \mathbf{d}} x \right). \end{aligned} \quad (3.12)$$

The proof for general $k \in \mathbf{Z}$ is the same, *mutatis mutandis*, with the contour \mathcal{C}_1 replaced by \mathcal{C}_k , the normalization constant $C_1^{(\mathbf{c}, \mathbf{d}, \beta)}$ by $C_k^{(\mathbf{c}, \mathbf{d}, \beta)}$ and the term $\Gamma(-s)$ in the numerator of the integrand by $\Gamma(1 - k - s)$. (Also see Remark 3.1 below.)

Case $L > M + 1$. In this case $p = L > q = M + 1$, the ODE (3.1) has an irregular singularity at $\zeta = 0$ and the solutions manifest Stokes' phenomenon. The function (3.4) is one such solution. (See Remark 3.2 below.) We now show that the series (2.50) is indeed its asymptotic series as $x \rightarrow 0$ in the sector (3.7).

First note again that the integrand in (3.4) has poles at $s = -k + 1, -k + 2, \dots$ with residues that give precisely the corresponding coefficient of x^s in (2.50). What must be shown is that the remainder of $\tilde{\phi}_k^{(\mathbf{c}, \mathbf{d}, \beta)}(x)$ minus a truncation of the series (2.50) to order $\mathcal{O}(x^N)$ is of order $o(x^N)$ as $x \rightarrow 0$ in the indicated sector, since this is the definition of Poincaré asymptotics.

To show this, consider the integral from (3.4):

$$F(\zeta) := \int_{\mathcal{C}_k} \frac{\Gamma(1 - k - s) \prod_{\ell=1}^L \Gamma\left(s + \frac{1}{\beta c_\ell}\right) \zeta^s}{\prod_{m=1}^M \Gamma\left(s - \frac{1}{\beta d_m}\right)} ds, \quad (3.13)$$

where we have set $\zeta = -\kappa_{\mathbf{c}, \mathbf{d}} x$ for brevity. We use the Stirling approximation in the following form

$$\ln \Gamma(s) = \left(s - \frac{1}{2}\right) \ln \left(s - \frac{1}{2}\right) - \left(s - \frac{1}{2}\right) + \mathcal{O}(1) \quad (3.14)$$

$$\ln \Gamma(1 - s) = -\left(s - \frac{1}{2}\right) \ln \left(s - \frac{1}{2}\right) + \left(s - \frac{1}{2}\right) - \ln \sin(\pi s) + \mathcal{O}(1). \quad (3.15)$$

Denoting the integrand of (3.13) $A_k(s)\zeta^s$, with $\zeta = -\kappa_{\mathbf{c},\mathbf{d}}x$, we can estimate its modulus as follows:

$$\ln(A_k(s)\zeta^s) = (L - M - 1)(s \ln s - s) - \ln \sin(\pi s) + s \ln \zeta + o(|s|). \quad (3.16)$$

We choose the contour of integration \mathcal{C}_k in (3.13) to tend to infinity in both positive and negative vertical directions and approach $\Re s = N + \frac{1}{2}$, $N \in \mathbb{R}$ in both. Concretely, we may take a contour that coincides with the vertical line $\Re s = N + \frac{1}{2}$ for $|\Im s|$ sufficiently large, but again passes to the left of the poles at $1 - k, 2 - k, \dots$ and to the right of those at $\{-j - \frac{1}{\beta_{c_l}}\}_{l=1, \dots, L, j \in \mathbb{N}}$. Along this contour the real part of (3.16) has the following asymptotic behaviour

$$\begin{aligned} \ln |A(s)\zeta^s| &= -(L - M - 1)|\Im s| \frac{\pi}{2} - \pi |\Im s| + |\Im s| \arg \zeta + o(|s|) \\ &= \frac{\pi}{2} (M - 1 - L) |\Im s| + |\Im s| \arg \zeta + o(|s|). \end{aligned} \quad (3.17)$$

Thus, since the modulus of the integrand tends to zero faster than any inverse power of $|s|$, the integrand is convergent as long as $|\arg \zeta| < \frac{\pi(L-M+1)}{2}$.

Using the Cauchy residue theorem we can retract (see Fig. 1) the contour \mathcal{C}_k to $\Re s = N + \frac{1}{2}$. In doing so we pick up finitely many residues at $s = -k + 1, -k + 2, \dots, N - 1, N$, which correspond to the truncation of the series (2.50) up to the power x^N included. It only remains to estimate the remaining integration $R(\zeta)$ (remainder term):

$$R(\zeta) = \int_{i\mathbb{R}+N+\frac{1}{2}} A_k(s)\zeta^s ds \quad (3.18)$$

The estimate (3.14) shows that the integral is well defined as long as $|\arg \zeta| < \frac{\pi(L-M+1)}{2}$. Furthermore the whole integral is easily estimated to be of the order $|\zeta|^{N+\frac{1}{2}}$ because

$$\left| \int_{i\mathbb{R}+N+\frac{1}{2}} A_k(s)\zeta^s ds \right| \leq |z|^{N+\frac{1}{2}} \int_{i\mathbb{R}+N+\frac{1}{2}} |A(s)| e^{\arg \zeta \Im s} ds \quad (3.19)$$

and the last integral can be uniformly estimated in any sector $|\arg \zeta| < \frac{\pi(L-M+1)}{2} - \epsilon$. This shows that ϕ_k has the formal series (2.40) as asymptotic series in the sense of Poincaré.




Figure 1: The contours of integration for the function (3.13) in the case $L > M + 1$.

Next, regardless of the relative value of L and M , applying the operator $\mathcal{D} + k - 1$ to $\tilde{\phi}_k^{(\mathbf{c}, \mathbf{d}, \beta)}(x)$ and evaluating under the integral sign it follows immediately that the sequence of equations (2.45) is satisfied.

Finally, the following computation shows that the sequence of eigenfunction equations (2.46) is satisfied by the $\tilde{\phi}_k^{(\mathbf{c}, \mathbf{d}, \beta)}(x)$'s, and hence, so are the recursion relations (2.44).

$$\begin{aligned}
& (\mathcal{D} + k - 1) \int_{\mathcal{C}_k} \frac{\Gamma(1 - k - s) \prod_{\ell=1}^L \Gamma\left(s + \frac{1}{\beta_{c_\ell}}\right) (-\kappa_{\mathbf{c}, \mathbf{d}x})^s}{\prod_{m=1}^M \Gamma\left(s - \frac{1}{\beta_{d_m}}\right)} ds. \\
&= \int_{\mathcal{C}_k} \frac{(s + k - 1) \Gamma(1 - k - s) \prod_{\ell=1}^L \Gamma\left(s + \frac{1}{\beta_{c_\ell}}\right) (-\kappa_{\mathbf{c}, \mathbf{d}x})^s}{\prod_{m=1}^M \Gamma\left(s - \frac{1}{\beta_{d_m}}\right)} ds \\
&= - \int_{\mathcal{C}_k} \frac{\Gamma(2 - k - s) \prod_{\ell=1}^L \Gamma\left(s + \frac{1}{\beta_{c_\ell}}\right) (-\kappa_{\mathbf{c}, \mathbf{d}x})^s}{\prod_{m=1}^M \Gamma\left(s - \frac{1}{\beta_{d_m}}\right)} ds \\
&= - \int_{\mathcal{C}_k} \frac{\Gamma(1 - k - s) \prod_{\ell=1}^L \Gamma\left(1 + s + \frac{1}{\beta_{c_\ell}}\right) (-\kappa_{\mathbf{c}, \mathbf{d}x})^{s+1}}{\prod_{m=1}^M \Gamma\left(1 + s - \frac{1}{\beta_{d_m}}\right)} ds \\
&= \kappa_{\mathbf{c}, \mathbf{d}x} \int_{\mathcal{C}_k} \frac{\Gamma(1 - k - s) \prod_{\ell=1}^L \Gamma\left(1 + s + \frac{1}{\beta_{c_\ell}}\right) (-\kappa_{\mathbf{c}, \mathbf{d}x})^s}{\prod_{m=1}^M \Gamma\left(1 + s - \frac{1}{\beta_{d_m}}\right)} ds
\end{aligned}$$

$$\begin{aligned}
&= \kappa_{\mathbf{c}, \mathbf{d}x} \int_{\mathcal{C}_k} \frac{\Gamma(1-k-s) \prod_{\ell=1}^L \left(s + \frac{1}{\beta_{c_\ell}}\right) \Gamma\left(s + \frac{1}{\beta_{c_\ell}}\right) (-\kappa_{\mathbf{c}, \mathbf{d}x})^s}{\prod_{m=1}^M \left(s - \frac{1}{\beta_{d_m}}\right) \Gamma\left(s - \frac{1}{\beta_{d_m}}\right)} ds \\
&= x \int_{\mathcal{C}_k} \frac{\Gamma(1-k-s) \prod_{\ell=1}^L (1 + \beta s c_\ell) \Gamma\left(s + \frac{1}{\beta_{c_\ell}}\right) (-\kappa_{\mathbf{c}, \mathbf{d}x})^s}{\prod_{m=1}^M (1 - \beta s d_m) \Gamma\left(s - \frac{1}{\beta_{d_m}}\right)} ds \\
&= x G_{\mathbf{c}, \mathbf{d}}(\beta \mathcal{D}) \int_{\mathcal{C}_k} \frac{\Gamma(1-k-s) \prod_{\ell=1}^L \Gamma\left(s + \frac{1}{\beta_{c_\ell}}\right) (-\kappa_{\mathbf{c}, \mathbf{d}x})^s}{\prod_{m=1}^M \Gamma\left(s - \frac{1}{\beta_{d_m}}\right)} ds, \tag{3.20}
\end{aligned}$$

where in the fourth line, the contour \mathcal{C}_k is understood to be chosen such that the shift $s \rightarrow s + 1$ does not traverse any poles. \square

Remark 3.1. For $L \leq M + 1$, we could equally well have replaced

$$G_{L, M+1}^{1, L} \left(\begin{matrix} 1 - \frac{1}{\beta_{c_1}}, \dots, 1 - \frac{1}{\beta_{c_L}} \\ 1 - k, 1 + \frac{1}{\beta_{d_1}}, \dots, 1 + \frac{1}{\beta_{d_M}} \end{matrix} \middle| -\kappa_{\mathbf{c}, \mathbf{d}x} \right) \tag{3.21}$$

in (3.4) by

$$G_{L, M+1}^{1, n} \left(\begin{matrix} 1 - \frac{1}{\beta_{c_1}}, \dots, 1 - \frac{1}{\beta_{c_L}} \\ 1 - k, 1 + \frac{1}{\beta_{d_1}}, \dots, 1 + \frac{1}{\beta_{d_M}} \end{matrix} \middle| (-1)^{L+n+1} \kappa_{\mathbf{c}, \mathbf{d}x} \right) \tag{3.22}$$

for any $0 \leq n \leq L$, with a suitably modified normalization constant, since these are all constant multiples of each other, as can be seen, e.g., from Slater's identity [37]:

$$\begin{aligned}
&G_{L, M+1}^{1, n} \left(\begin{matrix} 1 - \frac{1}{\beta_{c_1}}, \dots, 1 - \frac{1}{\beta_{c_L}} \\ 1 - k, 1 + \frac{1}{\beta_{d_1}}, \dots, 1 + \frac{1}{\beta_{d_M}} \end{matrix} \middle| \zeta \right) \\
&= \frac{\zeta^{1-k} \prod_{l=1}^n \Gamma(1-k + \frac{1}{\beta_{c_l}}) {}_L F_M \left(\begin{matrix} 1-k + \frac{1}{\beta_{c_1}}, \dots, 1-k + \frac{1}{\beta_{c_L}} \\ 1-k - \frac{1}{\beta_{d_1}}, \dots, 1-k - \frac{1}{\beta_{d_M}} \end{matrix} \middle| (-1)^{L+n+1} \zeta \right)}{\prod_{l=n+1}^L \Gamma(k - \frac{1}{\beta_{c_l}}) \prod_{m=1}^M \Gamma(1-k - \frac{1}{\beta_{d_m}})}, \tag{3.23}
\end{aligned}$$

The equality (3.8) also follows in this case from (3.23) with $n = L$.

For $n = L$, $k = 1$, this reproduces (3.12). For general k , it reproduces (3.8), with $\phi_k^{(\mathbf{c}, \mathbf{d}, \beta)}(x)$ defined by the series (2.50) which, up to normalization, is just x^{1-k} times the hypergeometric series for

$${}_L F_M \left(\begin{matrix} 1 - k + \frac{1}{\beta_{c_1}}, \dots, 1 - k + \frac{1}{\beta_{c_L}} \\ 1 - k - \frac{1}{\beta_{d_1}}, \dots, 1 - k - \frac{1}{\beta_{d_M}} \end{matrix} \middle| (-1)^{L+n} \kappa_{\mathbf{c}, \mathbf{d}x} \right). \tag{3.24}$$

Remark 3.2. If $L > M + 1$, we could also have replaced

$$G_{L,M+1}^{1,L} \left(\begin{matrix} 1 - \frac{1}{\beta_{c_1}}, \dots, 1 - \frac{1}{\beta_{c_L}} \\ 1 - k, 1 + \frac{1}{\beta_{d_1}}, \dots, 1 + \frac{1}{\beta_{d_M}} \end{matrix} \middle| -\kappa_{\mathbf{c}, \mathbf{d}x} \right) \quad (3.25)$$

in (3.4) by any suitably normalized linear combination of the linearly independent functions

$$G_{L,M+1}^{1,L-1} \left(\begin{matrix} 1 - \frac{1}{\beta_{c_1}}, \dots, \widehat{1 - \frac{1}{\beta_{c_i}}}, \dots, 1 - \frac{1}{\beta_{c_L}}, 1 - \frac{1}{\beta_{c_i}} \\ 1 - k, 1 + \frac{1}{\beta_{d_1}}, \dots, 1 + \frac{1}{\beta_{d_M}} \end{matrix} \middle| \kappa_{\mathbf{c}, \mathbf{d}x} \right), \quad i = 1, \dots, L, \quad (3.26)$$

which form a basis of solutions of the same differential equation and, within a normalization, have the same asymptotic series (2.50) within the angular sector

$$0 < \arg(\kappa_{\mathbf{c}, \mathbf{d}x}) < (L - M - 1) \frac{\pi}{2} \quad (3.27)$$

To see the relation between eq. (2.46) and the equations satisfied by the Meijer G -functions, we set

$$\zeta := -\kappa_{\mathbf{c}, \mathbf{d}x} \quad (3.28)$$

and rewrite (2.46) as

$$\zeta \frac{\prod_{l=1}^L (\mathcal{D} + \frac{1}{\beta_{c_l}})}{\prod_{m=1}^M (\mathcal{D} - \frac{1}{\beta_{d_m}})} \phi_k^{(\mathbf{c}, \mathbf{d}, \beta)} + (\mathcal{D} + k - 1) \phi_k^{(\mathbf{c}, \mathbf{d}, \beta)} = 0, \quad (3.29)$$

where

$$\mathcal{D} = \zeta \frac{d}{d\zeta} = x \frac{d}{dx}. \quad (3.30)$$

Multiplying (3.29) by ζ^{-1} and applying the operator $\prod_{m=1}^M (\mathcal{D} - \frac{1}{\beta_{d_m}})$ on the left gives

$$\prod_{l=1}^L (\mathcal{D} + \frac{1}{\beta_{c_l}}) \phi_k^{(\mathbf{c}, \mathbf{d}, \beta)} + \prod_{m=1}^M (\mathcal{D} - \frac{1}{\beta_{d_m}}) \zeta^{-1} (\mathcal{D} + k - 1) \phi_k^{(\mathbf{c}, \mathbf{d}, \beta)} = 0. \quad (3.31)$$

Now using the ladder relation

$$\left(\mathcal{D} - \frac{1}{\beta_{d_j}} \right) \zeta^{-1} = \zeta^{-1} \left(\mathcal{D} - \frac{1}{\beta_{d_j}} + 1 \right) \quad (3.32)$$

repeatedly to move ζ^{-1} in the second term in (3.31) to the left of the differential operator term and, finally, multiplying by ζ gives

$$\zeta \prod_{l=1}^L (\mathcal{D} + \frac{1}{\beta_{c_l}}) \phi_k^{(\mathbf{c}, \mathbf{d}, \beta)} + (\mathcal{D} + k - 1) \prod_{m=1}^M (\mathcal{D} - 1 - \frac{1}{\beta_{d_m}}) \phi_k^{(\mathbf{c}, \mathbf{d}, \beta)} = 0, \quad (3.33)$$

which is eq. (3.1) for

$$\begin{aligned}
p &= L, & q &= M + 1, & m &= 1, & n &= L \\
a_l &= 1 - \frac{1}{\beta c_l}, & l &= 1, \dots, L, \\
b_1 &= 1 - k, & b_{m+1} &= 1 + \frac{1}{\beta d_m}, & m &= 1, \dots, M.
\end{aligned} \tag{3.34}$$

In particular, for the case $L = 2$, $M = 0$, with three branch points, two of them, with branching profiles $(\mu^{(1)}, \mu^{(2)})$, weighted as in eq. (2.19), and a third unweighted one with profile μ , we have

$$\tilde{\phi}_k^{((c_1, c_2), \cdot, \beta)}(x) = C_k^{((c_1, c_2), \cdot, \beta)} G_{2,1}^{1,2} \left(1 - \frac{1}{\beta c_1} \quad 1 - \frac{1}{\beta c_2} \quad \middle| \quad \kappa_{\mathbf{c}, \mathbf{d}} x \right). \tag{3.35}$$

This corresponds to the enumeration of *Belyi curves* or, equivalently, *dessins d'enfants* [1, 47].

4 The τ -function $\tau^{(G, \beta)}(\mathbf{t})$ evaluated at power sums

As detailed in [4–6], $\tau^{(G, \beta)}(\mathbf{t})$ is the KP τ -function corresponding to the Grassmannian element spanned by the basis elements $\{\phi_k(1/z)\}_{k \in \mathbf{N}^+}$

$$w^{(G, \beta)} = \text{span}\{\phi_k(1/z)\}, \quad k \geq 1. \tag{4.1}$$

It follows [27] that, if $\tau^{(G, \beta)}(\mathbf{t})$ is evaluated at the trace invariants

$$\mathbf{t} = [X], \quad t_i = \frac{1}{i} \text{tr} X^i \tag{4.2}$$

of a diagonal $n \times n$ matrix

$$X := \text{diag}(x_1, \dots, x_n), \tag{4.3}$$

it is expressible as the ratio of $n \times n$ determinants

$$\tau^{(G, \beta)}([X]) = \frac{\prod_{i=1}^n x_i^{n-1} \det(\phi_i(x_j))_{1 \leq i, j \leq n}}{\prod_{i=1}^n \rho_{-i} \Delta(x)}, \tag{4.4}$$

where

$$\Delta(x) = \prod_{1 \leq i < j \leq n} (x_i - x_j) = \det(x_i^{n-j})_{1 \leq i, j \leq n} \tag{4.5}$$

is the Vandermonde determinant.

Remark 4.1. The evaluation $\tau^{(G,\beta)}([X])$ is precisely what was originally defined in [23] as a “hypergeometric function of matrix argument”. See also [40] for matrix integral representations of τ -functions.

It also follows from (2.45) that each $\{\phi_k^{(\mathbf{c},\mathbf{d},\beta)}(x)\}_{1 \leq k \leq n}$ may be expressed as a finite lower triangular linear combination of the powers of the Euler operator \mathcal{D} applied to $\phi_n^{(\mathbf{c},\mathbf{d},\beta)}(x)$

$$\phi_k(x) = \beta^{n-k} \mathcal{D}^{n-k} \phi_n(x) + \sum_{j=0}^{n-k-1} \Gamma_{kj} \mathcal{D}^j \phi_n(x), \quad k = 1, \dots, n. \quad (4.6)$$

where the Γ_{kj} 's are polynomials in β of lower degree with integer coefficients. Therefore

$$\tau^{(G,\beta)}([X]) = \gamma_n \left(\prod_{i=1}^n x_i^{n-1} \right) \frac{\det(\mathcal{D}^{i-1} \phi_n(x_j))_{1 \leq i,j \leq n}}{\Delta(x)}, \quad (4.7)$$

where

$$\gamma_n := \frac{\beta^{\frac{1}{2}n(n-1)}}{\prod_{i=1}^n \rho_{-i}}. \quad (4.8)$$

For rational weight generating functions $G = G_{\mathbf{c},\mathbf{d}}$, and any positive integer $n \in \mathbf{N}^+$, let

$$f_{(\mathbf{c},\mathbf{d},\beta,n)}(y) := \tilde{\phi}_n^{(\mathbf{c},\mathbf{d},\beta)}(e^y) = \int_{\mathcal{C}_n} A_n^{(\mathbf{c},\mathbf{d},\beta)}(s) e^{ys} ds, \quad (4.9)$$

$$A_n^{(\mathbf{c},\mathbf{d},\beta)}(s) := \frac{C_n^{(\mathbf{c},\mathbf{d},\beta)} \Gamma(1-n-s) \prod_{l=1}^L \Gamma\left(s + \frac{1}{\beta c_l}\right) (-\kappa_{\mathbf{c},\mathbf{d}})^s}{2\pi i \prod_{m=1}^M \Gamma\left(s - \frac{1}{\beta d_m}\right)}. \quad (4.10)$$

Defining the diagonal matrix

$$Y = \text{diag}(y_1, \dots, y_n) \quad (4.11)$$

by

$$X = e^Y, \quad Y = \ln(X), \quad x_i = e^{y_i}, \quad i = 1, \dots, n, \quad (4.12)$$

eq. (4.7) can be expressed as a ratio of Wronskian determinants

$$\tau^{(G_{\mathbf{c},\mathbf{d},\beta})}([X]) = \gamma_n \left(\prod_{i=1}^n x_i^{n-1} \right) \frac{\det\left(f_{(\mathbf{c},\mathbf{d},\beta,n)}^{(i-1)}(y_j)\right)_{1 \leq i,j \leq n}}{\Delta(e^y)}, \quad (4.13)$$

where

$$\Delta(e^y) := \prod_{1 \leq i < j \leq n} (e^{y_i} - e^{y_j}). \quad (4.14)$$

5 Matrix integral representation of $\tau^{(G_{\mathbf{c},\mathbf{d},\beta})}([X])$

Consider the generalized Brézin-Hikami [11] matrix integral

$$\mathbf{Z}_{d\mu_{(\mathbf{c},\mathbf{d},\beta,n)}}(X) = \int_{M \in \text{Nor}_{\mathcal{C}_n}^{n \times n}} d\mu_{(\mathbf{c},\mathbf{d},\beta,n)}(M) e^{\text{tr}YM}, \quad (5.1)$$

where

$$d\mu_{(\mathbf{c},\mathbf{d},\beta,n)}(M) := (\Delta(\zeta))^2 \det(A_n^{(\mathbf{c},\mathbf{d},\beta)}(M)) d\mu_0(U) \prod_{j=1}^n d\zeta_j \quad (5.2)$$

is a conjugation invariant measure on the space $\text{Nor}_{\mathcal{C}_n}^{n \times n}$ of $n \times n$ normal matrices

$$M = UZU^\dagger \in \text{Nor}_{\mathcal{C}_n}^{n \times n}, \quad U \in U(n), \quad Z = \text{diag}(\zeta_1, \dots, \zeta_n) \quad (5.3)$$

with eigenvalues $\zeta_i \in \mathbf{C}$ supported on the contour \mathcal{C}_n and $d\mu_0(U)$ is the Haar measure on $U(n)$.

Theorem 5.1. *The τ -function $\tau^{(G_{\mathbf{c},\mathbf{d},\beta})}(\mathbf{t})$, evaluated at the trace invariants $[X]$ of the externally coupled matrix X is given by*

$$\tau^{(G_{\mathbf{c},\mathbf{d},\beta})}([X]) = \frac{\beta^{\frac{1}{2}n(n-1)} (\prod_{i=1}^n x_i^{n-1}) \Delta(\ln(x))}{(\prod_{i=1}^n i!) \Delta(x)} \mathbf{Z}_{d\mu_{(\mathbf{c},\mathbf{d},\beta,n)}}(X), \quad (5.4)$$

where $\Delta(\ln(x)) = \Delta(y)$ is the Vandermonde determinant in the variables (y_1, \dots, y_n) .

Proof. Using the Harish-Chandra-Itzykson-Zuber identity [28, 32]

$$\int_{U \in U(n)} d\mu(U) e^{\text{tr}(YUZU^\dagger)} = \frac{(\prod_{i=1}^{n-1} i!) \det(e^{y_i \zeta_j})}{\Delta(y) \Delta(\zeta)} \quad (5.5)$$

to evaluate the angular integral gives

$$\begin{aligned} \mathbf{Z}_{d\mu_{(\mathbf{c},\mathbf{d},\beta,n)}}(X) &= \frac{(\prod_{i=1}^{n-1} i!)}{\Delta(y)} \prod_{i=1}^n \left(\int_{\mathcal{C}_n} d\zeta_i A_n^{(\mathbf{c},\mathbf{d},\beta)}(\zeta_i) \right) \Delta(\zeta) \det(e^{y_i \zeta_j})_{1 \leq i,j \leq n} \\ &= \frac{(\prod_{i=1}^n i!)}{\Delta(y)} \det \left(f_{(\mathbf{c},\mathbf{d},\beta,n)}^{(i-1)}(y_j) \right)_{1 \leq i,j \leq n}, \end{aligned} \quad (5.6)$$

where we have used the Andréiev identity [8] in the second line. Comparing eqs. (5.6) and (4.13), we obtain the matrix integral representation (5.4) of $\tau^{(G_{\mathbf{c},\mathbf{d},\beta})}([X])$. \square

Acknowledgements. This work was partially supported by the Natural Sciences and Engineering Research Council of Canada (NSERC) and the Fonds de recherche du Québec, Nature et technologies (FRQNT).

References

- [1] J. Ambjørn and L. Chekhov, “The matrix model for dessins d’enfants”, *Ann. Inst. Henri Poincaré D* **1**, 337-361 (2014).
- [2] J. Ambjørn and L. Chekhov, “A matrix model for hypergeometric Hurwitz numbers”, *Theor. Math. Phys.* **181**, 1486-1498 (2014).
- [3] J. Ambjørn and L. Chekhov, “Spectral curves for hypergeometric Hurwitz numbers”, arXiv:1806:07265
- [4] A. Alexandrov, G. Chapuy, B. Eynard and J. Harnad, “Weighted Hurwitz numbers and topological recursion: an overview”, *J. Math. Phys.* **59**, 081102: 1-20 (2018).
- [5] A. Alexandrov, G. Chapuy, B. Eynard and J. Harnad, “Fermionic approach to weighted Hurwitz numbers and topological recursion,” *Commun. Math. Phys.* **360** no.2, 777 (2018).
- [6] A. Alexandrov, G. Chapuy, B. Eynard and J. Harnad, “Weighted Hurwitz numbers and topological recursion” arXiv::1806.09738
- [7] A. Alexandrov, A. Mironov, A. Morozov and S. Natanzon, “ On KP-integrable Hurwitz functions”, *JHEP* **1411** 080 (2014).
- [8] C. Andréief, “Note sur une relation pour les intégrales définies des produits des fonctions”, *Mém. Soc. Sci. Phys. Nat. Bordeaux* **2**, 1-14 (1883).
- [9] H. Bateman and A. Erdélyi, “Higher Transcendental Functions”, Vol. I (PDF). New York: McGraw–Hill ((1953). (See § 5.3, “Definition of the G-Function”, p. 206.)
- [10] G. Borot, B. Eynard, M. Mulase and B. Safnuk, “A matrix model for simple Hurwitz numbers, and topological recursion”, *J. Geom. Phys.* **61**, 522–540 (2011).
- [11] E. Brézin and S. Hikami, “Correlations of nearby levels induced by a random potential”, *Nucl. Phys.* **B 479:3** 697-706 (1996).
- [12] V. Bouchard and M. Mariño, “Hurwitz numbers, matrix models and enumerative geometry”, *Proc. Symp. Pure Math.* **78**, 263-283 (2008).
- [13] B. Eynard, “Counting surfaces”, CRM Aisenstadt chair lectures, Progress in Mathematical Physics, 70. Birkhauser/Springer (2016).
- [14] B. Eynard and N. Orantin, “Algebraic methods in random matrices and enumerative geometry”, *Topical review J. Phys. A: Math. Theor.* **42**, 293001 (2009).

- [15] B. Eynard and N. Orantin, “Topological recursion in enumerative geometry and random matrices”, *J. Phys. A* **42**, 293001 (2009).
- [16] G. Frobenius, “Über die Charaktere der symmetrischen Gruppe”, *Sitzber. Preuss. Akad. Berlin*, 516-534 (1900).
- [17] G. Frobenius, “Über die Charakterische Einheiten der symmetrischen Gruppe”, *Sitzber. Akad. Wiss., Berlin*, 328-358 (1903). Gesammelte Abhandlung III, 244-274.
- [18] William Fulton and Joe Harris, “Representation Theory”, Graduate Texts in Mathematics, **129**, Springer-Verlag (N.Y., Berlin, Heidelberg, 1991), Chapt. 4, and Appendix A.
- [19] I.P. Goulden and D.M. Jackson, “The KP hierarchy, branched covers, and triangulations” *Adv. Math.* **219**, 932-951 (2008).
- [20] I. P. Goulden, M. Guay-Paquet and J. Novak, “Monotone Hurwitz numbers and the HCIZ Integral”, *Ann. Math. Blaise Pascal* **21**, 71-99 (2014).
- [21] M. Guay-Paquet and J. Harnad, “2D Toda τ -functions as combinatorial generating functions”, *Lett. Math. Phys.* **105**, 827-852 (2015).
- [22] M. Guay-Paquet and J. Harnad, “Generating functions for weighted Hurwitz numbers”, *J. Math. Phys.* **58**, 083503 (2017).
- [23] K. I. Gross and D. St. P. Richards, “Total positivity, spherical series, and hypergeometric functions of matrix argument”, *J. Approx. Theory*, **59**, no. 2, 224-246 (1989).
- [24] J. Harnad, “Weighted Hurwitz numbers and hypergeometric τ -functions: an overview”, *Proc. Symp. Pure Math.* **93**, 289-333 (2016).
- [25] J. Harnad, “Quantum Hurwitz numbers and Macdonald polynomials”, *J. Math. Phys.* **57** 113505 (2016).
- [26] J. Harnad, “Multispecies weighted Hurwitz numbers”, *SIGMA* **11** 097 (2015).
- [27] J. Harnad and F. Balogh, “ τ -functions and their applications”, *Mathematical physics monographs series*, Cambridge Univ. Press, Chapt. 5, Secs. 5.7, 5.12 (in press, 2019).
- [28] Harish-Chandra, “Differential operators on a semisimple Lie algebra”, *Amer. J. Math.* **79**, 87-120 (1957).
- [29] J. Harnad and A. Yu. Orlov, “Hypergeometric τ -functions, Hurwitz numbers and enumeration of paths”, *Commun. Math. Phys.* **338**, 267-284 (2015).
- [30] A. Hurwitz, “Über Riemann’sche Fläsche mit gegebenise Verzweigungspunkten”, *Math. Ann.* **39**, 1-61 (1891); *Matematische Werke I*, 321-384.
- [31] A. Hurwitz, “Über die Anzahl der Riemann’sche Fläsche mit gegebenise Verzweigungspunkten”, *Math. Ann.* **55**, 53-66 (1902); *Matematische Werke I*, 42-505.

- [32] C. Itzykson C and J.-B. Zuber “The planar approximation. II ”, *J. Math. Phys.* **21**, 411-421 (1980).
- [33] M. Kazarian and P. Zograf, “Virasoro constraints and topological recursion for Grothendieck’s dessin counting”, *Lett. Math. Phys.* **105** 1057-1084 (2015).
- [34] S. K. Lando and A.K. Zvonkin, *Graphs on Surfaces and their Applications*, Encyclopaedia of Mathematical Sciences, Volume **141**,
- [35] I. G. Macdonald, *Symmetric Functions and Hall Polynomials*, Clarendon Press, Oxford, (1995).
- [36] A. Mironov and A. Morozov, “Virasoro constraints for Kontsevich-Hurwitz partition function”, *JHEP* 02 024 (2009).
- [37] *Handbook of Mathematical Functions*, National Institute of Standards and Technology (NIST), U.S. Department of Commerce, (F. W. J. Olver, D. W. Lozier; R. F. Boisvert, C. W. Clark, eds.), Cambridge University Press (2010). ISBN 978-0-521-19225-5. MR 2723248.
- [38] A. Okounkov, “Toda equations for Hurwitz numbers”, *Math. Res. Lett.* **7**, 447–453 (2000).
- [39] A. Okounkov and R. Pandharipande, “Gromov-Witten theory, Hurwitz theory and completed cycles,” *Ann. Math.* **163**, 517 (2006).
- [40] A. Yu. Orlov, “New solvable matrix integrals”, *Int. J. Mod. Phys.* **19**, supp. 2 , 276-293 (2004).
- [41] A. Yu. Orlov and D. M. Scherbin, “Fermionic representation for basic hypergeometric functions related to Schur polynomials,” *Theor. Math. Phys.* **137**, 1574–1589 (2003).
- [42] A. Yu. Orlov and D. M. Scherbin, “Hypergeometric solutions of soliton equations”, *Theor. Math. Phys.* **128**, 906-926 (2001).
- [43] M. Sato, “Soliton equations as dynamical systems on infinite dimensional Grassmann manifolds”, RIMS, Kyoto Univ. *Kokyuroku* **439**, 30-46 (1981).
- [44] M. Sato and Y. Sato. “Soliton equations as dynamical systems on infinite dimensional Grassmann manifold”, in: Nonlinear PDE in Applied Science, Proc. U. S.-Japan Seminar, Tokyo 1982, Kinokuniya, Tokyo, 1983, pp. 259-271.
- [45] I. Schur, “Neue Begründung’ der Theorie der Gruppencharaktere”, *Sitzber. Akad. Wiss., Berlin*, 406-432 (1905).
- [46] G. Segal and G. Wilson, “Loop groups and equations of KdV type ”, *Publ. Math. IHÉS* **6**, 5-65 (1985).
- [47] P. Zograf, “Enumeration of Grothendieck’s dessins and KP hierarchy”, *Int. Math Res. Notices* **24**, 13533-13544 (2015).