

Time-resolved CCD photometry and time-series analysis of RR Lyrae type RR Gem star

V. A. Gohil and S.P. Bhatnagar

Department of Physics, M. K. Bhavnagar University, Bhavnagar - 364002, Gujarat, India;
spb@mkbhavuni.edu.in

Received 2019 March 24; accepted 2019 April 09

Abstract We present the results of a time-resolved photometric and time-series analysis of a RR Lyrae type star RR Gem. The main results are as follows: We found RR Gem's pulsation period, 0.39689 d, and its V and I mean magnitudes, 11.277 (V) and 11.063 (I) mag respectively. We confirm its variability type as RRab/BL because of its showing the Blažko effect and it also shows asymmetric light curves (steep ascending branches), periods from 0.3 to 1.0 days, and amplitudes from 0.3 to 2 mag. in V. They are fundamental mode pulsators.

Key words: Stars: individual: RR Gem — stars: variables: RR Lyr — stars: oscillations — stars: variables: RRAB/BL — stars: variables: Blazhko effect — techniques: photometric.

1 INTRODUCTION

RR Lyrae variables are older pulsating white giants with low metallicity. They are common in globular clusters-dense groups of old stars in the halos of galaxies. Like Cepheids, their pulsations are periodic. RR Lyraes have 0.5 solar mass and have a short pulsation period of 0.2 to 0.85 days and amplitude variations of 0.3 to 2 magnitudes. RR Lyrae stars are usually spectral class A. The main sequence and the HB(Horizontal Branch) do not intersect. RR Lyrae stars are on the HB, indeed where the instability strip intersects the HB. The HB stars have left the red giant branch and are characterized by helium fusion in their cores surrounded by a shell of hydrogen fusion. All of the RR Lyrae stars in a cluster have the same average apparent magnitude. In different clusters, the average apparent magnitude is different. This is because all RR Lyraes have about the same average absolute magnitude of +0.75.

RR¹ means that this is the RR variable in the Lyrae constellation, which are radially-pulsating giant A-F stars having amplitudes from 0.2 to 2 mag. in V band. Cases of variable light-curve shapes as well as variable periods are known. If their amplitudes are modified and also their periods, then they are called the "Blažko effect"(Blažko 1907)(denoted by the sub type BL). The majority of these stars belong to the spherical component of the Galaxy; they are present, sometimes in large numbers, in some globular clusters, where they are known as pulsating horizontal-branch stars. Like Cepheids, maximum expansion velocities of surface layers for these stars practically coincide with maximum light.

¹ <https://www.aavso.org/vsx/index.php?view=help.vartype&nolayout=1&abbrev=RR>

RRAB/BL² type variable stars are RR Lyrae variables with asymmetric light curves (steep ascending branches), periods from 0.3 to 1.0 days, and amplitudes from 0.5 to 2 mag. in V band. They are fundamental mode pulsators. BL indicates RR Lyrae stars showing the Blažko effect.

J. Jurcsik and her team (Jurcsik 1990, 1998; Jurcsik et al. 2005, 2006a,b,c, 2008a,b; Jurcsik 2009a; Jurcsik et al. 2009b,c) has done a remarkable work on the Blazhko behavior of RR Gem Star and in particular RR Lyr type stars showing Blažko effects. Most RR Lyrae stars repeat their light curves with remarkable regularity. About 30 % of the known galactic RR Lyrae stars, however, display cyclic modulation in shape and amplitude of the light curve over tens of pulsation cycles.

2 METHODOLOGY

2.1 Instrumentation

M. K. Bhavnagar University's Kumari Aanya Binoy Gaardi Observatory³ was used for this study. The Fully automated Observatory contains a 14 inch Celestron Schmidt-Cassegrain reflector type Telescope mounted on a German Equatorial mount. Telescope has an SBIG STF-7 CCD Camera along with SBIG CFW-8 filter wheel with Johnson-Cousins *UBVRI*⁴ photometric filters inbuilt into the CCD are attached on the telescope. Telescope is manual as well as fully computer controlled and telescope has two stepper motors for RA and DEC axis which is controlled by stepper motor controllers indigenously developed in collaboration with IUCAA, Pune(Inter University Center for Astronomy and Astrophysics). Open source software SCOPE is used to control the stepper motor controller and Cartes du Ciel(Sky Chart) was used for the CCD control as well as to give positions commands to SCOPE software which send command to stepper motor controller. Stepper motor controller performs the PWM and send electrical signals to RA and DEC stepper motors. There is an USB connection between CCD and PC for data as well as commands. Astronomy data is locally stored in Windows XP based PC. A GPS receiver is also connected with Windows PC which periodically updates time of the PC which in turn updates the DOC based PC's clock and thus the whole system's time gets updated periodically. Manual sliding rooftop is connected with a motor through pulley for opening and closing down the Observatory's roof.

2.2 Observations and data reduction

2.2.1 Differential Photometry

MaxIm DL⁵ photometric image analysis software was used for our data reduction work. Each and every time-series astronomical image was first calibrated for basic noise removal and then every continuous sequence was aligned and then photometry was performed on them.

For doing differential photometry(Rodríguez 2005) we needed our object star(RR Gem)⁶, one or more check stars and one reference star(standard star). Standard observatory procedures were used and used with minimum air-mass considerations.

For data reduction(calibration), flats field frames, dark frames and bias frames as well as light frames(See Figure 1.) were required so we programmed it into our MaxIm DL software and then it's done automatically according to our exposure time and other Johnson-Cousins *ubvri* filter value settings.

² <https://www.aavso.org/vsx/index.php?view=help.vartype&nolayout=1&abbrev=RRAB%2FBL>

³ <https://www.google.com/maps/place/Bhavnagar+Observatory/@21.7537555,72.1302003,18z/data=!4m5!3m4!1s0x395f509a719b3a2b:0x30504b9860ad6057!8m2!3d21.7542085!4d72.1303511?hl=en>

⁴ <http://www.aip.de/en/research/facilities/stella/instruments/data/johnson-ubvri-filter-curves>

⁵ <http://diffractionlimited.com/product/maxim-dl/>

⁶ <https://www.aavso.org/vsx/index.php?view=detail.top&oid=14304>




Fig. 1: Clear Filter Object, Comparison and Check Star Field-CLEAR Filter: 16-04-2014

For flat fielding we used daily twilight images and after all these we made master files of each of these and used them in our final calibration part.

In MaxIm DL we combined all the flats, darks and bias files and combined them into single standard master files according to different photometry filters. For calibration we used these master files. We first opened all the object star's light images and loaded the master calibration files and performed calibrate using calibrate all menu. By this we get all the files calibrated, then we again opened the calibrated light images and align them using the align tool.

Now for performing Photometry we again opened all the light images and using the photometry menu's differential photometry tool we performed the photometry on the light images by selecting each individual object star, reference star and check stars. Finally it gave us results in a Magnitude Vs JD(Jullian Day) graph in CSV or AAVSO Text format file.

2.2.2 Time-series analysis

Vstar(AAVSO)⁷ program was used for performing time-series analysis (Rodríguez 2005). We used DC DFT with Period Range tool and fed the proper values in it like low period as 0.1 d, high period as 0.5 d and resolution as 0.0001 and we got the periodogram with period 0.39689 d.

We downloaded the standard published data for our star from AAVSO data tool⁸ for the period of 10 months to get the bigger view of our star and performed the same above mentioned process and got the results(See Figures 2 and 3) then we also downloaded about 1 months data near to our observing period and analyzed the same and got this results(See Figures 4 and 5). By comparing this one month's data with our data(See Figures 6, 7 for (V) and 8, 9 for (I) respectively) we found it to be nearly perfectly matching with the magnitude as well as with the period. This ascertains that our observations and data reduction process is standardized.

⁷ <https://www.aavso.org/vstar>

⁸ <https://www.aavso.org/data-download>




Fig. 2: RR GEM 10 Months Data 2014 Johnson V Periodogram



Fig. 3: RR GEM 10 Months Phase Data 2014 Johnson V Phase Graph




Fig. 4: RR GEM 1 Months Data 2014 Johnson V Periodogram




Fig. 5: RR GEM 1 Months Phase Data 2014 Johnson V Phase Graph




Fig. 6: RR GEM 15 Days Our Data 2014 Johnson V Periodogram




Fig. 7: RR GEM 15 Days Phase Our Data 2014 Johnson V Phase Graph




Fig. 8: RR GEM 15 Days Our Data 2014 Johnson I Periodogram




Fig. 9: RR GEM 15 Days Phase Our Data 2014 Johnson I Phase Graph

According to AAVSO(VSX)⁹ its spectral type is A9-F6. It was discovered by L. Ceraski (Ceraski, L. 1903) and all these information is based on 09 Feb 1972 (HJD 2441357.205). Its AAVSO UID : 000-BBM-940. Its other names are AAVSO 0715+31, AN 13.1903, HIP 35667. Our data is from JD: 2456764.16483 (16 APR 2014) to JD: 2456801.11748 (23 MAY 2014) totaling 15 nights. During this time the mean magnitudes in visual filter and infrared filters were 11.277 (V) and 11.063 (I) respectively.

We also compared our results with other astronomical databases. The details are as shown in Table 1. As shown in Table 1, we compared our results with many astronomical databases like VSX, ASAS-SN¹⁰, CDS¹¹, GCVS¹² and General Variable Star Search Gateway¹³. We found our mean magnitude in V band: 11.277 mag and in I band: 11.063 mag, which is quite close to the published databases values

⁹ <https://www.aavso.org/vsx/index.php?view=search.top>

¹⁰ https://asas-sn.osu.edu/database/light_curves/195828

¹¹ <http://cdsportal.u-strasbg.fr/?target=rr%20gem>

¹² <http://www.sai.msu.su/gcvs/cgi-bin/search.cgi?search=RR+Gem>

¹³ <http://var2.astro.cz/gsg/index.php?star=rr+gem&all=yes&alldata=yes&oejv=yes&gcvs=yes&nsv=yes&brka=yes&meka=yes&czev=yes&bcbvs=yes&dssplate=yes&usecoords=GCVS&rezim=search.now>

Database	ID / Name	J2000.0		Type	V mag (Max)	V mag (Min)	Period (d)	Epoch	Spectral Class ^a
		(RA)	(DEC)						
VSX ^b	000-BBM-940	07:21:33.53	30:52:59.50	RRab/BL	10.62	11.99	0.3973106	2441357.205(HJD)	A9-F6
ASAS-SN ^c	ASASSN-VJ072133.66+305259.7	07:21:33.66	30:52:59.70	RRab	-	11.34	0.3992869	2457690.06(HJD)	-
CDS ^d	V* RR Gem	07:21:33.53	30:52:59.46	Variable Star of RR Lyr type	-	11.92	-	-	A8
GCVS ^e	RR Gem	07:21:33.53	30:52:59.50	RRab	10.62	11.99	0.3973106	41357.21	A9-F6
GVSSG ^f	RR Gem / 380010	07:21:33.50	30:52:59.50	RRab	10.62	11.99	0.3973106	41357.21(JD)	A9-F6

^a <https://www.handprint.com/ASTRO/specclass.html>^b Variable Star Index(AAVSO).^c The All Sky Automated Survey for SuperNovae.^d Strasbourg astronomical Data Center(CDS-Strasbourg).^e General Catalog of Variable Stars.^f General Variable Star Search Gateway.

Table 1: RR Gem in different Astronomical databases.

like 11.185 mag in V band with the difference between our results and VSX(AAVSO) with the minor difference of 0.092, which is quite close.

We also compared our period with the astronomical databases and found that our period 0.39689 d is very close to the published value 0.3973106 d VSX(AAVSO) with the minor difference of 0.0004206 d, which is also quite close.

2.2.3 Comparison and Check Stars

We used Check and Comparison Stars¹⁴ from Field photometry¹⁵ for RR Gem from AAVSO Variable Star Database:

Star type	AUID	J2000.0		Magnitudes				
		(RA)	(DEC)	(B)	(V)	(B - V)	(Rc)	(Ic)
Standard / Comparison Star	000-BJR-395	07:21:40.41	30:52:23.80	9.560(0.020)*	9.412(0.016)*	0.148(0.026)	9.322(0.020)*	9.219(0.021)*
Check Star	000-BBM-938	07:21:33.34	30:54:43	10.652(0.016)*	10.271(0.014)*	0.381(0.021)	10.043(0.017)*	9.826(0.019)*

Table 2: Comparison and Check Star Table for RR Gem.

The comparison / standard star and check stars were selected from the AAVSO's field photometry section. They were selected because they were very near, in the same CCD field. They were chosen because they were about the same magnitude as well as were within the CCD field. All the details are given in the table 2. Practically they should be near to your object star and also their magnitude as well as spectral class should also match. Comparison star should be a standard non-variable star and check star could be non-variable or variable for the aperture(differential) photometry. So, they were chosen accordingly and we found them to be of satisfactory nature as per our requirements.

3 RESULTS

RR Gem's mean magnitude for V was 11.277 (V) mag and for I was 11.063 (I) mag. We had also compared star's data from AAVSO Variable star database for near about the same time period and found its visible filter's mean magnitude value to be 11.185 V, which gives the difference of 0.092 V which is also quite negligible.

After performing time-series analysis it was found that the period of RR Gem star published in the VSX(AAVSO) database was 0.3973106 d and after our analysis we found it to be 0.39689 d, the difference between the two is 0.0004206 d which is quite low. Thus it establishes our analysis matches with the published literature values.

¹⁴ <https://www.aavso.org/apps/vsp/photometry/?fov=18.5&scale=F&star=rr+gem&orientation=ccd&maglimit=18.5&resolution=150&north=up&east=left&type=chart>¹⁵ <https://www.aavso.org/apps/vsp/photometry/?fov=18.5&scale=F&star=rr+gem&orientation=ccd&maglimit=18.5&resolution=150&north=up&east=left&type=photometry>

4 CONCLUSION

We have established that our observations reported in this paper are in very good agreement with the published data of the star RR Gem and that it is an RR Lyrae type star and its variability type is RRab/BL and is clearly showing the Blazhko effect. We also conclude that our observatory can be now used for future studies of variable stars and in particular will be useful for performing optical light curve data and related analysis for the upcoming transient alerts like GW and Gamma-ray bursts which require optical followups from ground based telescopes during the bright phases (at various geographic longitudes).

Acknowledgements We would like to acknowledge the M. K. Bhavnagar University for providing Astronomical research facility at Kumari Aanya Binoy Gardi Observatory and We would also like to thank, IUCAA(Inter University Center For Astronomy And Astrophysics), AAVSO(American Association for Variable Star Observers), VSX(Variable Star Index)(AAVSO).

References

- Blažko, 1907, AN, 175, 325
 Juresik J., Kovacs G., 1990, A & A, 312, 111
 Juresik J., Johanna, 1998, A & A, 333, 571
 Juresik J., Sódor, Á., Váradi, M., Szeidl, B., Washuettl, A., Weber, M., Dékány, I., Hurta, Zs., Lakatos, B., Posztobányi, K., 2005, A & A, 430, 1049
 Juresik J., Szeidl, B., Sódor, Á., Dékány, I., Hurta, Zs., Posztobányi, K., Vida, K., Váradi, M., Szing, A., 2006, AJ, 132, 61
 Juresik J., Sódor, Á., Váradi, M., Vida, K., Posztobányi, K., Szing, A., Hurta, Zs., Dekany, I., Washuettl, A., Vityi, N., 2006, IBVS, 5709, 1
 Juresik J., Szeidl, B., Váradi, M., Henden, A., Hurta, Zs., Lakatos, B., Posztobányi, K., Klagyivik, P., Sódor, Á., 2006, A & A, 445, 617
 Juresik J., Sódor, Á., Hurta, Zs., Kovari, Zs., Vida, K., Hajdu, G., Nagy, I., Dekany, I., Posztobányi, K., Koponyas, B., 2008, IBVS, 5844, 1
 Juresik J., Sódor, Á., Hurta, Zs., Váradi, M., Szeidl, B., Smith, H. A., Henden, A., Dkny, I., Nagy, I., Posztobányi, K., 2008, MNRAS, 391, 164
 Juresik J., 2009, CoAst, 159, 53
 Juresik J., Sódor, Á., Szeidl, B., Kolláth, Z., Smith, H. A., Hurta, Zs., Váradi, M., Henden, A., Dékny, I., Nagy, I., 2009, MNRAS, 393, 1553
 Juresik J., Sódor, Á., Szeidl, B., Hurta, Zs., Váradi, M., Posztobányi, K., Vida, K., Hajdu, G., Kővári, Zs., Nagy, I., 2009, MNRAS, 400, 1006
 P. Rodriguez-Gil, 2005, A & A, 431, 289