SEMIMARTINGALE DYNAMICS AND ESTIMATION FOR A SEMI-MARKOV CHAIN

Robert J. Elliott

University of South Australia, Adelaide, Australia and University of Calgary, Calgary, AB, Canada

April 23, 2019

Abstract

We consider a finite state discrete time process X. Without loss of generality the finite state space can be identified with the set of unit vectors $\{e_1, e_2, \ldots, e_N\}$ with $e_i = (0, \ldots, 0, 1, 0, \ldots, 0)' \in \mathbb{R}^N$. For a Markov chain the times the process stays in any state are geometrically distributed. This condition is relaxed for a semi-Markov chain. We first derive the semimartingale dynamics for a semi-Markov chain. We then consider the situation where the chain is observed in noise. We suggest how to estimate the occupation times in the states and derive filters and smoothers for quantities associated with the chain.

Acknowledgments. The support of NSERC and the Australian Research Council is appreciated.

1 Introduction

The topic of investigation is a finite state, discrete time process

 $X = \{X_k, k = 0, 1, ...\}$. We suppose, without loss of generality, that the state space of X is the set $S = \{e_1, e_2, ..., e_N\}$, $e_i = (0, ..., 0, 1, 0, ..., 0)' \in \mathbb{R}^N$, of unit vectors in \mathbb{R}^N . If X is a Markov chain, the times the chain spends in any state are geometrically distributed random variables. For a semi-Markov chain this condition is relaxed and the occupation times can have more general distributions. Semi-Markov chains are related to renewal processes and have been used in applications since their introduction over 60 years ago. Their more general occupation times often are a better fit for empirical data.

The two main contributions of this note are

- 1. the semimartingale dynamics for the semi-Markov chain
- 2. the consequent extension to semi-Markov chains of the filtering, smoothing and estimation results for hidden semi-Markov chains, once an integer estimate of the occupation time has been obtained.

Earlier references on semi-Markov processes include the books by Koski [6], Barbu and Limnios [1], and van der Hoek and Elliott [8]. References on filtering include Yu [9], Krishnamurthy, Moore and Chung [7] and Elliott, Limnios and Swishchuk [4].

Unfortunately the references [4] and [8] include a few typing errors.

2 Dynamics

All processes are defined on a probability space (Ω, \mathcal{F}, P) .

Our process of interest is the discrete time, finite state, time-homogeneous semi-Markov chain $X = \{X_k, 0 \leq k\}$. Without loss of generality the state space of the process can be identified with the set of unit vectors

$$S = \{e_1, e_2, \dots, e_N\}$$
$$e_i = (0, \dots, 0, 1, 0, \dots, 0)' \in \mathbb{R}^N.$$

A semi-Markov chain is similar to a Markov chain except that the time it spends in any state is no longer geometrically distributed. **Notation 2.1.** With $X_0 \in S$, we suppose either X_0 , or its distribution $p_0 = (p_0^1, p_0^2, \ldots, p_0^N)' \in \mathbb{R}^N$, is known. Write τ_1 for the first jump time of X, when $X_{\tau_1} \neq X_0$ and more generally

 τ_n for the n^{th} jump time.

 $\mathcal{F} = \{\mathcal{F}_k, k = 0, 1, 2, ...\}, \text{ where } \mathcal{F}_k = \sigma\{X_0, X_1, ..., X_k\}, \text{ is the filtra$ tion generated by X.

Definition 2.2. X is a semi-Markov chain if

$$P(X_{\tau_{n+1}} = e_j, \tau_{n+1} - \tau_n = m | \mathcal{F}_{\tau_n})$$

= $P(X_{\tau_{n+1}} = e_j, \tau_{n+1} - \tau_n = m | X_{\tau_n} = e_i)$
= $P(\tau_{n+1} - \tau_n = m | X_{\tau_{n+1}} = e_j, X_{\tau_n} = e_i) P(X_{\tau_{n+1}} = e_j | X_{\tau_n} = e_i)$
= $f_{ji}(m) p_{ji}$, say,

where

$$P(\tau_{n+1} - \tau_n = m | X_{\tau_{n+1}} = e_j, X_{\tau_n} = e_i) = f_{ij}(m)$$

and

$$P(X_{\tau_{n+1}} = e_j | X_{\tau_n} = e_i) = p_{ji}.$$

Assumption 2.3. We shall suppose $f_{ij}(m)$ does not depend on e_j and write

$$P(\tau_{n+1} - \tau_n = m | X_{\tau_{n+1}} = e_j, X_{\tau_n} = e_i) = P(\tau_{n+1} - \tau_n = m | X_{\tau_n} - e_i)$$
$$= \pi_i(m), \quad m = 1, 2, \dots$$

That is, for each $i, 1 \leq i \leq N, \{\pi_i(m), m = 1, 2, ...\}$ is a probability distribution on the positive integers.

Also, recall we assumed X is homogeneous in time so the above probabilities are independent of n.

Notation 2.4. Write

$$G_i(k) := P(\tau_{n+1} - \tau_n \le k | X_n = e_i)$$
$$= \sum_{m=1}^k \pi_i(m)$$

$$F_{i}(k) := P(\tau_{n+1} - \tau_{n} > k | X_{n} = e_{i})$$

= 1 - G_{i}(k).

Then

$$P(\tau_{n+1} = \tau_n + k + 1 | X_{\tau_n + k} = X_{\tau_n} = e_i)$$

= $P(\tau_{n+1} = \tau_n + k + 1 | \tau_{n+1} > \tau_n + k, X_{\tau_n} = e_i)$
= $\frac{\pi_i(k+1)}{F_i(k)} = \Delta^i(k)$, say.

Definition 2.5. For $1 \le i \le N$ define the process $h_k^i(X_k)$ by the recursion

$$h_0^i(X_0) = \langle X_0, e_i \rangle$$

$$h_k^i(X_k) = \langle X_k, e_i \rangle + \langle X_k, e_i \rangle \langle X_k, X_{k-1} \rangle h_{k-1}^i(X_{k-1}).$$

Then $h_k^i(X_k)$ measures the amount of time X has spent in state e_i since it last jumped to e_i .

If

$$h_k(X_k) = \sum_{i=1}^N h_k^i(X_k)$$

then $h_0(X_0) = 1$ and

$$h_k(X_k) = 1 + \langle X_k, X_{k-1} \rangle h_{k-1}(X_{k-1}).$$

Then $h_k(X_k)$ measures the amount of time since the last jump.

Lemma 2.6. Suppose $i \neq j$, $1 \leq i, j \leq N$. Then

$$P(X_{t+1} = e_j | X_t = e_i, h_t(X_t)) = p_{ji} \Delta^i (h_t^i(X_t)).$$

Proof. We first make some observations from the given data. As

$$X_t = e_i, \ h_t(X_t) = h_t^i(X_t).$$

Also

$$\tau_n = t - h_t(X_t) + 1$$
 so $X_t = X_{\tau_n} = e_i$.

Finally

$$\tau_{n+1} = t + 1.$$

Consequently

$$P(X_{t+1} = e_j | X_t = e_i, h_t(X_t))$$

= $P(X_{\tau_{n+1}} = e_j | X_{\tau_n} = e_i \text{ and } \tau_{n+1} = \tau_n + h_t(X_t))$
= $p_{ji}\Delta^i(h_t(X_t)).$

г			
L			
L			
L			
÷	-	-	-

Remark 2.7. We are assuming there is a jump from e_i to a different e_j , $i \neq j$, at time t + 1.

So, given there is a jump

$$\sum_{\substack{j=1\\j\neq i}}^n p_{ji} = 1.$$

Corollary 2.8. Under the same hypotheses, if there is not a jump at t + 1

$$P(X_{t+1} = e_i | X_t = e_i, h_t(X_t))$$

= $1 - \Delta^i (h_t^i(X_t))$
= $1 - \Delta^i h_t^i(X_t) \sum_{\substack{j=1 \ j \neq i}}^n p_{ji}.$

Notation 2.9. Write for k = 1, 2, ..., A(k) for the matrix with entries

$$a_{ii}(k) = 1 - \Delta^{i}(k)$$
$$a_{ji}(k) = p_{ji}\Delta^{i}(k).$$

Example 2.10. For N = 3

$$A(k) = \begin{pmatrix} 1 - \Delta^{1}(k) & p_{12}\Delta^{2}(k) & p_{13}\Delta^{3}(k) \\ p_{21}\Delta^{1}(k) & 1 - \Delta^{2}(k) & p_{23}\Delta^{3}(k) \\ p_{31}\Delta^{1}(k) & p_{32}\Delta^{2}(k) & 1 - \Delta^{3}(k) \end{pmatrix}.$$

Notation 2.11. Define the matrices:

 $\Pi = (p_{ij}, 1 \le i, j \le N)$

where $p_{ii} = -1$ and p_{ji} is, as above, for $i \neq j$.

$$D(k) = \operatorname{diag}(\Delta^{1}(k), \Delta^{2}(k), \dots, \Delta^{N}(k)).$$

Then

$$A(k) = I + \Pi D(k)$$

where I is the $N \times N$ identity matrix.

For the case when N = 3

$$\Pi = \begin{pmatrix} -1 & p_{12} & p_{13} \\ p_{21} & -1 & p_{23} \\ p_{31} & p_{32} & -1 \end{pmatrix} \qquad D(k) = \begin{pmatrix} \Delta^1(k) & 0 & 0 \\ 0 & \Delta^2(k) & 0 \\ 0 & 0 & \Delta^3(k) \end{pmatrix}.$$

A key result is the following representation of the semi-Markov chain X.

Theorem 2.12. The semi-Markov chain X has the following semi-martingale dynamics:

$$X_{k+1} = A(h_k(X_k))X_k + M_{k+1} \in \mathbb{R}^N$$

Here M_{k+1} is a martingale increment:

$$E[M_{k+1}|X_k, h_k(X_k)] = 0 \in \mathbb{R}^N$$

Proof. We need only remark that we can write $h_k(X_k)$ in $A(h_k(X_k))$ rather than the different occupation times $h^1(X_k), \ldots, h^N(X_k)$.

This is because the components of X_k are in effect indicator functions. If $X_k = e_i$ the product $A(h_k(X_k))e_i$ selects the i^{th} column of $A_k(h_k(X_k))$ and the h(k) will be that for $h_k^i(X_k)$.

3 Full Markov Dynamics

We now consider an infinite state space for the semi-Markov chain and give Markov dynamics.

Remark 3.1. The complete state of our semi-Markov chain X at a time k is given by the state of the chain X_k and the number of time steps h_k the chain has been in that state. Suppose

$$X_k = e_i$$
 and $h_k = r$.

Then at time k + 1 either $X_{k+1} = e_i$ and $h_{k+1} = r + 1$ or $X_{k+1} = e_j$, $j \neq i$, and $h_{k+1} = 1$.

That is the state space of the 'complete' process is $S \times \mathbb{N}$, where $\mathbb{N} \in \{1, 2, ...\}$ is the natural numbers. Then with the above notation

$$(e_i, r) \rightarrow (e_i, r+1)$$
 with probability $1 - \Delta^i(r)$

or

$$(e_i, r) \to (e_j, 1)$$
 with probability $p_{ji} \Delta^i(r)$.

Suppose N = 3 so $S = \{e_1, e_2, e_3\}$. A state space for this 3-state semi-Markov chain can be identified with countably many copies of S, so, for example $(e_1, 1)$ corresponds to the infinite vector (1, 0, 0, |0, ...)'.

The first three components of this infinite column vector correspond to the states $(e_1, 1), (e_2, 1), (e_3, 1)$. The second three correspond to $(e_1, 2), (e_2, 2), (e_3, 2)$ and so on. If at time 1 the chain is in state

$$(e_1, 1) = (1, 0, 0|0, 0, 0|0, 0, \dots)'$$

this can become either

$$(e_1, 2) = (0, 0, 0|1, 0, 0|0, 0, \dots)'$$

with probability $(1 - \Delta^1(1))$, or

$$(e_2, 1) = (0, 1, 0|0, 0, 0|0, 0, \dots)'$$

with probability $p_{21}\Delta^1(1)$, or

$$(e_3, 1) = (0, 0, 1|0, 0, 0|0, 0, \dots)'$$

with probability $p_{31}\Delta^1(1)$.

There is then an infinite matrix C which describes these transitions. In this 3 state case write

$$\Pi(i) = \begin{pmatrix} 0 & p_{12}\Delta^2(i) & p_{13}\Delta^3(i) \\ p_{21}\Delta^1(i) & 0 & p_{23}\Delta^3(i) \\ p_{31}\Delta^1(i) & p_{32}\Delta^2(i) & 0 \end{pmatrix}$$

and

$$D(i) = \begin{pmatrix} 1 - \Delta^{1}(i) & 0 & 0 \\ 0 & 1 - \Delta^{2}(i) & 0 \\ 0 & 0 & 1 - \Delta^{3}(i) \end{pmatrix} .$$

With 0 representing the 3×3 zero matrix

$$C = \begin{pmatrix} \Pi(1) & \Pi(2) & \Pi(3) & \cdots \\ D(1) & 0 & 0 & \cdots \\ 0 & D(2) & 0 & \cdots \\ \cdot & \cdot & \cdot & \cdots \\ \cdot & \cdot & \cdot & \cdots \end{pmatrix}.$$

If we write the enlarged vectors as \overline{X}_k then the semi-martingale dynamics of the semi-Markov chain can be written as

$$\overline{X}_{k+1} = C\overline{X}_k + \overline{M}_{k+1} \in \overline{S}$$

where \overline{S} is the set of all infinite unit vectors of the form $\{e_{ij}\}$ where $e_{ij} = \delta_{ij}$, the Kronecker delta function $1 \leq i, j \leq \infty$.

This gives

$$E[\overline{X}_{k+1}|\overline{X}_k] = C\overline{X}_k$$

and

$$E[\overline{X}_{k+1}|\overline{X}_0] = C^{k+1}\overline{X}_0.$$

However, the approximate estimates defined in Section 5.3 below are possibly of more use in practise.

4 Observation Dynamics

The Viterbi and smoothing results of [5] are now adapted to this situation.

We suppose the semi-Markov chain X is not observed directly. Instead there is an observation sequence $y = \{y_0, y_1, \ldots, y_k, \ldots\}$ where

$$y_k = c(X_k) + d(X_k)w_k \,.$$

 $\{w_k, k = 0, 1, 2, \}$ is a sequence of i.i.d. N(0, 1) random variables. $c(\cdot)$ and $d(\cdot)$ are known real valued functions. Note that any real function $g(X_k)$ takes only the finite number of values $g(e_1), g(e_2), \ldots, g(e_N)$.

Write $g_k = g(e_k)$ and $g = (g_1, g_2, \dots, g_N)' \in \mathbb{R}^N$. Then

$$g(X_k) = \langle g, X_k \rangle \,.$$

Consequently there are vectors $c = (c_1, c_2, \ldots, c_N)$, $d = (d_1, d_2, \ldots, d_N)$ such that

$$c(X_k) = \langle c, X_k \rangle$$
 and $d(X_k) = \langle d, X_k \rangle$.

We suppose $d_k > 0$ for $k = 1, \ldots, N$.

Remark 4.1. We suppose the observation process y is scalar valued. The extension to a vector y is immediate.

Reference Probability 4.2.

We suppose there is a second 'reference' probability measure , $\overline{P},$ under which

1. the process X is still a semi-Markov chain with dynamics

$$X_{k+1} = A(h(k))X_k + M_{k+1} \in \mathbb{R}^N,$$

2. the process $y = \{y_0, y_1, ...\}$ is a sequence of i.i.d. N(0, 1) random variables.

From \overline{P} we now construct the original probability P under which

1. the process X is a semi-Markov chain with dynamics

$$X_{k+1} = A(h(k))X_k + M_{k+1}$$

2. The process $w = (w_0, w_1, ...)$ is a sequence of i.i.d. N(0, 1) random variables where

$$w_k = \frac{y_k - \langle c, X_k \rangle}{\langle d, X_k \rangle}.$$

Definition 4.3. For k = 0, 1, 2, ... write

$$\lambda_{k} = \frac{\phi((y_{k} - \langle c, X_{k} \rangle) / \langle d, X_{k} \rangle)}{\langle d, X_{k} \rangle \phi(y_{k})}$$

where $\phi(x)$ is the N(0,1) density $\frac{e^{-x^2}}{\sqrt{2\pi}}$.

$$\Lambda_{0,k} := \prod_{\ell=0}^k \,\lambda_\ell \,.$$

Recall

$$\mathcal{F} = \sigma\{X_0, X_1, \dots, X_k\}$$

and write

$$\mathcal{Y}_k = \sigma\{y_0, y_1, \dots, y_k\}$$
$$\mathcal{G}_k = \sigma\{X_0, \dots, X_k, y_0, \dots, y_k\}.$$

We consider the related filtrations $\{\mathcal{F}_k\}, \{\mathcal{Y}_k\}$ and $\{\mathcal{G}_k\}$.

Definition 4.4. The original 'real world' probability P is defined in terms of \overline{P} by setting

$$\frac{dP}{d\overline{P}}\left|_{\mathcal{G}_{k}}=\Lambda_{0,k}\right.$$

We can then prove

Lemma 4.5. Under P X is a semi-Markov chain with $X_{k+1} = A(h(X_k))X_k + M_{k+1}$ and $\{w_k, k = 0, 1, ...\}$ is a sequence of i.i.d. N(0, 1) random variables where

$$w_k = (y_k - \langle c, X_k \rangle) / \langle d, X_k \rangle.$$

That is, under $P \ y_k = \langle c, X_k \rangle + \langle d, X_k \rangle w_k$.

Proof. For a proof see [3].

5 Filter for X

The basic problem now is to obtain a recursive estimate for

$$E[X_k|\mathcal{Y}_k].$$

From Bayes' rule, see [3], this is

$$\frac{\overline{E}[\Lambda_k X_k | \mathcal{Y}_k]}{\overline{E}[\Lambda_k | \mathcal{Y}_k]}$$

where \overline{E} denotes expectation with respect to \overline{P} . Write $\overline{q}_k = \overline{E}[\Lambda_k X_k | \mathcal{Y}_k] \in \mathbb{R}^N$.

$$\gamma_{j}(y_{k+1}) = \frac{\phi((y_{k+1} - \langle c, e_{j} \rangle) / \langle d, e_{j} \rangle)}{\langle d, e_{j} \rangle \phi(y_{k+1})}$$
$$B(y_{k+1}) = \begin{pmatrix} \gamma_{1}(y_{k+1}) & 0 \dots & 0\\ 0 & \gamma_{2}(y_{k+1}) & 0\\ \vdots & \ddots & \vdots\\ 0 & \dots & 0 & \gamma_{k}(y_{k+1}) \end{pmatrix}$$

•

Notation 5.1. Recall $h_k(X_k)$ is the number of time steps X has stayed in state X_k since the last jump.

We shall define below a \mathcal{Y}_k measurable integer estimate \hat{h}_k of $h_k(X_k)$. This will be used in the recursion for q.

Theorem 5.2. With \hat{h}_k a \mathcal{Y}_t -measurable integer estimate of $h_k(X_k)$ a recursive estimate for $\overline{q}_k = \overline{E}[\Lambda_k X_k | \mathcal{Y}_k]$ is given by

$$q_{k+1} = B(y_{k+1})A(\hat{h}_k)q_k; \ q_0 = p_0.$$

Proof.

$$\overline{q}_{k+1} = \overline{E}[\Lambda_{k+1}X_{k+1}|\mathcal{Y}_{k+1}] = \overline{E}[\Lambda_k\lambda_{k+1}(A(h_k(X_k))X_k + M_{k+1})|\mathcal{Y}_{k+1}] = \overline{E}[\Lambda_k\lambda_{k+1}A(h_k(X_k))X_k|\mathcal{Y}_{k+1}].$$
(5.1)

Suppose \hat{h}_k is a \mathcal{Y}_k -measurable estimate of $h_k(X_k)$ and substitute this in (5.1).

Then an estimate of (5.1) is

$$q_{k+1} := \overline{E}[\Lambda_k \lambda_{k+1} A(\widehat{h}_k) X_k | \mathcal{Y}_{k+1}] = B(y_{k+1} A(\widehat{h}_k) q_k$$
(5.2)

where q_k is the analogous recursive estimate of $\overline{E}[\Lambda_k X_k | \mathcal{Y}_k]$.

With $1 = (1, 1, \dots, 1)' \in \mathbb{R}^N$

$$\langle \overline{q}_{k+1}, 1 \rangle = \overline{E}[\Lambda_{k+1} | \mathcal{Y}_{k+1}]$$

so a normalized estimate of X_k given by \mathcal{Y}_k is

$$\frac{q_k}{\langle q_k, 1 \rangle}$$

Recursion For \hat{h} 5.3

Suppose \hat{h}_k is known and \mathcal{Y}_k measurable. The MAP estimate for X_k given \mathcal{Y}_k is the state e_{i^*} corresponding to the maximum value of q_k^i , $i = 1, 2, \ldots, N$, or one of those states in case of equality. The recursion (5.2) gives an estimate for q_{k+1} which in turn suggests a MAP estimate e_{j^*} for X_{k+1} . That is e_{j^*} corresponds to the maximum value of q_{k+1}^i , $i = 1, 2, \ldots, N$. Then if $i^* = j^*$ $\hat{h}_{k+1} = \hat{h}_k + 1$. Otherwise $\hat{h}_{k+1} = 1$.

These estimates for $h_k(X_k)$ provide corresponding estimates $A(\hat{h}_k)$ for the transition matrix $A(h_k(X_k))$. In turn this provides the required recursive estimates.

We noted in the representation of Theorem 2.12 that $h_k(X_k)$ could be used rather than $h_k^i(X_k)$. In the recursion of Theorem 5.2 the unnormalized probabilities $(q_k^1, q_k^2, \ldots, q_k^N)$ are used. Observing these could give \mathcal{Y}_k integer valued estimates $(\hat{h}_k^i, \ldots, \hat{h}_N)$ of $(h_k^1(X_k), \ldots, h_k^N(X_k))$. These can be used in the terms $a_{ji}(\hat{h}_k^i)$ of A.

6 Estimates

As stated earlier, the key contributions of this paper are:

- 1. the semi-Martingale dynamics of Theorem 2.12 for the semi-Markov chain.
- 2. the recursive estimation \hat{h}_k of the integer $h_k(X_k)$ which is then substituted in the dynamics.

This provides a recursive estimate of the state of the chain given the observations y.

However, steps 1) and 2) above then allow immediate recursive estimates for the following quantities:

$$N_k^{ji} = \sum_{\ell=1}^k \left\langle X_{\ell-1}, e_i \right\rangle \left\langle X_\ell, e_j \right\rangle$$

the number of jumps from e_i to e_j

$$J_k^i = \sum_{\ell=1}^k \left\langle X_{\ell-1}, e_i \right\rangle$$

the amount of time spent in state e_i sums of the form

$$G_k^i = \sum_{\ell=1}^k f(y_\ell) \left\langle X_{\ell-1}, e_i \right\rangle.$$

For example, consider the unnormalized estimate

$$\overline{\sigma}(N_k^{ji}X_k) := \overline{E}[\Lambda_k N_k^{ji}X_k | \mathcal{Y}_k].$$

A recursion for an approximation σ is given by

Lemma 6.1.

$$\sigma(N_{k+1}^{ji}X_{k+1}) = B(y_{k+1})A(\widehat{h}_k)\sigma(N_k^{ji}X_k) + a_{ji}(\widehat{h}_k)\langle q_k, e_i\rangle\gamma_j(y_{k+1})e_j$$

where q_k is given in Theorem 5.2,

Proof. We suppose q_k , $\sigma(N_k^{ji}X_k)$ and \hat{h}_k have been determined. Then

$$\overline{E}[\Lambda_{k+1}N_{k+1}^{ji}X_{k+1}|\mathcal{Y}_{k+1}]$$

$$\simeq \overline{E}[\Lambda_k\lambda_{k+1}(N_k^{ji}+\langle X_{k+1},e_j\rangle\langle X_k,e_i\rangle)X_{k+1}|\mathcal{Y}_{k+1}].$$

Using Theorem 2.12 this is

$$\overline{E}[\lambda_{k+1}\Lambda_k N_k^{ji}A(\widehat{h}_k)X_k|\mathcal{Y}_k] + \overline{E}[\lambda_{k+1}\Lambda_k \langle A(\widehat{h}_k)X_k, e_j \rangle \langle X_k, e_i \rangle)\mathcal{Y}_k]e_j$$

and the result follows.

A recursion for an approximation σ of

$$\overline{\sigma}(G_k^i) = \overline{E}[G_k^i X_k | \mathcal{Y}_k]$$

is given by

Lemma 6.2.

$$\sigma(G_{k+1}^{i}X_{k+1}) = B(y_{k+1})A(\widehat{h}_{k})\sigma(G_{k}^{i}X_{k}) + f(y_{k+1})\langle q_{k}, e_{i}\rangle B(y_{k+1})A(\widehat{h}_{k})e_{i}.$$

In applications we need f(y) = 1, y or y^2 . For example, with $f(y) = 1, G_k^i = J_k^i$ and we have an approximate recursion

$$\sigma(J_{k+1}^i X_{k+1}) = B(y_{k+1}) A(\widehat{h}_k) \sigma(J_k^i X_k) + \langle q_k, e_i \rangle B(y_{k+1}) A(\widehat{h}_k) e_i.$$

Remark 6.3. Now $\langle X_k, \underline{1} \rangle = 1$ for all k so $\langle \sigma(N_k^{ji}X_k), 1 \rangle$ gives an unnormal-ized estimate of $\sigma(N_k^{ji})$.

In turn these provide estimates such as

$$\widehat{a}_{ji}(\widehat{h}_k) = \frac{\sigma(N_k^{ji})}{\sigma(J_k^j)}, \quad \text{for} \quad i \neq j,$$

and for the other parameters of the model as in [3].

7 Smoothers

Suppose $0 \le k \le T$ and we know $\{y_0, y_1, \ldots, y_T\}$. We wish to find $E[X_k | \mathcal{Y}_T]$. Using Bayes' theorem again, see [3],

$$E[X_k|\mathcal{Y}_T] = \frac{\overline{E}[\Lambda_{0,T}X_k|\mathcal{Y}_T]}{\overline{E}[\Lambda_{0,T}|\mathcal{Y}_T]}$$

Write $\Lambda_{k+1,T} = \prod_{\ell=k+1}^{T} \lambda_{\ell}$. Then $\Lambda_{0,T} = \Lambda_{0,k} \Lambda_{k+1,T}$ and

$$\overline{E}[\Lambda_{0,T}X_k|\mathcal{Y}_T] = \overline{E}[\Lambda_{0,k}X_k\overline{E}[\Lambda_{k+1,T}|\mathcal{Y}_{T^v}\mathcal{F}_k]\mathcal{Y}_T].$$

The process $\{X_k, h_k(X_k)\}$ is Markov as shown in Theorem .

Therefore

$$\overline{E}[\Lambda_{k+1,T}|\mathcal{Y}_{T^v}\mathcal{F}_k] = \overline{E}[\Lambda_{k+1,T}|\mathcal{Y}_{T^n}\sigma\{X_k, h_k(X_k)\}].$$

Suppose we have a sequence of MAP \mathcal{Y}_T -measurable integer estimates \hat{h}_k of the $h_k(X_k)$.

Substituting these in the dynamics of Theorem 2.12 , we have approximate dynamics for X under which it becomes a Markov chain.

Definition 7.1. Write $v_{T,T}^i = 1$ and

$$v_{k,T}^i := \overline{E}[\Lambda_{k+1,T} | \mathcal{Y}_{T^v} \{ X_k = e_i \}_v \{ h_k(X_k) = \widehat{h}_k \}]$$

Set $v_{k,T} = (v_{k,T}^1, v_{k,T}^2, \dots, v_{k,T}^N)' \in \mathbb{R}^N.$

Theorem 7.2. The process v satisfies the backward dynamics

$$v_{k,T} = A'(\hat{h}_k)B(y_{k+1})v_{k+1,t}$$

$$v_{T,T} = (1, 1, \dots, 1)' \in \mathbb{R}^N.$$
(7.1)

Proof.

$$\begin{split} \langle v_{k,T}, e_i \rangle &= v_{k,T}^i = \overline{E}[\Lambda_{k+2,T}\lambda_{k+1} | \mathcal{Y}_{T^v} \{X_k = e_i\}_v \widehat{h}_k] \\ &= \sum_{j=1}^N \overline{E}[\langle X_{k+1}, e_j \rangle \Lambda_{k+2,T} | \mathcal{Y}_{T^v} \{X_k = e_i\}_v \widehat{h}_k] \gamma_j(y_{k+1}) \\ &= \sum_{j=1}^N \overline{E}[\langle X_{k+1}, e_j \rangle \overline{E}[\Lambda_{k+2,T} | \mathcal{Y}_{T^v} \{X_k = e_i\}_v \widehat{h}_{k^v} \{X_{k+1}e_j\} v \widehat{h}_{k+1}] \\ &\times |\mathcal{Y}_{T^v} \{X_k = e_j\}_v \widehat{h}_k] \gamma_j(y_{k+1}) \\ &= \sum_{j=1}^N \overline{E}[\langle X_{k+1}, e_j \rangle \langle v_{k+1,T}, e_j \rangle | \mathcal{Y}_{T^v} \{X_k = e_i\}_v \widehat{h}_k] \gamma_j(y_{k+1}) \\ &= \sum_{j=1}^N a_{ji}(\widehat{h}_k) \langle v_{k+1,T}, e_j \rangle \gamma_j(y_{k+1}). \end{split}$$

This gives the result of (7.1).

Theorem 7.4. An unnormalized smoothed estimate for X_k given observations $\{y_0, y_1, \ldots, y_T\}$ is

$$\widetilde{q}_{k,T} := \overline{E}[\Lambda_{0,T}X_k|\mathcal{Y}_T] = (\operatorname{diag}\langle q_k, e_i \rangle)v_{k,T}.$$

A normalized smoothed estimate is then

$$\frac{\widetilde{q}_{k,T}}{\langle \widetilde{q}_{k,T}, 1 \rangle} \, .$$

Proof.

$$\overline{E}[\Lambda_{0,T} \langle X_k, e_i \rangle | \mathcal{Y}_T] = \overline{E} \left[\Lambda_{0,T} \langle X_k, e_i \rangle \overline{E}[\Lambda_{k+1,T} | \mathcal{Y}_{T^v} \{ X_k = e_i \}_v \widehat{h}_k] \mathcal{Y}_{0,T} \right] \\ = \langle q_k, e_i \rangle \langle v_{k,T}, e_i \rangle.$$

Therefore,

$$\overline{E}[\Lambda_{0,T}X_k|\mathcal{Y}_{0,T}] = \sum_{i=1}^N \langle q_k, e_i \rangle \langle v_{k,T}, e_i \rangle e_i = (\operatorname{diag} \langle q_k, e_i \rangle) v_{k,T}.$$

8 Filtering the Full Markov Dynamics

We showed in Section 3 that the state space of our discrete time, finite state semi-Markov chain X could be identified with $\overline{S} = S \times \mathbb{N}$ where $S = \{e_1, e_2, \ldots, e_N\}$ and $\overline{N} = \{1, 2, 3, \ldots\}$.

We observed that by identifying this state space \overline{S} with countably many copies of S, considered as an infinite column vector, X becomes a Markov chain \overline{X} with transition matrix

$$C = \begin{pmatrix} \Pi(1) & \Pi(2) & \Pi(3) & \cdots \\ D(1) & 0 & 0 & \cdots \\ 0 & D(2) & 0 & \cdots \\ \cdot & \cdot & \cdot & \cdots \\ \cdot & \cdot & \cdot & \cdots \end{pmatrix}$$

 \mathbf{SO}

$$\overline{X}_{k+1} = C \, \overline{X}_k + \overline{M}_{k+1} \in \overline{S}$$

where

$$E[\overline{X}_{k+1}|\overline{X}_k] = C \overline{X}_k.$$

Suppose there is a scalar observation process $y = \{y_k, k = 1, 2, ...\}$ such that

$$y_k = \langle \overline{c}, \overline{X}_k \rangle + \langle \overline{d}, X_k \rangle w_k$$

where $w = \{w_k, k = 1, 2, ...\}$ is a sequence of i.i.d. N(0, 1) random variables.

Here $\overline{c} = (c_1, c_2, ...)$ and $\overline{d} = (d_1, d_i, ...)$. Results of [5] extend immediately to this countable state Markov chain so that, for example, if a reference measure \overline{P} is introduced, as in Section 3, then with

$$\overline{q}_{k} = \overline{E} \left[\Lambda_{k} \overline{X}_{k} | Y_{k} \right] \in R^{\infty}$$

$$\gamma_{j}(y_{k+1}) = \frac{\phi \left((y_{k+1} - \langle \overline{c}, e_{j} \rangle) / \langle \overline{d}, e_{j} \rangle \right)}{\langle \overline{d}, e_{j} \rangle \phi(y_{k+1})}$$
and
$$\overline{B}(y_{k+1}) = \operatorname{diag} \gamma_{i}(y_{k+1})$$

$$\overline{q}_{k+1} = \overline{B}(y_{k+1}) C \overline{q}_{k}.$$

Filtering and smoothing results for quantities such as N^{ij} , J^i and G^i are exactly as before, except vectors with countably many states are involved.

9 Conclusion

A semi-Markov chain is a generalization of a Markov chain where the time spent in any state no longer has a geometric distribution. We derive the semimartingale dynamics of a semi-Markov chain. Using integer estimates of the times spent in a state we extend the filter, smoothing and estimation results obtained in earlier work. In addition, by considering a countably infinite state space the semi-Markov chain is a Markov chain and all earlier results extend to this situation

References

- Barbu, V.S., and Limnios, N., Semi-Markov Chains and Hidden Semi-Markov Models Towards Applications: Their Use in Reliability Theory and DNA Analysis. Springer 2008.
- [2] Cinlar, E., Introduction to Stochastic Processes. Prentice Hall, Chapter 10, 1975.
- [3] Elliott, R.J., Aggoun, L, and Moore, J.B., Hidden Markov Models: Estimation and Control. Springer Verlag, 1995/2008.
- [4] Elliott, R.J., Limnios, N. and Swishchuk, A., Filtering hidden semi-Markov chains, Statistics and Probability Letters, 83 (2018), 2007-2014.
- [5] Elliott, R.J. and Malcolm, W.P., Date-recursive smoother formulae for partially observed discrete time Markov chains, Stochastic Analysis and Applications (2006), 579-597.
- [6] Koski, T., Hidden Markov Models for Bio-informatics. Kluwer Academic Publishers 2001.
- [7] Krishnamurthy, V., Moore, J.B. and Chung, S.-H., On hidden fractal model signal processing, Signal Processing 24 (1991, 177-192.

- [8] van der Hoek, J. and Elliott, R.J., Introduction to Hidden Semi-Markov Models, Cambridge U. Press, London Math. Society Lecture Note Series 445, 2018, 174pp.
- [9] Yu, S.-Z., Hidden semi-Markov models, Artifical Intelligence 174 (2010), 215-243.