

The Bijectivity of Mirror Functors on Tori

Kazushi Kobayashi*

Abstract

By the SYZ construction, a mirror pair (X, \check{X}) of a complex torus X and a mirror partner \check{X} of the complex torus X is described as the special Lagrangian torus fibrations $X \rightarrow B$ and $\check{X} \rightarrow B$ on the same base space B . Then, by the SYZ transform, we can construct a simple projectively flat bundle on X from each affine Lagrangian multi section of $\check{X} \rightarrow B$ with a unitary local system along it. However, there are ambiguities of the choices of transition functions of it, and this causes difficulties when we try to construct a functor between the symplectic geometric category and the complex geometric category. In this paper, we prove that there exists a bijection between the set of the isomorphism classes of their objects by solving this problem.

Contents

1	Introduction	1
2	Preparations	4
3	Complex geometry side	5
3.1	The definition of $E_{(r,A,r',\mathcal{U},p,q)}$	6
3.2	The isomorphism classes of $E_{(r,A,r',\mathcal{U},p,q)}$	14
4	Symplectic geometry side	18
4.1	The definition of $\mathcal{L}_{(r,A,p,q)}$	18
4.2	The isomorphism classes of $\mathcal{L}_{(r,A,p,q)}$	20
5	Main result	24

1 Introduction

Let X be an n -dimensional complex torus, and we denote by \check{X} a mirror partner of the complex torus X . For this mirror pair (X, \check{X}) , the homological mirror

*Department of Mathematics, Graduate School of Science, Osaka University, Toyonaka, Osaka, 560-0043, Japan. E-mail : k-kobayashi@cr.math.sci.osaka-u.ac.jp. 2010 Mathematics Subject Classification : 14J33 (primary), 14F05, 53D37 (secondary). Keywords : torus, homological mirror symmetry, SYZ transform.

symmetry conjecture [12], which is proposed by Kontsevich in 1994, states that there exists an equivalence

$$D^b(\text{Coh}(X)) \cong \text{Tr}(\text{Fuk}(\check{X}))$$

as triangulated categories. Here, $D^b(\text{Coh}(X))$ is the bounded derived category of coherent sheaves on X , and $\text{Tr}(\text{Fuk}(\check{X}))$ is the derived category of the Fukaya category $\text{Fuk}(\check{X})$ on \check{X} [4] obtained by the Bondal-Kapranov-Kontsevich construction [3], [12]. Historically this conjecture has been first studied when (X, \check{X}) is a pair of elliptic curves (see [19], [18], [1] etc.), and after that, when (X, \check{X}) is a pair of abelian varieties of higher dimension generalizing the case of elliptic curves [5] (see also [13]).

On the other hand, the SYZ construction [20] by Strominger, Yau, and Zaslow in 1996, proposes a way of constructing mirror pairs geometrically. By this construction, the mirror pair (X, \check{X}) is realized as the trivial special Lagrangian torus fibrations $\pi : X \rightarrow B$ and $\tilde{\pi} : \check{X} \rightarrow B$ on the same base space B which is homeomorphic to an n -dimensional real torus. Here, for each point $b \in B$, the special Lagrangian torus fibers $\pi^{-1}(b)$ and $\tilde{\pi}^{-1}(b)$ are related by the T-duality. In particular, it is expected that the homological mirror symmetry on the mirror pair (X, \check{X}) is realized by the SYZ transform (an analogue of the Fourier-Mukai transform) along the special Lagrangian torus fibers of $\pi : X \rightarrow B$ and $\tilde{\pi} : \check{X} \rightarrow B$.

Concerning the above discussions, we explain the purpose of this paper. For a given mirror pair (X, \check{X}) , we regard it as the trivial special Lagrangian torus fibrations $\pi : X \rightarrow B$ and $\tilde{\pi} : \check{X} \rightarrow B$ in the sense of the SYZ construction. First, in the symplectic geometry side, we consider the full subcategory $\text{Fuk}_{\text{aff}}(\check{X})$ of the Fukaya category $\text{Fuk}(\check{X})$ consisting of affine Lagrangian multi sections of $\tilde{\pi} : \check{X} \rightarrow B$ with unitary local systems along them (in this paper, we sometimes call $\text{Fuk}_{\text{aff}}(\check{X})$ simply the Fukaya category). Then, according to the discussions in [14] and [2], we can obtain a holomorphic vector bundle on X which admits a constant curvature connection from each object of $\text{Fuk}_{\text{aff}}(\check{X})$. This is called the SYZ transform. More precisely, the above constant curvature is expressed locally as

$$dz^t R d\bar{z} \cdot \text{id}, \tag{1}$$

where $z = (z_1, \dots, z_n)^t$ is the local complex coordinates of X , and R is a constant matrix of order n (actually, R is a Hermitian matrix of order n). On the other hand, for a holomorphic vector bundle on X with the Hermitian connection, if its curvature form is expressed locally as the form (1), such a holomorphic vector bundle admits a projectively flat structure (for example, see [11]). Therefore, we see that each object of $\text{Fuk}_{\text{aff}}(\check{X})$ is transformed to a projectively flat bundle on X , which in particular becomes simple. However, there are ambiguities of the choices of transition functions of it. In this paper, we consider a DG-category DG_X consisting of such simple projectively flat bundles with any compatible transition functions. We expect that this DG_X generates $D^b(\text{Coh}(X))$ though we do not discuss it in this paper. At least, it is known that it split-generates $D^b(\text{Coh}(X))$ when X is an abelian variety (cf. [17], [1]). In

this setting, when we fix a choice of transition functions of holomorphic vector bundles in DG_X , we obtain a map

$$\iota : \text{Ob}(Fuk_{\text{aff}}(\check{X})) \rightarrow \text{Ob}(DG_X)$$

by the SYZ transform. Then, for example, it is also shown in [5, Proposition 13.26] that the map ι induces an injection

$$\iota^{isom} : \text{Ob}^{isom}(Fuk_{\text{aff}}(\check{X})) \rightarrow \text{Ob}^{isom}(DG_X),$$

where $\text{Ob}^{isom}(DG_X)$ and $\text{Ob}^{isom}(Fuk_{\text{aff}}(\check{X}))$ denote the set of the isomorphism classes of holomorphic vector bundles in DG_X and the set of the isomorphism classes of objects of $Fuk_{\text{aff}}(\check{X})$, respectively. In the present paper, we prove that the map ι^{isom} is actually a bijection by constructing a natural map

$$F : \text{Ob}(DG_X) \rightarrow \text{Ob}(Fuk_{\text{aff}}(\check{X}))$$

whose direction is opposite to the direction of the map ι .

Of course, though we can also regard this result as the first step to prove the homological mirror symmetry conjecture on (X, \check{X}) , it is a stronger statement in the following sense. In general, for two A_∞ -categories \mathcal{C} and \mathcal{C}' , if there exists an A_∞ -equivalence $\mathcal{C} \cong \mathcal{C}'$, then it is known that there exists an equivalence $Tr(\mathcal{C}) \cong Tr(\mathcal{C}')$ as triangulated categories. Hence, in order to prove the homological mirror symmetry conjecture

$$Tr(DG_X) \cong Tr(Fuk_{\text{aff}}(\check{X}))$$

on (X, \check{X}) , it is enough to prove that there exists an A_∞ -equivalence

$$\mathcal{C}_X \cong \mathcal{C}_{\check{X}}$$

for some full subcategories $\mathcal{C}_X \subset DG_X$, $\mathcal{C}_{\check{X}} \subset Fuk_{\text{aff}}(\check{X})$ which generate the respective triangulated categories $Tr(\mathcal{C}_X) \cong Tr(DG_X)$, $Tr(\mathcal{C}_{\check{X}}) \cong Tr(Fuk_{\text{aff}}(\check{X}))$ (as such categories \mathcal{C}_X and $\mathcal{C}_{\check{X}}$, we would like to take the smallest possible category on each side, see also [13]). Thus, from the viewpoint of the proof of the homological mirror symmetry conjecture on (X, \check{X}) , if we can construct a bijection $\text{Ob}^{isom}(\mathcal{C}_X) \rightarrow \text{Ob}^{isom}(\mathcal{C}_{\check{X}})$ and prove the existence of an A_∞ -equivalence $\mathcal{C}_X \cong \mathcal{C}_{\check{X}}$, then the map ι^{isom} itself need not be bijective. Actually, our original motivation of proving the bijectivity of the map ι^{isom} is that we are interested in the DG-category DG_X itself and the corresponding geometric objects in the mirror dual side. In particular, in [9], exact triangles in $Tr(DG_X)$ consisting of objects of DG_X are studied by using the map F (see also [7]). Although [8] is closely related to [9], we are going to revise [8] by using the main result in the present paper. Furthermore, the map F is also employed in discussions in [10].

This paper is organized as follows. In section 2, we take a complex torus X , and explain the definition of a mirror partner \check{X} of the complex torus X . In section 3, we define a class of a certain kind of simple projectively flat bundles on X , and construct the DG-category DG_X consisting of those holomorphic

vector bundles. We also study the isomorphism classes of them in section 3. In section 4, we consider the Fukaya category $Fuk_{\text{aff}}(\check{X})$ consisting of affine Lagrangian multi sections of $\check{\pi} : \check{X} \rightarrow B$ with unitary local systems along them, and study the isomorphism classes of objects of $Fuk_{\text{aff}}(\check{X})$. In section 5, we explicitly construct the bijection $\text{Ob}^{isom}(DG_X) \rightarrow \text{Ob}^{isom}(Fuk_{\text{aff}}(\check{X}))$. This result is given in Theorem 5.1.

2 Preparations

In this section, we define a complex torus $T_{J=T}^{2n}$ and its mirror partner $\check{T}_{J=T}^{2n}$.

First, we define an n -dimensional complex torus $T_{J=T}^{2n}$ as follows. Let T be a complex matrix of order n such that $\text{Im}T$ is positive definite. We consider the lattice $2\pi(\mathbb{Z}^n \oplus T\mathbb{Z}^n)$ in \mathbb{C}^n and define

$$T_{J=T}^{2n} := \mathbb{C}^n / 2\pi(\mathbb{Z}^n \oplus T\mathbb{Z}^n).$$

Sometimes we regard the n -dimensional complex torus $T_{J=T}^{2n}$ as a $2n$ -dimensional real torus $\mathbb{R}^{2n} / 2\pi\mathbb{Z}^{2n}$. In this paper, we further assume that T is a non-singular matrix. Actually, in our setting described below, it turns out that the mirror partner of the complex torus $T_{J=T}^{2n}$ does not exist if $\det T = 0$. However, we can avoid this problem and discuss the homological mirror symmetry even if $\det T = 0$ by modifying the definition of the mirror partner of the complex torus $T_{J=T}^{2n}$ and the class of holomorphic vector bundles which we treat. This case is discussed in [10]. Here, we fix an $\varepsilon > 0$ small enough and let

$$O_{m_1 \cdots m_n}^{l_1 \cdots l_n} := \left\{ \begin{pmatrix} x \\ y \end{pmatrix} \in T_{J=T}^{2n} \mid \frac{2}{3}\pi(l_j - 1) - \varepsilon < x_j < \frac{2}{3}\pi l_j + \varepsilon, \right. \\ \left. \frac{2}{3}\pi(m_k - 1) - \varepsilon < y_k < \frac{2}{3}\pi m_k + \varepsilon, j, k = 1, \dots, n \right\}$$

be subsets of $T_{J=T}^{2n}$, where $l_j, m_k = 1, 2, 3$,

$$x := (x_1, \dots, x_n)^t, y := (y_1, \dots, y_n)^t,$$

and we identify $x_i \sim x_i + 2\pi$, $y_i \sim y_i + 2\pi$ for each $i = 1, \dots, n$. Sometimes we denote $O_{m_1 \cdots (m_k=m) \cdots m_n}^{l_1 \cdots (l_j=l) \cdots l_n}$ instead of $O_{m_1 \cdots m_n}^{l_1 \cdots l_n}$ in order to specify the values $l_j = l$, $m_k = m$. Then, $\{O_{m_1 \cdots m_n}^{l_1 \cdots l_n}\}_{l_j, m_k=1,2,3}$ is an open cover of $T_{J=T}^{2n}$, and we define the local coordinates of $O_{m_1 \cdots m_n}^{l_1 \cdots l_n}$ by

$$(x_1, \dots, x_n, y_1, \dots, y_n)^t \in \mathbb{R}^{2n}.$$

Furthermore, we locally express the complex coordinates $z := (z_1, \dots, z_n)^t$ of $T_{J=T}^{2n}$ by $z = x + Ty$.

Next, we define a mirror partner $\check{T}_{J=T}^{2n}$ of $T_{J=T}^{2n}$. From the viewpoint of the SYZ construction, we should describe $\check{T}_{J=T}^{2n}$ as the trivial special Lagrangian torus fibration. However, here, instead of constructing it, we consider the

2n-dimensional standard real torus $T^{2n} = \mathbb{R}^{2n}/2\pi\mathbb{Z}^{2n}$ with a modified (non standard) symplectic form. For each point $(x^1, \dots, x^n, y^1, \dots, y^n)^t \in T^{2n}$, we identify $x^i \sim x^i + 2\pi$, $y^i \sim y^i + 2\pi$, where $i = 1, \dots, n$. We also denote by $(x^1, \dots, x^n, y^1, \dots, y^n)^t$ the local coordinates in the neighborhood of an arbitrary point $(x^1, \dots, x^n, y^1, \dots, y^n)^t \in T^{2n}$. Furthermore, we use the same notation $(x^1, \dots, x^n, y^1, \dots, y^n)^t$ when we denote the coordinates of the covering space \mathbb{R}^{2n} of T^{2n} . For simplicity, we set

$$\check{x} := (x^1, \dots, x^n)^t, \check{y} := (y^1, \dots, y^n)^t.$$

We define a complexified symplectic form $\tilde{\omega}$ on T^{2n} by

$$\tilde{\omega} := d\check{x}^t(-T^{-1})^t d\check{y},$$

where $d\check{x} := (dx^1, \dots, dx^n)^t$ and $d\check{y} := (dy^1, \dots, dy^n)^t$. We decompose $\tilde{\omega}$ into

$$\tilde{\omega} = d\check{x}^t \text{Re}(-T^{-1})^t d\check{y} + \mathbf{i} d\check{x}^t \text{Im}(-T^{-1})^t d\check{y},$$

and define

$$\omega := \text{Im}(-T^{-1})^t, \quad B := \text{Re}(-T^{-1})^t.$$

Here, $\mathbf{i} = \sqrt{-1}$. Sometimes we identify the matrices ω and B with the 2-forms $d\check{x}^t \omega d\check{y}$ and $d\check{x}^t B d\check{y}$, respectively. Then, ω gives a symplectic form on T^{2n} . The closed 2-form B is often called the B-field. This complexified symplectic torus $(T^{2n}, \tilde{\omega} = d\check{x}^t(-T^{-1})^t d\check{y})$ is a mirror partner of the complex torus $T_{J=T}^{2n}$. Hereafter, we denote

$$\check{T}_{J=T}^{2n} := (T^{2n}, \tilde{\omega} = d\check{x}^t(-T^{-1})^t d\check{y})$$

for simplicity.

3 Complex geometry side

The purpose of this section is to explain the complex geometry side in the homological mirror symmetry setting on $(T_{J=T}^{2n}, \check{T}_{J=T}^{2n})$. In subsection 3.1, we define a class of holomorphic vector bundles

$$E_{(r,A,r',\mathcal{U},p,q)} \rightarrow T_{J=T}^{2n},$$

and construct a DG-category

$$DG_{T_{J=T}^{2n}}$$

consisting of these holomorphic vector bundles $E_{(r,A,r',\mathcal{U},p,q)}$. In particular, we first construct $E_{(r,A,r',\mathcal{U},p,q)}$ as a complex vector bundle, and then discuss when it becomes a holomorphic vector bundle in Proposition 3.1. In subsection 3.2, we study the isomorphism classes of holomorphic vector bundles $E_{(r,A,r',\mathcal{U},p,q)}$ by using the classification result of simple projectively flat bundles on complex tori by Matsushima [15] and Mukai [16].

3.1 The definition of $E_{(r,A,r',\mathcal{U},p,q)}$

We assume $r \in \mathbb{N}$, $A = (a_{ij}) \in M(n; \mathbb{Z})$, and $p, q \in \mathbb{R}^n$. First, we define $r' \in \mathbb{N}$ by using a given pair $(r, A) \in \mathbb{N} \times M(n; \mathbb{Z})$ as follows. By the theory of elementary divisors, there exist two matrices $\mathcal{A}, \mathcal{B} \in GL(n; \mathbb{Z})$ such that

$$\mathcal{A}\mathcal{A}\mathcal{B} = \begin{pmatrix} \tilde{a}_1 & & & & & & \\ & \ddots & & & & & \\ & & \tilde{a}_s & & & & \\ & & & 0 & & & \\ & & & & \ddots & & \\ & & & & & & 0 \end{pmatrix}, \quad (2)$$

where $\tilde{a}_i \in \mathbb{N}$ ($i = 1, \dots, s, 1 \leq s \leq n$) and $\tilde{a}_i | a_{i+1}$ ($i = 1, \dots, s-1$). Then, we define $r'_i \in \mathbb{N}$ and $a'_i \in \mathbb{Z}$ ($i = 1, \dots, s$) by

$$\frac{\tilde{a}_i}{r} = \frac{a'_i}{r'_i}, \quad \gcd(r'_i, a'_i) = 1,$$

where $\gcd(m, n) > 0$ denotes the greatest common divisor of $m, n \in \mathbb{Z}$. By using these, we set

$$r' := r'_1 \cdots r'_s \in \mathbb{N}. \quad (3)$$

This $r' \in \mathbb{N}$ is uniquely defined by a given pair $(r, A) \in \mathbb{N} \times M(n; \mathbb{Z})$, and it is actually the rank of $E_{(r,A,r',\mathcal{U},p,q)}$. Now, we define the transition functions of $E_{(r,A,r',\mathcal{U},p,q)}$ as follows (although the following notations are complicated, roughly speaking, the transition functions of $E_{(r,A,r',\mathcal{U},p,q)}$ in the cases of $x_j \mapsto x_j + 2\pi$, $y_k \mapsto y_k + 2\pi$ are given by $e^{\frac{i}{r}a_j y} V_j$, U_k , respectively, where $j, k = 1, \dots, n$). Let

$$\psi_{m_1 \cdots m_n}^{l_1 \cdots l_n} : \mathcal{O}_{m_1 \cdots m_n}^{l_1 \cdots l_n} \rightarrow \mathcal{O}_{m_1 \cdots m_n}^{l_1 \cdots l_n} \times \mathbb{C}^{r'}, \quad l_j, m_k = 1, 2, 3$$

be a smooth section of $E_{(r,A,r',\mathcal{U},p,q)}|_{\mathcal{O}_{m_1 \cdots m_n}^{l_1 \cdots l_n}}$. The transition functions of $E_{(r,A,r',\mathcal{U},p,q)}$ are non-trivial on

$$\begin{aligned} & \mathcal{O}_{m_1 \cdots m_n}^{(l_1=3) \cdots l_n} \cap \mathcal{O}_{m_1 \cdots m_n}^{(l_1=1) \cdots l_n}, \quad \mathcal{O}_{m_1 \cdots m_n}^{l_1(l_2=3) \cdots l_n} \cap \mathcal{O}_{m_1 \cdots m_n}^{l_1(l_2=1) \cdots l_n}, \dots, \quad \mathcal{O}_{m_1 \cdots m_n}^{l_1 \cdots (l_n=3)} \cap \mathcal{O}_{m_1 \cdots m_n}^{l_1 \cdots (l_n=1)}, \\ & \mathcal{O}_{(m_1=3) \cdots m_n}^{l_1 \cdots l_n} \cap \mathcal{O}_{(m_1=1) \cdots m_n}^{l_1 \cdots l_n}, \quad \mathcal{O}_{m_1(m_2=3) \cdots m_n}^{l_1 \cdots l_n} \cap \mathcal{O}_{m_1(m_2=1) \cdots m_n}^{l_1 \cdots l_n}, \dots, \\ & \mathcal{O}_{m_1 \cdots (m_n=3)}^{l_1 \cdots l_n} \cap \mathcal{O}_{m_1 \cdots (m_n=1)}^{l_1 \cdots l_n}, \end{aligned}$$

and otherwise are trivial. We define the transition function on $\mathcal{O}_{m_1 \cdots m_n}^{l_1 \cdots (l_j=3) \cdots l_n} \cap \mathcal{O}_{m_1 \cdots m_n}^{l_1 \cdots (l_j=1) \cdots l_n}$ by

$$\begin{aligned} & \psi_{m_1 \cdots m_n}^{l_1 \cdots (l_j=3) \cdots l_n} \Big|_{\mathcal{O}_{m_1 \cdots m_n}^{l_1 \cdots (l_j=3) \cdots l_n} \cap \mathcal{O}_{m_1 \cdots m_n}^{l_1 \cdots (l_j=1) \cdots l_n}} \\ & = e^{\frac{i}{r} a_j y} V_j \psi_{m_1 \cdots m_n}^{l_1 \cdots (l_j=1) \cdots l_n} \Big|_{\mathcal{O}_{m_1 \cdots m_n}^{l_1 \cdots (l_j=3) \cdots l_n} \cap \mathcal{O}_{m_1 \cdots m_n}^{l_1 \cdots (l_j=1) \cdots l_n}}, \end{aligned}$$

where $a_j := (a_{1j}, \dots, a_{nj}) \in \mathbb{Z}^n$ and $V_j \in U(r')$. Similarly, we define the transition function on $O_{m_1 \dots (m_k=3) \dots m_n}^{l_1 \dots l_n} \cap O_{m_1 \dots (m_k=1) \dots m_n}^{l_1 \dots l_n}$ by

$$\begin{aligned} & \psi_{m_1 \dots (m_k=3) \dots m_n}^{l_1 \dots l_n} \Big|_{O_{m_1 \dots (m_k=3) \dots m_n}^{l_1 \dots l_n} \cap O_{m_1 \dots (m_k=1) \dots m_n}^{l_1 \dots l_n}} \\ &= U_k \psi_{m_1 \dots (m_k=1) \dots m_n}^{l_1 \dots l_n} \Big|_{O_{m_1 \dots (m_k=3) \dots m_n}^{l_1 \dots l_n} \cap O_{m_1 \dots (m_k=1) \dots m_n}^{l_1 \dots l_n}}, \end{aligned}$$

where $U_k \in U(r')$. In the definitions of these transition functions, actually, we only treat $V_j, U_k \in U(r')$ which satisfy the cocycle condition, so we explain the cocycle condition below. When we define

$$\begin{aligned} & \psi_{m_1 \dots (m_k=3) \dots m_n}^{l_1 \dots (l_j=3) \dots l_n} \Big|_{O_{m_1 \dots (m_k=3) \dots m_n}^{l_1 \dots (l_j=3) \dots l_n} \cap O_{m_1 \dots (m_k=1) \dots m_n}^{l_1 \dots (l_j=1) \dots l_n}} \\ &= U_k \psi_{m_1 \dots (m_k=1) \dots m_n}^{l_1 \dots (l_j=3) \dots l_n} \Big|_{O_{m_1 \dots (m_k=3) \dots m_n}^{l_1 \dots (l_j=3) \dots l_n} \cap O_{m_1 \dots (m_k=1) \dots m_n}^{l_1 \dots (l_j=1) \dots l_n}} \\ &= (U_k) \left(e^{\frac{i}{r} a_j y} V_j \right) \psi_{m_1 \dots (m_k=1) \dots m_n}^{l_1 \dots (l_j=1) \dots l_n} \Big|_{O_{m_1 \dots (m_k=3) \dots m_n}^{l_1 \dots (l_j=3) \dots l_n} \cap O_{m_1 \dots (m_k=1) \dots m_n}^{l_1 \dots (l_j=1) \dots l_n}}, \end{aligned}$$

the cocycle condition is expressed as

$$V_j V_k = V_k V_j, \quad U_j U_k = U_k U_j, \quad \zeta^{-a_{kj}} U_k V_j = V_j U_k,$$

where ζ is the r -th root of 1 and $j, k = 1, \dots, n$. We define a set \mathcal{U} of unitary matrices by

$$\mathcal{U} := \left\{ V_j, U_k \in U(r') \mid V_j V_k = V_k V_j, \quad U_j U_k = U_k U_j, \quad \zeta^{-a_{kj}} U_k V_j = V_j U_k, \right. \\ \left. j, k = 1, \dots, n \right\}.$$

Of course, how to define the set \mathcal{U} relates closely to (in)decomposability of $E_{(r,A,r',\mathcal{U},p,q)}$. Here, we only treat the set \mathcal{U} such that $E_{(r,A,r',\mathcal{U},p,q)}$ is simple. Actually, we can take such a set $\mathcal{U} \neq \emptyset$ for any $(r, A, r') \in \mathbb{N} \times M(n; \mathbb{Z}) \times \mathbb{N}$, and this fact is discussed in Proposition 3.2. Furthermore, we define a connection $\nabla_{(r,A,r',\mathcal{U},p,q)}$ on $E_{(r,A,r',\mathcal{U},p,q)}$ locally as

$$\nabla_{(r,A,r',\mathcal{U},p,q)} := d - \frac{\mathbf{i}}{2\pi} \left(\left(\frac{1}{r} x^t A^t + \frac{1}{r} p^t \right) + \frac{1}{r} q^t T \right) dy \cdot I_{r'},$$

where $dy := (dy_1, \dots, dy_n)^t$ and d denotes the exterior derivative. In fact, $\nabla_{(r,A,r',\mathcal{U},p,q)}$ is compatible with the transition functions and so defines a global connection. Then, its curvature form $\Omega_{(r,A,r',\mathcal{U},p,q)}$ is expressed locally as

$$\Omega_{(r,A,r',\mathcal{U},p,q)} = -\frac{\mathbf{i}}{2\pi r} dx^t A^t dy \cdot I_{r'}, \quad (4)$$

where $dx := (dx_1, \dots, dx_n)^t$. In particular, this local expression (4) implies that holomorphic vector bundles $E_{(r,A,r',\mathcal{U},p,q)}$ are simple projectively flat bundles

(for example, the definition of projectively flat bundles is written in [11]). Moreover, the interpretation for these simple projectively flat bundles $E_{(r,A,r',\mathcal{U},p,q)}$ by using factors of automorphy is given in section 3 of [8]. Now, we consider the condition such that $E_{(r,A,r',\mathcal{U},p,q)}$ is holomorphic. We see that the following proposition holds.

Proposition 3.1. *For a given quadruple $(r, A, p, q) \in \mathbb{N} \times M(n; \mathbb{Z}) \times \mathbb{R}^n \times \mathbb{R}^n$, the complex vector bundle $E_{(r,A,r',\mathcal{U},p,q)} \rightarrow T_{J=T}^{2n}$ is holomorphic if and only if $AT = (AT)^t$ holds.*

Proof. A complex vector bundle is holomorphic if and only if the (0,2)-part of its curvature form vanishes, so we calculate the (0,2)-part of $\Omega_{(r,A,\mu,\mathcal{U})}$. It turns out to be

$$\Omega_{(r,A,r',\mathcal{U},p,q)}^{(0,2)} = \frac{\mathbf{i}}{2\pi r} d\bar{z}^t \{T(T - \bar{T})^{-1}\}^t A^t (T - \bar{T})^{-1} d\bar{z} \cdot I_{r'},$$

where $d\bar{z} := (d\bar{z}_1, \dots, d\bar{z}_n)^t$. Thus, $\Omega_{(r,A,r',\mathcal{U},p,q)}^{(0,2)} = 0$ is equivalent to that $\{T(T - \bar{T})^{-1}\}^t A^t (T - \bar{T})^{-1}$ is a symmetric matrix, i.e., $AT = (AT)^t$. \square

Concerning the above discussions, here, we mention the simplicity of holomorphic vector bundles $E_{(r,A,r',\mathcal{U},p,q)}$. The following proposition holds.

Proposition 3.2. *For each quadruple $(r, A, p, q) \in \mathbb{N} \times M(n; \mathbb{Z}) \times \mathbb{R}^n \times \mathbb{R}^n$, we can take a set $\mathcal{U} \neq \emptyset$ such that $E_{(r,A,r',\mathcal{U},p,q)}$ is simple.*

Proof. For a given pair $(r, A) \in \mathbb{N} \times M(n; \mathbb{Z})$, we can take two matrices $\mathcal{A}, \mathcal{B} \in GL(n; \mathbb{Z})$ which satisfy the relation (2). Then, note that $r' := r'_1 \cdots r'_s \in \mathbb{N}$ is uniquely defined in the sense of the relation (3). We fix such matrices $\mathcal{A}, \mathcal{B} \in GL(n; \mathbb{Z})$, and set

$$T' := \mathcal{B}^{-1} T \mathcal{A}^t.$$

By using this T' , we can consider the complex torus $T_{J=T'}^{2n} = \mathbb{C}^n / 2\pi(\mathbb{Z}^n \oplus T' \mathbb{Z}^n)$, and we locally express the complex coordinates $Z := (Z_1, \dots, Z_n)^t$ of $T_{J=T'}^{2n}$ by $Z = X + T'Y$, where $X := (X_1, \dots, X_n)^t$, $Y := (Y_1, \dots, Y_n)^t$. In particular, the complex torus $T_{J=T'}^{2n}$ is biholomorphic to the complex torus $T_{J=T}^{2n}$, and the biholomorphic map

$$\varphi : T_{J=T}^{2n} \xrightarrow{\sim} T_{J=T'}^{2n}$$

is actually given by

$$\varphi(z) = \mathcal{B}^{-1} z.$$

Furthermore, when we regard the complex manifolds $T_{J=T}^{2n}$ and $T_{J=T'}^{2n}$ as the real differentiable manifolds $\mathbb{R}^{2n} / 2\pi\mathbb{Z}^{2n}$, the biholomorphic map φ is regarded as the diffeomorphism

$$\begin{pmatrix} X \\ Y \end{pmatrix} = \varphi \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} \mathcal{B}^{-1} & O \\ O & (\mathcal{A}^{-1})^t \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}.$$

Now, we define a set $\mathcal{U}' \neq \emptyset$ as follows. First, we set

$$\tilde{V}_i := \begin{pmatrix} 0 & 1 & & \\ & & \ddots & \ddots \\ & & & 1 \\ 1 & & & 0 \end{pmatrix} \in U(r'_i), \quad \tilde{U}_i := \begin{pmatrix} 1 & & & \\ & \zeta_i & & \\ & & \ddots & \\ & & & \zeta_i^{r'_i-1} \end{pmatrix} \in U(r'_i),$$

where $\zeta_i := e^{\frac{2\pi i}{r'_i}}$, $i = 1, \dots, s$. By using these matrices \tilde{V}_i, \tilde{U}_i , we define

$$\begin{aligned} V'_1 &:= \tilde{V}_1 \otimes I_{r'_2} \otimes \cdots \otimes I_{r'_s} \in U(r'), \\ V'_2 &:= I_{r'_1} \otimes \tilde{V}_2 \otimes \cdots \otimes I_{r'_s} \in U(r'), \\ &\vdots \\ V'_s &:= I_{r'_1} \otimes \cdots \otimes I_{r'_{s-1}} \otimes \tilde{V}_s \in U(r'), \\ V'_{s+1} &:= I_{r'_1} \otimes \cdots \otimes I_{r'_s} \in U(r'), \\ &\vdots \\ V'_n &:= I_{r'_1} \otimes \cdots \otimes I_{r'_s} \in U(r'), \\ U'_1 &:= \tilde{U}_1^{-a'_1} \otimes I_{r'_2} \otimes \cdots \otimes I_{r'_s} \in U(r'), \\ U'_2 &:= I_{r'_1} \otimes \tilde{U}_2^{-a'_2} \otimes \cdots \otimes I_{r'_s} \in U(r'), \\ &\vdots \\ U'_s &:= I_{r'_1} \otimes \cdots \otimes I_{r'_{s-1}} \otimes \tilde{U}_s^{-a'_s} \in U(r'), \\ U'_{s+1} &:= I_{r'_1} \otimes \cdots \otimes I_{r'_s} \in U(r'), \\ &\vdots \\ U'_n &:= I_{r'_1} \otimes \cdots \otimes I_{r'_s} \in U(r'), \end{aligned}$$

and set

$$\mathcal{U}' := \left\{ V'_1, V'_2, \dots, V'_s, V'_{s+1}, \dots, V'_n, U'_1, U'_2, \dots, U'_s, U'_{s+1}, \dots, U'_n \in U(r') \right\} \neq \emptyset.$$

Then, we can construct the holomorphic vector bundle

$$E_{(r, \mathcal{AAB}, r', \mathcal{U}', \mathcal{A}_p, \mathcal{B}^t q)} \rightarrow T_{J=T'}^{2n},$$

and in particular, V'_j and U'_k are used in the definition of the transition functions of $E_{(r, \mathcal{AAB}, r', \mathcal{U}', \mathcal{A}_p, \mathcal{B}^t q)}$ in the X_j and Y_k directions, respectively ($j, k = 1, \dots, n$). For this holomorphic vector bundle $E_{(r, \mathcal{AAB}, r', \mathcal{U}', \mathcal{A}_p, \mathcal{B}^t q)}$, we can consider the pullback bundle $\varphi^* E_{(r, \mathcal{AAB}, r', \mathcal{U}', \mathcal{A}_p, \mathcal{B}^t q)}$ by the biholomorphic map φ , and we can regard the pullback bundle $\varphi^* E_{(r, \mathcal{AAB}, r', \mathcal{U}', \mathcal{A}_p, \mathcal{B}^t q)}$ as the holomorphic vector bundle

$$E_{(r, A, r', \mathcal{U}, p, q)} \rightarrow T_{J=T}^{2n}$$

by using a suitable set \mathcal{U} which is defined by employing the data $(\mathcal{U}', \mathcal{A}, \mathcal{B})$. In particular, since we can check $\mathcal{U} \neq \emptyset$ easily, in order to prove the statement of this proposition, it is enough to prove that $E_{(r, \mathcal{A}AB, r', \mathcal{U}', \mathcal{A}p, \mathcal{B}^t q)}$ is simple.

Let

$$\varphi^{r'_1 \cdots r'_s}$$

be an element in $\text{End}(E_{(r, \mathcal{A}AB, r', \mathcal{U}', \mathcal{A}p, \mathcal{B}^t q)})$. Since the rank of $E_{(r, \mathcal{A}AB, r', \mathcal{U}', \mathcal{A}p, \mathcal{B}^t q)}$ is $r' = r'_1 \cdots r'_s$, we can treat $\varphi^{r'_1 \cdots r'_s}$ as a matrix of order r' . Then, we can divide $\varphi^{r'_1 \cdots r'_s}$ as follows.

$$\varphi^{r'_1 \cdots r'_s} = \begin{pmatrix} \varphi_{11}^{r'_2 \cdots r'_s} & \cdots & \varphi_{1r'_1}^{r'_2 \cdots r'_s} \\ \vdots & \ddots & \vdots \\ \varphi_{r'_1 1}^{r'_2 \cdots r'_s} & \cdots & \varphi_{r'_1 r'_1}^{r'_2 \cdots r'_s} \end{pmatrix} = \left(\varphi_{k_1 l_1}^{r'_2 \cdots r'_s} \right)_{1 \leq k_1, l_1 \leq r'_1}.$$

Here, each $\varphi_{k_1 l_1}^{r'_2 \cdots r'_s}$ is a matrix of order $r'_2 \cdots r'_s$. Similarly, we can divide each $\varphi_{k_1 l_1}^{r'_2 \cdots r'_s}$ as follows.

$$\varphi_{k_1 l_1}^{r'_2 \cdots r'_s} = \begin{pmatrix} \left(\varphi_{k_1 l_1}^{r'_2 \cdots r'_s} \right)_{11}^{r'_3 \cdots r'_s} & \cdots & \left(\varphi_{k_1 l_1}^{r'_2 \cdots r'_s} \right)_{1r'_2}^{r'_3 \cdots r'_s} \\ \vdots & \ddots & \vdots \\ \left(\varphi_{k_1 l_1}^{r'_2 \cdots r'_s} \right)_{r'_2 1}^{r'_3 \cdots r'_s} & \cdots & \left(\varphi_{k_1 l_1}^{r'_2 \cdots r'_s} \right)_{r'_2 r'_2}^{r'_3 \cdots r'_s} \end{pmatrix} = \left(\left(\varphi_{k_1 l_1}^{r'_2 \cdots r'_s} \right)_{k_2 l_2}^{r'_3 \cdots r'_s} \right)_{1 \leq k_2, l_2 \leq r'_2}.$$

Here, each $\left(\varphi_{k_1 l_1}^{r'_2 \cdots r'_s} \right)_{k_2 l_2}^{r'_3 \cdots r'_s}$ is matrix of order $r'_3 \cdots r'_s$. By repeating the above steps, as a result, we can express $\varphi^{r'_1 \cdots r'_s}$ as

$$\left(\left(\left(\left(\left(\left(\varphi_{k_1 l_1}^{r'_2 \cdots r'_s} \right)_{k_2 l_2}^{r'_3 \cdots r'_s} \right) \cdots \right)_{k_{s-1} l_{s-1}}^{r'_s} \right) \right)_{k_s l_s} \right),$$

where $1 \leq k_i, l_i \leq r'_i$ ($i = 1, \dots, s$). Hereafter, we consider the local expression of each component

$$\left(\left(\left(\left(\left(\varphi_{k_1 l_1}^{r'_2 \cdots r'_s} \right)_{k_2 l_2}^{r'_3 \cdots r'_s} \right) \cdots \right)_{k_{s-1} l_{s-1}}^{r'_s} \right)_{k_s l_s} \right) \quad (5)$$

of $\varphi^{r'_1 \cdots r'_s}$. First, for each $k = 1, \dots, n$, by considering the transition functions of $E_{(r, \mathcal{A}AB, r', \mathcal{U}', \mathcal{A}p, \mathcal{B}^t q)}$ in the Y_k direction, we see that the morphism $\varphi^{r'_1 \cdots r'_s}$ must satisfy

$$\begin{aligned} & U'_k \cdot \varphi^{r'_1 \cdots r'_s}(X_1, \dots, X_n, Y_1, \dots, Y_n) \\ &= \varphi^{r'_1 \cdots r'_s}(X_1, \dots, X_n, Y_1, \dots, Y_k + 2\pi, \dots, Y_n) \cdot U'_k. \end{aligned} \quad (6)$$

Furthermore, the morphism $\varphi^{r'_1 \cdots r'_s}$ need to satisfy not only the relation (6) but also the Cauchy-Riemann equation

$$\bar{\partial}(\varphi^{r'_1 \cdots r'_s}) = 0. \quad (7)$$

Therefore, by the relations (6) and (7), we can give a local expression of the component (5) of $\varphi^{r'_1 \cdots r'_s}$ as follows.

$$\sum_{I_{k_1 l_1}^1, \dots, I_{k_s l_s}^s, I^{s+1}, \dots, I^n \in \mathbb{Z}} \lambda_{I_{k_1 l_1}^1, \dots, I_{k_s l_s}^s, I^{s+1}, \dots, I^n}^{k_1 l_1, \dots, k_s l_s} e^{\mathbf{i} \left(\frac{a'_1}{r'_1} (l_1 - k_1) + I_{k_1 l_1}^1, \dots, \frac{a'_s}{r'_s} (l_s - k_s) + I_{k_s l_s}^s, I^{s+1}, \dots, I^n \right)} T^{-1}(X + TY).$$

Here,

$$\lambda_{I_{k_1 l_1}^1, \dots, I_{k_s l_s}^s, I^{s+1}, \dots, I^n}^{k_1 l_1, \dots, k_s l_s} \in \mathbb{C}$$

is an arbitrary constant. Finally, for each $j = 1, \dots, n$, we consider the conditions on the transition functions of $E_{(r, AAB, r', U', Ap, B^t q)}$ in the X_j direction. By a direct calculation, if $j = 1, \dots, s$, we obtain

$$\begin{aligned} & \left(\cdots \left(\varphi_{k_1 l_1}^{r'_2 \cdots r'_s} \right) \cdots \right)_{k_j l_j}^{r'_{j+1} \cdots r'_s} (X_1, \dots, X_j + 2\pi, \dots, X_n, Y_1, \dots, Y_n) \\ &= \left(\cdots \left(\varphi_{k_1 l_1}^{r'_2 \cdots r'_s} \right) \cdots \right)_{(k_j+1)(l_j+1)}^{r'_{j+1} \cdots r'_s} (X_1, \dots, X_n, Y_1, \dots, Y_n), \end{aligned} \quad (8)$$

and if $j = s + 1, \dots, n$, we obtain

$$\begin{aligned} & \left(\cdots \left(\varphi_{k_1 l_1}^{r'_2 \cdots r'_s} \right) \cdots \right)_{k_s l_s} (X_1, \dots, X_j + 2\pi, \dots, X_n, Y_1, \dots, Y_n) \\ &= \left(\cdots \left(\varphi_{k_1 l_1}^{r'_2 \cdots r'_s} \right) \cdots \right)_{k_s l_s} (X_1, \dots, X_n, Y_1, \dots, Y_n). \end{aligned} \quad (9)$$

In particular, the relation (8) implies

$$\begin{aligned} & \left(\cdots \left(\varphi_{k_1 l_1}^{r'_2 \cdots r'_s} \right) \cdots \right)_{k_s l_s} (X_1, \dots, X_j + 2\pi r'_j, \dots, X_n, Y_1, \dots, Y_n) \\ &= \left(\cdots \left(\varphi_{k_1 l_1}^{r'_2 \cdots r'_s} \right) \cdots \right)_{k_s l_s} (X_1, \dots, X_n, Y_1, \dots, Y_n), \end{aligned} \quad (10)$$

and by using the relations (9) and (10), we have the condition

$$\begin{aligned} & \left(\frac{a'_1}{r'_1} (l_1 - k_1) + I_{k_1 l_1}^1, \dots, \frac{a'_s}{r'_s} (l_s - k_s) + I_{k_s l_s}^s, I^{s+1}, \dots, I^n \right) T'^{-1} \\ & \in \frac{\mathbb{Z}}{r'_1} \times \cdots \times \frac{\mathbb{Z}}{r'_s} \times \mathbb{Z} \times \cdots \times \mathbb{Z} \subset \mathbb{R}^n. \end{aligned} \quad (11)$$

Now, since $\text{Im}T'^{-1}$ is positive definite, the condition (11) turns out to be

$$\left(\frac{a'_1}{r'_1} (l_1 - k_1) + I_{k_1 l_1}^1, \dots, \frac{a'_s}{r'_s} (l_s - k_s) + I_{k_s l_s}^s, I^{s+1}, \dots, I^n \right) = 0. \quad (12)$$

Here, recall the relation (8). By the relation (8), it is enough to consider the condition (12) in the case $k_1 = \dots = k_s = 1$. We focus on the first component in the condition (12), i.e.,

$$\frac{a'_1}{r'_1}(l_1 - 1) + I_{1l_1}^1 = 0. \quad (13)$$

In the case $l_1 = 1$, the condition (13) turns out to be $I_{11}^1 = 0$, so by the relation (8), we see

$$\varphi_{11}^{r'_2 \cdots r'_s} = \varphi_{22}^{r'_2 \cdots r'_s} = \dots = \varphi_{r'_1 r'_1}^{r'_2 \cdots r'_s} \in M(r'_2 \cdots r'_s; \mathbb{C}).$$

We consider the cases $l_1 = 2, \dots, r'_1$. Note that $\gcd(r'_1, a'_1) = 1$ holds by the assumption. Therefore, we have

$$l_1 - 1 \in r'_1 \mathbb{Z}$$

by the condition (13). However, this fact contradicts the assumption $l_1 = 2, \dots, r'_1$. Thus, for each $l_1 = 2, \dots, r'_1$, we obtain

$$\varphi_{1l_1}^{r'_2 \cdots r'_s} = O,$$

and by using the relation (8) again, we also obtain

$$\varphi_{2(l_1+1)}^{r'_2 \cdots r'_s} = \varphi_{3(l_1+2)}^{r'_2 \cdots r'_s} = \dots = \varphi_{r'_1(l_1-1)}^{r'_2 \cdots r'_s} = O.$$

Similarly, by focusing on the second component in the condition (12), we see

$$\left(\varphi_{11}^{r'_2 \cdots r'_s} \right)_{11}^{r'_3 \cdots r'_s} = \left(\varphi_{11}^{r'_2 \cdots r'_s} \right)_{22}^{r'_3 \cdots r'_s} = \dots = \left(\varphi_{11}^{r'_2 \cdots r'_s} \right)_{r'_2 r'_2}^{r'_3 \cdots r'_s} \in M(r'_3 \cdots r'_s; \mathbb{C}),$$

and the other components $\left(\varphi_{11}^{r'_2 \cdots r'_s} \right)_{k_2 l_2}^{r'_3 \cdots r'_s}$ of the matrix $\varphi_{11}^{r'_2 \cdots r'_s}$ vanish. By repeating the above discussions, as a result, we have

$$\varphi_{r'_1}^{r'_2 \cdots r'_s} = \left(\left(\dots \left(\left(\varphi_{11}^{r'_2 \cdots r'_s} \right)_{11}^{r'_3 \cdots r'_s} \right) \dots \right)_{11}^{r'_s} \right)_{11} \cdot I_{r'_1},$$

where

$$\left(\left(\dots \left(\left(\varphi_{11}^{r'_2 \cdots r'_s} \right)_{11}^{r'_3 \cdots r'_s} \right) \dots \right)_{11}^{r'_s} \right)_{11} \in \mathbb{C}.$$

Thus, we can conclude

$$\text{End}(E_{(r, AAB, r', U', Ap, B^t q)}) \cong \mathbb{C}.$$

□

We define a DG-category

$$DG_{T_{J=T}^{2n}}$$

consisting of holomorphic vector bundles $(E_{(r,A,r',\mathcal{U},p,q)}, \nabla_{(r,A,r',\mathcal{U},p,q)})$. This definition is an extension of the case of a pair $(T_{J=T}^2, \bar{T}_{J=T}^2)$ of elliptic curves in [7] (see section 3) to the higher dimensional case, and it is also written in section 2 of [8]. The objects of $DG_{T_{J=T}^{2n}}$ are holomorphic vector bundles $E_{(r,A,r',\mathcal{U},p,q)}$ with $U(r')$ -connections $\nabla_{(r,A,r',\mathcal{U},p,q)}$. Of course, we assume $AT = (AT)^t$. Sometimes we simply denote $(E_{(r,A,\mu,\mathcal{U})}, \nabla_{(r,A,\mu,\mathcal{U})})$ by $E_{(r,A,r',\mathcal{U},p,q)}$. For any two objects

$$\begin{aligned} E_{(r,A,r',\mathcal{U},p,q)} &= (E_{(r,A,r',\mathcal{U},p,q)}, \nabla_{(r,A,r',\mathcal{U},p,q)}), \\ E_{(s,B,s',\mathcal{V},u,v)} &= (E_{(s,B,s',\mathcal{V},u,v)}, \nabla_{(s,B,s',\mathcal{V},u,v)}), \end{aligned}$$

the space of morphisms is defined by

$$\begin{aligned} &\text{Hom}_{DG_{T_{J=T}^{2n}}}(E_{(r,A,r',\mathcal{U},p,q)}, E_{(s,B,s',\mathcal{V},u,v)}) \\ &:= \Gamma(E_{(r,A,r',\mathcal{U},p,q)}, E_{(s,B,s',\mathcal{V},u,v)}) \otimes_{C^\infty(T_{J=T}^{2n})} \Omega^{0,*}(T_{J=T}^{2n}), \end{aligned}$$

where $\Omega^{0,*}(T_{J=T}^{2n})$ is the space of anti-holomorphic differential forms, and

$$\Gamma(E_{(r,A,r',\mathcal{U},p,q)}, E_{(s,B,s',\mathcal{V},u,v)})$$

is the space of homomorphisms from $E_{(r,A,r',\mathcal{U},p,q)}$ to $E_{(s,B,s',\mathcal{V},u,v)}$. The space of morphisms $\text{Hom}_{DG_{T_{J=T}^{2n}}}(E_{(r,A,r',\mathcal{U},p,q)}, E_{(s,B,s',\mathcal{V},u,v)})$ is a \mathbb{Z} -graded vector space, where the grading is defined as the degree of the anti-holomorphic differential forms. In particular, the degree r part is denoted

$$\text{Hom}_{DG_{T_{J=T}^{2n}}}^r(E_{(r,A,r',\mathcal{U},p,q)}, E_{(s,B,s',\mathcal{V},u,v)}).$$

We decompose $\nabla_{(r,A,r',\mathcal{U},p,q)}$ into its holomorphic part and anti-holomorphic part $\nabla_{(r,A,r',\mathcal{U},p,q)} = \nabla_{(r,A,r',\mathcal{U},p,q)}^{(1,0)} + \nabla_{(r,A,r',\mathcal{U},p,q)}^{(0,1)}$, and define a linear map

$$\text{Hom}_{DG_{T_{J=T}^{2n}}}^r(E_{(r,A,r',\mathcal{U},p,q)}, E_{(s,B,s',\mathcal{V},u,v)}) \rightarrow \text{Hom}_{DG_{T_{J=T}^{2n}}}^{r+1}(E_{(r,A,r',\mathcal{U},p,q)}, E_{(s,B,s',\mathcal{V},u,v)})$$

by

$$\psi \mapsto (2\nabla_{(s,B,s',\mathcal{V},u,v)}^{(0,1)})(\psi) - (-1)^r \psi (2\nabla_{(r,A,r',\mathcal{U},p,q)}^{(0,1)}).$$

We can check that this linear map is a differential. Furthermore, the product structure is defined by the composition of homomorphisms of vector bundles together with the wedge product for the anti-holomorphic differential forms. Then, these differential and product structure satisfy the Leibniz rule. Thus, $DG_{T_{J=T}^{2n}}$ forms a DG-category.

3.2 The isomorphism classes of $E_{(r,A,r',\mathcal{U},p,q)}$

In this subsection, we fix $r \in \mathbb{N}$, $A \in M(n; \mathbb{Z})$, and consider the condition such that $E_{(r,A,r',\mathcal{U},p,q)} \cong E_{(r,A,r',\mathcal{U}',p',q')}$ holds. Here,

$$p, q, p', q' \in \mathbb{R}^n$$

and

$$\mathcal{U} := \left\{ V_j, U_k \in U(r') \mid V_j V_k = V_k V_j, U_j U_k = U_k U_j, \zeta^{-a_{kj}} U_k V_j = V_j U_k, \right. \\ \left. j, k = 1, \dots, n \right\},$$

$$\mathcal{U}' := \left\{ V'_j, U'_k \in U(r') \mid V'_j V'_k = V'_k V'_j, U'_j U'_k = U'_k U'_j, \zeta^{-a_{kj}} U'_k V'_j = V'_j U'_k, \right. \\ \left. j, k = 1, \dots, n \right\}.$$

Furthermore, for each $j = 1, \dots, n$, we define $\xi_j, \theta_j, \xi'_j, \theta'_j \in \mathbb{R}$ by

$$e^{i\xi_j} = \det V_j, \quad e^{i\theta_j} = \det U_j, \quad e^{i\xi'_j} = \det V'_j, \quad e^{i\theta'_j} = \det U'_j,$$

and set

$$\xi := (\xi_1, \dots, \xi_n)^t, \quad \theta := (\theta_1, \dots, \theta_n)^t, \quad \xi' := (\xi'_1, \dots, \xi'_n)^t, \quad \theta' := (\theta'_1, \dots, \theta'_n)^t \in \mathbb{R}^n. \quad (14)$$

Now, in order to consider the condition such that $E_{(r,A,r',\mathcal{U},p,q)} \cong E_{(r,A,r',\mathcal{U}',p',q')}$ holds, we recall the following classification result of simple projectively flat bundles on complex tori by Matsushima and Mukai (see [15, Theorem 6.1], [16, Proposition 6.17 (1)], and note that the notion of semi-homogeneous vector bundles in [16] is equivalent to the notion of projectively flat bundles).

Theorem 3.3 (Matsushima [15], Mukai [16]). *For two simple projectively flat bundles E, E' over a complex torus \mathbb{C}^n/Γ ($\Gamma \subset \mathbb{C}^n$ is a lattice) which satisfy $(\text{rank} E, c_1(E)) = (\text{rank} E', c_1(E'))$, there exists a line bundle $L \in \text{Pic}^0(\mathbb{C}^n/\Gamma)$ such that*

$$E' \cong E \otimes L.$$

By using Theorem 3.3, we obtain the following theorem.

Theorem 3.4. *Two holomorphic vector bundles $E_{(r,A,r',\mathcal{U},p,q)}$, $E_{(r,A,r',\mathcal{U}',p',q')}$ are isomorphic to each other,*

$$E_{(r,A,r',\mathcal{U},p,q)} \cong E_{(r,A,r',\mathcal{U}',p',q')},$$

if and only if

$$(p+T^t q) - (p'+T^t q') \equiv \frac{r}{r'}(\theta - \theta') + T^t \frac{r}{r'}(\xi' - \xi) \pmod{2\pi r \left(\mathcal{A}^{-1} \begin{pmatrix} \mathbb{Z} \\ r'_1 \\ \vdots \\ \mathbb{Z} \\ r'_s \\ \mathbb{Z} \\ \vdots \\ \mathbb{Z} \end{pmatrix} \oplus T^t (\mathcal{B}^{-1})^t \begin{pmatrix} \mathbb{Z} \\ r'_1 \\ \vdots \\ \mathbb{Z} \\ r'_s \\ \mathbb{Z} \\ \vdots \\ \mathbb{Z} \end{pmatrix} \right)}$$

holds.

Proof. In order to prove the statement of this theorem, we again take the biholomorphic map

$$\varphi : T_{J=T}^{2n} \xrightarrow{\sim} T_{J=T'}^{2n}$$

in the proof of Proposition 3.2. Of course, we can also regard this biholomorphic map φ as the diffeomorphism $\mathbb{R}^{2n}/2\pi\mathbb{Z}^{2n} \xrightarrow{\sim} \mathbb{R}^{2n}/2\pi\mathbb{Z}^{2n}$ which is expressed locally as

$$\varphi \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} \mathcal{B}^{-1} & O \\ O & (\mathcal{A}^{-1})^t \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}.$$

Then, by the biholomorphicity of the map φ ,

$$E_{(r,A,r',\mathcal{U},p,q)} \cong E_{(r,A,r',\mathcal{U}',p',q')}$$

holds if and only if

$$(\varphi^{-1})^* E_{(r,A,r',\mathcal{U},p,q)} \cong (\varphi^{-1})^* E_{(r,A,r',\mathcal{U}',p',q')}$$

holds, so we consider the condition such that $(\varphi^{-1})^* E_{(r,A,r',\mathcal{U},p,q)} \cong (\varphi^{-1})^* E_{(r,A,r',\mathcal{U}',p',q')}$ holds. Now, by using the suitable sets $\tilde{\mathcal{U}}$ and $\tilde{\mathcal{U}}'$ (the definitions of $\tilde{\mathcal{U}}$ and $\tilde{\mathcal{U}}'$ depend on the data $(\mathcal{U}, \mathcal{A}, \mathcal{B})$ and $(\mathcal{U}', \mathcal{A}, \mathcal{B})$, respectively), we can regard $(\varphi^{-1})^* E_{(r,A,r',\mathcal{U},p,q)}$ and $(\varphi^{-1})^* E_{(r,A,r',\mathcal{U}',p',q')}$ as $E_{(r,\tilde{A},r',\tilde{\mathcal{U}},\tilde{p},\tilde{q})}$ and $E_{(r,\tilde{A},r',\tilde{\mathcal{U}}',\tilde{p}',\tilde{q}')}$, respectively, where

$$\tilde{A} := \mathcal{A}\mathcal{A}\mathcal{B} = \begin{pmatrix} \tilde{a}_1 & & & & & \\ & \ddots & & & & \\ & & \tilde{a}_s & & & \\ & & & 0 & & \\ & & & & \ddots & \\ & & & & & 0 \end{pmatrix}, \quad \tilde{p} := \mathcal{A}p, \quad \tilde{q} := \mathcal{B}^t q, \quad \tilde{p}' := \mathcal{A}p', \quad \tilde{q}' := \mathcal{B}^t q'.$$

Then, constant vectors $\xi, \theta, \xi', \theta' \in \mathbb{R}^n$ are also transformed to constant vectors

$$\tilde{\xi} := \mathcal{B}^t \xi, \quad \tilde{\theta} := \mathcal{A}\theta, \quad \tilde{\xi}' := \mathcal{B}^t \xi', \quad \tilde{\theta}' := \mathcal{A}\theta' \in \mathbb{R}^n,$$

respectively. For these holomorphic vector bundles $E_{(r,\tilde{A},r',\tilde{\mathcal{U}},\tilde{p},\tilde{q})}$ and $E_{(r,\tilde{A},r',\tilde{\mathcal{U}}',\tilde{p}',\tilde{q}')}$, by Theorem 3.3, we see that there exists a holomorphic line bundle $E_{(1,O,1,\mathcal{V},u,v)}$ such that

$$E_{(r,\tilde{A},r',\tilde{\mathcal{U}}',\tilde{p}',\tilde{q}')} \cong E_{(r,\tilde{A},r',\tilde{\mathcal{U}},\tilde{p},\tilde{q})} \otimes E_{(1,O,1,\mathcal{V},u,v)}.$$

Here,

$$\mathcal{V} := \left\{ e^{i\tau_1}, \dots, e^{i\tau_n}, e^{i\sigma_1}, \dots, e^{i\sigma_n} \in U(1) \right\}, \quad u, v \in \mathbb{R}^n,$$

and for simplicity, we set

$$\tau := (\tau_1, \dots, \tau_n)^t, \quad \sigma := (\sigma_1, \dots, \sigma_n)^t \in \mathbb{R}^n.$$

In particular, for each $j = 1, \dots, n$, we assume that $e^{i\tau_j}$ and $e^{i\sigma_j}$ are the transition functions of $E_{(1,O,1,\mathcal{V},u,v)}$ in the x_j direction and the transition function of $E_{(1,O,1,\mathcal{V},u,v)}$ in the y_j direction, respectively. Therefore, since

$$E_{(r,\bar{A},r',\tilde{\mathcal{U}},\tilde{p},\tilde{q})} \cong E_{(r,\bar{A},r',\tilde{\mathcal{U}}',\tilde{p}',\tilde{q}')} \quad (15)$$

holds if and only if

$$E_{(r,\bar{A},r',\tilde{\mathcal{U}},\tilde{p},\tilde{q})} \cong E_{(r,\bar{A},r',\tilde{\mathcal{U}},\tilde{p},\tilde{q})} \otimes E_{(1,O,1,\mathcal{V},u,v)} \quad (16)$$

holds, our first goal is to find the relation on the parameters $u, v, \tau, \sigma \in \mathbb{R}^n$ such that the relation (16) holds. By using Theorem 3.3 again, we see that there exists a holomorphic line bundle $E_{(1,O,1,\mathcal{V}',u',v')}$ such that

$$E_{(r,\bar{A},r',\tilde{\mathcal{U}},\tilde{p},\tilde{q})} \cong E_{(r,\bar{A},r',\mathcal{U}_0,p_0,q_0)} \otimes E_{(1,O,1,\mathcal{V}',u',v')},$$

where

$$\mathcal{U}_0 := \left\{ V'_1, \dots, V'_n, U'_1, \dots, U'_n \in U(r') \right\}, \quad p_0, q_0 \in \mathbb{R}^n.$$

Here, note that the definitions of $V'_1, \dots, V'_n, U'_1, \dots, U'_n \in U(r')$ are given in the proof of Proposition 3.2. Thus, since we can rewrite the relation (16) to

$$E_{(r,\bar{A},r',\mathcal{U}_0,p_0,q_0)} \otimes E_{(1,O,1,\mathcal{V}',u',v')} \cong \left(E_{(r,\bar{A},r',\mathcal{U}_0,p_0,q_0)} \otimes E_{(1,O,1,\mathcal{V}',u',v')} \right) \otimes E_{(1,O,1,\mathcal{V},u,v)},$$

as a result, it is enough to consider the condition such that

$$E_{(r,\bar{A},r',\mathcal{U}_0,p_0,q_0)} \cong E_{(r,\bar{A},r',\mathcal{U}_0,p_0,q_0)} \otimes E_{(1,O,1,\mathcal{V},u,v)}$$

holds. By a direct calculation, we can actually check that

$$E_{(r,\bar{A},r',\mathcal{U}_0,p_0,q_0)} \cong E_{(r,\bar{A},r',\mathcal{U}_0,p_0,q_0)} \otimes E_{(1,O,1,\mathcal{V},u,v)}$$

holds if and only if

$$u + T'^t v \equiv \sigma - T'^t \tau \pmod{2\pi \left(\begin{pmatrix} \frac{\mathbb{Z}}{r'_1} \\ \vdots \\ \frac{\mathbb{Z}}{r'_s} \\ \mathbb{Z} \\ \vdots \\ \mathbb{Z} \end{pmatrix} \oplus T'^t \begin{pmatrix} \frac{\mathbb{Z}}{r'_1} \\ \vdots \\ \frac{\mathbb{Z}}{r'_s} \\ \mathbb{Z} \\ \vdots \\ \mathbb{Z} \end{pmatrix} \right)} \quad (17)$$

holds, and in particular, we can regard the relation (17) as

$$(\tilde{p} + T'^t \tilde{q}) - \{(\tilde{p} + T'^t \tilde{q}) + r(u + T'^t v)\} \equiv \frac{r}{r'} \{\tilde{\theta} - (\tilde{\theta} + r' \sigma)\} + T'^t \frac{r}{r'} \{(\tilde{\xi} + r' \tau) - \tilde{\xi}\}$$

$$\pmod{2\pi r \left(\begin{pmatrix} \frac{\mathbb{Z}}{r'_1} \\ \vdots \\ \frac{\mathbb{Z}}{r'_s} \\ \mathbb{Z} \\ \vdots \\ \mathbb{Z} \end{pmatrix} \oplus T'^t \begin{pmatrix} \frac{\mathbb{Z}}{r'_1} \\ \vdots \\ \frac{\mathbb{Z}}{r'_s} \\ \mathbb{Z} \\ \vdots \\ \mathbb{Z} \end{pmatrix} \right)}. \quad (18)$$

Hence, we see that the relation (16) holds if and only if the relation (18) holds. Here, note that there exists an isomorphism

$$\det E_{(r, \tilde{A}, r', \tilde{\mathcal{U}}, \tilde{p}, \tilde{q})} \cong \det(E_{(r, \tilde{A}, r', \tilde{\mathcal{U}}, \tilde{p}, \tilde{q})} \otimes E_{(1, \mathcal{O}, 1, \mathcal{V}, u, v)})$$

with

$$\begin{aligned} (\tilde{p} + T'^t \tilde{q}) - \{(\tilde{p} + T'^t \tilde{q}) + r(u + T'^t v)\} &\equiv \frac{r}{r'} \{\tilde{\theta} - (\tilde{\theta} + r'\sigma)\} + T'^t \frac{r}{r'} \{(\tilde{\xi} + r'\tau) - \tilde{\xi}\} \\ &\left(\text{mod } \frac{r}{r'} 2\pi(\mathbb{Z}^n \oplus T'^t \mathbb{Z}^n) \right). \end{aligned}$$

Now, we consider the condition such that the relation (15) holds. One necessary condition for the relation (15) is that

$$\det E_{(r, \tilde{A}, r', \tilde{\mathcal{U}}, \tilde{p}, \tilde{q})} \cong \det E_{(r, \tilde{A}, r', \tilde{\mathcal{U}}', \tilde{p}', \tilde{q}')} \quad (19)$$

holds, and by a direct calculation, we can rewrite the relation (19) to the following :

$$(\tilde{p} + T'^t \tilde{q}) - (\tilde{p}' + T'^t \tilde{q}') \equiv \frac{r}{r'} (\tilde{\theta} - \tilde{\theta}') + T'^t \frac{r}{r'} (\tilde{\xi}' - \tilde{\xi}) \left(\text{mod } \frac{r}{r'} 2\pi(\mathbb{Z}^n \oplus T'^t \mathbb{Z}^n) \right). \quad (20)$$

Therefore, since the definition of $E_{(1, \mathcal{O}, 1, \mathcal{V}, u, v)}$ is given by

$$E_{(r, \tilde{A}, r', \tilde{\mathcal{U}}', \tilde{p}', \tilde{q}')} \cong E_{(r, \tilde{A}, r', \tilde{\mathcal{U}}, \tilde{p}, \tilde{q})} \otimes E_{(1, \mathcal{O}, 1, \mathcal{V}, u, v)},$$

and the relations (16) and (18) are equivalent, by considering the relation (20), we can give the condition such that the relation (15) holds as follows.

$$(\tilde{p} + T'^t \tilde{q}) - (\tilde{p}' + T'^t \tilde{q}') \equiv \frac{r}{r'} (\tilde{\theta} - \tilde{\theta}') + T'^t \frac{r}{r'} (\tilde{\xi}' - \tilde{\xi}) \left(\text{mod } 2\pi r \left(\begin{pmatrix} \mathbb{Z} \\ r'_1 \\ \vdots \\ \mathbb{Z} \\ r'_s \\ \mathbb{Z} \\ \vdots \\ \mathbb{Z} \end{pmatrix} \oplus T'^t \begin{pmatrix} \mathbb{Z} \\ r'_1 \\ \vdots \\ \mathbb{Z} \\ r'_s \\ \mathbb{Z} \\ \vdots \\ \mathbb{Z} \end{pmatrix} \right) \right).$$

Thus, by considering the pullback bundles $\varphi^* E_{(r, \tilde{A}, r', \tilde{\mathcal{U}}, \tilde{p}, \tilde{q})}$ and $\varphi^* E_{(r, \tilde{A}, r', \tilde{\mathcal{U}}', \tilde{p}', \tilde{q}')}$, we can conclude that

$$E_{(r, A, r', \mathcal{U}, p, q)} \cong E_{(r, A, r', \mathcal{U}', p', q')}$$

holds if and only if

$$(p + T^t q) - (p' + T^t q') \equiv \frac{r}{r'} (\theta - \theta') + T^t \frac{r}{r'} (\xi' - \xi) \left(\text{mod } 2\pi r (\mathcal{A}^{-1} \left(\begin{pmatrix} \mathbb{Z} \\ r'_1 \\ \vdots \\ \mathbb{Z} \\ r'_s \\ \mathbb{Z} \\ \vdots \\ \mathbb{Z} \end{pmatrix} \oplus T'^t (\mathcal{B}^{-1})^t \begin{pmatrix} \mathbb{Z} \\ r'_1 \\ \vdots \\ \mathbb{Z} \\ r'_s \\ \mathbb{Z} \\ \vdots \\ \mathbb{Z} \end{pmatrix} \right) \right)$$

holds. \square

Remark 3.5. *When we work over a pair $(T_{J=T}^2, \tilde{T}_{J=T}^2)$ of elliptic curves, Theorem 3.4 implies that there exists a one-to-one correspondence between the set of the isomorphism classes of holomorphic vector bundles $E_{(r,A,r',\mathcal{U},p,q)}$ and the set of the isomorphism classes of holomorphic line bundles $\det E_{(r,A,r',\mathcal{U},p,q)}$.*

4 Symplectic geometry side

In this section, we define the objects of the Fukaya category corresponding to holomorphic vector bundles $E_{(r,A,r',\mathcal{U},p,q)} \rightarrow T_{J=T}^{2n}$, and study the isomorphism classes of them. The discussions in this section are based on the SYZ construction (SYZ transform) [20] (see also [14], [2]).

4.1 The definition of $\mathcal{L}_{(r,A,p,q)}$

In this subsection, we define a class of pairs of affine Lagrangian submanifolds

$$L_{(r,A,p)}$$

in $\tilde{T}_{J=T}^{2n}$ and unitary local systems

$$\mathcal{L}_{(r,A,p,q)} \rightarrow L_{(r,A,p)}.$$

First, we recall the definition of objects of the Fukaya categories following [5, Definition 1.1]. Let (M, Ω) be a symplectic manifold (M, ω) together with a closed 2-form B on M . Here, we put $\Omega = \omega + \sqrt{-1}B$ (note $-B + \sqrt{-1}\omega$ is used in many of the literatures). Then, we consider pairs (L, \mathcal{L}) with the following properties :

$$L \text{ is a Lagrangian submanifold of } (M, \omega). \quad (21)$$

$$\mathcal{L} \rightarrow L \text{ is a line bundle together with a connection } \nabla^{\mathcal{L}} \text{ such that} \quad (22)$$

$$F_{\nabla^{\mathcal{L}}} = 2\pi\sqrt{-1}B|_L.$$

In this context, $F_{\nabla^{\mathcal{L}}}$ denotes the curvature form of the connection $\nabla^{\mathcal{L}}$. We define objects of the Fukaya category on (M, Ω) by pairs (L, \mathcal{L}) which satisfy the properties (21), (22).

Let us consider the following n -dimensional submanifold $\tilde{L}_{(r,A,p)}$ in \mathbb{R}^{2n} :

$$\tilde{L}_{(r,A,p)} := \left\{ \begin{pmatrix} \tilde{x} \\ \tilde{y} \end{pmatrix} \in \mathbb{R}^{2n} \mid \tilde{y} = \frac{1}{r}A\tilde{x} + \frac{1}{r^t}p \right\}.$$

We see that this n -dimensional submanifold $\tilde{L}_{(r,A,p)}$ satisfies the property (21), namely, $\tilde{L}_{(r,A,p)}$ becomes a Lagrangian submanifold in \mathbb{R}^{2n} if and only if $\omega A = (\omega A)^t$ holds. Then, for the covering map $\pi : \mathbb{R}^{2n} \rightarrow \tilde{T}_{J=T}^{2n}$,

$$L_{(r,A,p)} := \pi(\tilde{L}_{(r,A,p)})$$

defines a Lagrangian submanifold in $\tilde{T}_{j=T}^{2n}$. On the other hand, we can also regard the complexified symplectic torus $\tilde{T}_{j=T}^{2n}$ as the trivial special Lagrangian torus fibration $\tilde{\pi} : \tilde{T}_{j=T}^{2n} \rightarrow \mathbb{R}^n/2\pi\mathbb{Z}^n$, where \tilde{x} is the local coordinates of the base space $\mathbb{R}^n/2\pi\mathbb{Z}^n$ and \tilde{y} is the local coordinates of the fiber of $\tilde{\pi} : \tilde{T}_{j=T}^{2n} \rightarrow \mathbb{R}^n/2\pi\mathbb{Z}^n$. Then, we can interpret each affine Lagrangian submanifold $L_{(r,A,p)}$ in $\tilde{T}_{j=T}^{2n}$ as the affine Lagrangian multi section

$$s(\tilde{x}) = \frac{1}{r}A\tilde{x} + \frac{1}{r}p$$

of $\tilde{\pi} : \tilde{T}_{j=T}^{2n} \rightarrow \mathbb{R}^n/2\pi\mathbb{Z}^n$.

Remark 4.1. As explained in section 3, while $r' := r'_1 \cdots r'_s \in \mathbb{N}$ is the rank of $E_{(r,A,r',\mathcal{U},p,q)} \rightarrow T_{j=T}^{2n}$ (see the relations (2) and (3)), in the symplectic geometry side, this $r' \in \mathbb{N}$ is interpreted as follows. For the affine Lagrangian submanifold $L_{(r,A,p)}$ in $\tilde{T}_{j=T}^{2n}$ which is defined by a given data $(r, A, p) \in \mathbb{N} \times M(n; \mathbb{Z}) \times \mathbb{R}^n$, we regard it as the affine Lagrangian multi section $s(\tilde{x}) = \frac{1}{r}A\tilde{x} + \frac{1}{r}p$ of $\tilde{\pi} : \tilde{T}_{j=T}^{2n} \rightarrow \mathbb{R}^n/2\pi\mathbb{Z}^n$. Then, for each point $\tilde{x} \in \mathbb{R}^n/2\pi\mathbb{Z}^n$, we see

$$s(\tilde{x}) = \left\{ \left(\frac{1}{r}A\tilde{x} + \frac{1}{r}p + \frac{2\pi}{r}ABM_s \right) \in \tilde{\pi}^{-1}(\tilde{x}) \approx \mathbb{R}^n/2\pi\mathbb{Z}^n \mid \right. \\ \left. M_s = (m_1, \cdots, m_s, 0, \cdots, 0)^t \in \mathbb{Z}^n, 0 \leq m_i \leq r'_i - 1, i = 1, \cdots, s \right\},$$

and this indicates that $s(\tilde{x})$ consists of r' points. Thus, we can regard $r' \in \mathbb{N}$ as the multiplicity of $s(\tilde{x}) = \frac{1}{r}A\tilde{x} + \frac{1}{r}p$.

We then consider the trivial complex line bundle

$$\mathcal{L}_{(r,A,p,q)} \rightarrow L_{(r,A,p)}$$

with the flat connection

$$\nabla_{\mathcal{L}_{(r,A,p,q)}} := d - \frac{\mathbf{i}}{2\pi} \frac{1}{r} q^t d\tilde{x},$$

where $q \in \mathbb{R}^n$ is the unitary holonomy of $\mathcal{L}_{(r,A,p,q)}$ along $L_{(r,A,p)} \approx T^n$. We discuss the property (22) for this pair $(L_{(r,A,p)}, \mathcal{L}_{(r,A,p,q)})$:

$$\Omega_{\mathcal{L}_{(r,A,p,q)}} = d\tilde{x}^t B d\tilde{y} \Big|_{L_{(r,A,p)}}.$$

Here, $\Omega_{\mathcal{L}_{(r,A,p,q)}}$ is the curvature form of the flat connection $\nabla_{\mathcal{L}_{(r,A,p,q)}}$, i.e., $\Omega_{\mathcal{L}_{(r,A,p,q)}} = 0$. Hence, we see

$$d\tilde{x}^t B d\tilde{y} \Big|_{L_{(r,A,p)}} = \frac{1}{r} d\tilde{x}^t B A d\tilde{x} = 0,$$

so one has $BA = (BA)^t$. Note that $\omega A = (\omega A)^t$ and $BA = (BA)^t$ hold if and only if $AT = (AT)^t$ holds. Hereafter, for simplicity, we set

$$\mathcal{L}_{(r,A,p,q)} := (L_{(r,A,p)}, \mathcal{L}_{(r,A,p,q)}).$$

By summarizing the above discussions, we obtain the following proposition. In particular, the condition $AT = (AT)^t$ in the following proposition is also the condition such that a complex vector bundle $E_{(r,A,r',\mathcal{U},p,q)} \rightarrow T_{J=T}^{2n}$ becomes a holomorphic vector bundle (see Proposition 3.1).

Proposition 4.2. *For a given quadruple $(r, A, p, q) \in \mathbb{N} \times M(n; \mathbb{Z}) \times \mathbb{R}^n \times \mathbb{R}^n$, $\mathcal{L}_{(r,A,p,q)}$ gives an object of the Fukaya category on $\tilde{T}_{J=T}^{2n}$ if and only if $AT = (AT)^t$ holds.*

Definition 4.3. *We denote the full subcategory of the Fukaya category on $\tilde{T}_{J=T}^{2n}$ consisting of objects $\mathcal{L}_{(r,A,p,q)}$ which satisfy the condition $AT = (AT)^t$ by $\text{Fuk}_{\text{aff}}(\tilde{T}_{J=T}^{2n})$.*

4.2 The isomorphism classes of $\mathcal{L}_{(r,A,p,q)}$

The discussions in this subsection correspond to the discussions in subsection 3.2, so throughout this subsection, we fix $r \in \mathbb{N}$, $A \in M(n; \mathbb{Z})$, and consider the condition such that $\mathcal{L}_{(r,A,p,q)} \cong \mathcal{L}_{(r,A,p',q')}$ holds, where

$$p, q, p', q' \in \mathbb{R}^n.$$

We explain the definition of the equivalence of objects of the Fukaya categories (cf. [5, Definition 1.4]). Let us consider a pair (M, Ω) consisting of an even dimensional differentiable manifold M and a complexified symplectic form Ω . We take two objects $\mathcal{L} := (L, \mathcal{L})$, $\mathcal{L}' := (L', \mathcal{L}')$ of the Fukaya category $\text{Fuk}(M, \Omega)$, where L, L' are Lagrangian submanifolds in (M, Ω) , and $\mathcal{L} \rightarrow L$, $\mathcal{L}' \rightarrow L'$ are unitary local systems. Then, if there exists a symplectic automorphism $\Phi : (M, \Omega) \xrightarrow{\sim} (M, \Omega)$ such that

$$\begin{aligned} \Phi^{-1}(L') &= L, \\ \Phi^* \mathcal{L}' &\cong \mathcal{L}, \end{aligned}$$

we say that \mathcal{L} is isomorphic to \mathcal{L}' , and write $\mathcal{L} \cong \mathcal{L}'$.

We consider the isomorphism classes of $\mathcal{L}_{(r,A,p,q)}$, namely, we consider the condition such that $\mathcal{L}_{(r,A,p,q)} \cong \mathcal{L}_{(r,A,p',q')}$ holds as an analogue of Theorem 3.4. Actually, the following theorem holds.

Theorem 4.4. *Two objects $\mathcal{L}_{(r,A,p,q)}$, $\mathcal{L}_{(r,A,p',q')}$ are isomorphic to each other,*

$$\mathcal{L}_{(r,A,p,q)} \cong \mathcal{L}_{(r,A,p',q')},$$

if and only if

$$p \equiv p' \pmod{2\pi r A^{-1} \begin{pmatrix} \mathbb{Z} \\ r'_1 \\ \vdots \\ \mathbb{Z} \\ r'_s \\ \mathbb{Z} \\ \vdots \\ \mathbb{Z} \end{pmatrix}}, \quad q \equiv q' \pmod{2\pi r (\mathcal{B}^{-1})^t \begin{pmatrix} \mathbb{Z} \\ r'_1 \\ \vdots \\ \mathbb{Z} \\ r'_s \\ \mathbb{Z} \\ \vdots \\ \mathbb{Z} \end{pmatrix}}$$

hold.

Proof. First, in order to prove the statement of this theorem, we prepare some notations. Since we considered the complex torus $T_{J=T'}^{2n} = \mathbb{C}^n / 2\pi(\mathbb{Z}^n \oplus \mathcal{B}^{-1}T\mathcal{A}^t\mathbb{Z}^n)$ which is biholomorphic to the complex torus $T_{J=T}^{2n}$ in Theorem 3.4 (and Proposition 3.2), we take a mirror partner of the complex torus $T_{J=T'}^{2n}$. Here, we consider the complexified symplectic torus $\check{T}_{J=T'}^{2n} := T_{\check{\omega}=(-T'^{-1})^t}^{2n}$ as a mirror partner of the complex torus $T_{J=T}^{2n}$. We denote the local coordinates of $\check{T}_{J=T'}^{2n}$ by $(X^1, \dots, X^n, Y^1, \dots, Y^n)^t$, and set

$$\check{X} := (X^1, \dots, X^n)^t, \quad \check{Y} := (Y^1, \dots, Y^n)^t.$$

Let us consider a diffeomorphism $\phi : \check{T}_{J=T}^{2n} \xrightarrow{\sim} \check{T}_{J=T'}^{2n}$ which is expressed locally as

$$\begin{pmatrix} \check{X} \\ \check{Y} \end{pmatrix} = \phi \begin{pmatrix} \check{x} \\ \check{y} \end{pmatrix} = \begin{pmatrix} \mathcal{B}^{-1} & O \\ O & \mathcal{A} \end{pmatrix} \begin{pmatrix} \check{x} \\ \check{y} \end{pmatrix}.$$

By a direct calculation, we see

$$\phi^*(d\check{X}^t(-T'^{-1})^t d\check{Y}) = d\check{x}^t(-T^{-1})^t d\check{y},$$

where $d\check{X} := (dX^1, \dots, dX^n)^t$, $d\check{Y} := (dY^1, \dots, dY^n)^t$, so this diffeomorphism ϕ is a symplectomorphism.

Now, we define

$$\begin{aligned} (\phi^{-1})^* \mathcal{L}_{(r,A,p,q)} &:= ((\phi^{-1})^{-1}(L_{(r,A,p)}), (\phi^{-1})^* \mathcal{L}_{(r,A,p,q)}), \\ (\phi^{-1})^* \mathcal{L}_{(r,A,p',q')} &:= ((\phi^{-1})^{-1}(L_{(r,A,p')}), (\phi^{-1})^* \mathcal{L}_{(r,A,p',q')}), \end{aligned}$$

and let us consider the condition such that

$$(\phi^{-1})^* \mathcal{L}_{(r,A,p,q)} \cong (\phi^{-1})^* \mathcal{L}_{(r,A,p',q')}$$

holds. Namely, our first goal is to consider when it is possible to construct a symplectic automorphism $\Phi : \check{T}_{J=T'}^{2n} \xrightarrow{\sim} \check{T}_{J=T}^{2n}$ such that

$$\Phi^{-1}((\phi^{-1})^{-1}(L_{(r,A,p')})) = (\phi^{-1})^{-1}(L_{(r,A,p)}), \quad (23)$$

$$\Phi^*(\phi^{-1})^* \mathcal{L}_{(r,A,p',q')} \cong (\phi^{-1})^* \mathcal{L}_{(r,A,p,q)}. \quad (24)$$

We consider the condition (23). Since $r \in \mathbb{N}$, $A \in M(n; \mathbb{Z})$ are fixed, we can take the map

$$\Phi = \text{id}_{\check{T}_{J=T'}^{2n}}$$

as a symplectic automorphism $\Phi : \check{T}_{J=T'}^{2n} \xrightarrow{\sim} \check{T}_{J=T}^{2n}$ which satisfies the condition (23), in the case $(\phi^{-1})^{-1}(L_{(r,A,p)}) = (\phi^{-1})^{-1}(L_{(r,A,p')})$ only. Moreover,

$$(\phi^{-1})^{-1}(L_{(r,A,p)}) = (\phi^{-1})^{-1}(L_{(r,A,p')})$$

holds if and only if

$$\mathcal{A}p \equiv \mathcal{A}p' \pmod{2\pi r \begin{pmatrix} \frac{\mathbb{Z}}{r'_1} \\ \vdots \\ \frac{\mathbb{Z}}{r'_s} \\ \mathbb{Z} \\ \vdots \\ \mathbb{Z} \end{pmatrix}} \quad (25)$$

holds. Then, the condition (24) becomes

$$(\phi^{-1})^* \mathcal{L}_{(r,A,p,q)} \cong (\phi^{-1})^* \mathcal{L}_{(r,A,p',q')},$$

so hereafter, we consider when $(\phi^{-1})^* \mathcal{L}_{(r,A,p,q)} \cong (\phi^{-1})^* \mathcal{L}_{(r,A,p',q')}$ holds on $(\phi^{-1})^{-1}(L_{(r,A,p)}) = (\phi^{-1})^{-1}(L_{(r,A,p')})$ by computing an isomorphism

$$\psi : (\phi^{-1})^* \mathcal{L}_{(r,A,p,q)} \xrightarrow{\sim} (\phi^{-1})^* \mathcal{L}_{(r,A,p',q')}$$

explicitly. The morphism ψ need to satisfy the differential equation

$$\nabla_{(\phi^{-1})^* \mathcal{L}_{(r,A,p',q')}} \psi = \psi \nabla_{(\phi^{-1})^* \mathcal{L}_{(r,A,p,q)}}. \quad (26)$$

In particular, since the differential equation (26) turns out to be

$$\left(\frac{\partial \psi}{\partial X^1}, \dots, \frac{\partial \psi}{\partial X^n} \right) - \frac{\mathbf{i}}{2\pi r} ((q'^t \mathcal{B})_1 - (q^t \mathcal{B})_1, \dots, (q'^t \mathcal{B})_n - (q^t \mathcal{B})_n) \psi = 0, \quad (27)$$

by solving the differential equation (27), we obtain a solution

$$\psi(\tilde{X}) = \lambda e^{\frac{\mathbf{i}}{2\pi r} (q' - q)^t \mathcal{B} \tilde{X}}, \quad (28)$$

where $\lambda \in \mathbb{C}$ is an arbitrary constant. Furthermore, since $(\phi^{-1})^* \mathcal{L}_{(r,A,p,q)}$ and $(\phi^{-1})^* \mathcal{L}_{(r,A,p',q')}$ are trivial, this morphism ψ must satisfy

$$\begin{aligned} \psi(X^1, \dots, X^i + 2\pi r'_i, \dots, X^n) &= \psi(X^1, \dots, X^i, \dots, X^n) \quad (i = 1, \dots, s), \\ \psi(X^1, \dots, X^i + 2\pi, \dots, X^n) &= \psi(X^1, \dots, X^i, \dots, X^n) \quad (i = s + 1, \dots, n). \end{aligned}$$

By a direct calculation, we see that

$$\begin{aligned} \psi(X^1, \dots, X^i + 2\pi r'_i, \dots, X^n) &= e^{\frac{r'_i}{r} \mathbf{i} (q' - q)^t \mathcal{B}_i} \psi(X^1, \dots, X^i, \dots, X^n) \quad (i = 1, \dots, s), \\ \psi(X^1, \dots, X^i + 2\pi, \dots, X^n) &= e^{\frac{\mathbf{i}}{r} (q' - q)^t \mathcal{B}_i} \psi(X^1, \dots, X^i, \dots, X^n) \quad (i = s + 1, \dots, n) \end{aligned}$$

hold, where $\mathcal{B}_i := (\mathcal{B}_{1i}, \dots, \mathcal{B}_{ni})^t$, so we obtain

$$\begin{aligned} e^{\frac{r'_i}{r} \mathbf{i} (q' - q)^t \mathcal{B}_i} &= 1 \quad (i = 1, \dots, s), \\ e^{\frac{\mathbf{i}}{r} (q' - q)^t \mathcal{B}_i} &= 1 \quad (i = s + 1, \dots, n). \end{aligned}$$

Clearly, these relations are equivalent to

$$\mathcal{B}^t(q' - q) = 2\pi r N_{r'},$$

where

$$N_{r'} := \left(\frac{N_1}{r'_1}, \dots, \frac{N_s}{r'_s}, N_{s+1}, \dots, N_n \right)^t, \quad (N_1, \dots, N_s, N_{s+1}, \dots, N_n) \in \mathbb{Z}^n,$$

and then, by the formula (28), the isomorphism ψ is expressed locally as

$$\psi(\check{X}) = \lambda e^{iN_{r'}^t \check{X}}$$

with $\lambda \neq 0 \in \mathbb{C}$. Hence,

$$(\phi^{-1})^* \mathcal{L}_{(r,A,p,q)} \cong (\phi^{-1})^* \mathcal{L}_{(r,A,p',q')}$$

holds on $(\phi^{-1})^{-1}(L_{(r,A,p)}) = (\phi^{-1})^{-1}(L_{(r,A,p')})$ if and only if

$$\mathcal{B}^t q \equiv \mathcal{B}^t q' \pmod{2\pi r \begin{pmatrix} \frac{\mathbb{Z}}{r'_1} \\ \vdots \\ \frac{\mathbb{Z}}{r'_s} \\ \mathbb{Z} \\ \vdots \\ \mathbb{Z} \end{pmatrix}} \quad (29)$$

holds. Thus, by the relations (25) and (29), we can conclude that

$$\mathcal{L}_{(r,A,p,q)} \cong \mathcal{L}_{(r,A,p',q')}$$

holds if and only if

$$p \equiv p' \pmod{2\pi r \mathcal{A}^{-1} \begin{pmatrix} \frac{\mathbb{Z}}{r'_1} \\ \vdots \\ \frac{\mathbb{Z}}{r'_s} \\ \mathbb{Z} \\ \vdots \\ \mathbb{Z} \end{pmatrix}}, \quad q \equiv q' \pmod{2\pi r (\mathcal{B}^{-1})^t \begin{pmatrix} \frac{\mathbb{Z}}{r'_1} \\ \vdots \\ \frac{\mathbb{Z}}{r'_s} \\ \mathbb{Z} \\ \vdots \\ \mathbb{Z} \end{pmatrix}}$$

hold. □

Hence, by comparing Theorem 3.4 with Theorem 4.4, we can expect that the isomorphism classes of holomorphic vector bundles $E_{(r,A,r',\mathcal{U},p,q)} \rightarrow T_{j=T}^{2n}$ correspond to the isomorphism classes of objects $\mathcal{L}_{(r,A,p,q)}$ of the Fukaya category $Fuk_{\text{aff}}(\check{T}_{j=T}^{2n})$. Actually, by a direct calculation, we can check that there exists such a correspondence.

5 Main result

In this section, we prove that there exists a bijection between the set of the isomorphism classes of holomorphic vector bundles $E_{(r,A,r',\mathcal{U},p,q)} \rightarrow T_{J=T}^{2n}$ and the set of the isomorphism classes of objects $\mathcal{L}_{(r,A,p,q)}$ of the Fukaya category $Fuk_{\text{aff}}(\check{T}_{J=T}^{2n})$.

First, we prepare two notations. We denote the set of the isomorphism classes of objects of the DG-category $DG_{T_{J=T}^{2n}}$ (i.e., the set of the isomorphism classes of holomorphic vector bundles $E_{(r,A,r',\mathcal{U},p,q)}$) by

$$\text{Ob}^{isom}(DG_{T_{J=T}^{2n}}).$$

Similarly, we denote the set of the isomorphism classes of objects $\mathcal{L}_{(r,A,p,q)}$ of the Fukaya category $Fuk_{\text{aff}}(\check{T}_{J=T}^{2n})$ by

$$\text{Ob}^{isom}(Fuk_{\text{aff}}(\check{T}_{J=T}^{2n})).$$

Now, in order to state the main theorem, we define a map $F : \text{Ob}(DG_{T_{J=T}^{2n}}) \rightarrow \text{Ob}(Fuk_{\text{aff}}(\check{T}_{J=T}^{2n}))$ as follows. Clearly, we need four parameters r, A, p, q when we define objects $\mathcal{L}_{(r,A,p,q)}$ of $Fuk_{\text{aff}}(\check{T}_{J=T}^{2n})$. On the contrary, we need five parameters r, A, p, q, \mathcal{U} when we define objects $E_{(r,A,r',\mathcal{U},p,q)}$ of $DG_{T_{J=T}^{2n}}$. Hence, when we define a map $\text{Ob}(DG_{T_{J=T}^{2n}}) \rightarrow \text{Ob}(Fuk_{\text{aff}}(\check{T}_{J=T}^{2n}))$, we must transform not only the information about four parameters r, A, p, q but also the information about \mathcal{U} . For example, let us consider a map $\text{Ob}(DG_{T_{J=T}^{2n}}) \rightarrow \text{Ob}(Fuk_{\text{aff}}(\check{T}_{J=T}^{2n}))$ which is simply defined by

$$E_{(r,A,r',\mathcal{U},p,q)} \mapsto \mathcal{L}_{(r,A,p,q)}.$$

Then, this map does not induce a bijection between $\text{Ob}^{isom}(DG_{T_{J=T}^{2n}})$ and $\text{Ob}^{isom}(Fuk_{\text{aff}}(\check{T}_{J=T}^{2n}))$ unfortunately, and we can check it as follows. We set

$$T = \mathbf{i} \cdot I_2, \quad r = 2, \quad A = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad p = q = 0.$$

It is clear that $AT = (AT)^t$ holds and $r' = 2$. For this quadruple $(r, A, p, q) \in \mathbb{N} \times M(2; \mathbb{Z}) \times \mathbb{R}^2 \times \mathbb{R}^2$, we define mutually distinct \mathcal{U} and \mathcal{U}' by

$$\begin{aligned} \mathcal{U} &:= \left\{ V_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, U_1 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, V_2 = U_2 = I_2 \in U(2) \right\}, \\ \mathcal{U}' &:= \left\{ V'_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, U'_1 = \begin{pmatrix} \mathbf{i} & 0 \\ 0 & -\mathbf{i} \end{pmatrix}, V'_2 = U'_2 = I_2 \in U(2) \right\}. \end{aligned}$$

In this situation, we can easily check that $E_{(r,A,r',\mathcal{U},p,q)} \not\cong E_{(r,A,r',\mathcal{U}',p,q)}$ holds by using Theorem 3.4, and actually, the above map $E_{(r,A,r',\mathcal{U},p,q)} \mapsto \mathcal{L}_{(r,A,p,q)}$ sends both $E_{(r,A,r',\mathcal{U},p,q)}$ and $E_{(r,A,r',\mathcal{U}',p,q)}$ to the same object $\mathcal{L}_{(r,A,p,q)}$. Thus, by concerning these facts, here, we define a map

$$F : \text{Ob}(DG_{T_{J=T}^{2n}}) \rightarrow \text{Ob}(Fuk_{\text{aff}}(\check{T}_{J=T}^{2n}))$$

by

$$F(E_{(r,A,r',\mathcal{U},p,q)}) = \mathcal{L}_{(r,A,p-\frac{r}{r'}\theta, q+\frac{r}{r'}\xi)},$$

where $\xi, \theta \in \mathbb{R}^n$ denote the vectors associated to \mathcal{U} in the sense of the definition (14). The following is the main theorem of this paper.

Theorem 5.1. *The map F induces a bijection between $\text{Ob}^{isom}(DG_{T_{J=T}^{2n}})$ and $\text{Ob}^{isom}(Fuk_{\text{aff}}(\check{T}_{J=T}^{2n}))$.*

Proof. In this proof, for a given object $E_{(r,A,r',\mathcal{U},p,q)} \in \text{Ob}(DG_{T_{J=T}^{2n}})$, we denote by $\xi, \theta \in \mathbb{R}^n$ the vectors associated to \mathcal{U} in the sense of the definition (14). Similarly, for a given object $E_{(s,B,s',\mathcal{V},u,v)} \in \text{Ob}(DG_{T_{J=T}^{2n}})$, we denote by $\tau, \sigma \in \mathbb{R}^n$ the vectors associated to \mathcal{V} in the sense of the definition (14). We denote the induced map from the map F by

$$F^{isom} : \text{Ob}^{isom}(DG_{T_{J=T}^{2n}}) \rightarrow \text{Ob}^{isom}(Fuk_{\text{aff}}(\check{T}_{J=T}^{2n})).$$

Explicitly, it is defined by

$$F^{isom}([E_{(r,A,r',\mathcal{U},p,q)}]) = [F(E_{(r,A,r',\mathcal{U},p,q)})],$$

where, of course, $[E_{(r,A,r',\mathcal{U},p,q)}]$ and $[F(E_{(r,A,r',\mathcal{U},p,q)})]$ denote the isomorphism class of $E_{(r,A,r',\mathcal{U},p,q)}$ and the isomorphism class of $F(E_{(r,A,r',\mathcal{U},p,q)})$, respectively.

First, we check the well-definedness of the map F^{isom} . We take two arbitrary objects $E_{(r,A,r',\mathcal{U},p,q)}, E_{(s,B,s',\mathcal{V},u,v)} \in \text{Ob}(DG_{T_{J=T}^{2n}})$ and assume

$$E_{(r,A,r',\mathcal{U},p,q)} \cong E_{(s,B,s',\mathcal{V},u,v)}.$$

By considering the i -th Chern characters $ch_i(E_{(r,A,r',\mathcal{U},p,q)})$, $ch_i(E_{(s,B,s',\mathcal{V},u,v)})$ of the holomorphic vector bundles $E_{(r,A,r',\mathcal{U},p,q)}$, $E_{(s,B,s',\mathcal{V},u,v)}$ for each $i \in \mathbb{N}$, we see

$$ch_i(E_{(r,A,r',\mathcal{U},p,q)}) = ch_i(E_{(s,B,s',\mathcal{V},u,v)}). \quad (30)$$

We consider the equality (30) in the cases $i = 0, 1$. Then, we obtain

$$\begin{aligned} r' &= s', \\ \frac{r'}{r}A &= \frac{s'}{s}B, \end{aligned}$$

so one has

$$\frac{1}{r}A = \frac{1}{s}B.$$

Hence, we see that there exists a $k \in \mathbb{N}$ such that

$$s = kr, \quad (31)$$

$$B = kA. \quad (32)$$

Therefore, since we can regard $E_{(s,B,s',\mathcal{V},u,v)}$ as $E_{(kr,kA,r',\mathcal{V},u,v)} = E_{(r,A,r',\mathcal{V},\frac{1}{k}u,\frac{1}{k}v)}$, by Theorem 3.4, we see

$$p - \frac{r}{r'}\theta \equiv \frac{1}{k} \left(u - \frac{s}{s'}\sigma \right) \pmod{2\pi r \mathcal{A}^{-1} \begin{pmatrix} \frac{\mathbb{Z}}{r'_1} \\ \vdots \\ \frac{\mathbb{Z}}{r'_s} \\ \mathbb{Z} \end{pmatrix}}, \quad (33)$$

$$q + \frac{r}{r'}\xi \equiv \frac{1}{k} \left(v + \frac{s}{s'}\tau \right) \pmod{2\pi r (\mathcal{B}^{-1})^t \begin{pmatrix} \frac{\mathbb{Z}}{r'_1} \\ \vdots \\ \frac{\mathbb{Z}}{r'_s} \\ \mathbb{Z} \end{pmatrix}}. \quad (34)$$

Thus, by Theorem 4.4 and the relations (31), (32), (33), (34), we can conclude

$$\mathcal{L}_{(r,A,p-\frac{r}{r'}\theta,q+\frac{r}{r'}\xi)} \cong \mathcal{L}_{(s,B,u-\frac{s}{s'}\sigma,v+\frac{s}{s'}\tau)},$$

namely,

$$F(E_{(r,A,r',\mathcal{U},p,q)}) \cong F(E_{(s,B,s',\mathcal{V},u,v)}).$$

Next, we prove that F^{isom} is injective. We take two arbitrary objects $E_{(r,A,r',\mathcal{U},p,q)}, E_{(s,B,s',\mathcal{V},u,v)} \in \text{Ob}(DG_{T_{j=T}^{2n}})$ and assume

$$F(E_{(r,A,r',\mathcal{U},p,q)}) \cong F(E_{(s,B,s',\mathcal{V},u,v)}),$$

namely,

$$\mathcal{L}_{(r,A,p-\frac{r}{r'}\theta,q+\frac{r}{r'}\xi)} \cong \mathcal{L}_{(s,B,u-\frac{s}{s'}\sigma,v+\frac{s}{s'}\tau)}.$$

Then, we see that there exists a $k \in \mathbb{N}$ which satisfies the relations (31) and (32). Here, we take two matrices $\mathcal{A}, \mathcal{B} \in GL(n; \mathbb{Z})$ such that

$$\mathcal{A}\mathcal{B} = \begin{pmatrix} \tilde{a}_1 & & & & & & \\ & \ddots & & & & & \\ & & \tilde{a}_t & & & & \\ & & & 0 & & & \\ & & & & \ddots & & \\ & & & & & & 0 \end{pmatrix}, \quad (35)$$

where $\tilde{a}_i \in \mathbb{N}$ ($i = 1, \dots, t, 1 \leq t \leq n$) and $\tilde{a}_i | a_{i+1}$ ($i = 1, \dots, t-1$). Therefore,

since the relation (32) holds, we obtain

$$ABB = \mathcal{A}(kA)\mathcal{B} = \begin{pmatrix} k\tilde{a}_1 & & & & & & & \\ & \ddots & & & & & & \\ & & k\tilde{a}_t & & & & & \\ & & & 0 & & & & \\ & & & & \ddots & & & \\ & & & & & & & 0 \end{pmatrix}. \quad (36)$$

In particular, the relations (31), (32), (35), (36) imply

$$r' = s'. \quad (37)$$

Hence, by Theorem 4.4, the relations (33) and (34) hold. Now, note that we can regard $E_{(s,B,s',\mathcal{V},u,v)}$ as $E_{(kr,kA,r',\mathcal{V},u,v)} = E_{(r,A,r',\mathcal{V},\frac{1}{k}u,\frac{1}{k}v)}$ by the relations (31), (32), (37). Thus, by Theorem 3.4 and the relations (33) and (34), we see that

$$E_{(r,A,r',\mathcal{U},p,q)} \cong E_{(r,A,r',\mathcal{V},\frac{1}{k}u,\frac{1}{k}v)}$$

holds, and this relation indicates

$$E_{(r,A,r',\mathcal{U},p,q)} \cong E_{(s,B,s',\mathcal{V},u,v)}.$$

Finally, we prove that F^{isom} is surjective. We take an arbitrary quadruple $(r, A, p, q) \in \mathbb{N} \times M(n; \mathbb{Z}) \times \mathbb{R}^n \times \mathbb{R}^n$, and consider the element

$$[\mathcal{L}_{(r,A,p,q)}] \in \text{Ob}^{isom}(Fuk_{\text{aff}}(\check{T}_{J=T}^{2n})).$$

In particular, a representative of $[\mathcal{L}_{(r,A,p,q)}]$ is expressed as

$$\mathcal{L}_{(r,A,p+2\pi r A^{-1}M,q+2\pi r(B^{-1})^tN)}$$

by using a pair

$$(M, N) \in \begin{pmatrix} \mathbb{Z} \\ r'_1 \\ \vdots \\ \mathbb{Z} \\ r'_s \\ \mathbb{Z} \\ \vdots \\ \mathbb{Z} \end{pmatrix} \times \begin{pmatrix} \mathbb{Z} \\ r'_1 \\ \vdots \\ \mathbb{Z} \\ r'_s \\ \mathbb{Z} \\ \vdots \\ \mathbb{Z} \end{pmatrix}$$

(see Theorem 4.4). For the element $[\mathcal{L}_{(r,A,p,q)}]$, we consider the element

$$[E_{(r,A,r',\mathcal{U},p+\frac{r}{r'}\theta,q-\frac{r}{r'}\xi)}] \in \text{Ob}^{isom}(DG_{T_{J=T}^{2n}}),$$

where $\xi, \theta \in \mathbb{R}^n$ are the vectors associated to \mathcal{U} in the sense of the definition (14). Here, note that how to choose a set \mathcal{U} is not unique even if we fix a

quadruple $(r, A, p, q) \in \mathbb{N} \times M(n; \mathbb{Z}) \times \mathbb{R}^n \times \mathbb{R}^n$. Therefore, a representative of $[E_{(r, A, r', \mathcal{U}, p + \frac{r}{r'}\theta, q - \frac{r}{r'}\xi)]$ is expressed as

$$E_{(r, A, r', \mathcal{U}', p + \frac{r}{r'}\theta' + 2\pi r A^{-1}M, q - \frac{r}{r'}\xi' + 2\pi r(\mathcal{B}^{-1})^t N)}$$

by using a set \mathcal{U}' with the associated vectors $\xi', \theta' \in \mathbb{R}^n$ and a pair

$$(M, N) \in \begin{pmatrix} \frac{\mathbb{Z}}{r'_1} \\ \vdots \\ \frac{\mathbb{Z}}{r'_s} \\ \mathbb{Z} \\ \vdots \\ \mathbb{Z} \end{pmatrix} \times \begin{pmatrix} \frac{\mathbb{Z}}{r'_1} \\ \vdots \\ \frac{\mathbb{Z}}{r'_s} \\ \mathbb{Z} \\ \vdots \\ \mathbb{Z} \end{pmatrix}$$

(see Theorem 3.4). Then, by a direct calculation, we see

$$\begin{aligned} & F^{isom}([E_{(r, A, r', \mathcal{U}, p + \frac{r}{r'}\theta, q - \frac{r}{r'}\xi)]) \\ &= F^{isom}([E_{(r, A, r', \mathcal{U}', p + \frac{r}{r'}\theta' + 2\pi r A^{-1}M, q - \frac{r}{r'}\xi' + 2\pi r(\mathcal{B}^{-1})^t N)]) \\ &= [F(E_{(r, A, r', \mathcal{U}', p + \frac{r}{r'}\theta' + 2\pi r A^{-1}M, q - \frac{r}{r'}\xi' + 2\pi r(\mathcal{B}^{-1})^t N)})] \\ &= [\mathcal{L}_{(r, A, (p + \frac{r}{r'}\theta' + 2\pi r A^{-1}M) - \frac{r}{r'}\theta', (q - \frac{r}{r'}\xi' + 2\pi r(\mathcal{B}^{-1})^t N) + \frac{r}{r'}\xi')}] \\ &= [\mathcal{L}_{(r, A, p + 2\pi r A^{-1}M, q + 2\pi r(\mathcal{B}^{-1})^t N)}] \\ &= [\mathcal{L}_{(r, A, p, q)}]. \end{aligned}$$

This completes the proof. \square

Acknowledgment

I would like to thank Hiroshige Kajiura for various advices in writing this paper. I also would like to thank Masahiro Futaki and Atsushi Takahashi for helpful comments. Finally, I am grateful to the referee for useful suggestions. This work was supported by Grant-in-Aid for JSPS Research Fellow 18J10909.

References

- [1] M. Abouzaid, I. Smith, Homological mirror symmetry for the four-torus, *Duke Mathematical Journal*, 152.3 (2010), 373-440.
- [2] D. Arinkin, A. Polishchuk, Fukaya category and Fourier transform, *AMS IP STUDIES IN ADVANCED MATHEMATICS*, 2001, 23 : 261-274.
- [3] A. Bondal and M. Kapranov, Enhanced triangulated categories, *Math. USSR Sbornik* 70:93-107, 1991.

- [4] K. Fukaya, Morse homotopy, A^∞ -category, and Floer homologies, In: Proceedings of GARC Workshop on Geometry and Topology '93 (Seoul, 1993). Lecture Notes in Series, vol. 18, pp. 1-102. Seoul Nat. Univ., Seoul (1993).
- [5] K. Fukaya, Mirror symmetry of abelian varieties and multi-theta functions, *J. Alg. Geom.* 11, 393-512 (2002).
- [6] H. Kajiura, On some deformation of Fukaya categories. *Symplectic, Poisson, and Noncommutative Geometry*, 93-130, MSRI Publ. 62, Cambridge Univ. Press, New York, 2014.
- [7] K. Kobayashi, On exact triangles consisting of stable vector bundles on tori, *Differential Geometry and its Applications* 53 (2017) 268-292, arXiv : mathDG/1610.02821.
- [8] K. Kobayashi, Geometric interpretation for exact triangles consisting of projectively flat bundles on higher dimensional complex tori, arXiv : mathDG/1705.04007.
- [9] K. Kobayashi, On exact triangles consisting of projectively flat bundles on complex tori, doctoral thesis at Chiba University.
- [10] K. Kobayashi, Remarks on the homological mirror symmetry for tori, arXiv : mathDG/2004.05621.
- [11] S. Kobayashi, *Differential Geometry of Complex Vector Bundles*, Princeton University Press, 1987.
- [12] M. Kontsevich, Homological algebra of mirror symmetry, In *Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Zürich, 1994)*, Birkhäuser, 1995, pages 120-139, arXiv : math.AG/9411018.
- [13] M. Kontsevich, Y. Soibelman, Homological mirror symmetry and torus fibrations. In *Symplectic geometry and mirror symmetry (Seoul, 2000)*, pages 203-263. World Sci. Publishing, River Edge, NJ, 2001. math.SG/0011041.
- [14] N. C. Leung, S.-T. Yau, E. Zaslow, From special Lagrangian to Hermitian-Yang-Mills via Fourier-Mukai transform, *Adv. Theor. Math. Phys.* 4:13191341, 2000.
- [15] Y. Matsushima, Heisenberg groups and holomorphic vector bundles over a complex torus, *Nagoya Math. J.* Vol. 61 (1976), 161-195.
- [16] S. Mukai, Semi-homogeneous vector bundles on an abelian variety, *J. Math. Kyoto Univ.* 18 (1978), no. 2, 239-272.
- [17] D. O. Orlov, Remarks on generators and dimensions of triangulated categories, *Moscow Mathematical Journal* 9.1 (2009) : 143-149, arXiv : math.AG/0804.1163.

- [18] A. Polishchuk, A_∞ -structures on an elliptic curve, *Comm. Math. Phys.* 247, 527 (2004), arXiv : math.AG/0001048.
- [19] A. Polishchuk, E. Zaslow, Categorical mirror symmetry : the elliptic curve, *Adv. Theor. Math. Phys.* 2, 443-470 (1998), arXiv : math.AG/9801119.
- [20] A. Strominger, S.-T. Yau, and E. Zaslow, Mirror Symmetry is T-duality, *Nucl. Phys. B*, 479:243-259, 1996.