

On Some Problems Related to a Simplex and a Ball

Mikhail Nevskii*

May 5, 2019

Abstract

Let C be a convex body and let S be a nondegenerate simplex in \mathbb{R}^n . Denote by $\xi(C; S)$ the minimal $\tau > 0$ such that C is a subset of the simplex τS . By $\alpha(C; S)$ we mean the minimal $\tau > 0$ such that C is contained in a translate of τS . Earlier the author has proved the equalities $\xi(C; S) = (n+1) \max_{1 \leq j \leq n+1} \max_{x \in C} (-\lambda_j(x)) + 1$ (if $C \not\subset S$),

$\alpha(C; S) = \sum_{j=1}^{n+1} \max_{x \in C} (-\lambda_j(x)) + 1$. Here λ_j are linear functions called the basic Lagrange polynomials corresponding to S . In his previous papers, the author has investigated these formulae if $C = [0, 1]^n$. The present paper is related to the case when C coincides with the unit Euclidean ball $B_n = \{x : \|x\| \leq 1\}$, where $\|x\| = \left(\sum_{i=1}^n x_i^2 \right)^{1/2}$. We establish various relations for $\xi(B_n; S)$ and $\alpha(B_n; S)$, as well as we give their geometric interpretation.

Keywords: n -dimensional simplex, n -dimensional ball, homothety, absorption index

1 Preliminaries

Everywhere further $n \in \mathbb{N}$. An element $x \in \mathbb{R}^n$ is written in the form $x = (x_1, \dots, x_n)$. By definition,

$$\|x\| = \sqrt{(x, x)} = \left(\sum_{i=1}^n x_i^2 \right)^{1/2},$$

*Department of Mathematics, P.G. Demidov Yaroslavl State University, Sovetskaya str., 14, Yaroslavl, 150003, Russia orcid.org/0000-0002-6392-7618 mnevsk55@yandex.ru

$$B(x^{(0)}; \varrho) := \{x \in \mathbb{R}^n : \|x - x^{(0)}\| \leq \varrho\} \quad (x^{(0)} \in \mathbb{R}^n, \varrho > 0),$$

$$B_n := B(0; 1), \quad Q_n := [0, 1]^n, \quad Q'_n := [-1, 1]^n.$$

Let C be a convex body in \mathbb{R}^n . Denote by τC the image of C under the homothety with center of homothety in the center of gravity of C and ratio of homothety τ . For an n -dimensional nondegenerate simplex S , consider the value $\xi(C; S) := \min\{\sigma \geq 1 : C \subset \sigma S\}$. We call this number the *absorption index of S with respect to C* . Define $\alpha(C; S)$ as minimal $\tau > 0$ such that convex body C is a subset of the simplex τS . By $\text{ver}(G)$ we mean the set of vertices of convex polytope G .

Let $x^{(j)} = (x_1^{(j)}, \dots, x_n^{(j)})$, $1 \leq j \leq n+1$, be the vertices of simplex S . The matrix

$$\mathbf{A} := \begin{pmatrix} x_1^{(1)} & \dots & x_n^{(1)} & 1 \\ x_1^{(2)} & \dots & x_n^{(2)} & 1 \\ \vdots & \vdots & \vdots & \vdots \\ x_1^{(n+1)} & \dots & x_n^{(n+1)} & 1 \end{pmatrix}$$

is nondegenerate. By definition, put $\mathbf{A}^{-1} = (l_{ij})$. Linear polynomials $\lambda_j(x) = l_{1j}x_1 + \dots + l_{nj}x_n + l_{n+1,j}$ whose coefficients make up the columns of \mathbf{A}^{-1} have the property $\lambda_j(x^{(k)}) = \delta_j^k$, where δ_j^k is the Kronecker δ -symbol. We call λ_j the *basic Lagrange polynomials corresponding to S* . The numbers $\lambda_j(x)$ are barycentric coordinates of a point $x \in \mathbb{R}^n$ with respect to S . Simplex S is given by the system of linear inequalities $\lambda_j(x) \geq 0$. For more details about λ_j , see [3; Chapter 1].

The equality $\xi(C; S) = 1$ is equivalent to the inclusion $C \subset S$. If $C \not\subset S$, then

$$\xi(C; S) = (n+1) \max_{1 \leq j \leq n+1} \max_{x \in C} (-\lambda_j(x)) + 1. \quad (1)$$

(the proof was given in [2]; see also [3; § 1.3]). The relation

$$\max_{x \in C} (-\lambda_1(x)) = \dots = \max_{x \in C} (-\lambda_{n+1}(x)) \quad (2)$$

holds true if and only if the simplex $\xi(C; S)S$ is circumscribed around convex body C . In the case $C = Q_n$ equality (1) can be reduced to the form

$$\xi(Q_n; S) = (n+1) \max_{1 \leq j \leq n+1} \max_{x \in \text{ver}(Q_n)} (-\lambda_j(x)) + 1$$

and (2) is equivalent to the relation

$$\max_{x \in \text{ver}(Q_n)} (-\lambda_1(x)) = \dots = \max_{x \in \text{ver}(Q_n)} (-\lambda_{n+1}(x)). \quad (3)$$

For any C and S , we have $\xi(C; S) \geq \alpha(C; S)$. The equality $\xi(C; S) = \alpha(C; S)$ holds only in the case when the simplex $\xi(C; S)S$ is circumscribed around convex body C . This is equivalent to (2) and also to (3) when $C = Q_n$.

It was proved in [4] (see also [3; § 1.4]) that

$$\alpha(C; S) = \sum_{j=1}^{n+1} \max_{x \in C} (-\lambda_j(x)) + 1. \quad (4)$$

If $C = Q_n$, then this formula can be written in rather more geometric way:

$$\alpha(Q_n; S) = \sum_{i=1}^n \frac{1}{d_i(S)}. \quad (5)$$

Here $d_i(S)$ is the i th axial diameter of simplex S , i.e., the length of a longest segment in S parallel to the i th coordinate axis. Equality (5) was obtained in [11]. When $S \subset Q_n$, we have $d_i(S) \leq 1$. Therefore, for these simplices, (5) gives

$$\xi(Q_n; S) \geq \alpha(Q_n; S) = \sum_{i=1}^n \frac{1}{d_i(S)} \geq n. \quad (6)$$

Earlier the author established the equality

$$\frac{1}{d_i(S)} = \frac{1}{2} \sum_{j=1}^{n+1} |l_{ij}| \quad (7)$$

(see [2]). Being combined together, (5) and (7) yield

$$\alpha(Q_n; S) = \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^{n+1} |l_{ij}|. \quad (8)$$

Note that $\alpha(C; S)$ is invariant under parallel translation of the sets and for $\tau > 0$ we have $\alpha(\tau C; S) = \tau \alpha(C; S)$. Since $Q'_n = [-1, 1]^n$ is a translate of the cube $2Q_n$, after replacing Q_n with Q'_n we obtain from (8) an even simpler formula:

$$\alpha(Q'_n; S) = \sum_{i=1}^n \sum_{j=1}^{n+1} |l_{ij}|. \quad (9)$$

Let us define the value

$$\xi_n := \min\{\xi(Q_n; S) : S \text{ — } n\text{-мерный симплекс, } S \subset Q_n, \text{vol}(S) \neq 0\}.$$

Various estimates of ξ_n were obtained first by the author and then by the author and A. Yu. Ukhalov (e.g., see papers [1], [2], [5], [6], [7], [8], [12]

and book [3]). Always $n \leq \xi_n < n + 1$. Nowadays the precise values of ξ_n are known for $n = 2, 5, 9$ and also for the infinite set of odd n 's for any of which there exists an Hadamard matrix of order $n + 1$. If $n \neq 2$, then every known value of ξ_n is equal to n , whereas $\xi_2 = 1 + \frac{3\sqrt{5}}{5} = 2.34 \dots$. Still remains unknown is there exist an even n with the property $\xi_n = n$. There are some other open problems concerning the numbers ξ_n .

In this article, we will discuss the analogues of the above characteristics for a simplex and an Euclidean ball. Replacing a cube with a ball makes many questions much more simpler. However, geometric interpretation of general results has a certain interest also in this particular case. Besides, we will note some new applications of the basic Lagrange polynomials.

Numerical characteristics connecting simplices and subsets of \mathbb{R}^n have applications for obtaining various estimates in polynomial interpolation of functions defined on multidimensional domains. This approach and the corresponding analytic methods in details were described in [3]. Lately these questions have been managed to study also by computer methods (see, e. g., [5], [6], [8], [12]).

2 The value $\alpha(B_n; S)$

The *inradius* of an n -dimensional simplex S is the maximum of the radii of balls contained within S . The center of this unique maximum ball is called the *incenter* of S . The boundary of the maximum ball is a sphere that has a single common point with each $(n - 1)$ -dimensional face of S . By the *circumradius* of S we mean the minimum of the radii of balls containing S . The boundary of this unique minimal ball does not necessarily contain all the vertices of S . Namely, this is only when the center of the minimal ball lies inside the simplex.

The inradius r and the circumradius R of a simplex S satisfy the so-called *Euler inequality*

$$R \geq nr. \quad (10)$$

Equality in (10) takes place if and only if S is a regular simplex. Concerning the proofs of the Euler inequality, its history and generalizations, see, e. g., [10], [13], [14].

In connection with (10), let us remark an analogue to the following property being true for parallelotopes (see [11], [3; § 1.8]). *Let S be a nondegenerate simplex and let D, D^* be parallelotopes in \mathbb{R}^n . Suppose D^* is a homothetic copy of D with ratio $\tau > 1$. If $D \subset S \subset D^*$, then $\tau \geq n$.* This proposition holds true also for balls. In fact, the Euler inequality is equivalent to the following statement. *Suppose B is a ball with radius r_1 and B^* is a ball with*

radius r_2 . If $B \subset S \subset B^*$, then $r_1 \leq nr_2$. Equality takes place if and only if S is a regular simplex inscribed into B^* and B is the ball inscribed into S . Another equivalent form of these propositions is given by Theorem 2 (see the note after the proof of this theorem).

Let $x^{(1)}, \dots, x^{(n+1)}$ be the vertices and let $\lambda_1, \dots, \lambda_{n+1}$ be the basic Lagrange polynomials of a nondegenerate simplex $S \subset \mathbb{R}^n$ (see Section 1). In what follows Γ_j is the $(n-1)$ -dimensional hyperplane given by the equation $\lambda_j(x) = 0$, by Σ_j we mean the $(n-1)$ -dimensional face of S contained in Γ_j , symbol h_j denotes the height of S conducted from the vertex $x^{(j)}$ onto Γ_j , and r denotes the inradius of S . Define σ_j as $(n-1)$ -measure of Σ_j and put $\sigma := \sum_{j=1}^{n+1} \sigma_j$. Consider the vector $a_j := \{l_{1j}, \dots, l_{nj}\}$. This vector is orthogonal to Γ_j and directed into the subspace containing $x^{(j)}$. Obviously,

$$\lambda_j(x) = l_{1j}x_1 + \dots + l_{nj}x_n + l_{n+1,j} = (a_j, x) + l_{n+1,j} = (a_j, x) + \lambda_j(0).$$

Theorem 1. *The following equalities are true:*

$$\alpha(B_n; S) = \sum_{j=1}^{n+1} \left(\sum_{i=1}^n l_{ij}^2 \right)^{1/2}, \quad (11)$$

$$\alpha(B_n; S) = \sum_{j=1}^{n+1} \frac{1}{h_j}, \quad (12)$$

$$\alpha(B_n; S) = \frac{1}{r}, \quad (13)$$

$$\alpha(B_n; S) = \frac{\sigma}{n \text{vol}(S)}. \quad (14)$$

Proof. Let us obtain these pairwise-equivalent equalities from the top up to the bottom. First we note that

$$\alpha(B_n; S) = \sum_{j=1}^{n+1} \max_{x \in B} (-\lambda_j(x)) + 1. \quad (15)$$

Formula (15) is the particular case of (4) in the situation $C = B_n$. By the Cauchy inequality,

$$- \|a_j\| \|x\| \leq (a_j, x) \leq \|a_j\| \|x\|, \quad (16)$$

$$-\|a_j\| \|x\| - \lambda_j(0) \leq -\lambda_j(x) \leq \|a_j\| \|x\| - \lambda_j(0).$$

Both the upper and the lower bounds in (16) are reachable. This gives

$$\max_{x \in B_n}(-\lambda_j(x)) = \max_{\|x\| \leq 1}(-\lambda_j(x)) = \|a_j\| - \lambda_j(0).$$

Therefore,

$$\alpha(B_n; S) = \sum_{j=1}^{n+1} \max_{x \in B_n}(-\lambda_j(x)) + 1 = \sum_{j=1}^{n+1} \|a_j\| - \sum_{j=1}^{n+1} \lambda_j(0) + 1 = \sum_{j=1}^{n+1} \left(\sum_{i=1}^n l_{ij}^2 \right)^{1/2}.$$

We made use of the equality $\sum_{j=1}^{n+1} \lambda_j(0) = 1$. Since $\lambda_j(x^{(j)}) = 1$, we have

$$h_j = \text{dist}(x^{(j)}; \Gamma_j) = \frac{|\lambda_j(x^{(j)})|}{\|a_j\|} = \frac{1}{\|a_j\|} = \frac{1}{\left(\sum_{i=1}^n l_{ij}^2 \right)^{1/2}}.$$

Consequently,

$$\alpha(B_n; S) = \sum_{j=1}^{n+1} \left(\sum_{i=1}^n l_{ij}^2 \right)^{1/2} = \sum_{j=1}^{n+1} \frac{1}{h_j}.$$

We have obtained both (11) and (12).

Let us prove (13). The ball B_n is a subset of a translate of the simplex $\alpha(B_n; S)S$. This means that a translate of the ball $\frac{1}{\alpha(B_n; S)}B_n$ is contained in S . Since the maximum of the radii of balls being contained in S is equal to r , holds true $\frac{1}{\alpha(B_n; S)} \leq r$, i. e., $\alpha(B_n; S) \geq \frac{1}{r}$. To obtain the inverse inequality, denote by B' a ball of radius r inscribed into S . Then the ball $B_n = \frac{1}{r}B'$ is a subset of some translate of $\frac{1}{r}S$. Using the definition of $\alpha(B_n; S)$ we can write $\alpha(B_n; S) \leq \frac{1}{r}$. So, we have $\alpha(B_n; S) = \frac{1}{r}$.

Finally, in order to establish (14), it is sufficient to utilize (13) and the formula $\text{vol}(S) = \frac{1}{n}\sigma r$. The latter equality one can obtain from an ordinary formula for the volume of a simplex after subdividing S onto $n+1$ simplices in such a way that j th of these simplices has a vertex in the center of the inscribed ball and is supported on Σ_j . \square

Corollary 1. *We have*

$$\frac{1}{r} = \sum_{j=1}^{n+1} \frac{1}{h_j}.$$

For proving, it is sufficient to apply (12) and (13). It seems to be interesting that this geometric relation (which evidently can be obtained also in

a direct way) occurs to be equivalent to general formula for $\alpha(C; S)$ in the particular case when a convex body C coincide with an Euclidean unit ball.

Corollary 2. *The inradius r and the incenter z of a simplex S can be calculated by the following formulae:*

$$r = \frac{1}{\sum_{j=1}^{n+1} \left(\sum_{i=1}^n l_{ij}^2 \right)^{1/2}}, \quad (17)$$

$$z = \frac{1}{\sum_{j=1}^{n+1} \left(\sum_{i=1}^n l_{ij}^2 \right)^{1/2}} \sum_{j=1}^{n+1} \left(\sum_{i=1}^n l_{ij}^2 \right)^{1/2} x^{(j)}. \quad (18)$$

The tangent point of the ball $B(z; r)$ and facet Σ_k has the form

$$y^{(k)} = \frac{1}{\sum_{j=1}^{n+1} \left(\sum_{i=1}^n l_{ij}^2 \right)^{1/2}} \left[\sum_{j=1}^{n+1} \left(\sum_{i=1}^n l_{ij}^2 \right)^{1/2} x^{(j)} - \frac{1}{\left(\sum_{i=1}^n l_{ik}^2 \right)^{1/2}} (l_{1k}, \dots, l_{nk}) \right]. \quad (19)$$

Proof. Equality (17) follows immediately from (11) and (13). To obtain (18), let us remark that

$$r = \text{dist}(z; \Gamma_j) = \frac{|\lambda_j(z)|}{\|a_j\|}.$$

Since z lies inside S , each barycentric coordinate of this point $\lambda_j(z)$ is positive, i. e., $\lambda_j(z) = r\|a_j\|$. Consequently,

$$z = \sum_{j=1}^{n+1} \lambda_j(z) x^{(j)} = r \sum_{j=1}^{n+1} \|a_j\| x^{(j)}.$$

This coincides with (18). Finally, since vector $a_k = \{l_{1k}, \dots, l_{nk}\}$ is orthogonal to Σ_k and is directed from this facet inside the simplex, a unique common point of $B(z; r)$ and Σ_k has the form

$$y^{(k)} = z - \frac{r}{\|a_k\|} a_k = r \left(\sum_{j=1}^{n+1} \|a_j\| x^{(j)} - \frac{1}{\|a_k\|} a_k \right).$$

The latter is equivalent to (19). □

It is interesting to compare (11) with the formula (9) for $\alpha(Q'_n; S)$. Since B_n is a subset of the cube $Q'_n = [-1, 1]^n$, we have $\alpha(B_n; S) \leq \alpha(Q'_n; S)$. Analytically, this also follows from the estimate

$$\left(\sum_{i=1}^n l_{ij}^2 \right)^{1/2} \leq \sum_{i=1}^n |l_{ij}|.$$

For arbitrary $x^{(0)}$ and $\varrho > 0$, the number $\alpha(B(x^{(0)}; \varrho); S)$ can be calculated with the use of Theorem 1 and the equality $\alpha(B(x^{(0)}; \varrho); S) = \varrho \alpha(B_n; S)$.

If $S \subset Q_n$, then all the axial diameters $d_i(S)$ do not exceed 1 and (5) immediately gives $\alpha(Q_n; S) \geq n$. Moreover, the equality $\alpha(Q_n; S) = n$ holds when and only when each $d_i(S) = 1$. The following proposition expresses the analogues of these properties for simplices contained in a ball.

Theorem 2. *If $S \subset B_n$, then $\alpha(B_n; S) \geq n$. The equality $\alpha(B_n; S) = n$ holds true if and only if S is a regular simplex inscribed into B_n .*

Proof. By the definition of $\alpha(B_n; S)$, the ball B_n is contained in a translate of the simplex $\alpha(B_n; S)S$. Hence, some translate B' of the ball $\frac{1}{\alpha(B_n; S)}B_n$ is a subset of S . So, we have the inclusions $B' \subset S \subset B_n$. Since the radius of B' is equal to $\frac{1}{\alpha(B_n; S)}$, the inradius r and the circumradius R of S satisfy the inequalities $\frac{1}{\alpha(B_n; S)} \leq r$, $R \leq 1$. Making use of the Euler inequality $R \geq nr$, we can write

$$\frac{1}{\alpha(B_n; S)} \leq r \leq \frac{R}{n} \leq \frac{1}{n}. \quad (20)$$

Therefore, $\alpha(B_n; S) \geq n$.

The equality $\alpha(B_n; S) = n$ means that the left-hand value in (20) coincides with the right-hand one. Thus, all the inequalities in this chain turn into equalities. We obtain $R = 1$, $r = \frac{1}{n}$. Since in this case the Euler inequality (10) also becomes an equality, S is a regular simplex inscribed into B_n . Conversely, if S is a regular simplex inscribed into B_n , then $r = \frac{1}{n}$, i. e., $\alpha(B_n; S) = \frac{1}{r} = n$. \square

We see that Theorem 2 follows from the Euler inequality (10). In fact, these statements are equivalent. Indeed, suppose S is an arbitrary n -dimensional simple, r is the inradius and R is the circumradius of S . Let us denote by B the ball containing S and having radius R . Then some translate S' of the simplex $\frac{1}{R}S$ is contained in B_n . By Theorem 1, $\alpha(B_n; S')$ is the inverse to the inradius of S' , i. e., is equal to $\frac{R}{r}$. Now assume that Theorem 2 is true. Let us apply this theorem to the simplex $S' \subset B_n$. This gives $\alpha(B_n; S') = \frac{R}{r} \geq n$ and we have (10). Finally, if $R = nr$, then $\alpha(B_n; S') = n$. From Theorem 2 we obtain that both S' and S are regular simplices.

It follows from (6) that the minimum value of $\alpha(Q_n; S)$ for $S \subset Q_n$ also is equal to n . This minimal value corresponds to those and only those $S \subset Q_n$ for which every axial diameter $d_i(S)$ is equal to 1. The noted property is fulfilled for the maximum volume simplices in Q_n (see [3]), but not for the only these simplices, if $n > 2$.

3 The value $\xi(B_n; S)$

In this section, we will obtain the computational formula for the absorption index of a simplex S with respect to an Euclidean ball. We use the previous denotations.

Theorem 3. *Suppose S is a nondegenerate simplex in \mathbb{R}^n , $x^{(0)} \in \mathbb{R}^n$, $\varrho > 0$. If $B(x^{(0)}; \varrho) \not\subset S$, we have*

$$\xi(B(x^{(0)}; \varrho); S) = (n+1) \max_{1 \leq j \leq n+1} \left[\varrho \left(\sum_{i=1}^n l_{ij}^2 \right)^{1/2} - \sum_{i=1}^n l_{ij} x_i^{(0)} - l_{n+1,j} \right] + 1. \quad (21)$$

In particular, if $B_n \not\subset S$, then

$$\xi(B_n; S) = (n+1) \max_{1 \leq j \leq n+1} \left[\left(\sum_{i=1}^n l_{ij}^2 \right)^{1/2} - l_{n+1,j} \right] + 1. \quad (22)$$

Proof. Let us apply the general formula (1) in the case $C = B(x^{(0)}; \varrho)$. The Cauchy inequality yields

$$-\|a_j\| \|x - x^{(0)}\| \leq (a_j, x - x^{(0)}) \leq \|a_j\| \|x - x^{(0)}\|. \quad (23)$$

If $\|x - x^{(0)}\| \leq \varrho$, we see that

$$\begin{aligned} -\varrho \|a_j\| &\leq (a_j, x) - (a_j, x^{(0)}) \leq \varrho \|a_j\|, \\ -\lambda_j(x) &= -(a_j, x) - l_{n+1,j} \leq \varrho \|a_j\| - (a_j, x^{(0)}) - l_{n+1,j}. \end{aligned}$$

Since both the upper and the lower bounds in (23) are reachable,

$$\max_{\|x - x^{(0)}\| \leq \varrho} (-\lambda_j(x)) = \varrho \left(\sum_{i=1}^n l_{ij}^2 \right)^{1/2} - \sum_{i=1}^n l_{ij} x_i^{(0)} - l_{n+1,j}.$$

It follows that

$$\xi(B(x^{(0)}; \varrho); S) = (n+1) \max_{1 \leq j \leq n+1, \|x - x^{(0)}\| \leq \varrho} (-\lambda_j(x)) + 1 =$$

$$= (n+1) \max_{1 \leq j \leq n+1} \left[\varrho \left(\sum_{i=1}^n l_{ij}^2 \right)^{1/2} - \sum_{i=1}^n l_{ij} x_i^{(0)} - l_{n+1,j} \right] + 1,$$

and we obtain (21). Equality (22) appears from (21) for $x^{(0)} = 0, \varrho = 1$. \square

4 The equality $\beta_n = n$. Commentaries

Theorem 4. *If $S \subset B_n$, then $\xi(B_n; S) \geq n$. The equality $\xi(B_n; S) = n$ takes place if and only if S is a regular simplex inscribed into B_n .*

Proof. The statement immediately follows from Theorem 2 and the inequality $\xi(B_n; S) \geq \alpha(B_n; S)$. We give here also a direct proof without applying the Euler inequality that was used to obtain the estimate $\alpha(B_n; S) \geq n$.

First let S be a regular simplex inscribed into B_n . Then $\alpha(B_n; S) = n$ and the inradius of S is equal to $\frac{1}{n}$. Since the simplex $\xi(B_n; S)S$ is circumscribed around B_n , we have the equalities $\xi(S; B_n) = \alpha(S; B_n) = n$ and also relation (2) with $C = B_n$. It follows from (1) that for any $j = 1, \dots, n+1$

$$\max_{x \in B_n} (-\lambda_j(x)) = \frac{n-1}{n+1},$$

where λ_j are the basic Lagrange polynomials related to S .

Now suppose simplex S is contained in B_n but is not regular or is not inscribed into the ball. Denote the Lagrange polynomials of this simplex by μ_j . There exist a regular simplex S^* inscribed into B_n and an integer k such that S is contained in the strip $0 \leq \lambda_k(x) \leq 1$, the k th $(n-1)$ -dimensional faces of S and S^* are parallel, and S has not any common points with at least one of the boundary hyperplanes of this strip. Here λ_j are the basic Lagrange polynomials of S^* . The vertex $x^{(k)}$ of the simplex S^* does not lie in its k th facet. Assume u is a point of the boundary of B_n most distant from $x^{(k)}$. Then u is the maximum point of polynomial $-\lambda_k(x)$, i.e., $-\lambda_j(u) = \frac{n-1}{n+1}$. Consider the straight line connecting $x^{(k)}$ and u . Denote by y, z and t the intersection points of this line and pairwise parallel hyperplanes $\mu_k(x) = 1$, $\mu_k = 0$ and $\lambda_k(x) = 0$ respectively. We have

$$\|x^{(k)} - t\| \geq \|y - z\|, \quad \|t - u\| \leq \|z - u\|. \quad (24)$$

At least one of these inequalities is fulfilled in the strict form. The linearity of the basic Lagrange polynomials means that

$$\frac{\mu_k(z) - \mu_k(u)}{\mu_k(y) - \mu_k(z)} = \frac{\|z - u\|}{\|y - z\|}, \quad \frac{\lambda_k(t) - \lambda_k(u)}{\lambda_k(x^{(k)}) - \lambda_k(t)} = \frac{\|t - u\|}{\|x^{(k)} - t\|}.$$

Since $\mu_k(y) = 1$, $\mu_k(z) = 0$, $\lambda_k(x^{(k)}) = 1$, and $\lambda_k(t) = 0$, we get

$$-\mu_k(u) = \frac{\|z - u\|}{\|y - z\|} > \frac{\|t - u\|}{\|x^{(k)} - t\|} = -\lambda_k(u) = \frac{n-1}{n+1}.$$

We made use of (24) and took into account that at least one of the inequalities is strict. The application of (1) yields

$$\xi(B_n; S) = (n+1) \max_{1 \leq j \leq n+1} \max_{x \in B_n} (-\mu_j(x)) + 1 \geq (n+1)(-\mu_k(u)) + 1 > n.$$

Thus, if S is not regular simplex inscribed into B_n , then $\xi(B_n; S) > n$.

We see that each simplex $S \subset B_n$ satisfies the estimate $\xi(B_n; S) \geq n$. The equality takes place if and only if S is a regular simplex inscribed into B_n . \square

By analogy with the value $\xi_n = \min\{\xi(Q_n; S) : S \subset Q_n\}$ defined through the unit cube, let us introduce the similar numerical characteristic given by the unit ball:

$$\beta_n := \min\{\xi(B_n; S) : S - n\text{-мерный симплекс}, S \subset B_n, \text{vol}(S) \neq 0\}.$$

Many problems concerning ξ_n yet have not been solved. For example, $\xi_2 = 1 + \frac{3\sqrt{2}}{5}$ still remains the only accurate value of ξ_n for even n ; moreover, this value was discovered in a rather difficult way (see [3; Chapter 2]). Compared to ξ_n the problem on numbers β_n turns out to be trivial.

Corollary 3. *For any n , we have $\beta_n = n$. The only simplex $S \subset B_n$ extremal with respect to β_n is an arbitrary regular simplex inscribed into B_n .*

Proof. It is sufficient to apply Theorem 4. \square

The technique developed for a ball makes it possible to illustrate some results having been earlier got for a cube. Here we note a proof of the following known statement which differs from the proofs given in [3; §3.2] and [12].

Corollary 4. *If there exists an Hadamard matrix of order $n+1$, then $\xi_n = n$.*

Proof. It is known (see, e. g., [9]) that for these and only these n we can inscribe into Q_n a regular simplex S so that all the vertices of S will coincide with vertices of the cube. Let us denote by B the ball with radius $\frac{\sqrt{n}}{2}$ having the center in center of the cube. Clearly, Q_n is inscribed into B , therefore, the simplex is inscribed into the ball as well. Since S is regular, by Theorem 4 and by similarity reasons, we have $\xi(B; S) = n$. The inclusion $Q_n \subset B$ means that $\xi(Q_n; S) \leq \xi(B; S)$, i. e. $\xi(Q_n; S) \leq n$. From (6) it follows that the inverse

inequality $\xi(Q_n; S) \geq n$ is also true. Hence, $\xi(Q_n; S) = n$. Simultaneously (6) gives $\xi_n = \xi(Q_n; S) = n$. \square

This argument is based on the following fact: if S is a regular simplex with the vertices in vertices of Q_n , then the simplex nS absorbs not only the cube Q_n but also the ball B circumscribed around the cube. The corresponding absorption index n is the minimum possible both for the cube and the ball. In addition, we mention the following property.

Corollary 5. *Assume that $S \subset Q_n \subset nS$ and simplex S is not regular. Then $B \not\subset nS$.*

Proof. The inclusion $B \subset nS$ implies that $\xi(B; S) = n$. This way S is a regular simplex inscribed into the ball B . But since this is not so, B is not a subset of nS . \square

Simplices satisfying the condition of Corollary 5 exist at least for $n = 3, 5$, and 9 (see [12]).

The relations (6) mean that always $\xi_n \geq n$. Since $\xi_2 = 1 + \frac{3\sqrt{5}}{5} > 2$, there exist n 's such that $\xi_n > n$. Besides the cases when $n + 1$ is an Hadamard number, the equality $\xi_n = n$ is established for $n = 5$ and $n = 9$ (the extremal simplices in \mathbb{R}^5 and \mathbb{R}^9 are given in [12]). For all such dimensions holds true $\xi_n = \beta_n$, i. e., with respect to the minimum absorption index of an internal simplex, both the convex bodies, an n -dimensional cube and an n -dimensional ball, have the same behaviour.

The equality $\xi_n = n$ is equivalent to the existence of simplices satisfying the inclusions $S \subset Q_n \subset nS$. Some properties of such simplices (e. g., the fact that the center of gravity of S coincides with the center of the cube; see [7]) are similar to the properties of regular simplices inscribed into the ball. However, the problem to describe the set of all dimensions where exist those simplices, seems to be very difficult and nowadays is far from solution.

References

1. Nevskij, M. V., On a certain relation for the minimal norm of an interpolational projection, *Model. Anal. Inform. Sist.*, 2009, vol. 16, no. 1, pp. 24–43 (in Russian).
2. Nevskii, M. V. On a property of n -dimensional simplices, *Math. Notes*, 2010, vol. 87, no. 4, pp. 543–555.
3. Nevskii, M. V., *Geometricheskie ocenki v polinomialnoi interpoljacii* (Geometric Estimates in Polynomial Interpolation), Yaroslavl': Yarosl. Gos. Univ., 2012 (in Russian). .

4. Nevskii, M. V., On the minimal positive homothetic image of a simplex containing a convex body, *Math. Notes*, 2013, vol. 93, no. 3–4, pp. 470–478.
5. Nevskii, M. V., and Ukhalov, A. Yu., On numerical characteristics of a simplex and their estimates, *Model. Anal. Inform. Sist.*, 2016, vol. 23, no. 5, pp. 603–619 (in Russian). English transl.: *Aut. Control Comp. Sci.*, 2017, vol. 51, no. 7, pp. 757–769.
6. Nevskii, M. V., and Ukhalov, A. Yu., New estimates of numerical values related to a simplex, *Model. Anal. Inform. Sist.*, 2017, vol. 24, no. 1, pp. 94–110 (in Russian). English transl.: *Aut. Control Comp. Sci.*, 2017, vol. 51, no. 7, pp. 770–782.
7. Nevskii, M. V., and Ukhalov, A. Yu., On n -dimensional simplices satisfying inclusions $S \subset [0, 1]^n \subset nS$, *Model. Anal. Inform. Sist.*, 2017, vol. 24, no. 5, pp. 578–595 (in Russian). English transl.: *Aut. Control Comp. Sci.*, 2018, vol. 52, no. 7, pp. 667–679.
8. Nevskii, M. V., and Ukhalov, A. Yu., On minimal absorption index for an n -dimensional simplex, *Model. Anal. Inform. Sist.*, 2018, vol. 25, no. 1, pp. 140–150 (in Russian). English transl.: *Aut. Control Comp. Sci.*, 2018, vol. 52, no. 7, pp. 680–687.
9. Hudelson, M., Klee, V., and Larman, D., Largest j -simplices in d -cubes: some relatives of the Hadamard maximum determinant problem, *Linear Algebra Appl.*, 1996, vol. 241–243, pp. 519–598.
10. Klamkin M. S., and Tsifinis G. A., Circumradius–inradius inequality for a simplex, *Mathematics Magazine*, 1979, vol. 52, no. 1, pp. 20–22.
11. Nevskii, M., Properties of axial diameters of a simplex, *Discr. Comput. Geom.*, 2011, vol. 46, no. 2, pp. 301–312.
12. Nevskii, M., and Ukhalov A., Perfect simplices in \mathbb{R}^5 , *Beitr. Algebra Geom.*, 2018, vol. 59, no. 3, pp. 501–521.
13. Yang S., and Wang J., Improvements of n -dimensional Euler inequality, *Journal of Geometry*, 1994, vol. 51, pp. 190–195.
14. Vince A., A simplex contained in a sphere, *Journal of Geometry*, 2008, vol. 89, no. 1–2, pp. 169–178.