

ORTHODIAGONAL ANTI-INVOLUTIVE KOKOTSAKIS POLYHEDRA

IVAN EROFEEV AND GRIGORY IVANOV

ABSTRACT. We study the properties of Kokotsakis polyhedra of orthodiagonal anti-involutive type. Stachel conjectured that a certain resultant connected to a polynomial system describing flexion of a Kokotsakis polyhedron must be reducible. Izmistiev [3] showed that a polyhedron of the orthodiagonal anti-involutive type is the only possible candidate to disprove Stachel's conjecture. We show that the corresponding resultant is reducible, thereby confirming the conjecture. We do it in two ways: by factorization of the corresponding resultant and providing a simple geometric proof. We describe the space of parameters for which such a polyhedron exists and show that this space is non-empty. We show that a Kokotsakis polyhedron of orthodiagonal anti-involutive type is flexible and give explicit parameterizations in elementary functions and in elliptic functions of its flexion.

1. INTRODUCTION

A *Kokotsakis polyhedron* is a polyhedral surface in \mathbb{R}^3 which consists of one n -gon (the *base*), n quadrilaterals attached to every side of the n -gon, and n triangles placed between each pair of two consecutive quadrilaterals. Such surfaces are rigid in general. However, there exist special classes of flexible surfaces, that is, ones that allow isometric deformations. Flexible Kokotsakis polyhedra were studied by several authors: in the PhD thesis of Antonios Kokotsakis and in [4]; Nawratil [5] studied flexible Kokotsakis polyhedra with triangular base; and in [8] some classes of flexible Kokotsakis polyhedra with quadrilateral base were found. The question of complete classification for the case $n = 4$ was open. Using Bricard's approach [1], Stachel and Nawratil [6, 7, 9] with the use of the following trigonometric substitution




FIGURE 1. Notation of dihedral angles

$$(1) \quad z = \tan \frac{\varphi}{2}, \quad w_1 = \tan \frac{\psi_1}{2}, \quad u = \tan \frac{\theta}{2}, \quad w_2 = \tan \frac{\psi_2}{2},$$

expressed the dependencies between the dihedral angles φ , ψ_1 , θ , ψ_2 (see Figure 1) as a polynomial system in z , w_1 , u , w_2 :

$$(2) \quad \begin{aligned} P_1(z, w_1) &= 0, & P_2(z, w_2) &= 0, \\ P_3(u, w_1) &= 0, & P_4(u, w_2) &= 0. \end{aligned}$$

They studied this system with respect to different factorizations of the resultant $R_{12}(w_1, w_2)$ of P_1 and P_2 as polynomials in z , and they classified all classes of flexible Kokotsakis polyhedra with quadrilateral base for reducible R_{12} . Stachel conjectured that there is no flexible Kokotsakis polyhedron whose system has the irreducible resultant.

G.I. was supported by the Swiss National Science Foundation grant 200021.179133.

Considering a complexified version of system (2), Izmistiev [3] classified all possible classes of flexible polyhedra disregarding reality and embeddability. In particular, from the results of [3] it follows that there is only one possible candidate for a flexible Kokotsakis polyhedron with the irreducible resultant R_{12} . He called elements of this class Kokotsakis polyhedra of *orthodiagonal anti-involutive type* (OAI).

In this paper we study the properties of OAI Kokotsakis polyhedra. We find an explicit parameterization in elementary functions of all solutions of the system (2) for this type and show that there are solutions which may be embedded in the real three-dimensional space. We prove that the resultant R_{12} is always reducible, which confirms Stachel’s conjecture. In addition, we provide a simple geometric proof of this fact in Theorem 6 that directly follows from the results of [3].

The rest of the article is organized as follows. We give the definition of an OAI Kokotsakis polyhedron in Section 2 together with algebraic formulas for coefficients of the polynomial systems required for further analysis. However, we postpone an explanation of their geometric meaning to Section 6. Section 3 is devoted to the description of all solutions of the system (2) for an OAI Kokotsakis polyhedron, which is the system (10). In Theorem 2 of this Section, we give an explicit parameterization in elementary functions of all branches of the solution. In other words, here we come up with a description of all possible flexions of an OAI polyhedron when all planar angles are given. In Section 4 we study a four parameter system that describes planar angles of all flexible OAI Kokotsakis polyhedra. We give a full parameterization of the solution set of this system in terms of the angles of the base quadrilateral and one additional parameter τ in Theorem 4. Summarizing the results, we introduce in Section 5 an algorithm for constructing a flexible OAI Kokotsakis polyhedron for a given base quadrilateral without right angles. In addition, we present results from the numerical screening of the space of parameters. In Section 6, we discuss the geometry behind the definitions and equations that we use, show that there is a nice flattening effect in the case of the orthodiagonal anti-involutive surface, and give a simple proof of Stachel’s conjecture using it. In Appendix 8 we prove some technical results.

Remark 1.1. We formulate all the theorems in such a manner that the reader can understand them right after reading the definitions of Section 2. This allows to use formulas from these theorems and construct a flexible polyhedron without going into the details of the proofs.

2. DEFINITIONS

In this section we give an algebraic description of a Kokotsakis polyhedron of the orthodiagonal anti-involutive type. We do not discuss the nature of the written equations here. A comprehensive explanations can be found in [3]. However, we give a brief explanation in Section 6, where we show that Stachel’s conjecture is a simple consequence of the results from [3].




FIGURE 2. Planar angles in a Kokotsakis polyhedron.

Vertices of a Kokotsakis polyhedron and the values of its planar angles at the interior vertices are denoted as in Figure 2. Clearly, it is only a neighborhood of the *base* face $A_1A_2A_3A_4$ that matters: replacing, say, the vertex B_1 by any other point on the half-line A_1B_1 does not affect the flexibility or rigidity of the polyhedron.

For each of the four interior vertices A_1, A_2, A_3, A_4 consider the intersection of the cone of adjacent faces with a unit sphere centered at the vertex. This yields four spherical quadrilaterals Q_i with side lengths $\alpha_i, \beta_i, \gamma_i, \delta_i$ in this cyclic order. Let $\varphi, \psi_1, \theta,$ and ψ_2 be the exterior dihedral angles at edges A_1A_2, A_2A_3, A_3A_4 and A_4A_1 , respectively. Clearly, it suffices to parameterize these dihedral angles in order to describe a flexion of a Kokotsakis polyhedron.




FIGURE 3. Spherical quadrilateral Q

2.1. Orthodiagonal quadrilaterals. We call a spherical quadrilateral *orthodiagonal* if its diagonals are orthogonal. A spherical quadrilateral is called *elliptic* if its planar angles satisfy $\alpha \pm \beta \pm \gamma \pm \delta \neq 0 \pmod{2\pi}$.

Let Q be a spherical elliptic orthodiagonal quadrilateral with side lengths and angles denoted as in Figure 3. Then the orthodiagonality of Q is equivalent to the following identity:

$$(3) \quad \cos \alpha \cos \gamma = \cos \beta \cos \delta$$

(see Lemma 8.1 below).

By definition put $z = \tan \frac{\varphi}{2}$, $w = \tan \frac{\psi}{2}$. Then (z, w) satisfy the following equation (see Lemma 4.13 in [3]):

$$(4) \quad (z^2 + \lambda)(w^2 + \mu) = \nu zw, \quad z, w \in \mathbb{R}P^1 = \mathbb{R} \cup \infty,$$

where λ, μ are the so-called *involution factors* defined as follows

$$(5) \quad \lambda := \begin{cases} \frac{\tan \delta + \tan \alpha}{\tan \delta - \tan \alpha}, & \text{if } \alpha \neq \frac{\pi}{2} \text{ or } \delta \neq \frac{\pi}{2}, \\ \frac{\cos \beta + \cos \gamma}{\cos \beta - \cos \gamma}, & \text{if } \alpha = \delta = \frac{\pi}{2}; \end{cases}$$

$$(6) \quad \mu := \begin{cases} \frac{\tan \delta + \tan \gamma}{\tan \delta - \tan \gamma}, & \text{if } \gamma \neq \frac{\pi}{2} \text{ or } \delta \neq \frac{\pi}{2}, \\ \frac{\cos \beta + \cos \alpha}{\cos \beta - \cos \alpha}, & \text{if } \gamma = \delta = \frac{\pi}{2}; \end{cases}$$

and ν is given by

$$(7) \quad \nu := \begin{cases} \frac{(\lambda-1)(\mu-1)}{\cos \delta}, & \text{if } \delta \neq \frac{\pi}{2}, \\ 2(\mu-1) \tan \alpha, & \text{if } \delta = \gamma = \frac{\pi}{2}, \\ 2(\lambda-1) \tan \gamma, & \text{if } \delta = \alpha = \frac{\pi}{2}. \end{cases}$$

The involution factors λ, μ and ν are well-defined real numbers different from 0.

2.2. Orthodiagonal anti-involutive type. A Kokotsakis polyhedron belongs to the *orthodiagonal anti-involutive type*, if its planar angles satisfy the following conditions.

1) All quadrilaterals Q_i are orthodiagonal and elliptic:

$$\cos \alpha_i \cos \gamma_i = \cos \beta_i \cos \delta_i, \quad \alpha_i \pm \beta_i \pm \gamma_i \pm \delta_i \neq 0 \pmod{2\pi}.$$

2) The involution factors at common vertices are opposite:

$$(8) \quad \lambda_1 = -\lambda_2, \quad \mu_1 = -\mu_4, \quad \mu_2 = -\mu_3, \quad \lambda_3 = -\lambda_4.$$

3) The involution factors λ_i, μ_i and ν_i satisfy the following system:

$$(9) \quad \frac{\nu_1^2}{\lambda_1 \mu_1} = \frac{\nu_3^2}{\lambda_3 \mu_3}, \quad \frac{\nu_2^2}{\lambda_2 \mu_2} = \frac{\nu_4^2}{\lambda_4 \mu_4}, \quad \frac{\nu_1^2}{\lambda_1 \mu_1} + \frac{\nu_2^2}{\lambda_2 \mu_2} = 16.$$

By (4) and (8), variables (z, w_1, u, w_2) of a Kokotsakis polyhedron of the orthodiagonal anti-involutive type satisfy the system

$$(10) \quad \begin{aligned} P_1 &= (z^2 + \lambda_1)(w_1^2 + \mu_1) - \nu_1 z w_1 = 0, \\ P_2 &= (z^2 - \lambda_1)(w_2^2 - \mu_3) - \nu_2 z w_2 = 0, \\ P_3 &= (u^2 + \lambda_3)(w_2^2 + \mu_3) - \nu_3 u w_2 = 0, \\ P_4 &= (u^2 - \lambda_3)(w_1^2 - \mu_1) - \nu_4 u w_1 = 0. \end{aligned}$$

We are interested only in the *non-trivial* solutions of this system, that is, the one-parametric branches of the solution set chosen in a way that each of z, w_1, u, w_2 is not a constant.

Remark 2.1. System (9) expresses the proportionality of the resultants $R_{12}(w_1, w_2) = \text{res}_z(P_1, P_2)$ and $R_{34}(w_1, w_2) = \text{res}_u(P_3, P_4)$.

Remark 2.2. There is an error in the last equation of (9) in [3], Subsection 3.1.2.

2.3. Five-parametric system. The problem of description of all flexible Kokotsakis polyhedra of the orthodiagonal anti-involutive type is separated into two sub-problems in a natural way:

- to describe the planar angles $(\alpha_i, \beta_i, \gamma_i, \delta_i)$ such that the corresponding polyhedron is of the orthodiagonal anti-involutive type;
- to describe its flexion, i.e. the trajectory in dihedral angles space.

The latter is equivalent to solving system (10), which describes the configuration space of (z, w_1, u, w_2) , with given coefficients λ_i, μ_i, ν_i that satisfy (8) and (9). In Theorem 2 we give a complete parameterization with one variable of a solution set of this system.

It is more complicated to describe all admissible planar angles $(\alpha_i, \beta_i, \gamma_i, \delta_i)$. It is natural to consider the angles $(\delta_1, \delta_2, \delta_3, \delta_4)$ as parameters. Since $\delta_1 + \delta_2 + \delta_3 + \delta_4 = 2\pi$, it is a three parametric family. From the algebraic point of view, system (9) together with identities (8) and definition (7) is a system of three equations in four variables $(\lambda_1, \mu_1, \lambda_3, \mu_3)$. It is enough to introduce one more parameter to describe a solution, which we do in Section 4. On the other hand, given $(\delta_1, \delta_2, \delta_3, \delta_4)$ and $(\lambda_1, \mu_1, \lambda_3, \mu_3)$ together with (8) one can recover angles α_i, γ_i from

$$(11) \quad \tan \alpha_i = \frac{\lambda_i - 1}{\lambda_i + 1} \tan \delta_i, \quad \tan \gamma_i = \frac{\mu_i - 1}{\mu_i + 1} \tan \delta_i$$

in the case $\frac{\pi}{2} \notin \{\delta_1, \delta_2, \delta_3, \delta_4\}$, and compute β_i from

$$(12) \quad \cos \beta_i = \frac{\cos \alpha_i \cos \gamma_i}{\cos \delta_i},$$

if the right-hand side is in $(-1, 1)$. That is, the angles of the base quadrilateral and $(\lambda_1, \mu_1, \lambda_3, \mu_3)$ determine all the needed data to construct a Kokotsakis polyhedron.

It is not obvious that such a geometric construction exists even one finds all the planar angles of an OAI Kokotsakis polyhedron. However, the following result gives the answer to this question.

Definition 2.3. We consider the following *geometric assumptions* on $(\delta_1, \delta_2, \delta_3, \delta_4)$ and (λ_i, μ_i) .

1. $(\delta_1, \delta_2, \delta_3, \delta_4)$ are the angles of a quadrilateral without right angles, that is $\delta_i > 0$ and $\delta_i \neq \frac{\pi}{2}$, where $i = 1, 2, 3, 4$ and $\delta_1 + \delta_2 + \delta_3 + \delta_4 = 2\pi$.
2. $(\lambda_i, \mu_i)_1^4$ satisfy systems (8), (9) with ν_i given by (7).
3. The angles $\{\alpha_i, \gamma_i, \beta_i\}_1^4$ given by (11) and (12) are well-defined, that is, the right-hand side of (12) is in interval $(-1, 1)$.
4. Spherical quadrilaterals Q_i with sides $(\alpha_i, \beta_i, \gamma_i, \delta_i)$, where $1 \leq i \leq 4$, are elliptic.

Theorem 1. Let $(\delta_1, \delta_2, \delta_3, \delta_4)$ and $(\lambda_i, \mu_i)_1^4$ meet the geometric assumptions of Definition 2.3. Then a Kokotsakis polyhedron with spherical quadrilaterals Q_i with sides $(\alpha_i, \beta_i, \gamma_i, \delta_i)$ exists, it is flexible and all its possible flexion are parameterized as in Theorem 2.

In Theorem 4 of Section 4 we parameterize all $(\lambda_i, \mu_i)_1^4$ that satisfy systems (8), (9) with ν_i given by (7).

Remark 2.4. We find it interesting that if $(\delta_1, \delta_2, \delta_3, \delta_4)$ and $(\lambda_i, \mu_i)_1^4$ meet only first two of the geometric assumptions, a non-trivial solution of (10) still exists and is given by Theorem 2. However, the angles β_i might fail to be determined by (12) if the absolute value of the right-hand side of (12) is bigger than 1. This follows from Lemma 4.1 and a condition of the existence of non-trivial solution in Theorem 2.

3. THE REAL PART OF THE CONFIGURATION SPACE

The form of solution in real numbers of the system (10) depends on the signs of $(\lambda_1, \mu_1, \lambda_3, \mu_3)$. That is, it depends on the signs of the entries in the matrix

$$\begin{pmatrix} \lambda_1 & \mu_1 \\ \lambda_2 & \mu_2 \\ \lambda_3 & \mu_3 \\ \lambda_4 & \mu_4 \end{pmatrix} = \begin{pmatrix} \lambda_1 & \mu_1 \\ -\lambda_1 & -\mu_3 \\ \lambda_3 & \mu_3 \\ -\lambda_3 & -\mu_1 \end{pmatrix}.$$

We call arrangement of signs in this matrix a *sign pattern*. It will be shown below in Lemma 3.1 that there is a unique possible arrangement of signs up to re-enumeration of the vertices. Using this observation, we assume that

$$(13) \quad \lambda_1, \mu_1 > 0 \quad \text{and} \quad \lambda_3, \mu_3 < 0.$$

Theorem 2. Let λ_i, μ_i, ν_i satisfy systems (9), (8) and assumption (13). Then system (10) has a real non-trivial one-parametric set of solutions if and only if $\zeta_1 := \frac{|\nu_1|}{4\sqrt{\lambda_1\mu_1}} > 1$. The solution set has four branches, which, up to the symmetry $(z, w_1, u, w_2) \rightarrow -(z, w_1, u, w_2)$, are given by

$$(14) \quad \begin{cases} z = \operatorname{sgn} \nu_1 \cdot \sqrt{\lambda_1} \cdot F(t)F(t + \frac{\pi}{2}), \\ w_1 = \sqrt{\mu_1} \cdot F(t)F(t - \frac{\pi}{2}), \\ u = \operatorname{sgn} \nu_4 \cdot \sqrt{-\lambda_3} \cdot \frac{\sqrt{\zeta_1 + 1} \pm (G(t) + G(t + \frac{\pi}{2}))}{V(t) - V(t + \frac{\pi}{2})}, \\ w_2 = \sqrt{-\mu_3} \cdot \frac{\operatorname{sgn}(\nu_1\nu_2)\sqrt{\zeta_1 + 1} \pm \operatorname{sgn}(\nu_3\nu_4)(G(t) + G(t - \frac{\pi}{2}))}{V(t) - V(t - \frac{\pi}{2})}. \end{cases},$$

where

$$F(t) = \sin t \sqrt{\zeta_1 - 1} + \sqrt{1 + (\zeta_1 - 1) \sin^2 t};$$

$$G(t) = \sin t \sqrt{1 + (\zeta_1 - 1) \sin^2 t};$$

$$V(t) = \sin t \sqrt{1 + (\zeta_1 - 1) \cos^2 t},$$

$t \in [0, 2\pi)$ is a parameter and the choice of \pm in u and w_2 is simultaneous.

According to assumption (13) and by condition $\zeta_1 > 1$, we see that all values under square roots in the previous formulas are positive.

3.1. Uniqueness of the sign pattern. Due to system (9), we have some restrictions to possible signs of (λ_i, μ_i) .

Lemma 3.1. *System (8-10) allows unique sign pattern of (λ_i, μ_i) up to cyclic re-enumeration of vertices. The only possible sign pattern is*

$$(15) \quad \text{Sign} \begin{pmatrix} \lambda_1 & \mu_1 \\ \lambda_2 & \mu_2 \\ \lambda_3 & \mu_3 \\ \lambda_4 & \mu_4 \end{pmatrix} = \begin{pmatrix} + & + \\ - & + \\ - & - \\ + & - \end{pmatrix}.$$

Proof. We first show that for some vertex both $\lambda_i, \mu_i > 0$. By the third equation in (9), we see that either $\lambda_1\mu_1 > 0$ or $\lambda_2\mu_2 > 0$. If in this positive product both factors are also positive, we have found the sought vertex. If however both coefficients are negative, from the fact that in a sign pattern there are always exactly four pluses and four minuses (follows from (8)), we conclude that if in sign pattern there is a row of minuses, there should be a row of pluses.

As cyclic permutations just change the order of vertices of the base quadrilateral, we assume that $\lambda_1 > 0$ and $\mu_1 > 0$. In other words, known signs are:

$$\text{Sign} \begin{pmatrix} \lambda_1 & \mu_1 \\ \lambda_2 & \mu_2 \\ \lambda_3 & \mu_3 \\ \lambda_4 & \mu_4 \end{pmatrix} = \begin{pmatrix} + & + \\ - & * \\ * & * \\ * & - \end{pmatrix}$$

We rewrite equation $P_1(z, w_1) = 0$ of (10) in the following way:

$$\left(\frac{z}{\sqrt{\lambda_1}} + \frac{\sqrt{\lambda_1}}{z} \right) \left(\frac{w_1}{\sqrt{\mu_1}} + \frac{\sqrt{\mu_1}}{w_1} \right) = \frac{\nu_1}{\sqrt{\lambda_1\mu_1}}.$$

Since absolute values of both parentheses are greater or equal to 2, for this equation to have a one-parametric solution (and not discrete set of points), we obtain

$$\frac{\nu_1^2}{\lambda_1\mu_1} > 16.$$

By the third equation in (9)

$$\frac{\nu_2^2}{\lambda_2\mu_2} = 16 - \frac{\nu_1^2}{\lambda_1\mu_1} < 0.$$

Hence, $0 > \lambda_2\mu_2 = \lambda_1\mu_3$ or, equivalently, $\mu_3 < 0$. By the first equation in (9), we get that $\lambda_3 < 0$. Finally, the only possible sign pattern is (15). \square

3.2. Reduced system. To simplify equations of system (10) and (9), we scale the variables and consider a reduced system (after the re-enumeration of vertices so that $\lambda_1, \mu_1 > 0$ and $\lambda_3, \mu_3 < 0$). Making the following substitution in system (10):

$$(16) \quad \begin{aligned} f_1 = f_2 &= \frac{z}{\sqrt{\lambda_1}} \text{sgn } \nu_1, & f_3 = f_4 &= \frac{u}{\sqrt{-\lambda_3}} \text{sgn } \nu_4, \\ g_1 = g_4 &= \frac{w_1}{\sqrt{\mu_1}}, & g_2 = g_3 &= \frac{w_2}{\sqrt{-\mu_3}} \text{sgn}(\nu_1\nu_2), \\ \zeta_i &= \frac{|\nu_i|}{4\sqrt{|\lambda_i\mu_i|}} \quad \text{for } i = 1, 2, 4, & \zeta_3 &= \frac{\nu_3}{4\sqrt{\lambda_3\mu_3}} \text{sgn}(\nu_1\nu_2\nu_4); \end{aligned}$$

we get 4 equations:

$$(17) \quad \begin{cases} (f_1^2 + 1)(g_1^2 + 1) = 4\zeta_1 f_1 g_1, \\ (f_1^2 - 1)(g_3^2 + 1) = 4\zeta_2 f_1 g_3, \\ (f_3^2 - 1)(g_3^2 - 1) = 4\zeta_3 f_3 g_3, \\ (f_3^2 + 1)(g_1^2 - 1) = 4\zeta_4 f_3 g_1, \end{cases}$$

where ζ_1, ζ_2 and ζ_4 are positive.

System (9), which describes conditions for the proportionality of the resultants, can be rewritten in the form

$$(18) \quad \begin{cases} \zeta_1^2 = \zeta_3^2, \\ \zeta_2^2 = \zeta_4^2, \\ \zeta_1^2 - \zeta_2^2 = 1. \end{cases}$$

3.3. Equation with positive signature. Let us study the first equation in (17):

$$(19) \quad (f_1^2 + 1)(g_1^2 + 1) = 4\zeta_1 f_1 g_1.$$

Recall $\zeta_1 > 0$.

One can see that f_1 and g_1 are of the same sign, and there is a symmetry of the solution $(f_1, g_1) \rightarrow (-f_1, -g_1)$. We assume that $f_1, g_1 > 0$. Now we consider new coordinates $X = \log(f_1/g_1), Y = \log f_1 g_1$. In these coordinates, the equation takes the form:

$$(20) \quad \cosh X + \cosh Y = 2\zeta_1.$$

Thus, there is a one-parametric real solution set if and only if

$$(21) \quad \zeta_1 > 1,$$

which we have already shown in [Lemma 3.1](#).

Solution to Equation (20) can be parameterized in an explicit way. Let $\omega = \operatorname{arccosh} \zeta_1$, $\varrho = \sqrt{2} \sinh \frac{\omega}{2}$. Then we can parameterize solution of (20) as

$$X = \operatorname{arcsinh}(\varrho \cos t) \quad \text{and} \quad Y = \operatorname{arcsinh}(\varrho \sin t),$$

where $t \in [0, 2\pi)$, which, by direct calculations, leads to the following representation of g_1 and f_1 :

$$f_1 = F(t)F(t + \frac{\pi}{2}) \quad \text{and} \quad g_1 = F(t)F(t - \frac{\pi}{2}),$$

where

$$F(t) = \sqrt{(\zeta_1 - 1) \sin^2 t + 1} + \sqrt{\zeta_1 - 1} \sin t;$$

Moreover, the solution of (20) is bounded in the plane (X, Y) . This means that f_1 and g_1 cannot be equal 0 or $\pm\infty$. Or, speaking geometrically, if equation (19) describes the configuration space of the spherical quadrilateral of a Kokotsakis polyhedron, then the corresponding side quadrilaterals never cross the base plane (see [Lemma 6.5](#)).

3.4. Proof of [Theorem 2](#).

Proof. By the results of [Subsection 3.3](#) and by (21) in particular, we see that the real solution of the system is nonempty and non-trivial if and only if $\zeta_1 > 1$ (recall that ζ_1, ζ_2 and ζ_4 are positive). Also in [Subsection 3.3](#), the equation with positive sign signature was resolved, which leads to the explicit formulas for z and w_1 . One can prove that the formulas for u and w_2 from (14) are the solution of the system by the direct substitution. However, we want to show that we have found all branches of the solution. We can consider system (17) and variables (f_1, g_1, f_3, g_3) instead of (z, w_1, θ, w_2) . One can see that the solution of the first equation is given in [Subsection 3.3](#), the second equation is quadratic in g_3 , the fourth equation is quadratic in f_3 . Hence, we have two possible branches of solution for both f_3 and g_3 , and there are at most four branches of the solution up to the symmetry $(f_1, g_1, f_3, g_3) \rightarrow (-f_1, -g_1, -f_3, -g_3)$.

So the only difficulty is to check that we have chosen the correct branches to meet the third equation of the system. We claim that the latter obstacle implies that there are exactly two branches of the solution with positive f_1 and g_1 . Hence, we described all possible solution with positive f_1 and g_1 . Indeed, fixing f_1 , we fix $(g_3^2 + 1)/g_3$. Therefore, fixing f_1, f_3, g_1 , we fix $(g_3^2 - 1)/g_3$. But $(g_3^2 - 1)/g_3 = (g_3^2 + 1)/g_3 - 2/g_3$ changes its value if we change the branch for g_3 . Therefore, there is a unique branch of g_3 for each branch of f_3 . By the symmetry $(f_1, f_3, g_1, g_3) \rightarrow (-f_1, -g_1, -f_3, -g_3)$, we describe all branches of the solution. \square

3.5. Reducible resultant. As a linear substitution does not change the property of irreducibility of the resultant, we consider the resultant R_{12}^r of the first two equations of reduced system (17) as polynomials in f_1 . R_{12}^r is given by:

$$\begin{aligned} \frac{1}{4}R_{12}^r = & 1 + 2(1 - 2\zeta_1^2)g_1^2 + 2(1 + 2\zeta_2^2)g_3^2 + g_1^4 + 4(1 - 2\zeta_1^2 + 2\zeta_2^2)g_1^2g_3^2 + g_3^4 \\ & + 2(1 + 2\zeta_2^2)g_1^4g_3^2 + 2(1 - 2\zeta_1^2)g_1^2g_3^4 + g_1^4g_3^4. \end{aligned}$$

When $\zeta_1^2 - \zeta_2^2 = 1$ from (18) holds true, the resultant is the product of the following polynomials:

$$\frac{1}{4}R_{12}^r = \left(g_1^2g_3^2 + (\zeta_1 + \zeta_2)^2(g_1^2 - g_3^2) - 1\right) \left(g_1^2g_3^2 + (\zeta_1 - \zeta_2)^2(g_1^2 - g_3^2) - 1\right).$$

This can be checked by a direct calculation. Moreover, as we show in Lemma 6.6 below, the resultant is reducible if and only if $\zeta_1^2 - \zeta_2^2 = 1$.

3.6. Symmetries. Here we discuss some symmetries hidden in our equations.

First of all, the symmetry $(z, w_1, u, w_2) \rightarrow (-z, -w_1, -u, -w_2)$ or, equivalently, in terms of the dihedral angles $(\varphi, \psi_1, \theta, \psi_2) \rightarrow (-\varphi, -\psi_1, -\theta, -\psi_2)$, is just the symmetry with respect to a plane of the base quadrilateral. Clearly, it does not change a configuration.

Secondly, our equations allow us to make the following substitution for planar angles at vertex A_i of the base quadrilateral:

- (1) $(\alpha_i, \gamma_i) \rightarrow (\alpha_i - \pi, \gamma_i - \pi)$;
- (2) $(\alpha_i, \beta_i) \rightarrow (\alpha_i - \pi, \pi - \beta_i)$;
- (3) $(\gamma_i, \beta_i) \rightarrow (\gamma_i - \pi, \pi - \beta_i)$.

Clearly, the corresponding spherical quadrilateral remains elliptic, and the involution factors do not change. These symmetries may help with avoiding self-intersections of a polyhedron.

3.7. Elliptic parameterization. In [3] the author parameterized a solution of (4) in terms of the Jacobi elliptic sines. Using the results of that paper, we provide a sketch of the proof of the following result in which we give an elliptic parameterization of the solution set of system (10). A comprehensive description of the OAI flexible Kokotsakis polyhedra with the use of elliptic functions can be found in [2].

Theorem 3. *Let λ_i, μ_i, ν_i satisfy systems (9), (8) and assumption (13). Then system (10) has a real non-trivial one-parametric set of solutions if and only if $\zeta_1 := \frac{|\nu_1|}{4\sqrt{\lambda_1\mu_1}} > 1$. The solution set has four branches, which, up to the symmetry $(z, w_1, u, w_2) \rightarrow -(z, w_1, u, w_2)$, are given by*

$$(22) \quad \begin{cases} z = \operatorname{sgn} \nu_1 \cdot \sqrt{\lambda_1 k} \operatorname{sn}(K + it), \\ w_1 = \sqrt{\mu_1 k} \operatorname{sn}\left(K + \frac{iK'}{2} + it\right), \\ u = i \operatorname{sgn} \nu_4 \cdot \sqrt{-\lambda_3 k} \operatorname{sn}\left(K + \frac{iK'}{2} \pm \operatorname{sgn}(\nu_1\nu_2)\left(K - \frac{iK'}{2}\right) + it\right), \\ w_2 = i \operatorname{sgn}(\nu_1\nu_2) \cdot \sqrt{-\mu_3 k} \operatorname{sn}\left(K \mp \operatorname{sgn}(\nu_3\nu_4)\left(K - \frac{iK'}{2}\right) + it\right) \end{cases}$$

where k is the elliptic modulus given by $k = \left(\zeta_1 - \sqrt{\zeta_1^2 - 1}\right)^2$, K and $\frac{iK'}{2}$ are the quarter periods of elliptic sine with modulus k ; $t \in [0, 2K')$ is a parameter and the choice of \pm and \mp in u and w_2 is simultaneous.

Sketch of the proof. It was shown in [3] that the solution set of (4) has a parameterization of the form

$$z = p \operatorname{sn}(t, k), \quad w = q \operatorname{sn}(t + \tau, k),$$

where τ is a quarter-period of sn , that is, $\tau = nK + \frac{iK'}{2}$, and the amplitudes p, q belong to $\mathbb{R}_+ \cup i\mathbb{R}_+$. Moreover, there is a table with explicit formulas for (λ, μ, ν) as functions of (p, q, k) in Lemma 4.17 of [3]:

$$(\lambda, \mu, \nu) = \begin{cases} \left(\frac{p^2}{k}, \frac{q^2}{k}, \frac{2(1+k)}{k\sqrt{k}} pq \right), & \text{if } \tau = \frac{iK'}{2} \\ \left(\frac{p^2}{k}, \frac{q^2}{k}, -\frac{2(1+k)}{k\sqrt{k}} pq \right), & \text{if } \tau = 2K + \frac{iK'}{2} \\ \left(-\frac{p^2}{k}, -\frac{q^2}{k}, \frac{2i(1-k)}{k\sqrt{k}} pq \right), & \text{if } \tau = K + \frac{iK'}{2} \\ \left(-\frac{p^2}{k}, -\frac{q^2}{k}, -\frac{2i(1-k)}{k\sqrt{k}} pq \right), & \text{if } \tau = 3K + \frac{iK'}{2} \end{cases}$$

From this table, we see that the elliptic modulus k is a function of $\frac{\nu^2}{\lambda\mu}$. Since $k \in (0, 1)$ and by the choice of sign pattern (15), we get that the elliptic moduli of all four spherical quadrilaterals are the same in case of an OAI polyhedron. Then it is easy to recover what a quarter-period shift one can use in each equation of (17). \square

Remark 3.1. It follows that the OAI type of Kokotsakis polyhedra is a special class of the elliptic equimodular type in the notation of [3].

We note that the parameterizations from Theorem 3 cannot be transformed to the parameterizations given by Theorem 2 using linear substitution of the parameter t as the first one uses elliptic functions that are not elementary. However, under the substitution $t \rightarrow -\frac{K'}{\pi}t$, the parameterization given by Theorem 3 is quite close to the parameterization given by Theorem 2 for k small enough (see Figure 13 and Figure 14).

4. PLANAR ANGLES OF A FLEXIBLE POLYHEDRON

In this section, we study $(\delta_1, \delta_2, \delta_3, \delta_4)$ and (λ_i, μ_i) that meet the first two of the geometric assumptions. In the following theorem we manage to parameterize all such (λ_i, μ_i) . In fact, for solving system (9), we need to make it more symmetrical with a proper substitution, and we use new variables for this purpose. We introduce these substitutions in the theorem and explain their sense later.

Theorem 4. *Let all δ_i and (λ_i, μ_i) meet the first two of the geometric assumptions. Put*

$$(23) \quad s = \frac{\delta_1 - \delta_2 + \delta_3 - \delta_4}{4}, \quad x = \frac{\delta_1 - \delta_3}{2}, \quad y = \frac{\delta_2 - \delta_4}{2}.$$

Then there exists $\tau \in [0, 2\pi)$ and a proper choice of signs such that

$$(24) \quad \lambda_i = \frac{1 \pm \sqrt{1 - r_i^2}}{r_i}, \quad \mu_i = \frac{1 \pm \sqrt{1 - c_i^2}}{c_i} \quad \text{for } i = 1, 3,$$

where r_i, c_i are functions of (τ, x, y, s) with the symmetries

$$(25) \quad \begin{aligned} c_1(\tau, x, y, s) &= r_1(\tau + \pi, x, -y, s), \\ r_3(\tau, x, y, s) &= r_1(\tau, -x, -y, s), \\ c_3(\tau, x, y, s) &= r_1(\tau + \pi, -x, y, s). \end{aligned}$$

and r_1 is given by

$$(26) \quad r_1(\tau, x, y, s) = \frac{N + S\sqrt{L}}{2D},$$

where

$$\begin{aligned} S &= s_{10} \cos \tau + s_{01} \sin \tau, \\ L &= l_{20} \cos^2 \tau + l_{11} \sin \tau \cos \tau + l_{02} \sin^2 \tau, \\ N &= n_{20} \cos^2 \tau + n_{11} \sin \tau \cos \tau + n_{02} \sin^2 \tau, \\ D &= d_{20} \cos^2 \tau + d_{11} \sin \tau \cos \tau + d_{02} \sin^2 \tau, \end{aligned}$$

with

$$\begin{aligned} s_{10} &= \cos 2x + \cos 2y + 2 \cos 2s, \\ s_{01} &= \cos 2x - \cos 2y, \\ l_{20} &= (\cos 2x - \cos 2y)^2 + 8 \cos 2s (\cos 2x + \cos 2y), \\ l_{11} &= 2(\cos 2x - \cos 2y)(\cos 2x + \cos 2y + 2 \cos 2s), \\ l_{02} &= (\cos 2x + \cos 2y - 2 \cos 2s)^2, \\ n_{20} &= \sin(x+y)(\sin(x-3y) + \sin(3x-y) + 6 \sin(x-y+2s) - 2 \sin(x-y-2s)), \\ n_{11} &= 8(\sin^2(x+s) \cos^2(x-s) + \sin^2(y-s) \cos^2(y+s)), \\ n_{02} &= (\cos 2y - \cos 2x)(\cos 2x + \cos 2y - 2 \cos 2s), \\ d_{20} &= \cos(x-y)(\cos(3x+y) + \cos(x+3y) + 2 \cos(x+y-2s) + 4 \cos(x+3s) \cos(y-s)), \\ d_{11} &= \cos 4x - \cos 4y - 4 \sin(x+y) \sin(x-y+2s), \\ d_{02} &= \sin(x-y)(\sin(x+3y) - \sin(3x+y) + 2 \sin(x+y-2s) - 4 \cos(x+3s) \sin(y-s)). \end{aligned}$$

It is very intriguing that for each value of τ in this theorem such that all values under square roots are positive, system (10) has real non-trivial solutions.

Lemma 4.1. *Let δ_i and (λ_i, μ_i) meet the first two of the geometric assumptions. Then (15) is the sign pattern of (λ_i, μ_i) up to re-enumeration of vertices (16) and $\frac{\nu_j^2}{\lambda_j \mu_j} > 16$ for j such that $\lambda_j, \mu_j > 0$.*

4.1. A new symmetric system. System (9) can be rewritten in the following form:

$$(27) \quad \begin{cases} \frac{(\lambda_1 - 1)^2 (\mu_1 - 1)^2}{\lambda_1 \mu_1 \cos^2 \delta_1} + \frac{(\lambda_2 - 1)^2 (\mu_2 - 1)^2}{\lambda_2 \mu_2 \cos^2 \delta_2} = 16, \\ \frac{(\lambda_1 - 1)^2 (\mu_1 - 1)^2}{\lambda_1 \mu_1 \cos^2 \delta_1} = \frac{(\lambda_3 - 1)^2 (\mu_3 - 1)^2}{\lambda_3 \mu_3 \cos^2 \delta_3}, \\ \frac{(\lambda_2 - 1)^2 (\mu_2 - 1)^2}{\lambda_2 \mu_2 \cos^2 \delta_2} = \frac{(\lambda_4 - 1)^2 (\mu_4 - 1)^2}{\lambda_4 \mu_4 \cos^2 \delta_4}. \end{cases}$$

Make a substitution

$$(28) \quad r_i = \frac{2\lambda_i}{\lambda_i^2 + 1}, \quad c_i = \frac{2\mu_i}{\mu_i^2 + 1}, \quad a_i = \frac{1}{\cos^2 \delta_i},$$

and let

$$(29) \quad Z_i = a_i(r_i^{-1} - 1)(c_i^{-1} - 1) = 4 \frac{\nu_i^2}{\lambda_i \mu_i} = \frac{(\lambda_i - 1)^2 (\mu_i - 1)^2}{4 \lambda_i \mu_i \cos^2 \delta_i}.$$

Then system (27) is equivalent to the system of the following three equations $Z_1 + Z_2 = 4$, $Z_1 = Z_3$ and $Z_2 = Z_4$. Adding a linear dependent equation, system (27) can be rewritten in the following equivalent way

$$(30) \quad \frac{1}{2} Z_i = 1 - (-1)^i \tan \tau,$$

where τ is parameter.

System (8) takes the form:

$$(31) \quad r_1 = -r_2, \quad c_1 = -c_4, \quad c_2 = -c_3, \quad r_3 = -r_4.$$

The equations of system (30) are linear in both r_i or c_i . Since, by the definitions, λ_i, μ_i are real and $\nu_i \neq 0$, we obtain a restriction on the values of r_i, c_i :

$$(32) \quad -1 \leq r_i, c_i < 1.$$

It is clear that (r_i, c_i) has the same sign pattern as (λ_i, μ_i) and that substitution (28) is inverse to (24).

4.2. Proof of Theorem 4.

Proof. Restriction (32) yields positivity of radical expressions in (24). So, it suffices to prove identities (25) and (26). We start with the latter one.

Using exclusion technique to find $r_1(\tan \tau)$ from system (30) is equivalent to finding roots of an equation of the second degree with coefficients determined by the coefficients of the system (30). Indeed, from the equation $Z_i = \text{const}$, we see that functions $c_i(r_i)$ or $r_i(c_i)$ are linear fractional transformations. Since composition of two linear fractional transformations is a linear fractional transformation, finding $c_1(r_1), r_3(c_1)$ and $c_2(r_1), r_3(c_2)$, we obtain two different Möbius transformation functions for $r_3(r_1)$. Therefore, general solution has a form:

$$(33) \quad r_1(\tan \tau) = \frac{P(\tan \tau) \pm \sqrt{Q(\tan \tau)}}{R(\tan \tau)},$$

where P and R are polynomials of 2nd degree and Q is a polynomial of 4th degree. However, we noticed that Q always have a full square as a multiplier, when $\sum \delta_i = 2\pi$. Using our substitution (23), we managed to factorize Q , merge two branches in (33) and obtain (26). The explicit formulas for the coefficients were found and can be verified with the use of a system of computer algebra. We used Wolfram Mathematica 11 for this purpose.

Since equation (33) is obtained by the exclusion technique, it describes all possible r_1 . Note that it has two branches, however we have only one in (26). So we need explain how we managed to merge the branches. The radical expression in (33) equals to $\frac{S^2 L}{\cos^4 \tau}$. Since S is the only function that have linear terms in $\cos \tau$ and $\sin \tau$, substitution $\tau \rightarrow \tau + \pi$ changes the sign of S and preserves the values of N, L, D in (26), that is, it merges two branches of (33).

Let us explain identities (25). There are no more solutions as each equation of (30) is linear in either r_i or c_i . Apart from checking (25) with direct calculations, we provide the following arguments. The dependencies on x, y, s arise from re-enumeration of vertices, Explanation of a “phase shift” of τ is trickier. By the symmetry of the system (30), it is enough to explain a “phase shift” for c_1 . Fixing r_1 , we have exactly one possible value for c_1 by the equation $Z_1 = 2 - 2 \tan \tau$ and two possible branches in (33) rewritten for c_1 . The “phase shift” $\tau \rightarrow \tau + \pi$ corresponds to the right choice of the branch. \square

4.3. Proof of Lemma 4.1 and Theorem 1.

Proof of Lemma 4.1. By the first equation in (27), either r_1, c_1 or r_2, c_2 are both non-zero and have the same sign. Again, by Pigeonhole principle, it implies that there is a pair with positive numbers. Without loss of generality, let it be pair r_1, c_1 . By the second equation, the latter implies that r_3, c_3 has the same sign. We claim that they are negative, and, therefore, we have the case of the right sign pattern.

Assume the contrary that $r_3, c_3 > 0$. By the definition of Z_1 and system (30), we have

$$2 + 2 \tan \tau = Z_1 = \frac{1}{\cos^2 \delta_1} \left(\frac{1}{r_1} - 1 \right) \left(\frac{1}{c_1} - 1 \right) > 0,$$

or, equivalently, $2 > -2 \tan \tau$. Then, by the definition of Z_2 , we have

$$Z_2 = \frac{1}{\cos^2 \delta_2} (1/r_1 + 1)(1/c_3 + 1) > (1/r_1 + 1)(1/c_3 + 1) > 4 > 2 - 2 \tan \tau,$$

which contradicts system (30). Therefore, $r_3, c_3 < 0$. Then, by system (30) and (29), we obtain

$$\frac{\nu_1^3}{\lambda_3 \mu_3} = \frac{\nu_1^1}{\lambda_1 \mu_1} = \frac{1}{4} Z_3 = \frac{1}{4 \cos^2 \delta_3} \left(\frac{1}{|r_3|} + 1 \right) \left(\frac{1}{|c_3|} + 1 \right) > 1.$$

□

Theorem 1 is a direct consequence of this lemma.

Proof of Theorem 1. To proof the theorem we need to check the following

- spherical quadrilaterals with sides $(\alpha_i, \beta_i, \gamma_i, \delta_i)$ exist;
- system (10) has non-trivial solution.

The first assumption follows from **Lemma 8.2**, the second is guaranteed by **Lemma 4.1**. □

5. ALGORITHM

In the following Theorem, we summarize all needed steps to construct a flexible Kokotsakis polyhedron of the orthodiagonal anti-involutive type and to describe its flexion using given values of the angles of the base quadrilateral. We are always looking for angles in interval $(0, \pi)$. However, one can allow to the angles α_i and γ_i to be in $(0, \pi) \cup (\pi, 2\pi)$.

Theorem 5. *Given parameters $(\delta_1, \delta_2, \delta_3, \delta_4)$, the algorithm of constructing a flexible Kokotsakis polyhedron of the orthodiagonal anti-involutive type is the following:*

- (1) Check whether $(\delta_1, \delta_2, \delta_3, \delta_4)$ meet the first of the geometric assumptions, that is, they form a quadrilateral without right angles.
- (2) Calculate (x, y, s) using substitution (23).
- (3) Check whether there exists τ such that r_1, r_3, c_1, c_3 given by (26) and (25) satisfy inequality (32) (see **Subsection 5.1** for how this can be done). Set $r_2 = -r_1, r_4 = -r_3, c_2 = -c_3, c_4 = -c_1$.
- (4) Calculate the angles α_i and γ_i from

$$\tan \alpha_i = \sigma_i^\alpha \sqrt{\frac{1-r_i}{1+r_i}} \tan \delta_i \quad \text{and} \quad \tan \gamma_i = \sigma_i^\gamma \sqrt{\frac{1-c_i}{1+c_i}} \tan \delta_i,$$

where $\sigma_i^\bullet = \pm 1$ are chosen in a way to satisfy: $\sigma_1^\alpha \sigma_2^\alpha = \sigma_3^\alpha \sigma_4^\alpha = \sigma_1^\gamma \sigma_4^\gamma = \sigma_2^\gamma \sigma_3^\gamma = 1$. Select angles to be from 0 to π .

- (5) Check that $\frac{\cos \alpha_i \cos \gamma_i}{\cos \delta_i} \in (-1, 1)$. Find β_i using (12).
- (6) Check that obtained spherical quadrilaterals Q_i with sides $(\alpha_i, \beta_i, \gamma_i, \delta_i)$ are elliptic.
- (7) Find λ_i, μ_i, ν_i using formulas (5), (6) and (7).
- (8) Re-enumerate vertices to get (15) as a sign pattern of the system.
- (9) Find (z, w_1, u, w_2) using formulas of **Theorem 2**. Calculate dihedral angles using

$$(\varphi, \psi_1, \theta, \psi_2) = (2 \arctan z, 2 \arctan w_1, 2 \arctan u, 2 \arctan w_2).$$

Proof. If all requirements are fulfilled, an existence and flexibility of a Kokotsakis polyhedron constructing using this algorithm follows from **Theorem 1** and **Theorem 2**. □

Remark 5.1. We do not check whether the obtained surface is not self-intersecting. There remains a possibility that at every moment during flexion, there are two intersecting facets.

It's possible to derive a implicit formula for a set of (x, y, s) (or, equivalently, $(\delta_1, \delta_2, \delta_3, \delta_4)$) for which there exists τ such that $-1 < r_i, c_i < 1$ and the angles β_i are well-defined. However, the resulting formula is likely to be enormous and incomprehensible. We instead explain how to check the conditions and show the result of numerical screening in the next subsection.




FIGURE 4. Screening result. Due to the symmetry of the system, the set is centrally symmetric with respect to $(\frac{\pi}{4}, \frac{\pi}{4}, \frac{\pi}{4})$ and is symmetric with respect to the plane $x = y$.




FIGURE 5. Set of $(\delta_1, \delta_2, \delta_3)$ with existing flexible an OAI polyhedron. On the left: with convex base quadrilateral; on the right: with non-convex base quadrilateral ($\delta_4 > \pi$).

5.1. **Screening of the space of parameters $(\delta_1, \delta_2, \delta_3, \delta_4)$.** In [Theorem 5](#), one need to find a proper parameter τ for a given angles (x, y, s) (or, equivalently, $(\delta_1, \delta_2, \delta_3, \delta_4)$). We run a computer screening and found numerically the set of (x, y, s) for which there exists τ such that (λ_i, μ_i) are given by [Theorem 4](#) and $\{\delta_1, \delta_2, \delta_3, \delta_4\}$ meet the geometric assumptions. The result of our numerical computation is in [Figure 4](#), here we do not show surfaces on which at least one Q_i is not elliptic. [Figure 5](#) shows the result of screening in $(\delta_1, \delta_3, \delta_2)$ coordinates for convex and non-convex base quadrilaterals.

We claim that it can be done with an arbitrary precision or, probably, explicit formulas for the boundary can be found, as all the restrictions on $\tan \tau$ are polynomial.

More precisely,

$$(1) \frac{L(\tau, x, y, z)}{\cos^2 \tau} \geq 0 \text{ is clearly a polynomial in } \tan \tau.$$

- (2) $r_i < C = \text{const}$ (and others comparisons for r_i, c_i with constant C) is equivalent to a proper system of the following polynomial inequalities

$$\left(\frac{2CD - N}{S \cos \tau}\right)^2 \geq \frac{L}{\cos^2 \tau}, \quad \left(\frac{2CD - N}{S \cos \tau}\right)^2 \leq \frac{L}{\cos^2 \tau},$$

and comparison of $\frac{2CD-N}{S \cos \tau}$, $\frac{D}{\cos^2 \tau}$ and $\frac{S}{\cos \tau}$ with zero.

- (3) β_i is well-defined if and only if

$$\frac{\cos^2 \delta_i}{\cos^2 \alpha_i \cos^2 \gamma_i} \leq 1.$$

By $\frac{1}{\cos^2 \alpha} = 1 + \tan^2 \alpha$, the latter is

$$\cos^2 \delta_i \left(\frac{1 - r_i}{1 + r_i} \tan^2 \delta_i + 1\right) \left(\frac{1 - c_i}{1 + c_i} \tan^2 \delta_i + 1\right) \geq 1,$$

which is equivalent to a system of polynomial inequalities.

6. GEOMETRIC PROPERTIES

6.1. The configuration space of a spherical quadrilateral. Side lengths and angles of a spherical quadrilateral Q are denoted in Figure 6, and $z = \tan \frac{\varphi}{2}$, $w = \tan \frac{\psi}{2}$.




FIGURE 6. Spherical quadrilateral Q

As was discovered by Bricard [1], the equation of the configuration space of Q has the form

$$(34) \quad P(z, w) = a_{22}z^2w^2 + a_{20}z^2 + a_{02}w^2 + 2a_{11}zw + a_{00} = 0,$$

where the coefficients are trigonometric functions of the angles of Q .

By allowing z to take complex values, we arrive at a complex algebraic curve

$$(35) \quad C = \{(z, w) \in \mathbb{C}P^1 \times \mathbb{C}P^1 \mid P(z, w) = 0\}$$

with two coordinate projections

$$(36) \quad P_z : C \rightarrow \mathbb{C}P^1 \ni z \quad \text{and} \quad P_w : C \rightarrow \mathbb{C}P^1 \ni w.$$

If Q is of elliptic type then C is an elliptic curve. The projections in (36) are *two-fold branched covers with exactly 4 points in the branch set*. Since equation (34) is quadratic in one variable, one can easily find explicit formulas for points of the branch set by solving a quadratic equation. The complete classification of different classes of spherical quadrilaterals and their branch sets can be found in [3] Subsection 2.4 and Lemma 4.10.

Denote by

$$\begin{array}{ll} i: C \rightarrow C & j: C \rightarrow C \\ (z, w) \mapsto (z, w') & (z, w) \mapsto (z', w) \end{array}$$

the deck transformations of P_w and P_z . If z and w are real, we can give a geometric interpretation of these *involutions*: i and j act by folding the quadrilateral along one of its diagonals, see Figure 7. This means that realizable branch points correspond to a degenerate case when quadrilateral becomes a triangle. Hence, we conclude

Lemma 6.1. *Let Q be a spherical quadrilateral of elliptic type. We use the notation as in Figure 6 and put $z = \tan \frac{\varphi}{2}$. Then z belongs to the branch set of P_z if and only if $\chi = 0 \pmod{\pi}$.*




FIGURE 7. Involutions i and j on the configuration space of a quadrilateral.

6.2. Scissors-like linkage. Two adjacent vertices of the base quadrilateral share the common dihedral angles, which means the corresponding spherical quadrilaterals are *coupled* by means of the angle (see Figure 8). There are four such *scissors-like linkages* corresponding to edges A_1A_2 , A_2A_3 , A_3A_4 and A_4A_1 of the base quadrilateral in a Kokotsakis polyhedron. Isometric deformations of the polyhedron correspond to motions of these linkages. Algebraically speaking, these linkages are described by the rows and columns of system (2), and the system itself describes possible dihedral angles of the polyhedron. Moreover, a Kokotsakis polyhedron is flexible if and only if the system of polynomial equations (2) has a one-parameter family of solutions over the reals ([3], Lemma 2.2). One possible approach to find a solution is as follows:

- (1) consider a pair of opposite edges A_1A_2 and A_3A_4 together with corresponding linkages, which are described by systems $\{P_1 = 0, P_2 = 0\}$ and $\{P_3 = 0, P_4 = 0\}$, respectively;
- (2) exclude common variables z and u in the corresponding systems by computing the resultants $R_{12}(w_1, w_2) = \text{res}_z(P_1, P_2)$ and $R_{34}(w_1, w_2) = \text{res}_u(P_3, P_4)$.




FIGURE 8. Two coupled spherical quadrilaterals associated with the edge A_1A_2 . The two marked angles are required to stay equal during the deformation.

The polyhedron is flexible if and only if the algebraic sets $R_{12} = 0$ and $R_{34} = 0$ have a common irreducible component. This means that they are reducible or irreducible simultaneously. Stachel and Nawratil described all flexible classes for reducible R_{12} and R_{34} .

Remark 6.1. Geometrically speaking, the zero set of the resultant $R_{12}(w_1, w_2)$ gives an implicit dependence of the non-common angles of spherical quadrilaterals Q_1 and Q_2 , and R_{34} does the same for Q_3 and Q_4 . The polyhedron is flexible if and only if these “dependencies” have a common branch, that is, one can join the scissors-like linkages (Q_1, Q_2) and (Q_3, Q_4) during the corresponding flexion.

Izmestiev [3] considers the complexified configuration space of a spherical linkage (that is, the set of all complex solutions of the corresponding polynomial system). The following lemma is a direct consequence of the results of [3]. It follows from Lemma 5.1 and assertion (4) of Lemma 4.10 of that paper.

Lemma 6.2. *Let polynomial system $\{P_1(z, w_1) = 0 = P_2(z, w_2)\}$ describe a scissors-like linkage of two elliptic spherical quadrilaterals Q_1 and Q_2 . The resultant $R_{12}(w_1, w_2)$ is reducible if and only if the branch sets of w_1 and w_2 coincide.*

Remark 6.2. We note that Izmestiev understands the property of irreducibility of a coupling of two spherical quadrilaterals as the irreducibility of an algebraic set that describes the corresponding complexified configuration spaces. One should not confuse the properties of irreducibility of the resultant with the irreducibility of the coupling.

As was mentioned above, Izmestiev showed that there is only one possible candidate for a flexible Kokotsakis polyhedron with irreducible resultant R_{12} . In particular, all scissors-like linkages of such a polyhedron must be of a special type, which the author called anti-involutive coupling.

6.3. Anti-involutive coupling and its properties. A coupling of two orthodiagonal quadrilaterals is called *anti-involutive* if their involution factors at the common vertex are opposite, e.g. $\lambda_1 = -\lambda_2$ for two coupled spherical quadrilaterals associated with the edge A_1A_2 .

Lemma 6.3. *Let spherical orthodiagonal quadrilaterals Q_1 and Q_2 form a scissor-like linkage as in Figure 9 with the involution factors λ_1 and λ_2 in the common vertex. Then*

$$\begin{aligned} \lambda_1 = \lambda_2 &\Leftrightarrow \xi_1 = \xi_2, \xi'_1 = \xi'_2 \text{ during flexion} \\ \lambda_1 = -\lambda_2 &\Leftrightarrow \xi_1 = \xi'_2, \xi'_1 = \xi_2 \text{ during flexion} \end{aligned}$$




FIGURE 9. To the proof of Lemma 6.3

Proof. Equation (4) implies that the involutions of an orthodiagonal quadrilateral are given by

$$i(z, w) = (\lambda z^{-1}, w), \quad j(z, w) = (z, \mu w^{-1}).$$

Therefore, the condition $\lambda_1 = \lambda_2$ is equivalent to compatibility of the involutions of the coupled quadrilaterals at their common vertex:

$$i_1(z, w_1) = (z', w_1), \quad i_2(z, w_2) = (z'', w_2) \Rightarrow z' = z''.$$

On the other hand, the involution i_1 changes the angle of the first quadrilateral at the common vertex from $\xi_1 + \xi'_1$ to $\xi_1 - \xi'_1$, and the angle of the second quadrilateral from $\xi_2 + \xi'_2$ to $\xi_2 - \xi'_2$. The compatibility means that the angles at the common vertex remain equal after the involution: $\xi_1 - \xi'_1 = \xi_2 - \xi'_2$. Because $\xi_1 + \xi'_1 = \xi_2 + \xi'_2$, this is equivalent to $\xi_1 = \xi_2, \xi'_1 = \xi'_2$.

Observe that exchanging α and δ changes the sign of the involution factor. Thus, if we rotate the left half of the scissors linkage, then the condition $\lambda_1 = -\lambda_2$ transforms to $\lambda_1 = \lambda_2$. On the other hand, this exchanges the angles ξ_2 and ξ'_2 . This proves the second part of the lemma. \square

Lemma 6.4. *In an antiinvolutive ($\lambda_1 = -\lambda_2$) linkage of two orthodiagonal quadrilaterals, if one of the angles χ_1 and χ'_2 in Figure 10 becomes 0 or π , then the other one does as well.*

Proof. Indeed, $\chi_1 = 0 \pmod{\pi}$ if and only if $\xi_1 = 0 \pmod{\pi}$. \square




FIGURE 10. To the proof of Lemma 6.4.

6.4. Geometric properties of orthodiagonal anti-involutive type polyhedra. From the definition of an OAI polyhedron, we see that all four linkages of it are anti-involutive. Applying Lemma 6.4, we observe the following *flattening effect*.

Corollary 6.1. *In a flexible OAI Kokotsakis polyhedron the dihedral angles at the three bold edges in Figure 11 vanish or become π at the same time.*




FIGURE 11. A partial flattening of a flexible OAI polyhedron.

Using Theorem 2, it is quite easy to understand what kind of flattening has a flexible OAI Kokotsakis polyhedron.

Lemma 6.5. *Two dihedral angles of a flexible OAI Kokotsakis polyhedron never become 0 or π , the two other dihedral angles become 0 and π on each brunch of the solution given by Theorem 2*

Proof. We use the notation of [Section 3](#). If one of the angles $\varphi, \psi_1, \theta, \psi_2$ is 0 or π then the corresponding variable is 0 or ∞ . Since substitution [\(16\)](#) just scales the variables, the same happens to the corresponding variable of system [\(17\)](#). Let us consider it for simplicity. By [Subsection 3.3](#), the variables of the first equation of [\(17\)](#) (with positive signature) never become 0 or ∞ . Also, each branch of the solution of the whole system contains all solutions of the first equation up to the symmetry $(f_1, g_1) \rightarrow -(f_1, g_1)$. Moreover, since $\zeta_1 > 1$, then $f_1 = 1$ at two different points of the solution of the first equation. Therefore, by the second equation, g_3 becomes 0 or ∞ at these points on each branch of the solution of the whole system. It is easy to see that g_3 is different at this points. So it must be 0 and ∞ at these points, and it can have these values only at these points. A similar argument works for g_1 and f_3 . \square

Summarizing all results, we prove Stachel's conjecture.

Theorem 6. *Stachel's conjecture is true. The resultant R_{12} of a real flexible OAI Kokotsakis polyhedron is reducible.*

Proof. We use the notation of the first three sections. By [Lemma 6.2](#), it is enough to show that the branch sets for w_1 and w_2 coincide. By [Lemma 6.1](#), it suffices to prove that the dihedral angles at edges A_1B_1 and A_2B_2 (see [Figure 2](#)) become 0 or π simultaneously at four different points. This follows from [Corollary 6.1](#) and [Lemma 6.5](#). \square

Remark 6.3. It is easy to show that the resultant R_{12} is reducible by a direct calculation of the branch sets for w_1 and w_2 from the first two equations of system [\(10\)](#). The first equation of [\(8\)](#) and the third equation of [\(9\)](#) implies that the branch sets coincide. This observation works even for non-real polyhedra.

Lemma 6.6. *The resultant $R_{12}^*(g_1, g_3)$ of the first two equations of system [\(17\)](#) is reducible if and only if $\zeta_1^2 - \zeta_2^2 = 1$.*

Proof. It is easy to see that f_1 belongs to the branch set of g_1 if and only if

$$\frac{4\zeta_1 f_1}{f_1^2 + 1} = \pm 2.$$

From here, the branch set is

$$\left\{ \pm\zeta_1 \pm \sqrt{\zeta_1^2 - 1} \right\}.$$

Similar, the branch set of w_2 for the second equation is

$$\left\{ \pm\zeta_2 \pm \sqrt{\zeta_2^2 + 1} \right\}.$$

These sets coincide exactly when $\zeta_1^2 - \zeta_2^2 = 1$. \square

7. EXAMPLE

In this section we consider the construction of a flexible OAI Kokotsakis polyhedron with angles of the base quadrilateral:

$$\delta_1 = 1.36292, \quad \delta_2 = 1.41009, \quad \delta_3 = 1.80327, \quad \delta_4 = 2\pi - \delta_1 - \delta_2 - \delta_3 = 1.70691.$$

We selected $\tau = -\arctan 60 = -1.55413$. Following the procedure in [Theorem 5](#), we get

$$\begin{array}{llll} \alpha_1 = 1.34086, & \alpha_2 = 1.42575, & \alpha_3 = 1.69859, & \alpha_4 = 1.81798, \\ \gamma_1 = 1.15746, & \gamma_2 = 2.00166, & \gamma_3 = 1.4875, & \gamma_4 = 1.63656; \\ \beta_1 = 1.11122, & \beta_2 = 1.18397, & \beta_3 = 1.61684, & \beta_4 = 1.68958. \end{array}$$

In [Figure 12](#) we show the state of the given polyhedron when we observe the flattening effect.




FIGURE 12. Partial flattening.

In this case, we have $k = 0.0083$. And we observe the effect of a closeness of parameterization mentioned in Section 3. Hardly one can observe the difference between two parameterizations. However, they are different. And we can calculate numerically the difference




FIGURE 13. Angles $(\varphi, \psi_2, \theta, \psi_1)$ of t for a branch $s_g = -1, s_f = 1$ given by the parameterization of Theorem 2 and by the parameterization of Theorem 3 with linear substitution $t \rightarrow -\frac{K'}{\pi}t - 7.717$.




FIGURE 14. The differences between the dihedral angles in mentioned above parameterizations.

8. APPENDIX: PROPERTIES OF ORTHODIAGONAL QUADRILATERALS

Lemma 8.1. *The diagonals of a spherical quadrilateral with side lengths $\alpha, \beta, \gamma, \delta$ (in this cyclic order) are orthogonal if and only if its side lengths satisfy the relation*

$$\cos \alpha \cos \gamma = \cos \beta \cos \delta.$$

Proof. Take the intersection point of two diagonals of the quadrilateral. (A pair of opposite vertices determines a big circle. We take one of the two intersection points of these two big circles.) Denote the lengths of the segments between the vertices and the intersection point of the diagonals as shown in Figure 15. By the spherical Pythagorean theorem we have $\cos a \cos b = \cos \alpha$, $\cos b \cos c = \cos \beta$, $\cos c \cos d = \cos \gamma$, $\cos d \cos a = \cos \delta$. It follows that

$$\cos \alpha \cos \gamma = \cos a \cos b \cos c \cos d = \cos \beta \cos \delta.$$

□




FIGURE 15. To the proof of Lemma 8.1.

Lemma 8.2. *Let $\alpha, \beta, \gamma, \delta \in (0, \pi)$ satisfy $\cos \alpha \cos \gamma = \cos \beta \cos \delta$. Then there exists a spherical orthodiagonal quadrilateral with side lengths $\alpha, \beta, \gamma, \delta$.*

Proof. Assume that $\alpha + \beta$ is the minimum of the sum of two consecutive angles in $(\alpha, \beta, \gamma, \delta, \alpha)$. If $\delta = \alpha$ and $\beta = \gamma$, the construction is obvious. Otherwise, assume that $\delta > \alpha$. Put sides V_1V_2 and V_2V_3 with lengths α and β respectively on the same great circle. Then there is a vertex V_4 such that $V_1V_2V_4$ is a right angle and the length of V_4V_1 is δ . By the identity, the length of V_3V_4 is γ . □

REFERENCES

- [1] R. Bricard. Mémoire sur la théorie de l'octaèdre articulé. *Journal de Mathématiques pures et appliquées*, 3:113–148, 1897.
- [2] I. Izmetiev. Flexible Kokotsakis polyhedra based on orthodiagonal spherical quadrilaterals. In preparation.
- [3] I. Izmetiev. Classification of flexible Kokotsakis polyhedra with quadrangular base. *International Mathematics Research Notices*, 2017(3):715–808, 2016.
- [4] A. Kokotsakis. Über bewegliche polyeder. *Mathematische Annalen*, 107(1):627–647, 1933.
- [5] G. Nawratil. Flexible octahedra in the projective extension of the Euclidean 3-space. *J. Geom. Graph.*, 14(2):147–169, 2010.
- [6] G. Nawratil. Reducible compositions of spherical four-bar linkages with a spherical coupler component. *Mechanism and Machine Theory*, 46:725–742, 2011.
- [7] G. Nawratil. Reducible compositions of spherical four-bar linkages without a spherical coupler component. *Mechanism and Machine Theory*, 49:87–103, 2012.
- [8] R. Sauer and H. Graf. Über Flächenverbiegung in Analogie zur Verknickung offener Facettenfläche. *Mathematische Annalen*, 105(1):499–535, 1931.
- [9] H. Stachel. A kinematic approach to Kokotsakis meshes. *Comput. Aided Geom. Design*, 27(6):428–437, 2010.

CENTRE FOR SYNTHETIC AND SYSTEMS BIOLOGY,, SCHOOL OF BIOLOGICAL SCIENCES,, UNIVERSITY OF EDINBURGH, EDINBURGH EH9 3BF, UK

E-mail address: `ivan.erofeev@ed.ac.uk`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF FRIBOURG, CHEMIN DU MUSÉE 23, CH-1700 FRIBOURG PÉROLLES, SWITZERLAND

DEPARTMENT OF HIGHER MATHEMATICS, MOSCOW INSTITUTE OF PHYSICS AND TECHNOLOGY, INSTITUTSKII PEREULOK 9, DOLGOPRUDNY, MOSCOW REGION, 141700, RUSSIA.

E-mail address: `grigory.ivanov@unifr.ch`