

Scattering amplitude annihilators

Andrea Nützi^a and Michael Reiterer^b

^a*ETH Zurich, Switzerland*

^b*The Hebrew University of Jerusalem, Israel*

E-mail: andrea.nuetzi@math.ethz.ch, michael.reiterer@protonmail.com

ABSTRACT: Several second order differential operators are shown to annihilate the YM and GR tree scattering amplitudes. In particular we prove a conjecture of Loebbert, Mojaza and Plefka from their investigation of a hidden conformal symmetry in GR.

Contents

1	Introduction	1
2	Preliminaries	5
3	Kinematic variables	7
4	Recursion for the amplitudes	11
5	Some commutators	15
6	Proof of Theorem 1	16
7	Acknowledgments	24
A	Some tree amplitudes	24
B	Mathematica computer code	25
C	Additional formulas	26
D	Relation to Loebbert, Mojaza, Plefka [1]	28

1 Introduction

The scattering amplitudes of Yang-Mills (YM) and general relativity (GR) are interesting objects in physics. The dimension-neutral tree amplitudes are rational functions of many variables; see Appendix A for examples. As with other special functions, it is natural to ask what differential identities they satisfy. We show that a number of second order differential operators annihilate the dimension-neutral YM and GR amplitudes.

This paper is motivated by recent work of Loebbert, Mojaza and Plefka [1]. They investigated a potential hidden conformal symmetry of GR tree amplitudes in general spacetime dimension d . This led them to conjecture certain identities for the dimension-neutral amplitudes that they verified by explicit calculation for amplitudes with $n = 3, 4, 5, 6$ legs¹. They conjectured that they would continue to hold for $n \geq 7$. This paper contains:

- A reformulation of these identities as proper differential identities for the amplitudes. The formulation in [1] is more intricate, as discussed in Appendix D.

¹We thank F. Loebbert, M. Mojaza and J. Plefka for making available computer code to check the $n = 5$ case and for answering questions we had about their paper [1].

- A proof of these identities for all $n \geq 3$, by induction on n .

We do not take up the interesting theme of hidden conformal symmetry [1]. The identities conjectured in [1] and proved here do not readily imply identities for the amplitudes in specific dimensions such as $d = 4$, an interesting question that we leave open.

In this paper we exclusively work with the dimension-neutral version of the tree amplitudes; the spacetime dimension d will be absent. These amplitudes are rational functions on a complex vector space of dimension $2n(n-2)$ with simple poles along a constellation of hyperplanes. The coordinates on this vector space will be denoted k_{ij}, c_{ij}, e_{ij} with subscripts running over an index set² I with $|I| = n$. The following linear relations among the coordinates cut the dimension of this vector space down to $2n(n-2)$:

$$\begin{aligned} k_{ii} &= 0 & k_{ij} - k_{ji} &= 0 & \sum_{i \in I} k_{ij} &= 0 \\ c_{ii} &= 0 & & & \sum_{i \in I} c_{ij} &= 0 \\ e_{ii} &= 0 & e_{ij} - e_{ji} &= 0 & & \end{aligned} \tag{1.1}$$

The amplitudes are actually polynomial in the c_{ij} and e_{ij} variables. To obtain the amplitudes in d spacetime dimensions set $k_{ij} = k_i \cdot k_j$ and $c_{ij} = k_i \cdot \epsilon_j$ and $e_{ij} = (1 - \delta_{ij})\epsilon_i \cdot \epsilon_j$ where k_i and ϵ_i are d -dimensional vectors, the momentum and polarization of the i -th particle.

We need a workable definition of the amplitudes for general n . For the purpose of this paper, the amplitudes are defined by the recursion in Definition 11. This recursion is based on the factorization of residues formulated directly in k_{ij}, c_{ij}, e_{ij} variables³. On the face of it, it is neither obvious that this recursion admits a solution, nor that the solution is unique. A sketch of existence using Feynman rules is in Remark 2, but since we are unaware of a reference that spells this out in detail for the dimension-neutral amplitudes, we logically state existence as Assumption A below. Uniqueness is shown in Lemma 12.

Assumption A. *A sequence of rational functions satisfying the recursion in Definition 11 exists. We refer to them as the YM respectively GR amplitudes.*

In stating our main result below, we avoid standard notation for partial derivatives such as $\partial_{k_{ij}}, \partial_{c_{ij}}, \partial_{e_{ij}}$ because it is ambiguous how these act on functions on the vector space defined by (1.1). To illustrate, if $I = \{1, \dots, 5\}$ then c_{12} and $-c_{32} - c_{42} - c_{52}$ are two representatives of one the same function, but if we act on them with $\partial_{c_{12}}$ using a naive interpretation of the partial derivative, then we obtain conflicting results, 1 and 0 respectively. To fix this we introduce new notation $D_{k_{ij}}, D_{c_{ij}}, D_{e_{ij}}$ which are differential

²The YM amplitude requires a cyclic order on I , whereas the GR amplitude is permutation invariant and I is an unordered set. We will not mention this further in this introduction.

³The original recursion of this kind is BCFW recursion [2]. The recursion we use in this paper does not involve BCFW shifts nor decay conditions under such shifts, it is only required that the amplitudes factor properly. The proof that this determines the amplitudes uniquely is along the lines of [6], which is for $d = 4$, but is simpler in the present dimension-neutral case since one is on a vector space and all poles are along linear subspaces. It is useful in this paper to have the uniqueness statement in a simple form since the same argument (cf. Lemmas 12 and 13) also plays a key role in proof of the main theorem, Theorem 1.

operators that act unambiguously on functions on the vector space defined by (1.1). To illustrate, if $I = \{1, \dots, 5\}$ then we use $D_{c_{12}} = \partial_{c_{12}} - \frac{1}{4}(\partial_{c_{12}} + \partial_{c_{32}} + \partial_{c_{42}} + \partial_{c_{52}})$ which acts unambiguously. The $D_{k_{ij}}$ are linear combinations of the $\partial_{k_{ij}}$ with constant coefficients; the $D_{c_{ij}}$ are linear combinations of the $\partial_{c_{ij}}$ with constant coefficients; the $D_{e_{ij}}$ are linear combinations of the $\partial_{e_{ij}}$ with constant coefficients. Their complete definition is in Lemma 3, and we have also included computer code in Appendix B. We note that our definition of the D -operators is also a matter of convention; we have merely made a convenient choice of operators that span the $2n(n-2)$ -dimensional space of constant first order differential operators along the vector space defined by (1.1).

There are $2n+1$ first order differential operators that are well-known to annihilate the amplitudes: n of them express gauge invariance; n express homogeneity in the polarizations; and one expresses homogeneity jointly in all momenta. They are, respectively,

$$X_i = \sum_{j \in I} (k_{ji} D_{c_{ji}} + c_{ij} D_{e_{ij}}) \quad (1.2a)$$

$$Y_i = \sum_{j \in I} (c_{ji} D_{c_{ji}} + e_{ij} D_{e_{ij}}) - h \quad (1.2b)$$

$$Z = \sum_{i,j \in I} (k_{ij} D_{k_{ij}} + c_{ij} D_{c_{ij}}) - s \quad (1.2c)$$

where $h = 1$ and $s = 4 - n$ for YM respectively $h = 2$ and $s = 2$ for GR. Polynomiality in the c_{ij} and e_{ij} variables implies further obvious annihilators. There are, in addition, several second order annihilators as we show in this paper.

Theorem 1. *Define the YM and GR tree amplitude by the recursion in Definition 11 and make Assumption A. The differential operators A_i , B_i , C_i defined below annihilate the tree amplitudes for all $n = |I| \geq 3$ and $i \in I$:*

$$A_i = \sum_{j,k \in I} \left(\frac{1}{2} k_{jk} D_{c_{ji}} D_{c_{ki}} + c_{jk} D_{c_{ji}} D_{e_{ki}} + \frac{1}{2} e_{jk} D_{e_{ji}} D_{e_{ki}} \right) \quad (1.3a)$$

and

$$B_i = \sum_{j,k \in I} \left(c_{jk} D_{k_{ij}} D_{e_{ik}} + e_{jk} D_{c_{ij}} D_{e_{ik}} + k_{jk} D_{k_{ij}} D_{c_{ki}} + c_{kj} D_{c_{ij}} D_{c_{ki}} \right. \\ \left. - c_{kj} D_{k_{jk}} D_{e_{ji}} - e_{jk} D_{c_{jk}} D_{e_{ji}} - k_{jk} D_{k_{jk}} D_{c_{ji}} - c_{jk} D_{c_{ji}} D_{c_{jk}} \right) \quad (1.3b)$$

and $C_i = \tilde{C}_i - \frac{1}{n} \sum_{j \in I} \tilde{C}_j$ where

$$\tilde{C}_i = \sum_{j,k \in I} \left(\frac{1}{2} e_{jk} D_{c_{ij}} D_{c_{ik}} + c_{kj} D_{k_{ik}} D_{c_{ij}} + \frac{1}{2} k_{jk} D_{k_{ij}} D_{k_{ik}} \right. \\ \left. - e_{jk} D_{c_{ij}} D_{c_{jk}} - c_{kj} D_{k_{jk}} D_{c_{ij}} - c_{jk} D_{k_{ji}} D_{c_{jk}} - k_{jk} D_{k_{ji}} D_{k_{jk}} \right) \quad (1.3c)$$

The D -operators are defined in Lemma 3.

That B_i , C_i annihilate the GR amplitude was conjectured in [1], using a more intricate formulation that we discuss in Appendix D. The formulation of these identities directly as annihilating differential operators in Theorem 1 is therefore new⁴. Our proof of Theorem 1 is by induction on n and is summarized later in this introduction.

⁴It allows us, for instance, to compute a number of commutators in Section 5. They imply that every function annihilated by X_i , Y_i , Z , C_i is automatically annihilated by A_i , B_i .

Remark 1. The amplitudes in $d \geq 4$ spacetime dimensions are obtained from the dimension-neutral ones by setting $k_{ij} = k_i \cdot k_j$ and $c_{ij} = k_i \cdot \epsilon_j$ and $e_{ij} = (1 - \delta_{ij})\epsilon_i \cdot \epsilon_j$ where k_i and ϵ_i are elements of a d -dimensional complex vector space, and \cdot is a nondegenerate symmetric bilinear pairing⁵. The momentum vectors k_i are subject to $k_i \cdot k_i = 0$ and momentum conservation $\sum_i k_i = 0$, and the polarization vectors ϵ_i are subject to $k_i \cdot \epsilon_i = 0$. Assume $k_i \neq 0$ here, so ϵ_i lies in a subspace of dimension $d - 1$. For every i , the YM amplitude is linear in ϵ_i , the GR amplitude is quadratic in ϵ_i ; this is witnessed by the annihilator Y_i . For every $k \neq 0$ with $k \cdot k = 0$ abbreviate $P(k) = \{\epsilon \mid k \cdot \epsilon = 0\}/\mathbb{C}k$ which is a vector space of dimension $d - 2$, and observe that \cdot induces a nondegenerate symmetric bilinear pairing on $P(k)$. Then, separately for every i and fixed $k_i \neq 0$:

- As a function of ϵ_i , the YM amplitude descends to a linear form on $P(k_i)$.
- As a function of ϵ_i , the GR amplitude descends to a quadratic form on $P(k_i)$ ⁶.

This is witnessed by X_i and is known as gauge invariance. The amplitudes are also homogeneous jointly in all momenta, as witnessed by Z . In summary, the d -dimensional amplitudes are sections of certain vector bundles on the projective variety $k_i \cdot k_i = 0$ and $\sum_i k_i = 0$ ^{7,8,9}. In this paper we do not work with momenta k_i and polarizations ϵ_i . Our primary variables are k_{ij} , c_{ij} , e_{ij} and we exploit the fact that the relations (1.1) are linear.

Let us outline the proof of Theorem 1, which is by brute force by induction on n . It uses the recursive definition of the amplitudes, by which the amplitudes have simple poles along a constellation of hyperplanes, each residue being equal to a product of two lower amplitudes. Denoting by M_I the GR amplitude for index set I , schematically one has

$$\text{Res } M_I = U^2 M_{J \sqcup \{\bullet\}} M_{K \sqcup \{\bullet\}} \quad (1.4)$$

for every decomposition $I = J \sqcup K$ into two subsets of two or more elements, $|J| \geq 2$ and $|K| \geq 2$. The residue is taken along the locus where $k_J = \sum_{j \in J} k_j$ satisfies $k_J \cdot k_J = 0$. Translating to the coordinates we use, this locus is the hyperplane

$$\xi \stackrel{\text{def}}{=} \sum_{j,j' \in J} k_{jj'} = 0 \quad (1.5)$$

On the right hand side of (1.4), the bullet \bullet stands for one more particle. Properly defining the right hand side of (1.4) requires contracting the polarizations of the respective \bullet particles

⁵For instance, \mathbb{C}^d with standard pairing $z \cdot w = z_1 w_1 + \dots + z_d w_d$. The choice does not matter since all such vector spaces with nondegenerate pairing are isomorphic.

⁶Equivalently, a linear form on the symmetric tensor product $S^2 P(k_i)$. One can decompose this into the trace and the traceless part relative to the symmetric bilinear pairing on $P(k_i)$.

⁷Actually one only has a vector bundle away from the singular locus.

⁸It would be interesting to see whether the annihilators A_i , B_i , C_i or suitable linear combinations of them imply annihilators for the d -dimensional amplitudes. The latter would be differential operators on the vector bundle on which the amplitudes live, possibly taking values in another vector bundle.

⁹The fact that A_i annihilates is vacuous for YM, since A_i involves two derivatives with respect to the polarization of the i -th particle. It is not vacuous for GR amplitudes, in the sense that one can write down a rational function annihilated by the obvious annihilators listed before Theorem 1 but not by A_i .

in $M_{J \sqcup \{\bullet\}}$ and $M_{K \sqcup \{\bullet\}}$. This contraction is effected by U^2 , using the differential operator U in Definition 9. Suppose now we are at the n -th induction step. Within each induction step, one first shows that the A_i annihilate, then the B_i , then the C_i . The structure of the argument is analogous in each case, so suppose for concreteness that we want to show $C_i M_I = 0$ for some i . This is done in two steps:

1. Show that $C_i M_I$ has no pole. Namely, for every decomposition $I = J \sqcup K$ as above one must check that $C_i M_I$ has no pole along the hyperplane $\xi = 0$ in (1.5).
2. Show that the complete absence of poles in $C_i M_I$, together with other known properties of $C_i M_I$ such as homogeneity properties, imply $C_i M_I = 0$.

In Step 1, note that M_I has a simple ξ^{-1} type pole, so C_i being second order, the application $C_i M_I$ can naively have a third order pole, terms of type ξ^{-3} and ξ^{-2} and ξ^{-1} . While the ξ^{-3} term is easily seen to be absent, the absence of ξ^{-2} and ξ^{-1} terms is by a lengthy explicit calculation that exploits only the annihilators of $M_{J \sqcup \{\bullet\}}$ and $M_{K \sqcup \{\bullet\}}$ that are known by the induction hypothesis. In a nutshell, and using schematic notation again, in Step 1 one must check that $C_i(\frac{1}{\xi} U^2 M_{J \sqcup \{\bullet\}} M_{K \sqcup \{\bullet\}})$ has no pole. The computations establishing Step 1 are in the most technical lemma of this paper, Lemma 16. Step 2 relies on Lemma 13 (which is the same lemma that also establishes that the recursion determines the amplitudes uniquely) and on the commutators in Lemma 14. The YM case is entirely analogous.

The proof does not use explicit formulas for the amplitudes. Instead, the recursion for the amplitudes using (1.4) is effectively turned into a recursion for the annihilators.

One can ask what the full annihilator of the YM respectively GR tree amplitudes is. To every rational function one can associate its annihilator, a left ideal in the Weyl algebra of polynomial differential operators. Rational functions are holonomic, meaning their annihilator is as big as allowed by the Bernstein inequality¹⁰. In a Weyl algebra, every left ideal is finitely generated¹¹. Algorithms are available to determine the annihilator of a rational function, e.g. in Macaulay2 [3, 4], and while in principle such tools can be applied to tree amplitudes, this approach does not seem to be practically viable for general n .

It would be interesting to understand if there is a more conceptual explanation for these annihilators perhaps building on ideas in [1]; to inquire if there are corresponding identities for the amplitudes in specific dimensions such as $d = 4$; and to try to apply these annihilators to derive properties of the amplitudes.

2 Preliminaries

Notation. All vector spaces and algebras are over the complex numbers. For every finite-dimensional vector space X we denote:

¹⁰This says that over \mathbb{R} or \mathbb{C} , and for a proper left ideal $I \subseteq D$ in the Weyl algebra D in $x_1, \dots, x_m, \partial_1, \dots, \partial_m$, the dimension of D/I (defined to be the degree of a suitable Hilbert polynomial) is $\geq m$. Note that a rational function g/f always has m first order annihilators of the form $f(\partial_i g) - g(\partial_i f) - gf\partial_i \in D$ but they are in general of high polynomial degree. See also [5].

¹¹Somewhat surprisingly, by a theorem of Stafford, every left ideal in a Weyl algebra over \mathbb{R} or \mathbb{C} can in fact be generated by just two elements.

X^*	the dual vector space
R_X	the commutative algebra of polynomials $X \rightarrow \mathbb{C}$
D_X	the Weyl algebra on X
$\text{Frac}(R_X)$	the field of rational functions on X

These things are defined independently of coordinates. This is important because the spaces we encounter do not have a canonical coordinate system, and we work with a linearly dependent set of coordinates. Canonically,

$X^* \subseteq R_X$	the linear functionals, we refer to them as coordinates
$X \subseteq D_X$	first order constant coefficient differential operators

As vector spaces, $R_X \simeq SX^*$ and $D_X \simeq SX^* \otimes SX$ where S is the symmetric tensor algebra. Here SX are the constant coefficient differential operators.

Coordinate dependent definitions. Even though we do not commit to a particular basis, we recall the coordinate dependent definitions. By a coordinate we mean an element of X^* . A coordinate system is a basis $x_1, \dots, x_m \in X^*$. The polynomial ring is then

$$R_X = \mathbb{C}[x_1, \dots, x_m]$$

Let $\partial_1, \dots, \partial_m \in X$ be the dual basis. Then D_X is the associative algebra with identity generated by the variables $x_1, \dots, x_m, \partial_1, \dots, \partial_m$ modulo the two-sided ideal generated by

$$x_i x_j - x_j x_i \quad \partial_i \partial_j - \partial_j \partial_i \quad \partial_i x_j - x_j \partial_i - \delta_{ij}$$

Linear maps. If X, Y are vector spaces then every linear map $\alpha : X \rightarrow Y$ canonically induces several other linear maps:

$SX \rightarrow SY$	the push-forward, also denoted α
$Y^* \rightarrow X^*$	the adjoint map, denoted α^*
$R_Y \rightarrow R_X$	the pull-back, also denoted α^*
$\text{Frac}(R_Y) \rightarrow \text{Frac}(R_X)$	the pull-back, also denoted α^*

It neither induces a map $D_X \rightarrow D_Y$ nor $D_Y \rightarrow D_X$. We often find it convenient to specify a linear map by specifying the adjoint $Y^* \rightarrow X^*$.

Direct sums. For a direct sum of finite-dimensional vector spaces, there are canonical isomorphisms $R_{X \oplus Y} \simeq R_X \otimes R_Y$ and $D_{X \oplus Y} \simeq D_X \otimes D_Y$ that we frequently use. This uses the tensor product of algebras, where all elements of D_X commute with all elements of D_Y . Canonical inclusions such as $D_X \hookrightarrow D_X \otimes D_Y, \delta \mapsto \delta \otimes 1$ are sometimes used.

Index sets. Instead of a standard index set such as $\{1, \dots, n\}$ we work with finite index sets denoted I , hence n is replaced by $|I|$. For YM, a cyclic ordering of the elements of I is required. In many calculations, I is a disjoint union

$$I = J \sqcup K \tag{2.1a}$$

with $|J|, |K| \geq 2$ and often (2.1a) is assumed. Sometimes we need index sets with a distinguished element, always denoted \bullet . We abbreviate

$$J_\bullet = J \sqcup \{\bullet\} \quad K_\bullet = K \sqcup \{\bullet\} \quad (2.1b)$$

If J_\bullet, K_\bullet have a cyclic order then $J \sqcup K$ acquires a cyclic order¹². Conversely, if I has a cyclic order and (2.1a) respects this, then J_\bullet, K_\bullet inherit a cyclic order.

3 Kinematic variables

Here we define in detail the vector space on which the amplitudes live; the differential operators D used in Theorem 1; and notation that allows one to state the schematic factorization of residues in (1.4) in a precise way in Definition 11.

Definition 2. *For every finite set I with $|I| \geq 3$, consider first the complex ‘ambient’ vector space of dimension $3|I|^2$ with coordinate system*

$$k_{ij}, c_{ij}, e_{ij} \quad \text{where} \quad i, j \in I \quad (3.1)$$

Denote by $C(I)$ the linear subspace defined by the relations (1.1). In this paper, the elements (3.1) are understood to be elements of the dual space $C(I)^*$.

The dimension of $C(I)$ is

$$\underbrace{\frac{1}{2}n(n-3)}_k + \underbrace{n(n-2)}_c + \underbrace{\frac{1}{2}n(n-1)}_e = 2n(n-2)$$

where $n = |I|$. If $|I| = 3$ then the k_{ij} vanish identically as elements of $C(I)^*$. Note that the relations (1.1) do not refer to an ordering, hence there is a natural action of the group of permutations of I on the vector space $C(I)$.

The (3.1) are a linearly dependent set of coordinates, not a coordinate system on $C(I)$. Therefore we cannot define partial derivatives in the usual way. We work with the following linearly dependent set of constant coefficient differential operators.

Lemma 3. *There are unique*

$$D_{k_{ij}}, D_{c_{ij}}, D_{e_{ij}} \in C(I) \quad (3.2)$$

that as elements of the Weyl algebra $D_{C(I)}$ satisfy

$$\begin{aligned} [D_{k_{ij}}, k_{ab}] &= (1 - \delta_{ij})(1 - \delta_{ab})(\delta_{ia}\delta_{jb} + \delta_{ib}\delta_{ja} - \frac{1}{|I|-2}(\delta_{ia} + \delta_{ib} + \delta_{ja} + \delta_{jb}) + \frac{2}{(|I|-1)(|I|-2)}) \\ [D_{c_{ij}}, c_{ab}] &= (1 - \delta_{ij})(1 - \delta_{ab})(\delta_{ia}\delta_{jb} - \frac{1}{|I|-1}\delta_{jb}) \\ [D_{e_{ij}}, e_{ab}] &= (1 - \delta_{ij})(1 - \delta_{ab})(\delta_{ia}\delta_{jb} + \delta_{ib}\delta_{ja}) \end{aligned}$$

¹²If the elements of J_\bullet are $\bullet, a_1, \dots, a_{|J|}$ and the elements of K_\bullet are $\bullet, b_1, \dots, b_{|K|}$ listed in cyclic order, starting with \bullet for convenience, then on $J \sqcup K$ use the cyclic order $a_1, \dots, a_{|J|}, b_1, \dots, b_{|K|}$.

and such that all ‘mixed’ commutators are zero:

$$\begin{aligned} [D_{k_{ij}}, c_{ab}] &= 0 & [D_{c_{ij}}, k_{ab}] &= 0 & [D_{e_{ij}}, k_{ab}] &= 0 \\ [D_{k_{ij}}, e_{ab}] &= 0 & [D_{c_{ij}}, e_{ab}] &= 0 & [D_{e_{ij}}, c_{ab}] &= 0 \end{aligned}$$

They span $C(I)$. They satisfy

$$\begin{aligned} D_{k_{ii}} &= 0 & D_{k_{ij}} &= D_{k_{ji}} & \sum_i D_{k_{ij}} &= 0 \\ D_{c_{ii}} &= 0 & & & \sum_i D_{c_{ij}} &= 0 \\ D_{e_{ii}} &= 0 & D_{e_{ij}} &= D_{e_{ji}} & & \end{aligned} \tag{3.3}$$

Proof. The given commutators for $D_{k_{ij}}$ at first only determine an operator on the ambient vector space. But since $[D_{k_{ij}}, -]$ annihilates all left hand sides of the relations (1.1), we obtain a unique constant coefficient differential operator $\in C(I)$ as claimed. The rest of the lemma goes the same way. \square

We have introduced coordinates k_{ii} , c_{ii} , e_{ii} and derivatives $D_{k_{ii}}$, $D_{c_{ii}}$, $D_{e_{ii}}$ that are identically zero and in some sense superfluous. These phantom objects are nevertheless useful because they allow us to write sums as in (1.3).

Convention (2.1) is in force, $I = J \sqcup K$. To this decomposition of I we associate a coordinate ξ , and a vector Ξ transversal to the hyperplane $\xi = 0$. The dependence of these definitions on the decomposition $I = J \sqcup K$ is left implicit in the notation.




Figure 1: Decomposition of $C(I)$ associated to the decomposition $I = J \sqcup K$. The coordinates k^\perp , c^\perp , e^\perp are uniquely determined by being constant in the direction Ξ and coinciding with k , c , e respectively along $C(I)^\perp$. The derivatives D^\perp span $C(I)^\perp$.

Definition 4. Define $\xi \in C(I)^*$ and $\Xi \in C(I)$ by¹³

$$\xi = \sum_{j,j' \in J} k_{jj'} = \sum_{k,k' \in K} k_{kk'} \tag{3.4}$$

$$\Xi = \frac{1}{|J|(|J|-1)} \sum_{j,j' \in J} D_{k_{jj'}} - \frac{2}{|J||K|} \sum_{j \in J, k \in K} D_{k_{jk}} + \frac{1}{|K|(|K|-1)} \sum_{k,k' \in K} D_{k_{kk'}} \tag{3.5}$$

Also define

$$\begin{aligned} k_{jj'}^\perp &= k_{jj'} - \frac{1-\delta_{jj'}}{|J|(|J|-1)} \xi & D_{k_{jj'}}^\perp &= D_{k_{jj'}} - \frac{|K|(|K|-1)(1-\delta_{jj'})}{(|J|-1)(|I|-2)} \Xi \\ k_{jk}^\perp &= k_{kj}^\perp = k_{jk} + \frac{1}{|J||K|} \xi & D_{k_{jk}}^\perp &= D_{k_{kj}}^\perp = D_{k_{jk}} + \frac{(|K|-1)(|J|-1)}{(|J|-1)(|I|-2)} \Xi \end{aligned}$$

¹³One can equivalently define Ξ by $\Xi = \text{const} \cdot \sum_{j,j' \in J} D_{k_{jj'}}$ normalized so that $[\Xi, \xi] = 2$.

$$\begin{aligned}
k_{kk'}^\perp &= k_{kk'} - \frac{1-\delta_{kk'}}{|K|(|K|-1)}\xi & D_{k_{kk'}}^\perp &= D_{k_{kk'}} - \frac{|J|(|J|-1)(1-\delta_{kk'})}{(|I|-1)(|I|-2)}\Xi \\
c_{ii'}^\perp &= c_{ii'} & D_{c_{ii'}}^\perp &= D_{c_{ii'}} \\
e_{ii'}^\perp &= e_{ii'} & D_{e_{ii'}}^\perp &= D_{e_{ii'}}
\end{aligned}$$

for all $j, j' \in J$ and $k, k' \in K$ and $i, i' \in I = J \sqcup K$. Set $C(I)^\perp = \{\xi = 0\}$.

Lemma 5. *We have*

$$C(I) = \mathbb{C}\Xi \oplus C(I)^\perp$$

The D^\perp span $C(I)^\perp$. In the Weyl algebra $D_{C(I)}$,

$$\begin{aligned}
[\Xi, \xi] &= 2 & [\Xi, k_{jj'}] &= \frac{2(1-\delta_{jj'})}{|J|(|J|-1)} \\
[\Xi, k^\perp] &= [\Xi, c^\perp] = [\Xi, e^\perp] = 0 & [\Xi, k_{jk}] &= -\frac{2}{|J||K|} \\
[D^\perp, \xi] &= 0 & [\Xi, k_{kk'}] &= \frac{2(1-\delta_{kk'})}{|K|(|K|-1)}
\end{aligned}$$

for all $j, j' \in J$ and $k, k' \in K$. The $k^\perp, c^\perp, e^\perp$ and D^\perp satisfy the linear relations (1.1) and (3.3), with k replaced by k^\perp and so forth. They additionally satisfy

$$\sum_{j, j' \in J} k_{jj'}^\perp = 0 \quad \sum_{j, j' \in J} D_{k_{jj'}}^\perp = 0 \quad (3.6)$$

Same if the summation is over $k, k' \in K$ instead.

Proof. By direct calculation. \square

Let us pause to discuss what is needed to state the factorization (1.4) precisely. Essentially we need maps between the $\xi = 0$ hyperplane in $C(I)$ which is the locus of the pole under consideration on the one hand, and the spaces $C(J_\bullet)$ and $C(K_\bullet)$ where the two amplitudes live on the other hand. Here are more details:

- Since $C(I)$ lacks variables corresponding to the polarization of the \bullet particle, we use an extension $M = C(I) \oplus E$ where E provides the missing variables for \bullet .
- We define maps not only along $\xi = 0$ (now understood to be a hyperplane in M) but on all of M . This leads to two linear, surjective maps

$$\begin{array}{ccc}
& M & \\
\alpha_J \swarrow & & \searrow \alpha_K \\
C(J_\bullet) & & C(K_\bullet)
\end{array} \quad (3.7)$$

The extension to $\xi \neq 0$ is not needed to state the recursion in Definition 11, but it is exploited in the statement and proof of the technical Lemma 16.

- Along $\xi = 0$ the maps α_J and α_K are defined in the canonical way, by identifying the momenta k_i and polarizations ϵ_i with the corresponding ones in J_\bullet when $i \in J$ respectively K_\bullet when $i \in K$, and by translating this to the variables (3.1), cf. Remark 1. Together with the linear constraints, this defines the maps completely along $\xi = 0$. The extension to $\xi \neq 0$ is such that α_J and α_K are constant along Ξ .

- Finally, a differential operator U is introduced on M whose purpose is to contract the polarization of the particle \bullet in J_\bullet with the corresponding particle \bullet in K_\bullet .

In the remainder of this section, we spell these things out.

Definition 6 (Extension space). *Consider first the complex ‘ambient’ vector space of dimension $2|J| + 2|K| = 2|I|$ with coordinate system¹⁴*

$$c_{j\bullet}, e_{j\bullet}, c_{k\bullet}, e_{k\bullet} \quad \text{where} \quad j \in J, k \in K \quad (3.8)$$

Let E be the subspace defined by

$$\sum_{j \in J} c_{j\bullet} = 0 \quad \sum_{k \in K} c_{k\bullet} = 0$$

a linear subspace of dimension $2|J| + 2|K| - 2 = 2|I| - 2$. The elements (3.8) are understood to be elements of the dual space E^* . There are unique

$$D_{c_{j\bullet}}, D_{e_{j\bullet}}, D_{c_{k\bullet}}, D_{e_{k\bullet}} \in E$$

that as elements of the Weyl algebra D_E satisfy

$$\begin{aligned} [D_{c_{j\bullet}}, c_{j'\bullet}] &= \delta_{jj'} - \frac{1}{|J|} & [D_{c_{k\bullet}}, c_{k'\bullet}] &= \delta_{kk'} - \frac{1}{|K|} \\ [D_{e_{j\bullet}}, e_{j'\bullet}] &= \delta_{jj'} & [D_{e_{k\bullet}}, e_{k'\bullet}] &= \delta_{kk'} \end{aligned}$$

for $j, j' \in J$ and $k, k' \in K$ and all other ‘mixed’ commutators are zero¹⁵.

Definition 7 (Master space). *The master space corresponding to $I = J \sqcup K$ is*

$$M = C(I) \oplus E$$

with dimension $2n^2 - 2n - 2$, where $n = |I|$. Its Weyl algebra is therefore

$$D_M = D_{C(I)} \otimes D_E$$

We agree that elements of $D_{C(I)}$ and D_E are extended to D_M . For example, Ξ is also viewed as an element of D_M from now on. Set $M^\perp = C(I)^\perp \oplus E$, so

$$M = \mathbb{C}\Xi \oplus M^\perp \quad (3.9)$$

Lemma 8. *There is a unique linear map α_J whose adjoint $\alpha_J^* : C(J_\bullet)^* \rightarrow M^*$ maps¹⁶*

$$\begin{aligned} k_{jj'} &\mapsto k_{jj'}^\perp & c_{jj'} &\mapsto c_{jj'}^\perp & e_{jj'} &\mapsto e_{jj'}^\perp \\ k_{j\bullet}, k_{\bullet j} &\mapsto -\sum_{j' \in J} k_{j'j}^\perp & c_{\bullet j} &\mapsto -\sum_{j' \in J} c_{j'j}^\perp & & \\ e_{j\bullet}, e_{\bullet j} &\mapsto e_{j\bullet} & & & & \end{aligned}$$

for all $j, j' \in J$. Analogous for α_K . The maps α_J, α_K are surjective.

¹⁴In this instance, the bullet \bullet is nothing but suggestive decoration.

¹⁵That is, commutators mixing c and e or mixing J indices with K indices are zero.

¹⁶On the right hand side, the variables $k^\perp, c^\perp, e^\perp$ are in $C(I)^*$ whereas the variables $c_{j\bullet}, e_{j\bullet}$ are in E^* .

Proof. We have specified α_j^* on a set that spans $C(J_\bullet)^*$, hence it exists and is unique if it respects all relations. For instance, $\sum_{j' \in J} c_{j'j} + c_{\bullet j}$ is zero in $C(J_\bullet)^*$ and must be mapped to zero, which it is. More interestingly, $\sum_{j \in J} k_{\bullet j}$ is zero in $C(J_\bullet)^*$ and must be mapped to zero, and it is by (3.6). The last example shows that one must use k^\perp , not k , on M^* . Surjectivity follows from the explicit right-inverses constructed in Appendix C. \square

Definition 9 (Contraction operator). Define $U \in D_M$ by

$$U = \sum_{j \in J, k \in K} \left(k_{jk}^\perp D_{c_j \bullet} D_{c_k \bullet} + c_{jk}^\perp D_{c_j \bullet} D_{e_k \bullet} + c_{kj}^\perp D_{e_j \bullet} D_{c_k \bullet} + e_{jk}^\perp D_{e_j \bullet} D_{e_k \bullet} \right)$$

4 Recursion for the amplitudes

Here we define tree scattering amplitudes for YM, denoted A_I , and for GR, denoted M_I . The 3-point scattering amplitudes are polynomial.

Definition 10 (Base case). If $|I| = 3$ define $A_I, M_I \in R_{C(I)}$ by

$$\begin{aligned} A_I &= c_{12}e_{31} + c_{23}e_{12} + c_{31}e_{23} \\ M_I &= (A_I)^2 \end{aligned}$$

where 1, 2, 3 are shorthands for the three elements of I .

For higher amplitudes we state a recursion. That this recursion admits a solution at all is left as an assumption, Assumption A, but a rough sketch of a proof is in Remark 2. The purpose of this section is to state the recursion, and to show that if a solution exists, then it is unique. (See [6] for a self-contained treatment of amplitudes in $d = 4$.)

Roughly, the recursion says that the amplitudes are rational functions with only simple poles. The poles are along a known family of hyperplanes, and the residues are given recursively in terms of products of lower amplitudes. This is known as factorization of residues. The case $|I| = 3$ serves as the base case.

Define X_i, Y_i, Z as in (1.2) using the differential operators in Lemma 3.

Definition 11 (Recursion). By YM respectively GR tree amplitudes we mean a sequence of rational functions, one for every integer $n \geq 3$. Using an index set I with $n = |I|$, these rational functions are denoted, respectively,

$$A_I, M_I \in \text{Frac}(R_{C(I)})$$

It is understood that I must have a cyclic order for the YM amplitudes A_I but no order for GR amplitudes M_I . It is understood that I is merely an index, so if $I \simeq I'$ is a bijection, preserving the cyclic order in the case of YM, then $A_I \simeq A_{I'}$ respectively $M_I \simeq M_{I'}$ by relabeling coordinate indices. With these preliminaries, the following properties must hold:

- Base case. For $n = 3$ the amplitudes are given by Definition 10.
- Polynomiality in c and e . The amplitudes A_I , M_I are polynomial in the c - and e -variables. So they are polynomials in the c - and e -variables with coefficients that are rational functions of the k -variables.
- Permutation symmetry. A_I is invariant under cyclic permutations of I , whereas M_I is invariant under all permutations of I .
- Gauge invariance. For every $i \in I$, both A_I and M_I are annihilated by X_i .
- Polarization homogeneity. For every $i \in I$ define a grading on $C(I)^*$ by

$$|k_{ab}| = 0 \quad |c_{ab}| = \delta_{bi} \quad |e_{ab}| = \delta_{ai} + \delta_{bi}$$

for all $a \neq b$. Then A_I has degree 1 and M_I has degree 2. That is, they are annihilated by Y_i with $h = 1$ and $h = 2$ respectively.

- Momentum homogeneity. Define a grading on $C(I)^*$ by

$$|k_{ab}| = 2 \quad |c_{ab}| = 1 \quad |e_{ab}| = 0$$

for all $a \neq b$. Then A_I has degree $4 - n$ and M_I has degree 2. That is, they are annihilated by Z with $s = 4 - n$ and $s = 2$ respectively.

- Poles and residues. Both A_I and M_I have poles only along the subspaces $\xi = 0$ corresponding to decompositions $I = J \sqcup K$ with $|J|, |K| \geq 2$. In the case of A_I , there are poles only for decompositions that respect the cyclic order, thus a cyclic order is induced on J_\bullet and K_\bullet respectively. The poles are simple¹⁷ with residue

$$\text{Res}_{\xi=0} A_I = \pm \text{const} \cdot U \alpha_J^*(A_{J_\bullet}) \alpha_K^*(A_{K_\bullet}) \Big|_{\xi=0} \quad (4.1a)$$

$$\text{Res}_{\xi=0} M_I = \pm \text{const} \cdot U^2 \alpha_J^*(M_{J_\bullet}) \alpha_K^*(M_{K_\bullet}) \Big|_{\xi=0} \quad (4.1b)$$

as an identity in $\text{Frac}(R_M/\xi R_M)$. The operator U is in Definition 9. The specification of the residue is recursive because $|J_\bullet|, |K_\bullet| < |I|$.

In (4.1), the role of U respectively U^2 is to contract the polarizations of the \bullet particles that the amplitudes on the right hand side depend on.

Remark 2 (Existence, rough sketch of proof). The YM amplitudes A_I can be defined using color-ordered Feynman rules in the Lorenz-Feynman gauge, see e.g. [7], which yields rational functions of the products $k_i \cdot k_j$, $k_i \cdot \epsilon_j$, $\epsilon_i \cdot \epsilon_j$ as required. It is ‘well-known’ that this has all the properties in Definition 11, including gauge invariance and residue factorization. The translation of residue factorization to the variables (3.1) yields in particular

¹⁷So the amplitudes are in the principal fractional ideal generated by $1/(\prod \xi)$ where the product runs over all decompositions $I = J \sqcup K$ and ξ is defined by (3.4).

the operator U to carry out the sum over polarizations of the \bullet particles. Given the A_I , the GR amplitudes M_I can be defined using the field theoretical Kawai-Lewellen-Tye or KLT relations, see for instance [9], but then some properties in Definition 11 are not immediate, for instance permutation symmetry and the factorization of residues. These properties are discussed in some detail in the ‘purely field theoretical view’ of the KLT relations in [9], which exploits various known properties of the YM amplitudes and of the KLT kernels.

Lemma 12 (Uniqueness). *If amplitudes as in Definition 11 exist, then they are unique.*

Proof. The proof is not sensitive to the constants in (4.1), part of which is a matter of normalization. The proof is by induction on $n = |I|$. The $n = 3$ amplitudes are fixed. For $n \geq 4$, the residues of any two candidate amplitudes are the same by (4.1) and by the induction hypothesis. So the difference between any two candidates, call it $u \in \text{Frac}(R_{C(I)})$, has no poles and is regular, except perhaps where the poles intersect. But the union of all pairwise intersections of poles has codimension two, and by the Hartogs extension theorem (cf. [6] for more comments about this in the context of $d = 4$ amplitudes), u is actually globally regular. This means that $u \in R_{C(I)}$, a polynomial. Since u is annihilated by X_i and Y_i and Z , Lemma 13 below implies $u = 0$. \square

Lemma 13 below is not only used to prove uniqueness in Lemma 12, it will also play a key role in our proof of Theorem 1, hence the additional parameters p_i and q .

Lemma 13 (Vanishing lemma). *Suppose $n = |I| \geq 4$ and suppose X_i, Y_i, Z are as in (1.2), either with the parameters for YM or with the parameters for GR. Suppose $p_i \geq 0$ with $i \in I$ are integers with $\sum_{i \in I} p_i < n$, and $q \geq 0$ is another integer. Suppose a polynomial $u \in R_{C(I)}$ satisfies:*

$$X_i u = 0 \quad (Y_i + p_i)u = 0 \quad (Z + q)u = 0$$

for all $i \in I$. Then $u = 0$.

Proof. Distinguish the cases in Table 1. The schematic structure refers to u as a polynomial in the c, k variables with coefficients that are polynomials in the e variables. The schematic structure follows from $(Z + q)u = 0$. The polynomials $P(e)$ and $Q(e)$ are homogeneous. Their homogeneity degrees are given in terms of $p = \sum_{i \in I} p_i < n$ and follow from $\sum_{i \in I} (Y_i + p_i)u = 0$. Note that P is a single homogeneous polynomial in case 3, whereas P and Q are schematic for several homogeneous polynomials in cases 1 and 2. Since $p < n$ we have $\deg P > 0$, $\deg Q > 0$, in particular P, Q cannot be constant, so they are zero if their derivatives are zero. We discuss each case:

- Case 4: Here $u = 0$.
- Case 3: Here $P(e)$ is a single polynomial. Since $X_i u = 0$ for all i , we have

$$\sum_{j \in I} c_{ij} D_{e_{ij}} P = 0$$

case	situation	schematic structure of u	$\deg P$	$\deg Q$
1	GR, $q = 0$	$P(e)cc + Q(e)k$	$n - \frac{p}{2} - 1$	$n - \frac{p}{2}$
2	GR, $q = 1$	$P(e)c$	$n - \frac{p}{2} - \frac{1}{2}$	
3	GR, $q = 2$	$P(e)$	$n - \frac{p}{2}$	
4	GR, $q > 2$	0		
3	YM, $n = 4$ and $q = 0$	$P(e)$	$\frac{n}{2} - \frac{p}{2} = 2 - \frac{p}{2}$	
4	YM, $n > 4$ or $q > 0$	0		

Table 1: List of cases in the proof of Lemma 13.

There is no sum over i . For every fixed i the only relations between the $(c_{ij})_{j \in I}$ are $c_{ii} = 0$, which suffices to conclude $D_{e_{ij}}P = 0$. So $u = P = 0$.

- Case 2: Here $u = \sum_{a,b \in I} P_{ab}c_{ab}$ for some polynomials $P_{ab} = P_{ab}(e)$. Using (1.1) we may assume $\sum_a P_{ab} = 0$ and $P_{aa} = 0$. Since $X_i u = 0$ for all i ,

$$\sum_j k_{ji} P_{ji} + \sum_{a,b,j} c_{ab} c_{ij} D_{e_{ij}} P_{ab} = 0$$

Since the terms scale differently in c and k , they are separately zero. Continue with $\sum_j k_{ji} P_{ji} = 0$. There is no sum over i . Since $n \geq 4$, for every fixed i the only relations between the $(k_{ji})_{j \in I}$ are $k_{ii} = 0$ and $\sum_j k_{ji} = 0$, which implies $P_{ji} = 0$. So $u = 0$.

- Case 1: Here $u = \sum_{a,b,c,d} P_{abcd}c_{ab}c_{cd} + \sum_{a,b} Q_{ab}k_{ab}$. Using (1.1), we may assume $P_{abcd} = P_{cdab}$, $\sum_a P_{abcd} = 0$, $P_{aacd} = 0$, $Q_{ab} = Q_{ba}$, $\sum_a Q_{ab} = 0$, $Q_{aa} = 0$. Using $X_i u = 0$ we have $V_i + W_i = 0$ using the abbreviations

$$V_i = \sum_{a,b,j} (2k_{ji}c_{ab}P_{abji} + k_{ab}c_{ij}D_{e_{ij}}Q_{ab})$$

$$W_i = \sum_{a,b,c,d,j} c_{ab}c_{cd}c_{ij}D_{e_{ij}}P_{abcd}$$

Since they scale differently in c and k , we have $V_i = W_i = 0$.

- It follows from $V_i = 0$ that (reasoning as in case 2) if all P are zero then all derivatives of all Q are zero and then $Q = 0$, since $\deg Q > 0$. The problem is thus reduced to showing that all P are zero.
- From $(1 - \delta_{ai})D_{c_{ak}}V_i - (1 - \delta_{ak})D_{c_{ai}}V_k = 0$ we get (since this particular combination eliminates the Q terms using $D_{e_{ik}} = D_{e_{ki}}$):

$$(1 - \delta_{ai}) \sum_j k_{ji} P_{akji} - (1 - \delta_{ak}) \sum_j k_{jk} P_{aijk} = 0 \quad (4.2)$$

for all $a, i, k \in I$. By differentiating with respect to $D_{k_{cd}}$, one obtains linear identities with constant coefficients for the P .

- From $D_{c_{ab}}D_{c_{cd}}D_{c_{ef}}W_i = 0$ we get

$$(1 - \delta_{ab})(\delta_{ai} - \frac{1}{n-1})D_{e_{ib}}P_{cdef} + (1 - \delta_{cd})(\delta_{ci} - \frac{1}{n-1})D_{e_{id}}P_{efab} + (1 - \delta_{ef})(\delta_{ei} - \frac{1}{n-1})D_{e_{if}}P_{abcd} = 0 \quad (4.3)$$

for all $a, b, c, d, e, f, i \in I$.

If $n \geq 5$ then (4.3) alone implies that all derivatives of all P are zero¹⁸ so $P = 0$. If $n = 4$ then combining (4.2), (4.3) also yields $P = 0$. So $u = 0$.

□

5 Some commutators

Here we compute a number of commutators, showing for instance that the B_i appear in the commutators of type $[X_i, C_j]$. These commutators are also used to prove Theorem 1. In fact, Corollary 15 contains identities for $A_j f$, $B_j f$, $C_j f$ with f the amplitude, and this will be used together with Lemma 16 by which $A_j f$, $B_j f$, $C_j f$ are pole-free, and together with Lemma 13, to prove Theorem 1 which says that $A_j f$, $B_j f$, $C_j f$ vanish.

Lemma 14. *Let $|I| \geq 3$. In $D_{C(I)}$ we have the following identities:*

$$\begin{aligned} [X_i, A_j] &= -(1 - \delta_{ij}) \frac{1}{|I|-1} D_{e_{ij}} X_j \\ [X_i, B_j] &= -2\delta_{ij} A_i - (1 - \delta_{ij}) \left(\frac{2}{|I|-2} D_{c_{ji}} X_j - D_{c_{ij}} X_i - D_{e_{ij}} Y_i + \frac{1}{|I|-1} D_{e_{ij}} (Y_j + Z) \right) \\ [X_i, C_j] &= -\delta_{ij} B_i + \frac{1}{|I|} B_i + (1 - \delta_{ij}) (D_{k_{ij}} X_i + D_{c_{ji}} Y_i - \frac{1}{|I|-2} D_{c_{ji}} Z) \end{aligned}$$

as well as

$$\begin{aligned} [Y_i, A_j] &= -2\delta_{ij} A_j & [Z, A_j] &= 0 \\ [Y_i, B_j] &= -\delta_{ij} B_j & [Z, B_j] &= -B_j \\ [Y_i, C_j] &= 0 & [Z, C_j] &= -2C_j \end{aligned}$$

for all $i, j \in I$. All these identities hold for both YM and GR; recall that the parameters defining Y_i and Z are different in these two cases.

Proof. By direct calculation using equations (1.2), (1.3) and Lemma 3. □

Corollary 15. *Suppose f is either the YM amplitude, $f = A_I$, or the GR amplitude, $f = M_I$, as in Definition 11. Then for all $i, j \in I$:*

$$X_i(A_j f) = 0 \quad (Y_i + 2\delta_{ij})(A_j f) = 0 \quad Z(A_j f) = 0 \quad (5.1a)$$

If $A_j f = 0$ for all j then

$$X_i(B_j f) = 0 \quad (Y_i + \delta_{ij})(B_j f) = 0 \quad (Z + 1)(B_j f) = 0 \quad (5.1b)$$

If $B_j f = 0$ for all j then

$$X_i(C_j f) = 0 \quad Y_i(C_j f) = 0 \quad (Z + 2)(C_j f) = 0 \quad (5.1c)$$

Proof. Use Lemma 14 and the fact that $X_i f = Y_i f = Z f = 0$ for all i . □

¹⁸Use (4.3) viewed as linear homogeneous identities with constant coefficients for the $D_{e_{ij}} P_{abcd}$, simplified using the algebraic conditions on P and $D_{e_{ii}} = 0$, $D_{e_{ij}} = D_{e_{ji}}$.

6 Proof of Theorem 1

The annihilator of the YM amplitude A_I is a left ideal in the Weyl algebra $D_{C(I)}$, and likewise the annihilator of the GR amplitude M_I . The elements X_i, Y_i, Z are well-known to be in the annihilator¹⁹, and so are some constant coefficient operators of order 2 for YM, order 3 for GR, that witness polynomiality in the c and e variables. Theorem 1 asserts that A_i, B_i, C_i are also in the annihilator. The proof, given at the end of this section, is by induction on $|I|$. The induction step uses the following lemma.

Lemma 16 (Key technical lemma). *Let U be as in Definition 9. Suppose $I = J \sqcup K$ with $|J|, |K| \geq 2$ as before. Suppose Theorem 1 holds for the index sets J_\bullet and K_\bullet . Cyclic orders are understood in the case of YM. Then if*

$$\mathbf{O} \in D_{C(I)} \subseteq D_M$$

is one of A_i, B_i, C_i with $i \in I$, then both

$$\begin{aligned} & \mathbf{O} U \frac{\alpha_J^*(A_{J_\bullet}) \alpha_K^*(A_{K_\bullet})}{\xi} \\ & \mathbf{O} U^2 \frac{\alpha_J^*(M_{J_\bullet}) \alpha_K^*(M_{K_\bullet})}{\xi} \end{aligned}$$

are elements of $\text{Frac}(R_M)$ without pole along $\xi = 0$. (More precisely, they are in the localization of R_M at the codimension one prime ideal generated by ξ .)

Proof. The computation will be in the Weyl algebra, we will not directly work with rational functions. The first step is to split off the Ξ direction, using the direct sum decomposition (3.9). At the Weyl algebra level,

$$D_M = D_{\mathbb{C}\Xi} \otimes D_{M^\perp}$$

Here $D_{\mathbb{C}\Xi}$ is the Weyl algebra generated by ξ and Ξ , with $[\Xi, \xi] = 2$. Every element of $D_{\mathbb{C}\Xi}$ commutes with every element of D_{M^\perp} . For every \mathbf{O} we have

$$\mathbf{O} = \begin{pmatrix} 1 & \xi \end{pmatrix} \begin{pmatrix} S_{00} & S_{01} & S_{02} \\ S_{10} & S_{11} & S_{12} \end{pmatrix} \begin{pmatrix} 1 \\ \Xi \\ \Xi^2 \end{pmatrix}$$

for unique $S_{00}, \dots, S_{12} \in D_{M^\perp}$. This isolates all occurrences of ξ and Ξ . It is clear from (1.3) that at most one ξ and at most two Ξ appear. Set

$$\begin{aligned} S_0 &= S_{10} \\ S_1 &= S_{00} - 2S_{11} \\ S_2 &= S_{01} - 4S_{12} \\ S_3 &= S_{02} \end{aligned}$$

¹⁹In the logic of this paper, this is enshrined in Definition 11.

Note that Ξ commutes with $U \in D_{M^\perp}$ from Definition 9, and the pushforward of Ξ under both α_J and α_K is zero. Hence it suffices to show that

$$S_1 U \alpha_J^*(A_{J_\bullet}) \alpha_K^*(A_{K_\bullet}) = 0 \quad (6.1a)$$

$$S_2 U \alpha_J^*(A_{J_\bullet}) \alpha_K^*(A_{K_\bullet}) = 0 \quad (6.1b)$$

$$S_3 U \alpha_J^*(A_{J_\bullet}) \alpha_K^*(A_{K_\bullet}) = 0 \quad (6.1c)$$

for YM, corresponding (by definition of S_0, \dots, S_3) to the absence of $1/\xi$ and $1/\xi^2$ and $1/\xi^3$ terms respectively, and analogously it suffices to show that

$$S_1 U^2 \alpha_J^*(M_{J_\bullet}) \alpha_K^*(M_{K_\bullet}) = 0 \quad (6.2a)$$

$$S_2 U^2 \alpha_J^*(M_{J_\bullet}) \alpha_K^*(M_{K_\bullet}) = 0 \quad (6.2b)$$

$$S_3 U^2 \alpha_J^*(M_{J_\bullet}) \alpha_K^*(M_{K_\bullet}) = 0 \quad (6.2c)$$

for GR. Note that S_0 has dropped out of the computation since it cannot generate a pole along $\xi = 0$. Note that:

- $S_2 = S_3 = 0$ for $\mathbf{O} = A_i$ because it involves no k -derivative, hence no Ξ -derivatives. This also implies $S_{11} = 0$ in this case.
- $S_3 = 0$ for $\mathbf{O} = B_i$ because it involves no second k -derivatives, hence no second Ξ derivatives. This also implies $S_{12} = 0$ in this case.
- $S_3 = 0$ for $\mathbf{O} = C_i$ but this requires a calculation. It suffices to show that the analogous claim holds for \tilde{C}_i , that is for the following terms in (1.3c):

$$a \sum_{j,k \in I} k_{jk} D_{k_{ij}} D_{k_{ik}} + b \sum_{j,k \in I} k_{jk} D_{k_{ji}} D_{k_{jk}}$$

with $a = \frac{1}{2}$ and $b = -1$. Use Definition 4 to replace $k_{ab} = k_{ab}^\perp + \text{const}_{ab} \cdot \xi$ and $D_{k_{ab}} = D_{k_{ab}}^\perp + \text{const}_{ab} \cdot \Xi$ and keep only the k_{ab}^\perp respectively Ξ terms. Using the first equation in (3.6) and $2a + b = 0$, we get zero.

It now suffices to show

$$(6.1a), (6.2a) \text{ for } A_j, B_j, C_j \text{ for all } j \in J. \quad (6.3a)$$

$$(6.1b), (6.2b) \text{ for } B_j, C_j \text{ for all } j \in J. \quad (6.3b)$$

The restriction to $i = j \in J$ is new and is without loss of generality. It entails that the rest of this proof is not symmetric under exchanging J and K . The rest of this proof exploits the direct sum decomposition of M^\perp in (C.1), see also Corollary 19. At Weyl algebra level,

$$D_{M^\perp} = D_{\text{image } \pi} \otimes \underbrace{D_{\text{image } \beta_J}}_{\simeq D_{C(J_\bullet)}} \otimes \underbrace{D_{\text{image } \beta_K}}_{\simeq D_{C(K_\bullet)}}$$

The indicated isomorphisms are established by α_J, β_J and α_K, β_K . Call them

$$\gamma_J : D_{C(J_\bullet)} \rightarrow D_{\text{image } \beta_J}$$

$$\gamma_K : D_{C(K_\bullet)} \rightarrow D_{\text{image } \beta_K}$$

They are explicitly given by the formulas in Lemmas 8, 17, 18. Let $I_{\text{YM}} \subseteq D_{M^\perp}$ respectively $I_{\text{GR}} \subseteq D_{M^\perp}$ be the left ideals generated by (the difference between YM and GR is implicit in the parameters defining the Y and Z elements):

- The left ideal $I_\pi \subseteq D_{\text{image } \pi}$ generated by all partial derivatives. Equivalently, this is the annihilator of the constant functions on image π .
- The left ideal in $D_{\text{image } \beta_J}$ generated by the image under γ_J of the following elements, which are known annihilators of A_{J_\bullet} respectively M_{J_\bullet} :

$$A_j, A_\bullet, B_j, B_\bullet, C_j, C_\bullet, X_j, X_\bullet, Y_j, Y_\bullet, Z \in D_{C(J_\bullet)} \quad (6.4a)$$

and²⁰

$$\text{for YM: } D_{x_{j_1}\bullet} D_{y_{j_2}\bullet} \quad \text{for GR: } D_{x_{j_1}\bullet} D_{y_{j_2}\bullet} D_{z_{j_3}\bullet} \quad (6.4b)$$

for all $x, y, z \in \{c, e\}$ and all $j_1, j_2, j_3 \in J$.

- The left ideal in $D_{\text{image } \beta_K}$ generated by the image under γ_K of the following elements, which are known annihilators of A_{K_\bullet} respectively M_{K_\bullet} :

$$A_\bullet, B_\bullet, C_\bullet, X_\bullet, Y_\bullet, Z \in D_{C(K_\bullet)} \quad (6.5a)$$

and

$$\text{for YM: } D_{x_{k_1}\bullet} D_{y_{k_2}\bullet} \quad \text{for GR: } D_{x_{k_1}\bullet} D_{y_{k_2}\bullet} D_{z_{k_3}\bullet} \quad (6.5b)$$

for all $x, y, z \in \{c, e\}$ and all $k_1, k_2, k_3 \in K$.

It is part of the assumptions of this lemma, and a consequence of Definition 11, that (6.4) and (6.5) are known annihilators of A_{J_\bullet} , M_{J_\bullet} and A_{K_\bullet} , M_{K_\bullet} . More annihilators are known, but we will not need those. To show (6.3) it now suffices to show that (by construction of I_{YM} , I_{GR} and using $\alpha_J \pi = 0$, $\alpha_K \pi = 0$):

$$S_1 U \in I_{\text{YM}} \quad \text{for } \mathbf{O} = A_j, B_j, C_j \quad (6.6a)$$

$$S_2 U \in I_{\text{YM}} \quad \text{for } \mathbf{O} = B_j, C_j \quad (6.6b)$$

$$S_1 U^2 \in I_{\text{GR}} \quad \text{for } \mathbf{O} = A_j, B_j, C_j \quad (6.6c)$$

$$S_2 U^2 \in I_{\text{GR}} \quad \text{for } \mathbf{O} = B_j, C_j \quad (6.6d)$$

Thus the problem is reduced to one of checking membership in a left ideal in the Weyl algebra D_{M^\perp} . (For every fixed $|I|$ this can in principle be checked algorithmically using Gröbner bases. But we need to prove membership for all $|I|$.) To proceed, we use the canonical $D_{\text{image } \pi} / I_\pi \simeq R_{\text{image } \pi}$. Here $R_{\text{image } \pi}$ are the polynomials on image π . This gives a canonical map

$$\rho : D_{M^\perp} \rightarrow \underbrace{R_{\text{image } \pi}}_{\simeq D_{\text{image } \pi} / I_\pi} \otimes D_{\text{image } \beta_J} \otimes D_{\text{image } \beta_K}$$

²⁰These are also known annihilators. They witness polynomiality in the c and e variables related to ‘the polarization of particle \bullet ’ with degrees fixed by $Y_\bullet \in D_{C(J_\bullet)}$.

Decompose $R_{\text{image } \pi} = \bigoplus_{d \geq 0} (R_{\text{image } \pi})_d$ where $(R_{\text{image } \pi})_d$ are all polynomials that are homogeneous of degree d . Accordingly $\rho = \bigoplus_{d \geq 0} \rho_d$ where

$$\rho_d : D_{M^\perp} \rightarrow (R_{\text{image } \pi})_d \otimes D_{\text{image } \beta_J} \otimes D_{\text{image } \beta_K}$$

Clearly $\rho_d(S_1), \rho_d(S_2)$ can only be nonzero for $d = 0, 1$ because all \mathbf{O} in (1.3) have polynomial coefficients of order ≤ 1 . We claim that actually

$$\rho_1(S_1) = \rho_1(S_2) = 0 \tag{6.7}$$

for all $\mathbf{O} = A_j, B_j, C_j$ (an analogous statement fails for \tilde{C}_j). To see this, use the description of image π in Corollary 19. Here are some examples:

- Consider the term $\sum_{a,b \in I} e_{ab} D_{e_{aj}} D_{e_{bj}}$ in A_j . This term is already in D_M^\perp , since $e = e^\perp$ and $D_e = D_e^\perp$. If $a, b \in J$ or $a, b \in K$ then we get no contribution to ρ_1 since e_{ab}^\perp commutes with all elements of image π by Corollary 19. If $a \in K$ then $D_{e_{aj}}^\perp \in I_\pi$ and if $b \in K$ then $D_{e_{bj}}^\perp \in I_\pi$ and we also get no contribution to ρ_1 .
- Consider next $\sum_{a,b \in I} c_{ab} D_{c_{aj}} D_{e_{bj}}$ in A_j . By the same reasoning, it suffices to consider the sum over $a \in K, b \in J$. For $a \in K$, write $D_{c_{aj}} = D_{c_{aj}}^\perp$ as

$$D_{c_{aj}}^\perp = (D_{c_{aj}}^\perp - r) + r \quad \text{where} \quad r = \frac{1}{|K|} \sum_{k' \in K} D_{c_{k'j}}^\perp$$

The first term does not contribute to ρ_1 since $D_{c_{aj}}^\perp - r \in I_\pi$ by Corollary 19. The second does not contribute to ρ_1 since $\sum_{a \in K} c_{ab}^\perp = -\sum_{a \in J} c_{ab}^\perp$ which for $b \in J$ commutes with all elements of image π by Corollary 19.

- Terms involving k or D_k are more complicated. One must take into account $k = k^\perp + \text{const} \cdot \xi$ and $D_k = D_k^\perp + \text{const} \cdot \Xi$, in Definition 4.

Using (6.7) one can see that $\rho_d(S_1 U), \rho_d(S_2 U)$ can only be nonzero for $d = 0, 1$ and that $\rho_d(S_1 U^2), \rho_d(S_2 U^2)$ can only be nonzero for $d = 0, 1, 2$. In each of these cases, and for $\mathbf{O} = A_j, B_j, C_j$, Tables 2 and 3 list elements in (6.4) and (6.5) that suffice to prove membership in I_{YM} respectively I_{GR} , as in (6.6). The tag ‘poly’ subsumes all annihilators in (6.4b), (6.5b) that witness polynomiality.

We now discuss these tables in detail. The identities below are in

$$(R_{\text{image } \pi})_0 \otimes D_{\text{image } \beta_J} \otimes D_{\text{image } \beta_K} \simeq D_{\text{image } \beta_J} \otimes D_{\text{image } \beta_K} \tag{6.8}$$

To extract the various ‘Taylor coefficients’ we use the \tilde{D} defined in Corollary 19, understood here as mapping $(R_{\text{image } \pi})_d \rightarrow (R_{\text{image } \pi})_{d-1}$. We will not make explicit the ‘poly’ pieces and state some identities in the schematic form

$$a = b \pmod{\text{poly}}$$

which asserts that $a - b$ is in the left ideal of (6.8) generated, via γ_J and γ_K respectively, by (6.4b) and (6.5b). With these preliminaries, one row in Table 3 with $\mathbf{O} = C_j$ is proved

	d	elements from (6.4)	elements from (6.5)
$\rho_d(S_1 U)$ for $\mathbf{O} = A_j$	0		X_\bullet
	1	A_j, X_j, poly	
$\rho_d(S_2 U)$ for $\mathbf{O} = B_j$	0		X_\bullet
	1	X_j, poly	
$\rho_d(S_1 U)$ for $\mathbf{O} = B_j$	0		$B_\bullet, X_\bullet, \text{poly}$
	1	$B_j, X_j, Y_j, Z, \text{poly}$	Z, poly
$\rho_d(S_2 U)$ for $\mathbf{O} = C_j$	0		X_\bullet
	1	Z, poly	Z, poly
$\rho_d(S_1 U)$ for $\mathbf{O} = C_j$	0		$B_\bullet, X_\bullet, \text{poly}$
	1	$C_j, C_\bullet, Z, \text{poly}$	$C_\bullet, Z, \text{poly}$

Table 2: For YM, this table lists elements in (6.4) and (6.5) that suffice to prove membership in I_{YM} as in (6.6a), (6.6b). The list is not necessarily minimal.

	d	elements from (6.4)	elements from (6.5)
$\rho_d(S_1 U^2)$ for $\mathbf{O} = A_j$	0		A_\bullet
	1		X_\bullet
	2	A_j, X_j, poly	
$\rho_d(S_2 U^2)$ for $\mathbf{O} = B_j$	0		
	1		X_\bullet
	2	X_j, poly	
$\rho_d(S_1 U^2)$ for $\mathbf{O} = B_j$	0	X_\bullet	A_\bullet
	1		$B_\bullet, X_\bullet, \text{poly}$
	2	$B_j, X_j, Y_j, Z, \text{poly}$	Z, poly
$\rho_d(S_2 U^2)$ for $\mathbf{O} = C_j$	0		
	1		X_\bullet
	2	Z, poly	Z, poly
$\rho_d(S_1 U^2)$ for $\mathbf{O} = C_j$	0	A_\bullet, X_\bullet	A_\bullet
	1		$B_\bullet, X_\bullet, \text{poly}$
	2	$C_j, C_\bullet, Z, \text{poly}$	$C_\bullet, Z, \text{poly}$

Table 3: For GR, this table lists elements in (6.4) and (6.5) that suffice to prove membership in I_{GR} as in (6.6c), (6.6d). The list is not necessarily minimal.

by the identities

$$\begin{aligned}
\tilde{D}_{k_{j_1} k_1} \rho_1(S_2 U^2) &= 2D_{c_{j_1} \bullet} D_{c_{k_1} \bullet} D_{c_j \bullet} \gamma_K(X_\bullet) \\
\tilde{D}_{c_{j_1} k_1} \rho_1(S_2 U^2) &= 2D_{c_{j_1} \bullet} D_{e_{k_1} \bullet} D_{c_j \bullet} \gamma_K(X_\bullet) \\
\tilde{D}_{c_{k_1} j_1} \rho_1(S_2 U^2) &= 2D_{e_{j_1} \bullet} D_{c_{k_1} \bullet} D_{c_j \bullet} \gamma_K(X_\bullet) \\
\tilde{D}_{e_{j_1} k_1} \rho_1(S_2 U^2) &= 2D_{e_{j_1} \bullet} D_{e_{k_1} \bullet} D_{c_j \bullet} \gamma_K(X_\bullet)
\end{aligned} \tag{6.9}$$

in the space (6.8), they hold for all $j_1 \in J$ and $k_1 \in K$. It is essential here that derivatives such as $D_{e_{k_1} \bullet}$ are to the left of $\gamma_K(X_\bullet)$. On the other hand, since $\gamma_K(D_{e_{k_1} \bullet}) = D_{e_{k_1} \bullet}$ by Lemmas 17 and 18, it does not matter if this derivative is written inside or outside of γ_K .

We abbreviate

$$\tilde{D}_1 = \begin{pmatrix} \tilde{D}_{k_{j_1}k_1} \\ \tilde{D}_{c_{j_1}k_1} \\ \tilde{D}_{c_{k_1}j_1} \\ \tilde{D}_{e_{j_1}k_1} \end{pmatrix} \quad R_1 = \begin{pmatrix} D_{c_{j_1}\bullet} D_{c_{k_1}\bullet} \\ D_{c_{j_1}\bullet} D_{e_{k_1}\bullet} \\ D_{e_{j_1}\bullet} D_{c_{k_1}\bullet} \\ D_{e_{j_1}\bullet} D_{e_{k_1}\bullet} \end{pmatrix}$$

and

$$\tilde{D}_2 = \begin{pmatrix} \tilde{D}_{k_{j_1}k_1} & \tilde{D}_{k_{j_2}k_2} \\ \tilde{D}_{c_{j_1}k_1} & \tilde{D}_{c_{j_2}k_2} \\ \tilde{D}_{c_{k_1}j_1} & \tilde{D}_{c_{k_2}j_2} \\ \tilde{D}_{e_{j_1}k_1} & \tilde{D}_{e_{j_2}k_2} \\ \tilde{D}_{k_{j_1}k_1} & \tilde{D}_{c_{j_2}k_2} \\ \tilde{D}_{c_{j_1}k_1} & \tilde{D}_{c_{k_2}j_2} \\ \tilde{D}_{c_{k_1}j_1} & \tilde{D}_{e_{j_2}k_2} \\ \tilde{D}_{k_{j_1}k_1} & \tilde{D}_{c_{k_2}j_2} \\ \tilde{D}_{c_{j_1}k_1} & \tilde{D}_{e_{j_2}k_2} \\ \tilde{D}_{k_{j_1}k_1} & \tilde{D}_{e_{j_2}k_2} \end{pmatrix} \quad R_2 = \begin{pmatrix} D_{c_{j_1}\bullet} D_{c_{j_2}\bullet} D_{c_{k_1}\bullet} D_{c_{k_2}\bullet} \\ D_{c_{j_1}\bullet} D_{c_{j_2}\bullet} D_{e_{k_1}\bullet} D_{e_{k_2}\bullet} \\ D_{e_{j_1}\bullet} D_{e_{j_2}\bullet} D_{c_{k_1}\bullet} D_{c_{k_2}\bullet} \\ D_{e_{j_1}\bullet} D_{e_{j_2}\bullet} D_{e_{k_1}\bullet} D_{e_{k_2}\bullet} \\ D_{c_{j_1}\bullet} D_{c_{j_2}\bullet} D_{c_{k_1}\bullet} D_{e_{k_2}\bullet} \\ D_{c_{j_1}\bullet} D_{e_{j_2}\bullet} D_{e_{k_1}\bullet} D_{c_{k_2}\bullet} \\ D_{e_{j_1}\bullet} D_{e_{j_2}\bullet} D_{c_{k_1}\bullet} D_{e_{k_2}\bullet} \\ D_{c_{j_1}\bullet} D_{e_{j_2}\bullet} D_{c_{k_1}\bullet} D_{c_{k_2}\bullet} \\ D_{c_{j_1}\bullet} D_{e_{j_2}\bullet} D_{e_{k_1}\bullet} D_{e_{k_2}\bullet} \\ D_{c_{j_1}\bullet} D_{e_{j_2}\bullet} D_{c_{k_1}\bullet} D_{e_{k_2}\bullet} \end{pmatrix}$$

with the understanding that $j_1, j_2 \in J$ and $k_1, k_2 \in K$. The four identities (6.9) are now given, more succinctly, by

$$\tilde{D}_1 \rho_1(S_2 U^2) = 2 D_{c_{j\bullet}} R_1 \gamma_K(X_\bullet)$$

We now state all identities needed for (6.6). For YM, with reference to Table 2:

- If $\mathbf{O} = A_j$:

$$\begin{aligned} \rho_0(S_1 U) &= -\frac{|I|}{|I|-1} \gamma_J(D_{c_{\bullet j}}) D_{e_{j\bullet}} \gamma_K(X_\bullet) \\ \tilde{D}_1 \rho_1(S_1 U) &= R_1 \gamma_J(A_j) + \frac{|K|-1}{|I|-1} \gamma_J(D_{c_{\bullet j}}) R_1 \gamma_J(X_j) \quad \text{mod poly} \end{aligned}$$

- If $\mathbf{O} = B_j$:

$$\begin{aligned} \rho_0(S_2 U) &= \frac{(|J|-1)(|I|^2-|I|-|J|)}{|J|(|I|-1)(|I|-2)} D_{e_{j\bullet}} \gamma_K(X_\bullet) \\ \tilde{D}_1 \rho_1(S_2 U) &= -\frac{|K|(|K|-1)}{(|I|-1)(|I|-2)} R_1 \gamma_J(X_j) \quad \text{mod poly} \end{aligned}$$

and

$$\begin{aligned} \rho_0(S_1 U) &= D_{e_{j\bullet}} \gamma_K(B_\bullet) - \frac{1}{|J|(|I|-2)} \gamma_J(\\ &\quad (|I|-2)(|J|+1) D_{c_{j\bullet}} D_{c_{\bullet j}} \\ &\quad + |I|(|J|-1) D_{e_{j\bullet}} D_{k_{\bullet j}} \\ &\quad + (|I|-2) \sum_{j' \in J} D_{c_{j'\bullet}} D_{c_{j'j}}) \gamma_K(X_\bullet) \quad \text{mod poly} \end{aligned}$$

$$\begin{aligned}
\tilde{D}_1 \rho_1(S_1 U) &= R_1 \gamma_J(B_j) + \frac{2(|K|-1)}{|I|-2} \gamma_J(D_{k_{\bullet j}}) R_1 \gamma_J(X_j) \\
&\quad + \frac{|K|-1}{|I|-1} \gamma_J(D_{c_{\bullet j}}) R_1 \gamma_J(Y_j) \\
&\quad + \frac{|K|-1}{|I|-1} \gamma_J(D_{c_{\bullet j}}) R_1 \gamma_J(Z) \\
&\quad - \frac{|J|}{|I|-1} \gamma_J(D_{c_{\bullet j}}) R_1 \gamma_K(Z) \mod \text{poly}
\end{aligned}$$

- If $\mathbf{O} = C_j$:

$$\begin{aligned}
\rho_0(S_2 U) &= D_{c_{\bullet}} \gamma_K(X_{\bullet}) \\
\tilde{D}_1 \rho_1(S_2 U) &= -\frac{|K|(|K|-1)}{|I|(|I|-2)} R_1 \gamma_J(Z) + \frac{|K|(|J|-1)}{|I|(|I|-2)} R_1 \gamma_K(Z) \mod \text{poly}
\end{aligned}$$

and

$$\begin{aligned}
\rho_0(S_1 U) &= D_{c_{\bullet}} \gamma_K(B_{\bullet}) - \frac{1}{|J|} \gamma_J \left(\right. \\
&\quad \left. (|J|+1) D_{c_{\bullet}} D_{k_{\bullet j}} \right. \\
&\quad \left. + \sum_{j' \in J} D_{c_{j' \bullet}} D_{k_{jj'}} \right. \\
&\quad \left. - \sum_{j' \in J} D_{c_{j' \bullet}} D_{k_{\bullet j'}} \right) \gamma_K(X_{\bullet}) \mod \text{poly} \\
\tilde{D}_1 \rho_1(S_1 U) &= -\frac{|K|-1}{|I|} R_1 \gamma_J(C_{\bullet}) + R_1 \gamma_J(C_j) \\
&\quad + \frac{|K|-1}{|I|-2} \gamma_J(D_{k_{\bullet j}}) R_1 \gamma_J(Z) + \frac{|K|+1}{|I|} R_1 \gamma_K(C_{\bullet}) \\
&\quad - \frac{|J|-1}{|I|-2} \gamma_J(D_{k_{\bullet j}}) R_1 \gamma_K(Z) \mod \text{poly}
\end{aligned}$$

For GR, with reference to Table 3:

- If $\mathbf{O} = A_j$:

$$\begin{aligned}
\rho_0(S_1 U^2) &= 2 D_{e_{j \bullet}}^2 \gamma_K(A_{\bullet}) \\
\tilde{D}_1 \rho_1(S_1 U^2) &= -2 \frac{|I|}{|I|-1} \gamma_J(D_{c_{\bullet j}}) D_{e_{j \bullet}} R_1 \gamma_K(X_{\bullet}) \\
\tilde{D}_2 \rho_2(S_1 U^2) &= 2 R_2 \gamma_J(A_j) + 2 \frac{|K|-1}{|I|-1} \gamma_J(D_{c_{\bullet j}}) R_2 \gamma_J(X_j) \mod \text{poly}
\end{aligned}$$

- If $\mathbf{O} = B_j$:

$$\begin{aligned}
\rho_0(S_2 U^2) &= 0 \\
\tilde{D}_1 \rho_1(S_2 U^2) &= 2 \frac{(|J|-1)(|I|^2-|I|-|J|)}{|J|(|I|-1)(|I|-2)} D_{e_{j \bullet}} R_1 \gamma_K(X_{\bullet}) \\
\tilde{D}_2 \rho_2(S_2 U^2) &= -2 \frac{|K|(|K|-1)}{(|I|-1)(|I|-2)} R_2 \gamma_J(X_j) \mod \text{poly}
\end{aligned}$$

and

$$\begin{aligned}
\rho_0(S_1 U^2) &= -\frac{2}{|K|} D_{e_{j \bullet}} \sum_{k' \in K} (D_{c_{k' \bullet}})^2 \gamma_J(X_{\bullet}) + 4 D_{c_{j \bullet}} D_{e_{j \bullet}} \gamma_K(A_{\bullet}) \\
\tilde{D}_1 \rho_1(S_1 U^2) &= 2 D_{e_{j \bullet}} R_1 \gamma_K(B_{\bullet}) - \frac{2}{|J|(|I|-2)} \gamma_J \left(\right. \\
&\quad \left. (|I|-2)(|J|+1) D_{c_{j \bullet}} D_{c_{\bullet j}} \right. \\
&\quad \left. + |I|(|J|-1) D_{e_{j \bullet}} D_{k_{\bullet j}} \right. \\
&\quad \left. + (|I|-2) \sum_{j' \in J} D_{c_{j' \bullet}} D_{c_{j' j}} \right) R_1 \gamma_K(X_{\bullet}) \mod \text{poly}
\end{aligned}$$

$$\begin{aligned}
\tilde{D}_2 \rho_2(S_1 U^2) &= 2R_2 \gamma_J(B_j) + \frac{4(|K|-1)}{|I|-2} \gamma_J(D_{k_{\bullet j}}) R_2 \gamma_J(X_j) \\
&\quad + \frac{2(|K|-1)}{|I|-1} \gamma_J(D_{c_{\bullet j}}) R_2 \gamma_J(Y_j) \\
&\quad + \frac{2(|K|-1)}{|I|-1} \gamma_J(D_{c_{\bullet j}}) R_2 \gamma_J(Z) \\
&\quad - \frac{2|J|}{|I|-1} \gamma_J(D_{c_{\bullet j}}) R_2 \gamma_K(Z) \pmod{\text{poly}}
\end{aligned}$$

- If $\mathbf{O} = C_j$:

$$\begin{aligned}
\rho_0(S_2 U^2) &= 0 \\
\tilde{D}_1 \rho_1(S_2 U^2) &= 2D_{c_j \bullet} R_1 \gamma_K(X_\bullet) \\
\tilde{D}_2 \rho_2(S_2 U^2) &= -\frac{2|K|(|K|-1)}{|I|(|I|-2)} R_2 \gamma_J(Z) + \frac{2|K|(|J|-1)}{|I|(|I|-2)} R_2 \gamma_K(Z) \pmod{\text{poly}}
\end{aligned}$$

and

$$\begin{aligned}
\rho_0(S_1 U^2) &= -\frac{2}{|I|} \sum_{k' \in K} D_{c_{k' \bullet}}^2 \gamma_J(A_\bullet) - \frac{2}{|K|} D_{c_j \bullet} \sum_{k' \in K} D_{c_{k' \bullet}}^2 \gamma_J(X_\bullet) \\
&\quad + 2(D_{c_j \bullet}^2 - \frac{1}{|I|} \sum_{j' \in J} D_{c_{j' \bullet}}^2) \gamma_K(A_\bullet) \\
\tilde{D}_1 \rho_1(S_1 U^2) &= 2D_{j \bullet} R_1 \gamma_K(B_\bullet) - \frac{2}{|J|} \gamma_J(\\
&\quad (|J| + 1) D_{c_j \bullet} D_{k_{\bullet j}} \\
&\quad + \sum_{j' \in J} D_{c_{j' \bullet}} D_{k_{j j'}} \\
&\quad - \sum_{j' \in J} D_{c_{j' \bullet}} D_{k_{\bullet j'}}) R_1 \gamma_K(X_\bullet) \pmod{\text{poly}} \\
\tilde{D}_2 \rho_2(S_1 U^2) &= -\frac{2(|K|-1)}{|I|} R_2 \gamma_J(C_\bullet) + 2R_2 \gamma_J(C_j) \\
&\quad + \frac{2(|K|-1)}{|I|-2} \gamma_J(D_{k_{\bullet j}}) R_2 \gamma_J(Z) + \frac{2(|K|+1)}{|I|} R_2 \gamma_K(C_\bullet) \\
&\quad - \frac{2(|J|-1)}{|I|-2} \gamma_J(D_{k_{\bullet j}}) R_2 \gamma_K(Z) \pmod{\text{poly}}
\end{aligned}$$

These Weyl algebra identities are by direct calculation; this is algorithmically straightforward, best done using symbolic computation. They imply (6.6) hence Lemma 16. \square

Proof (of Theorem 1). The proof is by induction on $|I|$. The base case $|I| = 3$ is by direct calculation. As an example, if $I = \{1, 2, 3\}$ then

$$\tilde{C}_1 M_I = \tilde{C}_2 M_I = \tilde{C}_3 M_I = \text{const} \cdot e_{12} e_{23} e_{31}$$

It follows that $C_i M_I = 0$ for $i = 1, 2, 3$ as required. Let now $|I| \geq 4$. Let $f = A_I$ (YM) or $f = M_I$ (GR). For every $I = J \sqcup K$ with $|J|, |K| \geq 2$ write

$$f = (f - g) + g \quad g = \begin{cases} \text{const} \cdot U \frac{\alpha_J^*(A_{J_\bullet}) \alpha_K^*(A_{K_\bullet})}{\xi} & \text{for YM} \\ \text{const} \cdot U^2 \frac{\alpha_J^*(M_{J_\bullet}) \alpha_K^*(M_{K_\bullet})}{\xi} & \text{for GR} \end{cases}$$

Note that:

- By the recursion in Definition 11, the difference $f - g$ does not have a pole along $\xi = 0$ and therefore neither do $A_i(f - g)$, $B_i(f - g)$, $C_i(f - g)$.

- By the induction hypothesis, we can invoke Lemma 16 and conclude that also $A_i g$, $B_i g$, $C_i g$ do not have a pole along $\xi = 0$.

Hence $A_i f$, $B_i f$, $C_i f$ have no pole along $\xi = 0$ for every decomposition $I = J \sqcup K$, and therefore (by Hartogs extension as in the proof of Lemma 12) we have

$$A_i f, B_i f, C_i f \in R_{C(I)}$$

where $R_{C(I)}$ is the ring of polynomials. To show that they are actually zero:

- Use (5.1a) and Lemma 13 (with $u = A_i f$) to conclude that $A_i f = 0$.
- Then use (5.1b) and Lemma 13 (with $u = B_i f$) to conclude that $B_i f = 0$.
- Then use (5.1c) and Lemma 13 (with $u = C_i f$) to conclude that $C_i f = 0$.

□

7 Acknowledgments

M.R. is grateful to have received funding from ERC through grant agreement No. 669655.

A Some tree amplitudes

The expressions below are rational functions on the vector space $C(I)$ in Definition 2, defined using the relations (1.1). Not all symmetries are immediate from the formulas, for example the permutation invariance of the GR amplitudes is not. The expressions are up to an overall multiplicative constant.

- The YM amplitude A_I for $I = \{1, 2, 3\}$: $-c_{23}e_{12} + c_{21}e_{23} - c_{12}e_{31}$.
- The GR amplitude M_I for $I = \{1, 2, 3\}$: $(-c_{23}e_{12} + c_{21}e_{23} - c_{12}e_{31})^2$.
- The YM amplitude A_I for $I = \{1, 2, 3, 4\}$, with canonical cyclic order:

$$\begin{aligned} & -\frac{c_{14}c_{23}e_{12}}{k_{12}} + \frac{c_{13}c_{24}e_{12}}{k_{12}} - \frac{c_{14}c_{23}e_{12}}{k_{23}} - \frac{c_{12}c_{14}e_{13}}{k_{12}} - \frac{c_{12}c_{24}e_{13}}{k_{12}} + \frac{c_{12}c_{13}e_{14}}{k_{12}} + \frac{c_{12}c_{23}e_{14}}{k_{12}} + \frac{c_{14}c_{21}e_{23}}{k_{12}} \\ & + \frac{c_{21}c_{24}e_{23}}{k_{12}} - \frac{c_{13}c_{21}e_{24}}{k_{12}} - \frac{c_{21}c_{23}e_{24}}{k_{12}} - \frac{c_{12}c_{31}e_{34}}{k_{12}} + \frac{c_{21}c_{32}e_{34}}{k_{12}} + \frac{c_{14}c_{32}e_{13}}{k_{23}} + \frac{c_{12}c_{23}e_{14}}{k_{23}} - \frac{c_{13}c_{32}e_{14}}{k_{23}} \\ & + \frac{c_{14}c_{21}e_{23}}{k_{23}} + \frac{c_{21}c_{24}e_{23}}{k_{23}} + \frac{c_{24}c_{31}e_{23}}{k_{23}} - \frac{c_{21}c_{23}e_{24}}{k_{23}} - \frac{c_{23}c_{31}e_{24}}{k_{23}} + \frac{c_{21}c_{32}e_{34}}{k_{23}} + \frac{c_{31}c_{32}e_{34}}{k_{23}} - \frac{e_{34}e_{12}k_{23}}{k_{12}} \\ & - \frac{e_{14}e_{23}k_{12}}{k_{23}} - e_{34}e_{12} - e_{14}e_{23} + e_{13}e_{24} \end{aligned}$$

- The GR amplitude M_I for $I = \{1, 2, 3, 4\}$, as an unordered set:

$$\begin{aligned} & k_{32} \left(-e_{34}e_{12} - \frac{e_{34}k_{23}e_{12}}{k_{12}} - \frac{c_{14}c_{23}e_{12}}{k_{12}} + \frac{c_{13}c_{24}e_{12}}{k_{12}} - \frac{c_{14}c_{23}e_{12}}{k_{23}} - e_{14}e_{23} + e_{13}e_{24} \right. \\ & - \frac{c_{12}c_{14}e_{13}}{k_{12}} - \frac{c_{12}c_{24}e_{13}}{k_{12}} + \frac{c_{12}c_{13}e_{14}}{k_{12}} + \frac{c_{12}c_{23}e_{14}}{k_{12}} + \frac{c_{14}c_{21}e_{23}}{k_{12}} + \frac{c_{21}c_{24}e_{23}}{k_{12}} - \frac{c_{13}c_{21}e_{24}}{k_{12}} - \frac{c_{21}c_{23}e_{24}}{k_{12}} \\ & \left. - \frac{c_{12}c_{31}e_{34}}{k_{12}} + \frac{c_{21}c_{32}e_{34}}{k_{12}} + \frac{c_{14}c_{32}e_{13}}{k_{23}} + \frac{c_{12}c_{23}e_{14}}{k_{23}} - \frac{c_{13}c_{32}e_{14}}{k_{23}} + \frac{c_{14}c_{21}e_{23}}{k_{23}} + \frac{c_{21}c_{24}e_{23}}{k_{23}} + \frac{c_{24}c_{31}e_{23}}{k_{23}} \right) \end{aligned}$$

$$\begin{aligned}
& -\frac{c_{21}c_{23}e_{24}}{k_{23}} - \frac{c_{23}c_{31}e_{24}}{k_{23}} + \frac{c_{21}c_{32}e_{34}}{k_{23}} + \frac{c_{31}c_{32}e_{34}}{k_{23}} - \frac{e_{14}e_{23}k_{12}}{k_{23}} \Big) \\
& \cdot \left(e_{34}e_{12} - \frac{c_{13}c_{14}e_{12}}{k_{13}} - \frac{c_{13}c_{34}e_{12}}{k_{13}} + \frac{c_{14}c_{23}e_{12}}{k_{32}} - e_{13}e_{24} - e_{14}e_{32} \right. \\
& - \frac{e_{13}e_{24}k_{32}}{k_{13}} - \frac{c_{14}c_{32}e_{13}}{k_{13}} + \frac{c_{12}c_{34}e_{13}}{k_{13}} + \frac{c_{12}c_{13}e_{14}}{k_{13}} + \frac{c_{13}c_{32}e_{14}}{k_{13}} - \frac{c_{13}c_{21}e_{24}}{k_{13}} + \frac{c_{23}c_{31}e_{24}}{k_{13}} + \frac{c_{14}c_{31}e_{32}}{k_{13}} \\
& + \frac{c_{31}c_{34}e_{32}}{k_{13}} - \frac{c_{12}c_{31}e_{34}}{k_{13}} - \frac{c_{31}c_{32}e_{34}}{k_{13}} - \frac{c_{14}c_{32}e_{13}}{k_{32}} - \frac{c_{12}c_{23}e_{14}}{k_{32}} + \frac{c_{13}c_{32}e_{14}}{k_{32}} + \frac{c_{21}c_{23}e_{24}}{k_{32}} + \frac{c_{23}c_{31}e_{24}}{k_{32}} \\
& \left. + \frac{c_{14}c_{31}e_{32}}{k_{32}} + \frac{c_{21}c_{34}e_{32}}{k_{32}} + \frac{c_{31}c_{34}e_{32}}{k_{32}} - \frac{c_{21}c_{32}e_{34}}{k_{32}} - \frac{c_{31}c_{32}e_{34}}{k_{32}} - \frac{e_{14}e_{32}k_{13}}{k_{32}} \right)
\end{aligned}$$

The dimension-neutral YM amplitudes are discussed e.g. in [8], explicit expressions are at www.damtp.cam.ac.uk/user/crm66/SYM/pss.html. To get the dimension-neutral GR amplitudes one can use the Kawai-Lewellen-Tye or KLT relations, for a review see [9].

B Mathematica computer code

Here we provide definitions from Theorem 1 and Lemma 3 as Wolfram Mathematica computer code. The code is rudimentary and not practical except for very low n .

```
(* Set number of legs, n>=3 *)
n=3;

(* Linear relations *)
relations=Flatten[{Table[{k[i,i],c[i,i],e[i,i]},{i,1,n}],
Table[{k[i,j]-k[j,i],e[i,j]-e[j,i]},{i,1,n},{j,1,n}],
Table[Sum[{k[i,j],c[i,j]},{i,1,n},{j,1,n}]}];
normalForm=First[Solve[Thread[relations==0]]];

(* Definition of D operators *)
z=KroneckerDelta;
Dk[i_,j_][f_]:= (1-z[i,j])*Sum[(1-z[a,b])*(z[i,a]*z[j,b]+z[i,b]*z[j,a]
-1/(n-2)*(z[i,a]+z[i,b]+z[j,a]+z[j,b])+2/((n-1)*(n-2)))*D[f,k[a,b]],{a,1,n},{b,1,n}];
Dc[i_,j_][f_]:= (1-z[i,j])*Sum[(1-z[a,j])*(z[i,a]-1/(n-1))*D[f,c[a,j]],{a,1,n}];
De[i_,j_][f_]:= (1-z[i,j])*(D[f,e[i,j]]+D[f,e[j,i]]);

(* Definition of A,B,C operators *)
AOp[i_][f_]:=Sum[1/2*k[p,q]*Dc[p,i][Dc[q,i][f]]+c[p,q]*Dc[p,i][De[q,i][f]]
+1/2*e[p,q]*De[p,i][De[q,i][f]],{p,1,n},{q,1,n}];
BOp[i_][f_]:=Sum[c[p,q]*Dk[i,p][De[i,q][f]]+e[p,q]*Dc[i,p][De[i,q][f]]
+k[p,q]*Dk[i,p][Dc[q,i][f]]+c[q,p]*Dc[i,p][Dc[q,i][f]]
-c[q,p]*Dk[p,q][De[p,i][f]]-e[p,q]*Dc[p,q][De[p,i][f]]
-k[p,q]*Dk[p,q][Dc[p,i][f]]-c[p,q]*Dc[p,i][Dc[p,q][f]],{p,1,n},{q,1,n}];
COpTilde[i_][f_]:=Sum[1/2*e[p,q]*Dc[i,p][Dc[i,q][f]]+c[q,p]*Dk[i,q][Dc[i,p][f]]
+1/2*k[p,q]*Dk[i,p][Dk[i,q][f]]-e[p,q]*Dc[i,p][Dc[p,q][f]]
-c[q,p]*Dk[p,q][Dc[i,p][f]]-c[p,q]*Dk[p,i][Dc[p,q][f]]
-k[p,q]*Dk[p,i][Dk[p,q][f]],{p,1,n},{q,1,n}];
COp[i_][f_]:=COpTilde[i][f]-1/n*Sum[COpTilde[p][f],{p,1,n}];

(* Example: C_1 annihilates the GR amplitude with n=3 *)
M3=(-c[2,3]*e[1,2]+c[2,1]*e[2,3]-c[1,2]*e[3,1])^2;
COp[1][M3] /. normalForm // Simplify (* yields 0 *)
```

C Additional formulas

This is an extension of Section 3 that provides definitions and formulas that are useful in the proof of the technical Lemma 16. We extend (3.7) to four linear maps:

$$\begin{array}{ccc} & M & \\ \alpha_J \nearrow & & \nwarrow \alpha_K \\ C(J_\bullet) & & C(K_\bullet) \\ \beta_J \nwarrow & & \nearrow \beta_K \end{array}$$

The maps β_J and β_K are explicit right-inverses of α_J and α_K , so $\alpha_J\beta_J = \mathbb{1}$ and $\alpha_K\beta_K = \mathbb{1}$. Hence $\beta_J\alpha_J$ and $\beta_K\alpha_K$ are projections $M \rightarrow M$. The direction Ξ is in the kernel of these projections. We choose β_J and β_K so that their images are contained in M^\perp and so that $\alpha_K\beta_J = 0$ and $\alpha_J\beta_K = 0$. The result of this is a useful decomposition

$$M = \mathbb{C}\Xi \oplus \text{image } \pi \oplus \underbrace{\text{image } \beta_J}_{\simeq C(J_\bullet)} \oplus \underbrace{\text{image } \beta_K}_{\simeq C(K_\bullet)} \quad (\text{C.1})$$

$\underbrace{\hspace{10em}}_{=M^\perp}$

where $\pi = \mathbb{1}_{M^\perp} - \beta_J\alpha_J - \beta_K\alpha_K$ is a projection, $\pi^2 = \pi$, with $\alpha_J\pi = \alpha_K\pi = 0$.

Lemma 17. *The map α_J (see Lemma 8) satisfies, and is equivalently defined by,*

$$\begin{array}{ll} D_{k_{jj'}}^\perp \mapsto D_{k_{jj'}} - \frac{(|K|-1)(1-\delta_{jj'})}{|I|-2}(D_{k_{j\bullet}} + D_{k_{j'\bullet}}) & D_{e_{jj'}}^\perp \mapsto D_{e_{jj'}} \\ D_{k_{jk}}^\perp, D_{k_{kj}}^\perp \mapsto \frac{|J|-1}{|I|-2} D_{k_{j\bullet}} & D_{e_{jk}}^\perp, D_{e_{kj}}^\perp \mapsto 0 \\ D_{k_{kk'}}^\perp \mapsto 0 & D_{e_{kk'}}^\perp \mapsto 0 \\ D_{c_{jj'}}^\perp \mapsto D_{c_{jj'}} - \frac{(|K|-1)(1-\delta_{jj'})}{|I|-1} D_{c_{\bullet j'}} & D_{c_{j\bullet}} \mapsto D_{c_{j\bullet}} \\ D_{c_{jk}}^\perp \mapsto 0 & D_{e_{j\bullet}} \mapsto D_{e_{j\bullet}} \\ D_{c_{kj}}^\perp \mapsto \frac{|J|}{|I|-1} D_{c_{\bullet j}} & D_{c_{k\bullet}} \mapsto 0 \\ D_{c_{kk'}}^\perp \mapsto 0 & D_{e_{k\bullet}} \mapsto 0 \\ & \Xi \mapsto 0 \end{array}$$

for all $j, j' \in J$ and $k, k' \in K$. The elements on the left are in $M \subseteq D_M$. Those on the right are in $C(J_\bullet) \subseteq D_{C(J_\bullet)}$, using Lemma 3 for J_\bullet . Analogous for α_K .

Proof. Let $X \mapsto Y$ be any of these claimed assignments. We must check that $Y(y) = X(\alpha_J^* y)$ for all $y \in C(J_\bullet)^*$ where here $Y(y)$ means applying Y to y as a function²¹. Here α_J^* is given by Lemma 8. Consider for example the case $X = D_{k_{jk}}^\perp$ and $y = k_{ab} \in C(J_\bullet)^*$ with $a, b \in J$. Then

$$\begin{aligned} X(\alpha_J^*(y)) &= D_{k_{jk}}^\perp(k_{ab}^\perp) = D_{k_{jk}}^\perp(k_{ab}) = D_{k_{jk}}(k_{ab}) + \frac{(|K|-1)(|J|-1)}{(|I|-1)(|I|-2)} \Xi(k_{ab}) \\ &= (1 - \delta_{ab}) \left(-\frac{1}{|I|-2} (\delta_{ja} + \delta_{jb}) \right) + \frac{2}{(|I|-1)(|I|-2)} + \frac{(|K|-1)(|J|-1)}{(|I|-1)(|I|-2)} \frac{2(1-\delta_{ab})}{|J|(|J|-1)} \end{aligned}$$

²¹With respect to bases and their dual bases, the matrix for α_J is the transpose of the matrix for α_J^* defined in Lemma 8, but we do not work with bases.

using Lemmas 3 and 5. Similarly,

$$Y(y) = \frac{|J|-1}{|I|-2} D_{k_{j\bullet}}(k_{ab}) = \frac{|J|-1}{|I|-2} (1 - \delta_{ab}) \left(-\frac{1}{|J|-1} (\delta_{ja} + \delta_{jb}) + \frac{2}{|J|(|J|-1)} \right)$$

using Lemma 3 for J_\bullet . In this example we see that $X(\alpha_J^*(y)) = Y(y)$. \square

Lemma 18 (Right-inverse). *The map α_J is surjective. There exists a unique right inverse $\beta_J : C(J_\bullet) \rightarrow M$ whose image is the subspace spanned by:*

- All $D_{k_{jj'}}^\perp, D_{c_{jj'}}^\perp, D_{e_{jj'}}^\perp$ with $j, j' \in J$.
- All $D_{c_{j\bullet}}, D_{e_{j\bullet}}$ with $j \in J$.

This right-inverse β_J maps

$$\begin{aligned} D_{k_{jj'}} &\mapsto D_{k_{jj'}}^\perp - \frac{(|K|-1)(1-\delta_{jj'})}{|K|(|J|-1)} \sum_{j'' \in J} (D_{k_{jj''}}^\perp + D_{k_{j'j''}}^\perp) \\ D_{k_{j\bullet}}, D_{c_{j\bullet}} &\mapsto -\frac{|I|-2}{|K|(|J|-1)} \sum_{j' \in J} D_{k_{jj'}}^\perp \\ D_{c_{jj'}} &\mapsto D_{c_{jj'}}^\perp - \frac{(|K|-1)(1-\delta_{jj'})}{|J||K|} \sum_{j'' \in J} D_{c_{jj''}}^\perp \\ D_{c_{j\bullet}} &\mapsto -\frac{|I|-1}{|J||K|} \sum_{j' \in J} D_{c_{j'j}}^\perp \\ D_{c_{j\bullet}} &\mapsto D_{c_{j\bullet}} \\ D_{e_{jj'}} &\mapsto D_{e_{jj'}}^\perp \\ D_{e_{j\bullet}}, D_{e_{\bullet j}} &\mapsto D_{e_{j\bullet}} \end{aligned}$$

for all $j, j' \in J$. We have $\alpha_K \beta_J = 0$. Analogous for β_K with $\alpha_J \beta_K = 0$.

Proof. First note that these assignments define a map β_J because we have specified it on a set that spans $C(J_\bullet)$ and it respects all relations, for instance $\sum_{j' \in J} D_{c_{j'j}} + D_{c_{\bullet j}}$ is zero in $C(J_\bullet)$ and must be mapped to zero, which it is. Now $\alpha_J \beta_J = \mathbb{1}$ is by direct calculation using Lemma 17, in particular α_J is surjective and β_J is injective. The image of β_J is as claimed. The image determines the right-inverse, so it is unique as claimed. \square

Corollary 19. *The internal direct sum decomposition (C.1) holds. Also:*

- The extension space $E \subseteq M^\perp$ is contained in $\text{image } \beta_J \oplus \text{image } \beta_K$.
- The subspace $\text{image } \pi \subseteq M^\perp$ is the span of all

$$\begin{aligned} \tilde{D}_{k_{jk}} &= D_{k_{jk}}^\perp - \frac{1}{|J|} \sum_{j' \in J} D_{k_{j'k}}^\perp \\ &\quad - \frac{1}{|K|} \sum_{k' \in K} D_{k_{jk'}}^\perp + \frac{1}{|J||K|} \sum_{j' \in J, k' \in K} D_{k_{j'k'}}^\perp \\ \tilde{D}_{c_{jk}} &= D_{c_{jk}}^\perp - \frac{1}{|J|} \sum_{j' \in J} D_{c_{j'k}}^\perp \\ \tilde{D}_{c_{kj}} &= D_{c_{kj}}^\perp - \frac{1}{|K|} \sum_{k' \in K} D_{c_{k'j}}^\perp \\ \tilde{D}_{e_{jk}} &= D_{e_{jk}}^\perp \end{aligned}$$

with $j \in J, k \in K$. Its dimension is $(2|J|-1)(2|K|-1)$. In D_M its elements commute with all elements in the images of α_J^*, α_K^* in particular

$$k_{jj'}^\perp, c_{jj'}^\perp, e_{jj'}^\perp, k_{kk'}^\perp, c_{kk'}^\perp, e_{kk'}^\perp, c_{j\bullet}, e_{j\bullet}, c_{k\bullet}, e_{k\bullet} \in M^*$$

for all $j, j' \in J$ and $k, k' \in K$.

D Relation to Loebbert, Mojaza, Plefka [1]

The paper [1] investigates a potential hidden conformal symmetry for GR amplitudes in general spacetime dimension, and provides conceptual backing relating to soft dilatons. No attempt is made to review these aspects here. The purpose of this appendix is merely to clarify the relation between the conjecture for GR amplitudes in [1] and the new formulation in terms of the differential operators B_i, C_i in Theorem 1. (Here we do not discuss the more tentative conjecture for YM amplitudes in [1]. Nor do we discuss the operators A_i that are not part of the conjecture in [1].) The notation used in this appendix uses notation from [1], notation from this paper, and additional notation used only in this appendix.

- The conjecture we are referring to are equations (5.12) and (5.13) in [1] for general $n \geq 3$. The cases $n = 3, 4, 5, 6$ are separately discussed in [1].
- In [1], the primary differential operators are certain conformal generators. The special conformal generators are expressed in terms of operators²² F_i, G_i in [1] closely related to B_i, C_i in this paper respectively. In fact, superficially, the formulas in Appendix A of [1] coincide with the right hand sides of (1.3b), (1.3c). The symbols entering the formulas do however not have exactly the same meaning as we now discuss.
- In [1], ordinary partial derivatives $\partial_{s_{ij}}, \partial_{w_{ij}}, \partial_{e_{ij}}$ (the variables s, w, e in [1] correspond to k, c, e in this paper respectively) on the ambient vector space are used that are not compatible with the relations (1.1). Let us refer to A as the ambient vector space and to $C \subseteq A$ as the subspace given by (1.1). It is called ‘constraint surface’ in [1]. Note that $F_i, G_i : \text{Frac}(R_A) \rightarrow \text{Frac}(R_A)$. The conjecture in [1] is that the GR amplitudes are annihilated by $Y(F_i), Y(G_i)$ where, in notation used only in this appendix,

$$Y(X) = \sum_\pi R X \pi E \quad : \quad \text{Frac}(R_C) \rightarrow \text{Frac}(R_C)$$

where (all maps are linear):

- $E : \text{Frac}(R_C) \rightarrow \text{Frac}(R_A)$ extends functions to the ambient space, it is the pullback along a specific projection $A \rightarrow A$ with image C . This is referred to as ‘resolving the constraints’ in [1] and for the specific choice of E we refer to [1].
- $\pi : \text{Frac}(R_A) \rightarrow \text{Frac}(R_A)$ is a permutation of the index set I , and \sum_π is a sum over all. So $\sum_\pi \pi$ corresponds to symmetrization.

²²Actually in [1] the notation F_i, G_i is the result of applying certain operators to the amplitudes, but in this appendix we use this notation for the differential operators themselves.

- $X : \text{Frac}(R_A) \rightarrow \text{Frac}(R_A)$ is one of the partial differential operators of interest, F_i or G_i in the notation of [1], on the ambient space.
- $R : \text{Frac}(R_A) \rightarrow \text{Frac}(R_C)$ restricts functions from A to C ²³.
- Note that $Y(X)$ is not a partial differential operator on C because π is not local. But one can define a new, different operator $Y_{\text{loc}}(X)$ that has the virtue of being local in fact a polynomial differential operator on C ; and that nevertheless coincides with $Y(X)$ when acting on permutation invariant elements of $\text{Frac}(R_C)$. Namely²⁴

$$Y_{\text{loc}}(X) = \sum_{\pi} R X \pi E \pi^{-1} \quad : \quad \text{Frac}(R_C) \rightarrow \text{Frac}(R_C)$$

Clearly $Y(X)f = Y_{\text{loc}}(X)f$ for all rational functions $f \in \text{Frac}(R_C)$ that are permutation invariant, $\pi f = f$ for all π . But $Y_{\text{loc}}(X)$ is local, because $Y_{\text{loc}}(X) = \sum_{\pi} R X E_{\pi}$ where $E_{\pi} = \pi E \pi^{-1} : \text{Frac}(R_C) \rightarrow \text{Frac}(R_A)$ is simply the pullback along another projection $A \rightarrow A$ with image C .

- The GR amplitudes are permutation invariant. Therefore the conjecture in [1] that $Y(F_i)$, $Y(G_i)$ annihilate the GR amplitude is equivalent to $Y_{\text{loc}}(F_i)$, $Y_{\text{loc}}(G_i)$ annihilating the GR amplitude.
- The B_i , C_i in (1.3b), (1.3c) are, up to normalization, the operators $Y_{\text{loc}}(F_i)$, $Y_{\text{loc}}(G_i)$ but presented directly as differential operators on $C = C(I)$. A detailed translation to (1.3b), (1.3c) is omitted. There are terms proportional to the conformal dimension Δ in F_i , G_i , see [1], but they do not contribute to $Y_{\text{loc}}(F_i)$, $Y_{\text{loc}}(G_i)$ which therefore are independent of Δ .

References

- [1] Loebbert F., Mojaza M. and Plefka J., [JHEP 2018.5, 208 \(2018\)](#)
Hidden conformal symmetry in tree-level graviton scattering
- [2] Britto R., Cachazo F., Feng B., Witten E., [Phys. Rev. Lett. 94 \(18\), 181602 \(2005\)](#)
Direct proof of the tree-level scattering amplitude recursion relation in Yang-Mills theory
- [3] Grayson D.R. and Stillman M.E., www.math.uiuc.edu/Macaulay2/
Macaulay2, a software system for research in algebraic geometry
- [4] Castro-Jiménez F.J. and Leykin A., arxiv.org/abs/1110.0182 (2011)
Computing localizations iteratively
- [5] Zeilberger D., Journal of Computational and applied mathematics 32.3, 321-368 (1990)
A holonomic systems approach to special functions identities
- [6] Nützi A. and Reiterer M., arxiv.org/abs/1812.06454 (2018)
Scattering amplitudes in YM and GR as minimal model brackets and their recursive characterization
- [7] Dixon L., arxiv.org/abs/hep-ph/9601359 (1996)
Calculating scattering amplitudes efficiently

²³Actually R is only partially defined but we gloss over that since it is inconsequential here.

²⁴Here $\pi^{-1} : \text{Frac}(R_C) \rightarrow \text{Frac}(R_A)$.

- [8] Bjerrum-Bohr N.E.J., Bourjaily J.L., Damgaard P.H. and Feng B.,
Nuclear Physics B, 913, 964-986 (2016)
Analytic representations of Yang-Mills amplitudes
- [9] Søndergaard T., Advances in High Energy Physics (2012)
Perturbative gravity and gauge theory relations: a review