

# Geometric Estimates in Interpolation by Linear Functions on a Euclidean Ball

Mikhail Nevskii\*

May 9, 2019

## Abstract

Let  $B_n$  be the Euclidean unit ball in  $\mathbb{R}^n$  given by the inequality  $\|x\| \leq 1$ ,  $\|x\| := \left( \sum_{i=1}^n x_i^2 \right)^{\frac{1}{2}}$ . By  $C(B_n)$  we mean the space of continuous functions  $f : B_n \rightarrow \mathbb{R}$  with the norm  $\|f\|_{C(B_n)} := \max_{x \in B_n} |f(x)|$ . The symbol  $\Pi_1(\mathbb{R}^n)$  denotes the set of polynomials in  $n$  variables of degree  $\leq 1$ , i. e., the set of linear functions upon  $\mathbb{R}^n$ . Assume  $x^{(1)}, \dots, x^{(n+1)}$  are the vertices of an  $n$ -dimensional nondegenerate simplex  $S \subset B_n$ . The interpolation projector  $P : C(B_n) \rightarrow \Pi_1(\mathbb{R}^n)$  corresponding to  $S$  is defined by the equalities  $Pf(x^{(j)}) = f(x^{(j)})$ . Denote by  $\|P\|_{B_n}$  the norm of  $P$  as an operator from  $C(B_n)$  onto  $C(B_n)$ . We describe the approach in which  $\|P\|_{B_n}$  can be estimated from below via the volume of  $S$ .

Keywords: simplex, ball, linear interpolation, projector, norm, estimate

## 1 Main Definitions

We always suppose  $n \in \mathbb{N}$ . An element  $x \in \mathbb{R}^n$  will be written in the form  $x = (x_1, \dots, x_n)$ . By definition,

$$\|x\| := \left( \sum_{i=1}^n x_i^2 \right)^{\frac{1}{2}}, \quad B_n := \{x \in \mathbb{R}^n : \|x\| \leq 1\}, \quad Q_n := [0, 1]^n.$$

---

\*Department of Mathematics, P.G. Demidov Yaroslavl State University, Sovetskaya str., 14, Yaroslavl, 150003, Russia orcid.org/0000-0002-6392-7618 mnevsk55@yandex.ru

The notation  $L(n) \asymp M(n)$  means that there exist  $c_1, c_2 > 0$  not depending on  $n$  such that  $c_1 M(n) \leq L(n) \leq c_2 M(n)$ . By  $\Pi_1(\mathbb{R}^n)$  we mean the set of polynomials in  $n$  variables  $\leq 1$ , i.e., the set of linear functions upon  $\mathbb{R}^n$ .

Let  $S$  be a nondegenerate simplex in  $\mathbb{R}^n$  with vertices  $x^{(j)} = (x_1^{(j)}, \dots, x_n^{(j)})$ ,  $1 \leq j \leq n+1$ . Consider *the vertex matrix (or the node matrix)*

$$\mathbf{A} := \begin{pmatrix} x_1^{(1)} & \dots & x_n^{(1)} & 1 \\ x_1^{(2)} & \dots & x_n^{(2)} & 1 \\ \vdots & \vdots & \vdots & \vdots \\ x_1^{(n+1)} & \dots & x_n^{(n+1)} & 1 \end{pmatrix}.$$

We have the equality  $\text{vol}(S) = \frac{|\det(\mathbf{A})|}{n!}$ . Put  $\mathbf{A}^{-1} = (l_{ij})$ . Let us define  $\lambda_j$  as polynomials from  $\Pi_1(\mathbb{R}^n)$  whose coefficients form the columns of  $\mathbf{A}^{-1}$ , i.e.,  $\lambda_j(x) := l_{1j}x_1 + \dots + l_{nj}x_n + l_{n+1,j}$ . We call  $\lambda_j$  *the basic Lagrange polynomials* corresponding to this simplex. The numbers  $\lambda_j(x)$  are the barycentric coordinates of a point  $x \in \mathbb{R}^n$  with respect to  $S$ .

For a convex body  $\Omega \subset \mathbb{R}^n$ , denote by  $C(\Omega)$  the space of continuous functions  $f : \Omega \rightarrow \mathbb{R}$  with the uniform norm

$$\|f\|_{C(\Omega)} := \max_{x \in \Omega} |f(x)|.$$

We say that an interpolation projector  $P : C(\Omega) \rightarrow \Pi_1(\mathbb{R}^n)$  corresponds to a simplex  $S \subset \Omega$  if interpolation nodes of  $P$  coincide with vertices  $x^{(j)}$  of this simplex. Projector  $P$  is given by the equalities  $Pf(x^{(j)}) = f(x^{(j)})$ . We have the following analogue of the Lagrange interpolation formula:

$$Pf(x) = \sum_{j=1}^{n+1} f(x^{(j)}) \lambda_j(x). \quad (1)$$

Denote by  $\|P\|_\Omega$  the norm of  $P$  as an operator from  $C(\Omega)$  to  $C(\Omega)$ . It follows from (1) that

$$\|P\|_\Omega = \max_{x \in \Omega} \sum_{j=1}^{n+1} |\lambda_j(x)|.$$

If  $\Omega$  is a convex polytope in  $\mathbb{R}^n$  (e.g.,  $\Omega = Q_n$ ), this equality is equivalent to the formula

$$\|P\|_\Omega = \max_{x \in \text{ver}(\Omega)} \sum_{j=1}^{n+1} |\lambda_j(x)|,$$

where  $\text{ver}(\Omega)$  is the set of vertices of  $\Omega$ .

Let us define  $\theta_n(\Omega)$  as the minimal value of  $\|P\|_\Omega$  under the conditions  $x^{(j)} \in \Omega$ . In the case  $\Omega = Q_n$  various relations for the numbers  $\theta_n(\Omega)$ , including the equivalence  $\theta_n(\Omega) \asymp \sqrt{n}$ , were obtained by the author earlier. This results are systematized in [1]. Further on some estimates were improved (see [2], [3], [4] and references in these papers).

This paper deals with the case  $\Omega = B_n$ . We describe the approach when the norm of an interpolation projector  $P : C(B_n) \rightarrow \Pi_1(\mathbb{R}^n)$  can be estimated from below through the volume of the corresponding simplex. The essential feature of this approach is the application of the classical Legendre polynomials. We prove that  $\theta_n(B_n) \asymp \sqrt{n}$ . In other words, the interpolation projector corresponding to a regular simplex inscribed into the boundary sphere has the norm equivalent to the minimal possible.

## 2 Estimation of $\|P\|_{B_n}$ via the Volume of $S$

The *standardized Legendre polynomial of degree  $n$*  is the function

$$\chi_n(t) := \frac{1}{2^n n!} [(t^2 - 1)^n]^{(n)}$$

(the Rodrigues formula). For properties of  $\chi_n$ , see [6], [7]. The Legendre polynomials are orthogonal on  $[-1, 1]$  with respect to the weight  $w(t) = 1$ . It is known that  $\chi_n(1) = 1$ ; if  $n \geq 1$ , then  $\chi_n(t)$  strictly increases for  $t \geq 1$ . Denote by  $\chi_n^{-1}$  the function inverse to  $\chi_n$  on the halfline  $[1, +\infty)$ .

The appearance of Legendre polynomials in our questions is connected with such their property. For  $\gamma \geq 1$ , consider the set

$$E_{n,\gamma} := \left\{ x \in \mathbb{R}^n : \sum_{j=1}^n |x_j| + \left| 1 - \sum_{j=1}^n x_j \right| \leq \gamma \right\}.$$

In 2003 the author established that

$$\text{mes}_n(E_{n,\gamma}) = \frac{\chi_n(\gamma)}{n!}. \quad (2)$$

(the proof is also given in [1]). Utilizing this equality he managed to obtain the lower estimates for projector's norms related to linear interpolation on the unit cube  $Q_n$ . The following relations take place:

$$\theta_n(Q_n) \geq \chi_n^{-1} \left( \frac{1}{\nu_n} \right) = \chi_n^{-1} \left( \frac{n!}{h_n} \right). \quad (3)$$

Here  $\nu_n$  is the maximum volume of a simplex contained in  $Q_n$  and  $h_n$  is the maximum value of 0/1-determinant of order  $n$ . Applying the properties of

$\chi_n$  the author got from (3) some more visible inequalities, e. g.,  $\theta_n(Q_n) > \frac{1}{e}\sqrt{n-1}$ . This estimate occurred to be sharp concerning to  $n$  which led to the equivalence  $\theta_n(Q_n) \asymp \sqrt{n}$ .

Further on we will extend this approach to linear interpolation of functions given on the unit ball  $B_n$ . Denote  $\varkappa_n := \text{vol}(B_n)$ . By  $\sigma_n$  we mean the volume of a regular simplex inscribed into  $B_n$ .

**Theorem 1.** *Assume  $P : C(B_n) \rightarrow \Pi_1(\mathbb{R}^n)$  is an arbitrary interpolation projector. Then for the corresponding simplex  $S \subset B_n$  and the node matrix  $\mathbf{A}$  we have*

$$\|P\|_{B_n} \geq \chi_n^{-1} \left( \frac{\varkappa_n}{\text{vol}(S)} \right) = \chi_n^{-1} \left( \frac{n! \varkappa_n}{|\det(\mathbf{A})|} \right). \quad (4)$$

*Proof.* As known, a regular simplex which is inscribed into a ball has the maximum possible volume among all simplices being contained in this ball. Therefore,  $\text{vol}(S) = \frac{|\det(\mathbf{A})|}{n!} \leq \sigma_n$ . For each  $i = 1, \dots, n$ , let us subtract from the  $i$ th row of  $\mathbf{A}$  its  $(n+1)$ th row. Denote by  $\mathbf{B}$  the submatrix of order  $n$  which stands in the first  $n$  rows and columns of the result matrix. Then

$$|\det(\mathbf{B})| = |\det(\mathbf{A})| = n! \text{vol}(S) \leq n! \sigma_n.$$

In other words,

$$\frac{n! \sigma_n}{|\det(\mathbf{B})|} \geq 1. \quad (5)$$

Let  $x^{(j)}$  be the vertices and  $\lambda_j$  be the basic Lagrange polynomials of the simplex  $S$ . Since  $\lambda_1(x), \dots, \lambda_{n+1}(x)$  are the barycentric coordinates of a point  $x$ , we have

$$\|P\|_{B_n} = \max_{x \in B_n} \sum_{j=1}^{n+1} |\lambda_j(x)| = \max \left\{ \sum_{j=1}^{n+1} |\beta_j| : \sum_{j=1}^{n+1} \beta_j = 1, \sum_{j=1}^{n+1} \beta_j x^{(j)} \in B_n \right\}.$$

Let us replace  $\beta_{n+1}$  with the equal value  $1 - \sum_{j=1}^n \beta_j$ . The condition  $\sum_{j=1}^{n+1} \beta_j x^{(j)} \in B_n$  is equivalent to  $\sum_{j=1}^n \beta_j (x^{(j)} - x^{(n+1)}) \in B' := B_n - x^{(n+1)}$ . This means

$$\|P\|_{B_n} = \max \left\{ \sum_{j=1}^n |\beta_j| + \left| 1 - \sum_{j=1}^n \beta_j \right| \right\}. \quad (6)$$

Maximum in (6) is taken upon  $\beta_j$  such that  $\sum_{j=1}^n \beta_j (x^{(j)} - x^{(n+1)}) \in B'$ .

Consider the nondegenerate linear operator  $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$  which maps a point  $\beta = (\beta_1, \dots, \beta_n)$  into a point  $x = (x_1, \dots, x_n)$  according to the rule

$$x = F(\beta) := \sum_{j=1}^n \beta_j (x^{(j)} - x^{(n+1)}).$$

We have the matrix equality  $F(\beta) = (\beta_1, \dots, \beta_n)\mathbf{B}$ , where  $\mathbf{B}$  is the above introduced  $(n \times n)$ -matrix with the elements  $b_{ij} = x_j^{(i)} - x_j^{(n+1)}$ . Put

$$\gamma^* := \chi_n^{-1} \left( \frac{n! \varkappa_n}{|\det(\mathbf{B})|} \right).$$

Since  $\varkappa_n \geq \sigma_n$ , it follows from (5) that  $\gamma^*$  is defined correctly. Note the equality

$$\chi_n(\gamma^*) = \frac{n! \varkappa_n}{|\det(\mathbf{B})|}. \quad (7)$$

Now suppose  $1 \leq \gamma < \gamma^*$ ,

$$E_{n,\gamma} = \left\{ \beta = (\beta_1, \dots, \beta_n) \in \mathbb{R}^n : \sum_{j=1}^n |\beta_j| + \left| 1 - \sum_{j=1}^n \beta_j \right| \leq \gamma \right\}.$$

Let us show that  $B' \not\subset F(E_{n,\gamma})$ . It is sufficient to get the inequality  $\text{mes}_n(F(E_{n,\gamma})) < \text{mes}_n(B') = \varkappa_n$ . This is really so:

$$\begin{aligned} \text{mes}_n(F(E_{n,\gamma})) &< \text{mes}_n(F(E_{n,\gamma^*})) = |\det \mathbf{B}| \cdot \text{mes}_n(E_{n,\gamma^*}) = \\ &= |\det \mathbf{B}| \cdot \frac{\chi_n(\gamma^*)}{n!} = \varkappa_n. \end{aligned}$$

We have applied (2) and (7). Thus, for every  $\varepsilon > 0$ , there exists a point  $x^{(\varepsilon)}$  with the properties

$$x^{(\varepsilon)} = \sum_{j=1}^n \beta_j^{(\varepsilon)} (x^{(j)} - x^{(n+1)}) \in B', \quad \sum_{j=1}^n |\beta_j^{(\varepsilon)}| + \left| 1 - \sum_{j=1}^n \beta_j^{(\varepsilon)} \right| \geq \gamma^* - \varepsilon.$$

In view of (6) this gives  $\|P\|_{B_n} \geq \gamma^* - \varepsilon$ . Since  $\varepsilon > 0$  is an arbitrary, we obtain

$$\|P\| \geq \gamma^* = \chi_n^{-1} \left( \frac{n! \varkappa_n}{|\det(\mathbf{B})|} \right) = \chi_n^{-1} \left( \frac{n! \varkappa_n}{|\det(\mathbf{A})|} \right) = \chi_n^{-1} \left( \frac{\varkappa_n}{\text{vol}(S)} \right).$$

The theorem is proved. □

**Corollary 1.** *For each  $n$ ,*

$$\theta_n(B_n) \geq \chi_n^{-1} \left( \frac{\varkappa_n}{\sigma_n} \right). \quad (8)$$

*Proof.* Let  $P : C(B_n) \rightarrow \Pi_1(\mathbb{R}^n)$  be an arbitrary interpolation projector. Since the volume of the corresponding simplex is not greater than  $\sigma_n$ , inequality (4) yields

$$\|P\|_{B_n} \geq \chi_n^{-1} \left( \frac{\varkappa_n}{\text{vol}(S)} \right) \geq \chi_n^{-1} \left( \frac{\varkappa_n}{\sigma_n} \right).$$

This gives (8). □

It is known that

$$\varkappa_n = \frac{\pi^{\frac{n}{2}}}{\Gamma\left(\frac{n}{2} + 1\right)}, \quad \sigma_n = \frac{1}{n!} \sqrt{n+1} \left( \frac{n+1}{n} \right)^{\frac{n}{2}}, \quad (9)$$

$$\varkappa_{2k} = \frac{\pi^k}{k!}, \quad \varkappa_{2k+1} = \frac{2^{k+1}\pi^k}{(2k+1)!!} = \frac{2(k!)(4\pi)^k}{(2k+1)!}. \quad (10)$$

Therefore, the estimate (8) can be made more concrete.

**Corollary 2.** *For every  $n$ ,*

$$\theta_n(B_n) \geq \chi_n^{-1} \left( \frac{\pi^{\frac{n}{2}} n!}{\Gamma\left(\frac{n}{2} + 1\right) \sqrt{n+1} \left(\frac{n+1}{n}\right)^{\frac{n}{2}}} \right). \quad (11)$$

*If  $n = 2k$ , then (11) is equivalent to the inequality*

$$\theta_{2k}(B_{2k}) \geq \chi_{2k}^{-1} \left( \frac{\pi^k (2k)!}{k! \sqrt{2k+1} \left(\frac{2k+1}{2k}\right)^k} \right). \quad (12)$$

*For  $n = 2k + 1$  we have*

$$\theta_{2k+1}(B_{2k+1}) \geq \chi_{2k+1}^{-1} \left( \frac{2(k!)(4\pi)^k}{\sqrt{2k+2} \left(\frac{2k+2}{2k+1}\right)^{\frac{2k+1}{2}}} \right). \quad (13)$$

*Proof.* It is sufficient to apply (8), (9), and (10). □

**Corollary 3.** *Suppose  $P$  is a minimal interpolation projector, i. e.,  $\|P\|_{B_n} = \theta_n(B_n)$ . Then*

$$\text{vol}(S) \geq \frac{\varkappa_n}{\chi_n(\sqrt{n+1})} = \frac{\pi^{\frac{n}{2}}}{\Gamma\left(\frac{n}{2} + 1\right) \chi_n(\sqrt{n+1})}. \quad (14)$$

*Proof.* It was proved in [5] that  $\theta_n(B_n) \leq \sqrt{n+1}$ . Hence, if  $P$  is a minimal projector, then

$$\sqrt{n+1} \geq \|P\|_{B_n} \geq \chi_n^{-1} \left( \frac{\varkappa_n}{\text{vol}(S)} \right).$$

We have made use of (4). It remains to compare the boundary values and take into account (9).  $\square$

Since  $|\det(\mathbf{A})| = n! \text{vol}(S)$ , relation (14) also implies an estimate for the determinant of the node matrix corresponding to a minimal projector. Namely,  $|\det(\mathbf{A})| \geq \varepsilon_n$ , where  $\varepsilon_n$  is  $n!$  times more than the right-hand part of (14). Without giving the value of  $\varepsilon_n$  this restriction was used in [5].

### 3 The Relation $\theta_n(B_n) \asymp \sqrt{n}$

The Stirling formula  $n! = \sqrt{2\pi n} \left(\frac{n}{e}\right)^n e^{\frac{\zeta_n}{12n}}$ ,  $0 < \zeta_n < 1$ , yields

$$\sqrt{2\pi n} \left(\frac{n}{e}\right)^n < n! < \sqrt{2\pi n} \left(\frac{n}{e}\right)^n e^{\frac{1}{12n}}. \quad (15)$$

Also we will need the following estimates which were proved in [1, Section 3.4.2]:

$$\chi_{2k}^{-1}(s) > \left( \frac{(k!)^2 s}{(2k)!} \right)^{\frac{1}{2k}}, \quad \chi_{2k+1}^{-1}(s) > \left( \frac{(k+1)! k! s}{(2k+1)!} \right)^{\frac{1}{2k+1}}. \quad (16)$$

**Theorem 2.** *There exist a constant  $c > 0$  not depending on  $n$  such that*

$$\theta_n(B_n) > c\sqrt{n}. \quad (17)$$

*The inequality (17) takes place, e. g., with the constant*

$$c = \frac{\sqrt[3]{\pi}}{\sqrt{12e} \cdot \sqrt[6]{3}} = 0.2135\dots$$

*Proof.* First let  $n = 2k$  be even. Making use of (12) and (16), we get

$$\theta_{2k}(B_{2k}) \geq \chi_{2k}^{-1} \left( \frac{\pi^k (2k)!}{k! \sqrt{2k+1} \left(\frac{2k+1}{2k}\right)^k} \right) > \left( \frac{\pi^k (2k)! (k!)^2}{k! \sqrt{2k+1} \left(\frac{2k+1}{2k}\right)^k (2k)!} \right)^{\frac{1}{2k}} =$$

$$= \left( \frac{\pi^k k!}{\sqrt{2k+1} \left( \frac{2k+1}{2k} \right)^k} \right)^{\frac{1}{2k}}.$$

Let us estimate  $k!$  from below applying (15):

$$\theta_{2k}(B_{2k}) > \left( \frac{\pi^k \sqrt{2\pi k} \left( \frac{k}{e} \right)^k}{\sqrt{2k+1} \left( \frac{2k+1}{2k} \right)^k} \right)^{\frac{1}{2k}} = \sqrt{\frac{\pi}{e}} \left( \frac{2\pi k}{2k+1} \right)^{\frac{1}{4k}} \sqrt{\frac{2k}{2k+1}} \cdot \sqrt{k}.$$

Since  $n = 2k$ , for each even  $n$ ,

$$\theta_n(B_n) > \sqrt{\frac{\pi}{2e}} \left( \frac{\pi n}{n+1} \right)^{\frac{1}{2n}} \sqrt{\frac{n}{n+1}} \cdot \sqrt{n} > \sqrt{\frac{\pi}{3e}} \cdot \sqrt{n}. \quad (18)$$

We have utilized the inequalities  $\frac{\pi n}{n+1} > 1$ ,  $\frac{n}{n+1} > \frac{2}{3}$ . Remark that  $\sqrt{\frac{\pi}{3e}} = 0.6206\dots$

Now suppose  $n$  is odd, i. e.,  $n = 2k + 1$ . In this case (13) and (16) give

$$\begin{aligned} \theta_{2k+1}(B_{2k+1}) &\geq \chi_{2k+1}^{-1} \left( \frac{2(k!)(4\pi)^k}{\sqrt{2k+2} \left( \frac{2k+2}{2k+1} \right)^{\frac{2k+1}{2}}} \right) > \\ &> \left( \frac{2(k!)(4\pi)^k (k+1)!k!}{\sqrt{2k+2} \left( \frac{2k+2}{2k+1} \right)^{\frac{2k+1}{2}} (2k+1)!} \right)^{\frac{1}{2k+1}} = A^{\frac{1}{2k+1}} \cdot B^{\frac{1}{2k+1}}; \\ A &:= \frac{(k!)^2 (k+1)!}{(2k+1)!}, \quad B := \frac{\sqrt{2}(4\pi)^k}{\sqrt{k+1} \left( \frac{2k+2}{2k+1} \right)^{\frac{2k+1}{2}}}. \end{aligned}$$

From (15), it follows that

$$\begin{aligned} A &> \frac{2\pi k \left( \frac{k}{e} \right)^{2k} \cdot \sqrt{2\pi(k+1)} \left( \frac{k+1}{e} \right)^{k+1}}{\sqrt{2\pi(2k+1)} \left( \frac{2k+1}{e} \right)^{2k+1} e^{\frac{1}{12(2k+1)}}} = \\ &= 2\pi e^{-k - \frac{1}{12(2k+1)}} k^{2k+1} (k+1)^{k+1+\frac{1}{2}} (2k+1)^{-\frac{1}{2}-(2k+1)}, \\ A^{\frac{1}{2k+1}} &> (2\pi)^{\frac{1}{2k+1}} e^{-\frac{k}{2k+1} - \frac{1}{12(2k+1)^2}} k(k+1)^{\frac{k+1+\frac{1}{2}}{2k+1}} (2k+1)^{-\frac{1}{2(2k+1)}-1}. \end{aligned}$$

Also

$$\begin{aligned} (2\pi)^{\frac{1}{2k+1}} &> 1, \quad \frac{k}{2k+1} + \frac{1}{12(2k+1)^2} < \frac{1}{2}, \\ k(2k+1)^{-\frac{1}{2(2k+1)}-1} &= \frac{k}{2k+1} (2k+1)^{-\frac{1}{2(2k+1)}} > \frac{1}{3} (2k+1)^{-\frac{1}{2(2k+1)}} > \frac{1}{3\sqrt[6]{3}}. \end{aligned}$$



Consequently,

$$A^{\frac{1}{2k+1}} > \frac{1}{3\sqrt[6]{3}\sqrt{e}}(k+1)^{\frac{k+1+\frac{1}{2}}{2k+1}} > \frac{1}{3\sqrt[6]{3}\sqrt{e}}\sqrt{k+1}.$$

Now let us estimate  $B^{\frac{1}{2k+1}}$ :

$$\begin{aligned} B^{\frac{1}{2k+1}} &= \left( \frac{\sqrt{2}(4\pi)^k}{\sqrt{k+1} \left(\frac{2k+2}{2k+1}\right)^{\frac{2k+1}{2}}} \right)^{\frac{1}{2k+1}} = \\ &= (\sqrt{2})^{\frac{1}{2k+1}} (4\pi)^{\frac{k}{2k+1}} \sqrt{\frac{2k+1}{2k+2}} (k+1)^{-\frac{1}{2(2k+1)}} > \\ &> \sqrt[3]{4\pi} \cdot \sqrt{\frac{3}{4}} \cdot 2^{-\frac{1}{6}} = \sqrt{\frac{3}{2}} \cdot \sqrt[3]{\pi}. \end{aligned}$$

Therefore,

$$\theta_{2k+1}(B_{2k+1}) > A^{\frac{1}{2k+1}} \cdot B^{\frac{1}{2k+1}} > \sqrt{\frac{3}{2}} \cdot \frac{\sqrt[3]{\pi}}{3\sqrt[6]{3}\sqrt{e}} \cdot \sqrt{k+1} = \frac{\sqrt[3]{\pi}}{\sqrt{6e}\sqrt[6]{3}} \cdot \sqrt{k+1}.$$

Since  $n = 2k + 1$ , we have  $\sqrt{k+1} = \frac{1}{\sqrt{2}}\sqrt{2k+2} > \frac{1}{\sqrt{2}}\sqrt{n}$ . For every odd  $n$ ,

$$\theta_n(B_n) > \frac{\sqrt[3]{\pi}}{\sqrt{12e} \cdot \sqrt[6]{3}} \cdot \sqrt{n}. \quad (19)$$

The constant 0.2135... from the right-hand part of (19) is less than the constant from the above inequality (18) for even  $n$ . Hence, (19) is true for all  $n \in \mathbb{N}$ . This completes the proof.  $\square$

**Corollary 4.**  $\theta_n(B_n) \asymp \sqrt{n}$ .

*Proof.* In [5] it was proved that  $\theta_n(B_n) \leq \sqrt{n+1}$ . Consequently, the lower estimate  $\theta_n(B_n) > c\sqrt{n}$  is precise with respect to dimension  $n$ .  $\square$

**Corollary 5.** Assume  $P : C(B_n) \rightarrow \Pi_1(\mathbb{R}^n)$  is the interpolation projector whose nodes coincide with vertices of a regular simplex being inscribed into the boundary sphere  $\|x\| = 1$ . When  $\|P\|_{B_n} \asymp \theta_n(B_n)$ .

*Proof.* As it was shown in [5],  $\sqrt{n} \leq \|P\|_{B_n} \leq \sqrt{n+1}$ . It remains to utilize the previous corollary.  $\square$

Our results mean that any interpolation projector  $P$  corresponding to an inscribed regular simplex has the norm equivalent to the minimal possible. The equality  $\|P\|_{B_n} = \theta_n(B_n)$  remains proved only for  $1 \leq n \leq 4$ .

The author is grateful to A. Yu. Ukhlov for some useful computer calculations.

## References

1. Nevskii, M. V., *Geometricheskie ocenki v polinomialnoi interpoliacii* (Geometric Estimates in Polynomial Interpolation), Yaroslavl': Yarosl. Gos. Univ., 2012 (in Russian).
2. Nevskii, M. V., and Ukhalov, A. Yu., On numerical characteristics of a simplex and their estimates, *Model. Anal. Inform. Sist.*, 2016, vol. 23, no. 5, pp. 603–619 (in Russian). English transl.: *Aut. Control Comp. Sci.*, 2017, vol. 51, no. 7, pp. 757–769.
3. Nevskii, M. V., and Ukhalov, A. Yu., New estimates of numerical values related to a simplex, *Model. Anal. Inform. Sist.*, 2017, vol. 24, no. 1, pp. 94–110 (in Russian). English transl.: *Aut. Control Comp. Sci.*, 2017, vol. 51, no. 7, pp. 770–782.
4. Nevskii, M. V., and Ukhalov, A. Yu., On optimal interpolation by linear functions on an  $n$ -dimensional cube, *Model. Anal. Inform. Sist.*, 2018, vol. 25, no. 3, pp. 291–311 (in Russian). English transl.: *Aut. Control Comp. Sci.*, 2018, vol. 52, no. 7, pp. 828–842.
5. Nevskii, M. V., and Ukhalov, A. Yu., Linear interpolation on a Euclidean ball in  $\mathbb{R}^n$  (to appear).
6. Szego, G., *Orthogonal Polynomials*, New York: American Mathematical Society, 1959.
7. Suetin, P. K. *Klassicheskie ortogonal'nye mnogochleny*, Moscow: Nauka, 1979 (in Russian).