

A CLASS OF ANISOTROPIC EXPANDING CURVATURE FLOWS

WEIMIN SHENG AND CAIHONG YI

ABSTRACT. In this paper, we consider an expanding flow of smooth, closed, uniformly convex hypersurfaces in Euclidean R^{n+1} with speed $u^\alpha \sigma_k^\beta$ firstly, where u is support function of the hypersurface, $\alpha, \beta \in R^1$, and $\beta > 0$, σ_k is the k -th symmetric polynomial of the principle curvature radii of the hypersurface, k is an integer and $1 \leq k \leq n$. For $\alpha \leq 1 - k\beta$, $\beta > \frac{1}{k}$ we prove that the flow has a unique smooth and uniformly convex solution for all time, and converges smoothly after normalisation, to a sphere centered at the origin. Moreover, for $\alpha \leq 1 - k\beta$, $\beta > \frac{1}{k}$, we prove that the flow with the speed $f u^\alpha \sigma_k^\beta$ exists for all time and converges smoothly after normalisation to a soliton which is a solution of $f u^{\alpha-1} \sigma_k^\beta = c$ provided that f is a smooth positive function on S^n and satisfies that $(\nabla_i \nabla_j f^{\frac{1}{1+k\beta-\alpha}} + \delta_{ij} f^{\frac{1}{1+k\beta-\alpha}})$ is positive definite. When $\beta = 1$, our argument provides a proof to the well-known L_p Christoffel-Minkowski problem for the case $p \geq k + 1$ where $p = 2 - \alpha$, which is identify with Ivaki's recent result. Especially, we obtain the same result for $k = n$ without any constraint on smooth positive function f . Finally, we also give a counterexample for the two anisotropic expanding flows when $\alpha > 1 - k\beta$.

1. INTRODUCTION

Flows of convex hypersurfaces in \mathbb{R}^{n+1} by a class of speed functions which are homogeneous and symmetric in principal curvatures have been studied by many authors. Firey [15] first introduced the Gauss curvature flow as a model for the shape change of tumbling stones. In [23] Huisken considered the mean curvature flow. He showed that the flow has a unique smooth solution and the hypersurface converges to a round sphere if the initial hypersurface is closed and convex. Later, a range of flows with the speed of homogeneous of degree one in principal curvatures were established, see [12, 13, 3] etc. for example. For the curvature flow at the speed of α -power of the Gauss-Kronecker curvature, it was conjectured that the solution will converge to a round point along the flow for $\alpha > \frac{1}{n+2}$. Chow [12], Andrews [4], Choi and Daskalopoulos [9] gave the partial answers respectively. In [8], Brendle et al. finally resolved the conjecture for all $\alpha > \frac{1}{n+2}$ in all dimensions recently.

2010 *Mathematics Subject Classification.* 35K96, 53C44.

Key words and phrases. expanding flow, anisotropic flow, asymptotic behaviour.

The authors were supported by NSFC, grant no. 11571304.

For the problem on the existence of the prescribed polynomial of the principal curvature radii of the hypersurface, Urbas[29], Chow and Tsai[14], Gerhardt [16], Xia [31] studied the convergence for the flow with the speed of $F(\lambda_1, \dots, \lambda_n)$, where F is symmetric polynomial of the principal curvature radii λ_i of the hypersurface. Guan and Ma [18], Hu et al. [21], and Guan-Xia [20] etc. gave the proofs for a class of L_p Christoffel-Minkowski problems. On the other hand, as a nature extension, anisotropic flows usually provide alternative proofs and smooth category approach of the existence of solutions to elliptic PDEs arising in convex body geometry, see [30, 10, 19, 26, 24] etc.. In this paper, we consider two expanding flows of the convex hypersurfaces at the speeds of $u^\alpha \sigma_k^\beta(\lambda_1, \dots, \lambda_n)$ and $f u^\alpha \sigma_k^\beta(\lambda_1, \dots, \lambda_n)$ respectively, where u is the support function, f is a smooth positive function on S^n , $\alpha, \beta \in \mathbb{R}^1$, $\beta > \frac{1}{k}$ and $\sigma_k(\lambda_1, \dots, \lambda_n)$ is the k -th symmetric polynomial of the principal curvature radii of the hypersurface, k is an integer and $1 \leq k \leq n$. Generally, for the flow with the speed of high powers of curvatures, it is required that the initial hypersurface is uniformly convex and satisfies a suitable pinching conditions, so as to preserve the uniform convexity and converge to a sphere ([1],[6]). Here we prove the same result without any pinching conditions for the flow at the speed $u^\alpha \sigma_k^\beta$. Moreover, for $\alpha \leq 1 - k\beta$, and $\beta > \frac{1}{k}$ we prove that the solution to the flow which is moving at the speed $f u^\alpha \sigma_k^\beta$ exists for all time and converges smoothly after normalisation to a soliton which is a solution of $f u^{\alpha-1} \sigma_k^\beta = c$ if f is a smooth positive function on S^n and satisfies the condition that $(\nabla_i \nabla_j f^{\frac{1}{1+k\beta-\alpha}} + \delta_{ij} f^{\frac{1}{1+k\beta-\alpha}})$ is positive definite. We have the same result for $k = n$ without any constraint on positive smooth function f , which recovers the result of Chou and Wang [11]. When $\beta = 1$, the flow has been studied by Ivaki recently [24]. In this case the self-similar solution of the flow $\partial_t u = f u^\alpha \sigma_k^\beta$ is the solution to $f u^{\alpha-1} \sigma_k = c$ which is just the L_p Christoffel-Minkowski problem for $p = 2 - \alpha$.

Let M_0 be a smooth, closed and uniformly convex hypersurface in \mathbb{R}^{n+1} , and M_0 encloses the origin. We study the following anisotropic expanding curvature flow

$$(1.1) \quad \begin{cases} \frac{\partial X}{\partial t}(x, t) &= f(\nu) \langle X, \nu \rangle^\alpha \sigma_k^\beta(x, t) \nu, \\ X(x, 0) &= X_0(x), \end{cases}$$

where σ_k is the k -th elementary symmetric function for principal curvature radii, i.e

$$\sigma_k(\cdot, t) = \sum_{i_1 < \dots < i_k} \lambda_{i_1} \dots \lambda_{i_k},$$

λ_i is the principal curvature radii of hypersurface M_t , parametrized by $X(\cdot, t) : S^n \rightarrow \mathbb{R}^{n+1}$, and ν is the unit outer normal at $X(\cdot, t)$.

In this paper, we prove the following

Theorem 1.1. *Let M_0 be a smooth, closed and uniformly convex hypersurface in \mathbb{R}^{n+1} , $n \geq 2$, enclosing the origin. If $f \equiv 1$, $\alpha \in \mathbb{R}$, $\beta > \frac{1}{k}$ and $\alpha \leq 1 - k\beta$, k is an integer*

and $1 \leq k \leq n$. Then the flow (1.1) has a unique smooth and uniformly convex solution M_t for all time $t > 0$. After a proper rescaling $X \rightarrow \phi^{-1}(t)X$, where

$$(1.2) \quad \begin{cases} \phi(t) = e^{\gamma t}, & \text{if } \alpha = 1 - k\beta, \\ \phi(t) = (1 + (1 - k\beta - \alpha)\gamma t)^{\frac{1}{1-k\beta-\alpha}}, & \text{if } \alpha \neq 1 - k\beta. \end{cases}$$

and

$$\gamma = \sigma_k^\beta(1, \dots, 1)$$

the hypersurface $\widetilde{M}_t = \phi^{-1}M_t$ converges exponentially to a sphere centered at the origin in the C^∞ topology.

We denote by $u = \langle X, \nu \rangle$ the support function of M_t at X . When $f \equiv 1$, by a direct calculation, we obtain from the flow (1.1) that the support function u satisfies

$$(1.3) \quad \begin{cases} \frac{\partial u}{\partial t}(x, t) = u^\alpha \sigma_k^\beta, \\ u(., 0) = u_0. \end{cases}$$

Rescaling the hypersurface M_t in the way of Theorem 1.1, employing a new parameter τ , we get the normalised flow

$$(1.4) \quad \begin{cases} \frac{\partial u}{\partial \tau}(x, t) = u^\alpha \sigma_k^\beta - \gamma u, \\ u(., 0) = u_0. \end{cases}$$

where

$$\begin{aligned} \gamma &= \sigma_k^\beta(1, \dots, 1), \\ \tau &= \begin{cases} t & \text{if } \alpha = 1 - k\beta, \\ \frac{\log[(1 - \alpha - \beta k)\gamma t + 1]}{1 - \alpha - \beta k} & \text{if } \alpha \neq 1 - k\beta, \end{cases} \end{aligned}$$

For general function f , under the flow (1.1), the support function the support function u satisfies

$$(1.5) \quad \begin{cases} \frac{\partial u}{\partial t}(x, t) = f u^\alpha \sigma_k^\beta, \\ u(., 0) = u_0. \end{cases}$$

We define

$$\widetilde{u} = \left(\frac{|S^n|}{V_{k+1}(\underbrace{u, u, \dots, u}_{(k+1)\text{-times}})} \right)^{\frac{1}{k+1}} u,$$

where the definition of $V_{k+1}(u, u, \dots, u)$ see Section 2. Considering the normalised flow of (1.5) given by

$$(1.6) \quad \begin{cases} \partial_\tau u &= f u^\alpha \sigma_k^\beta - u \frac{\int_{S^n} f u^\alpha \sigma_k^{1+\beta} d\mu_{S^n}}{|S^n|}, \\ u(., 0) &= u_0. \end{cases}$$

where

$$\tau = \int_0^t \left(\frac{|S^n|}{V_{k+1}(u, u, \dots, u)} \right)^{\frac{1-k\beta-\alpha}{k+1}} ds.$$

In (1.6), we still use u instead of \tilde{u} . Consider the following functional which was introduced by Andrews in [5]

$$(1.7) \quad \mathcal{J}(u) = \int_{S^n} u \sigma_k (f u^{\alpha-1} \sigma_k^\beta)^{-\frac{1}{\beta}} d\mu.$$

When $\alpha \geq 1 - k\beta$, we will prove in Lemma 2.3 that $\mathcal{J}(u)$ is strictly decreasing along the flow(1.6) unless u solves the elliptic equation

$$(1.8) \quad f u^{\alpha-1} \sigma_k^\beta = c,$$

where c is a positive constant.

Theorem 1.2. *Let M_0 be a smooth, closed and uniformly convex hypersurface in R^{n+1} , $n \geq 2$, enclosing the origin. Suppose $\alpha, \beta \in R^1$, $\beta > \frac{1}{k}$ and $\alpha \leq 1 - k\beta$, k is an integer and $1 \leq k < n$, f is a smooth positive function on S^n and $\nabla_i \nabla_j f^{\frac{1}{1+k\beta-\alpha}} + \delta_{ij} f^{\frac{1}{1+k\beta-\alpha}}$ is positive definite. Then the flow (1.1) has a unique smooth and uniformly convex solution M_t for all time $t > 0$. After normalisation, the rescaled hypersurfaces \widetilde{M}_t converge smoothly to a smooth solution of (1.8), which is a minimiser of the functional (1.7).*

For $k = n$, we obtain the same result without constraint on f as follows.

Theorem 1.3. *Let M_0 be a smooth, closed and uniformly convex hypersurface in R^{n+1} , $n \geq 2$, enclosing the origin. Suppose $\alpha \in R^1$, $\beta > \frac{1}{n}$ and $\alpha \leq 1 - n\beta$. Then for any f smooth positive function on S^n , the flow (1.1) has a unique smooth and uniformly convex solution M_t for all time $t > 0$. After normalisation, the rescaled hypersurfaces \widetilde{M}_t converge smoothly to a smooth solution of (1.8), which is a minimiser of the functional (1.7).*

Remark 1.1. *When $\beta = 1$, our second flow (1.1) is just the one that Ivaki has studied recently [24]. In [24], Ivaki employed the functional in [20] to prove that the flow $\partial_t u = f u^\alpha \sigma_k$ has a unique smooth solution, and the rescaled flow converges smoothly to a homothetic self-similar solution which is a solution $f u^{\alpha-1} \sigma_k = c$ for $k < n$, $\alpha \leq 1 - k$ and positive function f satisfies that $(\nabla_i \nabla_j f^{\frac{1}{1+k-\alpha}} + \delta_{ij} f^{\frac{1}{1+k-\alpha}})$ is positive definite. When $k = n$, he obtained the same result without imposing any condition on f . It has been obtained by Chou and Wang in [11]. Before Ivaki [24], Hu et al. [21] proved the existence result for the L_p Christoffel Minkowski problem for $p = 2 - \alpha$: $f u^{\alpha-1} \sigma_k = 1$, $\alpha \leq 1 - k$, $1 \leq k \leq n-1$ for any positive function f satisfying $(\nabla_i \nabla_j f^{\frac{1}{1+k-\alpha}} + \delta_{ij} f^{\frac{1}{1+k-\alpha}}) \geq 0$. Under*

the condition that f is positive even function and satisfies $(\nabla_i \nabla_j f^{\frac{1}{1+k-\alpha}} + \delta_{ij} f^{\frac{1}{1+k-\alpha}}) \geq 0$, Guan and Xia in [20] obtained the existence result for $1 - k < \alpha < 1$.

We still denote by $u = \langle X, \nu \rangle$ the support function of M_t at X . We show the condition $\alpha \leq 1 - k\beta$ is necessary. In fact, by use of the method of [26, 27], we show

Theorem 1.4. *Suppose $\alpha > 1 - k\beta$, $\alpha, \beta \in \mathbb{R}$ and $\beta > 0$, k is an integer and $1 \leq k \leq n$. There exist a smooth, closed, uniformly convex hypersurface M_0 , such that under the flow (1.1),*

$$(1.9) \quad R(X(\cdot, t)) := \frac{\max_{S^n} u(\cdot, t)}{\min_{S^n} u(\cdot, t)} \rightarrow \infty, \quad \text{as } t \rightarrow T$$

for some $T > 0$.

This paper is organised as follows. In Section 2, we recall some properties of convex hypersurfaces and show that the functional (1.7) is strictly decreasing along the normalised flow (1.6) unless u satisfies the elliptic equation (1.8). In Section 3, we establish the a priori estimates, which ensure the long time existence of the normalized flows. In Section 4, we show that the flow (1.1) converge to the unit sphere (i.e. Theorem 1.1) and complete the proofs of Theorem 1.2 and Theorem 1.3. Finally, in Section 5, we prove Theorem 1.4.

2. PRELIMINARY

We recall some basic notations at first. Let M be a smooth, closed, uniformly convex hypersurface in R^{n+1} . Assume that M is parametrized by the inverse Gauss map $X : S^n \rightarrow M \subset R^{n+1}$ and encloses origin. The support function $u : S^n \rightarrow R^1$ of M is defined by

$$u(x) = \sup_{y \in M} \langle x, y \rangle.$$

The supremum is attained at a point $y = X(x)$, x is the outer normal of M at y . Hence

$$u(x) = \langle x, X(x) \rangle.$$

Let e_1, \dots, e_n be a smooth local orthonormal frame field on S^n , and ∇ the covariant derivative with respect to the standard metric e_{ij} on S^n . Denote by g_{ij} , g^{ij} , h_{ij} the metric, the inverse of the metric and the second fundamental form of M , respectively. Then the second fundamental form of M is given by (see e. g. [29])

$$h_{ij} = \nabla_i \nabla_j u + u e_{ij},$$

and h_{ij} is symmetric and satisfies the Codazzi equation

$$\nabla_i h_{jk} = \nabla_j h_{ik}.$$

By the Gauss-Weingarten formula

$$\nabla_i x = h_{jk} g^{kl} \nabla_l X,$$

we get

$$e_{ij} = \langle \nabla_i x, \nabla_j x \rangle = h_{ik} g^{kl} h_{jm} g^{ms} \langle \nabla_l X, \nabla_s X \rangle = g^{kl} h_{ik} h_{jl}.$$

Since M is uniformly convex, h_{ij} is invertible. Hence the principal curvature radii are the eigenvalues of the matrix

$$b_{ij} = h^{ik} g_{jk} = \nabla_{ij}^2 u + u \delta_{ij}.$$

Let $\sigma_k(A)$ be the k -th elementary symmetric function defined on the set \mathcal{M}_n of $n \times n$ matrices and $\sigma_k(A_1, \dots, A_k)$ be the complete polarization of σ_k for $A_i \in \mathcal{M}_n$, $i = 1, \dots, k$, i.e.

$$\sigma_k(A_1, \dots, A_k) = \frac{1}{k!} \sum_{i_1, \dots, i_k=1; j_1, \dots, j_k=1}^n \delta_{j_1, \dots, j_k}^{i_1, \dots, i_k} (A_1)_{i_1 j_1} \cdots (A_k)_{i_k j_k}.$$

Let Γ_k be Garding's cone

$$\Gamma_k = \{A \in \mathcal{M}_n : \sigma_i(A) > 0, i = 1, \dots, k\}.$$

For a function $u \in C^2(S^n)$, we denote by W_u the matrix

$$W_u := \nabla^2 u + uI.$$

In the case W_u is positive definite, the eigenvalue of W_u is the principal radii of a strictly convex hypersurface with support function u . Let $u^i \in C^2(S^n)$, $i = 1, \dots, n+1$. Set

$$V(u^1, u^2, \dots, u^{n+1}) := \int_{S^n} u^1 \sigma_n[W_{u^2}, \dots, W_{u^{n+1}}] d\mu,$$

$$V_{k+1}(u^1, u^2, \dots, u^{k+1}) := V(u^1, u^2, \dots, u^{k+1}, 1, \dots, 1).$$

Therefore we define the $k+1$ -th volume by

$$V_{k+1}(\underbrace{u, \dots, u}_{(k+1)\text{-times}}) := \int_{S^n} u \sigma_k[W_u, \dots, W_u] d\mu.$$

Next, we state the well-known Alexandrov-Fenchel inequality.

Lemma 2.1. ([28]) *Let $u^i \in C^2(S^n)$, $i = 1, 2, \dots, k$ be such that $u^i > 0$ and $W_{u^i} \in \Gamma_k$ for $i = 1, 2, \dots, k$. Then for any $v \in C^2(S^n)$, the Alexandrov-Fenchel inequality holds:*

$$V_{k+1}(v, u^1, \dots, u^k)^2 \geq V_{k+1}(v, v, u^2, \dots, u^k) V_{k+1}(u^1, u^1, u^2, \dots, u^k),$$

the equality holds if and only if $v = au^1 + \sum_{l=1}^{n+1} a_l x_l$ for some constants a, a_1, \dots, a_{n+1} .

We consider the flow (1.6). For convenience we still use t instead of τ to denote the time variable if no confusions arise, and we set

$$\rho = fu^{\alpha-1}\sigma_k^\beta, \quad \sigma[f] = \sigma_k[W_f, W_u, \dots, W_u]$$

$$\mathcal{Z}_p(u) = \int_{S^n} u\sigma_k(fu^{\alpha-1}\sigma_k^\beta)^p d\mu = \int_{S^n} u\sigma_k\rho^p d\mu, \quad \eta(t) = \frac{\int_{S^n} fu^\alpha\sigma_k^{1+\beta}d\mu}{|S^n|} = \frac{\mathcal{Z}_1}{|S^n|},$$

where $p \in R^1$. We mention the fact that $\int_{S^n} u\sigma_k d\mu = |S^n|$ here which comes from the scaling of \tilde{u} . Hence (1.6) can be written as $\frac{\partial u}{\partial t} = \rho u - u\frac{\mathcal{Z}_1}{|S^n|}$. By a direct calculation, we have

$$\begin{aligned} & \partial_t \mathcal{Z}_p(u) \\ &= \int_{S^n} (\rho u - u\frac{\mathcal{Z}_1}{|S^n|})\sigma_k\rho^p d\mu + \int_{S^n} ku\sigma[\rho u - u\frac{\mathcal{Z}_1}{|S^n|}]\rho^p d\mu \\ & \quad + \int_{S^n} p\rho^{p-1}u\sigma_k\left((\alpha-1)fu^{\alpha-2}(\rho u - u\frac{\mathcal{Z}_1}{|S^n|})\sigma_k^\beta + k\beta fu^{\alpha-1}\sigma_k^{\beta-1}\sigma[\rho u - u\frac{\mathcal{Z}_1}{|S^n|}]\right) d\mu \\ &= \mathcal{Z}_{1+p} - \frac{\mathcal{Z}_1\mathcal{Z}_p}{|S^n|} + k \int_{S^n} u\sigma[\rho u]\rho^p d\mu - k\frac{\mathcal{Z}_1\mathcal{Z}_p}{|S^n|} \\ & \quad + p(\alpha-1)\mathcal{Z}_{1+p} - p(\alpha-1)\frac{\mathcal{Z}_1\mathcal{Z}_p}{|S^n|} + k\beta p \int_{S^n} u\sigma[\rho u]\rho^p d\mu - k\beta p\frac{\mathcal{Z}_1\mathcal{Z}_p}{|S^n|} \\ &= (1+p(\alpha-1))\left(\mathcal{Z}_{1+p} - \frac{\mathcal{Z}_1\mathcal{Z}_p}{|S^n|}\right) + k(1+\beta p)\left(\int_{S^n} u\rho^p\sigma[\rho u]d\mu - \frac{\mathcal{Z}_1\mathcal{Z}_p}{|S^n|}\right). \end{aligned}$$

Lemma 2.2. *Suppose $\alpha \leq 1 - k\beta$, $\beta > \frac{1}{k}$, $\eta(t)$ is uniformly bounded.*

Proof. Since $\alpha \leq 1 - k\beta$, $\beta > \frac{1}{k}$, let $p = 1$, we have

$$\partial_t \mathcal{Z}_1 = (1 + (\alpha - 1))\left(\mathcal{Z}_2 - \frac{\mathcal{Z}_1\mathcal{Z}_1}{|S^n|}\right) + k(1 + \beta)\left(\int_{S^n} u\rho\sigma[\rho u]d\mu - \frac{\mathcal{Z}_1\mathcal{Z}_1}{|S^n|}\right).$$

Since h_{ij} satisfies the Codazzi equations, we have $\sum_i \nabla_i \sigma^{ij} = 0$ (see [2],[5]), and

$$\begin{aligned} \int_{S^n} u\rho\sigma_k[W_{\rho u}, W_u, \dots, W_u]d\mu &= \int_{S^n} u\rho\sigma[\rho u]d\mu \\ &= \int_{S^n} u\rho\sigma^{ij}(\nabla_i \nabla_j (u\rho) + \delta_{ij}u\rho)d\mu \\ &= \int_{S^n} u\rho\sigma^{ij}(h_{ij}\rho + 2\nabla_i u \nabla_j \rho + u\nabla_i \nabla_j \rho)d\mu \\ &= \mathcal{Z}_2 - \int_{S^n} u^2\sigma^{ij}\nabla_i \rho \nabla_j \rho d\mu. \end{aligned}$$

By the Alexandrov-Fenchel inequality in Lemma 2.1, we have

$$\begin{aligned} \left(\int_{S^n} u\psi\sigma[u]d\mu\right)^2 &\geq \int_{S^n} u\sigma[u]d\mu \int_{S^n} u\psi\sigma[u\psi]d\mu \\ (2.1) \quad &= |S^n|\left(\mathcal{Z}_2 - \int_{S^n} u^2\sigma^{ij}\nabla_i \psi \nabla_j \psi d\mu\right). \end{aligned}$$

Set $\psi = \rho$, the above inequality shows

$$\int_{S^n} u^2 \sigma^{ij} \nabla_i \rho \nabla_j \rho d\mu - \mathcal{Z}_2 + \frac{\mathcal{Z}_1^2}{|S^n|} \geq 0.$$

Hence

$$\begin{aligned} \partial_t \mathcal{Z}_1 &= \alpha \left(\mathcal{Z}_2 - \frac{\mathcal{Z}_1^2}{|S^n|} \right) - k(1 + \beta) \left(\int_{S^n} u^2 \sigma^{ij} \nabla_i \rho \nabla_j \rho d\mu - \mathcal{Z}_2 + \frac{\mathcal{Z}_1^2}{|S^n|} \right) \\ &\leq 0, \end{aligned}$$

where $\alpha \leq 1 - k\beta < 0$ and the Hölder inequality shows that $\mathcal{Z}_2 \geq \frac{\mathcal{Z}_1^2}{|S^n|}$.

$$\partial_t \eta = \frac{\partial_t \mathcal{Z}_1}{|S^n|} = \frac{1}{|S^n|} \left(\alpha \left(\mathcal{Z}_2 - \frac{\mathcal{Z}_1^2}{|S^n|} \right) - k(1 + \beta) \left(\int_{S^n} u^2 \sigma^{ij} \nabla_i \rho \nabla_j \rho d\mu - \mathcal{Z}_2 + \frac{\mathcal{Z}_1^2}{|S^n|} \right) \right) \leq 0.$$

Hence we obtain the uniform upper bound on $\eta(t)$. Next we prove the uniform lower bound. Set $\theta \leq -\frac{1}{\beta} < 0$, $\alpha \leq 1 - k\beta$, we have

$$\begin{aligned} \partial_t \mathcal{Z}_\theta(u) &= (1 + (\alpha - 1)\theta) \left(\mathcal{Z}_{1+\theta} - \frac{\mathcal{Z}_1 \mathcal{Z}_\theta}{|S^n|} \right) + k(1 + \theta\beta) \left(\int_{S^n} u \rho^\theta \sigma[\rho u] d\mu - \frac{\mathcal{Z}_1 \mathcal{Z}_\theta}{|S^n|} \right) \\ &= [1 + (\alpha - 1)\theta] \left(\mathcal{Z}_{\theta+1} - \frac{\mathcal{Z}_1 \mathcal{Z}_\theta}{|S^n|} \right) \\ &\quad + k(1 + \theta\beta) \left(\mathcal{Z}_{1+\theta} - \frac{4\theta}{(1 + \theta)^2} \int_{S^n} u^2 \sigma^{ij} \nabla_i (\rho^{\frac{1+\theta}{2}}) \nabla_j (\rho^{\frac{1+\theta}{2}}) - \frac{\mathcal{Z}_1 \mathcal{Z}_\theta}{|S^n|} \right). \end{aligned}$$

Set $\psi = \rho^{\frac{1+\theta}{2}}$ in the Alexandrov-Fenchel inequality (2.1), we obtain

$$\int_{S^n} u^2 \sigma^{ij} \nabla_i \rho^{\frac{1+\theta}{2}} \nabla_j \rho^{\frac{1+\theta}{2}} d\mu - \mathcal{Z}_{1+\theta} + \frac{\mathcal{Z}_{\frac{1+\theta}{2}}^2}{|S^n|} \geq 0.$$

Thus

$$\begin{aligned} \partial_t \mathcal{Z}_\theta(u) &= [1 + (\alpha - 1)\theta + k(1 + \theta\beta)] \left(\mathcal{Z}_{\theta+1} - \frac{\mathcal{Z}_1 \mathcal{Z}_\theta}{|S^n|} \right) - \frac{4\theta k(1 + \theta\beta)}{(1 + \theta)^2} \left(\mathcal{Z}_{1+\theta} - \frac{\mathcal{Z}_{\frac{1+\theta}{2}}^2}{|S^n|} \right) \\ &\quad - \frac{4\theta k(1 + \theta\beta)}{(1 + \theta)^2} \left(\int_{S^n} u^2 \sigma^{ij} \nabla_i \rho^{\frac{1+\theta}{2}} \nabla_j \rho^{\frac{1+\theta}{2}} d\mu - \mathcal{Z}_{1+\theta} + \frac{\mathcal{Z}_{\frac{1+\theta}{2}}^2}{|S^n|} \right) \\ &\leq 0 \end{aligned}$$

since $\theta < -\frac{1}{\beta}$, and by the Hölder inequality, we get $\mathcal{Z}_{1+\theta} \leq \frac{\mathcal{Z}_1 \mathcal{Z}_\theta}{|S^n|}$ and $\mathcal{Z}_{1+\theta} \geq \frac{\mathcal{Z}_{\frac{1+\theta}{2}}^2}{|S^n|}$. Hence, $\mathcal{Z}_\theta(u) \leq \mathcal{Z}_\theta(u_0)$. By the Hölder inequality again, we have

$$|S^n| = \int_{S^n} u \sigma_k d\mu \leq \left(\int_{S^n} f u^\alpha \sigma_k^{1+\beta} d\mu \right)^{\frac{-\theta}{1-\theta}} \left(\int_{S^n} u \sigma_k (f u^{\alpha-1} \sigma_k^\beta)^\theta d\mu \right)^{\frac{1}{1-\theta}} = \mathcal{Z}_1^{\frac{-\theta}{1-\theta}} \mathcal{Z}_\theta^{\frac{1}{1-\theta}}.$$

Therefore we get the uniform bound on $\eta(t)$. \square

Lemma 2.3. *The functional (1.7) is non-increasing along the normalised flow (1.6), and the equality holds if and only if M_t satisfies the elliptic equation (1.8).*

Proof. Since $\alpha \leq 1 - k\beta$, $\beta > \frac{1}{k}$, from the above calculation process, when $p = -\frac{1}{\beta}$, we have along the normalised flow (1.6)

$$\begin{aligned} \partial_t \mathcal{J}(u) = \partial_t \mathcal{Z}_{-\frac{1}{\beta}}(u) &= \frac{1 - \alpha + \beta}{\beta} \left(\mathcal{Z}_{1-\frac{1}{\beta}} - \frac{\mathcal{Z}_1 \mathcal{Z}_{-\frac{1}{\beta}}}{|S^n|} \right) \\ &\leq 0. \end{aligned}$$

The last inequality holds from the Hölder inequality, and the equality holds if and only if $f u^{\alpha-1} \sigma_k^\beta = c$, where c is a constant. \square

3. A PRIORI ESTIMATES

We first show the C^0 -estimate of the solution to (1.4).

Lemma 3.1. *Let $u(\cdot, t)$, $t \in [0, T)$, be a smooth, uniformly convex solution to (1.4). If $\alpha \leq 1 - k\beta$ and $\beta > \frac{1}{k}$, then there is a positive constant C depending only on α , β and the lower and upper bounds of $u(\cdot, 0)$ such that*

$$\frac{1}{C} \leq u(\cdot, t) \leq C.$$

Proof. Let $u_{\min}(t) = \min_{x \in S^n} u(\cdot, t) = u(x_t, t)$. For fixed time t , at the point x_t , we have

$$\nabla_i u = 0 \quad \text{and} \quad \nabla_{ij} u \geq 0,$$

then

$$\frac{du_{\min}}{dt} \geq \gamma u_{\min} (u_{\min}^{\alpha+k\beta-1} - 1).$$

Hence $u_{\min} \geq \min\{1, u_{\min}(0)\}$. Similarly, we have $u_{\max} \leq \max\{1, u_{\max}(0)\}$. \square

Lemma 3.2. *Let $n \geq 2$, $1 \leq k \leq n$, $\alpha \leq 1 - k\beta$, $\beta > \frac{1}{k}$, and $X(\cdot, t)$ be the solution to the normalised flow (1.6) which encloses the origin for $t \in [0, T)$. Then there is a positive constant C depending on the initial hypersurface and f , α, β , such that*

$$C^{-1} \leq u^{\alpha-1} \sigma_k^\beta \leq C.$$

Proof. Consider the auxiliary function

$$Q = f u^{\alpha-1} \sigma_k^\beta.$$

Since

$$(f u^\alpha \sigma_k^\beta)_{ij} = Q_{ij} u + Q_i u_j + Q_j u_i + Q u_{ij},$$

we get

$$\begin{aligned}
\partial_t Q &= (\alpha - 1)f^2 u^{2\alpha-2} \sigma_k^{2\beta} - \eta(\alpha - 1 + k\beta) f u^{\alpha-1} \sigma_k^\beta \\
&\quad + \beta f^2 u^{2\alpha-1} \sigma_k^{2\beta-1} \sum_i^n \sigma_k^{ii} + \beta f u^{\alpha-1} u^{\alpha-2} \sigma_k^{\beta-1} \sigma_k^{ij} (f u^\alpha \sigma_k^\beta)_{ij} \\
(3.1) &= (\alpha + k\beta - 1)Q^2 - \eta(\alpha + k\beta - 1)Q + \beta f u^\alpha \sigma_k^{\beta-1} \sigma_k^{ij} Q_{ij} + 2\beta f u^{\alpha-1} \sigma_k^{\beta-1} \sigma_k^{ij} Q_i u_j.
\end{aligned}$$

If $\alpha + k\beta - 1 \leq 0$, the sign of the coefficient of the highest order term Q^2 is negative. The sign of the coefficient of the lower order term Q is positive. So it is easy to see $C^{-1} \leq Q \leq C$, where C is the positive constant depending on $f, \alpha, \beta, \min_{S^n \times [0, T]} u$ and $\max_{S^n \times [0, T]} u$. \square

When $f = 1$, by use of (3.1), we can get $C^{-1} \leq Q \leq C$ for the flow (1.4). Then by Lemma 3.1, we have $C^{-1} \leq \sigma_k \leq C$. That is,

Corollary 3.3. *Let $n \geq 2$, $1 \leq k \leq n$, $\alpha \leq 1 - k\beta$, $\beta > \frac{1}{k}$ and $X(., t)$ be the solution to the normalised flow (1.4) which encloses the origin for $t \in [0, T)$. Then there is a positive constant C depending on the initial hypersurface and α, β , such that*

$$C^{-1} \leq \sigma_k \leq C.$$

Lemma 3.4. *Let $n \geq 2$, $1 \leq k \leq n$, $\alpha \leq 1 - k\beta$, $\beta > \frac{1}{k}$ and $X(., t)$ be the solution to the normalised flow (1.6) which encloses the origin for $t \in [0, T)$. Then there is a positive constant C depending on the initial hypersurface and f, α, β , such that $|\nabla \log u| \leq C$.*

Proof. Let $w = \log u$. Then we have

$$h_{ij} = u_{ij} + u \delta_{ij} = u(w_{ij} + w_i w_j + \delta_{ij})$$

Assume the auxiliary function $|\nabla u|^2 - Au^2 < 0$, for a positive constant $A > 0$ along the flow. Otherwise there is a point (x_{t_0}, t_0) where t_0 is the first time, such that $|\nabla u|^2 - Au^2 = 0$, $A > 0$ is a constant to be decided later. Hence at the point (x_{t_0}, t_0) , $\nabla_i |\nabla w|^2 = 0$. Choosing an orthonormal frame and rotating the the coordinate, such that $w_1 = |\nabla w|$, $w_i = 0$ for $i = 2, \dots, n$, and (w_{ij}) is diagonal. Then we get

$$(a_{ij}) := (w_{ij} + w_i w_j + \delta_{ij}) = \text{diag}(1 + w_1^2, 1 + w_{22}, \dots, 1 + w_{nn}),$$

and

$$\begin{aligned}
0 &\leq \partial_t(|\nabla u|^2 - Au^2) = 2u_i(fu^\alpha)_i\sigma_k^\beta + 2\beta fu^\alpha\sigma_k^{\beta-1}\sigma_k^{mn}\nabla_i h_{mn}u_i - 2Afu^{\alpha+1}\sigma_k^\beta \\
&\leq 2u_i(fu^\alpha)_i\sigma_k^\beta + 4k\beta fu^{\alpha+1}\sigma_k^\beta - 2\beta fu^{\alpha+2}\sigma_k^{\beta-1}\sum_i^n \sigma_k^{ii} - 2\beta fu^\alpha\sigma_k^{\beta-1}\sigma_k^{ij}h_{li}h_{lj} \\
&\quad + 2A(k\beta - 1)fu^{\alpha+1}\sigma_k^\beta - 2A\beta fu^{\alpha+2}\sigma_k^{\beta-1}\sum_i^n \sigma_k^{ii} + 2A\beta fu^\alpha\sigma_k^{\beta-1}\sigma_k^{ij}u_iu_j.
\end{aligned}$$

Substituting $u_i = uw_i$ and $w_1^2 = A$ into the above inequality, and denote $\sigma_k = \sigma_k(a_{ij})$, we have

$$0 \leq A(\alpha + k\beta - 1) + \sqrt{A}\frac{|\nabla f|}{f} + 2k\beta - (A + 1)^2\beta\frac{\sigma_k^{11}}{\sigma_k} - A\beta\frac{\sum_i^n \sigma_k^{ii}}{\sigma_k} + A^2\beta\frac{\sigma_k^{11}}{\sigma_k}$$

Hence we have

Case 1. $\alpha \leq 1 - k\beta$, then $A(1 - k\beta - \alpha) \leq \sqrt{A}\frac{|\nabla f|}{f} + 2k\beta$.

Case 2. $\alpha = 1 - k\beta$, then $AC_0\sigma_k^{-\frac{1}{k}} \leq A\beta\frac{\sum_i^n \sigma_k^{ii}}{\sigma_k} \leq \sqrt{A}\frac{|\nabla f|}{f} + 2k\beta$.

Since $\frac{\sum_i^n \sigma_k^{ii}}{\sigma_k} \geq C(n, k)\sigma_k^{-\frac{1}{k}}$ by the classic Newton-MacLaurin inequality [17], and the fact that $\sigma_k(a_{ij})$ is bounded by Lemma 3.2 for the Case 2. Let A be large enough we then get a contradiction. This completes the proof. \square

When $f = 1$, by use of the same argument in Lemma 3.4 and the result of Lemma 3.1. We have

Corollary 3.5. *Let $n \geq 2$, $1 \leq k \leq n$, $\alpha \leq 1 - k\beta$, $\beta > \frac{1}{k}$ and $X(\cdot, t)$ be the solution to the normalised flow (1.4) which encloses the origin for $t \in [0, T)$. Then there is a positive constant C depending on the initial hypersurface and α, β , such that $|\nabla u| \leq C$.*

Lemma 3.6. *Let $u(\cdot, t)$, $t \in [0, T)$, be a smooth, uniformly convex solution to (1.6). If $1 \leq k \leq n$, $\alpha \leq 1 - k\beta$, $\beta > \frac{1}{k}$, then there is a positive constant C_1 and C_2 depending only on f, α, β and the lower and upper bounds of $u(\cdot, 0)$ such that*

$$\frac{1}{C_1} \leq u(\cdot, t) \leq C_1, \quad \frac{1}{C_2} \leq \sigma_k \leq C_2.$$

Proof. Since for the normalised flow (1.6), $\int_{S^n} u\sigma_k d\mu = |S^n|$ is constant. From Lemma 3.2, there is a positive constant C , such that $C^{-1} \leq u^{\alpha-1}\sigma_k^\beta \leq C$. Hence we have

$$C^{-\frac{1}{\beta}}u_{\min}^{\frac{1-\alpha+\beta}{\beta}}(t) \leq \frac{\int_{S^n} u\sigma_k d\mu}{|S^n|} \leq C^{\frac{1}{\beta}}u_{\max}^{\frac{1-\alpha+\beta}{\beta}}(t)$$

Hence we obtain the uniform lower and upper bounds on u from Lemma 3.4. Then by Lemma 3.2, we get the uniform lower and upper bounds on σ_k . \square

Now we are going to estimate the upper and lower bounds of the principle curvature radii of the hypersurface M_t . We rewrite the equation (1.6) in the following form

$$\frac{\partial u}{\partial t} = fu^\alpha \sigma_k^\beta(u) - \eta u = \Phi L - \eta u,$$

where $L = F^{\beta k}$, $F = \sigma_k^{\frac{1}{k}}$, $\Phi = fu^\alpha$, and $h_{ij} = u_{ij} + u\delta_{ij}$.

Lemma 3.7. *Let $1 \leq k < n$, $\alpha, \beta \in R^1$, $\beta > \frac{1}{k}$ and $\alpha \leq 1 - k\beta$, $X(\cdot, t)$ be the solution to the normalised flow (1.6) for $t \in [0, T)$, which encloses the origin. Assume f is a smooth positive function on S^n and $(\nabla_i \nabla_j f^{\frac{1}{1+k\beta-\alpha}} + \delta_{ij} f^{\frac{1}{1+k\beta-\alpha}})$ is positive definite. Then there is a constant C depending only on f , α, β , $\min_{S^n \times [0, T)} u$ and $\max_{S^n \times [0, T)} u$, such that the principal curvature radii of $X(\cdot, t)$ are bounded from above and below*

$$\frac{1}{C} \leq \lambda_i(\cdot, t) \leq C$$

for all $t \in [0, T)$ and $i = 1, \dots, n$.

Proof. Suppose the maximum eigenvalue of the matrix $[\frac{h^{ij}}{u}]$ at time t is attained at the point x_t with unit eigenvector $\xi_t \in T_{x_t} S^n$. By a rotating the frame e_1, \dots, e_n at x_t , assume that at x_t we have $\xi_t = e_1$. At (x_t, t) , we have

$$\begin{aligned} \partial_t h^{11} &= \Phi L'_{kl} \nabla_k \nabla_l h^{11} - \Phi (h^{11})^2 (L + L'_{kl} h_{kl}) + \Phi (\text{tr} L') h^{11} + \eta h^{11} \\ &\quad - \Phi (h^{11})^2 (2L'_{km} h^{nl} + L''_{kl, mn}) \nabla_1 h_{kl} \nabla_1 h_{mn} - (h^{11})^2 (2\nabla_1 \Phi \nabla_1 L + L \nabla_1 \nabla_1 \Phi). \end{aligned}$$

Since $F = \sigma_k^{\frac{1}{k}}$ is concave and homogeneous of degree one, from [29]

$$(3.2) \quad F_{ij} h_{ij} = F,$$

$$(3.3) \quad (2F_{km} h^{nl} + F_{kl, mn}) \nabla_1 h_{kl} \nabla_1 h_{mn} \geq 2F^{-1} F_{kl} F_{mn} \nabla_1 h_{kl} \nabla_1 h_{mn},$$

we have

$$\begin{aligned} \partial_t h^{11} &= \Phi k\beta F^{k\beta-1} F_{kl} \nabla_k \nabla_l h^{11} - \Phi (h^{11})^2 (F^{k\beta} + k\beta F^{k\beta-1} F_{kl} h_{kl}) + \Phi k\beta F^{k\beta-1} \sum_i^n F_{ii} h^{11} \\ &\quad - \Phi (h^{11})^2 \left(2k\beta F^{k\beta-1} F_{km} h^{nl} + k\beta(k\beta - 1) F^{k\beta-2} F_{mn} F_{kl} \right. \\ &\quad \left. + k\beta F^{k\beta-1} F_{kl, mn} \right) \nabla_1 h_{kl} \nabla_1 h_{mn} - (h^{11})^2 (2\nabla_1 \Phi k\beta F^{k\beta-1} \nabla_1 F + F^{k\beta} \nabla_1 \nabla_1 \Phi) + \eta h^{11} \\ &\leq \Phi k\beta F^{k\beta-1} F_{kl} \nabla_k \nabla_l h^{11} - \Phi (h^{11})^2 (1 + k\beta) F^{k\beta} + \Phi k\beta F^{k\beta-1} \sum_i^n F_{ii} h^{11} + \eta h^{11} \\ &\quad - \Phi (h^{11})^2 k\beta(k\beta + 1) F^{k\beta-2} (\nabla_1 F)^2 - (h^{11})^2 (2k\beta F^{k\beta-1} \nabla_1 F \nabla_1 \Phi + F^{k\beta} \nabla_1 \nabla_1 \Phi). \end{aligned}$$

Since

$$2k\beta F^{k\beta-1} \nabla_1 F \nabla_1 \Phi \leq \Phi k\beta(k\beta+1) F^{k\beta} \left(\frac{\nabla_1 F}{F}\right)^2 + \frac{k\beta}{k\beta+1} F^{k\beta} \frac{(\nabla_1 \Phi)^2}{\Phi},$$

we have

$$\begin{aligned} \partial_t h^{11} &\leq \Phi k\beta F^{k\beta-1} F_{kl} \nabla_k \nabla_l h^{11} - \Phi F^{k\beta} (1+k\beta) (h^{11})^2 + \Phi k\beta F^{k\beta-1} \sum_i^n F_{ii} h^{11} + \eta h^{11} \\ &\quad + \frac{k\beta}{k\beta+1} F^{k\beta} \frac{(\nabla_1 \Phi)^2}{\Phi} (h^{11})^2 - F^{k\beta} \nabla_1 \nabla_1 \Phi (h^{11})^2. \end{aligned}$$

Since $\nabla_i \frac{h^{11}}{u} = 0$ and $\nabla_{ij} \frac{h^{11}}{u} \leq 0$,

$$\begin{aligned} \partial_t \frac{h^{11}}{u} &\leq -\Phi(1+k\beta) u F^{k\beta} \left(\frac{h^{11}}{u}\right)^2 + \frac{k\beta}{k\beta+1} u F^{k\beta} \frac{(\nabla_1 \Phi)^2}{\Phi} \left(\frac{h^{11}}{u}\right)^2 \\ &\quad - u F^{k\beta} \nabla_1 \nabla_1 \Phi \left(\frac{h^{11}}{u}\right)^2 + (k\beta-1) u^{-1} F^{k\beta} \Phi \frac{h^{11}}{u} + 2\eta \frac{h^{11}}{u} \\ &\leq u \Phi F^{k\beta} \left(\frac{h^{11}}{u}\right)^2 \left(\frac{k\beta}{1+k\beta} \left(\frac{f_1}{f}\right)^2 - \frac{f_{11}}{f} - \frac{2\alpha}{1+k\beta} \frac{u_1 f_1}{u f} + \frac{\alpha(1+k\beta-\alpha)}{1+k\beta} \left(\frac{u_1}{u}\right)^2 \right) \\ &\quad - \Phi(1+k\beta-\alpha) u F^{k\beta} \left(\frac{h^{11}}{u}\right)^2 + (k\beta-1-\alpha) u^{-1} F^{k\beta} \Phi \frac{h^{11}}{u} + 2\eta \frac{h^{11}}{u} \end{aligned}$$

Since $(f^{\frac{1}{1+k\beta-\alpha}})_{11} + f^{\frac{1}{1+k\beta-\alpha}} > 0$, substituting it into the above inequality, we have

$$\begin{aligned} \partial_t \frac{h^{11}}{u} &\leq u \Phi F^{k\beta} \left(\frac{h^{11}}{u}\right)^2 \left(-\frac{f_{11}}{f} + \frac{k\beta-\alpha}{1+k\beta-\alpha} \left(\frac{f_1}{f}\right)^2 - (1+k\beta-\alpha) \right. \\ &\quad \left. + \alpha \left(\sqrt{\frac{1+k\beta-\alpha}{1+k\beta}} \frac{u_1}{u} - \sqrt{\frac{1}{(1+k\beta)(1+k\beta-\alpha)}} \frac{f_1}{f} \right)^2 \right) \\ &\quad + (k\beta-1-\alpha) u^{-1} F^{k\beta} \Phi \frac{h^{11}}{u} + 2\eta \frac{h^{11}}{u} \end{aligned}$$

For $\beta > \frac{1}{k}$ and $\alpha \leq 1 - k\beta < 0$, we get $k\beta - 1 - \alpha \geq 2(k\beta - 1) > 0$. Hence

$$\partial_t \frac{h^{11}}{u} \leq -C_1 \left(\frac{h^{11}}{u}\right)^2 + C_2 \frac{h^{11}}{u},$$

that is, $h^{11} \leq C$, where C depends on the initial hypersurface, the minimum eigenvalue of $(\nabla_i \nabla_j f^{\frac{1}{1+k\beta-\alpha}} + \delta_{ij} f^{\frac{1}{1+k\beta-\alpha}})$, f , α , and β . Now together with Lemma 3.6, we get

$$(3.4) \quad C^{-1} \leq \lambda_i \leq C.$$

We therefore complete the proof. \square

Lemma 3.8. *Let $1 \leq k \leq n$, $\alpha, \beta \in R^1$, $\beta > \frac{1}{k}$ and $\alpha \leq 1 - k\beta$. Let $X(\cdot, t)$ be the solution to the normalised flow (1.4) for $t \in [0, T)$, which encloses the origin. Then*

there is a constant C depending only on the initial hypersurface and α, β , such that the principal curvature radii of $X(\cdot, t)$ are bounded from above and below

$$\frac{1}{C} \leq \lambda_i(\cdot, t) \leq C$$

for all $t \in [0, T)$ and $i = 1, \dots, n$.

Proof. We prove the lemma just as Lemma 3.7. Suppose the maximum eigenvalue of the matrix $[h^{ij}]$ at time t is attained at the point x_t with unit eigenvector $\xi_t \in T_{x_t}S^n$. By rotating the frame e_1, \dots, e_n at x_t , assume that at x_t we have $\xi_t = e_1$. At (x_t, t) , we have

$$\begin{aligned} \partial_t h^{11} &= \Phi k \beta F^{k\beta-1} F_{kl} \nabla_k \nabla_l h^{11} - \Phi (h^{11})^2 (F^{k\beta} + k\beta F^{k\beta-1} F_{kl} h_{kl}) + \Phi k \beta F^{k\beta-1} \sum_i^n F_{ii} h^{11} \\ &\quad - \Phi (h^{11})^2 \left(2k\beta F^{k\beta-1} F_{km} h^{nl} + k\beta(k\beta - 1) F^{k\beta-2} F_{mn} F_{kl} \right. \\ &\quad \left. + k\beta F^{k\beta-1} F_{kl, mn} \right) \nabla_1 h_{kl} \nabla_1 h_{mn} \\ &\quad - (h^{11})^2 (2\nabla_1 \Phi k \beta F^{k\beta-1} \nabla_1 F + F^{k\beta} \nabla_1 \nabla_1 \Phi) + \gamma h^{11} \\ &\leq -\Phi (h^{11})^2 F^{k\beta} + \gamma h^{11} - \Phi (h^{11})^2 k\beta(k\beta + 1) F^{k\beta-2} (\nabla_1 F)^2 \\ &\quad - (h^{11})^2 (2k\beta F^{k\beta-1} \nabla_1 F \nabla_1 \Phi + F^{k\beta} \nabla_1 \nabla_1 \Phi). \end{aligned}$$

Since

$$2k\beta F^{k\beta-1} \nabla_1 F \nabla_1 \Phi \leq \Phi k\beta(k\beta + 1) F^{k\beta} \left(\frac{\nabla_1 F}{F} \right)^2 + \frac{k\beta}{k\beta + 1} F^{k\beta} \frac{(\nabla_1 \Phi)^2}{\Phi},$$

we have

$$\begin{aligned} \partial_t h^{11} &\leq (\alpha - 1) u^\alpha \sigma_k^\beta (h^{11})^2 + (\gamma - \alpha u^{\alpha-1} \sigma_k^\beta) h^{11} \\ (3.5) \quad &\quad + \frac{\alpha(1 + k\beta - \alpha)}{k\beta + 1} u^{\alpha-2} (\nabla_1 u)^2 \sigma_k^\beta (h^{11})^2. \end{aligned}$$

Since $\beta > \frac{1}{k}$, $\alpha \leq 1 - k\beta < 0$, by (3.5) we have

$$\partial_t h^{11} \leq -C_0 (h^{11})^2 + C_1 h^{11}$$

where C_0 and C_1 are two positive constants which depend only on the initial hypersurface and α, β . Therefore we have

$$h^{11} \leq \max\{C_3, \max h^{11}(0)\}$$

where C_3 is also a positive constants depending only on the initial hypersurface and α, β . Now together with Lemma 3.3, we get

$$(3.6) \quad C^{-1} \leq \lambda_i \leq C.$$

□

Now we show Lemma 3.7 holds for any positive smooth function f when $k = n$, $\beta > \frac{1}{n}$ and $\alpha \leq 1 - n\beta$.

Lemma 3.9. *Let $n \geq 2$, $k = n$. If $\alpha \leq 1 - n\beta$, $\beta > \frac{1}{n}$ and $X(\cdot, t)$ be the solution to the normalised flow (1.6) which encloses the origin for $t \in [0, T)$. Then there is a constant C depending only on the initial hypersurface and f , α , β , such that the principal curvature radii of $X(\cdot, t)$ are bounded from above and below*

$$C^{-1} \leq \lambda_i(\cdot, t) \leq C$$

for all $t \in [0, T)$ and $i = 1, \dots, n$.

Proof. Consider the auxiliary function

$$w(\xi, t, \tau) = \log h^{\tau\tau} - \varepsilon \log u + \frac{M}{2}(u^2 + |\nabla u|^2),$$

where τ is a unit vector in the tangential space of S^n , while ε and M are large constants to be decided. Assume w achieve its maximum at (x_0, t_0) in the direction $\tau = (1, 0, \dots, 0)$. By a coordinate rotation, h_{ij} and h^{ij} are diagonal at this point. Then at the point (x_0, t_0) .

$$w = \log h^{11} - \varepsilon \log u + \frac{M}{2}(u^2 + |\nabla u|^2),$$

$$(3.7) \quad 0 = \nabla_i w = -h^{11} \nabla_i h_{11} - \varepsilon \frac{\nabla_i u}{u} + M u u_i + M \nabla_k u \nabla_{ki} u,$$

$$(3.8) \quad \begin{aligned} 0 \geq \nabla_{ij} w &= -h^{11} \nabla_{ij} h_{11} + 2h^{11} h^{kk} \nabla_1 h_{ik} \nabla_1 h_{kj} - (h^{11})^2 \nabla_i h_{11} \nabla_j h_{11} - \varepsilon \frac{\nabla_{ij} u}{u} \\ &+ \varepsilon \frac{\nabla_i u \nabla_j u}{u^2} + M u_i u_j + M u u_{ij} + M \nabla_{ki} u \nabla_{kj} u + M \nabla_k u \nabla_{kij} u \end{aligned}$$

Set $\Phi = f u^\alpha$, we have

$$\begin{aligned} 0 \leq \partial_t w &= -h^{11} \partial_t h_{11} - \varepsilon \frac{\partial_t u}{u} + M u \nabla_t u + M \nabla_k u \nabla_{kt} u \\ &= -h^{11} \left(\Phi_{11} \sigma_n^\beta + 2\beta \sigma_n^{\beta-1} \nabla_1 \Phi \nabla_1 \sigma_n + \beta(\beta-1) \Phi \sigma_n^{\beta-2} (\nabla_1 \sigma_n)^2 \right. \\ &\quad \left. + \beta \Phi \sigma_n^{\beta-1} \nabla_{11} \sigma_n - \eta h_{11} + \Phi \sigma_n^\beta \right) - \varepsilon \frac{\Phi \sigma_n^\beta - \eta u}{u} \\ &\quad + M u (\Phi \sigma_n^\beta - \eta u) + M u_k (\Phi \sigma_n^\beta - \eta u)_k \\ &= -h^{11} \left(\Phi_{11} \sigma_n^\beta + 2\beta \sigma_n^\beta \nabla_1 \Phi h^{ij} \nabla_1 h_{ij} + \beta(\beta-1) \Phi \sigma_n^\beta (h^{ij} \nabla_1 h_{ij})^2 \right. \\ &\quad \left. + \beta \Phi \sigma_n^\beta (h^{ij} \nabla_{ij} h_{11} + n - h_{11} \sum_i h^{ii} - h^{ii} h^{jj} (\nabla_1 h_{ij})^2 + (h^{ij} \nabla_1 h_{ij})^2) \right. \\ &\quad \left. - \eta h_{11} + \Phi \sigma_n^\beta \right) - \varepsilon \frac{\Phi \sigma_n^\beta - \eta u}{u} + M u (\Phi \sigma_n^\beta - \eta u) + M u_k (\Phi \sigma_n^\beta - \eta u)_k \end{aligned}$$

By (3.8) and multiplying $\Phi^{-1}\sigma_n^{-\beta}$ the two sides of the above inequality, we obtain

$$\begin{aligned}
0 &\leq -h^{11}\frac{\nabla_{11}\Phi}{\Phi} + h^{11}\left(\left(\frac{\nabla_1\Phi}{\Phi}\right)^2 + \beta^2(h^{ij}\nabla_1h_{ij})^2\right) - h^{11}\beta^2(h^{ij}\nabla_1h_{ij})^2 \\
&\quad + \beta\left(-2h^{11}h^{ij}h^{kk}\nabla_1h_{ik}\nabla_1h_{jk} + (h^{11})^2h^{ij}\nabla_ih_{11}\nabla_jh_{11} + n\varepsilon u^{-1}\right. \\
&\quad \left.- \varepsilon\sum_i h^{ii} - \varepsilon h^{ij}u_iu_ju^{-2} - Mh^{ij}h_{ik}h_{jk} + nMu - Mh^{ij}u_k\nabla_kh_{ij}\right) \\
&\quad - h^{11}(1+n\beta) + \beta\sum_i h^{ii} + \beta h^{11}h^{ii}h^{jj}(\nabla_1h_{ij})^2 + Mu + Mu_k\frac{\nabla_k\Phi}{\Phi} \\
&\quad + \beta Mh^{ij}u_k\nabla_kh_{ij} - \eta M\frac{u^2 + |\nabla u|^2}{\Phi\sigma_n^\beta} - \varepsilon u^{-1} + \eta\frac{\varepsilon + 1}{\Phi\sigma_n^\beta} \\
&\leq -2\beta h^{11}h^{ij}h^{kk}\nabla_1h_{ik}\nabla_1h_{kj} + \beta(h^{11})^2h^{ij}\nabla_ih_{11}\nabla_jh_{11} + \beta h^{11}h^{ii}h^{jj}(\nabla_1h_{ij})^2 \\
&\quad + \beta\frac{n\varepsilon}{u} - \beta\varepsilon\sum_i h^{ii} + \beta\sum_i h^{ii} + Mu + h^{11}\left(\frac{\nabla_1\Phi}{\Phi}\right)^2 \\
&\quad + M\beta nu - h^{11}\frac{\nabla_1\nabla_1\Phi}{\Phi} + Mu_k\frac{\nabla_k\Phi}{\Phi} - \eta\frac{Mr^2 - \varepsilon - 1}{\Phi\sigma_n^\beta} \\
&\leq C_0 - \beta(\varepsilon - 1)h^{11} + h^{11}\left(\frac{\nabla_1\Phi}{\Phi}\right)^2 - h^{11}\frac{\nabla_1\nabla_1\Phi}{\Phi} + Mu_k\frac{\nabla_k\Phi}{\Phi} - \eta\frac{Mr^2 - \varepsilon - 1}{\Phi\sigma_n^\beta},
\end{aligned}$$

where we use the Cauchy inequality $2\beta\frac{\nabla_1\Phi}{\Phi}h^{ij}\nabla_1h_{ij} \leq \left(\frac{\nabla_1\Phi}{\Phi}\right)^2 + \beta^2(h^{ij}\nabla_1h_{ij})^2$ for the second term.

Since $\nabla_k\Phi = \nabla_k(fu^\alpha)$, we obtain

$$\begin{aligned}
&h^{11}\left(\frac{\nabla_1\Phi}{\Phi}\right)^2 - h^{11}\frac{\nabla_1\nabla_1\Phi}{\Phi} + Mu_k\frac{\nabla_k\Phi}{\Phi} \\
&= h^{11}\left(\frac{f_1}{f} + \alpha\frac{u_1}{u}\right)^2 - h^{11}\left(\frac{\nabla_1\nabla_1f}{f} + 2\alpha\frac{\nabla_1f}{f}\frac{\nabla_1u}{u} + \alpha(\alpha - 1)\left(\frac{\nabla_1u}{u}\right)^2\right) \\
&\quad + M\left(\nabla_ku\frac{\nabla_kf}{f} + \alpha u^{-1}|\nabla u|^2\right) \\
&\leq Ch^{11} + CM
\end{aligned}$$

Choosing $M \geq \frac{\varepsilon+1}{\min_{r^*2}}$, the inequality becomes

$$0 \leq C_0 - \beta(\varepsilon - 1)h^{11} + Ch^{11} + CM.$$

By choosing ε large to get

$$0 \leq C_1 - C_2h^{11}.$$

That is, $h^{11} \leq C_3$, where C_3 is a constant depending only on f , α , β , $\min_{S^n \times [0,T]} u$ and $\max_{S^n \times [0,T]} u$. Hence the principal radii $\lambda_i \geq C_2, i = 1, \dots, n$. From Lemma 3.6, we

know $\sigma_n = \lambda_1 \cdots \lambda_n \leq C_1$. Therefore we get the C^2 estimate $C^{-1} \leq \lambda_i \leq C, i = 1, \dots, n$ for the solutions to the normalised flow (1.6). \square

From the estimates obtained in Lemmata 3.8, 3.7 and 3.9, we know that the equations (1.4) and (1.6) are uniformly parabolic. By the C^0 estimates (Lemmata 3.1 and 3.6), the gradient estimates (Lemma 3.5 and Corollary 3.4) and the C^2 estimates Lemmata 3.8, 3.7 and 3.9, and the Krylov's theory [25], we get the Hölder continuity of $\nabla^2 u$ and u_t . Then we can get higher order derivation estimates by the regularity theory of the uniformly parabolic equations. Therefore we get the long time existence and the uniqueness of the smooth solution to the normalized flows (1.4) and (1.6), respectively.

4. PROOF OF THEOREM 1.1, THEOREM 1.2 AND THEOREM 1.3

In this section we give the proof of Theorems 1.1 at first. In order to prove the convergence of the normalized flow (1.4), we require the following better gradient estimate.

Lemma 4.1. *Let $u(\cdot, t)$ be a smooth uniformly convex solution to the flow (1.4). If $\alpha \leq 1 - k\beta$, then there exist positive constants C and C_0 , depending only on the initial hypersurface and α, β , such that*

$$\max_{S^n} \frac{|\nabla u(\cdot, t)|}{u(\cdot, t)} \leq C e^{-C_0 t}$$

for all $t > 0$.

Proof. Let $w = \log u$. Then we have

$$h_{ij} = u_{ij} + u\delta_{ij} = e^w(w_{ij} + w_i w_j + \delta_{ij})$$

and

$$w_t = (e^w)^{\alpha+k\beta-1} \sigma_k^\beta([w_{ij} + w_i w_j + \delta_{ij}]) - \gamma.$$

Consider the auxiliary function

$$Q = \frac{1}{2} |\nabla w|^2.$$

At the point where Q attains its spatial maximum, we have

$$0 = \nabla_i Q = \sum w_l w_{li},$$

$$0 \geq \nabla_{ij} Q = \sum w_{li} w_{lj} + \sum w_l w_{lij},$$

and

$$\begin{aligned}
\partial_t Q &= \sum w_l w_{lt} \\
&= 2Q(\alpha + k\beta - 1)(e^w)^{\alpha+k\beta-1} \sigma_k^\beta \\
&\quad + \beta(e^w)^{\alpha+k\beta-1} \sigma_k^{\beta-1} \sigma_k^{ij} (w_l w_{ijl} + w_l w_{il} w_j + w_l w_i w_{jl}) \\
&= 2Q(\alpha + k\beta - 1)(e^w)^{\alpha+k\beta-1} \sigma_k^\beta + \beta(e^w)^{\alpha+k\beta-1} \sigma_k^{\beta-1} \sigma_k^{ij} w_l w_{ijl}.
\end{aligned}$$

By the Ricci identity,

$$\nabla_l w_{ij} = \nabla_j w_{li} + \delta_{il} w_j - \delta_{ij} w_l,$$

we get

$$\begin{aligned}
\partial_t Q &= 2(\alpha + k\beta - 1)(e^w)^{\alpha+k\beta-1} \sigma_k^\beta Q \\
&\quad + \beta(e^w)^{\alpha+k\beta-1} \sigma_k^{\beta-1} \sigma_k^{ij} \left(\sum_l w_l w_{lij} + w_i w_j - \delta_{ij} |\nabla w|^2 \right) \\
&\leq 2(\alpha + k\beta - 1)(e^w)^{\alpha+k\beta-1} \sigma_k^\beta Q + 2\beta(e^w)^{\alpha+k\beta-1} \sigma_k^{\beta-1} (\max_i \sigma_k^{ii} - \sum_i \sigma_k^{ii}) Q \\
&\leq -C_0 Q.
\end{aligned}$$

We therefore have $Q \leq C e^{-C_0 t}$, where C_0 and C are two positive constants which depends only on α , β and the geometry of M_0 . \square

Proof of Theorem 1.1. Case (i). $\alpha < 1 - k\beta$. Let $u(\cdot, t)$ be the solution. By rescaling M_0 if necessary, we assume

$$a := \min_{S^n} u(\cdot, 0) \leq 1 \leq \max_{S^n} u(\cdot, 0) =: b.$$

We also introduce two time-dependent functions

$$\begin{aligned}
u_1 &= [1 - (1 - a^{-q})e^{q\gamma t}]^{-\frac{1}{q}}, \\
u_2 &= [1 - (1 - b^{-q})e^{q\gamma t}]^{-\frac{1}{q}},
\end{aligned}$$

where $q = \alpha + k\beta - 1 < 0$. Both functions u_1 and u_2 are the solutions of (1.4), hence by the comparison principle, $u_1(t) \leq u(\cdot, t) \leq u_2(t)$. That is

$$(a^{-q} - 1)e^{q\gamma t} \leq u^{-q} - 1 \leq (b^{-q} - 1)e^{q\gamma t}.$$

Thus u converges to 1 exponentially. By the interpolation and the a priori estimates Lemmas 3.1 and 3.8, we see that $\|u(\cdot, t) - 1\|_{C^k(S^n)} \rightarrow 0$ exponentially. Hence M_t converges to the unit sphere centered at the origin.

Case (ii). $\alpha = 1 - k\beta$. By Lemma 4.1, we have that $\|\nabla u\| \rightarrow 0$ exponentially as $t \rightarrow \infty$. Hence by the interpolation and the a priori estimates, we can get that u converges exponentially to a constant in the C^∞ topology as $t \rightarrow \infty$. \square

Proofs of Theorems 1.2 and 1.3 . Recall Lemma 2.3. To complete the proof of Theorems 1.2 and 1.3, it suffices to show that the solution of (1.8) is unique.

Case1: $\alpha < 1 - k\beta$. Let u_1, u_2 be two smooth solutions of (1.8), i.e.

$$f u_1^{\alpha-1} \sigma_k^\beta(\nabla^2 u_1 + u_1 I) = c, f u_2^{\alpha-1} \sigma_k^\beta(\nabla^2 u_2 + u_2 I) = c.$$

Suppose $M = \frac{u_1}{u_2}$ attains its maximum at $X_0 \in S^n$, then at x_0 ,

$$\begin{aligned} 0 &= \nabla \log M = \frac{\nabla u_1}{u_1} - \frac{\nabla u_2}{u_2}, \\ 0 &\geq \nabla^2 \log M = \frac{\nabla^2 u_1}{u_1} - \frac{\nabla^2 u_2}{u_2}. \end{aligned}$$

Hence at x_0 , we get

$$1 = \frac{u_1^{\alpha-1} \sigma_k^\beta(\nabla^2 u_1 + u_1 I)}{u_2^{\alpha-1} \sigma_k^\beta(\nabla^2 u_2 + u_2 I)} = \frac{u_1^{\alpha-1+k\beta} \sigma_k^\beta(\frac{\nabla^2 u_1}{u_1} + I)}{u_2^{\alpha-1+k\beta} \sigma_k^\beta(\frac{\nabla^2 u_2}{u_2} + I)} \leq M^{\alpha-1+k\beta}.$$

Since $\alpha < 1 - k\beta$, $M(x_0) = \max_{S^n} M \leq 1$. Similarly one can show $\min_{S^n} M \geq 1$. Therefore $u_1 \equiv u_2$.

Case 2: $\alpha = 1 - k\beta$. We use the same method in [21] to get the solutions for equation(1.8) differ only by dilation. We omit the proof process here.

Hence we complete the proofs of the Theorems 1.2 and 1.3. \square

5. PROOF OF THEOREM 1.4

In this section we give the proof of Theorem 1.4. The methods for the cases that f being any smooth positive function and $f \equiv 1$ are the same , we just consider the case $f \equiv 1$. The calculation for the example of the flow (1.1) is similar to one in [26, 27], we give the brief proof.

By a simple calculation, we obtain the following expressions of the metric, the unit normal, the second fundamental forms and the support function of the hypersurface, according to the radial function r

$$(5.1) \quad g_{ij} = r^2 \delta_{ij} + r_i r_j,$$

$$(5.2) \quad n = \frac{r z - \nabla r}{\sqrt{r^2 + |\nabla r|^2}},$$

$$(5.3) \quad \tilde{h}_{ij} = \frac{-r r_{ij} + 2r_i r_j + r^2 \delta_{ij}}{\sqrt{r^2 + |\nabla r|^2}},$$

$$(5.4) \quad u = \frac{r^2}{\sqrt{r^2 + |\nabla r|^2}}.$$

For the convex body K and the hypersurface $M = \partial K$, consider the dual body K^* , by [22] we know $r^* = \frac{1}{u}$, and

$$(5.5) \quad K = \frac{\det \tilde{h}_{ij}}{\det g_{ij}}, \quad \frac{1}{\sigma_n^*} = \frac{\det e_{ij}}{\det \tilde{h}_{ij}^*}, \quad \frac{\det e_{ij}}{\det g_{ij}} = \frac{1}{r^{2n-2}(r^2 + |\nabla r|^2)}.$$

Since $r^* = \frac{1}{u}$,

$$\frac{\partial r^*}{\partial t} = -r^{*2-\alpha} \sigma_k^\beta ([r^{*-1} \delta_{ij} + 2r^{*-3} r_i^* r_j^* - r^{*-2} r_{ij}^*]).$$

Generally, we consider

$$(5.6) \quad \begin{cases} \frac{\partial r}{\partial t}(x, t) &= -r^{2-\alpha} \sigma_k^\beta ([r^{-1} \delta_{ij} + 2r^{-3} r_i r_j - r^{-2} r_{ij}]) (x, t) \\ r(\cdot, 0) &= r_0 \end{cases}$$

If $\alpha + k\beta - 1 > 0$, that is $2 - \alpha < k\beta + 1$. (1.9) is equivalent to

$$(5.7) \quad R(X(\cdot, t)) = \frac{\max_{S^n} r(\cdot, t)}{\min_{S^n} r(\cdot, t)} \rightarrow \infty,$$

as $t \rightarrow T$ for some $T > 0$. It is suffice to show that $\min_{S^n} r \rightarrow 0$ in finite time while $\max_{S^n} r$ remains positive. By the comparison principle, it suffices to construct a sub-solution $Y(\cdot, t)$

$$(5.8) \quad \begin{cases} \frac{\partial w}{\partial t}(x, t) &\geq -w^{2-\alpha} \sigma_k^\beta ([w^{-1} \delta_{ij} + 2w^{-3} w_i w_j - w^{-2} w_{ij}]) (x, t) \\ w(\cdot, 0) &= w_0 \end{cases}$$

such that $\min_{S^n} w(\cdot, t) \rightarrow 0$ but $\max_{S^n} w(\cdot, t)$ remains positive, as $t \rightarrow T$ for some finite time $T > 0$.

Lemma 5.1. *There is a sub-solution $Y(\cdot, t)$, where $t \in (-1, 0)$, to*

$$(5.9) \quad \begin{cases} \frac{\partial r}{\partial t} &= -ar^{2-\alpha} \sigma_k^\beta ([r^{-1} \delta_{ij} + 2r^{-3} r_i r_j - r^{-2} r_{ij}]) \\ r(\cdot, 0) &= r_0 \end{cases}$$

for a sufficiently large constant $a > 0$, such that $\min_{S^n} r(\cdot, t) \rightarrow 0$ but $\max_{S^n} r(\cdot, t)$ remains positive, as $t \rightarrow 0$.

Proof. The sub-solution is a family of closed convex hypersurfaces $\widehat{M}_t := Y(S^n, t)$. Near the origin, let \widehat{M}_t be the graph of a radial function

$$(5.10) \quad \psi(\rho, t) = \begin{cases} -|t|^\theta + |t|^{-\theta+\mu} \rho^2 & \text{if } \rho < |t|^\theta \\ -|t|^\theta - \frac{1-\mu}{1+\mu} |t|^{\theta(1+\mu)} + \frac{2}{1+\mu} \rho^{1+\mu} & \text{if } |t|^\theta \leq \rho \leq 1 \end{cases}$$

where $q = k\beta + 1 - \widehat{\alpha} > 0$, $\widehat{\alpha} = 2 - \alpha$, $\mu = \frac{q\theta-1}{k\beta\theta}$, and $\theta > \frac{1}{q}$ is a constant. It is easy to verify that ψ is strictly convex, and $\psi \in C^{1,1}(B_1(0))$.

If $Y(S^n, t)$ is a sub-solution for some $\widehat{\alpha}$, it is also a sub-solution for $\alpha' < \widehat{\alpha}$, so we prove the case when $q = k\beta + 1 - \widehat{\alpha}$ is very small. From the equalities (5.1), (5.3), (5.5) and Newton-Maclaurin inequality, by direct computation, we have

(i) if $0 \leq \rho \leq |t|^\theta$, then

$$r^{\widehat{\alpha}} \sigma_k^\beta ([r^{-1} \delta_{ij} + 2r^{-3} r_i r_j - r^{-2} r_{ij}]) \geq c |t|^{\theta \widehat{\alpha}} |t|^{k\beta(\mu-1)} = c |t|^{\theta-1},$$

$$\left| \frac{\partial}{\partial t} Y(p, t) \right| \leq \theta |t|^{\theta-1},$$

where $p = (x, \psi(|x|, t))$ is a point on the graph of ψ .

(ii) if $|t|^\theta \leq \rho \leq 1$, then

$$r^{\widehat{\alpha}} \sigma_k^\beta ([r^{-1} \delta_{ij} + 2r^{-3} r_i r_j - r^{-2} r_{ij}]) \geq c \rho^{\widehat{\alpha}} \rho^{k\beta(\mu-1)} = c |t|^{\theta-1},$$

$$\left| \frac{\partial}{\partial t} Y(p, t) \right| \leq \theta |t|^{\theta-1}.$$

Extending the graph of ψ to a closed convex hypersurface \widehat{M}_t , such that it is $C^{1,1}$ smooth, uniformly convex, rotationally symmetric. Moreover, assume that the ball $B_1(z)$ is contained in the interior of \widehat{M}_t , for all $t \in (-1, 0)$, where $z = (0, \dots, 0, 10)$ is a point on the x_{n+1} -axis. \square

For a given $\tau \in (-1, 0)$, let $B_1(z) \subset M_0 \subset \widehat{M}_\tau$, let M_t be a solution to the flow (5.9) with initial data M_0 . M_t touches the origin at $t = t_0$, for some $t_0 \in (\tau, 0)$. We assume $\tau \rightarrow 0$ closely.

On the other hand, let $\widetilde{X}(\cdot, t)$ be the solution to

$$\frac{\partial X}{\partial t} = -ba\widetilde{r}^{\widehat{\alpha}} \sigma_k^\beta \nu$$

with initial condition $\widetilde{X}(\cdot, \tau) = \partial B_1(z)$, where $b = 2^{\widehat{\alpha}} \sup\{|p|^{\widehat{\alpha}} : p \in M_t, \tau < t < t_0\}$ and $\widetilde{r} = |X - z|$. We can choose τ so small that the ball $B_{\frac{1}{2}}(z)$ is contained in the interior of $\widetilde{X}(\cdot, t)$ for all $t \in (\tau, t_0)$. By the comparison principle, we know that the ball $B_{\frac{1}{2}}(z)$ is contained in the interior of M_t for all $t \in (\tau, t_0)$. Hence, as $t \rightarrow t_0$, we have $\min r(\cdot, t) \rightarrow 0$ and $\max r(\cdot, t) > |z| = 10$. Hence (5.7) is proved for M_t .

For a large constant $a > 0$. Making the rescaling $\widetilde{M}_t = a^{-\frac{1}{4}} M_t$, \widetilde{M}_t solve the flow (5.6). Hence we complete the proof.

REFERENCES

- [1] Alessandrini, R.; Sinestrari, C.: Evolution of hypersurfaces by powers of the scalar curvature. Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) 9 (2010), no. 3, 541–571.
- [2] Andrews, B.: Entropy estimates for evolving hypersurfaces. Communications in Analysis and Geometry, (1994), 2(1), 267-275.
- [3] Andrews, B.: Gauss curvature flow: the fate of the rolling stones. Invent. Math. 138 (1999), no. 1, 151–161.
- [4] Andrews, B.: Contraction of convex hypersurfaces in Euclidean space. Calc. Var. Partial Differential Equations. 2 (1994), no. 2, 151–171.

- [5] Andrews, B.: Monotone quantities and unique limits for evolving convex hypersurfaces[J]. International Mathematics Research Notices,(1997), 20(20),1001-1031.
- [6] Andrews, B.; McCoy, J.: Convex hypersurfaces with pinched principal curvatures and flow of convex hypersurfaces by high powers of curvature. Trans. Amer. Math. Soc. 364 (2012), no. 7, 3427–3447.
- [7] Andrews, B; McCoy; Zhang, Y.: Contracting convex hypersurfaces by curvature. Calc. Var. Partial Differential Equations 47(2013), 611–665.
- [8] Brendle, S.; Choi, K.; Daskalopoulos, P.: Asymptotic behavior of flows by powers of the Gauss curvature. Acta Math. 219 (2017), no. 1, 1-16.
- [9] Choi, K.; Daskalopoulos, P.: Uniqueness of closed self-similar solutions to the Gauss curvature flow. ArXiv:1609.05487.
- [10] Chou, K.-S.; Wang, X.-J.: A logarithmic Gauss curvature flow and the Minkowski problem. Ann. Inst. H. Poincaré Anal. Non Linéaire 17 (2000), no. 6, 733–751.
- [11] Chou, K.-S.; Wang, X.-J.: The L_p Minkowski problem and the Minkowski problem in centroaffine geometry. Adv. in Math., 205 (2006), 33-83.
- [12] Chow, B.: Deforming convex hypersurfaces by the n -th root of the Gaussian curvature. J. Differential Geom. 22 (1985), no. 1, 117–138.
- [13] Chow, B.: Deforming convex hypersurfaces by the square root of the scalar curvature. Invent. Math. 87 (1987), no. 1, 63–82.
- [14] Chow, B.; D. H. Tsai.: Expansion of convex hypersurfaces by nonhomogeneous functions of curvature. Asian J. Math. 1(1997):769-784.
- [15] Firey, W. J.: Shapes of worn stones. Mathematika. 21 (1974), 1–11.
- [16] Gerhardt, C.: Non-scale-invariant inverse curvature flows in Euclidean space. Car. Var. Partial Differential Equations. 49(2014):471-489.
- [17] Guan, P. F.; Lin, C. S.: On equation $\det(u_{ij} + u\delta_{ij}) = u^p f$ on S^n . Preprint No 2000-7, NCTS in Tsing-Hua University,2000.
- [18] Guan, P. F.; Ma, X. N.: Christoffel-Minkowski problem I: convexity of solutions of a hessian equation. Invent. Math. 151(2003),553-577.
- [19] Guan, P. F.; Ni, L.: Entropy and a convergence theorem for Gauss curvature flow in high dimensions, J. Eur. Math. Soc. (in press).
- [20] Guan, P. F.; Xia, C.: L^p Christoffel-Minkowski problem: the case $1 < p < k + 1$. Cal. Var. Partial Differential Equations.57(2018), no.2,:69.
- [21] Hu, C. Q.; Ma, X. N.; Shen C.L.: On the Christoffel-Minkowski problem of Firey’s p - sum. Cal. Var. Partial Differential Equations.21(2004), no.2,137–155.
- [22] Huang, Y.; Lutwak, E.; Yang, D.; Zhang, G.: Geometric measures in the dual Brunn-Minkowski theory and their associated Minkowski problems. Acta Math. 216 (2016), no. 2, 325–388.
- [23] Huisken, G.: Flow by mean curvature of convex surfaces into spheres. J. Differential Geom. 20 (1984), no. 1, 237–266.
- [24] Ivaki, M.N.: Deforming a hypersurface by principal radii of curvature and support function. arXiv:1803.08470v1(2018).
- [25] Krylov, N.V.: Nonlinear elliptic and parabolic equations of the second order. D.Reidel Publishing Co.,Dordrecht,1987.xiv+462 pp.
- [26] Li, Q.-R.; Sheng, W.M.; Wang, X.-J.: Flow by Gauss curvature to the Aleksandrov and dual Minkowski problems. J. Eur. Math. Soc.(in press).
- [27] Li, Q.-R.; Sheng, W.M.; Wang, X.-J.: A class of anisotropic curvature type flow. Preprint.
- [28] Schneider, R.: Convex bodies: the Brunn-Minkowski theory. Cambridge University Press, Cambridge 1993.
- [29] Urbas, J.: An expansion of convex hypersurfaces. J. Differential Geom. 33 (1991), no. 1, 91–125.

- [30] Wang, X.J.: Existence of convex hypersurfaces with prescribed Gauss-Kronecker curvature. Trans. Amer. Math. Soc. 348 (1996), 4501–4524.
- [31] Xia, C.: Inverse anisotropic curvature flow from convex hypersurfaces. J Geom Anal. 27(2016), no. 3, 1–24.

WEIMIN SHENG: SCHOOL OF MATHEMATICAL SCIENCES, ZHEJIANG UNIVERSITY, HANGZHOU 310027, CHINA.

E-mail address: weimins@zju.edu.cn

CAIHONG YI: SCHOOL OF MATHEMATICAL SCIENCES, ZHEJIANG UNIVERSITY, HANGZHOU 310027, CHINA.

E-mail address: 11735001@zju.edu.cn