

# Global regularity for a class of 3D tropical climate model without thermal diffusion

Jinlu Li<sup>1\*</sup> and Yanghai Yu<sup>2†</sup>

<sup>1</sup>*School of Mathematics and Computer Sciences, Gannan Normal University, Ganzhou 341000, China*

<sup>2</sup>*School of Mathematics and Statistics, Anhui Normal University, Wuhu, Anhui, 241002, China*

May 14, 2019

---

**Abstract:** In this paper, we establish the global regularity for the 3D tropical climate model with fractional dissipation.

**Keywords:** Tropical climate model; Global regularity.

**MSC (2010):** 35D35; 76D03

---

## 1 Introduction

This paper focuses on the following 3D tropical climate model given by

$$\begin{cases} \partial_t u + u \cdot \nabla u + \mu \Lambda^{2\alpha} u + \nabla p + \operatorname{div}(v \otimes v) = 0, & x \in \mathbb{R}^3, t > 0, \\ \partial_t v + u \cdot \nabla v + v \cdot \nabla u + \nabla \theta + \nu \Lambda^{2\beta} v = 0, & x \in \mathbb{R}^3, t > 0, \\ \partial_t \theta + u \cdot \nabla \theta + \eta \Lambda^{2\gamma} \theta + \operatorname{div} v = 0, & x \in \mathbb{R}^3, t \geq 0, \\ (u, v, \theta)|_{t=0} = (u_0, v_0, \theta_0), & x \in \mathbb{R}^3. \end{cases} \quad (1.1)$$

here  $\mu, \nu, \eta, \alpha, \beta, \gamma$  are non-negative parameters,  $u = u(t, x)$  and  $v = v(t, x)$  stand for the barotropic mode and the first baroclinic mode of the vector velocity, respectively.  $\Pi = \Pi(t, x)$  and  $\theta = \theta(t, x)$  denote the scalar pressure and scalar temperature, respectively.  $\Lambda = \sqrt{-\Delta}$  is the Zygmund operator and the fractional power operator  $\Lambda^\delta = (-\Delta)^{\frac{\delta}{2}}$  is defined through Fourier transform as

$$\Lambda^\delta f(\xi) = \mathcal{F}^{-1} |\xi|^\delta \mathcal{F} f(\xi).$$

The original version of (1.1) without any fractional Laplacian terms was derived by Frierson–Majda–Pauluis [4] from the inviscid primitive equations with the aid of performing a Galerkin truncation to the hydrostatic Boussinesq equations, of which the first baroclinic mode had been originally used in some studies of tropical atmosphere. More relevant background on the tropical

---

\*E-mail: lijnlu@gnnu.cn

†E-mail: yuyanghai214@sina.com( Corresponding author)

climate model can be found in [6, 9, 10] and the references therein. From the mathematical point of view, the tropical climate model (1.1) are significant generalizations of the generalized magnetohydrodynamic (GMHD) equations which model the complex interaction between the fluid dynamic phenomena. In fact, when the temperature  $\theta \equiv \text{Constant}$ , (1.1) is reduced to the GMHD equations, namely,

$$\begin{cases} \partial_t u + u \cdot \nabla u + \nabla \Pi + \mu \Lambda^{2\alpha} u - v \cdot \nabla v = 0, & x \in \mathbb{R}^d, t > 0, \\ \partial_t v + u \cdot \nabla v + \nu \Lambda^{2\beta} v - v \cdot \nabla u = 0, & x \in \mathbb{R}^d, t > 0, \\ \operatorname{div} u = \operatorname{div} v = 0, & x \in \mathbb{R}^d, t \geq 0, \\ (u(0, x), v(0, x)) = (u_0(x), v_0(x)), & x \in \mathbb{R}^d, \end{cases} \quad (1.2)$$

here  $u = u(t, x)$  and  $v = v(t, x)$  denote the velocity and the magnetic field of the fluid, respectively. The mathematical studies on the tropical climate model (1.1) have attracted considerable attention recently from various authors and have motivated a large number of research papers. Here, we mainly recall some global regularity results which are more relative with our research in this field. Li–Titi [7] introduced a new quantity to bypass the obstacle caused by the absence of thermal diffusion and proved the global well-posedness of strong solutions for the 2D tropical climate model (1.1) with  $\alpha = \beta = 1$  and  $\mu > 0, \nu > 0, \eta = 0$ . Later, Ye [15] obtained the global regularity of a tropical climate model with the very weak dissipation of the barotropic ( $\alpha > 0, \beta = \gamma = 1$  and  $\mu, \nu, \eta > 0$ ) by the “weakly nonlinear” energy estimates approach and the maximal  $L_t^q L_x^p$  regularity for the heat kernel. Dong et al. [2] established the global regularity results for the 2D system (1.1) without thermal diffusion with  $\alpha + \beta = 2, \beta \in (1, \frac{3}{2}], \mu, \nu > 0$  and  $\alpha = 2, \mu > 0, \nu = \eta = 0$ , respectively. Recently, Dong et al. [3] established the global existence and regularity for the 2D system (1.1) ( $\mu, \nu, \eta > 0, \beta = 1$ ) with the fractional dissipation which are in two very broad ranges, namely,

$$\gamma \geq \frac{4 + \alpha - \sqrt{\alpha^2 + 8\alpha + 8}}{2}, \quad \text{if } 0 < \alpha < \frac{1}{2}; \quad \gamma \geq 1 - \alpha, \quad \text{if } \frac{1}{2} \leq \alpha \leq 1.$$

For the 3D case, Zhu [16] obtained the global regularity for the tropical climate model (1.1) with only fractional diffusion on barotropic mode, namely,  $\alpha \geq \frac{5}{2}, \mu > 0, \nu = \eta = 0$ . Naturally, we ask that how weak dissipation which the  $u$ -equations possess to ensure that the 3D system (1.1) has the global regularity? This is our principal goal of this paper. Our main result reads as follows

**Theorem 1.1** *Consider the system (1.1) with  $\alpha = \beta = \frac{5}{4}$  and  $\gamma = 0$ . Assume that  $(u_0, v_0, \theta_0) \in (H^2(\mathbb{R}^3) \cap \dot{B}_{2,1}^0(\mathbb{R}^3))^3$  and  $\operatorname{div} u_0 = 0$ . Then (1.1) has a unique global solution  $(u, v, \theta)$  satisfying for any  $T > 0$*

$$(u, v, \theta) \in (L^\infty(0, T; H^2))^3 \quad \text{and} \quad (u, v) \in L^2(0, T; H^{\frac{13}{4}}).$$

**Remark 1.1** *As mentioned above, Theorem 1.1 holds true for the 3D GMHD system.*

## 2 Proof of Theorem 1.1

Taking the  $L^2$  inner product of Eqs.(1.1)<sub>1</sub>–(1.1)<sub>3</sub> with  $u$ ,  $v$  and  $\theta$ , respectively, then using the fact  $\operatorname{div} u = 0$ , we have the following basic energy estimate

$$\|u\|_{L^2}^2 + \|v\|_{L^2}^2 + \|\theta\|_{L^2}^2 + 2 \int_0^T (\|\Lambda^{\frac{5}{4}}u\|_{L^2}^2 + \|\Lambda^{\frac{5}{4}}v\|_{L^2}^2) ds = \|u_0\|_{L^2}^2 + \|v_0\|_{L^2}^2 + \|\theta_0\|_{L^2}^2. \quad (2.3)$$

Applying  $\dot{\Delta}_j$  to (1.1)<sub>1</sub> and taking the  $L^2$  inner product of the resulting equation with  $\dot{\Delta}_j u$  yields

$$\frac{1}{2} \frac{d}{dt} \|\dot{\Delta}_j u\|_{L^2}^2 + c 2^{\frac{5}{2}j} \|\dot{\Delta}_j u\|_{L^2} \leq \|\dot{\Delta}_j u_0\|_{L^2} + \|\dot{\Delta}_j \operatorname{div}(u \otimes u)\|_{L^2} + \|\dot{\Delta}_j \operatorname{div}(v \otimes v)\|_{L^2}, \quad (2.4)$$

which implies

$$\|u\|_{L_T^\infty(\dot{B}_{2,1}^0)} + \|u\|_{L_T^1(\dot{B}_{2,1}^{\frac{5}{2}})} \lesssim \|u_0\|_{\dot{B}_{2,1}^0} + \int_0^T (\|u \otimes u\|_{\dot{B}_{2,1}^1} + \|v \otimes v\|_{\dot{B}_{2,1}^1}) ds. \quad (2.5)$$

Invoking the product estimate of two functions in homogeneous spaces (see Corollary 2.55 in [1]):

$$\|fg\|_{\dot{B}_{2,1}^{s+t}} \lesssim \|f\|_{\dot{H}^s} \|g\|_{\dot{H}^t} \quad \text{with} \quad -3/2 < s, t < 3/2 \quad \text{and} \quad s + t > 0 \quad (2.6)$$

Therefore, it follows from (2.3) and (2.5)

$$\|u\|_{L_T^\infty(\dot{B}_{2,1}^0)} + \|u\|_{L_T^1(\dot{B}_{2,1}^{\frac{5}{2}})} \lesssim \|u_0\|_{\dot{B}_{2,1}^0} + \int_0^T (\|u\|_{\dot{H}^{\frac{5}{4}}}^2 + \|v\|_{\dot{H}^{\frac{5}{4}}}^2) ds \leq C. \quad (2.7)$$

Taking the  $L^2$  inner product of Eqs.(1.1)<sub>1</sub>–(1.1)<sub>3</sub> with  $-\Delta u$ ,  $-\Delta v$  and  $-\Delta \theta$ , respectively, then using the fact  $\operatorname{div} u = 0$ , we have

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} (\|\nabla u\|_{L^2}^2 + \|\nabla v\|_{L^2}^2 + \|\nabla \theta\|_{L^2}^2) + \|\Lambda^{\frac{9}{4}}u\|_{L^2}^2 + \|\Lambda^{\frac{9}{4}}v\|_{L^2}^2 \\ &= \underbrace{\int_{\mathbb{R}^3} (u \cdot \nabla u) \cdot \Delta u dx}_{I_1} + \underbrace{\int_{\mathbb{R}^3} \operatorname{div}(v \otimes v) \cdot \Delta u dx}_{I_2} + \underbrace{\int_{\mathbb{R}^3} (v \cdot \nabla u) \cdot \Delta v dx}_{I_3} \\ & \quad + \underbrace{\int_{\mathbb{R}^3} (u \cdot \nabla v) \cdot \Delta v dx}_{I_4} + \underbrace{\int_{\mathbb{R}^3} (u \cdot \nabla \theta) \cdot \Delta \theta dx}_{I_5}, \end{aligned} \quad (2.8)$$

Next, we need to estimate the above five terms one by one.

Notice that the fact  $\operatorname{div}(v \otimes v) = v \cdot \nabla v + v \operatorname{div} v$ , we have

$$\begin{aligned} I_1 &\lesssim \|u\|_{L^6} \|\nabla u\|_{L^{\frac{12}{5}}} \|\Delta u\|_{L^{\frac{12}{5}}} \lesssim \|\nabla u\|_{L^2}^2 \|\Lambda^{\frac{5}{4}}u\|_{L^2}^2 + \varepsilon \|\Lambda^{\frac{9}{4}}u\|_{L^2}^2, \\ I_2 &\lesssim \|v\|_{L^6} \|\nabla v\|_{L^{\frac{12}{5}}} \|\Delta u\|_{L^{\frac{12}{5}}} \lesssim \|\nabla v\|_{L^2}^2 \|\Lambda^{\frac{5}{4}}v\|_{L^2}^2 + \varepsilon \|\Lambda^{\frac{9}{4}}u\|_{L^2}^2, \\ I_3 &\lesssim \|v\|_{L^6} \|\nabla u\|_{L^{\frac{12}{5}}} \|\Delta v\|_{L^{\frac{12}{5}}} \lesssim \|\nabla v\|_{L^2}^2 \|\Lambda^{\frac{5}{4}}u\|_{L^2}^2 + \varepsilon \|\Lambda^{\frac{9}{4}}v\|_{L^2}^2, \\ I_4 &\lesssim \|u\|_{L^6} \|\nabla v\|_{L^{\frac{12}{5}}} \|\Delta v\|_{L^{\frac{12}{5}}} \lesssim \|\nabla u\|_{L^2}^2 \|\Lambda^{\frac{5}{4}}v\|_{L^2}^2 + \varepsilon \|\Lambda^{\frac{9}{4}}v\|_{L^2}^2, \\ I_5 &= \int_{\mathbb{R}^3} \partial_k u^i \partial_i \theta \partial_k \theta dx \lesssim \|\nabla u\|_{L^\infty} \|\nabla \theta\|_{L^2}^2, \end{aligned}$$

where we have used the embedding  $\dot{H}^{\frac{1}{4}}(\mathbb{R}^3) \hookrightarrow L^{\frac{12}{5}}(\mathbb{R}^3)$  and the Young inequality.

Putting the above estimates together with (2.8), we obtain

$$\frac{d}{dt} \|\nabla u, \nabla v, \nabla \theta\|_{L^2}^2 + \|\Lambda^{\frac{9}{4}} u, \Lambda^{\frac{9}{4}} v\|_{L^2}^2 \lesssim \|\nabla u, \nabla v, \nabla \theta\|_{L^2}^2 (\|\Lambda^{\frac{5}{4}} u, \Lambda^{\frac{5}{4}} v\|_{L^2}^2 + \|\nabla u\|_{L^\infty}),$$

combining the embedding  $\dot{B}_{2,1}^{\frac{3}{2}}(\mathbb{R}^3) \hookrightarrow L^\infty(\mathbb{R}^3)$ , which implies

$$\sup_{0 \leq s \leq T} \|\nabla u, \nabla v, \nabla \theta\|_{L^2}^2 + \int_0^T \|\Lambda^{\frac{9}{4}} u\|_{L^2}^2 + \|\Lambda^{\frac{9}{4}} v\|_{L^2}^2 ds \leq C. \quad (2.9)$$

Multiplying (2.4) by  $2^j$  and summing over  $j \in \mathbb{Z}$  yields

$$\|u\|_{L_T^\infty(\dot{B}_{2,1}^1)} + \|u\|_{L_T^1(\dot{B}_{2,1}^{\frac{7}{2}})} \lesssim \|u_0\|_{\dot{B}_{2,1}^1} + \int_0^T (\|u \cdot \nabla u\|_{\dot{B}_{2,1}^1} + \|v \cdot \nabla v\|_{\dot{B}_{2,1}^1} + \|v \operatorname{div} v\|_{\dot{B}_{2,1}^1}) ds. \quad (2.10)$$

Using the product estimate (2.6) again, one has

$$\|u \cdot \nabla u\|_{\dot{B}_{2,1}^1} + \|v \cdot \nabla v\|_{\dot{B}_{2,1}^1} + \|v \operatorname{div} v\|_{\dot{B}_{2,1}^1} \lesssim \|u\|_{\dot{H}^{\frac{5}{4}}} \|u\|_{\dot{H}^{\frac{9}{4}}} + \|v\|_{\dot{H}^{\frac{5}{4}}} \|v\|_{\dot{H}^{\frac{9}{4}}}.$$

Therefore, we have from (2.10)

$$\|u\|_{L_T^\infty(\dot{B}_{2,1}^1)} + \|u\|_{L_T^1(\dot{B}_{2,1}^{\frac{7}{2}})} \lesssim \|u_0\|_{\dot{B}_{2,1}^1} + \int_0^T (\|u\|_{\dot{H}^{\frac{5}{4}}}^2 + \|v\|_{\dot{H}^{\frac{5}{4}}}^2 + \|u\|_{\dot{H}^{\frac{9}{4}}}^2 + \|v\|_{\dot{H}^{\frac{9}{4}}}^2) ds \leq C,$$

which gives

$$\int_0^T \|\nabla^2 u\|_{L^\infty} ds \leq C. \quad (2.11)$$

Applying  $\Delta$  to both sides of (1.1)<sub>1</sub>, (1.1)<sub>2</sub> and (1.1)<sub>3</sub>, taking the  $L^2$  inner product of the resulting equations with  $\Delta u$ ,  $\Delta v$  and  $\Delta \theta$ , respectively, then adding them together yields that

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \|\Delta u, \Delta v, \Delta \theta\|_2^2 + \|\Lambda^{\frac{13}{4}} u\|_2^2 + \|\Lambda^{\frac{13}{4}} v\|_2^2 \\ &= - \underbrace{\int_{\mathbb{R}^3} \Delta(u \cdot \nabla u) \cdot \Delta u dx}_{J_1} - \underbrace{\int_{\mathbb{R}^3} \Delta \operatorname{div}(v \otimes v) \cdot \Delta u dx}_{J_2} - \underbrace{\int_{\mathbb{R}^3} \Delta(u \cdot \nabla v) \cdot \Delta v dx}_{J_3} \\ & \quad - \underbrace{\int_{\mathbb{R}^3} \Delta(v \cdot \nabla u) \cdot \Delta v dx}_{J_4} - \underbrace{\int_{\mathbb{R}^3} \Delta(u \cdot \nabla \theta) \cdot \Delta \theta dx}_{J_5}, \end{aligned} \quad (2.12)$$

where we have used the fact

$$\int_{\mathbb{R}^3} \Delta \nabla \theta \cdot \Delta v dx + \int_{\mathbb{R}^3} \Delta \operatorname{div} v \cdot \Delta \theta dx = 0.$$

Next, we need to estimate the above four terms one by one as follows.

For the first term  $J_1$ , due to the divergence free condition  $\operatorname{div} u = 0$ , one has

$$J_1 = \int_{\mathbb{R}^3} \Delta u \cdot \nabla u \cdot \Delta u dx + \int_{\mathbb{R}^3} \partial_i u \cdot \partial_i \nabla u \cdot \Delta u dx \lesssim \|\nabla u\|_{L^\infty} \|\Delta u\|_{L^2}^2. \quad (2.13)$$

Similarly, we have

$$\begin{aligned} J_3 &= \int_{\mathbb{R}^3} \Delta u \cdot \nabla v \cdot \Delta v dx + \int_{\mathbb{R}^3} \partial_i u \cdot \partial_i \nabla v \cdot \Delta v dx \\ &\lesssim \|\Delta u\|_{L^\infty} \|\nabla v\|_{L^2} \|\Delta v\|_{L^2} + \|\nabla u\|_{L^\infty} \|\Delta v\|_{L^2}^2, \end{aligned} \quad (2.14)$$

$$\begin{aligned} J_5 &= \int_{\mathbb{R}^3} \Delta u \cdot \nabla \theta \cdot \Delta \theta dx + \int_{\mathbb{R}^3} \partial_i u \cdot \partial_i \nabla \theta \cdot \Delta \theta dx \\ &\lesssim \|\Delta u\|_{L^\infty} \|\nabla \theta\|_{L^2} \|\Delta \theta\|_{L^2} + \|\nabla u\|_{L^\infty} \|\Delta \theta\|_{L^2}^2, \end{aligned} \quad (2.15)$$

For the second term  $J_2$ , we decompose it into as

$$J_2 = \underbrace{\int_{\mathbb{R}^3} \Delta(v \cdot \nabla v) \cdot \Delta u dx}_{J_{21}} + \underbrace{\int_{\mathbb{R}^3} \Delta(v \operatorname{div} v) \cdot \Delta u dx}_{J_{22}}. \quad (2.16)$$

We estimate  $J_{21}$  as follows

$$\begin{aligned} J_{21} &\leq \int_{\mathbb{R}^3} \Delta v \cdot \nabla v \cdot \Delta u dx + \int_{\mathbb{R}^3} \partial_i v \cdot \partial_i \nabla v \cdot \Delta u dx + \int_{\mathbb{R}^3} v \cdot \nabla \Delta v \cdot \Delta u dx \\ &\lesssim \|\Delta u\|_{L^\infty} \|\nabla v\|_{L^2} \|\Delta v\|_{L^2} + \|v\|_{L^6} \|\Delta \nabla v\|_{L^{\frac{12}{5}}} \|\Delta u\|_{L^2} \\ &\lesssim \|\Delta u\|_{L^\infty} (\|\nabla v\|_{L^2}^2 + \|\Delta v\|_{L^2}^2) + \|\nabla v\|_{L^2}^2 \|\Delta u\|_{L^2}^2 + \varepsilon \|\Lambda^{\frac{13}{4}} v\|_{L^2}^2, \end{aligned}$$

Similarly, one has

$$J_{22} \lesssim \|\Delta u\|_{L^\infty} (\|\nabla v\|_{L^2}^2 + \|\Delta v\|_{L^2}^2) + \|\nabla v\|_{L^2}^2 \|\Delta u\|_{L^2}^2 + \varepsilon \|\Lambda^{\frac{13}{4}} v\|_{L^2}^2,$$

For the term  $J_4$ , we get

$$\begin{aligned} J_4 &\leq \int_{\mathbb{R}^3} \Delta v \cdot \nabla u \cdot \Delta v dx + \int_{\mathbb{R}^3} \partial_i v \cdot \partial_i \nabla u \cdot \Delta v dx + \int_{\mathbb{R}^3} v \cdot \nabla \Delta v \cdot \Delta v dx \\ &\lesssim \|\nabla u\|_{L^\infty} \|\Delta v\|_{L^2}^2 + \|\nabla^2 u\|_{L^\infty} \|\nabla v\|_{L^2} \|\Delta v\|_{L^2} + \|v\|_{L^6} \|\Delta \nabla v\|_{L^{\frac{12}{5}}} \|\Delta v\|_{L^2} \\ &\lesssim (\|\nabla u\|_{L^\infty} + \|\nabla^2 u\|_{L^\infty}) (\|\nabla v\|_{L^2}^2 + \|\Delta v\|_{L^2}^2) + \|\nabla v\|_{L^2}^2 \|\Delta u\|_{L^2}^2 + \varepsilon \|\Lambda^{\frac{13}{4}} v\|_{L^2}^2, \end{aligned} \quad (2.17)$$

Combining the above estimates with (2.12), one has

$$\frac{d}{dt} \|\Delta u, \Delta v, \Delta \theta\|_{L^2}^2 + \|\Lambda^{\frac{13}{4}} u, \Lambda^{\frac{13}{4}} v\|_{L^2}^2 \leq C \left(1 + \|\nabla u\|_\infty + \|\nabla^2 u\|_{L^\infty}\right) (1 + \|\Delta u, \Delta v, \Delta \theta\|_{L^2}^2),$$

which follows from (2.11) and Gronwall's Lemma that

$$\sup_{0 \leq s \leq T} \|\Delta u, \Delta v, \Delta \theta\|_{L^2}^2 + \int_0^T \|\Lambda^{\frac{13}{4}} u, \Lambda^{\frac{13}{4}} v\|_{L^2}^2 ds \leq C(T).$$

Thus, we complete the proof of Theorem 1.1.

## Acknowledgments

J. Li is supported by NSFC (No.11801090). Y. Yu is supported by NSF of Anhui Province (No.1908085QA05).

## References

- [1] H. Bahouri, J.-Y. Chemin, R. Danchin, *Fourier Analysis and Nonlinear Partial Differential Equations*, Grundlehren Math. Wiss., vol.343, Springer-Verlag, Berlin, Heidelberg, 2011.
- [2] B. Dong, W. Wang, J. Wu, H. Zhang, Global regularity results for the climate model with fractional dissipation, *Discrete Contin. Dyn. Syst. Ser. B* (2018). <https://doi.org/10.3934/dcdsb.2018102>
- [3] B. Dong, J. Wu, Z. Ye, Global regularity for a 2D tropical climate model with fractional dissipation, *J. Nonlinear Sci.*, (2018) <https://doi.org/10.1007/s00332-018-9495-5>
- [4] D. Frierson, A. Majda, O. Pauluis, Large scale dynamics of precipitation fronts in the tropical atmosphere, a novel relaxation limit, *Commun. Math. Sci.*, 2 (2004), 591–626.
- [5] J. Fan, H. Malaikah, S. Monaque, G. Nakamura, Y. Zhou, Global cauchy problem of 2D generalized MHD equations, *Monatsh Math.*, 175 (2014), 127–131.
- [6] A. Gill, Some simple solutions for heat-induced tropical circulation, *Q. J. R. Meteorol. Soc.*, 106 (1980), 447–462.
- [7] J. Li, E. Titi, Global well-posedness of strong solutions to a tropical climate model, *Discrete Contin. Dyn. Syst.*, 36 (2016), 4495–4516.
- [8] P. G. Lemari, *Recent Developments in the Navier–Stokes Problem*, CRC Research Notes in Mathematics, Vol 431, Boca Raton, FL: Chapman & Hall, 2002.
- [9] T. Matsuno, Quasi-geostrophic motions in the equatorial area, *J. Meteorol. Soc. Jpn.*, 44 (1966), 25–42.
- [10] A. Majda, J. Biello, The nonlinear interaction of barotropic and equatorial baroclinic Rossby waves, *J. Atmos. Sci.*, 60(2003), 1809–1821.
- [11] C. Ma, Z. Jiang, R. Wan, Local well-posedness for the tropical climate model with fractional velocity diffusion, *Kinet. Relat. Models*, 99 (2016), 551–570.
- [12] C. Ma, R. Wan, Spectral analysis and global well-posedness for a viscous tropical climate model with only a damp term, *Nonlinear Anal. Real World Appl.*, 39 (2018), 554–567.
- [13] R. Wan, Global small solutions to a tropical climate model without thermal diffusion, *J. Math. Phys.*, 57:2, 021507 (2016).
- [14] X. Wu, Y. Yu, Y. Tang, Global existence and asymptotic behavior for the 3D generalized Hall–MHD system, *Nonlinear Anal.*, 151 (2017), 41–50.
- [15] Z. Ye, Global regularity for a class of 2D tropical climate model, *J. Math. Anal. Appl.*, 446 (2017), 307–321.
- [16] M. Zhu, Global regularity for the tropical climate model with fractional diffusion on barotropic mode, *Appl. Math. Lett.*, 81 (2018), 99–104.