A FIBRATION THEOREM FOR COLLAPSING SEQUENCES OF ALEXANDROV SPACES

TADASHI FUJIOKA

ABSTRACT. Suppose a sequence M_j of Alexandrov spaces collapses to a space X with only weak singularities. Yamaguchi constructed a map $f_j : M_j \to X$ called an almost Lipschitz submersion for large j. We prove that if M_j has a uniform positive lower bound for the volumes of spaces of directions, which is sufficiently large compared to the weakness of singularities of X, then f_j is a locally trivial fibration. Moreover, we show some properties on the intrinsic metric and the volume of the fibers of f_j .

1. INTRODUCTION

Let M_j be a sequence of *n*-dimensional Alexandrov spaces with curvature $\geq \kappa$ and diameter $\leq D$. It is well-known that M_j has a convergent subsequence in the Gromov-Hausdorff distance and that the limit space X is also an Alexandrov space of dimension $\leq n$ and with curvature $\geq \kappa$. The main problem of the convergence theory of Alexandrov spaces is to determine the relation between the topology and geometry of X and those of M_j with large j.

According to Perelman's stability theorem, if dim X = n, then M_j is homeomorphic to X ([8], [5]). The case dim X < n is called a collapse. In this case, Yamaguchi [16] proved that if both M_j and X are Riemannian manifolds, then there exists a locally trivial fibration $f_j: M_j \to X$, which is an almost Riemannian submersion. This result can be generalized to the case when both M_j and X have only weak singularities ([2, 9.13]). Furthermore, Yamaguchi [17] also proved that even if M_j is a general Alexandrov space, if X has only weak singularities, then there exists a map $f_j: M_j \to X$ called an almost Lipschitz submersion, which is a generalization of an almost Riemannian submersion.

To state Yamaguchi's almost Lipschitz submersion theorem, we introduce some notation. We denote by δ a positive number less than some constant depending only on n and κ , and by $\varkappa(\delta)$ a positive function depending only on n and κ such that $\varkappa(\delta) \to 0$ as $\delta \to 0$. For a positive number a, we denote by c(a) a positive constant depending only on n, κ and a, which is much smaller than a.

Theorem 1.1 ([17, 0.2]). Let k < n and let X be a k-dimensional Alexandrov space with curvature $\geq \kappa$ such that every point has a (k, δ) -strainer with length $> \ell$. Let M be an n-dimensional Alexandrov space with curvature $\geq \kappa$ that is μ -close to X in the Gromov-Hausdorff distance. Suppose $\mu < c(\ell\delta^2) \ll \ell\delta^2$. Then, there exists

Date: June 18, 2021.

²⁰¹⁰ Mathematics Subject Classification. 53C20, 53C23.

Key words and phrases. Alexandrov spaces, collapse, fibration, strainers, almost regular maps.

a $\varkappa(\delta)$ -almost Lipschitz submersion $f: M \to X$ in the following sense:

$$\left|\frac{|f(x)f(y)|}{|xy|} - \sin\inf_{z} \angle yxz\right| < \varkappa(\delta)$$

for any $x, y \in M$, where the infimum is taken over all $z \in f^{-1}(f(x))$.

Yamaguchi conjectured that the map f is actually a locally trivial fibration. Rong and Xu [11] showed that it is true if each fiber of f is a topological manifold (without boundary) of codimension k. Xu [14] (cf. [15]) also proved that f is a Hurewicz fibration. Their proofs are based on the construction of neighborhood retractions to the fibers, which can be applied to a wider class of maps called e^{ε} -Lipschitz and co-Lipschitz maps. We prove Yamaguchi's conjecture from a different point of view, in the following case:

Theorem 1.2. Under the conditions of Theorem 1.1, let ε be a lower bound for the volume of the space of directions at each point of M. Suppose in addition $\delta < c(\varepsilon) \ll \varepsilon$. Then, the map f is a locally trivial fibration.

In the case of collapse of codimension one, the additional assumption is always satisfied.

Corollary 1.3. Under the conditions of Theorem 1.1, if k = n - 1, then f is a locally trivial fibration whose fiber is homeomorphic to a circle or a closed interval.

Let us explain how to prove Theorem 1.2. Let $\{(a_i, b_i)\}_{i=1}^k$ be a (k, δ) -strainer at $p \in X$, where $k = \dim X$. Then, the distance map $\varphi = (|a_1 \cdot |, \dots, |a_k \cdot |) : X \to \mathbb{R}^k$ gives a local chart around p. Let $\hat{\varphi} : M \to \mathbb{R}^k$ and $\hat{p} \in M$ be natural lifts of φ and p, respectively. Then, $\varphi^{-1} \circ \hat{\varphi}$ gives a local map between neighborhoods of \hat{p} and p. The global map f in Theorem 1.1 is constructed by gluing such local maps. We prove that f satisfies

(1.1)
$$\left| \left(\varphi \circ f(x) - \varphi \circ f(y) \right) - \left(\hat{\varphi}(x) - \hat{\varphi}(y) \right) \right| < \varkappa(\delta) |xy|$$

for any x, y near \hat{p} . Roughly speaking, the differential of $\varphi \circ f$ is close to that of $\hat{\varphi}$. This inequality allows f to inherit the properties of $\varphi^{-1} \circ \hat{\varphi}$.

In [8], Perelman developed the theory of noncritical maps and proved that a proper noncritical map is a locally trivial fibration (here, the definition of noncriticality includes an assumption on the volume of spaces of directions). In particular, the above $\hat{\varphi}$ is a locally trivial fibration near \hat{p} under the additional assumption of Theorem 1.2. We slightly modify the definition of noncritical maps in terms of the inequality (1.1) and show that $\varphi \circ f$ is also a locally trivial fibration. From this point of view, the general case of Yamaguchi's conjecture is reduced to the problem of proving Perelman's fibration theorem for noncritical maps without the assumption on the volume of spaces of directions (see also Remark 5.16).

Next, we discuss the fibers of the map f in Theorem 1.1 (not in Theorem 1.2). Fibers of almost regular maps such as $\hat{\varphi}$ were studied in [2, §11–12]. The inequality (1.1) enables us to apply the arguments there to $\varphi \circ f$. It is also known that the fundamental group of the homotopy fiber of f contains a nilpotent subgroup whose index is uniformly bounded above ([4], [17], [6], [14], [15]). Here, we show some metric properties of the fibers of f. Note that the diameters of the fibers of f are very small (less than a constant multiple of the Gromov-Hausdorff distance μ between M and X). Let vol_m denote the m-dimensional Hausdorff measure.

Theorem 1.4. Let $f: M \to X$ be the map of Theorem 1.1. Let F_p denote the fiber $f^{-1}(p)$ over $p \in X$.

(1) The induced intrinsic metric of F_p is almost isometric to the original one, that is,

$$|xy|_{F_p} < (1 + \varkappa(\delta))|xy|$$

for any $x, y \in F_p$, where $| , |_{F_p}$ denotes the induced intrinsic metric of F_p . (In particular, we can use both metrics below.)

(2) The Hausdorff dimension of F_p is n-k. Moreover, we have

$$0 < \operatorname{vol}_{n-k} F_p < C(\operatorname{diam} F_p)^{n-k}$$

where C is a positive constant depending only on n and κ .

(3) Fix $p \in X$. Then, for any $q \in X$ sufficiently close to p, we have

$$\left|\frac{\operatorname{vol}_{n-k} F_q}{\operatorname{vol}_{n-k} F_p} - 1\right| < \varkappa(\delta).$$

The property (1) was known for fibers of almost regular maps such as $\hat{\varphi}$ ([2, 11.11]). Thus, it also holds for the fibers of $\varphi \circ f$ satisfying the inequality (1.1). This result was stated in the first version of [14], but has been deleted in the second version (due to an oversight as explained in the abstract on arXiv). Regarding (2), it was only known that the topological dimension of fibers of almost regular maps such as $\hat{\varphi}$ is no greater than n - k ([2, 11.8]). The left inequality in (2) was conjectured in [7, 4.2]. The author does not know whether the volume of F_p is actually continuous in (3).

Remark 1.5. Strictly speaking, we should not assert that our results (Theorem 1.2, Corollary 1.3 and Theorem 1.4) hold for the almost Lipschitz submersion constructed by Yamaguchi [17]. In fact, we construct the map f of Theorem 1.1 in a different way from [17] and show the inequality (1.1) for this new map but not for the original one. However, both constructions are essentially the same and (1.1) actually holds for Yamaguchi's map. See Theorem 4.1 and Remark 4.3.

The organization of this paper is as follows: In §2, we introduce some notation and conventions which will be used throughout this paper. In §3, we recall some basic facts on Alexandrov spaces, especially strainers, and list a few lemmas from [8] for later use. In §4, we construct the map f of Theorem 1.1 and show the inequality (1.1). In §5, we generalize Perelman's fibration theorem for noncritical maps in terms of the inequality (1.1) and prove Theorem 1.2 and Corollary 1.3. Most of the contents of this section are slight modifications of those of [8, §3]. In §6, we define the notion of almost regular maps in terms of the inequality (1.1) and study fibers of them (note that our definition of almost regular maps is different from that of [2]). In §6.1, we prove Theorem 1.4(1). In §6.2, we prove the left inequality of Theorem 1.4(2), and in §6.3, we prove the right inequality. In §6.4, we prove Theorem 1.4(3).

2. NOTATION AND CONVENTIONS

The dimension of Alexandrov spaces is usually denoted by n. The lower curvature bound κ of Alexandrov spaces is fixed and omitted unless otherwise stated. A positive integer k is usually no greater than n and is often less than n. We

always assume that a lower bound ℓ for the lengths of strainers is no greater than 1 especially when $\kappa < 0$ (indeed, all our arguments using strainers are local).

We denote by c and C various small and large positive constants, respectively. Unless otherwise stated, such constants depend only on n and κ . If they depend on additional parameters, it will be indicated explicitly, like $c(\varepsilon)$.

We always assume that a positive number δ is smaller than some constant c_0 depending only on n and κ . We denote by $\varkappa(\delta)$ various positive functions such that $\varkappa(\delta) \to 0$ as $\delta \to 0$. Unless otherwise stated, \varkappa depends only on n and κ . In this case, we often assume that $\varkappa(\delta)$ is also smaller than c_0 .

In §3.2, §5 and §6.3, we use another positive number ε . In these sections, \varkappa may depend additionally on ε . Furthermore, we assume that δ is smaller than some constant $c_0(\varepsilon)$ depending only on n, κ and ε , which is much smaller than any other $c(\varepsilon)$ appearing in these sections. We often assume that $\varkappa(\delta)$ is also smaller than $c_0(\varepsilon)$.

The *m*-dimensional Hausdorff measure is denoted by vol_m . We use the standard Euclidean norm on \mathbb{R}^k except in §5, where we use the maximum norm instead.

3. Preliminaries

We first recall some basic facts on Alexandrov spaces. See [2] or [1] for more details.

Let M be an n-dimensional Alexandrov space with curvature $\geq \kappa$. For a geodesic triangle $\triangle pqr$ in M with vertices p, q and r, we denote by $\tilde{\triangle}pqr$ a geodesic triangle with the same sidelengths in the simply-connected complete surface of constant curvature κ . Then, by the definition of an Alexandrov space, the natural correspondence from $\triangle pqr$ to $\tilde{\triangle}pqr$ is nonexpanding. Let $\angle qpr$ denote the angle of $\triangle pqr$ at p and $\tilde{\angle}qpr$ the corresponding angle of $\tilde{\triangle}pqr$. Then, the Alexandrov convexity implies that $\angle qpr \geq \tilde{\angle}qpr$.

For $p \in M$, we denote by Σ_p the space of directions at p. Then, Σ_p is an (n-1)dimensional compact Alexandrov space with curvature ≥ 1 . For a point $q \in M$, we denote by $q'_p \in \Sigma_p$ one of the directions of shortest paths from p to q. Furthermore, for a closed subset $A \subset M$, we denote by $A'_p \subset \Sigma_p$ the set of all directions of shortest paths from p to A. We sometimes use the notation $Q'_p \subset \Sigma_p$ to denote the set of all directions of shortest paths from p to q by regarding Q as $\{q\}$.

The class of all Alexandrov spaces with dimension $\leq n$, curvature $\geq \kappa$ and diameter $\leq D$ is compact with respect to the Gromov-Hausdorff distance. Furthermore, the class of all pointed Alexandrov spaces with dimension $\leq n$ and curvature $\geq \kappa$ is compact with respect to the pointed Gromov-Hausdorff topology.

3.1. Strainers. Let M be an n-dimensional Alexandrov space.

Definition 3.1. A point $p \in M$ is said to be (k, δ) -strained if there exists k pairs $\{(a_i, b_i)\}_{i=1}^k$ of points in M such that

$$\begin{split} \widetilde{\ \ } a_i p b_i > \pi - \delta, \ \widetilde{\ \ } a_i p a_{i'}, \widetilde{\ \ } a_i p b_{i'}, \widetilde{\ \ } b_i p b_{i'} > \pi/2 - \delta \end{split}$$

for all $1 \leq i \neq i' \leq k$. The collection $\{(a_i, b_i)\}_{i=1}^k$ is called a (k, δ) -strainer at p. The number $\min_i\{|a_ip|, |b_ip|\}$ is called the *length* of this strainer.

We now describe a few basic properties of strainers and strained points.

Lemma 3.2. Let (a,b) be a $(1,\delta)$ -strainer at $p \in M$ with length $> \ell$. Then, we have

$$\left|\tilde{\angle}axy - \angle axy\right| < \varkappa(\delta)$$

for any $x, y \in B(p, \ell \delta)$ and any shortest paths xa, xy.

See [2, 5.6] or [1, 10.8.13] for the proof.

Let $S^k(\Sigma)$ be the k-fold spherical suspension over a space Σ of curvature ≥ 1 (see [2, 4.3.1] for the definition). Note that it is isometric to the spherical join of Σ and the unit sphere \mathbb{S}^{k-1} of dimension k-1. Thus, $S^k(\Sigma)$ contains an isometric copy of \mathbb{S}^{k-1} (we identify them). Let $\{(\xi_i, \eta_i)\}_{i=1}^k$ be a collection of pairs of points in $\mathbb{S}^{k-1} \subset S^k(\Sigma)$ such that

$$\xi_i \eta_i | = \pi, \quad |\xi_i \xi_{i'}| = |\xi_i \eta_{i'}| = |\eta_i \eta_{i'}| = \pi/2$$

for all $1 \leq i \neq i' \leq k$. We call such a collection an *orthogonal k-frame* of $S^k(\Sigma)$. Conversely, if a space of curvature ≥ 1 has such a collection, then it is isometric to a k-fold spherical suspension (see [1, 10.4.3]).

The space of directions at a strained point is close to a suspension in the following sense:

Lemma 3.3. Let $p \in M$ be a (k, δ) -strained point with a strainer $\{(a_i, b_i)\}_{i=1}^k$. Let $(A_i)'_p$, $(B_i)'_p$ denote the set of all directions of shortest paths from p to a_i , b_i , respectively. Then, there exists a $\varkappa(\delta)$ -approximation $\Sigma_p \to S^k(\Sigma)$ which sends $\{((A_i)'_p, (B_i)'_p)\}_{i=1}^k$ to an orthogonal k-frame of $S^k(\Sigma)$, where Σ is a space of curvature ≥ 1 and dimension $\leq n - k - 1$ (possibly empty). In particular, if k = n, then Σ_p is $\varkappa(\delta)$ -close to \mathbb{S}^{n-1} in the Gromov-Hausdorff distance.

The proof is by contradiction (see [3, 3.2] for instance).

Remark 3.4. Furthermore, if k < n, then Σ is nonempty. Indeed, \mathbb{S}^{k-1} cannot be a limit of a collapsing sequence of Alexandrov spaces of curvature ≥ 1 . This follows, for instance, from the fact that such a limit space either has diameter no greater than $\pi/2$ or contains a proper extremal subset ([10, 3.2]). In particular, if k = n - 1, then Σ_p is $\varkappa(\delta)$ -close to \mathbb{S}^{n-1} or the closed unit hemisphere \mathbb{S}^{n-1}_+ .

Let $0 < \varepsilon < 1$. Let X and Y be metric spaces. A map $f: X \to Y$ is called an ε -almost isometry if it is surjective and $||f(x)f(y)|/|xy|-1| < \varepsilon$ for any $x, y \in X$. Let U be an open subset of X. A map $f: U \to Y$ is called an ε -open map if for any $x \in U$ and $v \in Y$ such that $\overline{B}(x, \varepsilon^{-1}|f(x)v|) \subset U$, there exists $y \in U$ such that f(y) = v and $\varepsilon |xy| \leq |f(x)v|$.

For a (k, δ) -strainer $\{(a_i, b_i)\}_{i=1}^k$ in M, we call $f = (|a_1 \cdot |, \dots, |a_k \cdot |) : M \to \mathbb{R}^k$ the *distance coordinate* associated with this strainer. The above two lemmas imply the following:

Proposition 3.5. Let $f : M \to \mathbb{R}^k$ be the distance coordinate associated with a (k, δ) -strainer at p with length $> \ell$. Then,

- (1) if k = n, then f is a $\varkappa(\delta)$ -almost isometry from $B(p, \ell\delta)$ to an open subset of \mathbb{R}^n ;
- (2) if k < n, then f is a $(1 + \varkappa(\delta))$ -Lipschitz and $(1 \varkappa(\delta))$ -open map on $B(p, \ell\delta)$.

For the proof of (1), see [2, 9.4] or [1, 10.9.16]. For the proof of (2), see [3, 3.3] for instance.

3.2. Lemmas from Perelman's paper. We list a few lemmas on spaces of curvature > 1 from [8, §2], which will be used in §5 and §6.3.

Let Σ be an (n-1)-dimensional Alexandrov space with curvature ≥ 1 . Here we always assume that δ is sufficiently small compared with ε (see §2).

Lemma 3.6 ([8, 2.2]). Σ cannot contain n + 2 compact subsets $\{A_i\}_{i=1}^{n+2}$ such that $|A_iA_i| > \pi/2 - \delta$ for any $i \neq j$ and $|A_1A_i| > \pi/2 + \varepsilon$ for any $i \geq 3$.

- **Lemma 3.7** ([8, 2.3, 2.4]). (1) Let $\{A_i\}_{i=1}^{k+2}$ be k+2 compact subsets $(0 \le k \le n-1)$ of Σ such that $|A_iA_j| > \pi/2 \delta$ for any $i \ne j$ and $|A_1A_i| > \pi/2 + \varepsilon$ for any $i \ge 2$. Then, there is a point $\xi \in \Sigma$ such that $|\xi A_i| = \pi/2$ for any $i \ge 3$ and $|\xi A_1| > \pi/2 + c(\varepsilon)$, $|\xi A_2| < \pi/2 c(\varepsilon)$.
 - (2) (1) holds true if we replace the assumption $|A_1A_2| > \pi/2 + \varepsilon$ by $|A_1A_2| > \pi/2 \delta$ and the conclusion $|\xi A_1| > \pi/2 + c(\varepsilon)$ by $|\xi A_1| > \pi/2 \varkappa(\delta)$.
 - (3) Under the assumptions of (1), there is a point $\xi \in \Sigma$ such that $|\xi A_i| = \pi/2$ for any $i \ge 3$ and $|\xi A_2| > \pi/2 + c(\varepsilon)$.

Lemma 3.8 ([8, 2.5.2]). Let A be a subset of Σ . Let $A(\pi/2 \pm a)$ denote the set of points $x \in \Sigma$ such that $||Ax| - \pi/2| \leq a$. Then, the number of ω -discrete points contained in $A(\pi/2 \pm a)$ is at most Ca/ω^{n-1} for any $0 < \omega \leq a$.

4. Construction of a global map

In this section, we construct the global map f of Theorem 1.1 and prove the inequality (1.1).

Theorem 4.1 (cf. [17, 0.2]). Let X be a k-dimensional Alexandrov space such that every point has a (k, δ) -strainer with length $> \ell$. Let M be an n-dimensional Alexandrov space and $g: M \to X$ a μ -approximation. Suppose $\mu < c(\ell\delta^2) \ll \ell\delta^2$. Then, there exists a map $f: M \to X$, which is C μ -close to g, satisfying the following property:

Let $\{(a_i, b_i)\}_{i=1}^k$ be a (k, δ) -strainer at $p \in X$ such that $|a_i p| = \ell \delta$ for all i. Let $\varphi : X \to \mathbb{R}^k$ be its distance coordinate and $\hat{\varphi} : M \to \mathbb{R}^k$ a lift of φ . Then, we have

(4.1)
$$\left| \left(\varphi \circ f(x) - \varphi \circ f(y) \right) - \left(\hat{\varphi}(x) - \hat{\varphi}(y) \right) \right| < \varkappa(\delta) |xy|$$

for any $x, y \in B(\hat{p}, \ell \delta^2)$, where $\hat{p} \in M$ denotes a lift of p (i.e. $|pg(\hat{p})| < \mu$).

Remark 4.2. The inequality (4.1) immediately implies that f is a $\varkappa(\delta)$ -almost isometry when k = n (cf. [2, 9.8], [12, 3.1], [13]) and is a $(1 - \varkappa(\delta))$ -open, $\varkappa(\delta)$ -almost Lipschitz submersion when k < n. See Propositions 6.5 and 6.6.

Remark 4.3. The inequality (4.1) actually holds for the almost Lipschitz submersion constructed by Yamaguchi [17] (use [17, 4.6, 4.13]). His construction is based on an embedding of X to the Hilbert space $L^2(X)$ of all L^2 -functions on X and the existence of a "tubular neighborhood" of the image of X. Here we give a more direct proof. Our construction is a generalization of that of the almost isometry in [12, 3.1] when k = n. However, both constructions are based on the same idea of gluing local distance coordinates.

Proof of Theorem 4.1. We denote by $\hat{a} \in M$ a lift of $a \in X$ with respect to g. Set $r := \ell \delta^2$. Let $\{p_j\}_{j=1,2,\ldots}$ be a maximal r/2-discrete net of X (possibly infinite).

Take a (k, δ) -strainer $\{(\alpha_i^j, \beta_i^j)\}_{i=1}^k$ with length $> \ell$ for each p_j and let a_i^j, b_i^j be points on shortest paths $p_j \alpha_j^i, p_j \beta_j^j$ at distance $\ell \delta$ from p_j , respectively. Set

$$\begin{aligned} \varphi_j &:= (|a_1^j \cdot |, \dots, |a_k^j \cdot |), \\ \hat{\varphi}_j &:= (|\hat{a}_1^j \cdot |, \dots, |\hat{a}_k^j \cdot |), \end{aligned} \qquad U_j &:= B(p_j, r), \quad \lambda U_j &:= B(p_j, \lambda r), \\ \hat{U}_j &:= B(\hat{p}_j, r), \quad \lambda \hat{U}_j &:= B(\hat{p}_j, \lambda r) \end{aligned}$$

for $\lambda > 0$. Since $\mu \ll r$, $\{\hat{U}_j\}_j$ is a covering of M and $\{(\hat{a}_i^j, \hat{b}_i^j)\}_{i=1}^k$ is a $(k, \varkappa(\delta))$ strainer for \hat{p}_j . By Proposition 3.5, φ_j is a $\varkappa(\delta)$ -almost isometry from $10U_j$ to an
open subset of \mathbb{R}^k and $\hat{\varphi}_j$ is a $(1 + \varkappa(\delta))$ -Lipschitz map on $3\hat{U}_j$. Hence, we can
define $\varphi_j^{-1} \circ \hat{\varphi}_j$ on $3\hat{U}_j$. Note that $|\varphi_j^{-1} \circ \hat{\varphi}_j, g| < C\mu$. We take an average of them
to obtain a global map. Define $f_j : \bigcup_{i'=1}^j \hat{U}_{i'} \to X$ inductively as follows:

$$f_1 := \varphi_1^{-1} \circ \hat{\varphi}_1 : \hat{U}_1 \to X$$

and for $j \geq 2$,

$$f_j := \begin{cases} f_{j-1} & \text{on} & \bigcup_{j'=1}^{j-1} \hat{U}_{j'} \setminus 2\hat{U}_j \\ \varphi_j^{-1} \circ ((1-\chi_j)\varphi_j \circ f_{j-1} + \chi_j \hat{\varphi}_j) & \text{on} & \bigcup_{j'=1}^{j-1} \hat{U}_{j'} \cap (2\hat{U}_j \setminus \hat{U}_j) \\ \varphi_j^{-1} \circ \hat{\varphi}_j & \text{on} & \hat{U}_j. \end{cases}$$

Here $\chi_j := \chi(|\hat{p}_j \cdot |/r)$, where $\chi : [0, \infty) \to [0, 1]$ is a smooth function such that $\chi \equiv 1$ on [0, 1] and $\chi \equiv 0$ on $[2, \infty]$. Note that χ_j is L/r-Lipschitz for some constant L.

Then, it easily follows that $|f_j, g| < C\mu$ by the induction on j (in particular, the above definition of f_j works). Note that the number of the induction steps at each point in the domain of f_j is uniformly bounded above. Indeed, since $\{p_j\}_j$ is r/2-discrete and $\mu \ll r$, the multiplicity of the covering $\{2\hat{U}_j\}_j$ is bounded above by some constant depending only on k. We define $f(x) := f_{N_x}(x)$ for $x \in M$, where $N_x := \max\{j \mid x \in 2\hat{U}_j\}$.

Now, we show the inequality (4.1). Let p, \hat{p} , φ , $\hat{\varphi}$ be as in the assumption. Set $\lambda \hat{U} := B(\hat{p}, \lambda r)$ as above. We prove by the induction on j that

(4.2)
$$\left| (\varphi \circ f_j(x) - \varphi \circ f_j(y)) - (\hat{\varphi}(x) - \hat{\varphi}(y)) \right| < \varkappa(\delta) |xy|$$

for $x, y \in 3\hat{U} \cap \text{Dom}(f_j)$, where $\text{Dom}(f_j) = \bigcup_{j'=1}^j \hat{U}_{j'}$. Note that we may assume |xy| < r since $|f_j, g| < C\mu$ and $\mu \ll r$.

First, we prove the inequality (4.2) for the special case $\varphi = \varphi_j$ and $p = p_j$. The base case j = 1 is trivial. Suppose $j \ge 2$. Let us consider the case $x, y \in (2\hat{U}_j \setminus \hat{U}_j) \cap \text{Dom}(f_j)$ (the other cases are similar). Then, we have

$$\begin{aligned} (\varphi_j \circ f_j(x) - \varphi_j \circ f_j(y)) &- (\hat{\varphi}_j(x) - \hat{\varphi}_j(y)) \\ &= (1 - \chi_j(x))(\varphi_j \circ f_{j-1}(x) - \hat{\varphi}_j(x)) - (1 - \chi_j(y))(\varphi_j \circ f_{j-1}(y) - \hat{\varphi}_j(y)) \\ &= (1 - \chi_j(x))((\varphi_j \circ f_{j-1}(x) - \varphi_j \circ f_{j-1}(y)) - (\hat{\varphi}_j(x) - \hat{\varphi}_j(y))) \\ &- (\chi_j(x) - \chi_j(y))(\varphi_j \circ f_{j-1}(y) - \hat{\varphi}_j(y)). \end{aligned}$$

The norm of the first term of the last formula is less than $\varkappa(\delta)|xy|$ by the induction hypothesis. The same is true for the second term since χ_j is L/r-Lipschitz and $|\varphi_j \circ f_{j-1}, \hat{\varphi}_j| < C\mu$, where $\mu \ll r$.

Next, we consider the general case. Lemma 3.2 implies that the inequality (4.2) is equivalent to

(4.3)
$$\left| \cos \angle a_i f_j(x) f_j(y) \cdot \frac{|f_j(x) f_j(y)|}{|xy|} - \cos \angle \hat{a}_i xy \right| < \varkappa(\delta)$$

for all $1 \leq i \leq k$. By the induction hypothesis, we may assume that $x, y \in 3\hat{U} \cap \text{Dom}(f_j) \cap 3\hat{U}_j$ (recall |xy| < r). Fix $1 \leq i \leq k$. First, by the strainer $\{(a_{i'}^j, b_{i'}^j)\}_{i'=1}^k$ at $f_j(x)$ and Lemma 3.3, $\Sigma_{f_j(x)}$ is $\varkappa(\delta)$ -close to \mathbb{S}^{k-1} in the Gromov-Hausdorff distance. Thus, we have

$$\left|\cos \angle a_i f_j(x) f_j(y) - \sum_{i'=1}^k \cos \angle a_i f_j(x) a_{i'}^j \cdot \cos \angle a_{i'}^j f_j(x) f_j(y)\right| < \varkappa(\delta).$$

Next, since $|f_j, g| < C\mu$ and $\mu \ll r$, we have $|\tilde{\angle}a_i f_j(x) a_{i'}^j - \tilde{\angle}\hat{a}_i x \hat{a}_{i'}^j| < \varkappa(\delta)$ for all $1 \leq i' \leq k$. Recall that $a_{i'}^j, b_{i'}^j$ are on the shortest paths $p_j \alpha_{i'}^j, p_j \beta_{i'}^j$ at distance $\ell \delta$ from p_j , where $\{(\alpha_{i'}^j, \beta_{i'}^j)\}_{i'=1}^k$ is a (k, δ) -strainer with length $> \ell$ at p_j . Hence, by using Lemma 3.2 twice, we have

$$\left| \angle a_i f_j(x) a_{i'}^j - \angle \hat{a}_i x \hat{a}_{i'}^j \right| < \varkappa(\delta)$$

for all $1 \leq i' \leq k$. Finally, by the strainer $\{(\hat{a}_{i'}^j, \hat{b}_{i'}^j)\}_{i'=1}^k$ at x and Lemma 3.3 again, Σ_x is $\varkappa(\delta)$ -close to a k-fold spherical suspension $S^k(\Sigma)$. Furthermore, this approximation sends $(\hat{a}_i)'_x$ into the $\varkappa(\delta)$ -neighborhood of \mathbb{S}^{k-1} in $S^k(\Sigma)$ since the previous inequality implies that $\sum_{i'=1}^k \cos^2 \angle \hat{a}_i x \hat{a}_{i'}^j \approx \sum_{i'=1}^k \cos^2 \angle a_i f_j(x) a_{i'}^j \approx 1$ (roughly speaking, $(\hat{a}_i)'_x$ is a "horizontal direction"). Therefore, we have

$$\left|\cos \angle \hat{a}_i xy - \sum_{i'=1}^{\kappa} \cos \angle \hat{a}_i x \hat{a}_{i'}^j \cdot \cos \angle \hat{a}_{i'}^j xy \right| < \varkappa(\delta).$$

Combining the above three inequalities with the inequality (4.3) for the special case $\varphi = \varphi_j$ and $p = p_j$, we obtain the general one.

5. Modification of Perelman's fibration theorem

In this section, we prove Theorem 1.2 and Corollary 1.3. We generalize the notion of noncritical maps introduced by Perelman [8] in terms of the inequality (4.1) and prove the fibration theorem [8, 1.4.1] for such generalized noncritical maps. Although the proof is almost the same as the original one, we give the details because of the difficulty of the original proof. Hence, most of the contents of this section are just slight modifications of those of [8, §3].

Here we use another positive number ε in addition to δ . Note that \varkappa in this section may depend on ε . Furthermore, we assume that δ is much smaller than ε and every $c(\varepsilon)$ in this section and that so is $\varkappa(\delta)$. We use the maximum norm on \mathbb{R}^k unless otherwise stated. See §2 for the notation and conventions.

The following definition is the key to generalizing the fibration theorem. For a point p and subsets A, B in an Alexandrov space, $\angle ApB$ denotes the comparison angle at p of the comparison triangle with sidelengths |pA|, |pB| and |AB| if it exists; otherwise $\angle := 0$.

Definition 5.1 (cf. [8, 3.1]). Let U be an open subset of an Alexandrov space M. A map $f = (f_1, \ldots, f_k) : U \to \mathbb{R}^k$ is said to be (ε, δ) -noncritical (in the generalized sense) at $p \in U$ if it satisfies the following conditions:

8

(1) Each f_i satisfies the following inductive condition: there exists a function $g_i: U \to \mathbb{R}$ such that

(5.1)
$$|(f_i(x) - f_i(y)) - (g_i(x) - g_i(y))| < \delta |xy|$$

for any $x, y \in U$, and

$$g_i = \inf_{\gamma} g_{i\gamma}, \quad g_{i\gamma} = \varphi_{i\gamma}(|A_{i\gamma} \cdot |) + \sum_{l=1}^{i-1} \varphi_{i\gamma}^l(f_l(\cdot)) + c_{i\gamma},$$

where $c_{i\gamma} \in \mathbb{R}$, $A_{i\gamma}$ are compact subsets of M, $\varphi_{i\gamma}$ and $\varphi_{i\gamma}^l$ have right and left derivatives, $\varphi_{i\gamma}^l$ are ε^{-1} -Lipschitz functions, and $\varphi_{i\gamma}$ are increasing functions such that $\varphi_{i\gamma}(0) = 0$ and $\varepsilon |x-y| \leq |\varphi_{i\gamma}(x) - \varphi_{i\gamma}(y)| \leq \varepsilon^{-1} |x-y|$.

(2) The sets of indices $\Gamma_i(p) := \{\gamma \mid g_i(p) = g_{i\gamma}(p)\}$ satisfy $\#\Gamma_i(p) < \varepsilon^{-1}$. Furthermore, there exists $\rho = \rho(p) > 0$ such that for all i

$$g_i(x) < g_{i\gamma}(x) - \rho$$

for $x \in B(p, \rho)$ and $\gamma \notin \Gamma_i(p)$ (we assume $B(p, \rho) \subset U$).

- (3) $\angle A_{i\alpha}pA_{j\beta} > \pi/2 \delta$ for all $i \neq j, \alpha \in \Gamma_i(p), \beta \in \Gamma_j(p)$.
- (4) There exists a point $w = w(p) \in M$ such that $\tilde{\angle}A_{i\gamma}pw > \pi/2 + \varepsilon$ for all i and $\gamma \in \Gamma_i(p)$.

We have added the inequality (5.1) to the original definition. In particular, each f_i is not necessarily defined by distance functions unlike g_i and does not always have directional derivatives. However, the inequality (5.1) guarantee that their difference quotients are almost equal. In case $f_i \equiv g_i$, our definition coincides with the original one. Note that each g_i is defined by f_l $(1 \le l \le i - 1)$ but not by g_l (this is necessary to prove Proposition 5.11 below).

The purpose of this section is to prove the following generalized fibration theorem:

Theorem 5.2 (cf. [8, 1.4.1]). Let U be a domain of an n-dimensional Alexandrov space M such that $\operatorname{vol}_{n-1} \Sigma_p > \varepsilon$ for any $p \in U$. If a map $f: U \to \mathbb{R}^k$ is proper and (ε, δ) -noncritical in the generalized sense for some $\delta \ll \varepsilon$, then it is a locally trivial fibration.

Remark 5.3. The term "noncritical" (not " (ε, δ) -noncritical") in the original statement [8, 1.4.1] includes the assumption on the volume of spaces of directions (see [8, 3.7] or Definition 5.14 below). However, we do not use it here to emphasize the dependence of the fibration theorem on the volume of spaces of directions.

Let us first recall the structure of the proof of the original fibration theorem. It consists of two parts: the first half is its geometric part, which is presented in [8, §3], and the second half is its topological part, which is presented in [8, §1]. Furthermore, as remarked in [8, 1.3], all the arguments in the topological part are based only on the properties of noncritical maps established in the geometric part. In particular, we do not have to go back to the definition of noncritical maps in the topological part. Thus, in order to prove the generalized fibration theorem, it is sufficient to verify that all the propositions about noncritical maps in the geometric part hold true for the generalized ones. Indeed, the same proofs work well by using the inequality (5.1). Nevertheless, we give the details to explain how and where our modification works because the original proofs are very complicated.

From here (until Definition 5.14), we proceed in parallel with [8, §3]. We first prove the openness of a generalized noncritical map (note that the set of generalized noncritical points of a given map is clearly open). The following proof is a good demonstration of the usefulness of the inequality (5.1) and is actually the only place where we use it in the proof of the generalized fibration theorem (see also Remark 5.6 below).

Proposition 5.4 (cf. [8, 3.2]). Let U be an open subset of an n-dimensional Alexandrov space M. Let $f: U \to \mathbb{R}^k$ be (ε, δ) -noncritical in the generalized sense at any point of U. Then, $k \leq n$ and f is $c(\varepsilon)$ -open (and is locally $c(\varepsilon)^{-1}$ -Lipschitz). Furthermore, if k = n, then f is a local bi-Lipschitz homeomorphism.

We need the following elementary lemma to prove the $c(\varepsilon)$ -openness, which is a metric version of [8, 2.1.1]. The proof is easy and will be omitted.

Lemma 5.5 (cf. [8, 2.1.1]). Let $f : U \to \mathbb{R}^k$ be a continuous (not necessarily differentiable) map defined on an open subset of an Alexandrov space. Let $\|\cdot\|$ be an arbitrary norm on \mathbb{R}^k . Suppose that for any $x \in U$ and $v \in \mathbb{R}^k$ with $f(x) \neq v$, there exists a point $y \in U$ with $y \neq x$ such that

$$||f(y) - v|| - ||f(x) - v|| \le -\varepsilon |xy|.$$

Then, f is ε -open with respect to the norm $\|\cdot\|$.

Proof of Proposition 5.4. The inequality $k \leq n$ immediately follows from Lemma 3.6 and Definition 5.1(3),(4). Let us prove the $c(\varepsilon)$ -openness of f. We show that for any $p \in U$ and $1 \leq i \leq k$, there exist $q_i^{\pm} \in U$ arbitrarily close to p such that

(5.2)
$$\begin{aligned} |f_j(q_i^{\pm}) - f_j(p)| &< \varkappa(\delta) |pq_i^{\pm}| & \text{for } j < i, \\ c(\varepsilon) |pq_i^{\pm}| &< \pm (f_j(q_i^{\pm}) - f_j(p)) < c(\varepsilon)^{-1} |pq_i^{\pm}| & \text{for } j = i, \\ |f_j(q_i^{\pm}) - f_j(p)| &< c(\varepsilon)^{-1} |pq_i^{\pm}| & \text{for } j > i. \end{aligned}$$

Let us first explain how to finish the proof, assuming the above. Define a norm $||v|| := \sum_{j=1}^{k} c_j(\varepsilon) |v_j|$ for $v = (v_1 \dots, v_k) \in \mathbb{R}^k$, where $c_j(\varepsilon) \ll c_{j-1}(\varepsilon)$ (the choices of $c_j(\varepsilon)$ depend on the constants $c(\varepsilon)$ in the inequalities (5.2) so that the following argument holds). By Lemma 5.5, it suffices to show that for any $p \in U$ and $v \in \mathbb{R}^k$ with $f(p) \neq v$, there is $q \in U$ with $q \neq p$ such that

$$||f(q) - v|| - ||f(p) - v|| \le -c(\varepsilon)|pq|.$$

Without loss of generality, we may assume $f_i(p) < v_i$ for some *i*. Choose q_i^+ so close to *p* that $f_i(q_i^+) < v_i$ and put $q := q_i^+$. Then, the inequalities (5.2) and the suitable choices of $c_j(\varepsilon)$ yield the desired inequality (note $\varkappa(\delta) \ll c_j(\varepsilon)$).

Now, we prove the inequalities (5.2). Set $A'_j(p) := \bigcup_{\gamma \in \Gamma_j(p)} (A_{j\gamma})'_p$ for each $1 \leq j \leq k$. Fix $1 \leq i \leq k$. Then, by Definition 5.1(3),(4) and Lemma 3.7(1),(3), we get $\xi_i^{\pm} \in \Sigma_p$ such that

$$\pm (|A'_i(p)\xi_i^{\pm}| - \pi/2) > c(\varepsilon), \quad |A'_j(p)\xi_i^{\pm}| = \pi/2$$

for any $j \neq i$. Choose $q_i^{\pm} \in U$ near p such that $(q_i^{\pm})'_p$ is sufficiently close to ξ_i^{\pm} . We show the inequalities (5.2) by the induction on j. Here is an outline. Fix j and suppose that (5.2) hold for j' < j. Then, by the definition of g_j (see Definition 5.1(1)), the inequality (5.2) for j with f_j replaced by g_j holds. Together with the

10

inequality (5.1), this implies (5.2) for j. Let us demonstrate this in the case where k = 3, i = 2 and the sign is +. Put $\xi := \xi_2^+$ and $q := q_2^+$. First,

$$g_1(q) - g_1(p) = \min_{\gamma \in \Gamma_1(p)} \left\{ \varphi_{1\gamma}(|A_{1\gamma}q|) - \varphi_{1\gamma}(|A_{1\gamma}p|) \right\}$$
$$= \min_{\gamma \in \Gamma_1(p)} \left\{ -\varphi'_{1\gamma}(|A_{1\gamma}p|) \cos |(A_{1\gamma})'_p \xi| \cdot |pq| + o(|pq|) \right\},$$

where $\varphi'_{1\gamma}$ denotes the right or left derivative of $\varphi_{1\gamma}$. Since $\varphi'_{1\gamma} \ge 0$ and $|A'_1(p)\xi| = \pi/2$, we have

$$|g_1(q) - g_1(p)| < \delta |pq|$$

provided q is sufficiently close to p. Then, the inequality (5.1) implies

$$|f_1(q) - f_1(p)| < 2\delta |pq|$$

Second,

$$g_{2}(q) - g_{2}(p) = \min_{\gamma \in \Gamma_{2}(p)} \{ \varphi_{2\gamma}(|A_{2\gamma}q|) - \varphi_{2\gamma}(|A_{2\gamma}p|) + \varphi_{2\gamma}^{1}(f_{1}(q)) - \varphi_{2\gamma}^{1}(f_{1}(p)) \} = \min_{\gamma \in \Gamma_{2}(p)} \{ -\varphi_{2\gamma}'(|A_{2\gamma}p|) \cos |(A_{2\gamma})_{p}'\xi| \cdot |pq| + o(|pq|) + \varphi_{2\gamma}^{1}(f_{1}(q)) - \varphi_{2\gamma}^{1}(f_{1}(p)) \}.$$

Since $\varphi_{2\gamma}^1$ is ε^{-1} -Lipschitz, the inequality obtained in the previous step yields $|\varphi_{2\gamma}^1(f_1(q)) - \varphi_{2\gamma}^1(f_1(p))| < \varkappa(\delta)|pq|$. Furthermore, since $\varepsilon \leq \varphi_{2\gamma}' \leq \varepsilon^{-1}$ and $|A_2'(p)\xi| > \pi/2 + c(\varepsilon)$, we have

$$c(\varepsilon)|pq| < g_2(q) - g_2(p) < c(\varepsilon)^{-1}|pq|$$

provided q is sufficiently close to p (note $\varkappa(\delta) \ll c(\varepsilon)$). Then, the inequality (5.1) implies

$$c(\varepsilon)|pq| < f_2(q) - f_2(p) < c(\varepsilon)^{-1}|pq|.$$

Finally,

$$g_{3}(q) - g_{3}(p) = \min_{\gamma \in \Gamma_{3}(p)} \Big\{ \varphi_{3\gamma}(|A_{3\gamma}q|) - \varphi_{3\gamma}(|A_{3\gamma}p|) + \sum_{l=1,2} \big(\varphi_{3\gamma}^{l}(f_{l}(q)) - \varphi_{3\gamma}^{l}(f_{l}(p)) \big) \Big\}.$$

Since both $\varphi_{3\gamma}$ and $\varphi_{3\gamma}^l$ are ε^{-1} -Lipschitz, we have

$$|g_3(q) - g_3(p)| < c(\varepsilon)^{-1} |pq|.$$

Then, the inequality (5.1) implies

$$|f_3(q) - f_3(p)| < c(\varepsilon)^{-1} |pq|.$$

This completes the proof of the inequalities (5.2) in our special case. The general case is similar. Note that the local $c(\varepsilon)^{-1}$ -Lipschitzness of f also follows from a similar inductive argument.

Next, we consider the case k = n. It suffices to show that f is injective near each $p \in U$. Let $\rho = \rho(p) > 0$ and $w = w(p) \in M$ be as in Definition 5.1 and take $0 < r < \min_{i,\gamma \in \Gamma_i(p)} \{\rho, \delta | wp |, \delta | A_{i\gamma}p |\}$. Suppose there exist two distinct points $x, y \in B(p, r)$ such that f(x) = f(y). We may assume $|wx| \leq |wy|$. In particular, we have $\tilde{\angle}wxy > \pi/2 - \varkappa(\delta)$. On the other hand, by the definition of noncriticality, we have $\tilde{\angle}A_{i\gamma}xw > \pi/2 + c(\varepsilon)$ and $\tilde{\angle}A_{i\gamma}xA_{j\beta} > \pi/2 - \varkappa(\delta)$ for all $i \neq j$ and $\gamma \in \Gamma_i(p), \beta \in \Gamma_j(p)$. Furthermore, we show $\tilde{\angle}A_{i\gamma}xy > \pi/2 - \varkappa(\delta)$ for all i and $\gamma \in \Gamma_i(x)$ (note $\Gamma_i(x) \subset \Gamma_i(p)$). Then, these inequalities contradict Lemma 3.6 for Σ_x since $\varkappa(\delta) \ll c(\varepsilon)$. Let $\gamma \in \Gamma_i(x)$. We may assume $|A_{i\gamma}x| > |A_{i\gamma}y|$; otherwise, we have $\tilde{\angle}A_{i\gamma}xy > \pi/2 - \varkappa(\delta)$. Then, we have

$$egin{aligned} g_i(x) - g_i(y) &\geq g_{i\gamma}(x) - g_{i\gamma}(y) \ &= arphi_{i\gamma}(|A_{i\gamma}x|) - arphi_{i\gamma}(|A_{i\gamma}y|) \ &\geq arepsilon(|A_{i\gamma}x| - |A_{i\gamma}y|), \end{aligned}$$

since f(x) = f(y) and $\varphi_{i\gamma}$ is an increasing function with co-Lipschitz constant ε . On the other hand, we have $|g_i(x) - g_i(y)| < \delta |xy|$ by the inequality (5.1). Together with the above inequality, this implies $\tilde{\angle}A_{i\gamma}xy > \pi/2 - \varkappa(\delta)$.

Remark 5.6. The same arguments as in the above proof show the following two properties. Let $f: U \to \mathbb{R}^k$ be (ε, δ) -noncritical in the generalized sense at $p \in U$.

- Let ξ ∈ Σ_p be a direction such that |A'_i(p)ξ| = π/2 for all i, where A'_i(p) = U_{γ∈Γ_i(p)}(A_{iγ})'_p. Then, for any q ∈ U sufficiently close to p such that q'_p is sufficiently close to ξ, we have |f(p)f(q)| < κ(δ)|pq|.
 Let 0 < r < min_{i,γ∈Γ_i(p)}{ρ(p), δ|A_{iγ}p|}. Then, for any x, y ∈ B(p, r) with
- (2) Let $0 < r < \min_{i,\gamma \in \Gamma_i(p)} \{\rho(p), \delta | A_{i\gamma}p| \}$. Then, for any $x, y \in B(p, r)$ with $|f(x)f(y)| < \delta |xy|$, we have $\tilde{\angle}A_{i\gamma}xy > \pi/2 \varkappa(\delta)$ for all i and $\gamma \in \Gamma_i(x)$.

Indeed, (1) follows from the same inductive argument as in the proof of the inequalities (5.2) for j < i. (2) follows from the same argument as in the proof of the local injectivity when k = n (note that the assumption k = n is not needed here and that the condition f(x) = f(y) used in the above proof can be weakened to $|f(p)f(q)| < \delta |pq|$). Hereafter, we will often use these properties as well as Proposition 5.4. Furthermore, as long as we use them, we do not need the inequality (5.1) anymore in the proof of the generalized fibration theorem.

From now on, we consider the case k < n unless otherwise stated. The next proposition is actually unnecessary for the proof of the fibration theorem, but is shown here (it is used to prove the stability theorem [8, 4.3]).

Proposition 5.7 (cf. [8, 3.3]). Let $f: U \to \mathbb{R}^k$ be (ε, δ) -noncritical in the generalized sense at $p \in U$. Let $\Pi := f^{-1}(f(p)), \rho_0 := \delta \min_{i,\gamma \in \Gamma_i(p)} \{\rho(p), |w(p)p|, |A_{i\gamma}p|\}$ and $q, r \in \Pi \cap B(p, \rho_0)$. Then, there exists a curve in Π connecting q and r of length $\leq c(\varepsilon)|qr|$.

Proof. We may assume that $|wq| \leq |wr|$. Then, we have $|w'_q, r'_q| > \pi/2 - \varkappa(\delta)$. By Remark 5.6(2), we have $|(A_{i\gamma})'_q, r'_q| > \pi/2 - \varkappa(\delta)$ for all i and $\gamma \in \Gamma_i(q)$. Moreover, by the definition of noncriticality, we have $|(A_{i\gamma})'_q, w'_q| > \pi/2 + c(\varepsilon)$ and $|(A_{i\gamma})'_q, (A_{j\beta})'_q| > \pi/2 - \varkappa(\delta)$ for all $i \neq j$ and $\gamma \in \Gamma_i(q), \beta \in \Gamma_j(q)$. Set $A'_i(q) := \bigcup_{\gamma \in \Gamma_i(q)} (A_{i\gamma})'_q$. Then, applying Lemma 3.7(2) to $w'_q, r'_q, A'_1(q), \ldots, A'_k(q)$, we get a direction $\xi \in \Sigma_q$ such that

$$|r'_{a}\xi| < \pi/2 - c(\varepsilon), \quad |A'_{i}(q)\xi| = \pi/2$$

for all *i*. Choose q_1 near q such that $(q_1)'_q$ is close to ξ . Then, the first inequality above implies $|rq_1| \leq |rq| - c(\varepsilon)|qq_1|$. Furthermore, the second inequalities imply $|f(q)f(q_1)| < \varkappa(\delta)|qq_1|$ (see Remark 5.6(1)). By the $c(\varepsilon)$ -openness of f, we obtain

 $q_2 \in \Pi$ near q_1 such that $c(\varepsilon)|q_1q_2| \leq |f(q_1)f(q)|$ (cf. [8, 2.1.3]). Therefore, we have

$$\begin{aligned} |rq_2| &\leq |rq_1| + |q_1q_2| \\ &\leq |rq| - c(\varepsilon)|qq_1| + c(\varepsilon)^{-1}\varkappa(\delta)|qq_1| \\ &\leq |rq| - c(\varepsilon)|qq_2|. \end{aligned}$$

Now, the desired curve is obtained by taking a limit of broken geodesics.

The following setting will be used in all the arguments below.

Setting 5.8 (cf. [8, 3.4]). Let $f: U \to \mathbb{R}^k$ be (ε, δ) -noncritical in the generalized sense at $p \in U$, where U is an open subset of an n-dimensional Alexandrov space M and k < n. Assume $\operatorname{vol}_{n-1} \Sigma_p \ge \varepsilon$. Let $w = w(p) \in M$ be as in Definition 5.1. Then, the Bishop-Gromov inequality implies that $\operatorname{vol}_{n-1} B(w'_p, \varepsilon/2) \ge c(\varepsilon)$. Let $0 < \omega < \delta$. Choose points $\{w_{\alpha}\}_{\alpha=1}^{N} \subset M$ near p such that

- $N \ge L/\omega^{n-1}$, where $L = c(\varepsilon)$;
- $\tilde{\angle} w_{\alpha} p w_{\beta} > \omega \ (1 \le \alpha \ne \beta \le N);$
- $\tilde{\angle} w_{\alpha} p A_{i\gamma} > \pi/2 + \varepsilon/2 \ (1 \le \alpha \le N, \ 1 \le i \le k, \ \gamma \in \Gamma_i(p)).$

Let V be a small neighborhood of p such that for any $x \in V$

- $\tilde{\angle} w_{\alpha} x w_{\beta} > \omega \ (1 \le \alpha \ne \beta \le N);$
- $\tilde{\angle} w_{\alpha} x A_{i\gamma} > \pi/2 + \varepsilon/2 \ (1 \le \alpha \le N, \ 1 \le i \le k, \ \gamma \in \Gamma_i(p));$
- $\tilde{\angle}A_{i\alpha}xA_{j\beta} > \pi/2 \delta \ (1 \le i \ne j \le k, \ \alpha \in \Gamma_i(p), \ \beta \in \Gamma_j(p));$ $|xp| < \min\{\rho(p), \delta|pw_\alpha|, \delta|pA_{i\gamma}| \ | \ 1 \le \alpha \le N, 1 \le i \le k, \gamma \in \Gamma_i(p)\}.$

Define a function $\sigma: V \to \mathbb{R}$ by

$$\sigma(x) := \frac{1}{N} \sum_{\alpha=1}^{N} |w_{\alpha}x|.$$

We first prove the following two lemmas under the above setting.

Lemma 5.9 (cf. [8, 3.4 Assertion 1]). Under Setting 5.8, let $x, y \in V$ be such that $|f(x)f(y)| < \delta |xy|$. Then, one of the following holds:

- (1) $(f, |x, \cdot|)$ is $(c(\varepsilon), \varkappa(\delta))$ -noncritical in the generalized sense at y;
- (2) $\sigma(y) \sigma(x) > c(\varepsilon)|xy|$.

Proof. Observe that the conditions (1)–(3) of Definition 5.1 for (f, |x|) at y are clearly satisfied (see Remark 5.6(2)). The rest of the proof is exactly the same as the original one. Let $a = c(\varepsilon)$ and $b = c(\varepsilon)$ be constants such that $a \ll b \ll L$. Suppose that (2) does not hold for this a. Then, the mean value of $\cos |(w_{\alpha})'_{u}, X'_{u}|$ is less than 2a, where X'_y denotes the set of all directions of shortest paths from y to x. On the other hand, by Lemma 3.8, the number of α such that $||(w_{\alpha})'_{y}, X'_{y}| - \pi/2| \leq 1$ b is less than $Cb/\omega^{n-1} \ll N$. Hence, there exists α such that $|(w_{\alpha})'_{u}, X'_{u}| >$ $\pi/2 + b$; otherwise, the mean value of $\cos|(w_{\alpha})'_{y}, X'_{y}|$ is greater than c(b) > 2a, a contradiction. Thus, a point w(y) on a shortest path $w_{\alpha}y$ sufficiently close to y satisfies the condition (4) of Definition 5.1 for $(f, |x \cdot |)$ at y.

Lemma 5.10 (cf. [8, 3.4 Assertion 2]). Under Setting 5.8, let $x, y \in V$ be such that $|f(x)f(y)| < \delta |xy|$ and assume that x is a local maximum point of $\sigma|_{f^{-1}(f(x))}$. Then, there exists α such that $\tilde{\angle} w_{\alpha} y x > \pi/2 + c(\varepsilon)$.

Proof. It suffices to show that $|(W_{\alpha})'_{x}, y'_{x}| < \pi/2 - c(\varepsilon)$ for some α , where $(W_{\alpha})'_{x}$ denotes the set of all directions of shortest paths from x to w_{α} . Let $a = c(\varepsilon)$ and $b = c(\varepsilon)$ be constants such that $a \ll b \ll L$. Suppose $|(W_{\alpha})'_{x}, y'_{x}| \ge \pi/2 - a$ for all α . Let \mathcal{A}_{1} (resp. \mathcal{A}_{2}) be the set of indices α such that $|(W_{\alpha})'_{x}, y'_{x}| \ge \pi/2 + a$ (resp. $> \pi/2 + a$). Set $W'(x) := \bigcup_{\alpha \in \mathcal{A}_{2}} (W_{\alpha})'_{x}, A'_{i}(x) := \bigcup_{\gamma \in \Gamma_{i}(p)} (\mathcal{A}_{i\gamma})'_{x}$. Note that $|\mathcal{A}'_{i}(x), y'_{x}| > \pi/2 - \varkappa(\delta)$ by Remark 5.6(2). Then, applying Lemma 3.7(1) to $W'(x), y'_{x}, A'_{1}(x), \ldots, A'_{k}(x)$, we get a direction $\xi \in \Sigma_{x}$ such that

 $|W'(x)\xi| > \pi/2 + c(a), \quad |A'_i(x)\xi| = \pi/2$

for all *i*. Let \mathcal{A}_3 be the set of indices $\alpha \in \mathcal{A}_2$ such that $|(W_{\alpha})'_x \xi| > \pi/2 + b$. Note that $\#\mathcal{A}_1 \leq Ca/\omega^{n-1}$ and $\#\mathcal{A}_3 \geq (L - Ca - Cb)/\omega^{n-1}$ (see Lemma 3.8). Choose x_1 near x such that $(x_1)'_x$ is close to ξ . Then, we have

$$\sigma(x_1) \ge \sigma(x) + N^{-1} (\#\mathcal{A}_3 \cdot c(b) - \#\mathcal{A}_1) |xx_1|$$

$$\ge \sigma(x) + N^{-1} \omega^{-(n-1)} ((L - Ca - Cb) \cdot c(b) - Ca) |xx_1|$$

$$\ge \sigma(x) + c(\varepsilon) |xx_1|$$

since $a \ll b \ll L$ and $N \leq C/\omega^{n-1}$. Furthermore, by Remark 5.6(1), we have $|f(x_1)f(x)| < \varkappa(\delta)|xx_1|$. Thus, by the $c(\varepsilon)$ -openness of f, we can find $x_2 \in f^{-1}(f(x))$ near x such that $c(\varepsilon)|x_1x_2| \leq \varkappa(\delta)|xx_1|$ (cf. [8, 2.1.3]). Therefore, we have

$$\sigma(x_2) \ge \sigma(x_1) - |x_1 x_2| \ge \sigma(x) + c(\varepsilon)|x x_1| - c(\varepsilon)^{-1} \varkappa(\delta)|x x_1| > \sigma(x).$$

This contradicts the local maximality of $\sigma|_{f^{-1}(f(x))}$ at x.

Let $f : U \to \mathbb{R}^k$ be (ε, δ) -noncritical in the generalized sense at $p \in U$. We say that f is (ε, δ) -complementable (in the generalized sense) at p if there exists a function f_{k+1} defined on a neighborhood of p such that (f, f_{k+1}) is (ε, δ) -noncritical in the generalized sense at p.

The next proposition is the key to the proof of the fibration theorem. For $v \in \mathbb{R}^k$ and r > 0, we denote by $I^k(v, r)$ the closed *r*-neighborhood of v in \mathbb{R}^k with respect to the maximum norm (recall that we use the maximum norm in this section).

Proposition 5.11 (cf. [8, 3.5]). Let $f: U \to \mathbb{R}^k$ be (ε, δ) -noncritical in the generalized sense at $p \in U$ and assume $\operatorname{vol}_{n-1} \Sigma_p \geq \varepsilon$. Suppose that f is not $(c(\varepsilon), \varkappa(\delta))$ -complementable at p. Then, for sufficiently small R > 0, there exists a function h defined on $U_1 = f^{-1}(I^k(f(p), \delta^5 R)) \cap \overline{B}(p, R)$ such that

- (1) $h(U_1) = [0, R]$ and h(x) = |px| if |px| > R/2;
- (2) f is injective on $S = h^{-1}(0)$;
- (3) f is $(c(\varepsilon), \varkappa(\delta))$ -complementable at any point of $U_1 \setminus S$;
- (4) (f,h) is $(c(\varepsilon,\varkappa(\delta))$ -noncritical in the generalized sense at any point $x \in U_1$ such that $|f(x)f(S)| < \frac{1}{3}\delta^5 h(x)$.

Proof. The proof is completely the same as the original one. Let R > 0 be so small that U_1 is contained in the neighborhood V of Setting 5.8. Let $M = c(\varepsilon)$ be the constant of Lemma 5.9(2) and define a compact set S by

$$S := \left\{ x \in U_1 \mid \begin{array}{c} \sigma(x) - \sigma(y) \ge M |xy| \text{ for all } y \in U_1 \\ \text{satisfying } |f(x)f(y)| < \delta |xy| \end{array} \right\}$$

Then, f is $(c(\varepsilon), \varkappa(\delta))$ -complementable at any point of $U_1 \setminus S$ by Lemma 5.9. In particular, $p \in S$ by the assumption. Furthermore, the $c(\varepsilon)$ -openness of f implies

that $|f(x)f(y)| \geq M_1|xy|$ for any $x, y \in S$, where $M_1 = c(\varepsilon)$. (Indeed, assume f is M_2 -open, where $M_2 = c(\varepsilon)$, and let $M_1 \ll MM_2$. Suppose that there exist $x, y \in S$ such that $|f(x)f(y)| < M_1|xy|$. Then, we can find $x_1 \in f^{-1}(f(x))$ such that $|x_1y| \ll M|xy|$. Together with the definition of S, this implies $\sigma(x) > \sigma(y)$. Similarly $\sigma(x) < \sigma(y)$, a contradiction.) In particular, f is injective on S and $S \subset \overline{B}(p, M_1^{-1}\delta^5R)$.

Let us define the function h. Choose a sequence S_j of finite subsets of S such that

- $S_0 := \{p\};$
- $S_j \supset S_{j-1}$ and $f(S_j)$ is a maximal $\delta^{j+5}R$ -discrete net in f(S). Then, we define $h(x) := \inf\{h_\gamma(x) \mid p_\gamma \in \bigcup_j S_j\}$ by

$$h_{\gamma}(x) := \varphi_{\delta^{j+1}R}(|p_{\gamma}x|) + \sum_{l=1}^{k} \frac{10}{M_{1}} |f_{l}(x) - f_{l}(p_{\gamma})|$$

for $p_{\gamma} \in S_j \setminus S_{j-1}$ $(j \ge 1)$ and

$$h_{\gamma}(x) := \min\left\{\varphi_{\delta R}(|px|) + \sum_{l=1}^{k} \frac{10}{M_{1}}|f_{l}(x) - f_{l}(p)|, \ \frac{1}{2}\varphi_{\frac{R}{2}}(|px|) + \frac{R}{4}\right\}$$

for $p_{\gamma} = p$, where

$$\varphi_r(a) := \begin{cases} a, & a \le r \\ 2a - r, & a \ge r. \end{cases}$$

Note that the above definition of h is exactly the same as in the original proof and that it has the same form as the function q_i in Definition 5.1(1).

Then, it easily follows that $h^{-1}(0) = S$, $h(U_1) = [0, R]$ and h(x) = |px| if |px| > R/2. To verify the conclusion (4), we need the following lemma.

Lemma 5.12 ([8, 3.5 Assertion 3]). For $x \in U_1 \setminus S$, set $\Gamma(x) := \{\gamma \mid h_{\gamma}(x) = h(x)\}$. Suppose $|f(x)f(S)| < \frac{1}{3}\delta^5h(x)$. Then, there exists $s \in S$ such that

$$|f(x)f(s)| < \delta^2 |xs|, \quad |p_\gamma s| < \delta |xs|$$

for all $\gamma \in \Gamma(x)$. Moreover, we have $\#\Gamma(x) < c^{-1}(\varepsilon)$.

We omit the proof since it is based only on the definition of h and the M_1 -co-Lipschitzness of f on S, and does not depend on the definition of noncriticality (see the original proof).

Let us show the conclusion (4). Let $x \in U_1 \setminus S$ and $s \in S$ be as above. Then, by Lemma 5.10, there exists α such that $\tilde{\angle} w_{\alpha} xs > \pi/2 + c(\varepsilon)$. Together with the second inequality in the above lemma, this implies $\tilde{\angle} w_{\alpha} xp_{\gamma} > \pi/2 + c(\varepsilon)$ for all $\gamma \in \Gamma(x)$. Furthermore, the two inequalities in the above lemma together with Remark 5.6(2) imply that $\tilde{\angle} A_{i\beta} xp_{\gamma} > \pi/2 - \varkappa(\delta)$ for all $\beta \in \Gamma_i(x)$ and $\gamma \in \Gamma(x)$. Thus, all the conditions of Definition 5.1 are satisfied for (f, h) at x.

The last proposition is a refinement of the previous one on each fiber.

Proposition 5.13 (cf. [8, 3.6]). Let $f: U \to \mathbb{R}^k$ be (ε, δ) -noncritical in the generalized sense at $p \in U$ and assume $\operatorname{vol}_{n-1} \Sigma_p \geq \varepsilon$. Let R > 0 be so small that $U_1 = f^{-1}(I^k(f(p), \delta^5 R)) \cap \overline{B}(p, R)$ is contained in the neighborhood V of Setting 5.8. Suppose that

• for any $x \in U_1$ such that $\delta R \leq |px| \leq R$, we have $\sigma(p) - \sigma(x) \geq M|px|$, where $M = c(\varepsilon)$ is the constant of Lemma 5.9(2) (in particular, this is weaker than the condition that f is not $(c(\varepsilon), \varkappa(\delta))$ -complementable at p). Then, for any $v \in I^k(f(p), \delta^5 R)$, there exists a function $h_v : U_1 \to [0, R]$ and a point $o_v \in U_1 \cap f^{-1}(v)$ such that

- (1) for $x \in U_1 \cap f^{-1}(v)$, we have $h_v(x) = R \Leftrightarrow |px| = R$ and $h_v(x) = 0 \Leftrightarrow x = o_v$:
- (2) (f, h_v) is $(c(\varepsilon), \varkappa(\delta))$ -noncritical in the generalized sense on $U_1 \cap f^{-1}(v) \setminus \{o_v\}$.

Proof. The proof is completely the same as the original one. Let o_v be a maximum point of $\sigma|_{U_1 \cap f^{-1}(v)}$. Then, by the assumption and the $c(\varepsilon)$ -openness of f, we have $|po_v| < \delta R$. (Indeed, suppose the contrary; then we have $\sigma(p) \ge \sigma(o_v) + M|po_v|$. On the other hand, we can find $q \in f^{-1}(v)$ such that $|pq| \ll |po_v|$. Thus, we have $\sigma(q) > \sigma(o_v)$, a contradiction.) Define

$$h_v(x) := \min\left\{\varphi_{\delta R}(|o_v x|), \ \frac{1}{2}\varphi_{\frac{R}{2}}(|px|) + \frac{R}{4}\right\},\$$

where $\varphi_r(a) = \max\{a, 2a - r\}$ as before. Then, (1) is clear. Let us show (2). Let $x \in U_1 \cap f^{-1}(v) \setminus \{o_v\}$. Then, $(f, |o_v \cdot|)$ is $(c(\varepsilon), \varkappa(\delta))$ -noncritical at x by Lemma 5.10 and Remark 5.6(2). Similarly, $(f, |p \cdot|)$ is $(c(\varepsilon), \varkappa(\delta))$ -noncritical at x if $|px| \ge \delta R$ by the assumption and Lemma 5.9. On the other hand, $\varphi_{\delta R}(|o_v x|) < \frac{1}{2}\varphi_{\frac{R}{2}}(|px|) + \frac{R}{4}$ if $|px| < \delta R$ since $|po_v| < \delta R$. Thus, (f, h_v) is $(c(\varepsilon), \varkappa(\delta))$ -noncritical at x.

We have now shown all the propositions in [8, §3] for generalized noncritical maps. Finally, we define the term "noncritical (in the generalized sense)" (not " (ε, δ) -noncritical") as follows:

Definition 5.14 (cf. [8, 3.7]). Let U be a domain in an n-dimensional Alexandrov space M such that $\varepsilon_0 = \inf_{p \in U} \operatorname{vol}_{n-1} \Sigma_p > 0$. A map $f: U \to \mathbb{R}^k$ $(0 \le k \le n+1)$ is said to be *noncritical (in the generalized sense)* at p if it is (ε, δ) -noncritical in the generalized sense at p for some $\varepsilon < \varepsilon_0$ and $\delta < \Delta_{n,k}(\varepsilon)$, where $\Delta_{n,k}(\varepsilon)$ is defined by reverse induction on k in such a way that (ε, δ) -noncritical maps $f: U \to \mathbb{R}^k$ with $\varepsilon < \varepsilon_0$ and $\delta < \Delta_{n,k}(\varepsilon)$ satisfy all the propositions above (including lemmas) and the pairs $(c(\varepsilon), \varkappa(\delta))$ appearing in them satisfy $\varkappa(\delta) < \Delta_{n,k+1}(c(\varepsilon))$.

Then, all the properties of noncritical maps listed in [8, 1.3] are also true for this definition. As pointed out at the beginning of this discussion, the remaining topological part [8, 1.4–1.5] of the proof of the fibration theorem is based only on those properties. Thus, the generalized fibration theorem follows in exactly the same way. This completes the proof of Theorem 5.2.

Proof of Theorem 1.2 and Corollary 1.3. Theorem 1.2 follows from Theorem 4.1 and Theorem 5.2. Indeed, the map $\varphi \circ f$ in Theorem 4.1 is $(c, \varkappa(\delta))$ -noncritical in the generalized sense on $B(\hat{p}, \ell \delta^2)$ (we can find w of Definition 5.1(4) by Proposition 3.5). Note that $(\varphi \circ f)^{-1}(B(\varphi(p), \ell \delta^2/2))$ has a compact closure in $B(\hat{p}, \ell \delta^2)$, since f is $C\mu$ -close to the μ -approximation g, where $\mu \ll \ell \delta^2$ (we use the standard Euclidean metric here). Therefore, by Theorem 5.2, it is homeomorphic to the product $f^{-1}(p) \times B(\varphi(p), \ell \delta^2/2)$, where the second component is given by $\varphi \circ f$. Moreover, Corollary 1.3 follows from Lemma 3.3 and Remark 3.4. Indeed, if k = n-1, then $\operatorname{vol}_{n-1} \Sigma_p > c$ for any $p \in M$. It easily follows from Proposition 5.7 and the topological part [8, 1.4-1.5] of the proof of the fibration theorem that the fiber is homeomorphic to a circle or a closed interval. \square

Problem 5.15. Is it possible to prove the fibration theorem for noncritical maps without the assumption on the volume of spaces of directions? If possible, then probably the map f of Theorem 4.1 is a locally trivial fibration.

Remark 5.16. A similar modification does not work for Perelman's another proof of the fibration theorem in [9], which requires no assumptions on the volume of spaces of directions. This is because the regularity of maps defined in [9] is stronger than the above noncriticality in that it does not include an error δ as in Definition 5.1. For example, a map $f = (|A_1 \cdot |, \dots, |A_k \cdot |) : M \to \mathbb{R}^k$, where A_i are compact subsets of M, is called ε -regular at $p \in M$ if it satisfies

- (1) $\angle ((A_i)'_p, (A_j)'_p) > \pi/2 + \varepsilon$ for all $i \neq j$; (2) there exists a direction $\xi \in \Sigma_p$ such that $\angle ((A_i)'_p, \xi) > \pi/2 + \varepsilon$ for all i.

In addition, f is simply called *regular* at p if it is ε -regular at p for some $\varepsilon > 0$. Then, the fibration theorem in [9] states that every proper regular map defined on a domain of an Alexandrov space is a locally trivial fibration. However, we cannot weaken the condition (1) to $\angle ((A_i)'_p, (A_j)'_p) > \pi/2 - \delta$ for any $\delta \ll \varepsilon$. The reason is as follows. There is a counterpart [9, 1.3] of Propositions 5.11, 5.13 asserting that if $g: M \to \mathbb{R}^k$ is regular and incomplementable at p, then there exists a nonpositive function g_{k+1} defined on a neighborhood of p such that (g, g_{k+1}) is regular on the complement of $g_{k+1}^{-1}(0)$. Nevertheless, unlike Propositions 5.11, 5.13, even if the given map g is ε -regular at p, the new map (g, g_{k+1}) is not uniformly $c(\varepsilon)$ -regular. Thus, if we modify the condition (1) as above, the given error δ may be too big compared with the regularity of the new map (g, g_{k+1}) . The same problem occurs if we try to generalize the definition of regularity by using inequality (4.1) like Definition 5.1.

We conclude this section with the following stability theorem for fibrations. Note that the stability theorem [8, 4.3] for framed subsets can be generalized like the fibration theorem just by replacing "noncritical map" with "noncritical map in the generalized sense." Indeed, the proof of the stability theorem is based only on the properties of noncritical maps established in $[8, \S3]$ like the topological part of the proof of the fibration theorem.

Corollary 5.17. Let X, M be Alexandrov spaces as in Theorem 1.2 and suppose in addition that they are compact. Let M be an n-dimensional Alexandrov space sufficiently close to M (in particular, M also satisfies the assumption of Theorem 1.2; see [2, 7.14]). Let $f: M \to X$ and $\tilde{f}: \tilde{M} \to X$ be the fibrations constructed in Theorem 4.1 (where the approximation $\tilde{g}: \tilde{M} \to X$ is assumed to be obtained from the approximations $g: M \to X$ and $\Psi: M \to \tilde{M}$). Then, there exists a homeomorphism $\Phi: M \to \tilde{M}$ close to Ψ respecting f, that is, $f = \tilde{f} \circ \Phi$.

Proof. We give only an outline. Let $\varphi: X \to \mathbb{R}^k$ be the distance coordinate around $p \in X$ as in Theorem 4.1. Then, $\varphi \circ f$ and $\varphi \circ \tilde{f}$ are noncritical maps in the generalized sense on some neighborhoods of lifts of p in M and \tilde{M} , respectively. By the generalized fibration theorem, $(\varphi \circ f)^{-1}(I^k(\varphi(p), \rho))$ is homeomorphic to $f^{-1}(p) \times I^k(\varphi(p), \rho)$ for sufficiently small $\rho > 0$, where the second component is given by $\varphi \circ f$ and the fiber $f^{-1}(p)$ is an MCS-space in the sense of [8, 1.1].

Furthermore, by the generalized stability theorem mentioned above, there exists a homeomorphism $\Phi_p : (\varphi \circ f)^{-1}(I^k(\varphi(p),\rho)) \to (\varphi \circ \tilde{f})^{-1}(I^k(\varphi(p),\rho))$ close to Ψ respecting $\varphi \circ f$, provided that \tilde{M} is sufficiently close to M. Take a finite cover of M by such product neighborhoods with respect to f. Then, the desired homeomorphism is constructed by the same gluing argument as in the proof of Complement to Theorem B in [8, §1]. \Box

6. PROPERTIES OF THE FIBERS

In this section, we study the properties of the fibers of the map f of Theorem 4.1. Note that the assumption on the volume of spaces of directions as in Theorem 1.2 is not required here. We first remark that the diameters of the fibers of f are very small:

Remark 6.1. Let $f : M \to X$ be the map of Theorem 4.1 and $p \in X$. Since f is $C\mu$ -close to the μ -approximation g, the fiber $f^{-1}(p)$ is contained in the $C\mu$ -neighborhood of a lift $\hat{p} \in M$ of p. Note that $\mu < c(\ell\delta^2) \ll \ell\delta^2$.

In view of the inequality (4.1), we mainly deal with the following class of maps in this section:

Definition 6.2. Let M be an Alexandrov space and $p \in M$. Let $\{(a_i, b_i)\}_{i=1}^k$ be a (k, δ) -strainer at p with length $> \ell$ and $g : M \to \mathbb{R}^k$ its distance coordinate. Suppose a map $f : B(p, \ell) \to \mathbb{R}^k$ satisfies

(6.1)
$$|(f(x) - f(y)) - (g(x) - g(y))| < \delta |xy|$$

for any $x, y \in B(p, \ell)$. Then, we call f a δ -almost regular map associated with the strainer $\{(a_i, b_i)\}_{i=1}^k$.

While the domain of f may seem to be too large, it is useful for simplicity and is sufficient for our applications.

Remark 6.3. The above definition of an almost regular map is different from that in [2, 11.7] (the original definition generalizes the distance coordinate of a strainer from a different point of view). However, as in the case of noncritical maps, all the claims about almost regular maps in [2] hold for the above ones. This is why we use the same term.

Remark 6.4. Let $f: M \to X$ be the map of Theorem 4.1 and let $p, \hat{p}, \varphi, \hat{\varphi}$ be as in Theorem 4.1. Then, by the inequality (4.1), $\varphi \circ f$ is a $\varkappa(\delta)$ -almost regular map on $B(\hat{p}, \ell\delta^2)$ associated with a $(k, \varkappa(\delta))$ -strainer at \hat{p} with length $> \ell\delta^2$.

The following proposition is a generalization of Proposition 3.5:

Proposition 6.5. Let M be an n-dimensional Alexandrov space and $p \in M$. Let $f : B(p, \ell) \to \mathbb{R}^k$ be a δ -almost regular map associated with a (k, δ) -strainer at p with length $> \ell$. Then,

- (1) if k = n, then f is a $\varkappa(\delta)$ -almost isometry from $B(p, \ell\delta)$ to an open subset of \mathbb{R}^n ;
- (2) if k < n, then f is a $(1 + \varkappa(\delta))$ -Lipschitz and $(1 \varkappa(\delta))$ -open map on $B(p, \ell\delta)$.

The proof is an easy application of the inequality (6.1) (the $(1 - \varkappa(\delta))$ -openness follows from [3, 3.1] since the distance coordinate g satisfies the assumption of [3, 3.1] on $B(p, \ell\delta)$ and so does f).

In particular, by Remark 6.4, the map f of Theorem 4.1 is a $\varkappa(\delta)$ -almost isometry onto X when k = n, and is a $(1 + \varkappa(\delta))$ -Lipschitz and $(1 - \varkappa(\delta))$ -open map when k < n (note that f is $C\mu$ -close to the μ -approximation g globally, where $\mu \ll \ell \delta^2$).

6.1. Intrinsic metric of the fibers. In this section, we prove Theorem 1.4(1). We first show that an almost regular map is an almost Lipschitz submersion near the strained point. Recall that for a subset A of an Alexandrov space and $p \in A$, the space of directions of A at p is defined as the subset of Σ_p consisting of all limit points $\lim_{i\to\infty} (p_i)'_p$, where $p_i \in A$ converges to p.

Proposition 6.6 (cf. [17]). Let M be an n-dimensional Alexandrov space, $p \in M$ and k < n. Let $f : B(p, \ell) \to \mathbb{R}^k$ be a δ -almost regular map associated with a (k, δ) -strainer $\{(a_i, b_i)\}_{i=1}^k$ at p with length $> \ell$. Then, f is a $\varkappa(\delta)$ -almost Lipschitz submersion on $B(p, \ell\delta)$ in the following sense:

$$\left|\frac{|f(x)f(y)|}{|xy|} - \sin\angle(y'_x, V_x)\right| < \varkappa(\delta),$$

for any $x, y \in B(p, \ell \delta)$ and any direction y'_x , where V_x denotes the space of directions of $f^{-1}(f(x))$ at x (and is nonempty).

In particular, by Remark 6.4, the map f of Theorem 4.1 is locally (indeed, globally) a $\varkappa(\delta)$ -almost Lipschitz submersion when k < n.

Remark 6.7. The above definition of an almost Lipschitz submersion is slightly stronger than that in [17] (see Theorem 1.1).

Proof of Proposition 6.6. Let $x, y \in B(p, \ell \delta)$. Then, by Lemma 3.2 and the inequality (6.1), we have

$$\left|\frac{|f(x)f(y)|^2}{|xy|^2} - \sum_{i=1}^k \cos^2 \angle a_i xy\right| < \varkappa(\delta).$$

By Lemma 3.3, there exists a $\varkappa(\delta)$ -approximation from Σ_x to a k-fold spherical suspension $S^k(\Sigma)$, where Σ has curvature ≥ 1 and is nonempty (see Remark 3.4). Let $\tilde{\Sigma} \subset \Sigma_x$ be a subset corresponding to $\Sigma \subset S^k(\Sigma)$ via this approximation. Then, we have

$$\left|\sum_{i=1}^{k} \cos^{2} \angle a_{i} x y - \sin^{2} \angle (y'_{x}, \tilde{\Sigma})\right| < \varkappa(\delta).$$

Thus, it suffices to show that the Hausdorff distance between $\tilde{\Sigma}$ and V_x is less than $\varkappa(\delta)$. The above inequalities immediately imply that V_x is contained in the $\varkappa(\delta)$ -neighborhood of $\tilde{\Sigma}$. On the other hand, let $\xi \in \tilde{\Sigma}$. Choose a point z near x such that z'_x is sufficiently close to ξ . Then, the above inequalities imply that $|f(x)f(z)| < \varkappa(\delta)|xz|$. By the $(1 - \varkappa(\delta))$ -openness of f, we can find $w \in f^{-1}(f(x))$ such that $(1 - \varkappa(\delta))|wz| \leq |f(x)f(z)|$. In particular, we have $|wz| < \varkappa(\delta)|xz|$. Since z can be chosen arbitrarily close to x, we obtain $\angle zxw < \varkappa(\delta)$. This completes the proof.

Now, we prove Theorem 1.4(1). By Remarks 6.1 and 6.4, it suffices to show the following:

Corollary 6.8 (cf. [2, 11.11]). Let M be an n-dimensional Alexandrov space, $p \in M$ and k < n. Let $f : B(p, \ell) \to \mathbb{R}^k$ be a δ -almost regular map associated with a (k, δ) strainer at p with length $> \ell$. Then, for any $x, y \in f^{-1}(f(p)) \cap B(p, \ell\delta)$, there exists a curve in $f^{-1}(f(p))$ connecting x and y of length $< (1 + \varkappa(\delta))|xy|$.

Proof. By Proposition 6.6, for any $x, y \in f^{-1}(f(p)) \cap B(p, \ell\delta)$, there exists $z \in f^{-1}(f(p))$ arbitrarily close to x such that $\angle yxz < \varkappa(\delta)$. In particular, the first variation formula implies that $|yz| < |yx| - (1 - \varkappa(\delta))|xz|$. Thus, the desired curve is obtained by taking a limit of broken geodesics.

6.2. Lower bound for the volume of the fibers. In this section, we prove the left inequality of Theorem 1.4(2). By Remark 6.4, it suffices to show the following:

Proposition 6.9. Let M be an n-dimensional Alexandrov space, $p \in M$ and k < n. Let $f : B(p, \ell) \to \mathbb{R}^k$ be a δ -almost regular map associated with a (k, δ) -strainer $\{(a_i, b_i)\}_{i=1}^k$ at p with length $> \ell$. Suppose $\operatorname{vol}_n(B(p, D)) > v$. Then, we have

 $\operatorname{vol}_{n-k} f^{-1}(f(p)) > c(D, v, \ell).$

Remark 6.10. Since f is $(1 - \varkappa(\delta))$ -open on $B(p, \ell\delta)$, we have $\operatorname{vol}_{n-k} f^{-1}(u) > c(D, v, \ell)$ for any $u \in B(f(p), \ell\delta/2)$

From now, we fix sufficiently small δ depending only on n and κ (we determine it later). We first give a lower bound for the diameter of the fiber:

Lemma 6.11. Under the same assumption as Proposition 6.9, there exists a point $q \in f^{-1}(f(p))$ such that $\rho < |pq| < \ell \delta$, where $\rho = c(D, v, \ell)$.

Proof. We argue by contradiction. Suppose that there exists a sequence of *n*-dimensional Alexandrov spaces (M_j, p_j) with vol_n $B(p_j, D) > v$ and δ-almost regular maps $f_j : B(p_j, \ell) \to \mathbb{R}^k$ associated with (k, δ) -strainers $\{(a_i^j, b_j^j)\}_{i=1}^k$ at p_j with lengths > ℓ such that diam $f_j^{-1}(f_j(p_j)) \cap B(p_j, \ell \delta) \to 0$. For simplicity, we assume that $|p_j a_i^j|$, $|p_j b_i^j|$ are uniformly bounded above. Since vol_n $B(p_j, D) > v$, we may assume that (M_j, p_j) converges to an Alexandrov space (M, p) of dimension *n* (note that it is different from (M, p) in the statement of Lemma 6.11; so is *q* below). Furthermore, since lengths > ℓ , we may assume that $\{(a_i^i, b_i^j)\}_{i=1}^k$ at *p*. Then, by Lemma 3.3, Σ_p is $\varkappa(\delta)$ -close to a *k*-fold spherical suspension $S^k(\Sigma)$, where Σ is a space of curvature ≥ 1. Notice that Σ is nonempty since $k < n = \dim M$ (see Remark 3.4). Therefore, there exists $q \in M$ near *p* such that $|g(p)g(q)| < \varkappa(\delta)|pq|$, where *g* denotes the distance coordinate associated with the strainer $\{(a_i^j, b_i^j)\}_{i=1}^k$. By the inequality (6.1), we have $|f_j(p_j)f_j(q_j)| < \varkappa(\delta)|p_jq_j|$. Hence, by the $(1 - \varkappa(\delta))$ -openness, we can find $\hat{q}_j \in f_j^{-1}(f_j(p_j))$ such that $|\hat{q}_jq_j| < \varkappa(\delta)|p_jq_j|$. In particular, we have $|p_j\hat{q}_j| > |pq|/2$ if δ is small enough. This contradicts the assumption that diam $f_j^{-1}(f_j(p_j)) \cap B(p_j, \ell\delta) \to 0$.

Proof of Proposition 6.9. Let q be as in Lemma 6.11 and let r be the midpoint of a shortest path connecting p and q. Put $a_{k+1} := p$ and $b_{k+1} := q$. Then, by the inequality (6.1) and Lemma 3.2, we see that $\{(a_i, b_i)\}_{i=1}^{k+1}$ is a $(k+1, \varkappa(\delta))$ -strainer at r with length $> \rho/2$. Furthermore, we have $|f(p)f(r)| < \varkappa(\delta)|pr|$. Thus, by

20

the $(1 - \varkappa(\delta))$ -openness, we can find $s \in f^{-1}(f(p))$ such that $|ps| < \varkappa(\delta)|pr|$. Then, $\{(a_i, b_i)\}_{i=1}^{k+1}$ is also a $(k+1, \varkappa(\delta))$ -strainer for s with length $> \rho/3$. Hence, $(f, |a_{k+1} \cdot |)$ is a $\varkappa(\delta)$ -almost regular map around s. Repeating this argument (n-k)-times, we get $\hat{p} \in f^{-1}(f(p))$ and $h = (|a_{k+1} \cdot |, \ldots, |a_n \cdot |)$ such that (f, h) is a $\varkappa(\delta)$ -almost regular map associated with a $(n, \varkappa(\delta))$ -strainer at \hat{p} with length $> \hat{\ell} = c(D, v, \ell)$. Thus, by Proposition 6.5(2), (f, h) is a $\varkappa(\delta)$ -almost isometry from $B(\hat{p}, \hat{\ell}\delta)$ to an open subset of \mathbb{R}^n . Therefore, the restriction of h to $f^{-1}(f(p))$ gives a $\varkappa(\delta)$ -almost isometry from a neighborhood of \hat{p} in $f^{-1}(f(p))$ to an $(\hat{\ell}\delta/2)$ -ball in \mathbb{R}^{n-k} . This completes the proof (fix small δ such that the last $\varkappa(\delta)$ is less than 1/2).

Remark 6.12. Let $f: B(p, \ell) \to \mathbb{R}^k$ be a δ -almost regular map associated with a (k, δ) -strainer at p with length $> \ell$. Set $F := f^{-1}(f(p)) \cap B(p, \ell \delta)$. Suppose that $x \in F$ has an $(n - k, \delta)$ -strainer $\{(a_{k+i}, b_{k+i})\}_{i=1}^{n-k}$ such that $a_{k+i}, b_{k+i} \in F$. Then, the same argument as above shows that the restriction of the distance coordinate of this strainer to F gives a $\varkappa(\delta)$ -almost isometry from a neighborhood of x in F to an open subset in \mathbb{R}^{n-k} . We can also prove that the complement of the set of all such points x in F has Hausdorff dimension at most n - k - 1. The proof is similar to that of [2, 10,6]. See also Lemma 6.25.

6.3. Upper bound for the volume of the fibers. In this section, we prove the right inequality of Theorem 1.4(2). The proof is based on the theory of noncritical maps by Perelman [8] and the rescaling technique for collapsing sequences by Ya-maguchi [18]. We always assume that δ is much smaller than ε and $c(\varepsilon)$ and that so is $\varkappa(\delta)$ (see §2).

We consider the following regularity of maps in this section:

Definition 6.13. Let $f: M \to \mathbb{R}^k$ be a map defined on an Alexandrov space M. For positive numbers ε , δ and ρ , we say that f is $(\varepsilon, \delta, \rho)$ -noncritical at $p \in M$ if there exists a map $g = (|a_1 \cdot |, \ldots, |a_k \cdot |) : M \to \mathbb{R}^k$, where $a_i \in M$, such that

- (1) $|(f(x) f(y)) (g(x) g(y))| < \delta |xy|$ for any $x, y \in B(p, \rho)$;
- (2) $|a_i p| > \rho$ and $\tilde{\angle} a_i p a_{i'} > \pi/2 \delta$ for all $i \neq i'$;
- (3) there exists $w \in M$ such that $|wp| > \rho$ and $\tilde{\angle}a_i pw > \pi/2 + \varepsilon$ for all *i*.

Remark 6.14. This definition is a special case of Definition 5.1 (except for the existence of ρ). In particular, if f is $(\varepsilon, \delta, \rho)$ -noncritical at p, then $k \leq n$ and it is $c(\varepsilon)$ -open on $B(p, \rho\delta)$ (see Proposition 5.4).

Remark 6.15. Let $f : B(p, \ell) \to \mathbb{R}^k$ be a δ -almost regular map associated with a (k, δ) -strainer $\{(a_i, b_i)\}_{i=1}^k$ at p with length $> \ell$. Then, it is $(c, \delta, c\ell)$ -noncritical at p. Indeed, we can find $w \in M$ such that $|wp| > c\ell$ and $\tilde{\angle}a_i pw > \pi/2 + c$ by Proposition 3.5.

For a (compact) metric space X, we denote by $\beta_{\nu}(X)$ the maximal possible number of ν -discrete points in X. Furthermore, for simplicity, we introduce a notation

$$v_m(X) := \sup_{0 < \nu < 1} \nu^m \beta_\nu(X)$$

for $m \geq 0$. Clearly, $\operatorname{vol}_m(X) \leq C(m) \cdot v_m(X)$, where vol_m denotes the *m*-dimensional Hausdorff measure and C(m) is a constant depending only on *m*.

We prove the following proposition in this section:

Proposition 6.16. Let M be an n-dimensional Alexandrov space and $k \leq n$. For a map $f : M \to \mathbb{R}^k$ and $p \in M$, set $F := f^{-1}(f(p))$. Let $F(\varepsilon, \delta, \rho)$ be the set of all $(\varepsilon, \delta, \rho)$ -noncritical points of f in F. Then, for any sufficiently small $\delta > 0$ (depending only on $n, \kappa, D, \varepsilon$ and ρ), we have

$$v_{n-k}(F(\varepsilon,\delta,\rho)\cap B(p,D)) < C(D,\varepsilon,\rho).$$

In particular,

$$\operatorname{vol}_{n-k}(F(\varepsilon,\delta,\rho) \cap B(p,D)) < C(D,\varepsilon,\rho).$$

Proof of the right inequality of Theorem 1.4(2). Let $f : M \to X$ be the map of Theorem 4.1 and let $p, \hat{p}, \varphi, \hat{\varphi}$ be as in Theorem 4.1. Then, by the inequality (4.1) and Remark 6.15, $\varphi \circ f$ is $(c, \varkappa(\delta), c\ell\delta^2/2)$ -noncritical on $B(\hat{p}, \ell\delta^2/2)$. Furthermore, the diameter of the fiber $f^{-1}(p)$ is less than $C\mu$, where $\mu \ll \ell\delta^2$ (see Remark 6.1). In particular, every point of $f^{-1}(p)$ is a $(c, \varkappa(\delta), c\ell\delta^2/2)$ -noncritical point of $\varphi \circ f$. Thus, rescaling M by the reciprocal of the diameter of $f^{-1}(p)$ and applying Proposition 6.16, we obtain the desired estimate. \Box

Proof of Proposition 6.16. We prove it by reverse induction on k. First, consider the base case k = n. The following argument is the same as the second half of the proof of Proposition 5.4. Let $x \in F(\varepsilon, \delta, \rho)$ and let $a_i, w \in M$ be as in Definition 6.13. Suppose there exists $y \in B(x, \rho\delta)$ such that f(x) = f(y) and $x \neq y$. If $|wx| \geq |wy|$, then by the definition of $(\varepsilon, \delta, \rho)$ -noncriticality, we have

$$\begin{split} \tilde{\angle} a_i y a_{i'} > \pi/2 - \varkappa(\delta), \quad \tilde{\angle} a_i y w > \pi/2 + c(\varepsilon), \\ \tilde{\angle} a_i y x > \pi/2 - \varkappa(\delta), \quad \tilde{\angle} w y x > \pi/2 - \varkappa(\delta) \end{split}$$

for all $1 \leq i \neq i' \leq n$. This contradicts Lemma 3.6 for Σ_y . We also get a contradiction when $|wx| \leq |wy|$. Therefore, the set $F(\varepsilon, \delta, \rho)$ is $\rho\delta$ -discrete. In particular, the cardinality of $F(\varepsilon, \delta, \rho) \cap B(p, D)$ is bounded above by some constant $C(D, \varepsilon, \rho)$ (note that we can take $\delta = c(\varepsilon)$ so that Lemma 3.6 holds in the above argument).

Next, consider the case k < n. We argue by contradiction. Thus, it is sufficient to prove the following:

Proposition 6.17. Suppose a sequence (M_j, p_j) of n-dimensional Alexandrov spaces converges to an l-dimensional Alexandrov space (X, p) in the pointed Gromov-Hausdorff topology. Let $f_j : M_j \to \mathbb{R}^k$ and $F_j := f_j^{-1}(f_j(p_j))$. Then, for any positive numbers D, ε, ρ and $\delta_j \to 0$, we have

$$\liminf_{j \to \infty} v_{n-k}(F_j(\varepsilon, \delta_j, \rho) \cap B(p_j, D)) < \infty,$$

where $F_j(\varepsilon, \delta_j, \rho)$ denotes the set of all $(\varepsilon, \delta_j, \rho)$ -noncritical points of f_j in F_j .

We prove it by reverse induction on l, the dimension of the limit space. Note that $k \leq l \leq n$ (the left inequality follows from Lemma 3.6; in particular, $l \geq 1$). By the compactness of the limit set of $F_j(\varepsilon, \delta_j, \rho) \cap B(p_j, D)$, the proof is reduced to the following local version:

Lemma 6.18. Suppose $x_j \in F(\varepsilon, \delta_j, \rho) \cap B(p_j, D)$ converges to $x \in \overline{B}(p, D)$. Then, there exists d > 0 (independent of j) such that

$$\liminf_{j \to \infty} v_{n-k}(F_j(\varepsilon, \delta_j, \rho) \cap B(x_j, d)) < \infty.$$

Let x_j and x be as above. Since it is sufficient to prove Proposition 6.17 for smaller ε , we may assume $\operatorname{vol}_{l-1} \Sigma_x \geq \varepsilon$. By the $(\varepsilon, \delta_j, \rho)$ -noncriticality of f_j at x_j , there exist $g_j = (|a_1^j \cdot |, \ldots, |a_k^j \cdot |) : M_j \to \mathbb{R}^k$ and $w_j \in M_j$ satisfying the conditions of Definition 6.13. Passing to a subsequence, we may assume that w_j converges to $w \in X$. The following argument is similar to Setting 5.8. Let $0 < \omega < \delta$, where $\delta \ll \varepsilon$ is different from δ_j . Since $\operatorname{vol}_{l-1} \Sigma_x \geq \varepsilon$, there exists an ω -discrete set $\{\xi_\alpha\}_{\alpha=1}^N$ in the $\varepsilon/100$ -neighborhood of $w'_x \in \Sigma_x$ such that $N \geq L/\omega^{l-1}$, where $L = c(\varepsilon)$. Let $w_\alpha \in M$ be a point near x in the direction ξ_α and $w^j_\alpha \in M_j$ a lift of w_α (i.e. $w^j_\alpha \to w_\alpha$). Then, we have

$$\tilde{\angle} w_{\alpha} x w_{\alpha'} > \omega, \quad \liminf_{j \to \infty} \tilde{\angle} a_i^j x_j w_{\alpha}^j > \pi/2 + \varepsilon/2$$

for all $1 \le \alpha \ne \alpha' \le N$ and $1 \le i \le k$ (note that $|a_i^j x_j|$ may go to infinity in the second inequality). Let $0 < r < \delta \min_{\alpha} \{\rho, |xw_{\alpha}|\}$ be sufficiently small. Then, for any $y \in B(x, 2r)$ and $y_j \in B(x_j, 2r)$, we have

$$\tilde{\angle} w_{\alpha} y w_{\alpha'} > \omega, \quad \liminf_{j \to \infty} \tilde{\angle} a_i^j y_j w_{\alpha}^j > \pi/2 + \varepsilon/3$$

for all $1 \leq \alpha \neq \alpha' \leq N$ and $1 \leq i \leq k$. Define $\sigma: X \to \mathbb{R}$ by

$$\sigma := \frac{1}{N} \sum_{\alpha=1}^{N} |w_{\alpha} \cdot |.$$

The maps f_j are uniformly Lipschitz on $B(x_j, \rho)$ by Definition 6.13(1). Hence, we may assume that the normalized map $f_j(\cdot) - f_j(x_j)$ converges to some f: $B(x, \rho) \to \mathbb{R}^k$. Note that $g_j(\cdot) - g_j(x_j)$ also converges to f by Definition 6.13(1) since $\delta_j \to 0$. Set

$$F^+ := \{ y \in B(x,\rho) \mid f^i(y) \ge f^i(x) = 0, \ 1 \le i \le k \},\$$

where f^i denotes the *i*-th component of f.

Claim 6.19 (cf. [8, 3.4 Assertion 1]). One of the following holds:

- (1) There exist $a_x \in X$ and $\rho_x > 0$ such that $(f_j, |a_x^j \cdot |)$ is $(c(\varepsilon), \varkappa(\delta), \rho_x)$ -noncritical at x_j for sufficiently large j, where $a_x^j \in M_j$ is a lift of a_x .
- (2) The restriction of σ to $F^+ \cap \overline{B}(x, r)$ has a strict maximum value at x. More precisely, we have

$$\sigma(x) \ge \sigma(y) + c(\varepsilon)|xy|$$

for any $y \in F^+ \cap \overline{B}(x, r)$.

Proof. The proof is the same as that of [8, 3.4 Assertion 1] (see Lemma 5.9). Take $a = c(\varepsilon)$ and $b = c(\varepsilon)$ such that $a \ll b \ll L$. Suppose that (2) does not hold for this a. Let $y \in F^+ \cap \overline{B}(x,r)$ be such that $\sigma(x) < \sigma(y) + a|xy|$. Then, we have $\frac{1}{N} \sum_{\alpha=1}^{N} \cos \angle w_{\alpha} xy < 2a$ (where $\angle w_{\alpha} xy$ is the minimum angle between xw_{α} and xy). On the other hand, $N \ge L/\omega^{l-1}$ and the number of α such that $|\angle w_{\alpha} xy - \pi/2| \le b$ is less than Cb/ω^{l-1} by Lemma 3.8. Hence, there exists α such that $\angle w_{\alpha} xy > \pi/2 + b$; otherwise, $\frac{1}{N} \sum_{\alpha=1}^{N} \cos \angle w_{\alpha} xy > c(b) > 2a$, a contradiction. Choose a point w_x on a shortest path xw_{α} so close to x that $\angle w_x xy > \pi/2 + b$. Take a lift w_x^j of w_x on a shortest path $x_j w_{\alpha}^j$ (we may assume $x_j w_{\alpha}^j$ converges to xw_{α}). Set $a_x := y$ and let $a_x^j \in M_j$ be a lift of a_x . Then, we have

$$\angle a_i^j x_j a_x^j > \pi/2 - \varkappa(\delta), \quad \angle a_x^j x_j w_x^j > \pi/2 + b$$

for all $1 \leq i \leq k$ and sufficiently large j (the first inequality follows from $a_x \in F^+$). The other conditions of noncriticality of $(f_j, |a_x^j \cdot |)$ at x_j are obviously satisfied. \Box

We first prove Lemma 6.18 in the case of Claim 6.19(1). Recall that Proposition 6.16 for k + 1 holds by the induction hypothesis. Let $0 < d < \rho_x \delta$. Then, $(f_j, |a_x^j \cdot |)$ is $(c(\varepsilon), \varkappa(\delta), \rho_x/2)$ -noncritical on $B(x_j, 2d)$. In particular, it is $c(\varepsilon)$ -open on $B(x_j, 2d)$. Given a ν -discrete set in $F_j \cap B(x_j, d)$, split it into $2d/\delta\nu$ classes so that the difference of the distances from a_x^j to any two points in the same class is no more than $\delta\nu$. Then, for each class, we can take a corresponding $\nu/2$ -discrete set in a fiber of $(f_j, |a_x^j \cdot |)$ by the $c(\varepsilon)$ -openness. Proposition 6.16 for k + 1 implies that the number of such $\nu/2$ -discrete points is less than $C(\varepsilon)\nu^{-(n-k-1)}$ (we have to apply Proposition 6.16 to the rescaled space $\rho_x^{-1}M_j$ so that the choice of δ depends only on ε because ρ_x depends on δ). Thus, Lemma 6.18 follows.

From now on, we consider the case of Claim 6.19(2).

Subclaim 6.20 (cf. [8, 3.4 Assertion 2]). In the case of Claim 6.19(2), for any $y \in F^+ \cap B(x, r) \setminus \{x\}$, there exists α such that $\tilde{\angle} w_{\alpha} y x > \pi/2 + c(\varepsilon)$.

Proof. Take $a = c(\varepsilon) \ll L$. It suffices to show that $\angle w_{\alpha}xy \leq \pi/2 - a$ for some α (where $\angle w_{\alpha}xy$ is the minimum angle). Suppose that $\angle w_{\alpha}xy > \pi/2 - a$ for all α . Then, by Lemma 3.8, the number of α such that $\angle w_{\alpha}xy \leq \pi/2 + a$ is less than Ca/ω^{l-1} . Since $N \geq L/\omega^{l-1}$ and $a \ll L$, we have

$$\sigma'(y'_x) \ge \frac{1}{N} \sum_{\alpha=1}^N -\cos \angle w_\alpha x y > c(a).$$

Recall that f is the limit of $g_j(\cdot) - g_j(x_j)$. Since g_j^i , the *i*-th component of g_j , are uniformly λ -concave near x_j , so is f^i near x (where $\lambda > 0$ depends only on κ and ρ). Hence, for any point z (close to x) on a shortest path xy, we have

$$f^{i}(z) \ge f^{i}(x) - \frac{\lambda}{2}|xz||yz| \ge f^{i}(x) - \frac{\lambda r}{2}|xz|$$

since $y \in F^+$. Furthermore, f is $c(\varepsilon)$ -open near x since it is a limit of $c(\varepsilon)$ -maps. Therefore, we can find a point $\hat{z} \in F^+ \cap B(x,r)$ such that $c(\varepsilon)|\hat{z}z| \leq (\lambda r/2)|xz|$. Since x is the maximum point of σ on $F^+ \cap \overline{B}(x,r)$, we have

$$\sigma(x) \ge \sigma(\hat{z}) \ge \sigma(z) - |\hat{z}z| \ge \sigma(z) - c(\varepsilon)^{-1} \frac{\lambda r}{2} |xz|.$$

Since z is arbitrary on xy, we obtain $\sigma'(y'_x) \leq c(\varepsilon)^{-1}\lambda r/2$. This contradicts $\sigma'(y'_x) > c(a)$ because $r < \rho \delta \ll c(\varepsilon)$.

Define $\sigma_j: M_j \to \mathbb{R}$ by

$$\sigma_j := \frac{1}{N} \sum_{\alpha=1}^N |w_{\alpha}^j \cdot|$$

and set

$$F_j^+ := \{ y_j \in B(x_j, \rho) \mid f_j^i(y_j) \ge f_j^i(x_j), \ 1 \le i \le k \},\$$

where f_j^i denotes the *i*-th component of f_j . Let \hat{x}_j be a maximum point of σ_j on $F_j^+ \cap \bar{B}(x_j, r)$. Then, Claim 6.19(2) implies that \hat{x}_j converges to x.

Subclaim 6.21 (cf. [9, 3.9]). Indeed, $\hat{x}_j \in F_j$ for large j.

Proof. Fix large j and suppose the contrary. Then, $f_j^{i_0}(\hat{x}_j) > f_j^{i_0}(x_j)$ for some i_0 . Let A'_i (resp. W') be the set of all directions of shortest paths from \hat{x}_j to a_i^j (resp. w_{α}^j for all α). Then, by Lemma 3.7(1), there exists $\xi \in \Sigma_{\hat{x}_j}$ such that

$$\angle(\xi,A_{i_0}')<\pi/2-c(\varepsilon),\quad \angle(\xi,A_i')=\pi/2,\quad \angle(\xi,W')>\pi/2+c(\varepsilon)$$

for all $i \neq i_0$. Let $y \in M_j$ be a point near \hat{x}_j in the direction ξ such that

$$f_{j}^{i_{0}}(y) > f_{j}^{i_{0}}(x_{j}), \quad |g_{j}^{i}(y) - g_{j}^{i}(\hat{x}_{j})| < \delta |y\hat{x}_{j}|, \quad \sigma_{j}(y) > \sigma_{j}(\hat{x}_{j}) + c(\varepsilon)|y\hat{x}_{j}|$$

for all $i \neq i_0$. In particular, the middle inequality above and the inequality of Definition 6.13(1) imply that

$$f_j^i(y) \ge f_j^i(\hat{x}_j) - (\delta + \delta_j)|y\hat{x}_j| \ge f_j^i(x_j) - 2\delta|y\hat{x}_j|$$

for all $i \neq i_0$. Therefore, by the $c(\varepsilon)$ -openness of f_j , we can find a point $\hat{y} \in F_j^+ \cap B(x_j, r)$ such that $c(\varepsilon)|\hat{y}y| \leq 2\delta |y\hat{x}_j|$. Then, we have

$$\sigma_j(\hat{y}) \ge \sigma_j(y) - |\hat{y}y| \ge \sigma_j(\hat{x}_j) + (c(\varepsilon) - 2\delta c(\varepsilon)^{-1})|y\hat{x}_j|$$

This contradicts the choice of \hat{x}_j because $\delta \ll c(\varepsilon)$.

Claim 6.22 (cf. [18, 3.2]). In the above situation, one of the following holds:

- (1) There exists a subsequence of $\{j\}$ such that for any $y_j \in F_j^+ \cap B(\hat{x}_j, r) \setminus \{\hat{x}_j\}$, there is α such that $\tilde{\angle} w_{\alpha}^j y_j \hat{x}_j > \pi/2 + c(\varepsilon)$.
- (2) There exists a sequence $d_j \to 0$ of positive numbers such that (i) for any $u \in E^+ \cap B(\hat{a}, m) \setminus \overline{B}(\hat{a}, d_j)$ there is a such that $\tilde{f}(m)$
 - (i) for any $y_j \in F_j^+ \cap B(\hat{x}_j, r) \setminus \overline{B}(\hat{x}_j, d_j)$, there is α such that $\tilde{\angle} w_{\alpha}^j y_j \hat{x}_j > \pi/2 + c(\varepsilon);$
 - (ii) for any limit (\hat{X}, \hat{x}) of the rescaled spaces $(\frac{1}{d_j}M_j, \hat{x}_j)$, we have dim $\hat{X} \ge l+1$, where $l = \dim X$.

In particular, if l = n, then (1) holds. Note that when (1) (resp. (2)) holds, $(f_j, |\hat{x}_j \cdot |)$ is $(c(\varepsilon), \varkappa(\delta), s/2)$ -noncritical on $F_j \cap \partial B(\hat{x}_j, s)$ for any $s \in (0, r)$ (resp. $s \in (d_j, r)$).

Proof. Let $a_1 = c(\varepsilon)$ be the constant in Subclaim 6.20. Take $a = c(\varepsilon)$ such that $a \ll L$ and $a \le a_1$. Suppose that (1) does not hold for this a. Then, for any large j, there exists $y_j \in F_j^+ \cap B(\hat{x}_j, r) \setminus {\hat{x}_j}$ such that $\tilde{\angle} w_{\alpha}^j y_j \hat{x}_j \le \pi/2 + a$ for all α . Let \hat{y}_j be a farthest point from \hat{x}_j among such y_j and set $d_j := |\hat{x}_j \hat{y}_j|$. Then, (2)(i) is trivial. Moreover, Subclaim 6.20 implies that $d_j \to 0$ since $a \le a_1$.

Now, we prove (2)(ii). Suppose $(\frac{1}{d_j}M_j, \hat{x}_j)$ converges to an Alexandrov space (\hat{X}, \hat{x}) of nonnegative curvature. Passing to a subsequence, we may assume that \hat{y}_j converges to $\hat{y} \in \hat{X}$. We may further assume that shortest paths $\hat{x}_j \hat{y}_j$ and $\hat{x}_j w_{\alpha}^j$ converge to a shortest path $\hat{x}\hat{y}$ and a ray γ_{α} starting from \hat{x} , respectively. Let $v_j, v_{\alpha}^j \in \Sigma_{\hat{x}_j}$ and $v, v_{\alpha} \in \Sigma_{\hat{x}}$ denote the directions of them, respectively. Then, by the monotonicity of angles, we have

$$\angle (v_{\alpha}, v_{\alpha'}) \ge \lim_{j \to \infty} \check{\angle} w_{\alpha}^{j} \hat{x}_{j} w_{\alpha'}^{j} \ge \omega,$$

$$\angle (v, v_{\alpha}) \ge \lim_{j \to \infty} \check{\angle} \hat{y}_{j} \hat{x}_{j} w_{\alpha}^{j} = \pi - \lim_{j \to \infty} \check{\angle} \hat{x}_{j} \hat{y}_{j} w_{\alpha}^{j} \ge \pi/2 - a$$

for all $1 \leq \alpha \neq \alpha' \leq N$. Suppose (2)(ii) does not hold, i.e. dim $\hat{X} = l$. Then, an argument similar to the first paragraph of the proof of Subclaim 6.20 implies that

(6.2)
$$\frac{1}{N}\sum_{\alpha=1}^{N} -\cos \angle (v, v_{\alpha}) > c(a)$$

On the other hand, by an argument similar to the second paragraph of the proof of Subclaim 6.20, we show that

(6.3)
$$\sigma'_j(v_j) \le c(\varepsilon)^{-1} \left(\frac{\lambda d_j}{2} + 2\delta_j\right),$$

where $\lambda > 0$ is a constant depending only on κ and ρ such that g_j^i are uniformly λ -concave near x_j . Fix large j and let z be an arbitrary point on a shortest path $\hat{x}_j \hat{y}_j$. Then, by the λ -concavity of g_j^i , we have

$$\frac{|\hat{y}_j z|}{d_j} (g_j^i(z) - g_j^i(\hat{x}_j)) + \frac{|\hat{x}_j z|}{d_j} (g_j^i(z) - g_j^i(\hat{y}_j)) \ge -\frac{\lambda}{2} |\hat{x}_j z| |\hat{y}_j z|.$$

Moreover, since $\hat{x}_j, \hat{y}_j \in F_j^+$, by the inequality of Definition 6.13(1), we have

$$f_j^i(z) - f_j^i(x_j) \ge \begin{cases} f_j^i(z) - f_j^i(\hat{x}_j) \ge (g_j^i(z) - g_j^i(\hat{x}_j)) - \delta_j |\hat{x}_j z|, \\ f_j^i(z) - f_j^i(\hat{y}_j) \ge (g_j^i(z) - g_j^i(\hat{y}_j)) - \delta_j |\hat{y}_j z|. \end{cases}$$

Combining the three inequalities above, we obtain

$$f_j^i(z) - f_j^i(x_j) \ge -\frac{\lambda}{2} |\hat{x}_j z| |\hat{y}_j z| - 2\delta_j \frac{|\hat{x}_j z| |\hat{y}_j z|}{d_j}$$
$$\ge -\left(\frac{\lambda d_j}{2} + 2\delta_j\right) |\hat{x}_j z|.$$

Furthermore, since f_j is $c(\varepsilon)$ -open near x_j , we can find a point $\hat{z} \in F_j^+ \cap B(x_j, r)$ such that $c(\varepsilon)|\hat{z}z| \leq (\lambda d_j/2 + 2\delta_j)|\hat{x}_j z|$. Since \hat{x}_j is the maximum point of σ_j on $F_i^+ \cap \bar{B}(x_j, r)$, we have

$$\sigma_j(\hat{x}_j) \ge \sigma(\hat{z}) \ge \sigma(z) - |\hat{z}z| \ge \sigma(z) - c(\varepsilon)^{-1} \left(\frac{\lambda d_j}{2} + 2\delta_j\right) |\hat{x}_j z|.$$

Since z is arbitrary on $\hat{x}_j \hat{y}_j$, we obtain the desired inequality (6.3).

Therefore, applying the first variation formula to the left-hand side of the inequality (6.3) and passing to the limit, we obtain

$$\frac{1}{N}\sum_{\alpha=1}^{N} - \cos \angle (v, v_{\alpha}) \le 0$$

by the lower semicontinuity of angles (choose v_{α}^{j} to be closest to v_{j} among all directions from \hat{x}_{j} to w_{α}^{j} so that the first variation formula holds). This contradicts the inequality (6.2).

Now, we give the proof of Lemma 6.18 in the case of Claim 6.19(2). Recall that Proposition 6.16 for k + 1 and Proposition 6.17 for k and $l' \ge l + 1$ hold by the induction hypothesis. The proof is divided into the two cases of Claim 6.22.

First, suppose Claim 6.22(1) holds. In this case, Lemma 6.18 follows from Proposition 6.17 for k + 1 as in the case of Claim 6.19(1). Consider a ν -discrete set in $F_j \cap B(\hat{x}_j, r) \setminus B(\hat{x}_j, \nu/2)$. Note that $(f_j, |\hat{x}_j \cdot |)$ is $(c(\varepsilon), \varkappa(\delta), \nu/3)$ -noncritical (and

hence $c(\varepsilon)$ -open) on a small neighborhood of $F_j \cap B(\hat{x}_j, r) \setminus B(\hat{x}_j, \nu/2)$. Split the ν -discrete set into $r/\delta\nu$ classes so that the difference of the distances from \hat{x}_j to any two points in the same class is no more than $\delta\nu$. Then, for each class, we can take a corresponding $\nu/2$ -discrete set of $F_j \cap \partial B(\hat{x}_j, s)$ for some $s \in (\nu/2, r)$ by the $c(\varepsilon)$ -openness. Proposition 6.16 for k + 1 implies that the number of such $\nu/2$ -discrete points is less than $C(\varepsilon)\nu^{-(n-k-1)}$ (apply Proposition 6.16 to the rescaled space $s^{-1}M$). Thus, Lemma 6.18 follows. In particular, Proposition 6.17 for l = n holds (the base case of the reverse induction on l).

Next, suppose Claim 6.22(2) holds. Then, the same argument as in the previous case shows that $v_{n-k}(F_j \cap B(\hat{x}_j, r) \setminus \bar{B}(\hat{x}_j, d_j))$ is uniformly bounded above. On the other hand, Proposition 6.17 for $l' \geq l+1$ applied to $\frac{1}{d_j}M_j$ implies that $v_{n-k}(F_j(\varepsilon, \delta_j, d_j) \cap \bar{B}(\hat{x}_j, d_j))$ is uniformly bounded above for some subsequence. This completes the proof.

6.4. Almost continuity of the volume of the fibers. In this section, we prove Theorem 1.4(3). The proof is similar to that of the continuity of the volume of Alexandrov spaces under the Gromov-Hausdorff convergence ([2, 10.8], [12, 3.5], [17, 0.6]). By Remarks 6.1 and 6.4, it suffices to prove the following:

Proposition 6.23. Let M be an n-dimensional Alexandrov space, $p \in M$ and k < n. Let $f : B(p, \ell) \to \mathbb{R}^k$ be a δ -almost regular map associated with a (k, δ) -strainer $\{(a_i, b_i)\}_{i=1}^k$ at p with length $> \ell$. For simplicity, assume that $f^{-1}(\overline{B}(f(p), r))$ is contained in $B(p, \ell\delta)$ for some $0 < r \ll \ell\delta$. Set $F_v := f^{-1}(v)$ for $v \in B(f(p), r)$. Then, for any sufficiently close $u, v \in B(f(p), r)$, we have

$$\left|\frac{\operatorname{vol}_{n-k} F_u}{\operatorname{vol}_{n-k} F_v} - 1\right| < \varkappa(\delta).$$

Let F_v be as above. Let m be a positive integer and θ , ρ positive numbers. We denote by $F_v(m, \theta, \rho)$ the set of all points in F_v having an (m, θ) -strainer $\{(a_{k+i}, b_{k+i})\}_{i=1}^m$ with length $> \rho$ such that $a_{k+i}, b_{k+i} \in F_v$. The above proposition immediately follows from the following two lemmas:

Lemma 6.24. For any small $\rho > 0$, if $u, v \in B(f(p), r)$ are sufficiently close, then there exists a $\varkappa(\delta)$ -almost isometry from $F_v(n-k, \delta, \rho)$ to an open subset of F_u .

Proof. Since the proof is similar to that of Theorem 4.1 in the noncollapsing case, we only outline it. Assume $\rho \ll |f^{-1}(\bar{B}(f(p),r)), \partial B(p,\ell\delta)|$ and $|uv| \ll \rho\delta$. Fix $x \in F_v(n-k,\delta,\rho)$ and let $\{(a_{k+i},b_{k+i})\}_{i=1}^{n-k}$ be an $(n-k,\delta)$ -strainer at x with length $> \rho$ such that $a_{k+i}, b_{k+i} \in F_v$. Then, the inequality (6.1) implies that $\{(a_i,b_i)\}_{i=1}^n$ is an $(n,\varkappa(\delta))$ -strainer at x with length $> \rho$. Hence, by Proposition 6.5(1), there exists a $\varkappa(\delta)$ -almost isometry from $B(x,\rho\delta)$ to an open subset of \mathbb{R}^n whose first k components coincide with f. Therefore, a translation with respect to the fcoordinate gives a $\varkappa(\delta)$ -almost isometry from $B(x,\rho\delta/2) \cap F_v$ to an open subset of F_u , provided $|uv| \ll \rho\delta$. Gluing such local maps as in the proof of Theorem 4.1, we obtain a global almost isometry. \Box

Lemma 6.25 (cf. [2, 10.9]). Let $1 \le m \le n - k$ and $\theta, \rho > 0$. Then, for any $0 < \nu < \rho$, we have

$$\nu^{n-k}\beta_{\nu}(F_v \setminus F_v(m,\theta,\rho)) < \varkappa_{\theta}(\rho),$$

where $\varkappa_{\theta}(\rho)$ is a positive function depending only on n, κ and θ such that $\varkappa_{\theta}(\rho) \to 0$ as $\rho \to 0$. In particular,

$$\operatorname{vol}_{n-k}(F_v \setminus F_v(n-k,\delta,\rho)) < \varkappa_{\delta}(\rho).$$

Proof. The proof is carried out by the induction on m as in [2, 10.9]. Let $\rho_1 \gg \rho$ and $\theta_1 = c(\theta) \ll \theta$. Take a maximal $(\rho_1\theta_1)$ -discrete net $\{x_\alpha\}_{\alpha=1}^N$ of $F_v(m-1,\theta_1,\rho_1)$. Then, we have $N < C(\rho_1\theta_1)^{-(n-k)}$ by Proposition 6.16. Fix α and let $\{(a_{k+i},b_{k+i})\}_{i=1}^{m-1}$ be an $(m-1,\theta_1)$ -strainer at x_α with length $> \rho_1$ such that $a_{k+i}, b_{k+i} \in F_v$. Divide $B(x_\alpha,\rho_1\theta_1) \cap F_v$ into $(2\rho_1/\rho)^{m-1}$ classes $\{D_{\alpha\beta}\}_{\beta}$ so that if x, y belong to the same class, then we have $||a_{k+i}x| - |a_{k+i}y|| \leq \rho\theta_1$ for all $1 \leq i \leq m-1$. By the argument in [2, 10.5], if there exist sufficiently many ρ -discrete points (depending on θ) in $D_{\alpha\beta}$, then we can choose three of them so that they form a $(1,\theta)$ -strainer (and a strained point) with length $> \rho$. Therefore, the number of ρ -discrete points in $D_{\alpha\beta} \setminus F_v(m,\theta,\rho)$ is less than $C(\theta)$. Hence, we can cover $F_v(m-1,\theta_1,\rho_1) \setminus F_v(m,\theta,\rho)$ by at most $C(\theta)\rho_1^{-(n-k)}(\rho_1/\rho)^{m-1}$ balls of radius ρ (recall $\theta_1 = c(\theta)$). By Proposition 6.16 again, the number of ν -discrete points in each ρ -ball is less than $C(\rho/\nu)^{n-k}$ for any $0 < \nu < \rho$. Thus, we have

$$\nu^{n-k}\beta_{\nu}(F_{v}(m-1,\theta_{1},\rho_{1})\setminus F_{v}(m,\theta,\rho)) < C(\theta)\left(\frac{\rho}{\rho_{1}}\right)^{n-k-m+1}$$

for any $0 < \nu < \rho$. Together with the induction hypothesis and a suitable choice of ρ_1 , this yields the desired estimate.

Proof of Proposition 6.23. By Proposition 6.9 and Remark 6.10, we may assume $\operatorname{vol}_{n-k} F_v > c(p)$ for any $v \in B(f(p), r)$. Let $\rho > 0$ be so small that $\varkappa_{\delta}(\rho) \ll c(p)$. Then, by the above two lemmas, for sufficiently close $u, v \in B(f(p), r)$, we have

$$(1 + \varkappa(\delta)) \operatorname{vol}_{n-k} F_u \ge \operatorname{vol}_{n-k} F_v(n-k,\delta,\rho)$$

$$\ge \operatorname{vol}_{n-k} F_v - \varkappa_\delta(\rho) \ge (1 - \varkappa(\delta)) \operatorname{vol}_{n-k} F_v.$$

This completes the proof.

Acknowledgment

The author would like to thank Prof. Takao Yamaguchi for his advice and encouragement. He is also grateful to the referee for the careful reading and useful comments.

References

- D. Burago, Y. Burago and S. Ivanov, A course in metric geometry, Grad. Stud. Math., 33, Amer. Math. Soc., Providence, RI, 2001.
- [2] Yu. Burago, M. Gromov and G. Perel'man, A.D. Alexandrov spaces with curvature bounded below, Uspekhi Mat. Nauk 47 (1992), no. 2(284), 3–51, 222; translation in Russian Math. Surveys 47 (1992), no. 2, 1–58.
- [3] T. Fujioka, Regular points of extremal subsets in Alexandrov spaces, preprint, arXiv:1905.05480, 2019.
- [4] K. Fukaya and T. Yamaguchi, The fundamental groups of almost nonnegatively curved manifolds, Ann. of Math. 136 (1992), 253–333.
- [5] V. Kapovitch, Perelman's stability theorem, Surv. Differ. Geom., XI, 103–136, Int. Press, Somerville, MA, 2007.
- [6] V. Kapovitch, A. Petrunin and W. Tuschmann, Nilpotency, almost nonnegative curvature, and the gradient flow on Alexandrov spaces, Ann. of Math. 171 (2010), 343–373.

- [7] A. Mitsuishi and T. Yamaguchi, Obtuse constants of Alexandrov spaces, J. Math. Soc. Japan 71 (2019), no. 4, 1081–1103.
- [8] G. Perelman, Alexandrov's spaces with curvatures bounded from below II, preprint, 1991.
- G. Ya. Perel'man, Elements of Morse theory on Aleksandrov spaces, Algebra i Analiz 5 (1993), no. 1, 232–241; translation in St. Petersburg Math. J. 5 (1994), no. 1, 205–213.
- [10] G. Perelman, Collapsing with no proper extremal subsets, Comparison geometry, 149–154, Math. Sci. Res. Inst. Publ., 30, Cambridge Univ. Press, Cambridge, 1997.
- X. Rong and S. Xu, Stability of e^ϵ-Lipschitz and co-Lipschitz maps in Gromov-Hausdorff topology, Adv. Math. 231 (2012), 774–797.
- [12] T. Shioya, Mass of rays in Alexandrov spaces of nonnegative curvature, Comment. Math. Helv. 69 (1994), no. 2, 208–228.
- [13] Y. Wang, X. Su and H. Sun, A new proof of almost isometry theorem in Alexandrov geometry with curvature bounded below, Asian J. Math. 17 (2013), no. 4, 715–728.
- [14] S. Xu, Homotopy lifting property of an e^{ϵ} -Lipschitz and co-Lipschitz map, preprint, arXiv:1211.5919, 2013.
- [15] S. Xu and X. Yao, Margulis lemma and Hurewicz fibration theorem on Alexandrov spaces, preprint, arXiv:1902.10973, 2019.
- [16] T. Yamaguchi, Collapsing and pinching under a lower curvature bound, Ann. of Math. 133 (1991), no. 2, 317–357.
- [17] T. Yamaguchi, A convergence theorem in the geometry of Alexandrov spaces, Actes de la table ronde de géométrie différentielle, 601–642, Sémin. Congr., 1, Soc. Math. France, Paris, 1996.
- [18] T. Yamaguchi, Collapsing and essential coverings, preprint, arXiv:1205.0441, 2012.

DEPARTMENT OF MATHEMATICS, KYOTO UNIVERSITY, KITASHIRAKAWA, KYOTO 606-8502, JAPAN *Email address*: tfujioka@math.kyoto-u.ac.jp