# 1ST EIGENVALUE PINCHING FOR CONVEX HYPERSURFACES IN A RIEMANNIAN MANIFOLD

YINGXIANG HU, SHICHENG XU

ABSTRACT. Let  $M^n$  be a closed convex hypersurface lying in a convex ball B(p, R) of the ambient (n + 1)-manifold  $N^{n+1}$ . We prove that, by pinching Heintze-Reilly's inequality via sectional curvature upper bound of B(p, R), 1st eigenvalue and mean curvature of M, not only M is Hausdorff close and almost isometric to a geodesic sphere  $S(p_0, R_0)$  in N, but also its enclosed domain is  $C^{1,\alpha}$ -close to a geodesic ball of constant curvature.

# 1. INTRODUCTION

As a natural extension of Reilly's result [24] in  $\mathbb{R}^{n+1}$ , Heintze [12] proved that a closed submanfold  $M^n$  (including high codimension) lying in a convex ball B(p, R) with  $R \le \frac{\pi}{4\sqrt{\delta}}$  of  $N^{n+m}$ , whose sectional curvature  $K_N \le \delta$ , satisfies

(1.1) 
$$\lambda_1(M) \le \begin{cases} n\delta + \frac{n}{|M|} \int_M |H|^2, & \text{for } \delta \ge 0; \\ n\delta + n||H||_{\infty}^2, & \text{for } \delta < 0, \end{cases}$$

where  $\lambda_1(M)$  is the first nonzero eigenvalue of the Laplace-Beltrami operator on M,  $||H||_{\infty} = \max_M |H|$  and |M| its area. Here  $\frac{\pi}{\sqrt{\delta}}$  is viewed to be  $\infty$  if  $\delta \le 0$ . Equality holds in (1.1) if and only if M is minimally immersed in some geodesic sphere of N.

Let  $s_{\delta}$  be the usual  $\delta$ -sine function defined on  $[0, \frac{\pi}{\sqrt{\delta}})$  by

$$s_{\delta}(r) := \begin{cases} \frac{1}{\sqrt{\delta}} \sin(\sqrt{\delta}r), & \text{if } \delta > 0; \\ r, & \text{if } \delta = 0; \\ \frac{1}{\sqrt{-\delta}} \sinh(\sqrt{-\delta}r), & \text{if } \delta < 0. \end{cases}$$

and  $s_{\delta}^{-1}$  be its inverse function. Let  $\omega_n$  be the volume of *n*-sphere of radius 1 in  $\mathbb{R}^{n+1}$ . As observed in [14], if equality holds in (1.1) for an immersed hypersurface  $M^n$  ( $n \ge 2$ ) in  $N^{n+1}$ , then *M* encloses a geodesic ball of constant curvature  $\delta$ . We prove in [14] that if (1.1) almost holds for  $||H||_{\infty}^2$ , i.e.,

(1.2) 
$$n(\delta + ||H||_{\infty}^2) \le \lambda_1(M)(1+\epsilon),$$

then *M* is Hausdorff close to a geodesic sphere  $S(p_0, R_0)$ , and  $B(p_0, R_0)$  is almost to be of constant  $\delta$ , provided with  $R \leq \frac{\pi}{8\sqrt{\delta}}$ ,  $|M| \leq \omega_n s_{\delta}^n(\frac{\pi}{4\sqrt{\delta}})$ , and a rescaling invariant control on the mean curvature and area of *M*:

(1.3)

 $|M|^{\frac{1}{n}}||H||_{\infty} \le A.$ 

Date: May 15, 2019.

<sup>2010</sup> Mathematics Subject Classification. 53C20, 53C21, 53C24.

Key words and phrases. 1st eigenvalue, mean curvature, upper curvature bound, quantitative rigidity, convex hypersurfaces.

According to [14], (1.3) can be viewed as a relative non-collapsing condition for hypersurfaces, i.e.,

$$\frac{|B(x_0, r) \cap M|}{|M|} \ge C(n, \delta)r^n, \quad \text{for any } x_0 \in M \text{ and } 0 < r \le \min\left\{ \text{diam}_N(M), \frac{\pi}{2\sqrt{\delta}} \right\},$$

where diam<sub>N</sub>(M) is the extrinsic diameter of M as a subset in N.

Therefore, typical examples that do not satisfies (1.3) contain the boundary  $\partial U_r$  of *r*-neighborhoods of a high co-dimensional submanifold *X* (not a point) with  $r \ll \operatorname{diam}_N X$ .

In this paper, we prove that if  $\partial U$  is convex, then  $\partial U$  still cannot satisfy pinching condition (1.2). We use  $\varkappa(\epsilon | R, \delta, \cdots)$  to denote a positive function on variables  $\epsilon, R, \delta, \cdots$  that converges to 0 as  $\epsilon \to 0$  with other quantities  $R, \delta, \cdots$  fixed.

**Theorem 1.1.** Let *n* be an integer  $\geq 2$  and let  $N^{n+1}$  be a complete Riemannian manifold with  $\mu \leq K_N \leq \delta$ . Let  $M^n$  be an embedded convex hypersurface in a convex geodesic ball B(p, R) of *N*. If  $\delta > 0$  we further assume

(1.4) 
$$R \le \frac{\pi}{8\sqrt{\delta}}, \quad |M| \le \omega_n s_{\delta}^n \left(\frac{\pi}{4\sqrt{\delta}}\right).$$

*Then* (1.2) *with*  $\epsilon < \epsilon_0 = \epsilon_0(\mu, \delta, n, R)$  *implies that* 

- (1.1.1) *M* is  $C \epsilon^{\frac{1}{2(2n+1)}} s_{\delta}(R_0)$ -Hausdorff-close and  $C \epsilon^{\frac{1}{2(2n+1)}}$ -almost isometric to a geodesic sphere  $S(p_0, R_0)$ , where  $R_0 = s_{\delta}^{-1}(\frac{1}{\sqrt{\delta + \|H\|_{\infty}^2}})$ , and  $C = C(n, \delta, \mu, R)$  is a positive constant;
- (1.1.2) *M* is  $\varkappa(\epsilon | R, \delta, \mu, n, \alpha)$ - $C^{1,\alpha}$  close to a round sphere of constant curvature  $1/(s_{\delta}(R_0))^2$ with  $0 \le \alpha < 1$ ;
- (1.1.3) The enclosed domain  $\Omega$  by M is  $\varkappa(\epsilon | R, \delta, \mu, n, \alpha) C^{1,\alpha}$ -close to a ball of constant curvature  $\delta$ .

*Here*  $\epsilon_0$ , *C*,  $\varkappa$  *above do not depend on R when*  $\delta \ge 0$ .

By Theorem 1.1, one cannot obtain a collapsed metric g via perturbing the interior of a geodesic ball  $\Omega$  of constant curvature  $\delta$  to a new metric  $g_{\delta}$  whose sectional curvature's upper bound is close to  $\delta$ .

We point it out that the extrinsically relative non-collapsing condition (1.3) are required in all earlier known results via pinching (1.3). Here for Theorem 1.1 we also prove that convexity of *M* plus the pinching condition (1.2) implies (1.3) holds for some universal *A*.

At present quantitative rigidity results with respect to the upper curvature bound  $\delta$  are rarely known. There are relative more rigidity results for convex domains to be of constant curvature via information along their boundary and interior's lower (Ricci) curvature bound (e.g., [25, Theorems 1, 2], [18, Theorem 1.2], etc.). Compared to Theorem 1.1, few of their quantitative versions are known.

We refer to [14], [11], [1], [7], and [13] for related results in manifolds and space forms. Let us give the main idea in proving Theorem 1.1. If there is a universal constant  $A = A(n, \delta, \mu)$  such that (1.3) holds for a convex hypersurface *M* in Theorem 1.1, then it is a direct corollary of the following and Cheeger-Gromov's convergence Theorem ( [4, 5], [10], [15], [9], [21]) that conclusions in Theorem 1.1 hold.

**Theorem 1.2** ([14]). Let *n* be an integer  $\geq 2$ , and let  $N^{n+1}$  be a complete Riemannian manifold with  $\mu \leq K_N \leq \delta$ . Let  $M^n$  be an immersed hypersurface in a convex geodesic ball B(p, R) of N, where R and |M| satisfy (1.4) when  $\delta > 0$ . If for  $q \geq \overline{q} > n$  and A > 0, we have

(1.5)  $|M|^{\frac{1}{n}} ||H||_{\infty} \le A, \quad and \quad |M|^{\frac{1}{n}} ||B||_{q} \le A,$ 

2

where  $||B||_q = \left(\frac{1}{|M|}\int |B|^q\right)^{1/q}$  is the normalized  $L^q$  norm of the 2nd fundamental form B of M. Then (1.2) with  $\epsilon < \epsilon_1(A, \bar{q}, R, \delta, \mu, n)$  implies that

- (1.2.1) *M* is  $C_1 \epsilon^{\frac{1}{2(2n+1)}} s_{\delta}(R_0)$ -Hausdorff-close to a geodesic sphere  $S(p_0, R_0)$  in *N*, where  $R_0 = s_{\delta}^{-1}(\frac{1}{\sqrt{\delta + ||H||_{c_0}^2}});$
- (1.2.2)  $B(p_0, R_0)$  is  $\varkappa(\epsilon) C^{1,\alpha}$ -close to a ball of constant curvature  $\delta$  for any  $0 \le \alpha < 1$ ; (1.2.3) *M* is embedded and  $C_2 \epsilon^{\min\{\frac{1}{2(2n+1)}, \frac{q-n}{2(q-n+qn)}\}}$ -almost isometric to  $S(p_0, R_0)$ .

Here constants  $C_1, C_2$  and  $\epsilon_1$  depend on  $n, \delta, \mu, \bar{q}, R, A$ , and the function  $\varkappa(\epsilon)$  also depends on  $n, \delta, \mu, \bar{q}, R, A, \alpha$ , where the dependence on R can be dropped when  $\delta \ge 0$ .

In order to obtain  $A(n, \mu, \delta)$  above, we argue by contradiction. Suppose there is a sequence of convex domains  $\Omega_i$  such that  $|\partial \Omega_i|^{\frac{1}{n}} ||H_{\partial \Omega_i}||_{\infty} \to \infty$ . We will prove that  $\lambda_1(\partial \Omega_i)$  admits a universal upper bound after a rescaling such that the area  $|\partial \Omega_i| = 1$ . On the other hand, by pinching condition (1.2),  $||H_{\partial \Omega_i}||_{\infty}$  has a uniform upper bound, a contradiction.

We will use gradient flows of semi-concave functions on Alexandrov spaces [2] to derive a universal ratio upper bound on the intrinsic diameter of  $\partial \Omega_i$ , and then apply Cheeger-Colding's eigenvalue convergence theorem [6] on  $\partial \Omega_i$  to obtain an upper bound of  $\lambda_1$ .

## 2. Proof of Theorem 1.1

In the following we will apply some results on Alexandrov spaces with curvature bounded below. The references of this section are [2], [23].

Roughly speaking, a finite-dimensional *Alexandrov space X of curv*  $\geq \mu$  is a locally compact length metric space on which Toponogov comparison theorem for triangles holds just as that on manifolds of sectional curvature  $\geq \mu$ .

A locally Lipschitz function  $f : U \subset X \to \mathbb{R}$  is called  $\lambda$ -concave, if U is an open domain and there is a real number  $\lambda$  such that for any unit-speed minimal geodesic  $\gamma$  in U,  $f \circ \gamma(t) - \frac{\lambda}{2}t^2$  is concave.

The gradient flow  $\Phi_t$  of a  $\lambda$ -concave function is well-defined, and is  $e^{\lambda t}$ -Lipschitz (see [23, Lemma 2.1.4(i)]). *Extremal subsets* are those subsets which are invariant under gradient flows of all  $\lambda$ -concave functions [19, 23]. The boundary of an Alexandrov space of curv  $\geq \mu$  is an extremal subset.

A *quasigeodesic* in X is a unit-speed curve  $\gamma$  such that for any point p in X, the function  $f(t) = \int_0^{d(p,\gamma(t))} s_\mu(s) ds$  satisfies  $f'' \leq 1 - \mu f$  in the barrier sense. By its definition, limit of quasigeodesics is also a quasigeodesic, see [20].

By the generalized Lieberman lemma [22, Theorem 1.1], any shortest geodesic in an extremal subset is a quasigeodesic in the ambient space.

We say that a sequence  $(X_i, d_i)$  of metric spaces *GH-converges* to (X, d) in Gromov-Hausdorff topology, if there are  $\epsilon_i$ -isometries  $\psi_i : X_i \to X$  with  $\epsilon_i \to 0$ , i.e., for any  $x, y \in X_i, |d_i(x, y) - d(\psi_i(x), \psi_i(y))| \le \epsilon_i$ , and  $\epsilon_i$ -neighborhood of  $\psi_i(X_i)$  covers *X*.

The following is the main technical lemma in this paper to derive a bound of intrinsic diameter of M, which is required in applying a compactness argument under Gromov-Hausdorff topology.

**Lemma 2.1.** Let  $\Omega$  be a convex domain of diameter  $\leq D$  with a smooth boundary  $M = \partial \Omega$ in  $N^{n+1}$  whose sectional curvature  $K_N \geq \mu$ . Then the intrinsic diameter of M,

 $\operatorname{diam}(M) \le C(n, \mu, D) \operatorname{diam}(\Omega),$ 

where  $C(n, \mu, D)$  is a positive constant depending on  $n, \mu$  and D.

*Proof.* Let us argue by contradiction. Assume the contrary, then there is a sequence  $\Omega_i$  of convex domains of diameter  $\leq d$ , where the ratio diam $(M_i)/$  diam $(\Omega_i) \rightarrow \infty$ .

Note that the closure  $\overline{\Omega}_i$  of  $\Omega_i$  is an Alexandrov space with curv  $\geq \mu$  of Hausdorff dimension n + 1, and its boundary  $M_i = \partial \Omega_i$  is an extremal subset of  $\Omega_i$ . By passing to a subsequence, we may assume that  $\overline{\Omega}_i$  *GH*-converges to an Alexandrov space X with curv  $\geq \mu$ .

Case 1. The Hausdorff dimension of X equals n + 1. That is,  $\Omega_i$  is a non-collapsing sequence. Then Lemma 2.1 is a direct corollary of [22, Theorem 1.2], whose special case is that the boundary  $\partial \Omega_i = M_i$  with the intrinsic metric *GH*-converges to  $\partial X$ , the boundary of X.

Indeed, by the *GH*-convergence,  $d \ge \lim_{i\to\infty} \operatorname{diam}(\Omega_i) = \operatorname{diam}(X) > 0$ . And thus the intrinsic diameter  $\operatorname{diam}(\partial X) = \lim_{i\to\infty} \operatorname{diam}(M_i) = \infty$ . On the other hand, by Gauss equation,  $\partial \Omega_i$  is an Alexandrov space with curv  $\ge \mu$ . Hence  $\partial X$  is also an Alexandrov space with the same lower curvature bound. By the compactness of  $\partial X$ ,  $\operatorname{diam}(\partial X) < \infty$ , a contradiction.

Case 2. The Hausdorff dimension of X < n + 1. By passing to a rescaled converging subsequence, we assume that diam $(\Omega_i) \rightarrow 1$ .

Let  $p_i, q_i$  be two points in  $M_i$  such that the intrinsic distance  $d_{M_i}(p_i, q_i) = \text{diam}(M_i)$ , and let  $\gamma_i = [p_iq_i]$  be a minimal geodesic in  $\Omega_i$  connecting  $p_i = \gamma_i(0)$  and  $q_i$  and realizing their distance  $d(p_i, q_i)$  in  $\Omega_i$ . By Ascoli theorem and passing to a subsequence, we may assume  $\gamma_i$  converges to a minimal geodesic  $\gamma$  in X, which connects p and q. Then  $p_i \rightarrow p, q_i \rightarrow q$ , and the length of geodesics,  $L(\gamma_i) \rightarrow L(\gamma) = d(p, q)$ .

(i)  $0 < d(p,q) \le 1$ . Let us consider the distance function dist<sub>p<sub>i</sub></sub> to point  $p_i$ . Then it converges to dist<sub>p</sub>. Let  $\alpha_i$  be a gradient curve of dist<sub>p<sub>i</sub></sub> such that  $\alpha_i(0) \in M_i$  and  $\alpha_i(0)$  converges to  $\gamma(\epsilon) \ne p$  as  $\overline{\Omega}_i$  GH-converges to X. Since  $\Omega_i$  is collapsing, such  $\alpha_i$  always exists for any  $\epsilon > 0$  and  $\alpha_i$  lies in  $M_i$ .

Since  $\gamma$  is the gradient curve of dist<sub>*p*</sub>, by [23, Lemma 2.1.5], the gradient curve  $\alpha_i$  of dist<sub>*p*<sub>i</sub></sub> and  $L(\alpha_i)$  converges to  $\gamma|_{[\epsilon,L(\gamma)]}$  and  $L(\gamma)$  respectively. Therefore, the gradient curve  $\alpha_i$  has length  $\leq L(\gamma) + \epsilon$  for all *i* large.

Let  $\hat{p}_i$  and  $\hat{q}_i$  be the endpoints of  $\alpha_i$ , then for  $\epsilon \to 0$  we have  $\hat{p}_i = \alpha_i(0) \to p$  and  $\hat{q}_i \to q$ . Now by the convexity of  $\Omega_i$ , we are able to use the gradient flow  $\Phi_\epsilon$  of dist $_{p_i}$  (resp. dist $_{q_i}$ ) by a definite time  $\leq 2\epsilon$  to push the minimal geodesic  $[\hat{q}_i q_i]$  (resp.  $[\hat{p}_i p_i]$ ) to a curve  $\beta_{q_i}$  (resp.  $\beta_{p_i}$ ) in  $M_i$ , whose length  $\leq e^{C(\mu) \cdot \frac{\epsilon}{d(p,q)}} \epsilon$ . Since the joint curve by  $\alpha_i$ ,  $\Phi_t(p_i)$ ,  $\Phi_t(\hat{p}_i)$ ,  $\Phi_t(\hat{q}_i)$ ,  $\beta_{p_i}$  and  $\beta_{q_i}$  gives rise to a curve of uniformly bounded length connecting  $p_i$  and  $q_i$  in  $M_i$ , a contradiction to  $d_{M_i}(p_i, q_i) \to \infty$ .

(ii) d(p,q) = 0, i.e., p = q. Since diam(X) = 1, there is a point  $z \in X$  such that  $d(z,p) = \frac{1}{2}$ . Let  $z_i \in \Omega_i$  be a sequence of points converging to z. Then for all i large, the gradient flow  $\Phi_2$  of dist<sub> $z_i$ </sub> pushes the minimal geodesic  $[p_iq_i]$  in  $\Omega_i$  to a curve  $\alpha$  in  $M_i$  with length  $e^{C(\mu)}d(p_i,q_i)$ . Then the gradient curves  $\Phi_t(p_i)$  ( $t \in [0,1]$ ),  $\alpha$  and  $\Phi_t(q_i)$  ( $t \in [0,1]$ ) form a curve connecting  $p_i$  and  $q_i$  in  $M_i$ , whose length is bounded by  $2 + e^{C(\mu)}d(p_i,q_i) \rightarrow 2$ . This contradicts to that  $d_{M_i}(p_i,q_i) \rightarrow \infty$ .

*Remark* 2.2. Note that for closed convex hypersurfaces in  $\mathbb{R}^n$ , diam $(M) \le \frac{\pi}{2}$  diam $(\Omega)$ , see [16, 17], cf. [3, Page 43]. It is interesting if one can give a sharp estimate on the constant in Lemma 2.1.

Proof of Theorem 1.1.

4

Since *M* is the boundary of a convex domain  $\Omega$ , |B| is bounded by |H|. By Theorem 1.2, it suffices to show that, via pinching (1.2),  $|M|^{\frac{1}{n}} ||H||_{\infty}$  has a uniform upper bound.

Let us argue by contradiction. Let  $\Omega_i$  be a sequence of convex domains in  $N_i$  such that  $|\partial \Omega_i|^{\frac{1}{n}} ||H_{\partial \Omega_i}||_{\infty} \to \infty$ .

By Gauss equaton, the convexity of  $\Omega_i$  implies that  $\partial \Omega_i$  is also a Riemannian manifold of sectional curvature  $\geq \mu$ . By Lemma 2.1, we have

$$\operatorname{diam}(\partial \Omega_i) \leq C(n, \mu, R) \operatorname{diam}(\Omega_i).$$

By Bishop's volume comparison theorem, the volume  $|\partial \Omega_i|$  is also uniformly bounded from above.

By the discussion above, we may rescale  $\Omega_i$  such that  $|\partial \Omega_i| = 1$  and the rescaled sectional curvature of  $\Omega_i$  still admits a uniform lower bound. By passing to a subsequence,  $\partial \Omega_i$  *GH*-converges to *Y*. According to Cheeger-Colding [6], the 1st-eigenvalues  $\lambda_1(\partial \Omega_i)$ converge to  $\lambda_1(Y)$ , the 1st-eigenvalue of *Y*, which by [8] is well-defined. Then the pinching condition

$$n(\delta + ||H_{\partial\Omega_i}||_{\infty}^2) \le \lambda_1(\partial\Omega_i)(1+\epsilon) \to \lambda_1(Y)(1+\epsilon)$$

implies that  $||H_{\partial\Omega_i}||_{\infty}$  is bounded uniformly. This contradicts to the choice of  $\Omega_i$ .

Acknowledgements. The first author was supported by China Postdoctoral Science Foundation (No.2018M641317). The second author was supported partially by National Natural Science Foundation of China [11871349], [11821101], by research funds of Beijing Municipal Education Commission and Youth Innovative Research Team of Capital Normal University.

### References

- E. Aubry and J.-F. Grosjean, Spectrum of hypersurfaces with small extrinsic radius or large λ<sub>1</sub> in Euclidean spaces, Journal of Functional Analysis, 271 (2016), Issue 5, 1213-1242; arXiv:1210.5690 [math.DG]
- [2] Yu. Burago, M. Gromov, and G. Perelman, A.D. Aleksandrov spaces with curvatures bounded below, (Russian) Uspekhi Mat. Nauk 47 (1992), no. 2(284), 3-51, 222; translation in Russian Math. Surveys 47 (1992), no. 2, 1-58.
- [3] Yu. D. Burago and V. A. Zalgaller, Geometric inequalities, Springer, 1988, 331 pp.
- [4] J. Cheeger, Comparison and finiteness theorems for Riemannian manifolds. PhD thesis, Princeton University, 1967.
- [5] J. Cheeger, *Finiteness theorems for Riemannian manifolds*, American Journal of Mathematics, 1970, 92(1):61-74.
- [6] J. Cheeger and T.H. Colding, On the structure of spaces with Ricci curvature bounded below, II, J. Differential Geom. 54(2000), 13-35; On the structure of spaces with Ricci curvature bounded below, III, J. Differential Geom. 54(2000), 37-74.
- [7] B. Colbois and J.-F. Grosjean, A pinching theorem for the first eigenvalue of the Laplacian on hypersurfaces of the Euclidean space, Comment. Math. Helv. 82(2007), 175-195.
- [8] K. Fukaya, Collapsing of Riemannian manifolds and eigenvalues of Laplace operator, Invent. math. 87 (1987), 517-547.
- [9] R. E. Green and H. Wu, *Lipschitz convergence of Riemannian manifolds*, Pacific J. Math., 1988, 131:119-141.
- [10] M. Gromov, Structures mètriques pour les variètès riemanniennes. (French) [Metric structures for Riemann manifolds]. Textes Mathématiques [Mathematical Texts], 1. CEDIC/Fernand Nathan, Paris, 1981.
- [11] J.-F. Grosjean, J. Roth, Eigenvalue pinching and application to the stability and the almost umbilicity of hypersurfaces, Math. Z. 271(2012), 469-488.
- [12] E. Heintze, *Extrinsic upper bound for*  $\lambda_1$ , Math. Ann. **280**(1988), 389-402.
- [13] Y. Hu and H.W. Xu, An eigenvalue pinching theorem for compact hypersurfaces in a sphere, J. Geom. Anal. 27, (2017), 2472-2489.

### YINGXIANG HU, SHICHENG XU

- [14] Y. Hu and S.C. Xu, *Recognizing shape via 1st eigenvalue, mean curvature and upper curvature bound*, preprint, 2019.
- [15] A. Kasue, A convergence theorem for Riemannian manifolds and some applications, Nagoya Math. J., 1989, 114:21–51.
- [16] N. P. Makuha, A connection between the inner and outer diameters of a general closed convex surface. (Russian) Ukrain. Geometr. Sb. Vyp. 2, 49-51, 1966.
- [17] N. P. Makuha, Supplement to the paper "A connection between the inner and outer diameters of a general convex surface". (Russian) Ukrain. Geometr. Sb. Vyp. 4, 37-39, 1967.
- [18] P. Miao and X. Wang, Boundary effect of Ricci curvature, J. Diff. Geom., 103(1), 59-82, 2016.
- [19] G. Perelman and A. M. Petrunin, Extremal subsets in Aleksandrov spaces and the generalized Liberman theorem, Algebra i Analiz, 5(1):242-256, 1993.
- [20] G. Perelman and A. Petrunin, Quasigeodesics and gradient curves in Alexandrov spaces, preprint, 1994.
- [21] S. Peters, *Convergence of Riemannian manifolds*, Compositio Mathematica, 1987, 62(1):3-16.
- [22] A. Petrunin, Applications of quasigeodesics and gradient curves. In Grove, Karsten (ed.) et al., Comparison geometry. Cambridge: Cambridge University. Math. Sci. Res. Inst. Publ. 30, 203-219. 1997.
- [23] A. Petrunin, Semiconcave functions in Alexandrov's geometry, Surveys in differential geometry. Vol. XI, 137-201, Surv. Differ. Geom., 11, Int. Press, Somerville, MA, 2007.
- [24] R. Reilly, On the first eigenvalue of the Laplacian for compact submanifolds of Euclidean space, Comment. Math. Helv. 52(1977), 525-533.
- [25] V. Schroeder and M. Strake, Rigidity of convex domains in manifolds with nonnegative Ricci and sectional curvature, Comment. Math. Helvetici 64 (1989) 173-186.

YAU MATHEMATICAL SCIENCES CENTER, TSINGHUA UNIVERSITY, BEIJING, CHINA *E-mail address*: huyingxiang@mail.tsinghua.edu.cn

MATHEMATICS DEPARTMENT, CAPITAL NORMAL UNIVERSITY, BEIJING, CHINA *E-mail address*: shichengxu@gmail.com

6