

FROBENIUS MANIFOLDS AND A NEW CLASS OF EXTENDED AFFINE WEYL GROUPS $\widetilde{W}^{(k,k+1)}(A_l)$

DAFENG ZUO

ABSTRACT. We present a new class of extended affine Weyl groups $\widetilde{W}^{(k,k+1)}(A_l)$ for $1 \leq k < l$ and obtain an analogue of Chevalley-type theorem for their invariants. We further show the existence of Frobenius manifold structures on the orbit spaces of $\widetilde{W}^{(k,k+1)}(A_l)$ and also construct Landau–Ginzburg superpotentials for these Frobenius manifold structures.

CONTENTS

1. Introduction	2
2. A new class of extended affine Weyl groups $\widetilde{W}^{(k,k+1)}(A_l)$	5
3. A flat pencil of metrics on the orbit space \mathcal{M}	8
4. Frobenius manifold structures on the orbit space \mathcal{M}	15
4.1. The change of coordinates	15
4.2. Flat coordinates of the metric $\eta^{ij}(z)$	16
4.3. Frobenius manifold structures on the orbit space \mathcal{M}	19
4.4. Examples	25
5. The group $\widetilde{W}^{(k,k+1)}(A_l)$ and the Hurwitz space $\mathbb{M}_{k,l-k+1,1}$	30
6. Conclusions	34
References	35

Date: May 27, 2019.

2000 Mathematics Subject Classification. Primary 53D45; Secondary 32M10.

Key words and phrases. Extended affine Weyl group, Frobenius manifold.

1. INTRODUCTION

E.Witten, R.Dijkgraaf, E.Verlinde and H.Verlinde ([7, 8]) in 1990's introduced a remarkable system of partial differential equations, i.e., WDVV equations of associativity, on two dimensional topological field theory (briefly 2dTFT). In order to understand a geometrical foundation of 2dTFT on the bases of WDVV equations, B.Dubrovin ([9, 10]) extended the Atiyah's axioms of 2dTFT ([6]) and invented a nice geometrical object, that is, Frobenius manifold designed as a coordinate-free formulation of WDVV equations.

For an arbitrary n -dimensional Frobenius manifold, B.Dubrovin ([10]) defined a monodromy group, which acts on n -dimensional linear space and can be regarded as (an extension of) a group generated by reflections. The Frobenius manifold itself can be identified with the orbit space of the group in the sense to be specified for each class of monodromy groups, which gives a clue to understanding of Frobenius manifold structure on the orbit space. It was shown by B.Dubrovin ([10, 11]) that any finite Coxeter group can serve as a monodromy group of a polynomial Frobenius manifold, that is to say, the potential is a polynomial with respect to the flat coordinates t^1, \dots, t^l . Furthermore, he put forward the following conjecture, "*Any massive polynomial Frobenius manifold with positive invariant degrees is isomorphic to the orbit space of a finite Coxeter group*", which was proved by C.Hertling ([16]). Besides these, in [13, 18] it has been shown that there are l different Frobenius manifold structures on the orbit spaces of the Coxeter groups B_l and D_l . Especially, we also proved in [18] that the corresponding potentials are meromorphic along the divisors $t^k = 0$ for a given $1 \leq k \leq l - 2$, where t^1, \dots, t^l are the flat coordinates of the Frobenius manifold.

Let R be an irreducible reduced root system defined in an l -dimensional Euclidean space V with Euclidean inner product $(\ , \)$, we fix a basis of simple roots $\alpha_1, \dots, \alpha_l$ and denote by α_j^\vee , $j = 1, 2, \dots, l$ the corresponding coroots. The Weyl group $W(R)$ is generated by the reflections $\mathbf{x} \mapsto \mathbf{x} - (\alpha_j^\vee, \mathbf{x})\alpha_j$, $\forall \mathbf{x} \in V$, $j = 1, \dots, l$. The semi-direct product of $W(R)$ by the lattice of coroots yields the affine Weyl group $W_a(R)$ that acts on V by the affine transformations

$$\mathbf{x} \mapsto w(\mathbf{x}) + \sum_{j=1}^l m_j \alpha_j^\vee, \quad w \in W, \quad m_j \in \mathbb{Z}.$$

We denote by $\omega_1, \dots, \omega_l$ the fundamental weights defined by the relations

$$(\omega_i, \alpha_j^\vee) = \delta_{ij}, \quad i, j = 1, \dots, l.$$

B.Dubrovin and Y.Zhang in [12] (also [19]) defined an *extended affine Weyl group* $\widetilde{W}^{(k)}(R)$, which acts on the extended space $V \oplus \mathbb{R}$ and is generated by the transformations

$$x = (\mathbf{x}, x_{l+1}) \mapsto (w(\mathbf{x}) + \sum_{j=1}^l m_j \alpha_j^\vee, x_{l+1}), \quad w \in W, \quad m_j \in \mathbb{Z},$$

and

$$x = (\mathbf{x}, x_{l+1}) \mapsto (\mathbf{x} + \gamma \omega_k, x_{l+1} - \gamma).$$

Here $1 \leq k \leq l$, $\gamma = 1$ except for the cases when $R = B_l, k = l$ and $R = F_4, k = 3$ or $k = 4$, in these three cases $\gamma = 2$. For a particular choice of a simple root α_k , they proved an analogue of Chevalley theorem for their invariants. On the orbit space of $\widetilde{W}^{(k)}(R)$, they constructed a Frobenius manifold structure whose potential is a weighted homogeneous polynomial of $t^1, \dots, t^{l+1}, e^{t^{l+1}}$, where t^1, \dots, t^{l+1} are the flat coordinates of the Frobenius manifold.

Observe that for the root system of type A_l , there is in fact no restrictions on the choice of α_k . However, for the root systems of type $B_l, C_l, D_l, E_6, E_7, E_8, F_4, G_2$ there is only one choice for each. In [14] Slodowy pointed out that the Chevalley type theorem is a consequence of the results of Looijenga and Wirthmüller [2, 3, 5], and in fact it holds true for any choice of the base element α_k . A natural question is that “*Whether the geometric structures that were revealed by Dubrovin-Zhang’s construction also exist on the orbit spaces of the extended affine Weyl groups for an arbitrary choice of α_k ?*” Our recent work in [19] is to give an affirmative answer to this question for the root systems of type B_l, C_l and also for D_l . We show, by fixing another integer $0 \leq m \leq l - k$, that on the corresponding orbit spaces there also exist Frobenius manifold structures with potentials $F(t)$ that are weighted homogeneous polynomials w.r.t $t^1, \dots, t^{l+1}, \frac{1}{t^{l-m}}, \frac{1}{t^l}, e^{t^{l+1}}$. We also construct Landau–Ginzburg (briefly LG) superpotentials for these Frobenius manifold structures.

In this paper we will present a new extension of affine Weyl groups denoted by $\widetilde{W}^{(k,k+1)}(R)$, which is different from those in [12, 19], and study Frobenius manifold structures on the corresponding orbit spaces $\mathcal{M}^{(k,k+1)}(R)$, where the new extended affine Weyl groups $\widetilde{W}^{(k,k+1)}(R)$ act on the extended space $V \oplus \mathbb{R}^2$ generated by the transformations

$$x = (\mathbf{x}, x_{l+1}, x_{l+2}) \mapsto (w(\mathbf{x}) + \sum_{j=1}^l m_j \alpha_j^\vee, x_{l+1}, x_{l+2}), \quad w \in W, \quad m_j \in \mathbb{Z},$$

and

$$x = (\mathbf{x}, x_{l+1}, x_{l+2}) \mapsto (\mathbf{x} + \gamma \omega_k, x_{l+1} - \gamma, x_{l+2})$$

and

$$x = (\mathbf{x}, x_{l+1}, x_{l+2}) \mapsto (\mathbf{x} + \gamma \omega_{k+1}, x_{l+1}, x_{l+2} - \gamma)$$

Here $1 \leq k \leq l-1$, $\gamma = 1$ except for the cases when $R = B_l, k = l$ and $R = F_4, k = 3$ or $k = 4$, in these three cases $\gamma = 2$.

By a direct verification, we could not obtain any flat pencil of metrics and Frobenius manifold structures on the orbit spaces $\mathcal{M}^{(1,2)}(B_2)$, $\mathcal{M}^{(1,2)}(C_2)$ and $\mathcal{M}^{(1,2)}(G_2)$ etc. We thus have to restrict our study to the type A_l case, i.e. $\mathcal{M}^{(k,k+1)}(A_l)$ and will show that (see Theorem 4.5)

Main Theorem 1. *For any fixed integer $1 \leq k < l$, there exists a unique Frobenius manifold structure of charge $d = 1$ on the orbit space $\mathcal{M}^{(k,k+1)}(A_l) \setminus \{\tilde{y}_{l+1} = 0\} \cup \{\tilde{y}_{l+2} = 0\}$ of $\widetilde{W}^{(k,k+1)}(A_l)$ such that*

- (1) *the invariant flat metric and the intersection form of the Frobenius manifold structure coincide with the metrics $(\eta^{ij}(y))$ in (3.18) and $(g^{ij}(y))$ in (3.5) respectively;*
- (2) *the unity and the Euler vector fields have the form*

$$e = \frac{\partial}{\partial y^k} + \frac{\partial}{\partial y^{k+1}} \quad (1.1)$$

and

$$E = \sum_{\alpha=1}^l d_\alpha y^\alpha \frac{\partial}{\partial y^\alpha} + \frac{1}{k} \frac{\partial}{\partial y^{l+1}} + \frac{1}{l-k} \frac{\partial}{\partial y^{l+2}}, \quad (1.2)$$

where d_1, \dots, d_l are defined in (2.13);

- (3) *in the flat coordinates t^1, \dots, t^{l+2} of the metric (3.18) defined on certain covering of $\mathcal{M}^{(k,k+1)}(A_l)$ the potential of the Frobenius manifold structure is of the form $F(t) = \widehat{F}(t) + \frac{1}{2}(t^{k+1})^2 \log(t^{k+1})$, where $\widehat{F}(t)$ is a weighted homogeneous polynomial in $t^1, t^2, \dots, t^{l+2}, e^{t^{l+1}}, e^{t^{l+2}-t^{l+1}}$.*

On the orbit space of the extended affine Weyl group $\widetilde{W}^{(k)}(A_l)$, an alternative construction of the Frobenius manifold structure was given in [12]. This structure was given in terms of a LG superpotential construction. In particular, it was shown that $\widetilde{W}^{(k)}(A_l)$ describes the monodromy of roots of trigonometric polynomials - the superpotential - with a given bidegree being of the form

$$\lambda(\varphi) = e^{i k \varphi} + a_1 e^{i(k-1)\varphi} + \dots + a_{l+1} e^{i(k-l-1)\varphi}, \quad a_{l+1} \neq 0.$$

A natural question is that

“Whether a similar construction about the Frobenius manifold structure exists on the orbit space $\mathcal{M}^{(k,k+1)}(A_l) \setminus \{\tilde{y}_{l+1} = 0\} \cup \{\tilde{y}_{l+2} = 0\}$ of the extended affine Weyl group $\widetilde{W}^{(k,k+1)}(A_l)$?”

Let $\mathbb{M}_{k,l-k+1,1}$ be the space of a particular class of LG superpotentials consisting of trigonometric-Laurent series of one variable with tri-degree $(k+1, l-k, 1)$, these being functions of the form

$$\lambda(\varphi) = (e^{i\varphi} - a_{l+2})^{-1} (e^{i(k+1)\varphi} + a_1 e^{ik\varphi} + \cdots + a_{l+1} e^{i(k-l)\varphi}), \quad a_{l+1} a_{l+2} \neq 0,$$

where $a_j \in \mathbb{C}$ for $j = 1, \dots, l+2$. The space $\mathbb{M}_{k,l-k+1,1}$ carries a natural structure of Frobenius manifold. Its invariant inner product η and the intersection form g of two vectors ∂', ∂'' tangent to $\mathbb{M}_{k,l-k+1,1}$ at a point $\lambda(\varphi)$ can be defined by the formulae (5.2) and (5.3). We will show that (see Theorem 5.1)

Main Theorem 2. *The Frobenius manifolds $\mathcal{M}^{(k,k+1)}(A_l) \setminus \{\tilde{y}_{l+1} = 0\} \cup \{\tilde{y}_{l+2} = 0\}$ and $\mathbb{M}_{k,l-k+1,1}$ are locally isomorphic.*

2. A NEW CLASS OF EXTENDED AFFINE WEYL GROUPS $\widetilde{W}^{(k,k+1)}(A_l)$

To keep self-contained, we recall some known facts about Weyl groups of type A_l , see [4] for details. Let \mathbb{R}^{l+1} be a $(l+1)$ -dimensional Euclidean space with Euclidean inner product (\cdot, \cdot) and an orthonormal basis $\epsilon_1, \dots, \epsilon_{l+1}$. Let A_l be an irreducible reduced root system in the hyperplane $V = \left\{ \sum_{s=1}^{l+1} v_s \epsilon_s \in \mathbb{R}^{l+1} \mid \sum_{s=1}^{l+1} v_s = 0 \right\}$.

We fix a basis

$$\alpha_1 = \epsilon_1 - \epsilon_2, \dots, \alpha_l = \epsilon_l - \epsilon_{l+1}$$

of simple roots. The corresponding coroots are $\alpha_j^\vee = \alpha_j$ for $j = 1, \dots, l$. The Weyl group $W = W(A_l)$ is generated by the reflections

$$\mathbf{x} \mapsto \mathbf{x} - (\alpha_j^\vee, \mathbf{x}) \alpha_j, \quad \forall \mathbf{x} \in V, \quad j = 1, \dots, l. \quad (2.1)$$

W acts on V by permutations of the coordinates v_1, \dots, v_{l+1} . The basic W -invariant Fourier polynomials coincide with the elementary symmetric functions

$$y_j(\mathbf{x}) = \sigma_j(e^{2\pi i v_1}, \dots, e^{2\pi i v_{l+1}}), \quad j = 1, \dots, l. \quad (2.2)$$

Definition 2.1. *For any fixed integer $1 \leq k < l$, we call $\widetilde{W} = \widetilde{W}^{(k,k+1)}(A_l)$ to be an extended affine Weyl group of type A if it acts on*

$$\widetilde{V} = V \oplus \mathbb{R}^2$$

generated by the transformations

$$x = (\mathbf{x}, x_{l+1}, x_{l+2}) \mapsto (w(\mathbf{x}) + \sum_{j=1}^l m_j \alpha_j^\vee, x_{l+1}, x_{l+2}), \quad w \in W, \quad m_j \in \mathbb{Z}, \quad (2.3)$$

and

$$x = (\mathbf{x}, x_{l+1}, x_{l+2}) \mapsto (\mathbf{x} + \omega_k, x_{l+1} - 1, x_{l+2}), \quad (2.4)$$

and

$$x = (\mathbf{x}, x_{l+1}, x_{l+2}) \mapsto (\mathbf{x} + \omega_{k+1}, x_{l+1}, x_{l+2} - 1). \quad (2.5)$$

Coordinates x_1, \dots, x_l may be introduced on the space V via the expression

$$\mathbf{x} = x_1 \alpha_1^\vee + \dots + x_l \alpha_l^\vee. \quad (2.6)$$

That is to say,

$$v_1 = x_1, \quad v_i = x_j - x_{j-1}, \quad v_{l+1} = -x_l, \quad j = 2, \dots, l. \quad (2.7)$$

Definition 2.2. $\mathcal{A} = \mathcal{A}^{(k,k+1)}(A_l)$ is the ring of all \widetilde{W} -invariant Fourier polynomials of $x_1, \dots, x_l, \frac{1}{l+1}x_{l+1}, \frac{1}{l+1}x_{l+2}$ that are bounded in the following limit conditions

$$\mathbf{x} = \mathbf{x}^0 - i\omega_k \tau, \quad x_{l+1} = x_{l+1}^0 + i\tau, \quad x_{l+2} = x_{l+2}^0, \quad \tau \rightarrow +\infty \quad (2.8)$$

and

$$\mathbf{x} = \mathbf{x}^0 - i\omega_{k+1} \tau, \quad x_{l+1} = x_{l+1}^0, \quad x_{l+2} = x_{l+2}^0 + i\tau, \quad \tau \rightarrow +\infty \quad (2.9)$$

for any $x^0 = (\mathbf{x}^0, x_{l+1}^0, x_{l+2}^0)$.

Conditions (2.8) and (2.9) are essential for this construction as did in [12, 19]. For simplicity, we introduce a set of numbers

$$d_{j,k} := (\omega_j, \omega_k) = \begin{cases} \frac{j(l-k+1)}{l+1}, & j = 1, \dots, k, \\ \frac{k(l-j+1)}{l+1}, & j = k+1, \dots, l \end{cases} \quad (2.10)$$

and define the following Fourier polynomials

$$\begin{aligned} \widetilde{y}_j(x) &= e^{2\pi i(d_{j,k} x_{l+1} + d_{j,k+1} x_{l+2})} y_j(\mathbf{x}), \quad j = 1, \dots, l, \\ \widetilde{y}_{l+1}(x) &= e^{2\pi i x_{l+1}}, \quad \widetilde{y}_{l+2}(x) = e^{2\pi i x_{l+2}}. \end{aligned} \quad (2.11)$$

Lemma 2.3. ([12]) For any fixed integer $1 \leq k_r \leq l$, we have

$$y_j(\mathbf{x}) = e^{2\pi d_{j,k_r} \tau} [y_j^{0,r}(\mathbf{x}^0) + \mathcal{O}(e^{-2\pi\tau})], \quad \tau \rightarrow +\infty, \quad \text{for } j = 1, \dots, l,$$

where $\mathbf{x} = \mathbf{x}^0 - i\omega_{k_r} \tau$ and

$$y_j^{0,r}(\mathbf{x}^0) = \frac{1}{n_j} \sum_{w \in W, (w(\omega_j) - \omega_j, \omega_{k_r}) = 0} e^{2\pi i (w(\omega_j), \mathbf{x}^0)},$$

where $n_j = \#\{w \in W | e^{2\pi i (w(\omega_j), \mathbf{x})} = e^{2\pi i (\omega_j, \mathbf{x})}\}$. Moreover, the Fourier polynomials $y_1^{0,r}(\mathbf{x}^0), \dots, y_l^{0,r}(\mathbf{x}^0)$ are algebraically independent.

From these explicit expressions in (2.11) and Lemma 2.3, it is not difficult to see that $\widetilde{y}_j(x) \in \mathcal{A}$ for $j = 1, \dots, l+2$. Furthermore, we have

Theorem 2.4. (*Chevalley-type theorem*) *The ring \mathcal{A} is isomorphic to the ring of polynomials of $\tilde{y}_1(x), \dots, \tilde{y}_{l+2}(x)$.*

Proof. Observe that $\tilde{y}_1(x), \dots, \tilde{y}_{l+2}(x)$ are algebraically independent. So in order to prove the theorem, we only need to show that any element $f(x)$ of the ring \mathcal{A} can be represented as a polynomial of $\tilde{y}_1(x), \dots, \tilde{y}_{l+2}(x)$. From the invariance with respect to \widetilde{W} , it follows that $f(x)$ can be represented as a polynomial of $\tilde{y}_1(x), \dots, \tilde{y}_{l+1}(x), \tilde{y}_{l+2}(x), \tilde{y}_{l+1}^{-1}(x), \tilde{y}_{l+2}^{-1}(x)$. It suffices to show that in $f(x)$ there are no negative powers of $\tilde{y}_{l+1}(x)$ and $\tilde{y}_{l+2}(x)$.

Assume that

$$f(x) = \sum_{s \geq -S} \tilde{y}_{l+1}^s \sum_{t \in \Lambda} \tilde{y}_{l+2}^t Q_{s,t}(\tilde{y}_1(x), \dots, \tilde{y}_l(x)) \quad (2.12)$$

for a positive integer S and the polynomial $Q_{-S,t_0}(\tilde{y}_1(x), \dots, \tilde{y}_l(x))$ does not vanish identically, where $t_0 = \min\{t \in \Lambda \mid Q_{-S,t} \text{ does not vanish identically}\}$ and Λ is a finite subset of \mathbb{Z} . With the use of Lemma 2.3, in the limit (2.8) the function $f(x)$ behaves as

$$f(x) = e^{2\pi S\tau - 2\pi i S x_{l+1}^0} \sum_{t \in \Lambda} e^{2\pi i t x_{l+2}^0} [Q_{s,t}(\tilde{y}_1^{0,1}(x^0), \dots, \tilde{y}_l^{0,1}(x^0)) + \mathcal{O}(e^{-2\pi\tau})],$$

where $\tilde{y}_j^{0,1}(x^0) = e^{2\pi i(d_{j,k}x_{l+1}^0 + d_{j,k+1}x_{l+2}^0)} y_j^{0,1}(\mathbf{x}^0)$ for $j = 1, \dots, l$. In order to assure the function $f(x)$ bounded for $\tau \mapsto +\infty$, it is necessary to have

$$Q_{-S,t_0}(\tilde{y}_1^{0,1}(x^0), \dots, \tilde{y}_l^{0,1}(x^0)) \equiv 0,$$

which is a contradiction with the algebraic independence of $\tilde{y}_1^{0,1}(x^0), \dots, \tilde{y}_l^{0,1}(x^0)$. This means that there are no negative powers of $\tilde{y}_{l+1}(x)$. Similarly one can show that there are no negative powers of $\tilde{y}_{l+2}(x)$. This completes the proof of the theorem. \square

Corollary 2.5. *The function \deg defined as*

$$\begin{aligned} \deg \tilde{y}_{l+1} &= \frac{1}{k}, & \deg \tilde{y}_{l+2} &= \frac{1}{l-k}, \\ d_j := \deg \tilde{y}_j &= \frac{d_{j,k}}{k} + \frac{d_{j,k+1}}{l-k} = \begin{cases} \frac{j}{k}, & j = 1, \dots, k, \\ \frac{l-j+1}{l-k}, & j = k+1, \dots, l \end{cases} \end{aligned} \quad (2.13)$$

determines on \mathcal{A} a structure of graded polynomial ring. Especially,

$$d_k = d_{k+1} = 1 > d_s, \quad s \neq k, k+1. \quad (2.14)$$

The numbers d_1, \dots, d_{l+2} with $d_{l+1} = d_{l+2} = 0$ satisfy a duality relation. For any given integer k , we denote $A_l \setminus \{\alpha_k, \alpha_{k+1}\} = \mathcal{R}_1 \cup \mathcal{R}_2$, where $\mathcal{R}_1 = \{\alpha_1, \dots, \alpha_{k-1}\}$

and $\mathcal{R}_2 = \{\alpha_{k+2}, \dots, \alpha_l\}$. On each component we have an involution $j \mapsto j^*$ given by the reflection with respect to the center of the component. Let us define

$$k^* = l + 1, (k + 1)^* = l + 2, (l + 2)^* = k + 1, (l + 1)^* = k, \quad (2.15)$$

then

$$d_j + d_{j^*} = 1, \quad j = 1, \dots, l + 2. \quad (2.16)$$

3. A FLAT PENCIL OF METRICS ON THE ORBIT SPACE \mathcal{M}

Let us denote $\mathcal{M}^{(k,k+1)} := \widetilde{V} \otimes \mathbb{C}/\widetilde{W}$, called the *orbit space* of the extended Weyl group \widetilde{W} . We define an indefinite flat metric $(dx_i, dx_j)^\sim$ on $\widetilde{V}_{\mathbb{C}} = \widetilde{V} \otimes_{\mathbb{R}} \mathbb{C}$ where \widetilde{V} is the orthogonal direct sum of V and \mathbb{R}^2 . Here V is endowed with the W -invariant Euclidean metric ¹

$$(dx_a, dx_b)^\sim = \frac{1}{4\pi^2}(\omega_a, \omega_b), \quad 1 \leq a, b \leq l \quad (3.1)$$

and \mathbb{R}^2 is endowed with the metric

$$(dx_{l+1}, dx_{l+1})^\sim = -\frac{\tau_{11}}{4\pi^2}, \quad (dx_{l+1}, dx_{l+2})^\sim = -\frac{\tau_{12}}{4\pi^2}, \quad (dx_{l+2}, dx_{l+2})^\sim = -\frac{\tau_{22}}{4\pi^2}, \quad (3.2)$$

where

$$\begin{pmatrix} \tau_{11} & \tau_{12} \\ \tau_{12} & \tau_{22} \end{pmatrix} = \begin{pmatrix} d_{k,k} & d_{k,k+1} \\ d_{k+1,k} & d_{k+1,k+1} \end{pmatrix}^{-1} = \begin{pmatrix} \frac{k+1}{k} & -1 \\ -1 & \frac{l-k+1}{l-k} \end{pmatrix}.$$

The set of generators for the ring \mathcal{A} are defined by (2.11). They form a system of global coordinates on $\mathcal{M}^{(k,k+1)}$. We now introduce a system of local coordinates on $\mathcal{M}^{(k,k+1)}$ as follows

$$y^1 = \widetilde{y}_1, \dots, y^l = \widetilde{y}_l, y^{l+1} = \log \widetilde{y}_{l+1} = 2\pi i x_{l+1}, y^{l+2} = \log \widetilde{y}_{l+2} = 2\pi i x_{l+2}. \quad (3.3)$$

They live on the universal covering $\widetilde{\mathcal{M}}$ of \mathcal{M} , where $\mathcal{M} := \mathcal{M}^{(k,k+1)} \setminus \{\widetilde{y}_{l+1} = 0\} \cup \{\widetilde{y}_{l+2} = 0\}$. The projection

$$\text{Pr} : \widetilde{V} \rightarrow \widetilde{\mathcal{M}}, \quad (x_1, \dots, x_{l+2}) \mapsto (y^1, \dots, y^{l+2}) \quad (3.4)$$

induces a symmetric bilinear form on $T^*\widetilde{\mathcal{M}}$

$$(dy^i, dy^j)^\sim \equiv g^{ij}(y) := \sum_{a,b=1}^{l+2} \frac{\partial y^i}{\partial x_a} \frac{\partial y^j}{\partial x_b} (dx_a, dx_b)^\sim. \quad (3.5)$$

¹As is common in the Frobenius manifold literature, we use the word metric to denote a complex-valued, symmetric, non-degenerate, bilinear form.

Lemma 3.1. *The matrix entries $g^{ij}(y)$ of (3.5) are weighted homogeneous polynomials in $y^1, \dots, y^l, e^{y^{l+1}}, e^{y^{l+2}}$ of the degree $\deg g^{ij}(y) = \deg y^i + \deg y^j$, here $\deg y^{l+1+\nu} = d_{l+1+\nu} = 0$ for $\nu = 0, 1$. The matrix $(g^{ij}(y))$ does not degenerate outside the Pr-images of the hyperplanes*

$$\{(\mathbf{x}, x_{l+1}, x_{l+2}) | (\beta, \mathbf{x}) = m \in \mathbb{Z}, \forall x_{l+1}, \forall x_{l+2}\}, \quad \beta \in \Phi^+,$$

where Φ^+ is the set of the all positive roots.

Proof. With the use of (3.5), (2.10) and (2.13), we obtain

$$\begin{aligned} g^{j,l+1}(y) &= \zeta_j d_j y^j, & g^{j,l+2}(y) &= (1 - \zeta_j) d_j y^j, & j &= 1, \dots, l, \\ g^{l+1,l+1}(y) &= \frac{k+1}{k}, & g^{l+1,l+2}(y) &= -1, & g^{l+2,l+2}(y) &= \frac{l-k+1}{l-k}, \end{aligned} \quad (3.6)$$

where $\zeta_j = \begin{cases} 1, & 1 \leq j \leq k, \\ 0, & k+1 \leq j \leq l. \end{cases}$ Also, for $1 \leq i, j \leq l$ we have

$$g^{ij}(y) = c_{ij} y^i y^j + \frac{1}{4\pi^2} \sum_{p,q=1}^l \frac{\partial y^i}{\partial x_p} \frac{\partial y^j}{\partial x_q} (\omega_p, \omega_q), \quad c_{ij} = (\zeta_j d_{i,k} + (1 - \zeta_j) d_{i,k+1}) d_j, \quad (3.7)$$

which are Fourier polynomials invariant with respect to \widetilde{W} and bounded in the limits (2.8) and (2.9). It follows from Theorem 2.4 and (3.5) that $g^{ij}(y)$ are weighted homogeneous polynomials in $y^1, \dots, y^l, e^{y^{l+1}}, e^{y^{l+2}}$ of the degree $\deg g^{ij}(y) = \deg y^i + \deg y^j$.

Observe that the Jacobian of the projection map (3.4) is given by

$$\begin{aligned} \det \left(\frac{\partial y^j}{\partial x_a} \right) &= -4\pi^2 e^{2\pi i \sum_{j=1}^l (d_{j,k} x_{l+1} + d_{j,k+1} x_{l+2})} \det \left(\frac{\partial y_j(\mathbf{x})}{\partial x_p} \right) \\ &= c e^{2\pi i \sum_{j=1}^l (d_{j,k} x_{l+1} + d_{j,k+1} x_{l+2})} J(\mathbf{x}), \end{aligned} \quad (3.8)$$

where $J(\mathbf{x}) = e^{-\sum_{\alpha \in \Phi^+} \pi i(\alpha, \mathbf{x})} \prod_{\beta \in \Phi^+} (e^{2\pi i(\beta, \mathbf{x})} - 1)$ and c is a nonzero constant ([4]). So

the projection map (3.4) is a local diffeomorphism outside the above hyperplanes, which assures the nondegeneracy of $(g^{ij}(y))$. \square

Lemma 3.2. *For $k \leq i, j \leq k+1$, the term $y^i y^j$ only possibly appears in $g^{ij}(y)$ and $g^{ji}(y)$ with the coefficient $c_{ij} - d_{i,j}$, where $c_{ij} = (\zeta_j d_{i,k} + (1 - \zeta_j) d_{i,k+1}) d_j$ and $d_{i,j} = (\omega_i, \omega_j)$.*

Proof. From (3.7), we have for $k \leq i, j \leq k+1$

$$g^{ij}(y) = e^{2\pi i[(d_{i,k} + d_{j,k})x_{l+1} + (d_{i,k+1} + d_{j,k+1})x_{l+2}]} (c_{ij} y_i(\mathbf{x}) y_j(\mathbf{x}) + \beta_{ij}(\mathbf{x})), \quad (3.9)$$

where

$$\begin{aligned}\beta_{ij}(\mathbf{x}) &= \frac{1}{4\pi^2} \sum_{p,q=1}^l \frac{\partial y_i(\mathbf{x})}{\partial x_p} \frac{\partial y_j(\mathbf{x})}{\partial x_q}(\omega_p, \omega_q) \\ &= -\frac{1}{n_i n_j} \sum_{w, w' \in W} e^{2\pi i(w(\omega_i) + w'(\omega_j), \mathbf{x})}(w(\omega_i), w'(\omega_j)).\end{aligned}$$

Now we use the standard partial ordering of the weights (see the page 69 in [1])

$$\omega \succ \omega' \quad \text{iff} \quad \omega - \omega' = \sum_{m=1}^l c_m \alpha_m$$

for some nonnegative integers c_1, \dots, c_l . In this case, we will write $e^{2\pi i(\omega, \mathbf{x})} \succ e^{2\pi i(\omega', \mathbf{x})}$. All the terms in the W -invariant Fourier polynomials $\beta_{ij}(\mathbf{x})$ are strictly less than $e^{2\pi i(\omega_i + \omega_j, \mathbf{x})}$ except the terms $-d_{i,j} e^{2\pi i(\omega_i + \omega_j, \mathbf{x})}$. So the term $y^i y^j$ possibly appears in $g^{ij}(y)$ and $g^{ji}(y)$ with the coefficient $c_{ij} - d_{i,j}$.

Observe that

$$\frac{g^{kk}(y)}{y^k y^{k+1}} = e^{2\pi i(\frac{k}{l+1}x_{l+1} - \frac{l-k}{l+1}x_{l+2})} \left(\frac{c_{kk} y_k(\mathbf{x})}{y_{k+1}(\mathbf{x})} + \frac{\beta_{kk}(\mathbf{x})}{y_k(\mathbf{x}) y_{k+1}(\mathbf{x})} \right)$$

and

$$\frac{g^{kk}(y)}{(y^{k+1})^2} = e^{2\pi i(\frac{k}{l+1}x_{l+1} - \frac{l-k}{l+1}x_{l+2})} \frac{c_{kk}(y_k(\mathbf{x}))^2 + \beta_{kk}(\mathbf{x})}{(y_{k+1}(\mathbf{x}))^2}.$$

So $y^k y^{k+1}$ and $(y^{k+1})^2$ do not appear in $g^{kk}(y)$. Similarly, we could prove the other cases. □

Lemma 3.3. *Denote*

$$e = \varsigma_1 \frac{\partial}{\partial y^k} + \varsigma_2 \frac{\partial}{\partial y^{k+1}}, \quad \varsigma_1, \varsigma_2 \in \mathbb{R}, \quad (3.10)$$

then for $1 \leq i, j \leq l+2$,

$$\mathcal{L}_e(\mathcal{L}_e g^{ij}(y)) = 0, \quad (3.11)$$

where \mathcal{L}_e is the Lie derivative along the vector field e .

Proof. According to the weighted homogeneity and (2.14) and (3.6), it suffices to show that $k \leq i, j \leq k+1$,

$$\frac{\partial^2}{\partial y^k \partial y^k} g^{ij}(y) = \frac{\partial^2}{\partial y^{k+1} \partial y^{k+1}} g^{ij}(y) = \frac{\partial^2}{\partial y^k \partial y^{k+1}} g^{ij}(y) = 0. \quad (3.12)$$

It follows from

$$c_{k(k+1)} = d_{k,k+1}, \quad c_{(k+\nu)(k+\nu)} = d_{k+\nu, k+\nu}$$

that $g^{k+\nu, k+\nu}(y)$ does not contain $(y^{k+\nu})^2$ for $\nu = 0, 1$, and $g^{k, k+1}(y)$ ($= g^{k+1, k}(y)$) does not contain $y^k y^{k+1}$. Combining with Lemma 3.2, we obtain the desired (3.12) and complete the proof of this lemma. \square

Corollary 3.4. For $1 \leq i, j \leq l+2$,

$$g^{ij}(\dots, y^k + \varsigma_1 \lambda, y^{k+1} + \varsigma_2 \lambda, \dots)$$

are linear in the parameter λ .

Suppose Σ is the discriminant of $\mathcal{M}^{(k, k+1)}$, i.e., $\Sigma = \{y \mid \det(g^{ij}(y)) = 0\}$, then on $\mathcal{M}^{(k, k+1)} \setminus \Sigma$ the matrix $(g^{ij}(y))$ is invertible. The inverse matrix $(g_{ij}(y))$ determines a flat metric on $\mathcal{M}^{(k, k+1)} \setminus \Sigma$. Let us now compute the coefficients of the correspondent Levi-Civita connection ∇ for the metric $g_{ij}(y)$. It is convenient to consider the contravariant components of the connection

$$\Gamma_m^{ij}(y) = (dy^i, \nabla_m dy^j),$$

which are related to the standard Christoffel coefficients by the formula

$$\Gamma_m^{ij}(y) = -g^{is}(y)\Gamma_{sm}^j(y).$$

For the contravariant components, we have the following formulae

$$\Gamma_m^{ij}(y) dy^m = \frac{\partial y^i}{\partial x_a} \frac{\partial^2 y^j}{\partial x_b \partial x_r} (dx_a, dx_b) \sim dx_r \quad (3.13)$$

and

$$2g^{sm}(y)\Gamma_m^{ij}(y) = g^{im}(y) \frac{\partial g^{js}(y)}{\partial y^m} + g^{sm}(y) \frac{\partial g^{ji}(y)}{\partial y^m} - g^{jm}(y) \frac{\partial g^{is}(y)}{\partial y^m} \quad (3.14)$$

and

$$\frac{\partial g^{ij}(y)}{\partial y^m} = \Gamma_m^{ij}(y) + \Gamma_m^{ji}(y). \quad (3.15)$$

Lemma 3.5. $\Gamma_m^{ij}(y)$ are weighted homogeneous polynomials in $y^1, \dots, y^l, e^{y^{l+1}}, e^{y^{l+2}}$ of the degree $\deg \Gamma_m^{ij}(y) = \deg y^i + \deg y^j - \deg y^m$.

Proof. By using (3.13) and (3.8), we can represent

$$(\Gamma_1^{ij}(y), \dots, \Gamma_{l+2}^{ij}(y)) \left(\frac{\partial y^r}{\partial x_a} \right) = \left(\frac{\partial y^i}{\partial x_a} \frac{\partial^2 y^j}{\partial x_b \partial x_1} (dx_a, dx_b) \sim, \dots, \frac{\partial y^i}{\partial x_a} \frac{\partial^2 y^j}{\partial x_b \partial x_{l+2}} (dx_a, dx_b) \sim \right),$$

and

$$\Gamma_m^{ij}(y) = e^{2\pi i \sum_{\nu=0}^1 (d_{i, k+\nu} + d_{j, k+\nu} - d_{m, k+\nu}) x_{l+1+\nu}} \frac{P_m^{ij}(\mathbf{x})}{J(\mathbf{x})}, \quad (3.16)$$

where $d_{l+1+\nu, k+\nu} = 0$ for $\nu = 0, 1$ and $P_m^{ij}(\mathbf{x})$ is certain Fourier polynomial in x_1, \dots, x_l . As discussed the Lemma 2.2 in [12], $P_m^{ij}(\mathbf{x})$ is anti-invariant with

respect to Weyl group W and divisible by $J(\mathbf{x})$. We thus know $\Gamma_m^{ij}(y) \in \mathcal{A}$, whose homogeneity property is obvious. \square

Lemma 3.6. *For $1 \leq i, j \leq l+2$, we have*

$$\mathcal{L}_e(\mathcal{L}_e \Gamma_m^{ij}(y)) = 0. \quad (3.17)$$

Equivalently, $\Gamma_m^{ij}(\dots, y^k + \varsigma_1 \lambda, y^{k+1} + \varsigma_2 \lambda, \dots)$ are linear in the parameter λ .

Proof. By the degrees, it suffices to show that

$$\mathcal{L}_e(\mathcal{L}_e \Gamma_{l+i}^{k+\nu, k+t}(y)) = 0, \quad i = 1, 2, \quad \nu, t = 0, 1.$$

Observe that

$$\begin{aligned} \Gamma_{l+i}^{k+\nu, k+\nu}(y) &= \frac{\partial x_r}{\partial y^{l+i}} \frac{\partial y^{k+\nu}}{\partial x_p} \frac{\partial^2 y^{k+\nu}}{\partial x_q \partial x_r} (dx_p, dx_q) \sim \\ &= \frac{\partial y^{k+\nu}}{\partial x_p} \frac{\partial}{\partial y^{l+i}} \left(\frac{\partial y^{k+\nu}}{\partial x_q} \right) (dx_p, dx_q) \sim \\ &= \frac{1}{2} \frac{\partial}{\partial y^{l+i}} g^{k+\nu, k+\nu}(y), \end{aligned}$$

and using (3.11), then

$$\mathcal{L}_e(\mathcal{L}_e \Gamma_{l+i}^{k+\nu, k+\nu}(y)) = 0$$

for $\nu = 0, 1$ and $i = 1, 2$.

By choosing $s = k, i = k, j = k+1$ in (3.14) and using (3.6), we get

$$\begin{aligned} y^k \Gamma_{l+1}^{k, k+1}(y) &= g^{k, l+1}(y) \Gamma_{l+1}^{k, k+1}(y) + g^{k, l+2}(y) \Gamma_{l+2}^{k, k+1}(y) \\ &= g^{km}(y) \frac{\partial g^{k+1, k}(y)}{\partial y^m} - \frac{1}{2} g^{k+1, m}(y) \frac{\partial g^{kk}(y)}{\partial y^m} - \sum_{j=1}^l g^{kj}(y) \Gamma_j^{k, k+1}(y). \end{aligned}$$

Repeat using the degrees and Lemma 3.2 and Lemma 3.3, we thus conclude

$$\mathcal{L}_e(\mathcal{L}_e \Gamma_{l+1}^{k, k+1}(y)) = 0.$$

Furthermore, with the help of (3.15), we have $\mathcal{L}_e(\mathcal{L}_e \Gamma_{l+1}^{k+1, k}(y)) = 0$.

Similarly, by choosing $s = k+1, i = k, j = k+1$ in (3.14) and using (3.6), (3.16) and (3.15), we have

$$\begin{aligned} y^{k+1} \Gamma_{l+2}^{k, k+1}(y) &= g^{k, l+1}(y) \Gamma_{l+1}^{k, k+1}(y) + g^{k+1, l+2}(y) \Gamma_{l+2}^{k, k+1}(y) \\ &= \frac{1}{2} g^{km}(y) \frac{\partial g^{k+1, k+1}(y)}{\partial y^m} - \sum_{j=1}^l g^{k+1, j}(y) \Gamma_j^{k, k+1}(y) \end{aligned}$$

and

$$\Gamma_{l+2}^{k+1, k}(y) = \frac{\partial g^{k+1, k}}{\partial y^{l+2}} - \Gamma_{l+2}^{k, k+1}(y).$$

So

$$\mathcal{L}_e(\mathcal{L}_e\Gamma_{l+2}^{k,k+1}(y)) = \mathcal{L}_e(\mathcal{L}_e\Gamma_{l+2}^{k+1,k}(y)) = 0.$$

This completes the proof of the lemma. \square

Lemma 3.7. *Setting*

$$\eta^{ij}(y) = \mathcal{L}_e g^{ij}(y) \quad (3.18)$$

and denoting $\mathcal{R}_{k,k+1} = \mathcal{R}_1 \cup \mathcal{R}_2$, then we have

(1) If α_i and α_j belong to different components of $\mathcal{R}_{k,k+1}$, then $\eta^{ij}(y) = 0$;

(2) The block $\eta_{(t)} = (\eta^{ij}(y))|_{\alpha_i, \alpha_j \in \mathcal{R}_t}$ of the matrix $(\eta^{ij}(y))$ corresponding to any branch \mathcal{R}_t has triangular form. The antidiagonal elements of $\eta_{(t)}$ consists of the constant numbers η^{ii^*} for $\alpha_i \in \mathcal{R}_t$, where $t = 1, 2$;

(3) $\eta^{a,l+1}(y) = \varsigma_1 \delta_{a,k}$, $\eta^{a,l+2}(y) = \varsigma_2 \delta_{a,k+1}$ for $a = 1, \dots, l+2$.

Proof. (1) Let $\alpha_i \in \mathcal{R}_1$ and $\alpha_j \in \mathcal{R}_2$, i.e. $1 \leq i < k$ and $k+1 < j \leq l$. As discussed above, if $\beta_{ij}(\mathbf{x})$ as a polynomial in $y_1(\mathbf{x}), \dots, y_l(\mathbf{x})$ contains a monomial $y_1^{p_1} \cdots y_l^{p_l}$ with $p_k = 1$, then

$$\omega_i + \omega_j = p_1 \omega_1 + \cdots + p_l \omega_l + \sum_{s=1}^l q_s \alpha_s \quad (3.19)$$

for some nonnegative integers q_1, \dots, q_l . We multiply (3.19) by ω_1 and obtain

$$l+1+k-i-j - \sum_{s \neq k} p_s(l-s-1) = (l+1)q_1.$$

Since $k-i-j < 0$, then we have $q_1 = 0$ and

$$l-i+1-(j-k) = \sum_{s \neq k} p_s(l-s-1)$$

which yields that $p_s = 0$ for $s = 1, \dots, i$. So (3.19) becomes

$$\omega_i + \omega_j = p_{i+1} \omega_{i+1} + \cdots + p_l \omega_l + \sum_{s=2}^l q_s \alpha_s. \quad (3.20)$$

We multiply (3.20) by $\alpha_1, \dots, \alpha_i$ and get $q_2 = \cdots = q_i = 0$, $q_{i+1} = -1$, which contradicts nonnegativity of q 's. So in this case $\frac{\partial}{\partial y^k} g^{ij}(y) = 0$. Similarly, one can show that $\frac{\partial}{\partial y^{k+1}} g^{ij}(y) = 0$. We thus complete the proof of the first statement.

(2) Observe that in any component of $\mathcal{R}_{k,k+1}$, the numbers d_i are distinct and ordered monotonically and $\deg \eta^{ij}(y) = d_i + d_j - d_k$. We thus conclude that $\eta^{ij}(y) = 0$ when $d_i + d_j = d_k$, and $\eta^{ij}(y) = \text{constant}$ when $d_i + d_j = d_k$ which happens if the labels i and j are dual to each other in the sense of (2.16).

(3) Obviously, the third statement follows from (3.6). \square

Proposition 3.8. *If $\varsigma_1 \varsigma_2 \neq 0$, then the determinant of $(\eta^{ij}(y))$ is a nonzero constant.*

Proof. By using Lemma 3.7, we know that

$$\det(\eta^{ij}(y)) = (-1)^{\frac{k(k+1)+(l-k)(l-k+1)}{2}} \prod_{i=1}^l \eta^{ii*}. \quad (3.21)$$

It suffices to show that η^{ii*} are nonzero constants for $i = 1, \dots, l$.

For a fixed $1 \leq i < k$, with the use of (3.9) we obtain

$$\begin{aligned} \eta^{ii*} &= \eta^{i(k-i)}(y) = \mathcal{L}_e g^{i(k-i)}(y) \\ &= e^{2\pi i(d_{k,k}x_{l+1} + d_{k,k+1}x_{l+2})} \mathcal{L}_e (c_{i(k-i)} y_i(\mathbf{x}) y_{k-i}(\mathbf{x}) + \beta_{i(k-i)}(\mathbf{x})) \\ &= e^{2\pi i(d_{k,k}x_{l+1} + d_{k,k+1}x_{l+2})} \mathcal{L}_e \beta_{i(k-i)}(\mathbf{x}) \\ &= \frac{\partial}{\partial y_k(\mathbf{x})} \beta_{i(k-i)}(\mathbf{x}) + e^{2\pi i(\frac{k}{l+1}x_{l+1} - \frac{l-k}{l+1}x_{l+2})} \frac{\partial}{\partial y_{k+1}(\mathbf{x})} \beta_{i(k-i)}(\mathbf{x}). \end{aligned}$$

Since η^{ii*} is a constant, we thus have

$$\frac{1}{\varsigma_1} \eta^{ii*} = \frac{\partial}{\partial y_k(\mathbf{x})} \beta_{i(k-i)}(\mathbf{x}) = \frac{1}{4\pi^2} \frac{\partial}{\partial y_k(\mathbf{x})} \sum_{p,q=1}^l \frac{\partial y_i(\mathbf{x})}{\partial x_p} \frac{\partial y_{k-i}(\mathbf{x})}{\partial x_q} (\omega_p, \omega_q)$$

which coincides with the nonzero constant $\eta^{i(k-i)}$ used in the case $\widetilde{W}^{(k)}(A_l)$ (e.g., please see the Corollary 2.3 in [12]).

Similarly, for a fixed $k+1 < i \leq l$, we have

$$\begin{aligned} \frac{1}{\varsigma_2} \eta^{ii*} &= \eta^{i(l+k+2-i)}(y) = \mathcal{L}_e g^{i(l+k+2-i)}(y) \\ &= e^{2\pi i(d_{k+1,k}x_{l+1} + d_{k+1,k+1}x_{l+2})} \mathcal{L}_e (c_{i(l+k+2-i)} y_i(\mathbf{x}) y_{k-i}(\mathbf{x}) + \beta_{i(k-i)}(\mathbf{x})) \\ &= e^{2\pi i(d_{k+1,k}x_{l+1} + d_{k+1,k+1}x_{l+2})} \mathcal{L}_e \beta_{i(l+k+2-i)}(\mathbf{x}) \\ &= e^{2\pi i(\frac{l-k}{l+1}x_{l+2} - \frac{k}{l+1}x_{l+1})} \frac{\partial}{\partial y_k(\mathbf{x})} \beta_{i(k-i)}(\mathbf{x}) + \frac{\partial}{\partial y_{k+1}(\mathbf{x})} \beta_{i(l+k+2-i)}(\mathbf{x}) \\ &= \frac{\partial}{\partial y_{k+1}(\mathbf{x})} \beta_{i(l+k+2-i)}(\mathbf{x}) = \frac{1}{4\pi^2} \frac{\partial}{\partial y_{k+1}(\mathbf{x})} \sum_{p,q=1}^l \frac{\partial y_i(\mathbf{x})}{\partial x_p} \frac{\partial y_{l+k+2-i}(\mathbf{x})}{\partial x_q} (\omega_p, \omega_q) \end{aligned}$$

which is exactly the nonzero constant $\eta^{i(l+k+2-i)}$ used in the case $\widetilde{W}^{(k+1)}(A_l)$ ([12]).

Observe that $\eta^{kk*} = \varsigma_1$ and $\eta^{(k+1)(k+1)*} = \varsigma_2$ in Lemma 3.7, we thus complete the proof of this proposition. \square

According to Lemma D.1 in [10] (or see Lemma 3.3 in [19]) and using Lemma 3.7, Proposition 3.8 and Lemma 3.6, we have

Theorem 3.9. $g^{ij}(y)$ and $\eta^{ij}(y)$ form a flat pencil of metrics, i.e., the metric

$$g^{ij}(y) + \lambda \eta^{ij}(y)$$

is flat for arbitrary λ and the Levi-Civita connection for this metric has the form $\Gamma_m^{ij}(y) + \lambda \gamma_m^{ij}(y)$. Here $\gamma_m^{ij}(y) := \mathcal{L}_e \Gamma_m^{ij}(y)$ are the contravariant components of the Levi-Civita connection for the metric $(\eta^{ij}(y))$.

Without loss of generality, in what follows we take $\varsigma_1 = \varsigma_2 = 1$ unless otherwise stated.

4. FROBENIUS MANIFOLD STRUCTURES ON THE ORBIT SPACE \mathcal{M}

In this section we want to describe Frobenius manifold structures on the orbit space \mathcal{M} of $\widetilde{W}^{(k,k+1)}(A_l)$ for $1 \leq k < l$.

4.1. The change of coordinates. In order to do this, we firstly make the change of coordinates

$$\begin{aligned} z^j &= y^j, & j &= 1, \dots, k-1, k+2, \dots, l, \\ z^k &= y^k, & z^{k+1} &= y^{k+1} - y^k, \\ z^{l+1} &= y^{l+2}, & z^{l+2} &= y^{l+1} + y^{l+2}, \end{aligned} \quad (4.1)$$

such that

$$e = \frac{\partial}{\partial y^k} + \frac{\partial}{\partial y^{k+1}} = \frac{\partial}{\partial z^k}. \quad (4.2)$$

Let us denote

$$g^{ij}(z) = \sum_{a,b=1}^{l+2} \frac{\partial z^i}{\partial y^a} \frac{\partial z^j}{\partial y^b} (dy^a, dy^b)^\sim, \quad \eta^{ij}(z) = \sum_{a,b=1}^{l+2} \frac{\partial z^i}{\partial y^a} \frac{\partial z^j}{\partial y^b} \eta^{ab}(y) \quad (4.3)$$

and $\Gamma_m^{ij}(z)$ and $\gamma_m^{ij}(z)$ are the contravariant components of the Levi-Civita connection for the metric $g^{ij}(z)$ and $\eta^{ij}(z)$. Under the simple change of coordinates (4.1), it is easy to know that

- (1) $g^{ij}(z)$ and $\Gamma_m^{ij}(z)$ are weighted homogeneous polynomials in $z^1, \dots, z^{l+2}, e^{z^{l+1}}, e^{z^{l+2}-z^{l+1}}$ of the degrees $\deg g^{ij}(z) = \deg z^i + \deg z^j$ and

$$\deg \Gamma_m^{ij}(z) = \deg z^i + \deg z^j - \deg z^m, \quad (4.4)$$

where $\deg z^i = d_i$. Moreover, $g^{ij}(z)$ and $\Gamma_m^{ij}(z)$ are at most linear in z^k and

$$\begin{aligned} g^{k,l+1}(z) &= 0, & g^{k+1,l+1}(z) &= z^{k+1} + z^k, \\ g^{k,l+2}(z) &= z^k, & g^{k+1,l+2}(z) &= z^{k+1}, \\ g^{l+1,l+1}(z) &= \frac{l-k+1}{l-k}, & g^{l+1,l+2}(z) &= \frac{1}{l-k}, \\ g^{l+2,l+2}(z) &= \frac{l}{k(l-k)}. \end{aligned} \quad (4.5)$$

(2) $\eta^{ij}(z) = \frac{\partial g^{ij}(z)}{\partial z^k}$ and $\gamma_m^{ij}(z) = \frac{\partial \Gamma_m^{ij}(z)}{\partial z^k}$. Especially,

$$\eta^{i,l+1}(z) = \delta_i^{k+1}, \quad \eta^{i,l+2}(z) = \delta_i^k, \quad i = 1, \dots, l+2. \quad (4.6)$$

So we rename $k^* = l+2$ and $(k+1)^* = l+1$, and also have

$$\eta^{ii^*}(z) = \eta^{ii^*}, \quad i \neq k, k+1, \quad \eta^{kk^*}(z) = \eta^{(k+1)(k+1)^*}(z) = 1.$$

(3) $g^{ij}(z)$ and $\eta^{ij}(z)$ form a flat pencil of metrics.

(4) $\eta_{ij}(z)$ are weighted homogeneous polynomials in $z^1, \dots, z^l, e^{z^{l+2}-z^{l+1}}, e^{z^{l+1}}$, where $(\eta^{ij}(z))$ is the inverse matrix of $(\eta^{ij}(z))$.

4.2. Flat coordinates of the metric $\eta^{ij}(z)$. In this subsection, we want to describe flat coordinates of the metric $\eta^{ij}(z)$.

Lemma 4.1. *For $\nu, s, t = 0, 1$, we have*

$$\frac{\partial}{\partial z^{k+\nu}} \eta^{k+s, k+t}(z) = 0. \quad (4.7)$$

Proof. Observe that

$$\begin{aligned} \frac{\partial}{\partial z^{k+1}} \eta^{k, k+1}(z) &= \frac{\partial}{\partial z^{k+1}} \frac{\partial}{\partial z^k} \sum_{a,b=1}^{l+2} \frac{\partial z^k}{\partial y^a} \frac{\partial z^{k+1}}{\partial y^b} g^{ab}(y) \\ &= \frac{\partial}{\partial y^{k+1}} \left(\frac{\partial}{\partial y^k} + \frac{\partial}{\partial y^{k+1}} \right) (g^{k,k+1}(y) - g^{kk}(y)) = 0 \end{aligned}$$

which follows from (3.12). The other cases are similar. \square

Lemma 4.2. *For $\nu = 0, 1$, we have*

$$\gamma_j^{i, l+1+\nu}(z) = 0, \quad i, j = 1, \dots, l+2. \quad (4.8)$$

Proof. With the use of (4.4) and $\gamma_j^{i, l+1+\nu}(z) = \frac{\partial}{\partial z^k} \Gamma_j^{i, l+1+\nu}(z)$, we obtain

$$\deg \gamma_j^{i, l+1+\nu}(z) = d_i - d_j - d_k < 0, \quad i.e., \quad \gamma_j^{i, l+1+\nu}(z) = 0 \quad (4.9)$$

except the cases $\gamma_{l+1+\mu}^{k+\sigma, l+1+\nu}(z)$ for $\nu, \mu, \sigma = 0, 1$. So it suffices to show that

$$\gamma_{l+1+\mu}^{k+\sigma, l+1+\nu}(z) = 0, \quad \nu, \mu, \sigma = 0, 1. \quad (4.10)$$

Since $\gamma_m^{ij}(z)$ are the contravariant components of the Levi-Civita connection for the metric $g^{ij}(z)$, then

$$2\eta^{sm}(z)\gamma_m^{ij}(z) = \eta^{im}(z)\frac{\partial\eta^{js}(z)}{\partial z^m} + \eta^{sm}(z)\frac{\partial\eta^{ji}(z)}{\partial z^m} - \eta^{jm}(z)\frac{\partial\eta^{is}(z)}{\partial z^m}. \quad (4.11)$$

By choosing $i = k + \sigma$ and $j = l + 1 + \nu$ in (4.11), it follows from (4.9) that

$$\begin{aligned} 2\eta^{s, l+1}(z)\gamma_{l+1}^{k+\sigma, l+1+\nu}(z) + 2\eta^{s, l+2}(z)\gamma_{l+2}^{k+\sigma, l+1+\nu}(z) &= \eta^{k+\sigma, m}(z)\frac{\partial\eta^{l+1+\nu, s}(z)}{\partial z^m} \\ + \eta^{sm}(z)\frac{\partial\eta^{l+1+\nu, k+\sigma}(z)}{\partial z^m} - \eta^{l+1+\nu, m}(z)\frac{\partial\eta^{k+\sigma, s}(z)}{\partial z^m}. \end{aligned} \quad (4.12)$$

Taking $s = k$ and $s = k + 1$ respectively in (4.12) and with the help of (4.6) and (4.7), we get the desired identities (4.10). \square

Theorem 4.3. *There exist flat coordinates of the metric $(\eta^{ij}(z))$ in the form*

$$\begin{aligned} t^\alpha &= z^\alpha + h^\alpha(z^1, \dots, \widehat{z^\alpha}, \dots, z^l, e^{z^{l+1}}, e^{z^{l+2}-z^{l+1}}), \\ t^{l+1} &= z^{l+1}, \quad t^{l+2} = z^{l+2}, \quad \alpha = 1, \dots, l, \end{aligned} \quad (4.13)$$

where h^α are weighted homogeneous polynomials in $z^1, \dots, z^{\alpha-1}, z^{\alpha+1}, \dots, z^l, e^{z^{l+1}}, e^{z^{l+2}-z^{l+1}}$ of degree d_α defined in (2.13).

Proof. Local existence of the coordinates t^1, \dots, t^{l+2} follows from flatness of the metric $(\eta^{ij}(z))$. The flat coordinates $t = t(z)$ are to be found from the following system

$$\frac{\partial^2 t}{\partial z^i \partial z^j} + \eta_{is}(z)\gamma_j^{sm}(z)\frac{\partial t}{\partial z^m} = 0, \quad i, j = 1, \dots, l+2. \quad (4.14)$$

The system (4.14) can be written as linear differential equations

$$\frac{\partial w_i}{\partial z^j} - \gamma_{ij}^m(z)w_m = 0, \quad w_i = \frac{\partial t}{\partial z^i}.$$

This is an overdetermined holonomic system. So the space of solutions has dimension $l+2$. Observe that those coefficients in (4.14) are weighted homogeneous polynomials in $z^1, \dots, z^l, e^{z^{l+1}}, e^{z^{l+2}-z^{l+1}}$. From (4.8), it follows that

$$t^{l+1} = z^{l+1}, \quad t^{l+2} = z^{l+2}$$

are two solutions of (4.14). We choose remaining solutions

$$t^\alpha = z^\alpha + h^\alpha(z^1, \dots, \widehat{z^\alpha}, \dots, z^l, e^{z^{l+1}}, e^{z^{l+2}-z^{l+1}}), \quad \alpha = 1, \dots, l$$

in such a way that

$$\frac{\partial t^\alpha}{\partial z^\beta}(0, \dots, 0, 0, 0) = \delta_\beta^\alpha, \quad \alpha, \beta = 1, \dots, l. \quad (4.15)$$

These solutions t^α are power series in $z^1, \dots, z^l, e^{z^{l+1}}, e^{z^{l+2}-z^{l+1}}$. The system (4.14) is invariant with respect to the transformation

$$z^\alpha \mapsto c_0^{d_\alpha} z^\alpha, \quad z^{l+1} \mapsto z^{l+1} + \frac{1}{l-k} \log(c_0), \quad z^{l+2} \mapsto z^{l+2} + \frac{l}{k(l-k)} \log(c_0)$$

for any positive constant c_0 . This yields that t^α are weighted homogeneous in $z^1, \dots, z^l, e^{z^{l+1}}, e^{z^{l+2}-z^{l+1}}$ of the same degree $d_\alpha > 0$. Thus h^α are weighted homogeneous polynomials in $z^1, \dots, z^{\alpha-1}, z^{\alpha+1}, \dots, z^l, e^{z^{l+1}}, e^{z^{l+2}-z^{l+1}}$ of degree d_α . \square

Corollary 4.4. *In the flat coordinates t^1, \dots, t^{l+2} , the entries of the metric $(\eta^{ij}(t))$ have the form*

$$\eta^{ij} := \eta^{ij}(t) = \begin{cases} \eta^{ii^*}(y), & j = i^*, \\ 0, & j \neq i^*, \end{cases}$$

for $i, j = 1, \dots, l+2$. Especially,

$$\eta^{i,l+1}(t) = \delta_i^{k+1}, \quad \eta^{i,l+2}(t) = \delta_i^k, \quad i = 1, \dots, l+2.$$

The entries of the matrix $(g^{ij}(t))$ and the Christoffel symbols $\Gamma_m^{ij}(t)$ are weighted homogeneous polynomials of $t^1, \dots, t^l, e^{t^{l+1}}, e^{t^{l+2}-t^{l+1}}$ of degrees $d_i + d_j$ and $d_i + d_j - d_m$ respectively. In particular,

$$\begin{aligned} g^{\alpha,l+2}(t) &= d_\alpha t^\alpha, \quad 1 \leq \alpha \leq l, & g^{l+1,l+1}(t) &= \frac{l-k+1}{l-k}, \\ g^{l+1,l+2}(t) &= \frac{1}{l-k}, & g^{l+2,l+2}(t) &= \frac{l}{k(l-k)}, \end{aligned} \quad (4.16)$$

and

$$\Gamma_j^{l+2,i}(t) = d_j \delta_j^i, \quad 1 \leq i, j \leq l+2 \quad (4.17)$$

and

$$g^{k+1,l+1}(t) = t^k + t^{k+1} + g_0(t) \quad (4.18)$$

for certain weighted homogenous polynomial $g_0(t)$ in $t^1, \dots, t^{k-1}, t^{k+2}, \dots, t^l, e^{t^{l+1}}, e^{t^{l+2}-t^{l+1}}$ of degree 1.

Proof. In the flat coordinates t^1, \dots, t^{l+2} , using (4.15) and Theorem 4.3 we have

$$\frac{\partial}{\partial t^k} = \frac{\partial}{\partial z^k}, \quad \eta^{ij}(t) = \frac{\partial g^{ij}(t)}{\partial t^k}.$$

Thus the first statement of this corollary follows from the fact that the linear part of t^α is z^α .

By definition, we easily obtain (4.16). The identity (4.18) follows from (4.5) and (4.13). It remains to prove (4.17). Notice that $t^{l+2} = 2\pi i(x_{l+1} + x_{l+2})$ and

$$\begin{aligned}
\sum_{j=1}^{l+2} \Gamma_j^{l+2,i}(t) dt^j &= \sum_{p,q,r=1}^{l+2} \frac{\partial t^{l+2}}{\partial x_p} \frac{\partial^2 t^i}{\partial x_q \partial x_r} (dx_p, dx_q)^\sim dx_r \\
&= \sum_{r=1}^{l+2} \sum_{p,q=l+1}^{l+2} \frac{\partial t^{l+2}}{\partial x_p} \frac{\partial^2 t^i}{\partial x_q \partial x_r} (dx_p, dx_q)^\sim dx_r \\
&= -4\pi^2 \sum_{p=l+1}^{l+2} [d_{i,k}(dx_p, dx_{l+1})^\sim + d_{i,k+1}(dx_p, dx_{l+2})^\sim] dt^i \\
&= \left(\frac{d_{i,k}}{k} + \frac{d_{i,k+1}}{l-k} \right) dt^i = d_i dt^i.
\end{aligned}$$

This completes the proof of the corollary. \square

4.3. Frobenius manifold structures on the orbit space \mathcal{M} . Now we are ready to describe the Frobenius manifold structures on the orbit space of the extended affine Weyl group $\widetilde{W}^{(k,k+1)}(A_l)$.

Theorem 4.5. *For any fixed integer $1 \leq k < l$, there exists a unique Frobenius manifold structure of charge $d = 1$ on the orbit space \mathcal{M} of $\widetilde{W}^{(k,k+1)}(A_l)$ such that the potential $F(t) = \widehat{F}(t) + \frac{1}{2}(t^{k+1})^2 \log(t^{k+1})$, where $\widehat{F}(t)$ is a weighted homogeneous polynomial in $t^1, t^2, \dots, t^{l+2}, e^{t^{l+1}}, e^{t^{l+2}-t^{l+1}}$, satisfying*

- (1) *the unity vector field e coincides with $\frac{\partial}{\partial t^k}$;*
- (2) *the Euler vector field has the form*

$$E = \sum_{\alpha=1}^l d_\alpha t^\alpha \frac{\partial}{\partial t^\alpha} + \frac{1}{l-k} \frac{\partial}{\partial t^{l+1}} + \frac{l}{k(l-k)} \frac{\partial}{\partial t^{l+2}}, \quad (4.19)$$

where d_1, \dots, d_l are defined in (2.13);

- (3) *the invariant flat metric and the intersection form of the Frobenius manifold structure coincide respectively with the metrics η^{ij} and $g^{ij}(t)$ on \mathcal{M} .*

Proof. The idea of the proof is similar to that of [12], i.e., using the theory of flat pencils of metrics ([10]).

Let $\Gamma_\gamma^{\alpha\beta}(t)$ be the coefficients of the Levi-Civita connection for the metric $(,)^\sim$ in the coordinates t^1, \dots, t^{l+2} . According to Proposition D.1 of [10] one can represent these functions as

$$\Gamma_\gamma^{\alpha\beta}(t) = \eta^{\alpha\epsilon} \partial_\epsilon \partial_\gamma f^\beta(t) \quad (4.20)$$

for some functions $f^\beta(t)$. From the weighted homogeneity of $\Gamma_\gamma^{\alpha\beta}(t)$ and Corollary 4.4, one has

$$\partial_\alpha \partial_\gamma (\mathcal{L}_E f^\beta - (1 + d_\beta) f^\beta) = 0$$

for any α, β . So

$$\mathcal{L}_E f^\beta(t) = (d_\beta + 1) f^\beta(t) + A_\sigma^\beta t^\sigma + B^\beta \quad (4.21)$$

for some constants A_σ^β, B^β . Doing a transformation

$$f^\beta(t) \mapsto \tilde{f}^\beta(t) = f^\beta(t) + R_\lambda^\beta t^\lambda + Q^\beta$$

all the coefficients A_σ^β, B^β in (4.21) can be killed except $A_{k+\nu}^{l+1+\tau}$, for $\nu, \tau = 0, 1$. Indeed, after the transformation,

$$\begin{aligned} \mathcal{L}_E \tilde{f}^\beta(t) &= (d_\beta + 1) \tilde{f}^\beta(t) + \sum_{\gamma=1}^l [R_\gamma^\beta (d_\gamma - 1 - d_\beta) + A_\gamma^\beta] t^\gamma \\ &\quad + [A_{l+1}^\beta - (1 + d_\beta) R_{l+1}^\beta] t^{l+1} + [A_{l+2}^\beta - (1 + d_\beta) R_{l+2}^\beta] t^{l+2} \\ &\quad + \frac{1}{l-k} R_{l+1}^\beta + \frac{l}{k(l-k)} R_{l+2}^\beta + B^\beta - (1 + d_\beta) Q^\beta \end{aligned}$$

The function $\tilde{f}^\beta(t)$ does still satisfy (4.20). Choosing

$$\begin{aligned} R_{l+1}^\beta &= \frac{1}{1 + d_\beta} A_{l+1}^\beta, \quad R_{l+2}^\beta = \frac{1}{1 + d_\beta} A_{l+2}^\beta, \\ Q^\beta &= \frac{1}{1 + d_\beta} \left[B^\beta + \frac{1}{l-k} R_{l+1}^\beta + \frac{l}{k(l-k)} R_{l+2}^\beta \right], \end{aligned}$$

one kills the constant term in the r.h.s. of (4.21) and the term linear in t^{l+1} and t^{l+2} . In order to kill other linear terms, putting

$$R_\gamma^\beta = \frac{1}{d_\beta + d_{\gamma^*}} A_\gamma^\beta,$$

where γ^* is the index dual to γ in the sense of duality defined in the above subsection. We can do this unless $d_\beta = d_{\gamma^*} = 0$. The last equation holds only for

$$\beta = l + 1 + \tau, \quad \gamma = k + \nu, \quad \nu, \tau = 0, 1.$$

So all linear terms can be killed except $A_{k+\nu}^{l+1+\tau}$ for $\nu, \tau = 0, 1$ in (4.21). Thus for $\beta \neq l + 1, l + 2$ the polynomials $f^\beta(t)$ can be assumed to be homogeneous of the degree $d_\beta + 1$.

Next we want to show that for $1 \leq \beta \leq l$ the functions $f^\beta(t)$ are polynomials in $t^1, \dots, t^l, e^{t^{l+1}}, e^{t^{l+2}-t^{l+1}}$. We already know that this is true for the Christoffel

coefficients $\Gamma_\gamma^{\alpha\beta}(t)$. Denoting

$$\eta_{\alpha\epsilon}\Gamma_{l+i}^{\epsilon\beta}(t) = \sum_{m=0}^{M_i} \sum_{n=0}^{N_i} C_{\alpha,m,n}^{\beta,i} e^{m t^{l+1} + n(t^{l+2} - t^{l+1})} \equiv \partial_\alpha \partial_{l+i} f^\beta(t), \quad i = 1, 2,$$

where the coefficients $C_{\alpha,m,n}^{\beta,i}$ are polynomials in t^1, \dots, t^l and M_i, N_i are certain positive integers. From the compatibility condition

$$\partial_{l+i} (\partial_\alpha \partial_{l+j} f^\beta(t)) = \partial_\alpha (\partial_{l+i} \partial_{l+j} f^\beta(t)), \quad i, j = 1, 2,$$

one obtains

$$\partial_\alpha C_{l+i,0,0}^{\beta,j} = 0, \quad \alpha = 1, \dots, l, \quad i, j = 1, 2.$$

So $C_{l+i,0,0}^{\beta,j}$ are constants. But $\partial_{l+i} \partial_{l+j} f^\beta(t)$ must be a weighted homogeneous polynomial in $t^1, \dots, t^l, e^{t^{l+1}}, e^{t^{l+2} - t^{l+1}}$ of the positive degree $1 + d_\beta$. Thus

$$C_{l+i,0,0}^{\beta,j} = 0, \quad i, j = 1, 2.$$

and

$$\partial_\alpha \partial_{l+i} f^\beta(t) = \sum_{m=1}^{M_i} \sum_{n=1}^{N_i} C_{\alpha,m,n}^{\beta,i} e^{m t^{l+1} + n(t^{l+2} - t^{l+1})}, \quad i = 1, 2. \quad (4.22)$$

From the compatibility condition

$$\partial_{l+i} (\partial_{l+j} \partial_{l+j} f^\beta(t)) = \partial_{l+j} (\partial_{l+i} \partial_{l+j} f^\beta(t)) = \partial_{l+j} (\partial_{l+j} \partial_{l+i} f^\beta(t)),$$

for $i, j = 1, 2$, one gets $M := M_1 = M_2$ and $N := N_1 = N_2$ and

$$C_{l+1,m,n}^{\beta,1} = \frac{m-n}{m} C_{l+2,m,n}^{\beta,1} = \frac{m-n}{m} C_{l+1,m,n}^{\beta,2} = \frac{(m-n)^2}{m^2} C_{l+2,m,n}^{\beta,2}. \quad (4.23)$$

It follows from (4.22) and (4.23) that

$$f^\beta(t) = \sum_{m=1}^M \sum_{n=1}^N \frac{1}{m^2} C_{l+2,m,n}^{\beta,2} e^{m t^{l+1} + n(t^{l+2} - t^{l+1})} + t^{l+1} D_1^\beta + t^{l+2} D_2^\beta + D_0^\beta$$

for some new polynomials $D_i^\beta = D_i^\beta(t^1, \dots, t^l)$, $i = 0, 1, 2$. Since $\partial_\alpha \partial_\gamma f^\beta(t)$ must not contain terms linear in t^{l+1} and t^{l+2} , these polynomials D_i^β are at most linear in t^1, \dots, t^l . Using the homogeneity of $f^\beta(t)$, so $D_i^\beta = 0$, $i = 1, 2$ and

$$f^\beta(t) = \sum_{m=1}^M \sum_{n=1}^N \frac{1}{m^2} C_{l+2,m,n}^{\beta,2} e^{m t^{l+1} + n(t^{l+2} - t^{l+1})} + D_0^\beta, \quad \beta = 1, \dots, l.$$

The coefficients $\Gamma_\gamma^{\alpha\beta}(t)$ must also satisfy the conditions

$$g^{\alpha\sigma} \Gamma_\sigma^{\beta\gamma} = g^{\beta\sigma} \Gamma_\sigma^{\alpha\gamma}. \quad (4.24)$$

For $\alpha = l + 2$, it follows from (4.20), (4.24) and (4.17) that

$$\mathcal{L}_E(\eta^{\beta\epsilon} \partial_\epsilon f^\gamma) = d_\gamma g^{\beta\gamma}. \quad (4.25)$$

Notice that the right side has no summation w.r.t the index γ . Because of $\deg f^\gamma = d_\gamma + 1$, then

$$\deg(\eta^{\beta\epsilon}\partial_\epsilon f^\gamma) = d_\beta + d_\gamma, \quad \gamma \neq l+1, l+2.$$

Thus,

$$(d_\gamma + d_\beta)\eta^{\beta\epsilon}\partial_\epsilon f^\gamma = d_\gamma g^{\beta\gamma}, \quad \gamma \neq l+1, l+2. \quad (4.26)$$

Putting

$$F^\gamma = \frac{1}{d_\gamma} f^\gamma, \quad \gamma \neq l+1, l+2$$

and using (4.26), one has

$$\eta^{\beta\epsilon}\partial_\epsilon F^\gamma = \eta^{\gamma\epsilon}\partial_\epsilon F^\beta, \quad 1 \leq \gamma, \beta \leq l. \quad (4.27)$$

From (4.27) it follows that a function $F = F(t)$ exists such that

$$F^\gamma = \eta^{\gamma\epsilon}\partial_\epsilon F, \quad 1 \leq \gamma \leq l. \quad (4.28)$$

The dependence of F on t^k is not determined from (4.28). However, putting $\beta = l+2$ in (4.26), one obtains

$$\partial_k F^\gamma = t^\gamma, \quad 1 \leq \gamma \leq l. \quad (4.29)$$

By using (4.28) and (4.29), one gets

$$\begin{aligned} \partial_{l+2}(\partial_k F) &= t^k, \quad \partial_{l+1}(\partial_k F) = t^{k+1} = \sum_{\alpha=1}^l \eta_{l+1,\alpha} t^\alpha, \\ \partial_\gamma(\partial_k F) &= \sum_{\alpha=1}^l \eta_{\gamma\alpha} t^\alpha, \quad \gamma \neq k, k+1, l+1, l+2. \end{aligned}$$

Notice that $(l+1)^* = k+1$, hence

$$\partial_k F = t^k t^{l+2} + \frac{1}{2} \sum_{\alpha, \beta \neq k, l+2} \eta_{\alpha\beta} t^\alpha t^\beta + \mu(t^k)$$

for some function $\mu(t^k)$. Shifting $F(t) \mapsto F(t) + \int \mu(t^k) dt^k$ one can kill this function, and the equations in (4.28) still hold true due to $\eta^{ik} = \delta_{i,l+2}$. So one has the representation

$$F(t) = \frac{1}{2} (t^k)^2 t^{l+2} + \frac{1}{2} t^k \sum_{\alpha, \beta \neq k, l+2} \eta_{\alpha\beta} t^\alpha t^\beta + G(t) \quad (4.30)$$

with some $G(t)$ independent on t^k .

From the definition (4.28) of F and the weighted homogeneity of f^γ , $\gamma \neq l+1, l+2$, it follows that

$$\mathcal{L}_E F(t) = 2F(t) + \rho(t^k, t^{k+1}) \quad (4.31)$$

for certain unknown function $\rho(t^k, t^{k+1})$. By using (4.30) and the duality condition of the degrees, one has

$$\begin{aligned} \mathcal{L}_E F(t) &= (t^k)^2 t^{l+2} + t^k \sum_{\alpha, \beta \neq k, l+2} \eta_{\alpha\beta} t^\alpha t^\beta \\ &\quad + \mathcal{L}_E G(t) + \frac{l}{2k(l-k)} (t^k)^2 + \frac{1}{l-k} t^k t^{k+1}. \end{aligned} \quad (4.32)$$

From (4.31) and (4.32), one gets

$$\mathcal{L}_E G(t) = 2G(t) + \rho(t^k, t^{k+1}) - \frac{l}{2k(l-k)} (t^k)^2 - \frac{1}{l-k} t^k t^{k+1}.$$

But $\mathcal{L}_E G(t)$ does not depend on t^k , thus there exists an unknown function $\varphi(t^{k+1})$ with $\deg \varphi(t^{k+1}) = 2$, which does not depend on t^k , such that

$$\varphi(t^{k+1}) = s\rho(t^k, t^{k+1}) - \frac{l}{2k(l-k)} (t^k)^2 - \frac{1}{l-k} t^k t^{k+1}$$

and

$$\mathcal{L}_E G(t) = 2G(t) + \varphi(t^{k+1}) + c_0,$$

for a constant c_0 . Killing the constant by a shift, it follows that

$$\mathcal{L}_E G(t) = 2G(t) + \varphi(t^{k+1}), \quad (4.33)$$

and $G(t) = G(t^1, \dots, t^{k-1}, t^{k+1}, \dots, t^{l+1}, e^{t^{l+1}}, e^{t^{l+2}-t^{l+1}})$ is a weighted homogeneous function of the degree 2. The above conditions determine this function uniquely. By integrating polynomials, it yields that G is a polynomial in $t^1, \dots, t^{k-1}, t^{k+2}, \dots, t^{l+1}, e^{t^{l+1}}, e^{t^{l+2}-t^{l+1}}$. But because of the existence of $\varphi(t^{k+1})$, G may have some terms which are not polynomials in t^{k+1} .

Substituting $F(t)$ into (4.25), one gets

$$g^{\alpha\beta} = \mathcal{L}_E F^{\alpha\beta}, \quad F^{\alpha\beta} = \eta^{\alpha\epsilon} \eta^{\beta\delta} \partial_\epsilon \partial_\delta F. \quad (4.34)$$

For $\alpha = l+1, \beta = l+2$, the equation (4.34) reads

$$\mathcal{L}_E \frac{\partial^2 F(t)}{\partial t^k \partial t^{k+1}} = \frac{1}{l-k}. \quad (4.35)$$

For $\alpha = l+2, \beta = l+2$ the equation, (4.34) reads

$$\mathcal{L}_E \frac{\partial^2 F(t)}{\partial t^k \partial t^k} = \frac{l}{k(l-k)}. \quad (4.36)$$

For $\alpha = \beta = l+1$, the equation (4.34) reads

$$\mathcal{L}_E \frac{\partial^2 F(t)}{\partial t^{k+1} \partial t^{k+1}} = \frac{l-k+1}{l-k}. \quad (4.37)$$

By using (4.32) and (4.33), we get

$$\mathcal{L}_E \frac{\partial^2 G(t)}{\partial t^{k+1} \partial t^{k+1}} = \frac{l-k+1}{l-k}$$

and

$$\partial_{k+1}^2 \varphi(t^{k+1}) = \partial_{k+1}^2 \mathcal{L}_E G(t) - 2\partial_{k+1}^2 G(t) = \frac{l-k+1}{l-k}. \quad (4.38)$$

Notice that $\deg \varphi = 2$ and integrating (4.38) to obtain

$$\varphi(t^{k+1}) = \frac{l-k+1}{2(l-k)} (t^{k+1})^2. \quad (4.39)$$

By using (4.28) and (4.30), one can write

$$G(t) = H_0(t) + \psi(t^{k+1}) + a_0 t^{l+1} (t^{k+1})^2, \quad \deg \psi(t^{k+1}) = 2. \quad (4.40)$$

Here a_0 is a constant and $H_0(t)$ is a weighted homogeneous polynomial in $t^1, \dots, t^{k-1}, t^{k+1}, \dots, t^{l+1}, e^{t^{l+1}}, e^{t^{l+2}-t^{l+1}}$ of degree 2 with $\mathcal{L}_E H_0(t) = 2H_0(t)$. In particular, $H_0(t)$ is at most linear in t^{k+1} . By substituting (4.39) and (4.40) into (4.33), one has

$$t^{k+1} \frac{\partial \psi(t^{k+1})}{\partial t^{k+1}} = 2\psi(t^{k+1}) + \left(\frac{k+1}{2k} - \frac{a_0}{k} \right) (t^{k+1})^2. \quad (4.41)$$

The solutions of (4.41) are

$$\psi(t^{k+1}) = c_1 (t^{k+1})^2 + \left(\frac{l-k+1}{2(l-k)} - \frac{a_0}{l-k} \right) (t^{k+1})^2 \log(t^{k+1}), \quad (4.42)$$

where c_1 is an integral constant.

For $\alpha = k+1, \beta = l+1$, the equation (4.34) reads

$$g^{k+1, l+1}(t) = \mathcal{L}_E \frac{\partial^2 F(t)}{\partial t^{k+1} \partial t^{l+1}} = t^k + 2a_0 t^{k+1} + \mathcal{L}_E \frac{\partial^2 H_0(t)}{\partial t^{k+1} \partial t^{l+1}}. \quad (4.43)$$

By comparing (4.18) with (4.43), one gets

$$a_0 = \frac{1}{2}, \quad \mathcal{L}_E \frac{\partial^2 H_0(t)}{\partial t^{k+1} \partial t^{l+1}} = g_0(t).$$

Thus,

$$G(t) = H(t) + \frac{1}{2} (t^{k+1})^2 \log(t^{k+1}),$$

where $H(t) = H_0(t) + c_1 (t^{k+1})^2 + \frac{1}{2} (t^{k+1})^2 t^{l+1}$.

In a word, we show that the existence of a unique weighted homogeneous polynomial

$$H(t) := H(t^1, \dots, t^{k-1}, t^{k+1}, \dots, t^{l+1}, e^{t^{l+1}}, e^{t^{l+2}-t^{l+1}})$$

of degree 2 such that the function

$$F = \frac{1}{2} (t^k)^2 t^{l+2} + \frac{1}{2} t^k \sum_{\alpha, \beta \neq k, l+2} \eta_{\alpha\beta} t^\alpha t^\beta + \frac{1}{2} (t^{k+1})^2 \log(t^{k+1}) + H(t) \quad (4.44)$$

satisfies the equations

$$g^{\alpha\beta} = \mathcal{L}_E F^{\alpha\beta}, \quad \Gamma_\gamma^{\alpha\beta} = d_\beta c_\gamma^{\alpha\beta}, \quad \alpha, \beta, \gamma = 1, \dots, l+2, \quad (4.45)$$

where $c_\gamma^{\alpha\beta} = \frac{\partial F^{\alpha\beta}}{\partial t^\gamma}$.

Obviously, the function F satisfies the equations

$$\frac{\partial^3 F}{\partial t^k \partial t^i \partial t^j} = \eta_{ij}, \quad i, j = 1, \dots, l+2 \quad (4.46)$$

and the quasi-homogeneity condition

$$\mathcal{L}_E F = 2F + \frac{l}{2k(l-k)} (t^k)^2 + \frac{1}{l-k} t^k t^{k+1} + \frac{l-k+1}{2(l-k)} (t^{k+1})^2. \quad (4.47)$$

From the properties of a flat pencil of metrics [10], it follows that F also satisfies associativity equations of WDVV

$$c_m^{ij} c_q^{mp} = c_m^{ip} c_q^{mj} \quad (4.48)$$

for any set of fixed indices i, j, p, q . Now the theorem follows from above properties of the function F and the simple identity $\mathcal{L}_E e = -e$. This completes the proof of the theorem. \square

4.4. Examples. We end this section by giving some examples to illustrate the above construction. For the brevity, instead of t^1, \dots, t^{l+2} we will denote the flat coordinates of the metric η^{ij} by t_1, \dots, t_{l+2} , and also denote $\partial_i = \frac{\partial}{\partial t_i}$ in this subsection.

Example 4.6. Let \widetilde{W} be the extended affine Weyl group $\widetilde{W}^{(1,2)}(A_2)$, then

$$d_{1,1} = \frac{2}{3}, \quad d_{1,2} = d_{2,1} = \frac{1}{3}, \quad d_{2,2} = \frac{2}{3}$$

and

$$\begin{aligned} y^1 &= e^{\frac{2\pi i}{3}(2x_3+x_4)} (e^{2\pi i x_1} + e^{2\pi i x_2} + e^{-2\pi i(x_1+x_2)}), \\ y^2 &= e^{\frac{2\pi i}{3}(x_3+2x_4)} (e^{-2\pi i x_1} + e^{-2\pi i x_2} + e^{2\pi i(x_1+x_2)}), \\ y^3 &= 2i\pi x_3, \quad y^4 = 2i\pi x_4. \end{aligned}$$

The metric $(,)^\sim$ has the form

$$((dx_a, dx_b)^\sim) = \frac{1}{4\pi^2} \begin{pmatrix} \frac{2}{3} & \frac{1}{3} & 0 & 0 \\ \frac{1}{3} & \frac{2}{3} & 0 & 0 \\ 0 & 0 & -2 & 1 \\ 0 & 0 & 1 & -2 \end{pmatrix}.$$

We thus have

$$(g^{ij}(y)) = \left(\sum_{a,b=1}^4 \frac{\partial y^i}{\partial x_a} \frac{\partial y^j}{\partial x_b} (dx_a, dx_b)^\sim \right) = \begin{pmatrix} 2y^2e^{y^3} & 3e^{y^3+y^4} & y^1 & 0 \\ 3e^{y^3+y^4} & 2y^1e^{y^4} & 0 & y^2 \\ y^1 & 0 & 2 & -1 \\ 0 & y^2 & -1 & 2 \end{pmatrix}$$

and

$$(\eta^{ij}(y)) = (\mathcal{L}_e g^{ij}(y)) \begin{pmatrix} 2e^{y^3} & 0 & 1 & 0 \\ 0 & 2e^{y^4} & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}, \quad e = \frac{\partial}{\partial y^1} + \frac{\partial}{\partial y^2}.$$

The flat coordinates for the metric $(\eta^{ij}(y))$ are

$$t_1 = y^1 - e^{y^3}, \quad t_2 = -y^1 + y^2 + e^{y^3} - e^{y^4}, \quad t_3 = y^4, \quad t_4 = y^3 + y^4.$$

The potential has the expression

$$F = \frac{1}{2} t_1^2 t_4 + t_1 t_2 t_3 + \frac{1}{2} t_2^2 t_3 + e^{t_4} - t_2 e^{t_3} + t_2 e^{t_4 - t_3} + \frac{1}{2} t_2^2 \log t_2$$

and the unit vector field is

$$e = \partial_1$$

and the Euler vector field is given by

$$E = t_1 \partial_1 + t_2 \partial_2 + \partial_3 + 2\partial_4.$$

Example 4.7. Let \widetilde{W} be the extended affine Weyl group $\widetilde{W}^{(1,2)}(A_3)$, then

$$d_{1,1} = \frac{3}{4}, \quad d_{1,2} = \frac{1}{2}, \quad d_{1,3} = \frac{1}{4}, \quad d_{2,2} = 1, \quad d_{2,3} = \frac{1}{2}, \quad d_{3,3} = \frac{3}{4}$$

and

$$y^1 = e^{\frac{\pi i}{2}(3x_4+2x_5)} \sum_{j=1}^4 \xi_a, \quad y^2 = e^{\pi i(x_4+2x_5)} \sum_{1 \leq a < b \leq 4} \xi_a \xi_b, \\ y^3 = e^{\frac{\pi i}{2}(x_4+2x_5)} \sum_{1 \leq a < b < c \leq 4} \xi_a \xi_b \xi_c, \quad y^4 = 2i\pi x_4, \quad y^5 = 2i\pi x_5,$$

where $\xi_j = e^{2\pi i(x_j - x_{j-1})}$ for $j = 1, 2, 3$ and $x_0 = 0$, $\xi_4 = e^{-2\pi i x_3}$. The metric $(,)^\sim$ has the form

$$((dx_i, dx_j)^\sim) = \frac{1}{4\pi^2} \begin{pmatrix} \frac{3}{4} & \frac{1}{2} & \frac{1}{4} & 0 & 0 \\ \frac{1}{2} & 1 & \frac{1}{2} & 0 & 0 \\ \frac{1}{4} & \frac{1}{2} & \frac{3}{4} & 0 & 0 \\ 0 & 0 & 0 & -2 & \frac{1}{2} \\ 0 & 0 & 0 & \frac{1}{2} & -\frac{3}{2} \end{pmatrix}.$$

We thus have

$$(g^{ij}(y)) = \begin{pmatrix} 2y^2e^{y^4} & 3y^3e^{y^4+y^5} & 4e^{y^4+y^5} & y^1 & 0 \\ 3y^3e^{y^4+y^5} & 2y^1y^3e^{y^5} + 4e^{2y^5+y^4} & 3y^1e^{y^5} & 0 & y^2 \\ 4e^{y^4+y^5} & 3y^1e^{y^5} & -\frac{1}{2}(y^3)^2 + 2y^2 & 0 & \frac{1}{2}y^3 \\ y^1 & 0 & 0 & 2 & -1 \\ 0 & y^2 & \frac{1}{2}y^3 & -1 & \frac{3}{2} \end{pmatrix}$$

and

$$(\eta^{ij}(y)) = (\mathcal{L}_e g^{ij}(y)) = \begin{pmatrix} 2e^{y^4} & 0 & 0 & 1 & 0 \\ 0 & 2y^3e^{y^5} & 3e^{y^5} & 0 & 1 \\ 0 & 3e^{y^5} & 2 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \end{pmatrix}, \quad e = \frac{\partial}{\partial y^1} + \frac{\partial}{\partial y^2}.$$

To write down the flat coordinates, we first introduce the following variables

$$z^1 = y^1, \quad z^2 = y^2 - y^1, \quad z^3 = y^3, \quad z^4 = y^5, \quad z^5 = y^4 + y^5,$$

and obtain

$$(\eta^{ij}(z)) = \begin{pmatrix} 2e^{z^5-z^4} & -2e^{z^5-z^4} & 0 & 0 & 1 \\ -2e^{z^5-z^4} & 2e^{z^5-z^4} + 2z^3e^{z^4} & 3e^{z^4} & 1 & 0 \\ 0 & 3e^{z^4} & 2 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \end{pmatrix},$$

then the flat coordinates for the metric $(\eta^{ij}(z))$ are given by

$$\begin{aligned} t_1 &= z^1 - e^{z^5-z^4}, & t_2 &= z^2 - z^3e^{z^4} + e^{z^5-z^4} + e^{2z^4}, \\ t_3 &= z^3 - e^{z^4}, & t_4 &= z^4, & t_5 &= z^5. \end{aligned}$$

The potential has the expression

$$F = \frac{1}{2}t_1^2t_5 + t_1t_2t_4 + \frac{1}{2}t_2^2t_4 + \frac{1}{4}t_3^2t_2 + \frac{1}{4}t_3^2t_1 - \frac{1}{96}t_3^4 \\ + t_3e^{t_5} - t_2t_3e^{t_4} + t_2e^{t_5-t_4} + \frac{1}{2}t_2e^{2t_4} + \frac{1}{2}\mathbf{t}_2^2 \log(\mathbf{t}_2)$$

and the unit vector field is

$$e = \partial_1$$

and the Euler vector field is given by

$$E = t_1\partial_1 + t_2\partial_2 + \frac{1}{2}t_3\partial_3 + \frac{1}{2}\partial_4 + \frac{3}{2}\partial_5.$$

Example 4.8. Let \widetilde{W} be the extended affine Weyl group $\widetilde{W}^{(2,3)}(A_3)$, then

$$d_{1,1} = \frac{3}{4}, \quad d_{1,2} = \frac{1}{2}, \quad d_{1,3} = \frac{1}{4}, \quad d_{2,2} = 1, \quad d_{2,3} = \frac{1}{2}, \quad d_{3,3} = \frac{3}{4}$$

and

$$y^1 = e^{\frac{\pi i}{2}(2x_4+x_5)} \sum_{j=1}^4 \xi_a, \quad y^2 = e^{\pi i(2x_4+x_5)} \sum_{1 \leq a < b \leq 4} \xi_a \xi_b, \\ y^3 = e^{\frac{\pi i}{2}(2x_4+3x_5)} \sum_{1 \leq a < b < c \leq 4} \xi_a \xi_b \xi_c, \quad y^4 = 2i\pi x_4, \quad y^5 = 2i\pi x_5,$$

where $\xi_j = e^{2\pi i(x_j - x_{j-1})}$ for $j = 1, 2, 3$ and $x_0 = 0$, $\xi_4 = e^{-2\pi i x_3}$. The metric $(,)^\sim$ has the form

$$((dx_i, dx_j)^\sim) = \frac{1}{4\pi^2} \begin{pmatrix} \frac{3}{4} & \frac{1}{2} & \frac{1}{4} & 0 & 0 \\ \frac{1}{2} & 1 & \frac{1}{2} & 0 & 0 \\ \frac{1}{4} & \frac{1}{2} & \frac{3}{4} & 0 & 0 \\ 0 & 0 & 0 & -\frac{3}{2} & 1 \\ 0 & 0 & 0 & 1 & -2 \end{pmatrix}.$$

We thus have

$$(g^{ij}(y)) = \begin{pmatrix} -\frac{1}{2}(y^1)^2 + 2y^2 & 3y^3e^{y^4} & 4e^{y^4+y^5} & \frac{1}{2}y^1 & 0 \\ 3y^3e^{y^4} & 2y^1y^3e^{y^4} + 4e^{y^5+2y^4} & 3y^1e^{y^4+y^5} & y^2 & 0 \\ 4e^{y^4+y^5} & 3y^1e^{y^4+y^5} & 2y^2e^{y^5} & 0 & y^3 \\ \frac{1}{2}y^1 & y^2 & 0 & \frac{3}{2} & -1 \\ 0 & 0 & y^3 & -1 & 2 \end{pmatrix}$$

and

$$(\eta^{ij}(y)) = (\mathcal{L}_e g^{ij}(y)) = \begin{pmatrix} 2 & 3e^{y^4} & 0 & 0 & 0 \\ 3e^{y^4} & 2y^1 e^{y^4} & 0 & 1 & 0 \\ 0 & 0 & 2e^{y^5} & 0 & 1 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \end{pmatrix}, \quad e = \frac{\partial}{\partial y^2} + \frac{\partial}{\partial y^3}.$$

To write down the flat coordinates, we first introduce the following variables

$$z^1 = y^1, \quad z^2 = y^2, \quad z^3 = y^3 - y^2, \quad z^4 = y^5, \quad z^5 = y^4 + y^5,$$

and obtain

$$(\eta^{ij}(z)) = \begin{pmatrix} 2 & 3e^{z^5-z^4} & -3e^{z^5-z^4} & 0 & 0 \\ 3e^{z^5-z^4} & 2e^{z^5-z^4} z^1 & -2z^1 e^{z^5-z^4} & 0 & 1 \\ -3e^{z^5-z^4} & -2z^1 e^{z^5-z^4} & 2z^1 e^{z^5-z^4} + 2e^{z^4} & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \end{pmatrix},$$

then the flat coordinates for the metric $(\eta^{ij}(z))$ are given by

$$\begin{aligned} t_1 &= z^1 - e^{z^5-z^4}, & t_2 &= z^2 - z^1 e^{z^5-z^4} + e^{2z^5-2z^4}, \\ t_3 &= z^3 - e^{z^4} + z^1 e^{z^5-z^4} - e^{2(z^5-z^4)}, & t_4 &= z^4, & t_5 &= z^5. \end{aligned}$$

The potential has the expression

$$\begin{aligned} F &= \frac{1}{2} t_2^2 t_5 + t_2 t_3 t_4 + \frac{1}{2} t_3^2 t_4 + \frac{1}{4} t_1^2 t_2 - \frac{1}{96} t_1^4 \\ &+ t_1 e^{t_5} - t_3 e^{t_4} + t_1 t_3 e^{t_5-t_4} - \frac{1}{2} e^{2t_5-2t_4} t_3 + \frac{1}{2} \mathbf{t}_3^2 \log(\mathbf{t}_3) \end{aligned}$$

and the unit vector field is

$$e = \partial_1$$

and the Euler vector field is given by

$$E = \frac{1}{2} t_1 \partial_1 + t_2 \partial_2 + t_3 \partial_3 + \partial_4 + \frac{3}{2} \partial_5.$$

5. THE GROUP $\widetilde{W}^{(k,k+1)}(A_l)$ AND THE HURWITZ SPACE $\mathbb{M}_{k,l-k+1,1}$

In this section we want to show that the space $\mathbb{M}_{k,l-k+1,1}$ as a Frobenius manifold is isomorphic to the orbit space \mathcal{M} of the extended affine Weyl group $\widetilde{W}^{(k,k+1)}(A_l)$, where $\mathcal{M} = \mathcal{M}^{(k,k+1)} \setminus \{\widetilde{y}_{l+1} = 0\} \cup \{\widetilde{y}_{l+2} = 0\}$ for $1 \leq k < l$.

Let $\mathbb{M}_{k,l-k+1,1}$ be the space of a particular class of LG superpotentials consisting of trigonometric-Laurent series of one variable with tri-degree $(k+1, l-k, 1)$, these being functions of the form

$$\lambda(\varphi) = (e^{i\varphi} - a_{l+2})^{-1}(e^{i(k+1)\varphi} + a_1 e^{ik\varphi} + \cdots + a_{l+1} e^{i(k-l)\varphi}), \quad a_{l+1} a_{l+2} \neq 0, \quad (5.1)$$

where $a_j \in \mathbb{C}$ for $j = 1, \dots, l+2$. For brevity, we denote $m = l-k$ in this section. According to [10], the space $\mathbb{M}_{k,m+1,1}$ is a simple Hurwitz space and carries a natural structure of Frobenius manifold. The invariant inner product η and the intersection form g of two vectors ∂', ∂'' tangent to $M_{k,m+1,1}$ at a point $\lambda(\varphi)$ can be defined by the following formulae

$$\widetilde{\eta}(\partial', \partial'') = (-1)^{k+1} \sum_{|\lambda| < \infty} \operatorname{res}_{d\lambda=0} \frac{\partial'(\lambda(\varphi)d\varphi)\partial''(\lambda(\varphi)d\varphi)}{d\lambda(\varphi)}, \quad (5.2)$$

and

$$\widetilde{g}(\partial', \partial'') = - \sum_{|\lambda| < \infty} \operatorname{res}_{d\lambda=0} \frac{\partial'(\log \lambda(\varphi)d\varphi)\partial''(\log \lambda(\varphi)d\varphi)}{d \log \lambda(\varphi)}. \quad (5.3)$$

In these formulae, the derivatives $\partial'(\lambda(\varphi)d\varphi)$ etc. are to be calculated keeping φ fixed. The formulae (5.2) and (5.3) uniquely determine multiplication of tangent vectors on $\mathbb{M}_{k,m+1,1}$ assuming that the Euler vector field E has the form

$$E = \sum_{j=1}^{l+1} \frac{j}{k} a_j \frac{\partial}{\partial a_j} + \frac{1}{k} a_{l+2} \frac{\partial}{\partial a_{l+2}}. \quad (5.4)$$

For any tangent vectors ∂', ∂'' and ∂''' to $M_{k,m+1,1}$, one has

$$c(\partial', \partial'', \partial''') = \widetilde{\eta}(\partial' \cdot \partial'', \partial''') = - \sum_{|\lambda| < \infty} \operatorname{res}_{d\lambda=0} \frac{\partial'(\lambda(\varphi)d\varphi)\partial''(\lambda(\varphi)d\varphi)\partial'''(\lambda(\varphi)d\varphi)}{d\lambda(\varphi)d\varphi}. \quad (5.5)$$

The canonical coordinates u_1, \dots, u_{l+2} for this multiplication are the critical values of $\lambda(\varphi)$ and

$$\partial_{u_\alpha} \cdot \partial_{u_\beta} = \delta_{\alpha\beta} \partial_{u_\alpha}, \quad \text{where} \quad \partial_{u_\alpha} = \frac{\partial}{\partial u_\alpha}. \quad (5.6)$$

We start with factorizing

$$\lambda(\varphi) = (e^{i\varphi} - e^{i\varphi_{l+2}})^{-1} e^{-im\varphi} \prod_{b=1}^{l+1} (e^{i\varphi} - e^{i\varphi_b}). \quad (5.7)$$

Theorem 5.1. *Let $\mathfrak{f} : \mathcal{M} \rightarrow \mathbb{M}_{k,m+1,1}$ be induced by the map*

$$(x_1, \dots, x_{l+2}) \mapsto (\varphi_1, \dots, \varphi_{l+2}) \quad (5.8)$$

with

$$\begin{aligned} \varphi_1 &= 2\pi(\rho + x_1), & \varphi_j &= 2\pi(\rho + x_j - x_{j+1}), \quad j = 2, \dots, l, \\ \varphi_{l+1} &= 2\pi(\rho - x_l), & \varphi_{l+2} &= 2\pi x_{l+1}, \end{aligned} \quad (5.9)$$

where $\rho = \frac{m+1}{l+1}x_{l+1} + \frac{m}{l+1}x_{l+2}$. Then \mathfrak{f} is an m -fold covering map, which is also a local isomorphism between the Frobenius manifolds \mathcal{M} and $\mathbb{M}_{k,m+1,1}$.

Proof. With the use of (5.1) and (5.7), one obtains

$$a_j = (-1)^j \sigma_j(e^{i\varphi_1}, \dots, e^{i\varphi_{l+1}}), \quad a_{l+2} = e^{i\varphi_{l+2}}, \quad j = 1, \dots, l+1.$$

From the formulae (2.2), (2.11) and (5.9), it follows

$$a_j = \begin{cases} (-1)^j y^j, & j = 1, \dots, k, \\ (-1)^j y^j e^{(j-k)y^{l+1} + (j-k-1)y^{l+2}}, & j = k+1, \dots, l, \\ (-1)^{l+1} e^{(m+1)y^{l+1} + my^{l+2}}, & j = l+1, \\ e^{y^{l+1}}, & j = l+2. \end{cases} \quad (5.10)$$

Then the map $\mathfrak{f} : (\tilde{y}_1, \dots, \tilde{y}_{l+2}) \mapsto (a_1, \dots, a_{l+2})$ is given by

$$\begin{aligned} a_j &= (-1)^j \tilde{y}_j, \quad j = 1, \dots, k, \\ a_{k+s} &= (-1)^{k+s} \tilde{y}_{k+s} \tilde{y}_{l+1}^s \tilde{y}_{l+2}^{s-1}, \quad s = 1, \dots, l-k, \\ a_{l+1} &= (-1)^{l+1} \tilde{y}_{l+1}^{m+1} \tilde{y}_{l+2}^m, \quad a_{l+2} = \tilde{y}_{l+1} \end{aligned} \quad (5.11)$$

and the Jacobian of \mathfrak{f} is proportional to $\tilde{y}_{l+1}^{m+1} \tilde{y}_{l+2}^{m-1}$. So \mathfrak{f} is an m -fold covering map.

Now let us proceed to prove that \mathfrak{f} is a local isomorphism between the two Frobenius manifolds. By using (5.11), it is easy to check that the Euler vector fields (5.4) and (4.19) coincide. So it suffices to prove that the intersection form (5.3) coincides with the intersection form of the orbit space, and the metric (5.2) coincides with the metric (3.18).

Let us denote the roots of $\lambda'(\varphi)$ by ψ_γ , $\gamma = 1, \dots, l+2$, then

$$\lambda'(\varphi) = (k+1)i(e^{i\varphi} - e^{i\varphi_{l+2}})^{-2} e^{-im\varphi} \prod_{\gamma=1}^{l+2} (e^{i\varphi} - e^{i\psi_\gamma}). \quad (5.12)$$

We define $u_\alpha = \lambda(\psi_\alpha)$, $\alpha = 1, \dots, l+2$, then

$$\partial_{u_\alpha} \lambda(\varphi)|_{\varphi=\psi_\beta} = \delta_{\alpha\beta}. \quad (5.13)$$

Using (5.7), (5.13) and the Lagrange interpolation formula one obtains

$$\partial_{u_\alpha} \lambda(\varphi) = \frac{ie^{i\psi_\alpha} \lambda'(\varphi)}{(e^{i\varphi} - e^{i\psi_\alpha}) \lambda''(\psi_\alpha)}. \quad (5.14)$$

It follows from (5.7) and (5.14) that

$$\frac{ie^{i\psi_\alpha} \lambda'(\varphi)}{(e^{i\varphi} - e^{i\psi_\alpha}) \lambda''(\psi_\alpha)} = \lambda(\varphi) \frac{i\partial_{u_\alpha} \varphi_{l+2} e^{i\varphi_{l+2}}}{e^{i\varphi} - e^{i\varphi_{l+2}}} - \sum_{s=1}^{l+1} \frac{i\partial_{u_\alpha} \varphi_s e^{i\varphi_s} \lambda(\varphi)}{e^{i\varphi} - e^{i\varphi_s}}. \quad (5.15)$$

Putting $\varphi = \varphi_b$ in (5.15) for $b = 1, \dots, l+1$, then

$$\partial_{u_\alpha} \varphi_b = -\frac{ie^{i\psi_\alpha}}{(e^{i\varphi_b} - e^{i\psi_\alpha}) \lambda''(\psi_\alpha)}, \quad b = 1, \dots, l+1. \quad (5.16)$$

Let us rewrite (5.15) as

$$\frac{ie^{i\psi_\alpha} \lambda'(\varphi)}{(e^{i\varphi} - e^{i\psi_\alpha}) \lambda''(\psi_\alpha) \lambda(\varphi)^2} = \frac{i\partial_{u_\alpha} \varphi_{l+2} e^{i\varphi_{l+2}}}{(e^{i\varphi} - e^{i\varphi_{l+2}}) \lambda(\varphi)} - \sum_{b=1}^{l+1} \frac{i\partial_{u_\alpha} \varphi_b e^{i\varphi_b}}{(e^{i\varphi} - e^{i\varphi_b}) \lambda(\varphi)}. \quad (5.17)$$

Also, putting $\varphi = \varphi_{l+2}$ in (5.17) one gets

$$\partial_{u_\alpha} \varphi_{l+2} = -\frac{ie^{i\psi_\alpha}}{(e^{i\varphi_{l+2}} - e^{i\psi_\alpha}) \lambda''(\psi_\alpha)}. \quad (5.18)$$

We denote

$$\varpi_1 = x_1, \quad \varpi_j = x_j - x_{j-1}, \quad \varpi_{l+1} = x_{l+1}, \quad \varpi_{l+2} = x_{l+2} \quad (5.19)$$

for $j = 2, \dots, l$. With the help of (5.9), (5.16) and (5.18), one has

$$\begin{aligned} \partial_{u_\alpha} \varpi_\beta &= \frac{e^{i\psi_\alpha}}{2\pi i (e^{i\varphi_\beta} - e^{i\psi_\alpha}) \lambda''(\psi_\alpha)} \\ &\quad - \frac{m+1}{l+1} \partial_{u_\alpha} \varpi_{l+1} - \frac{m}{l+1} \partial_{u_\alpha} \varpi_{l+2}, \\ \partial_{u_\alpha} \varpi_{l+1} &= \frac{e^{i\psi_\alpha}}{2\pi i (e^{i\varphi_{l+1}} - e^{i\psi_\alpha}) \lambda''(\psi_\alpha)}, \\ \partial_{u_\alpha} \varpi_{l+2} &= \sum_{b=1}^{l+1} \frac{e^{i\psi_\alpha}}{2\pi m i (e^{i\varphi_b} - e^{i\psi_\alpha}) \lambda''(\psi_\alpha)} \\ &\quad - \frac{(m+1)e^{i\psi_\alpha}}{2\pi m i (e^{i\varphi_{l+2}} - e^{i\psi_\alpha}) \lambda''(\psi_\alpha)}. \end{aligned} \quad (5.20)$$

From (5.2), (5.3) and (5.13) one gets

$$\tilde{\eta}_{\alpha\beta} = \tilde{\eta}(\partial_{u_\alpha} \partial_{u_\beta}) = (-1)^{k+1} \frac{\delta_{\alpha\beta}}{\lambda''(\psi_\alpha)} \quad (5.21)$$

and

$$\tilde{g}_{\alpha\beta} = \tilde{g}(\partial_{u_\alpha} \partial_{u_\beta}) = -\frac{\delta_{\alpha\beta}}{u_\alpha \lambda''(\psi_\alpha)}. \quad (5.22)$$

It follows from (5.10) that the vector field $e = \frac{\partial}{\partial y^k} + \frac{\partial}{\partial y^{k+1}}$ in the coordinates a_1, \dots, a_{l+2} coincides with $e = (-1)^k (\frac{\partial}{\partial a_k} - a_{l+2} \frac{\partial}{\partial a_{k+1}})$. We shift

$$a_k \mapsto a_k + c, \quad a_{k+1} \mapsto a_{k+1} - ca_{l+2}$$

which produces the corresponding shift

$$u_i \mapsto u_i + c, \quad i = 1, \dots, l+2$$

of the critical values. This shift does not change the critical points ψ_α neither the values of the second derivative $\lambda''(\psi_\alpha)$. So

$$\mathcal{L}_e \tilde{g}^{\alpha\beta} = \mathcal{L}_e (-u_\alpha \lambda''(\psi_\alpha) \delta_{\alpha\beta}) = (-1)^{k+1} \lambda''(\psi_\alpha) \delta_{\alpha\beta} = \tilde{\eta}^{\alpha\beta}.$$

Finally we want to show that the bilinear form (5.3) in the coordinates x_1, \dots, x_{l+2} coincides with the form defined in (3.19) and (3.2). We shall use the following identity

$$\begin{aligned} & \sum_{\alpha=1}^{l+2} \frac{u_\alpha e^{2i\psi_\alpha}}{(e^{i\varphi_a} - e^{i\psi_\alpha})(e^{i\varphi_b} - e^{i\psi_\alpha}) \lambda''(\psi_\alpha)} \\ &= \sum_{\alpha=1}^{l+2} \operatorname{res}_{\varphi=\psi_\alpha} \frac{\lambda(\varphi) u_\alpha e^{2i\varphi}}{(e^{i\varphi} - e^{i\varphi_a})(e^{i\varphi} - e^{i\varphi_b}) \lambda'(\varphi)} \\ &= \begin{cases} \delta_{ab} - \frac{1}{k}, & 1 \leq a, b \leq l+1, \\ -\frac{1}{k}, & 1 \leq a \leq l+1, b = l+2, \\ -1 - \frac{1}{k}, & a = b = l+2, \end{cases} \end{aligned} \quad (5.23)$$

which follows from the explicit form of $\lambda(\varphi)$. With the use of (5.20), (5.22) and (5.23), then

$$\begin{aligned} (d\varpi_{l+1}, d\varpi_{l+1}) &= \sum_{\alpha=1}^{l+2} \frac{1}{\tilde{g}_{\alpha\alpha}(u)} \partial_{u_\alpha} \varpi_{l+1} \partial_{u_\alpha} \varpi_{l+1} \\ &= \frac{1}{4\pi^2} \sum_{\alpha=1}^{l+2} \frac{u_\alpha e^{2i\psi_\alpha}}{(e^{i\varphi_{l+2}} - e^{i\psi_\alpha})^2 \lambda''(\psi_\alpha)} = -\frac{1}{4\pi^2} \frac{k+1}{k} \end{aligned}$$

and

$$\begin{aligned}
(d\varpi_{l+1}, d\varpi_{l+2}) &= \sum_{\alpha=1}^{l+2} \frac{1}{\tilde{g}_{\alpha\alpha}(u)} \partial_{u_\alpha} \varpi_{l+1} \partial_{u_\alpha} \varpi_{l+2} \\
&= \frac{1}{4\pi^2 m} \sum_{\alpha=1}^{l+2} \sum_{b=1}^{l+1} \frac{u_\alpha e^{2i\psi_\alpha}}{(e^{i\varphi_b} - e^{i\psi_\alpha})(e^{i\varphi_{l+2}} - e^{i\psi_\alpha}) \lambda''(\psi_\alpha)} \\
&\quad - \frac{m+1}{4\pi^2 m} \sum_{\alpha=1}^{l+2} \frac{u_\alpha e^{2i\psi_\alpha}}{(e^{i\varphi_{l+2}} - e^{i\psi_\alpha})^2 \lambda''(\psi_\alpha)} \\
&= \frac{1}{4\pi^2 m} \sum_{b=1}^{l+1} \left(-\frac{1}{k}\right) + \frac{(m+1)(k+1)}{4\pi^2 m k} = \frac{1}{4\pi^2}
\end{aligned}$$

and

$$\begin{aligned}
(d\varpi_{l+2}, d\varpi_{l+2}) &= \sum_{\alpha=1}^{l+2} \frac{1}{\tilde{g}_{\alpha\alpha}(u)} \partial_{u_\alpha} \varpi_{l+2} \partial_{u_\alpha} \varpi_{l+2} \\
&= \frac{1}{4\pi^2 m^2} \sum_{a,b=1}^{l+1} \left(\delta_{ab} - \frac{1}{k}\right) - \frac{2(m+1)}{4\pi^2 m^2} \sum_{a=1}^{l+1} \left(-\frac{1}{k}\right) + \frac{(m+1)^2}{4\pi^2 m^2} \left(-1 - \frac{1}{k}\right) \\
&= -\frac{1}{4\pi^2} \frac{m+1}{m}
\end{aligned}$$

and for $b = 1, \dots, l$,

$$(d\varpi_b, d\varpi_{l+1}) = (d\varpi_b, d\varpi_{l+2}) = 0$$

and for $1 \leq s, b \leq l$,

$$(d\varpi_a, d\varpi_b) = \sum_{\alpha=1}^{l+2} \frac{1}{\tilde{g}_{\alpha\alpha}(u)} \partial_{u_\alpha} \varpi_a \partial_{u_\alpha} \varpi_b = \frac{1}{4\pi^2} \left(\delta_{ab} - \frac{1}{l+1}\right).$$

By using (5.19) and the above explicit forms, it is easy to verify that the intersection form (5.3) coincides with (3.1) and (3.2). This completes the proof of the theorem. \square

6. CONCLUSIONS

We have presented a new class of extended affine Weyl groups $\widetilde{W}^{(k,k+1)}(A_l)$ for $1 \leq k < l$. On the orbit spaces $\mathcal{M}^{(k,k+1)}(A_l) \setminus \{\tilde{y}_{l+1} = 0\} \cup \{\tilde{y}_{l+2} = 0\}$, we have shown the existence of Frobenius manifold structures and constructed LG superpotentials for these Frobenius manifold structures. Besides these, there are still some open problems deserved further study.

- Is it possible to obtain an explicit realization of the integrable hierarchies associated with the Frobenius manifolds on the orbit space of $\widetilde{W}^{(k,k+1)}(A_l)$? Perhaps this problem is related to the works in [23] or [24] about rational reductions of the 2D-Toda hierarchy, or in [25, 26] about the finite Toda lattice of CKP type when $k = 1$ and $l = 2$.
- How about the almost dual structure of the resulting Frobenius manifold structures? ([17, 27])
- Whether the resulting Frobenius manifolds could be regarded as Frobenius submanifolds in Strachan's sense ([15]) of certain infinite-dimensional Frobenius manifolds ([20, 21, 22]) or not?

Acknowledgments. The author is grateful to Professor Youjin Zhang for bringing me the attention of this project and helpful discussions. This work is partially supported by NSFC (No.11671371, No.11871446) and Wu Wen-Tsun Key Laboratory of Mathematics, USTC, CAS.

REFERENCES

- [1] J.E.Humphryd, *Introduction to Lie algebras and representation theory*, Graduate texts in mathematics 9, Springer-Verlage, New York, Heidelberg, Berlin, 1972.
- [2] E.Looijenga, *Root systems and elliptic curves*, Inventiones math., 38 (1976) 17–32.
- [3] E.Looijenga, *Invariant theory of generalized root systems*, Inventiones math., 61 (1980) 1–32.
- [4] N.Bourbaki, *Groupes et Algèbres de Lie, Chapitres 4, 5 et 6*, Masson, Paris-New York-Barcelone-Milan-Mexico-Rio de Janeiro, 1981.
- [5] K. Wirthmüller, *Torus embeddings and deformations of simple space curves*, Acta Mathematica 157 (1986) 159–241.
- [6] M.F.Atiyah, *Topological quantum field theories*, Publ. Math. I.H.E.S., 68 (1988) 175.
- [7] E.Witten, *On the structure of the topological phase of two-dimensional gravity*, Nucl.Phys.B 340(1990) 281–332.
- [8] R.Dijkgraaf, E.Verlinde, and H.Verlinde, *Topological strings in $d < 1$* , Nuclear Phys. B 352(1991) 59–86.
- [9] B.Dubrovin, *Integrable systems in topological field theory*, Nucl. Phys. B 379 (1992) 627–689.
- [10] B.Dubrovin, *Geometry of 2D topological field theories*, In:Springer Lecture Notes in Math. 1620(1996) 120–348.
- [11] B.Dubrovin, *Differential geometry of the space of orbits of a Coxeter group*, 181–211, Surv.Diff.Geom., IV, Int.Press, 1998.
- [12] B. Dubrovin and Y.Zhang, *Extended affine Weyl groups and Frobenius manifolds*, Compositio Mathematica 111(1998) 167–219.
- [13] M.Bertola, *Jacobi groups, Jacobi forms and their applications*, Phd. dissertation. 1999 (SISSA).

- [14] P. Slodowy, *A remark on a recent paper by B. Dubrovin and Y. Zhang*, Preprint 1997.
- [15] I.A.B. Strachan, *Frobenius submanifolds*, J. Geom. Phys. 38 (3-4) (2001) 285–307.
- [16] C.Hertling, *Frobenius manifolds and moduli spaces for singularities*. Cambridge Tracts in Mathematics, 151 Cambridge University Press, Cambridge, 2002.
- [17] B.Dubrovin, *On almost duality for Frobenius manifolds*, Geometry, topology, and mathematical physics, 75–132, Amer. Math. Soc. Transl. Ser. 2, 212, 2004.
- [18] D.Zuo, *Frobenius manifolds associated to B_l and D_l , revisited*, International Mathematics Research Notices, Vol. 2007, Article ID rnm020, 25 pages.
- [19] B.Dubrovin, I.A.B.Strachan, Y.Zhang and D.Zuo, *Extended affine Weyl groups of BCD-type: their Frobenius manifolds and Landau–Ginzburg superpotentials*, To appear in Adv.Math., (arXiv:1510.08690).
- [20] G. Carlet, B. Dubrovin and L.P. Mertens, *Infinite-dimensional Frobenius manifolds for $2+1$ integrable systems*, Math. Ann. 349 (1) (2011) 75–115.
- [21] C-Z.Wu and D.Xu, *A class of infinite-dimensional Frobenius manifolds and their submanifolds*, Int. Math. Res. Not. IMRN (19) (2012) 4520–4562.
- [22] C.-Z. Wu and D.Zuo, *Infinite-dimensional Frobenius manifolds underlying the Toda lattice hierarchy*, Adv. Math. 255 (2014) 487–524.
- [23] T. Milanov, Y. Shen and H.-H. Tseng, *Gromov–Witten theory of Fano orbifold curves, Gamma integral structures and ADE-Toda Hierarchies*, Geom. Topol. **20** (2016), 2135–2218.
- [24] A.Brini,G.Carlet,S.Romano and P.Rossi, *Rational reductions of the 2D-Toda hierarchy and mirror symmetry*, J. Eur. Math. Soc., 19 (2017) no. 3, 835–880.
- [25] X.-K. Chang, X.-B. Hu and S.-H. Li, *Degasperis-Procesi peakon dynamical system and finite Toda lattice of CKP type*, Nonlinearity, 31 (2018) 4746–4775.
- [26] S.-H. Li, *Private communications*, May 2019.
- [27] R.Stedman and I.A.B.Strachan, *Extended V-systems and almost-duality for extended affine Weyl orbit spaces*, arXiv:1612.03667.

SCHOOL OF MATHEMATICAL SCIENCE, UNIVERSITY OF SCIENCE AND TECHNOLOGY OF CHINA, HEFEI 230026, P.R.CHINA

E-mail address: dfzuo@ustc.edu.cn