

Wiener index and Harary index on pancyclic graphs

Huicai Jia ^{*}; Hongye Song [†]

Abstract

Wiener index and Harary index are two classic and well-known topological indices for the characterization of molecular graphs. Recently, Yu et al. [14] established some sufficient conditions for a graph to be pancyclic in terms of the edge number, the spectral radius and the signless Laplacian spectral radius of the graph. In this paper, we give some sufficient conditions for a graph being pancyclic in terms of the Wiener index, the Harary index, the distance spectral radius and the Harary spectral radius of a graph.

AMS Classification: 05C50

Key words: pancyclic graphs; Wiener index; distance spectral radius; Harary index; Harary spectral radius

1 Introduction

All graphs considered here are finite, undirected and connected simple graphs. Let G be a graph with vertex set $V(G) = \{v_1, v_2, \dots, v_n\}$ and edge set $E(G)$. Let $N_G(v)$ denote the neighbour set of v in G . Denote by d_i or $d(v_i)$ the degree of vertex v_i . Let (d_1, d_2, \dots, d_n) be the degree sequence of G , where $d_1 \leq d_2 \leq \dots \leq d_n$. Then $d_1 = \delta$ is called the minimum degree of G . We use $d_G(v_i, v_j)$ to denote the distance between vertices v_i and v_j . The *union* of simple graphs G and H is the graph $G \cup H$ with vertex set $V(G) \cup V(H)$ and edge set $E(G) \cup E(H)$. If G and H are disjoint, we refer to their union as a disjoint union, and denote it by $G + H$. The disjoint union of k graphs G is denoted by kG . By starting with a disjoint union of G and H and adding edges joining every vertex of G to every vertex of H , one can obtain the *join* of G and H , denoted by $G \vee H$. Let \overline{G} denote the complement of G .

A pancyclic graph is a graph that contains cycles of all possible lengths from three up to the number of vertices in the graph. Pancyclic graphs are a generalization of Hamiltonian graphs, which have a cycle of the maximum possible length.

The distance matrix $D(G)$ is defined so that (i, j) -entry, d_{ij} , is equal to $d_G(v_i, v_j)$. The *Wiener index* of a molecular graph was introduced by and named by Wiener [13] in 1947. It

^{*}College of Science, Henan University of Engineering, Zhengzhou, Henan 451191, China; School of Mathematics, Renmin University of China, Beijing, 100872, China. Email: jhc607@163.com

[†]School of General Education, Beijing International Studies University, Beijing, China. Email: songhongye@bisu.edu.cn

is defined as the sum of distances between all pairs of vertices of a connected graph G , i.e., $W(G) = \sum_{u,v \in V(G)} d_G(u,v)$. Let $D_i(G)$ and $D_v(G)$ denote the sum of row i of $D(G)$ and the row sum of $D(G)$ corresponding to vertex v , respectively. Then

$$W(G) = \frac{1}{2} \sum_{v \in V(G)} D_v(G) = \frac{1}{2} \sum_{i=1}^n D_i(G).$$

The *distance spectral radius* of G is the largest eigenvalue of $D(G)$, denoted by $\rho(G)$.

The Harary index of a graph G , denoted by $H(G)$, has been introduced independently by Ivanciuc et al. [5] and Plavšić et al. [12] in 1993 for the characterization of molecular graphs. The *Harary index* $H(G)$ is defined as the sum of reciprocals of distances between all pairs of vertices of a connected graph G , i.e. $H(G) = \sum_{u,v \in V(G)} \frac{1}{d_G(u,v)}$. Note that in any disconnected graph G , the distance is infinite between any two vertices from two distinct components. Therefore its reciprocal can be viewed as zero. Thus we can define validly the Harary index of disconnected graph G as follows: $H(G) = \sum_{i=1}^k H(G_i)$, where G_1, G_2, \dots, G_k are the components of G . We often use $\hat{D}_i(G)$ or $\hat{D}_{v_i}(G)$ to denote $\sum_{v_j \in V(G)} \frac{1}{d_G(v_i, v_j)}$. Then

$$H(G) = \frac{1}{2} \sum_{v_i \in V(G)} \hat{D}_{v_i}(G) = \frac{1}{2} \sum_{i=1}^n \hat{D}_i(G).$$

The Harary matrix $RD(G)$ of G , which is initially called the reciprocal distance matrix and introduced by [5], is an $n \times n$ matrix whose (i, j) -entry is equal to $\frac{1}{d_{ij}}$ if $i \neq j$ and 0 otherwise. The *Harary spectral radius* of G is the largest eigenvalue of $RD(G)$, denoted by $\rho^*(G)$. Note that in any disconnected graph G , $\rho^*(G) = \max\{\rho^*(G_i) | 1 \leq i \leq k\}$.

The problem of deciding whether a given graph is Hamiltonian is NP-complete. Many reasonable sufficient or necessary conditions were given. Recently, spectral graph theory has been applied to the problem. Some related sufficient spectral conditions on the Wiener index and the Harary index for a graph to be Hamiltonian, traceable and Hamiltonian-connected have been given in [1, 2, 3, 6, 7, 8, 9, 10, 11, 15, 16].

In this paper, we consider the problem of deciding whether a given graph is pancyclic. Yu et al. [14] established some sufficient conditions for a graph to be pancyclic in terms of the edge number, the spectral radius and the signless Laplacian spectral radius of the graph. Motivated by these results, we present some sufficient conditions for a graph to be pancyclic in terms of the Wiener index, the Harary index, the distance spectral radius and the Harary spectral radius of a graph, respectively.

2 Preliminaries

Before giving the proof of our theorems, we introduce some fundamental lemmas and properties in this section.

Let $\text{NP} = \{K_2 \vee (K_{n-4} + 2K_1), K_5 \vee 6K_1, K_3 \vee (K_2 + 3K_1), K_3 \vee (K_{1,4} + K_1), K_3 \vee (K_{1,3} + K_2), (K_2 \vee 2K_1) \vee 5K_1, K_4 \vee 5K_1, K_{1,2} \vee 4K_1, K_2 \vee (K_{1,3} + K_1), K_3 \vee 4K_1\}$.

Lemma 2.1 ([14]) *Let G be a connected graph on $n \geq 5$ vertices and m edges with minimum degree $\delta \geq 2$. If $m \geq \binom{n-2}{2} + 4$, then G is a pancyclic graph unless $G \in \mathbb{NP}$ or G is a bipartite graph.*

Lemma 2.2 ([4]) *Let G be a connected graph on n vertices. Then $\rho(G) \geq \frac{2W(G)}{n}$, and the equality holds if and only if the row sums of $D(G)$ are all equal.*

Lemma 2.3 ([10]) *Let G be a graph on n vertices. Then $\rho^*(G) \geq \frac{2H(G)}{n}$, and the equality holds if and only if the row sums of $RD(G)$ are all equal.*

Lemma 2.4 *Let G be a bipartite graph on n vertices. Then $W(G) > \frac{n^2+3n-14}{2}$.*

Proof. Let G be a bipartite graph of order n whose vertices are divided into two disjoint and independent sets V_1 and V_2 , $|V_1| = a$, ($1 \leq a \leq n-1$), $|V_2| = n-a$.

$$\begin{aligned}
W(G) &= \frac{1}{2} \sum_{i \in V(G)} D_i(G) \\
&\geq \frac{1}{2} \left[\sum_{i \in V_1} (d_i + 3(n-a-d_i) + 2(a-1)) + \sum_{j \in V_2} (d_j + 3(a-d_j) + 2(n-a-1)) \right] \\
&= \frac{1}{2} \left[\sum_{i \in V_1} (3n-a-2-2d_i) + \sum_{j \in V_2} (2n+a-2-2d_j) \right] \\
&= -a^2 + na + n^2 - n - \sum_{i \in V(G)} d_i \\
&= -a^2 + na + n^2 - n - 2 \\
&\geq -a^2 + na + n^2 - n - 2a(n-a) = a^2 - na + n^2 - n \\
&\geq \frac{3n^2 - 4n}{4}. \tag{1}
\end{aligned}$$

By manipulation, we have $\frac{3n^2-4n}{4} > \frac{n^2+3n-14}{2}$. Then $W(G) > \frac{n^2+3n-14}{2}$. The proof of Lemma 2.4 is completed.

Lemma 2.5 *Let G be a bipartite graph on n vertices. Then $H(\overline{G}) > \frac{5n^2-23n+28}{2(n-1)}$ for $n \geq 8$.*

Proof. Let G be a bipartite graph of order $n \geq 8$ whose vertices are divided into two disjoint and independent sets V_1 and V_2 , $|V_1| = a$, ($1 \leq a \leq n-1$), $|V_2| = n-a$.

Case 1. G is not a complete bipartite graph.

$$\begin{aligned}
H(\overline{G}) &= \frac{1}{2} \sum_{i=1}^n \hat{D}_i(\overline{G}) = \frac{1}{2} \sum_{v \in V(G)} [d_{\overline{G}}(v) + \frac{1}{2}(n-1-d_{\overline{G}}(v))] \\
&= \frac{1}{2} \sum_{v \in V(G)} \left[\frac{n-1}{2} + \frac{d_{\overline{G}}(v)}{2} \right] = \frac{n(n-1)}{4} + \frac{1}{4} \sum_{v \in V(G)} d_{\overline{G}}(v) \\
&= \frac{n(n-1)}{4} + \frac{1}{4} \sum_{v \in V(G)} (n-1-d_G(v)) \\
&= \frac{n(n-1)}{2} - \frac{m}{2} \\
&> \frac{n(n-1)}{2} - \frac{1}{2}a(n-a) \\
&= \frac{a^2 - na + n^2 - n}{2} \\
&\geq \frac{3n^2 - 4n}{8}
\end{aligned}$$

Through calculation, we have $\frac{3n^2-4n}{8} > \frac{5n^2-23n+28}{2(n-1)}$. Then $H(\overline{G}) > \frac{5n^2-23n+28}{2(n-1)}$ for $n \geq 3$.

Case 2. G is a complete bipartite graph.

Then $G = K_{a,n-a}$ and

$$\begin{aligned}
H(\overline{G}) &= \frac{1}{2} \sum_{i=1}^n \hat{D}_i(\overline{G}) = \frac{1}{2} \left[\sum_{v \in V_1} (a-1) + \sum_{v \in V_2} (n-a-1) \right] \\
&= \frac{1}{2} [a(a-1) + (n-a)(n-a-1)] \\
&= a^2 - na + \frac{n^2 - n}{2} \\
&\geq \frac{n^2 - 2n}{4}
\end{aligned}$$

By manipulation, we have $\frac{n^2-2n}{4} > \frac{5n^2-23n+28}{2(n-1)}$ for $n \geq 8$. Then $H(\overline{G}) > \frac{5n^2-23n+28}{2(n-1)}$ for $n \geq 8$. The proof of Lemma 2.5 is completed.

3 Wiener index, Harary index on pancyclic graphs

Theorem 3.1 *Let G be a connected graph of order $n \geq 5$ with minimum degree $\delta \geq 2$. If $W(G) \leq \frac{n^2+3n-14}{2}$, then G is a pancyclic graph unless $G \in \mathbb{NP}$.*

Proof. Suppose that G is not a pancyclic graph. Then

$$\begin{aligned}
W(G) &= \frac{1}{2} \sum_{i=1}^n D_i(G) \geq \frac{1}{2} \sum_{i=1}^n (d_i + 2(n-1-d_i)) \\
&= \frac{1}{2} \sum_{i=1}^n (2(n-1) - d_i) \\
&= n(n-1) - m. \tag{2}
\end{aligned}$$

Note that $W(G) \leq \frac{n^2+3n-14}{2}$, we have $m \geq n(n-1) - \frac{n^2+3n-14}{2} = \binom{n-2}{2} + 4$. By Lemma 2.1, we obtain that $G \in \mathbb{NP}$ or G is a bipartite graph. By a direct calculation, for all $G \in \mathbb{NP}$, $W(G) \leq \frac{n^2+3n-14}{2}$. If G is a bipartite graph, by Lemma 2.4, we obtain $W(G) > \frac{n^2+3n-14}{2}$. This completes the proof.

Theorem 3.2 *Let G be a connected graph of order $n \geq 5$ with minimum degree $\delta \geq 2$, and \overline{G} be a connected graph. If $W(\overline{G}) \geq \frac{n^3-6n^2+23n-28}{2}$, then G is a pancyclic graph unless G is a bipartite graph which does not contain complete bipartite graph.*

Proof. Suppose that G is not a pancyclic graph. Then

$$\begin{aligned} W(\overline{G}) &= \frac{1}{2} \sum_{i=1}^n D_i(\overline{G}) \leq \frac{1}{2} \sum_{v \in V(G)} [d_{\overline{G}}(v) + (n-1)(n-1-d_{\overline{G}}(v))] \\ &= \frac{1}{2} \sum_{v \in V(G)} [(n-1)^2 + (2-n)d_{\overline{G}}(v)] \\ &= \frac{1}{2}n(n-1)^2 - \frac{n-2}{2} \sum_{v \in V(G)} (n-1-d_G(v)) \\ &= \frac{n(n-1)}{2} + (n-2)m. \end{aligned}$$

Note that $W(\overline{G}) \geq \frac{n^3-6n^2+23n-28}{2}$, we have $m \geq \frac{n^3-7n^2+24n-28}{2(n-2)} = \binom{n-2}{2} + 4$. By Lemma 2.1, we obtain that $G \in \mathbb{NP}$ or G is a bipartite graph. Note that for all $G \in \mathbb{NP}$, \overline{G} is disconnected, and G is a complete bipartite graph, \overline{G} is also disconnected. The proof of Theorem 3.2 is completed.

Theorem 3.3 *Let G be a connected graph of order $n \geq 5$ with minimum degree $\delta \geq 2$. If $H(G) \geq \frac{n^2-3n+7}{2}$, then G is a pancyclic graph unless $G \in \mathbb{NP}$.*

Proof. Suppose that G is not a pancyclic graph. Then

$$\begin{aligned} H(G) &= \frac{1}{2} \sum_{i=1}^n \hat{D}_i(G) \leq \frac{1}{2} \sum_{i=1}^n (d_i + \frac{1}{2}(n-1-d_i)) \\ &= \frac{n(n-1)}{4} + \frac{m}{2}. \end{aligned}$$

Note that $H(G) \geq \frac{n^2-3n+7}{2}$, we have $m \geq n^2 - 3n + 7 - \frac{n(n-1)}{2} = \binom{n-2}{2} + 4$. By Lemma 2.1, we obtain that $G \in \mathbb{NP}$ or G is a bipartite graph. By a direct calculation, for all $G \in \mathbb{NP}$, $H(G) \geq \frac{n^2-3n+7}{2}$. If G is a bipartite graph whose vertices are divided into two disjoint and

independent sets V_1 and V_2 , $|V_1| = a$, ($1 \leq a \leq n-1$), $|V_2| = n-a$. We have

$$\begin{aligned}
H(G) &= \frac{1}{2} \sum_{i \in V(G)} \hat{D}_i(G) = \frac{1}{2} \left(\sum_{i \in V_1} \hat{D}_i(G) + \sum_{j \in V_2} \hat{D}_j(G) \right) \\
&\leq \frac{1}{2} \left[\sum_{i \in V_1} \left(d_i + \frac{1}{3}(n-a-d_i) + \frac{1}{2}(a-1) \right) + \sum_{j \in V_2} \left(d_j + \frac{1}{3}(a-d_j) + \frac{1}{2}(n-a-1) \right) \right] \\
&= \frac{2a^2 - 2na + 3n^2 - 3n}{12} + \frac{1}{3} \sum_{i \in V(G)} d_i = \frac{2a^2 - 2na + 3n^2 - 3n}{12} + \frac{2m}{3} \\
&\leq \frac{2a^2 - 2na + 3n^2 - 3n}{12} + \frac{2a(n-a)}{3} = \frac{-2a^2 + 2na + n^2 - n}{4} \leq \frac{3n^2 - 2n}{8}.
\end{aligned}$$

Note that $\frac{3n^2-2n}{8} < \frac{n^2-3n+7}{2}$. We complete the proof.

Theorem 3.4 *Let G be a connected graph on $n \geq 8$ vertices with minimum degree $\delta \geq 2$. If $H(\overline{G}) \leq \frac{5n^2-23n+28}{2(n-1)}$, then G is a pancyclic graph unless $G \in \{K_5 \vee 6K_1, K_3 \vee (K_2 + 3K_1), K_3 \vee (K_{1,4} + K_1), K_3 \vee (K_{1,3} + K_2), (K_2 \vee 2K_1) \vee 5K_1, K_4 \vee 5K_1\}$.*

Proof. Suppose that G is not a pancyclic graph. Then

$$\begin{aligned}
H(\overline{G}) &= \frac{1}{2} \sum_{i=1}^n \hat{D}_i(\overline{G}) \geq \frac{1}{2} \sum_{v \in V(G)} \left[d_{\overline{G}}(v) + \frac{1}{n-1}(n-1-d_{\overline{G}}(v)) \right] \\
&= \frac{1}{2} \sum_{v \in V(G)} \left[1 + \frac{n-2}{n-1} d_{\overline{G}}(v) \right] \\
&= \frac{n}{2} + \frac{n-2}{2(n-1)} \sum_{v \in V(G)} (n-1-d_G(v)) \\
&= \frac{n(n-1)}{2} - \frac{n-2}{2(n-1)} \sum_{v \in V(G)} d_G(v) \\
&= \frac{n(n-1)}{2} - \frac{n-2}{n-1} m. \tag{3}
\end{aligned}$$

Note that $H(\overline{G}) \leq \frac{5n^2-23n+28}{2(n-1)}$, we have $m \geq \frac{n(n-1)^2}{2(n-2)} - \frac{5n^2-23n+28}{2(n-2)} = \binom{n-2}{2} + 4$. By Lemma 2.1, we obtain that $G \in \mathbb{NP}$ or G is a bipartite graph. Note that $n \geq 8$, $H(\overline{K_2 \vee (K_{n-4} + 2K_1)}) \geq \frac{n^2-n-8}{4} > \frac{5n^2-23n+28}{2(n-1)}$. However, by a direct calculation, for all $G \in \mathbb{NP} \setminus K_2 \vee (K_{n-4} + 2K_1)$, $H(\overline{G}) \leq \frac{5n^2-23n+28}{2(n-1)}$. If G is a bipartite graph, by Lemma 2.4, we have $H(\overline{G}) > \frac{5n^2-23n+28}{2(n-1)}$ for $n \geq 8$. So we obtain this theorem.

Theorem 3.5 *Let G be a connected graph of order $n \geq 5$ with minimum degree $\delta \geq 2$. If $\rho(G) \leq n+3 - \frac{14}{n}$, then G is a pancyclic graph unless $G \in \{K_2 \vee (K_{1,3} + K_1), K_3 \vee 4K_1\}$.*

Proof. Assume that G is not a pancyclic graph. By Lemma 2.2 and (2) in the proof of Theorem 3.1, we have

$$\rho(G) \geq \frac{2W(G)}{n} \geq \frac{2}{n} [n(n-1) - m] = 2(n-1) - \frac{2}{n} m.$$

Since $\rho(G) \leq n + 3 - \frac{14}{n}$, we have $m \geq \binom{n-2}{2} + 4$. According to Lemma 2.1, $G \in \text{NP}$ or G is a bipartite graph.

For $G = K_2 \vee (K_{n-4} + 2K_1)$, let $X = (x_1, x_2, \dots, x_n)^T$ be the eigenvector corresponding to $\rho(G)$, where x_i ($1 \leq i \leq n-4$) corresponds to the vertex of degree $n-3$, x_j ($n-3 \leq j \leq n-2$) corresponds to the vertex of degree 2 and x_l ($n-1 \leq l \leq n$) corresponds to the vertex of degree $n-1$. Without loss of generality, let $x := x_i, 1 \leq i \leq n-4$; $y := x_j, n-1 \leq j \leq n$; $z := x_l, n-1 \leq l \leq n$. As $\rho X = D(G)X$, we have

$$\begin{cases} \rho x = (n-5)x + 4y + 2z, \\ \rho y = 2(n-4)x + 2y + 2z, \\ \rho z = (n-4)x + 2y + z. \end{cases}$$

It follows that $\rho(K_2 \vee (K_{n-4} + 2K_1))$ is the largest root of the equation

$$\rho^3 - (n-2)\rho^2 - (7n-23)\rho - 2(n-3) = 0.$$

Let $f(x) = \rho^3 - (n-2)\rho^2 - (7n-23)\rho - 2(n-3)$, then $n \geq 5$, $f(n+3 - \frac{14}{n}) = -2n^2 + 16n - 6 - \frac{476}{n} + \frac{2156}{n^2} - \frac{2744}{n^3} < 0$. Hence $\rho(K_2 \vee (K_{n-4} + 2K_1)) > n + 3 - \frac{14}{n}$ for $n \geq 5$. By a direct calculation, we have the following Table 1.

G	$\rho(G)$	$n + 3 - \frac{14}{n}$
$K_5 \vee 6K_1$	13.245	12.727
$K_3 \vee (K_2 + 3K_1)$	9.5947	9.25
$K_3 \vee (K_{1,4} + K_1)$	10.8341	10.444
$K_3 \vee (K_{1,3} + K_2)$	10.7672	10.444
$(K_2 \vee 2K_1) \vee 5K_1$	10.7624	10.444
$K_4 \vee 5K_1$	10.6325	10.444
$K_{1,2} \vee 4K_1$	8.2736	8
$K_2 \vee (K_{1,3} + K_1)$	8	8
$K_3 \vee 4K_1$	8	8

Table 1. The distance spectral radius of some graphs

If G is a bipartite graph, by (1) the proof of Lemma 2.4, we have $W(G) \geq \frac{3n^2-4n}{4}$, then $\rho(G) \geq \frac{2W(G)}{n} \geq \frac{3n-4}{2} > n + 3 - \frac{14}{n}$. This completes the proof of Theorem 3.5.

Theorem 3.6 *Let G be a connected graph on $n \geq 8$ vertices with $\delta \geq 2$. If $\rho^*(\overline{G}) \leq \frac{5n^2-23n+28}{n(n-1)}$, then G is a pancyclic graph.*

Proof. Suppose that G is not a pancyclic graph. From Lemma 2.3 and (3) in the proof of Theorem 3.4, we have

$$\rho^*(\overline{G}) \geq \frac{2H(\overline{G})}{n} \geq n-1 - \frac{2(n-2)m}{n(n-1)}.$$

Since $\rho^*(\overline{G}) \leq \frac{5n^2-23n+28}{n(n-1)}$, we have $m \geq \binom{n-2}{2} + 4$. According to Lemma 2.1, we obtain $G \in \text{NP}$ or G is a bipartite graph.

For $G = K_2 \vee (K_{n-4} + 2K_1)$, let $X = (x_1, x_2, \dots, x_n)^T$ be the eigenvector corresponding to $\rho^*(\overline{G})$, where x_i ($1 \leq i \leq n-4$) corresponds to the vertex of degree 2, x_j ($n-3 \leq j \leq n-2$) corresponds to the vertex of degree $n-3$ and x_l ($n-1 \leq l \leq n$) corresponds to the vertex of degree 0. Without loss of generality, let $x := x_i, 1 \leq i \leq n-4$; $y := x_j, n-1 \leq j \leq n$; $z := x_l, n-1 \leq l \leq n$. As $\rho X = RD(G)X$, we have

$$\begin{cases} \rho^* x = \frac{1}{2}(n-5)x + 2y + 0z, \\ \rho^* y = (n-4)x + y + 0z. \end{cases}$$

It follows that $\rho(K_2 \vee (K_{n-4} + 2K_1))$ is the largest root of the equation

$$2(\rho^*)^2 - (n-3)\rho^* + 11 - 3n = 0.$$

Then $\rho^* = \frac{n-3+\sqrt{n^2+18n-79}}{4} > \frac{5n^2-23n+28}{n(n-1)}$ for $n \geq 5$. By a direct calculation, we have the following Table 2.

G	$\rho^*(\overline{G})$	$\frac{5n^2-23n+28}{n(n-1)}$
$K_5 \vee 6K_1$	5	3.4546
$K_3 \vee (K_2 + 3K_1)$	4.0663	2.9286
$K_3 \vee (K_{1,4} + K_1)$	4.4115	3.1389
$K_3 \vee (K_{1,3} + K_2)$	4.5455	3.1389
$(K_2 \vee 2K_1) \vee 5K_1$	4	3.1389
$K_4 \vee 5K_1$	4	3.1389
$K_{1,2} \vee 4K_1$	3	2.667
$K_2 \vee (K_{1,3} + K_1)$	3.4575	2.667
$K_3 \vee 4K_1$	3	2.667

Table 2. The Harary spectral radius of complements of some graphs

Clearly, for $G \in \mathbb{NP}$, $\rho^*(\overline{G}) > \frac{5n^2-23n+28}{n(n-1)}$.

If G is a bipartite graph of order $n \geq 8$ whose vertices are divided into two disjoint and independent sets V_1 and V_2 , $|V_1| = a, (1 \leq a \leq n-1), |V_2| = n-a$.

Case 1. G is not a complete bipartite graph.

By Lemma 2.3 and the proof of Lemma 2.5, we have $\rho^*(\overline{G}) \geq \frac{3n-4}{4} > \frac{5n^2-23n+28}{n(n-1)}$ for $n \geq 3$.

Case 2. G is a complete bipartite graph.

Then $G = K_{a,n-a}$.

By Lemma 2.3 and the proof of Lemma 2.5, we have $\rho^*(\overline{G}) \geq \frac{n-2}{2} > \frac{5n^2-23n+28}{n(n-1)}$ for $n \geq 8$. Hence we obtain the theorem.

4 Conclusions

We consider the problem of deciding whether a given graph is pancyclic. We present some sufficient conditions for a graph to be pancyclic in terms of the Wiener index, the Harary index, the distance spectral radius and the Harary spectral radius of a graph, respectively.

References

- [1] Feng, L, Zhu, X, Liu, W: Wiener index, Harary index and graph properties, *Discrete Appl. Math.* **223**, 72-83 (2017)
- [2] Hua, H, Wang, M: On Harary index and traceable graphs, *MATCH Commun. Math. Comput. Chem.* **70**, 297-300 (2013)
- [3] Hua, H, Ning, B: Wiener Index, Harary Index and Hamiltonicity of Graphs, *MATCH Commun. Math. Comput. Chem.* **78**, 153-162 (2017)
- [4] Indulal, G: Sharp bounds on the distance spectral radius and the distance energy of graphs, *Linear Algebra Appl.* **430**, 106-113 (2009)
- [5] Ivanciuc, O, Balaban, T, Balaban, A: Reciprocal distance matrix, related local vertex invariants and topological indices, *J. Math. Chem.* **12**, 309-318 (1993)
- [6] Jia, H, Liu, R, Du, X: Wiener index and Harary index on Hamilton-connected and traceable graphs, *Ars Combinatoria* **141**, 53-62 (2018)
- [7] Kuang, M, Huang, G, Deng, H: Some sufficient conditions for Hamiltonian property in terms of Wiener-type invariants, *Proceedings Mathematical Sciences* **126**, 1-9 (2016)
- [8] Li, R: Wiener index and some Hamiltonian properties of graphs, *International Journal of Mathematics and Soft Computing* **5**, 11-16 (2015)
- [9] Li, R: Harary index and some Hamiltonian properties of graphs, *AKCE International Journal of Graphs and Combinatorics* **12**, 64-69 (2015)
- [10] Liu, R, Du, X, Jia, H: Some observations on Harary index and traceable graphs, *MATCH Commun. Math. Comput. Chem.* **77**, 195-208 (2017)
- [11] Liu, R, Du, X, Jia, H: Wiener index on traceable and Hamiltonian graphs, *Bull. Aust. Math. Soc.* **94**, 362-372 (2016)
- [12] Plavšić, D, Nikolić, S, Trinajstić, N, Mihalić, Z: On the Harary index for the characterization of chemical graphs, *J. Math. Chem.* **12**, 235-250 (1993)
- [13] Wiener, H: Structural determination of paraffin boiling points, *Journal of the American Chemical Society* **69**, 17-20 (1947)
- [14] Yu, G, Yu, T, Shu, A, Xia, X: Some sufficient conditions on pancyclic graphs, *arXiv: 1809.09897v1 [math.CO] 26 Sep 2018*.
- [15] Yang, L: Wiener index and traceable graphs, *Bull. Aust. Math. Soc.* **88**, 380-383 (2013)
- [16] Zeng, T: Harary index and Hamiltonian property of graphs, *MATCH Commun. Math. Comput. Chem.* **70**, 645-649 (2013)