

On extremal results of multiplicative Zagreb indices of trees with given distance k -domination number

Fazal Hayat

School of Mathematical Sciences, South China Normal University,

Guangzhou 510631, PR China

E-mail: fhayatmaths@gmail.com

Abstract

The first multiplicative Zagreb index Π_1 of a graph G is the product of the square of every vertex degree, while the second multiplicative Zagreb index Π_2 is the product of the products of degrees of pairs of adjacent vertices. In this paper, we give sharp lower bound for Π_1 and upper bound for Π_2 of trees with given distance k -domination number, and characterize those trees attaining the bounds.

Key Words: first multiplicative Zagreb index, second multiplicative Zagreb index, trees, distance k -domination number.

Mathematics Subject Classification (2010): 05C05, 05C35, 05C69

1 Introduction

In this article we consider only simple, undirected and connected graphs. Let G be a graph with vertex set $V(G)$ and edge set $E(G)$. The degree of $v \in V(G)$, denoted by $d_G(v)$, is the number of vertices in G adjacent to v , and the neighborhood of v is the set $N_G(v) = \{w \in V(G) : vw \in E(G)\}$. Evidently, $|N_G(v)| = d_G(v)$. A vertex

with degree one is called pendent vertex. The distance between any two vertices u and v of a graph G is denoted by $d_G(u, v)$. The maximum distance from a vertex $v \in V(G)$ to all other vertices of G is called eccentricity of v in G . The diameter of a graph G is the maximum eccentricity of all vertices of G .

A graph G that has n vertices and $n - 1$ edges is called a tree. As usual, by P_n and S_n we denote the path and the star on n vertices, respectively.

The first and the second Zagreb indices are among the oldest topological molecular descriptors, see [5]. They are defined as follows:

$$M_1(G) = \sum_{u \in V(G)} d(u)^2 \text{ and } M_2(G) = \sum_{uv \in E(G)} d(u)d(v).$$

Many interesting properties of them may be found in [2, 6, 7, 15, 20, 21].

In 2010, Todeschini et al. [4, 13] put forward the multiplicative Zagreb indices as follows:

$$\Pi_1(G) = \prod_{u \in V(G)} d(u)^2 \text{ and } \Pi_2(G) = \prod_{uv \in E(G)} d(u)d(v).$$

It is easily seen that $\Pi_2(G) = \prod_{uv \in E(G)} d(u)d(v) = \prod_{u \in V(G)} d(u)^{d(u)}$. Some properties for the multiplicative Zagreb indices have been established, see [3, 8, 9, 10, 16, 17].

For a positive integer k , a set $D \subseteq V(G)$ is said to be distance k -dominating set of G if for every vertex $u \in V(G) \setminus D$, $d_G(u, v) \leq k$ for some vertex $v \in D$. The minimum cardinality among all distance k -dominating set of G is called the distance k -domination number of G , denoted by $\gamma_k(G)$. A distance 1-dominating set of G is known as a dominating set of G and the distance 1-domination number of G is just the classical domination number of G .

Borovicanin and Furtula [1] presented sharp lower and upper bounds on Zagreb indices of trees in terms of domination number, and Wang et al. [18] found sharp lower and upper bounds on multiplicative Zagreb indices of trees in terms of domination number. Recently, Pei and Pan [12] investigated the connection between the Zagreb indices and the distance k -domination number of trees.

Motivated by the above results, in this paper, we study the multiplicative Zagreb indices of trees in terms of distance k -domination number. We provide sharp lower

bound for Π_1 and upper bound for Π_2 in terms of distance k -domination number of a tree, and characterize those trees for which the bounds are attained.

2 Preliminaries

In this section, we present some propositions, definitions and lemmas which are helpful in our main results.

Lemma 2.1. [3] *Let T be a tree of order $n > 5$ such that $T \not\cong P_n, S_n$. Then*

$$\Pi_1(S_n) < \Pi_1(T) < \Pi_1(P_n) \text{ and } \Pi_2(P_n) < \Pi_2(T) < \Pi_2(S_n).$$

Let T be a tree and $uv \in E(T)$ a non-pendent edge of T . Assume that $T - uv = T_1 \cup T_2$ with vertex $u \in V(T_1)$ and $v \in V(T_2)$. Let T_{uv} be the tree obtained by identifying the vertex u of T_1 and the vertex v of T_2 and attaching a pendent vertex w to this vertex.

Lemma 2.2. [19] *Let T be a tree with a non-pendent edge uv . Then*

$$\Pi_1(T_{uv}) < \Pi_1(T) \text{ and } \Pi_2(T_{uv}) > \Pi_2(T).$$

Lemma 2.3. [19] *Let u and v be two distinct vertices in a graph G . Let u_1, \dots, u_r be pendent neighbors of u and v_1, \dots, v_t pendent neighbors of v . Define $G' = G - \{vv_1, \dots, vv_t\} + \{uv_1, \dots, uv_t\}$ and $G'' = G - \{uu_1, \dots, uu_r\} + \{vu_1, \dots, vu_r\}$. Then*

$$\max\{\Pi_1(G'), \Pi_1(G'')\} < \Pi_1(G)$$

and

$$\min\{\Pi_1(G'), \Pi_1(G'')\} > \Pi_2(G).$$

Lemma 2.4. [11] *Let G be a connected graph of order n with $n \geq k + 1$. Then*
 $\gamma_k(G) \leq \lfloor \frac{n}{k+1} \rfloor$.

Lemma 2.5. [12] *Let T be a tree of order n with maximum degree Δ and distance k -domination number $\gamma_k \geq 2$. Then $\Delta \leq n - k\gamma_k$.*

Lemma 2.6. [14] *Let T be a tree on $(k+1)n$ vertices. Then $\gamma_k(T) = n$ if and only if one of the following conditions holds:*

- (1) T is any tree on $k+1$ vertices;
- (2) $T = R \circ k$ for some tree R on $n \geq 1$ vertices, where $R \circ k$ is the graph obtained by taking one copy of R and $|V(R)|$ copies of the path P_{k-1} of length $k-1$ and then joining the i th vertex of R to exactly one end vertex in the i th copy of P_{k-1} .

Lemma 2.7. *Let T be a tree of order n . Let*

$$f(T) = \prod_{w \in V(T)} (d_T(w) + 1).$$

Then $f(T) \geq 2^{n-1}n$ with equality if and only if $T \cong S_n$.

Proof. If $n = 1$, it is obviously. Suppose that $n \geq 2$ and that the result is true for a tree of order $n-1$. Let u be a pendent vertex of T , being adjacent to vertex v . By induction assumption,

$$f(T - u) \geq 2^{n-2}(n-1)$$

i.e.,

$$d_T(v) \prod_{w \in V(T) \setminus \{u,v\}} (d_T(w) + 1) \geq 2^{n-2}(n-1)$$

with equality if and only if $T - u \cong S_{n-1}$. Now we have

$$\begin{aligned} f(T) &= (d_T(u) + 1)(d_T(v) + 1) \prod_{w \in V(T) \setminus \{u,v\}} (d_T(w) + 1) \\ &= 2(d_T(v) + 1) \prod_{w \in V(T) \setminus \{u,v\}} (d_T(w) + 1) \\ &\geq 2(d_T(v) + 1) \cdot \frac{2^{n-2}(n-1)}{d_T(v)} \\ &\geq 2^{n-1}n. \end{aligned}$$

with equalities if and only if $d_T(v) = n-1$ and $T - u \cong S_{n-1}$, i.e., $T \cong S_n$. □

Lemma 2.8. *Let T be a tree of order n . Let*

$$h(T) = \prod_{w \in V(T)} (d_T(w) + 1)^{d_T(w)+1}.$$

Then $h(T) \leq 4^{n-1}n^n$ with equality if and only if $T \cong S_n$.

Proof. If $n = 1$, it is obviously. Suppose that $n \geq 2$ and that the result is true for a tree of order $n - 1$. Let u be a pendent vertex of T , being adjacent to vertex v . By induction assumption,

$$h(T - u) \leq 4^{n-2}(n - 1)^{n-1}$$

i.e.,

$$(d_T(v))^{d_T(v)} \prod_{w \in V(T) \setminus \{u, v\}} (d_T(w) + 1)^{d_T(w)+1} \leq 4^{n-2}(n - 1)^{n-1}$$

with equality if and only if $T - u \cong S_{n-1}$. Now we have

$$\begin{aligned} h(T) &= (d_T(u) + 1)^{d_T(u)+1} \cdot (d_T(v) + 1)^{d_T(v)+1} \cdot \left(\prod_{w \in V(T) \setminus \{u, v\}} (d_T(w) + 1)^{d_T(w)+1} \right) \\ &= 4(d_T(v) + 1)^{d_T(v)+1} \cdot \prod_{w \in V(T) \setminus \{u, v\}} (d_T(w) + 1)^{d_T(w)+1} \\ &\leq 4(d_T(v) + 1)^{d_T(v)+1} \cdot \frac{4^{n-2}(n - 1)^{n-1}}{(d_T(v))^{d_T(v)}} \\ &\leq 4^{n-1}n^n. \end{aligned}$$

with equalities if and only if $d_T(v) = n - 1$ and $T - u \cong S_{n-1}$, i.e., $T \cong S_n$. \square

For a graph G with $S \subset V(G)$, let $N_G(S) = \cup_{v \in S} N_G(v)$.

Lemma 2.9. *Let T be a tree with minimum value of first multiplicative Zagreb index or maximum value of second multiplicative zagreb index among all n -vertex trees with distance k -domination number γ_k . Let*

$$B_T = \{x \in V(T) | d_T(x) = 1 \text{ and } \gamma_k(T - x) = \gamma_k(T)\}.$$

If $B_T \neq \emptyset$, then $|N_T(B_T)| = 1$.

Proof. Suppose that $|N_T(B_T)| \geq 2$, say $u, v \in N_T(B_T)$. If $u' \notin D$ for some pendent neighbor u' of u , then $D \setminus \{u'\} \cup \{u\}$ is a distance k -dominating set of T . So we may assume that no pendent neighbor of u and v is in D . Define $T' = T - vv' + uv'$ and $T'' = T - uu' + vv'$, where u' (v' , respectively) is a pendent neighbor of u (v , respectively). Then $\gamma_k(T) = \gamma_k(T') = \gamma_k(T'')$. By Lemma 2.3,

$$\max\{\Pi_1(T'), \Pi_1(T'')\} < \Pi_1(T)$$

and

$$\min\{\Pi_1(T'), \Pi_1(T'')\} > \Pi_2(T).$$

a contradiction. Hence $|N_T(B_T)| = 1$. □

3 Main results

In this section, we present sharp lower bounds of first multiplicative Zagreb index and upper bounds for second multiplicative zagreb index of a tree of order n with distance k -domination number γ_k .

A tree is starlike if it contains at most one vertex of degree at least three. Obviously, a starlike tree is either a path or a tree with exactly one vertex of degree at least three. In the latter case, it consists of pendent paths at common vertex.

Definition 3.1. For positive integers n , k and s with $n \geq (k+1)s$, define $T_{n,k,s}$ to be a starlike tree with maximum degree $n - ks$, and if it is not a path, then it has one pendent path of length $k - 1$, $s - 1$ pendent paths of length k and $n - (k+1)s$ paths of length 1.

Note that

$$\Pi_1(T_{n,k,s}) = 4^{ks-1}(n - ks)^2 \text{ and } \Pi_2(T_{n,k,s}) = 4^{ks-1}(n - ks)^{n-ks}.$$

As mentioned earlier, sharp lower bounds on first multiplicative Zagreb index and upper bounds on the second multiplicative Zagreb index of an n -vertex tree with distance 1-domination number have been given in [18], so we only consider $k \geq 2$.

Definition 3.2. If $P = v_0v_1 \dots v_d$ is a diametric path of tree T of order n , then denote by T_i the component of $T - v_{i-1}v_i - v_iv_{i+1}$ containing v_i for $i = 1, \dots, d - 1$.

Definition 3.3. Denoted by $T_{n,k,2}^a$ the tree formed from the path $P_{2k+2} = v_0v_1 \dots v_{2k+1}$ by joining $n - 2(k+1)$ pendent vertices to v_a , where $a \in \{1, \dots, k\}$.

For a graph G , it is obvious that $\gamma_k(G) \leq \gamma_1(G)$ for $k \geq 2$. Note also that $\gamma_1(S_n) = 1$. Thus, by Lemma 2.1, we have the following result.

Theorem 3.1. *Let T be an n -vertex tree and $\gamma_k(T) = 1$. Then $\Pi_1(T) \geq n^2$ and $\Pi_2(T) \leq n^n$. Either equality holds if and only if $T \cong S_n$.*

Theorem 3.2. *Let T be a tree of order n with $\gamma_k(T) = 2$. Then*

$$\Pi_1(T) \geq 4^{2k-1}(n-2k)^2 \text{ and } \Pi_2(T) \leq 4^{2k-1}(n-2k)^{n-2k}.$$

Either equality holds if and only if $T \cong T_{n,k,2}^a$ with $a \in \{1, \dots, k\}$.

Proof. Let T be a tree of order n with distance k -domination number 2 that minimize the first multiplicative Zagreb index and maximize the second multiplicative Zagreb index respectively.

Let $P = v_0 \dots v_d$ be a diametric path of T . If $d \leq 2k$, then $\{v_{\lfloor \frac{d}{2} \rfloor}\}$ is a distance k -dominating set of T , a contradiction. If $d \geq 2k + 2$, for $i \in \{1, \dots, d-2\}$, $T_{v_i v_{i+1}}$ is a tree of order n with distance k -domination number 2, by Lemma 2.2, we have $\Pi_1(T_{v_i v_{i+1}}) < \Pi_1(T)$, and $\Pi_2(T_{v_i v_{i+1}}) > \Pi_2(T)$, also a contradiction. Hence $d = 2k + 1$.

If T_i is not a star for some $i \in \{1, \dots, 2k\}$, then as above, $T_{v_i v_{i+1}}$ is a tree of order n with distance k -domination number 2 such that $\Pi_1(T_{v_i v_{i+1}}) < \Pi_1(T)$, and $\Pi_2(T_{v_i v_{i+1}}) > \Pi_2(T)$, a contradiction. Thus each T_i for $i \in \{1, \dots, d-1\}$ is a star with center v_i . Now by Lemma 2.3, for some $a \in \{1, \dots, k\}$, $T \cong T_{n,k,2}^a$.

By direct calculation, $\Pi_1(T) = \Pi_1(T_{n,k,2}^a) = 4^{2k-1}(n-2k)^2$ and $\Pi_2(T) = \Pi_2(T_{n,k,2}^a) = 4^{2k-1}(n-2k)^{n-2k}$ for $a \in \{1, \dots, k\}$. \square

Lemma 3.1. *Let T be a tree of order n with distance k -domination number $\gamma_k \geq 3$. If $n = (k+1)\gamma_k$, then*

$$\Pi_1(T) \geq 4^{k\gamma_k-1}(\gamma_k)^2 \text{ and } \Pi_2(T) \leq 4^{k\gamma_k-1}(\gamma_k)^{\gamma_k}.$$

Either equality holds if and only if $T \cong T_{n,k,\gamma_k}$.

Proof. By Lemma 2.6, we have $T = R \circ k$ for some tree R on γ_k vertices. For

$w \in V(R)$, $d_R(w) = d_T(w) - 1$. Thus

$$\begin{aligned}\Pi_1(T) &= \prod_{w \in V(R)} d_T^2(w) \prod_{z \in V(T) \setminus V(R)} d_T^2(z) \\ &= \left(\prod_{w \in V(R)} (d_R(w) + 1)^2 \right) (2^2)^{(k-1)\gamma_k} \\ &= 4^{(k-1)\gamma_k} \cdot f^2(R),\end{aligned}$$

where $f(R) = \prod_{w \in V(R)} (d_R(w) + 1)$. By Lemma 2.7, $f^2(R) \geq (2^{\gamma_k-1} \cdot \gamma_k)^2$ with equality if and only if $R \cong S_{\gamma_k}$. Therefore

$$\Pi_1(T) \geq 4^{k\gamma_k-1} (\gamma_k)^2$$

with equality if and only if $T = R \circ k$ with $R \cong S_{\gamma_k}$, i.e., $T \cong T_{n,k,\gamma_k}$.

Similarly,

$$\begin{aligned}\Pi_2(T) &= \prod_{w \in V(R)} (d_T(w))^{d_T(w)} \prod_{z \in V(T) \setminus V(R)} (d_T(z))^{d_T(z)} \\ &= \left(\prod_{w \in V(R)} (d_R(w) + 1)^{d_R(w)+1} \right) 4^{(k-1)\gamma_k} \\ &= 4^{(k-1)\gamma_k} h(T),\end{aligned}$$

where $h(T) = \prod_{w \in V(R)} (d_R(w) + 1)^{d_R(w)+1}$. By Lemma 2.8, $h(T) \leq 4^{\gamma_k-1} (\gamma_k)^{\gamma_k}$ with equality if and only if $R \cong S_{\gamma_k}$. Therefore

$$\Pi_2(T) \leq 4^{k\gamma_k-1} (\gamma_k)^{\gamma_k}$$

with equality if and only if $T = R \circ k$ with $R \cong S_{\gamma_k}$, i.e., $T \cong T_{n,k,\gamma_k}$.

□

Lemma 3.2. *Let T be a tree of order n with $\gamma_k(T) = 3$, then*

$$\Pi_1(T) \geq 4^{3k-1} (n - 3k)^2 \text{ and } \Pi_2(T) \leq 4^{3k-1} (n - 3k)^{n-3k}.$$

Either equality holds if and only if $T \cong T_{n,k,3}$.

Proof. By Lemmas 2.4 and 3.1, we have $n \geq (k+1)\gamma_k$, and the result holds for $n = (k+1)\gamma_k$. We present our proof by induction on n . Suppose that $n > 3(k+1)$ and the result is true for $n-1$.

Let $P = v_0 \dots v_d$ be a diametric path and D be a minimum distance k -dominating set of T . We claim that $d \geq 2k+2$, for otherwise, $\{v_k, v_{k+1}\}$ is a distance k -dominating set, a contradiction. We may choose distance k -dominating set D of cardinality $\gamma_k(G)$ with $\{v_k, v_{d-k}\} \subseteq D$ such that $(\cup_{a=0}^k V(T_a) \setminus \{v_k\}) \cap D = \emptyset$ and $(\cup_{a=d-k}^d V(T_a) \setminus \{v_{d-k}\}) \cap D = \emptyset$.

Let $v_0 = w_1, v_d = w_2, \dots, w_m$ be all the pendent vertices of T and $B_T = \{w_i | 1 \leq i \leq m, \gamma_k(T - w_i) = \gamma_k(T)\}$.

We claim that $|B_T| \geq 1$.

Suppose that $B_T = \emptyset$. Then $\gamma_k(T - w_i) = \gamma_k(T) - 1$ for $1 \leq i \leq m$.

For some $i \in \{1, \dots, k, d-k, \dots, d-1\}$, if $d_T(v_i) \geq 3$, then $V(T_i) \cap \{w_3, \dots, w_m\} \neq \emptyset$. As $\{v_k, v_{d-k}\} \in D$, we have $\gamma_k(T - z) = \gamma_k(T)$ for $z \in V(T_i) \cap \{w_3, \dots, w_m\}$, a contradiction. It follows that $d_T(v_i) = 2$ for $i \in \{1, \dots, k, d-k, \dots, d-1\}$.

As $d_T(v_1) = 2$, we have $\gamma_k(T - v_0) = \gamma_k(T) - 1$. Note that $d_T(v_1, v_{k+1}) = k$ and $(\cup_{a=0}^k V(T_a) \setminus \{v_k\}) \cap D = \emptyset$. Thus $v_{k+1} \in D$. Similarly, $v_{d-k-1} \in D$. If $d > 2k+2$, then $v_k, v_{k+1}, v_{d-k-1}, v_{d-k}$ are all distinct, a contradiction to the fact that $\gamma_k(T) = 3$. Hence $d = 2k+2$ and $D = \{v_k, v_{k+1}, v_{d-k}\}$.

If $d_T(v_{k+1}) = 2$, then $T \cong P_{2k+3}$ and $\{v_k, v_{d-k}\}$ is a distance k -dominating set, a contradiction. Then $d_T(v_{k+1}) \geq 3$ and thus $m \geq 3$. If $m > 3$, then $\gamma_k(T - w_i) = \gamma_k(T)$ for some $i \in \{1, \dots, m\}$, which is impossible. Hence $m = 3$. Thus T_{k+1} is a path with end vertices v_{k+1} and w_3 . By the definition of distance k -domination number, we have $d(v_{k+1}, w_3) = k$. It follows that $|V(T)| = 3(k+1)$, which is contradiction. This proves our Claim.

Now by our claim and Lemma 2.9, we have $|N_T(B_T)| = 1$.

Let w be a pendent vertex such that $\gamma_k(T - w) = \gamma_k(T)$ and z being a unique vertex adjacent to x . Then by lemma 2.5, we have $d_T(z) \leq n - k\gamma_k$.

$$\frac{d_T(z)}{d_T(z) - 1} \geq \frac{n - k\gamma_k}{n - 1 - k\gamma_k}$$

with equality if and only if $d_T(z) = n - k\gamma_k$. Note that

$$\Pi_1(T) = \Pi_1(T - w) \cdot \frac{d_T(z)^2}{(d_T(z) - 1)^2}.$$

By induction hypothesis, we have

$$\begin{aligned} \Pi_1(T) &\geq 4^{k\gamma_k-1}(n-1-k\gamma_k)^2 \cdot \left(\frac{d_T(z)}{d_T(z)-1}\right)^2 \\ &\geq 4^{k\gamma_k-1}(n-k\gamma_k)^2 \end{aligned}$$

with equalities if and only if $T - w \cong T_{n-1,k,\gamma_k}$ and $d_T(z) = \Delta = n - k\gamma_k$, i.e., $T \cong T_{n,k,\gamma_k}$.

Let

$$g(t) = \frac{x^x}{(x-1)^{x-1}}.$$

Obviously, $g'(x) = g(x) \log \frac{x}{x-1} > 0$ for $x > 1$. Thus $g(x)$ is strictly increasing for $x > 1$, implying that $g(d_T(z)) \leq g(n - k\gamma_k)$ with equality if and only if $d_T(z) = n - k\gamma_k$. Similarly as above,

$$\Pi_2(T) = \Pi_2(T - w) \cdot \frac{(d_T(z))^{d_T(z)}}{(d_T(z) - 1)^{d_T(z)-1}} = \Pi_2(T - w)g(d_T(z)),$$

and by induction hypothesis, we have

$$\begin{aligned} \Pi_2(T) &\leq 4^{k\gamma_k-1}(n-1-k\gamma_k)^{n-1-k\gamma_k} g(d_T(z)) \\ &\leq 4^{k\gamma_k-1}(n-1-k\gamma_k)^{n-1-k\gamma_k} g(n-k\gamma_k) \\ &= 4^{k\gamma_k-1}(n-k\gamma_k)^{n-k\gamma_k} \end{aligned}$$

with equalities holds if and only if $T - w \cong T_{n-1,k,\gamma_k}$ and $d_T(z) = \Delta = n - k\gamma_k$, i.e., $T \cong T_{n,k,\gamma_k}$. \square

Let D be a distance k -dominating set of a graph G . Let $N_G^a(v)$ be the set of vertices with distance a from v . A vertex $v \in V(G)$ is called a private k -neighbor of u with respect to D if $\cup_{a=0}^k N_G^a(v) \cap D = \{u\}$, that is $d_G(v, u) \leq k$ and $d_G(v, x) \geq k+1$, for any vertex $x \in D \setminus \{u\}$.

Theorem 3.3. *Let T be a tree of order n with distance k -domination number $\gamma_k \geq 3$. Then*

$$\Pi_1(T) \geq 4^{k\gamma_k-1}(n - k\gamma_k)^2 \text{ and } \Pi_2(T) \leq 4^{k\gamma_k-1}(n - k\gamma_k)^{n-k\gamma_k}.$$

Either equality holds if and only if $T \cong T_{n,k,\gamma_k}$.

Proof. Let $P = v_0 \dots v_d$ be a diametric path of T . Define $B_T = \{w \in V(T) \mid d_T(w) = 1 \text{ and } \gamma_k(T-w) = \gamma_k(T)\}$. We may choose distance k -dominating set D of cardinality γ_k with $\{v_k, v_{d-k}\} \subseteq D$ such that $(\cup_{a=0}^k V(T_a) \setminus \{v_k\}) \cap D = \emptyset$ and $(\cup_{a=d-k}^d V(T_a) \setminus \{v_{d-k}\}) \cap D = \emptyset$.

If $B_T = \emptyset$ then for $i = 0, d$, $\gamma_k(T-v_i) = \gamma_k(T) - 1$. If $B_T \neq \emptyset$, then by Lemma 2.9, $|N_T(B_T)| = 1$. If $v_0, v_d \in B_T$, then since $d - 1 > 1$, we have $\{v_1, v_{d-1}\} \subseteq |N_T(B_T)|$ and thus $|N_T(B_T)| > 1$, a contradiction. Thus, in either case, we may assume that $\gamma_k(T - v_0) = \gamma_k(T) - 1$, and thus $\{v_k, v_{k+1}, v_{d-k}\} \subseteq D$.

For $i = 1, \dots, k$, if some T_i is not a star with center v_i , then applying Lemma 2.2 for a non-pendent edge e of T_i to obtain a tree T_e , we have $\Pi_1(T_e) < \Pi_1(T)$ and $\Pi_2(T_e) > \Pi_2(T)$. Thus, we may assume that T_i is a star with center v_i for all $i = 1, \dots, k$.

If there are at least two vertices, say v_i and v_j with $1 \leq i < j \leq k$, with degree at least 3 in T , then by Lemma 2.3 we may find a tree T' by moving the pendent edges at v_i to v_j or vice versa such that $\Pi_1(T') < \Pi_1(T)$ and $\Pi_2(T') > \Pi_2(T)$. So we may assume that there is at most one vertex among vertices v_1, \dots, v_k with degree at least 3. That is, among vertices v_1, \dots, v_k , either each vertex has degree 2 or exactly one vertex, say v_{i_0} has at least one pendent neighbor, where $1 \leq i_0 \leq k$. Let

$$\begin{aligned} T'' &= T - \{v_{i_0}z \mid z \in N_T(v_{i_0}) \setminus \{v_{i_0-1}, v_{i_0+1}\}\} \\ &\quad + \{v_{k+1}z \mid z \in N_T(v_{i_0}) \setminus \{v_{i_0-1}, v_{i_0+1}\}\}. \end{aligned}$$

Let s is number of pendent edges at v_{i_0} . If $s = 0$, then $T = T''$.

Suppose that $s \geq 1$. Then for $i = 1, \dots, k$, $d_{T''}(v_i) = 2$ and D is minimum distance k -dominating set of T'' . Let $PN_{k,D}(z)$ be the set of all private k -neighbors of z with respect to D in T'' . Then for any $z \in \cup_{a=0}^k N_{T''}^a(v_k) \setminus \{v_0, \dots, v_k\}$, $d_{T''}(z, v_{k+1}) \leq k$. It follows that $D \setminus \{v_k\}$ is a distance k -dominating set of the

tree $T'' - \{v_0, \dots, v_k\}$. Also, $PN_{k,D}(v_{k+1}) \subseteq V(T'') \setminus \{v_0, \dots, v_k\}$. It shows that $D \setminus \{v_k\}$ is a minimum distance k -dominating set of $T'' - \{v_0, \dots, v_k\}$. Thus $\gamma_k(T'' - \{v_0, \dots, v_k\}) = \gamma_k - 1$ and $\gamma_k(T'' - \{v_0, \dots, v_{k-1}\}) = \gamma_k - 1$.

Let $\Gamma = \prod_{u \in V(T) \setminus \{v_{i_0}, v_{k+1}\}} d_T^2(u)$ and $\Phi = \prod_{u \in V(T) \setminus \{v_{i_0}, v_{k+1}\}} (d_T(u))^{d_T(u)}$. Then

$$\begin{aligned} \Pi_1(T) - \Pi_1(T'') &= ((2+s)^2 d_T^2(v_{k+1}) - 4(d_T(v_{k+1}) + s)^2) \Gamma \\ &= s(d_T(v_{k+1}) - 2)(s(d_T(v_{k+1}) + 2) + 4d_T(v_{k+1})) \Gamma \\ &\geq 0, \end{aligned}$$

and thus $\Pi_1(T) \geq \Pi_1(T'')$ with equality if and only if $d_T(v_{k+1}) = 2$. Also

$$\begin{aligned} \Pi_2(T'') - \Pi_2(T) &= \left(4(d_T(v_{k+1}) + s)^{d_T(v_{k+1})+s} - (s+2)^{s+2} (d_T(v_{k+1}))^{d_T(v_{k+1})} \right) \Phi \\ &= F(s) \Phi \\ &\geq 0, \end{aligned}$$

where $F(s) = 4(d_T(v_{k+1}) + s)^{d_T(v_{k+1})+s} - (s+2)^{s+2} (d_T(v_{k+1}))^{d_T(v_{k+1})}$. It is easy to check that $F(s)$ is an increasing function for $s \geq 0$. Thus $\Pi_2(T'') \geq \Pi_2(T)$ with equality if and only if $d_T(v_{k+1}) = 2$.

Now we have shown that $\Pi_1(T) \geq \Pi_1(T'')$ and $\Pi_2(T'') \geq \Pi_2(T)$ with either equality if and only if $s = 0$ (i.e., $T = T''$) or $d_T(v_{k+1}) = 2$.

In the following, we prove that, $\Pi_1(T'') \geq 4^{k\gamma_k-1} (n - k\gamma_k)^2$ and $\Pi_2(T'') \leq 4^{k\gamma_k-1} (n - k\gamma_k)^{n-k\gamma_k}$ with either equality if and only if $T'' \cong T_{n,k,\gamma_k}$.

By Lemma 3.2, the result holds for $n \geq (k+1)\gamma_k$ and $\gamma_k = 3$. Suppose that $\gamma_k \geq 4$, and the result is true for $n \geq (k+1)(\gamma_k - 1)$.

Note that $\gamma_k(T'' - \{v_0, \dots, v_k\}) = \gamma_k - 1$ and $|V(T'' - \{v_0, \dots, v_k\})| = n - k - 1 >$

$(k+1)(\gamma_k-1)$. Then

$$\begin{aligned}
\Pi_1(T'') &= \Pi_1(T'' - \{v_0, \dots, v_k\}) \cdot \left(\frac{d_{T''}(v_{k+1})}{(d_{T''}(v_{k+1}) - 1)} \right)^2 \cdot \prod_{i=0}^k d_{T''}^2(v_i) \\
&\geq \Pi_1(T''_{n-k-1, k, \gamma_k-1}) \cdot \left(\frac{d_{T''}(v_{k+1})}{(d_{T''}(v_{k+1}) - 1)} \right)^2 \cdot 4^k \\
&= 4^{k(\gamma_k-1)-1} \cdot (n-k-1-k(\gamma_k-1))^2 \cdot \left(\frac{d_{T''}(v_{k+1})}{(d_{T''}(v_{k+1}) - 1)} \right)^2 \cdot 4^k \\
&\geq 4^{k\gamma_k-1} \cdot (n-1-k\gamma_k)^2 \cdot \frac{(n-k\gamma_k)^2}{(n-1-k\gamma_k)^2} \\
&= 4^{k\gamma_k-1} (n-k\gamma_k)^2
\end{aligned}$$

with equalities if and only if $T'' - \{v_0, \dots, v_k\} \cong T''_{n-k-1, k, \gamma_k-1}$, and $d_{T''}(v_{k+1}) = \Delta = n - k\gamma_k$. Recall that for $i = 1, \dots, k$, $d_{T''}(v_i) = 2$. Thus $\Pi_1(T'') \geq 4^{k\gamma_k-1} (n - k\gamma_k)^2$ with equality if and only if $T'' \cong T_{n, k, \gamma_k}$.

Similarly, we have

$$\begin{aligned}
\Pi_2(T'') &= \Pi_2(T'' - \{v_0, \dots, v_k\}) \cdot \frac{(d_{T''}(v_{k+1}))^{d_{T''}(v_{k+1})}}{(d_{T''}(v_{k+1}) - 1)^{d_{T''}(v_{k+1})-1}} \cdot \prod_{i=0}^k (d_{T''}(v_i))^{d_{T''}(v_i)} \\
&\leq \Pi_2(T''_{n-k-1, k, \gamma_k-1}) \cdot 4^k \cdot \frac{(d_{T''}(v_{k+1}))^{d_{T''}(v_{k+1})}}{(d_{T''}(v_{k+1}) - 1)^{d_{T''}(v_{k+1})-1}} \\
&= 4^{k(\gamma_k-1)-1} (n-k-1-k(\gamma_k-1))^{n-k-1-k(\gamma_k-1)} \cdot 4^k \\
&\quad \cdot \frac{(d_{T''}(v_{k+1}))^{d_{T''}(v_{k+1})}}{(d_{T''}(v_{k+1}) - 1)^{d_{T''}(v_{k+1})-1}} \\
&\leq 4^{k\gamma_k-1} (n-k\gamma_k)^{n-k\gamma_k}
\end{aligned}$$

with equalities if and only if $T'' - \{v_0, \dots, v_k\} \cong T''_{n-k-1, k, \gamma_k-1}$, and $d_{T''}(v_{k+1}) = \Delta = n - k\gamma_k$. Also for $i = 1, \dots, k$, $d_{T''}(v_i) = 2$. Thus $\Pi_2(T'') \leq 4^{k\gamma_k-1} (n - k\gamma_k)^{n-k\gamma_k}$ with equality if and only if $T \cong T_{n, k, \gamma_k}$.

Now we conclude that $\Pi_1(T) \geq \Pi_1(T'') \geq 4^{k\gamma_k-1} (n - k\gamma_k)^2$ with equality in the first inequality if and only if $T = T''$ or $T \not\cong T''$ and $d_T(v_{k+1}) = 2$, and with equality in the second inequality if and only if $T'' \cong T_{n, k, \gamma_k}$. We show that if $T \not\cong T''$ and $d_T(v_{k+1}) = 2$, then $T'' \not\cong T_{n, k, \gamma_k}$. Otherwise, say $T'' = T_{n, k, \gamma_k}$. As $d_{T''}(v_{k+1}) = n - k\gamma_k$, there are $n - (k+1)\gamma_k$ pendent edges at v_{k+1} in T'' . By the above

argument, $d_T(v_{k+1}) = \gamma_k > 2$, a contradiction. Therefore $\Pi_1(T) \geq 4^{k\gamma_k-1}(n - k\gamma_k)^2$ with equality if and only if $T \cong T_{n,k,\gamma_k}$. Similarly, $\Pi_2(T) \leq 4^{k\gamma_k-1}(n - k\gamma_k)^{n-k\gamma_k}$ with equality if and only if $T \cong T_{n,k,\gamma_k}$. \square

References

- [1] B. Borovicanin, B. Furtula, On extremal Zagreb indices of trees with given domination number, *Appl. Math. Comput* 279 (2016) 208–218.
- [2] I. Gutman, On the origin of two degree based topological indices, *Bull. Acad. Serbe Sci. Arts. (Cl. Sci. Math.)* 146 (2014) 39-52.
- [3] I. Gutman, Multiplicative Zagreb indices of trees. *Bull, Soc. Math. Banja Luka* 18 (2011)17-23.
- [4] I. Gutman, B. Furtula (Eds.), *Novel Molecular Structure Descriptors Theory and Applications*, University Kragujevac, Kragujevac, 2010, pp. 72-100.
- [5] I. Gutman, N. Trinajstić, Graph theory and molecular orbitals Total π -electron energy of alternant hydrocarbons, *Chem. Phys. Lett.* 17(1972) 535–538.
- [6] S.M. Hosamani, I. Gutman, Zagreb indices of transformation graphs and total transformation graphs, *Appl. Math. Comput* 247 (2014) 1156-1160.
- [7] Y. Hu, X. Li, Y. Shi, T. Xu, I. Gutman, On molecular graphs with smallest and greatest zeroth order general Randi index, *MATCH Commun. Math. Comput. Chem* 54 (2005) 425-434.
- [8] A. Iranmanesh, M.A. Hosseinzadeh, I. Gutman, On multiplicative Zagreb indices of graphs, *Iran. J. Math. Chem.* 3 (2012) 145–154.
- [9] R. Kazemi, Note on the multiplicative Zagreb indices, *Discrete Appl. Math.* 198 (2016) 147–154.

- [10] J. Liu, Q. Zhang, Sharp upper bounds for multiplicative Zagreb indices, *MATCH Commun. Math. Comput. Chem.* 68 (2012) 231–240.
- [11] A. Meir, J.M. Moon, Relations between packing and covering numbers of a tree, *Pac. J. Math.* 61(1975) 225-233.
- [12] L. Pei, X. Pan, Extremal values on Zagreb indices of trees with given distance k -domination number, *J. Inequal. Appl.* (2018) Paper No. 16, 17 pp.
- [13] R. Todeschini, V. Consonni, New local vertex invariants and molecular descriptors based on functions of the vertex degrees, *MATCH Commun. Math. Comput. Chem* 64 (2010) 359-372.
- [14] J. Topp. L. Volkmann, On packing and covering numbers of graphs, *Discrete Math.* 96 (1991) 229-238.
- [15] F. Wang, F. Belardo, A lower bound for the first Zagreb index and its application, *MATCH Commun. Math. Comput. Chem.* 74 (2015) 35-56.
- [16] S. Wang, B. Wei, Multiplicative Zagreb indices of k -trees, *Discrete Appl. Math.* 180 (2015) 168–175.
- [17] S. Wang, C. Wang, L. Chen, J.B. Liu, On extremal multiplicative Zagreb indices of trees with given number of vertices of maximum degree, *Discrete Appl. Math.* 227 (2017) 166–173.
- [18] S. Wang, C. Wang, J.-B. Liu, On extremal multiplicative Zagreb indices of trees with given domination number, *Appl. Math. Comput* 332 (2018) 338–350.
- [19] K. Xu, H. Hua, A unified approach to extremal multiplicative Zagreb indices for trees, unicyclic and bicyclic graphs, *MATCH Commun. Math. Comput. Chem.* 68 (2012) 241-256.
- [20] B. Zhou, Remarks on Zagreb indices, *MATCH Commun. Math. Comput. Chem.* 57 (2007) 591-596.

- [21] B. Zhou, I. Gutman, Further properties of Zagreb indices, MATCH Commun. Math. Comput. Chem. 54 (2005) 233-239.