

Centralizers of elements of infinite order in plane Cremona groups

ShengYuan Zhao

Abstract

Let \mathbf{K} be an algebraically closed field. The Cremona group $\mathrm{Cr}_2(\mathbf{K})$ is the group of birational transformations of the projective plane $\mathbb{P}_{\mathbf{K}}^2$. We carry out an overall study of centralizers of elements of infinite order in $\mathrm{Cr}_2(\mathbf{K})$ which leads to a classification of embeddings of \mathbf{Z}^2 into $\mathrm{Cr}_2(\mathbf{K})$, as well as a classification of maximal non-torsion abelian subgroups of $\mathrm{Cr}_2(\mathbf{K})$ when $\mathrm{char}(\mathbf{K}) = 0$.

Mathematics Subject Classification 2010: 14E07

Contents

1	Introduction	1
2	Elements which are not base-wandering Jonquières twists	6
2.1	Loxodromic elements	6
2.2	Halphen twists	7
2.3	Elliptic elements of infinite order	8
2.4	Jonquières twists with trivial action on the base	11
3	Base-wandering Jonquières twists	13
3.1	Algebraically stable maps	14
3.2	Elliptic elements in the Jonquières group	18
3.3	The group $\mathrm{Cent}_0(f)$	21
3.4	Persistent indeterminacy points	23
3.5	Local analysis around a fiber	26
4	Proofs of the main results	34
5	References	39

1 Introduction

Let \mathbf{K} be an algebraically closed field. The *plane Cremona group* $\mathrm{Cr}_2(\mathbf{K})$ is the group of birational transformations of the projective plane $\mathbb{P}_{\mathbf{K}}^2$. It is isomorphic to the group of \mathbf{K} -algebra automorphisms of $\mathbf{K}(X_1, X_2)$, the function field of $\mathbb{P}_{\mathbf{K}}^2$. Using a system

of homogeneous coordinates $[x_0; x_1; x_2]$, a birational transformation $f \in \text{Cr}_2(\mathbf{K})$ can be written as

$$[x_0 : x_1 : x_2] \mapsto [f_0(x_0, x_1, x_2) : f_1(x_0, x_1, x_2) : f_2(x_0, x_1, x_2)]$$

where f_0, f_1, f_2 are homogeneous polynomials of the same degree without common factor. This degree does not depend on the system of homogeneous coordinates. We call it the *degree* of f and denote it by $\deg(f)$. Geometrically it is the degree of the pull-back by f of a general projective line. Birational transformations of degree 1 are homographies and form $\text{Aut}(\mathbb{P}_{\mathbf{K}}^2) = \text{PGL}_3(\mathbf{K})$, the group of automorphisms of the projective plane.

Four types of elements. Following the work of M. Gizatullin, S. Cantat, J. Diller and C. Favre, we can classify an element $f \in \text{Cr}_2(\mathbf{K})$ into exactly one of the four following types according to the growth of the sequence $(\deg(f^n))_{n \in \mathbf{N}}$ (See [DF01] Theorems 0.2, 0.3):

1. The sequence $(\deg(f^n))_{n \in \mathbf{N}}$ is bounded, f is birationally conjugate to an automorphism f' of a rational projective surface X and a positive iterate of f' lies in the connected component of the identity of the automorphism group $\text{Aut}(X)$. We call f an *elliptic element*.
2. The sequence $(\deg(f^n))_{n \in \mathbf{N}}$ grows linearly, f preserves a unique pencil of rational curves and f is not conjugate to an automorphism of any rational projective surface. We call f a *Jonquière's twist*.
3. The sequence $(\deg(f^n))_{n \in \mathbf{N}}$ grows quadratically, f is conjugate to an automorphism of a rational projective surface preserving a unique elliptic fibration. We call f a *Halphen twist*.
4. The sequence $(\deg(f^n))_{n \in \mathbf{N}}$ grows exponentially and f is called *loxodromic*.

If every element of a subgroup $G \subset \text{Cr}_2(\mathbf{K})$ is elliptic and if G is conjugate to a group of automorphisms of a projective rational surface, then we call G an *elliptic group*.

The standard reference [DF01] is written for $\mathbf{K} = \mathbf{C}$ but the same proof works over an algebraically closed field \mathbf{K} of characteristic different from 2 and 3. The only problem with characteristics 2 and 3 is that the important ingredient [Giz80] does not deal with quasi-elliptic fibrations. This minor issue has been clarified in [CD12a] and [CGL] so that the above classification holds in arbitrary characteristic (cf. [CD12a] Section 4.1 and [CGL] Sections 4.3, 4.4).

The Jonquière's group Fix an affine chart of \mathbb{P}^2 with coordinates (x, y) . The *Jonquière's group* $\text{Jonq}(\mathbf{K})$ is the subgroup of the Cremona group of all transformations of the form

$$(x, y) \mapsto \left(\frac{ax + b}{cx + d}, \frac{A(x)y + B(x)}{C(x)y + D(x)} \right) \quad (1)$$

where

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{PGL}_2(\mathbf{K}), \quad \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \text{PGL}_2(\mathbf{K}(x)).$$

In other words, $\text{Jonq}(\mathbf{K})$ is the group of all birational transformations of $\mathbb{P}^1 \times \mathbb{P}^1$ permuting the fibers of the projection onto the first factor; it is isomorphic to the semi-direct product $\text{PGL}_2(\mathbf{K}) \ltimes \text{PGL}_2(\mathbf{K}(x))$. The Jonquières group is defined only if a system of coordinates is chosen; a different choice of the affine chart yields a conjugation by an element of $\text{PGL}_3(\mathbf{K})$. We can directly check with Formula (1) that the sequence $(\deg(f^n))_{n \in \mathbf{N}}$ grows at most linearly. Thus elements of $\text{Jonq}(\mathbf{K})$ are either elliptic or Jonquières twists. We denote by $\text{Jonq}_0(\mathbf{K})$ the normal subgroup of $\text{Jonq}(\mathbf{K})$ that preserves fiberwise the rational fibration, i.e. the subgroup of those transformations of the form $(x, y) \mapsto \left(x, \frac{A(x)y+B(x)}{C(x)y+D(x)}\right)$; it is isomorphic to $\text{PGL}_2(\mathbf{K}(x))$. A Jonquières twist of the Jonquières group will be called a *base-wandering Jonquières twist* if its action on the base of the rational fibration has infinite order.

If $\mathbf{K} = \overline{\mathbf{F}_p}$ is the algebraic closure of a finite field, then \mathbf{K}, \mathbf{K}^* and $\text{PGL}_2(\mathbf{K})$ are all torsion groups. Thus, if $\mathbf{K} = \overline{\mathbf{F}_p}$ then base-wandering Jonquières twists do not exist. When $\text{char}(\mathbf{K}) = 0$, or when $\text{char}(\mathbf{K}) = p > 0$ and $\mathbf{K} \neq \overline{\mathbf{F}_p}$, there exist base-wandering Jonquières twists.

The group of automorphisms of a Hirzebruch surface will be systematically considered as a subgroup of the Jonquières group in the following way:

$$\text{Aut}(\mathbb{F}_n) = \left\{ (x, y) \mapsto \left(\frac{ax+b}{cx+d}, \frac{y+t_0+t_1x+\dots+t_nx^n}{(cx+d)^n} \right) \mid \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{GL}_2(\mathbf{K}), t_0, \dots, t_n \in \mathbf{K} \right\}. \quad (2)$$

Main results.

Theorem 1.1 *Let Γ be a subgroup of $\text{Cr}_2(\mathbf{K})$ which is isomorphic to \mathbf{Z}^2 . Then there are a projective rational surface X and a birational map $\phi : \mathbb{P}^2 \dashrightarrow X$ such that $\Gamma' = \phi\Gamma\phi^{-1}$ has a pair of generators (f, g) that fits in one of the following (mutually exclusive) situations:*

1. f, g are elliptic elements, $\Gamma' \subset \text{Aut}(X)$.
2. f, g are Halphen twists preserving a same elliptic fibration on X , and $\Gamma' \subset \text{Aut}(X)$.
3. one or both of the f, g are Jonquières twists, and there exist $m, n \in \mathbf{N}^*$ such that the finite index subgroup of Γ' generated by f^m and g^n is in an 1-dimensional torus over $\mathbf{K}(x)$ in $\text{Jonq}_0(\mathbf{K}) = \text{PGL}_2(\mathbf{K}(x))$;
4. f is a base-wandering Jonquières twist and g is elliptic. In some affine chart, we can write f, g in one of the following forms:
 - g is $(x, y) \mapsto (\alpha x, \beta y)$ and f is $(x, y) \mapsto (\eta(x), yR(x^k))$ where $\alpha, \beta \in \mathbf{K}^*$, $\alpha^k = 1, R \in \mathbf{K}(x), \eta \in \text{PGL}_2(\mathbf{K}), \eta(\alpha x) = \alpha\eta(x)$ and η has infinite order;
 - (only when $\text{char}(\mathbf{K}) = 0$) g is $(x, y) \mapsto (\alpha x, y+1)$ and f is $(x, y) \mapsto (\eta(x), y+R(x))$ where $\alpha \in \mathbf{K}^*, R \in \mathbf{K}(x), R(\alpha x) = R(x), \eta \in \text{PGL}_2(\mathbf{K}), \eta(\alpha x) = \alpha\eta(x)$ and η has infinite order.

Remark 1.2 When \mathbf{K} is the algebraic closure of a finite field, the above list can be shortened since there is no elliptic elements of infinite order nor base-wandering Jonquières twists.

Remark 1.3 From Theorem 1.1 it is not difficult to see that (see Section 4 for a proof), when Γ is isomorphic to \mathbf{Z}^2 , the degree function $deg : \Gamma \rightarrow \mathbf{N}$ is governed by the word length function with respect to some generators in the following sense. In the first case of the above theorem it is bounded. In the second case it is up to a bounded term a positive definite quadratic form over \mathbf{Z}^2 . In the third case the degree of $f^i \circ g^j$ is dominated by $deg(f)i + deg(g)j$ if i, j are positive. In the fourth case the degree function is up to a bounded term $f^i \circ g^j \mapsto c|i|$ for some $c \in \mathbf{Q}_+$.

Theorem 1.1 is based on several known results. The main new feature is the following result (see Theorem 3.2 for a more precise reformulation).

Corollary 1.4 *Let $G \subset \text{Jonq}(\mathbf{K})$ be a subgroup isomorphic to \mathbf{Z}^2 . Suppose that the action of G on the base of the rational fibration is faithful. Then G is an elliptic subgroup.*

We state it as a corollary (in cases 2, 3, 4 of Theorem 1.1 the action on the base is not faithful) but we will see that it is rather an intermediate step to prove Theorem 1.1.

A *maximal abelian subgroup* is an abelian subgroup which is not strictly contained in any other abelian subgroup. Over the field of complex numbers, finite abelian subgroups of $\text{Cr}_2(\mathbf{C})$ have been classified in [Bla07] and maximal uncountable abelian subgroups of $\text{Cr}_2(\mathbf{C})$ have been studied in [Dés06b] (Theorems 1.5, 1.6). When $\text{char}(\mathbf{K}) = 0$ we will use Theorem 1.1 to classify maximal abelian subgroups of $\text{Cr}_2(\mathbf{K})$ which contain at least one element of infinite order, see Theorem 4.1. Our classification is more precise than the results in [Dés06b].

Another theorem we obtain from Theorem 1.1 is the following:

Theorem 1.5 *Assume $\text{char}(\mathbf{K}) = 0$. Let $f \in \text{Cr}_2(\mathbf{K})$ be an element of infinite order. If the centralizer of f is not virtually abelian, then f is an elliptic element and a power of f is conjugate to an automorphism of \mathbb{A}^2 of the form $(x, y) \mapsto (x, y + 1)$ or $(x, y) \mapsto (x, \beta y)$ with $\beta \in \mathbf{K}^*$.*

Previously known results. Let us begin with the group of polynomial automorphism of the affine plane $\text{Aut}(\mathbb{A}^2)$. It can be seen as a subgroup of $\text{Cr}_2(\mathbf{K})$. It is the amalgamated product of the group of affine automorphisms with the so called *elementary group*

$$\text{El}(\mathbf{K}) = \{(x, y) \mapsto (\alpha x + \beta, \gamma y + P(x)) \mid \alpha, \beta, \gamma \in \mathbf{K}, \alpha\beta \neq 0, P \in \mathbf{K}[x]\}.$$

Let \mathbf{K} be the field of complex numbers. S. Friedland and J. Milnor showed in [FM89] that an element of $\text{Aut}(\mathbb{A}_{\mathbf{C}}^2)$ is either conjugate to an element of $\text{El}(\mathbf{C})$ or to a generalized Hénon map, i.e. a composition $f_1 \circ \dots \circ f_n$ where the f_i are Hénon maps of the form $(x, y) \mapsto (y, P_i(y) - \delta_i x)$ with $\delta_i \in \mathbf{C}^*$, $P_i \in \mathbf{C}[y]$, $deg(P_i) \geq 2$. S. Lamy and C. Bisi showed in [Lam01] and [Bis04] that the centralizer in $\text{Aut}(\mathbf{C}^2)$ of a generalized Hénon map is finite by cyclic, and that of an element of $\text{El}(\mathbf{C})$ is uncountable (see also [Bis08])

for partial extensions to higher dimension). Note that, when viewed as elements of $\text{Cr}_2(\mathbf{C})$, a generalized Hénon map is loxodromic and an element of $\text{El}(\mathbf{C})$ is elliptic.

As regards the Cremona group, centralizers of loxodromic elements are known to be finite by cyclic (S. Cantat [Can11] Theorem 5.1, J. Blanc-S. Cantat [BC16] Corollary 4.7). Centralizers of Halphen twists are virtually abelian of rank at most 8 (M. Gizatullin [Giz80] Proposition 7, S. Cantat [Can11] Proposition 5.2). When \mathbf{K} is the field of complex numbers, centralizers of elliptic elements of infinite order are completely described by J. Blanc-J. Déserti in [BD15] Lemmas 2.7, 2.8 and centralizers of Jonquières twists in $\text{Jonq}_0(\mathbf{K})$ are completely described by D. Cerveau-J. Déserti in [CD12b]. Centralizers of base-wandering Jonquières twists are also studied in [CD12b] but they were not fully understood, for example the results in loc. cit. are not sufficient for classifying pairs of Jonquières twists generating a copy of \mathbf{Z}^2 . Thus, in order to obtain a classification of embeddings of \mathbf{Z}^2 in $\text{Cr}_2(\mathbf{K})$, we need a detailed study of centralizers of base-wandering Jonquières twists, which is the main task of this article. Regarding the elements of finite order and their centralizers in $\text{Cr}_2(\mathbf{K})$, the problem is of a rather different flavour and we refer the readers to [Bla07], [DI09], [Ser10], [Ure] and the references therein.

Remark 1.6 There is a topology on $\text{Cr}_2(\mathbf{K})$, called Zariski topology, which is introduced by M. Demazure and J-P. Serre in [Dem70] and [Ser10]. Note that the Zariski topology does not make $\text{Cr}_2(\mathbf{K})$ an infinite dimensional algebraic group (cf. [BF13]). With respect to the Zariski topology, the centralizer of any element of $\text{Cr}_2(\mathbf{K})$ is closed (J-P. Serre [Ser10]). When K is a local field, J. Blanc and J-P. Furter construct in [BF13] an Euclidean topology on $\text{Cr}_2(\mathbf{K})$ which when restricted to $\text{PGL}_3(\mathbf{K})$ coincides with the Euclidean topology of $\text{PGL}_3(\mathbf{K})$; centralizers are also closed with respect to the Euclidean topology. In particular the intersection of the centralizer of an element in $\text{Cr}_2(\mathbf{K})$ with an algebraic subgroup G of $\text{Cr}_2(\mathbf{K})$ is a closed subgroup of G , with respect to the Zariski topology of G (and with respect to the Euclidean topology when the later is present).

Comparison with other results. S. Smale asked in the '60s if, in the group of diffeomorphisms of a compact manifold, the centralizer of a generic diffeomorphism consists only of its iterates. There has been a lot of work on this question, see for example [BCW09] for an affirmative answer in the C^1 case. Similar phenomena also appear in the group of germs of 1-dimensional holomorphic diffeomorphisms at $0 \in \mathbf{C}$ ([É81]). See the introduction of [CD12b] for more references in this direction. With regard to $\text{Cr}_2(\mathbf{K})$, it is known that loxodromic elements form a Zariski dense subset of $\text{Cr}_2(\mathbf{K})$ (cf. [Xie15], [BD05]) and that their centralizers coincide with the cyclic group formed by their iterates up to finite index (cf. [BC16]). Centralizers of general Jonquières twists are also finite by cyclic (Remark 3.5).

One may compare our classification of \mathbf{Z}^2 in $\text{Cr}_2(\mathbf{K})$ to the following two theorems where the situations are more rigid. The first can be seen as a continuous counterpart and is proved by F. Enriques [Enr93] and M. Demazure [Dem70], the second can be seen as a torsion counterpart and is proved by A. Beauville [Bea07]:

1. *If \mathbf{K}^{*r} embeds as an algebraic subgroup into $\text{Cr}_2(\mathbf{K})$, then $r \leq 2$; if $r = 2$ then*

the embedding is conjugate to an embedding into the group of diagonal matrices Δ in $\mathrm{PGL}_3(\mathbf{K})$.

2. If $p \geq 5$ is a prime number different from the characteristic of \mathbf{K} and if $(\mathbf{Z}/p\mathbf{Z})^r$ embeds into $\mathrm{Cr}_2(\mathbf{K})$, then $r \leq 2$; if $r = 2$ then the embedding is conjugate to an embedding into the group of diagonal matrices Δ in $\mathrm{PGL}_3(\mathbf{K})$.

The classification of \mathbf{Z}^2 in $\mathrm{Cr}_2(\mathbf{K})$ is a very natural special case of the study of finitely generated subgroups of $\mathrm{Cr}_2(\mathbf{K})$; and information on centralizers can be useful for studying homomorphisms from other groups into $\mathrm{Cr}_2(\mathbf{K})$, see for example [Dés06a]. We refer the reader to the surveys [Fav10],[Can18] for representations of finitely generated groups into $\mathrm{Cr}_2(\mathbf{K})$ and [CX18] for general results in higher dimension.

Acknowledgement. I would like to address my warmest thanks to my supervisor Serge Cantat for initiating me into Cremona groups, for numerous discussions, for his constant support and for encouraging me to write this paper. I would also like to thank JunYi Xie for helpful discussions on related topics ranging from proof details to general background. Many thanks to the anonymous referee for her/his careful reading and for her/his comments that improve largely the exposition of the paper.

2 Elements which are not base-wandering Jonquières twists

This section contains a quick review of some scattered results about centralizers from [Can11],[BD15],[CD12b],[BC16]. Some of the proofs are reproduced, because the original proofs were written over \mathbf{C} on the one hand, and because we will need some by-products of the proofs on the other hand.

2.1 Loxodromic elements

Theorem 2.1 ([BC16] Corollary 4.7) *Let $f \in \mathrm{Cr}_2(\mathbf{K})$ be a loxodromic element. The infinite cyclic group generated by f is a finite index subgroup of the centralizer of f in $\mathrm{Cr}_2(\mathbf{K})$.*

Proof We provide a proof which is simpler than [BC16]. See [Can11] and [Can18] for the technical tools used in this proof. The Cremona group $\mathrm{Cr}_2(\mathbf{K})$ acts faithfully by isometries on an infinite dimensional hyperbolic space \mathbb{H} and the action of a loxodromic element is loxodromic in the sense of hyperbolic geometry. In particular there is a unique f -invariant geodesic $Ax(f)$ on which f acts by translation and the translation length is $\log(\lim_{n \rightarrow \infty} \deg(f^n)^{1/n})$. The centralizer $\mathrm{Cent}(f)$ preserves $Ax(f)$ and by considering translation lengths we get a morphism $\phi : \mathrm{Cent}(f) \rightarrow \mathbf{R}$. We claim that the image of ϕ is discrete thus cyclic. Let us see first how the conclusion follows from the claim. Let $x \in \mathbb{H}$ be a point which corresponds to an ample class and let y be an arbitrary point on $Ax(f)$. An element of the kernel $\mathrm{Ker}(\phi)$ fixes a point in $Ax(f)$ and thus fixes

$Ax(f)$ pointwise because it commutes with f . Therefore for any element g of $\text{Ker}(\phi)$ the distance $d(x, g(x))$ is bounded by $2d(x, y)$. This implies that $\text{Ker}(\phi)$ is a subgroup of $\text{Cr}_2(\mathbf{K})$ of bounded degree. If $\text{Ker}(\phi)$ were infinite then its Zariski closure G in $\text{Cr}_2(\mathbf{K})$ would be an algebraic subgroup of strictly positive dimension contained, after conjugation, in the automorphism group of a rational surface. As $\text{Cent}(f)$ is Zariski closed, the elements of G commute with f . The orbits of a one-parameter subgroup of G would form an f -invariant pencil of curves. This contradicts the fact that f is loxodromic. Consequently $\text{Ker}(\phi)$ is finite and hence $\text{Cent}(f)$ is finite by cyclic.

Now let us prove the claim that the image of ϕ is discrete. This follows directly from a spectral gap property for translation lengths of loxodromic elements proved in [BC16]. We give here an easier direct proof found with S. Cantat. Suppose by contradiction that there is a sequence $(g_n)_n$ of distinct elements of $\text{Cent}(f)$ whose translation lengths on $Ax(f)$ tend to 0 when n goes to infinity. Without loss of generality, we can suppose the existence of a point y on $Ax(f)$ and a real number $\varepsilon > 0$ such that $\forall n, d(y, g_n(y)) < \varepsilon$. Let $x \in \mathbb{H}$ be an element which corresponds to an ample class. Then it follows that

$$\forall n, d(x, g_n(x)) \leq d(x, y) + d(y, g_n(y)) + d(g_n(y), g_n(x)) < 2d(x, y) + \varepsilon =: d,$$

i.e. the sequence $(g_n)_n$ has bounded degree d . Up to extracting a subsequence, we can assume that all the g_n has the same degree k with $1 < k \leq d$. Elements of degree k of the Cremona group form a quasi-projective variety $\text{Cr}_2^k(\mathbf{K})$. JunYi Xie proved in [Xie15] that for any $0 < \lambda < \log(k)$, the loxodromic elements of $\text{Cr}_2^k(\mathbf{K})$ whose translation lengths are greater than λ form a Zariski open dense subset of $\text{Cr}_2^k(\mathbf{K})$. Thus the g_n give rise to a strictly ascending chain of Zariski open subsets of $\text{Cr}_2^k(\mathbf{K})$, contradicting the noetherian property of Zariski topology. This finishes the proof. Note that [Xie15] is also used to prove the spectral gap property in [BC16]. \square

2.2 Halphen twists

We only recall here the final arguments of the proofs.

Theorem 2.2 ([Giz80] and [Can11] Proposition 4.7) *Let $f \in \text{Cr}_2(\mathbf{K})$ be a Halphen twist. The centralizer $\text{Cent}(f)$ of f in $\text{Cr}_2(\mathbf{K})$ contains a finite index abelian subgroup of rank less than or equal to 8.*

Proof Being a Halphen twist, the birational transformation f is up to conjugation an automorphism of a projective rational surface and preserves a relatively minimal elliptic fibration. An iterate of f preserves any fiber of this elliptic fibration (cf. [Giz80] Proposition 7). Thus the f -invariant fibration is unique. As a consequence $\text{Cent}(f)$ acts by automorphisms preserving this fibration. It is proved in [Giz80] (see [CGL] Sections 4.3, 4.4 for a clarification in characteristics 2 and 3) that the automorphism group of a rational minimal elliptic surface has a finite index abelian subgroup of rank less than 8. \square

2.3 Elliptic elements of infinite order

In this section we reproduce a part of [BD15]; we follow the original proofs (for $\text{char}(\mathbf{K}) = 0$) in loc. cit. and some extra details are added in case $\text{char}(\mathbf{K}) > 0$.

The proof of the following proposition in [BD15] translates word by word to the case of positive characteristic. It is based on a G -Mori-program for rational surfaces due to J. Manin [Man67] and V. Iskovskih [Isk79].

Proposition 2.3 ([BD15] Proposition 2.1) *Let S be a smooth rational surface over \mathbf{K} . Let $f \in \text{Aut}(S)$ be an automorphism of infinite order whose action on $\text{Pic}(S)$ is of finite order. Then there exists a birational morphism $S \rightarrow X$ where X is a Hirzebruch surface \mathbb{F}_n ($n \neq 1$) or the projective plane \mathbb{P}^2 , which conjugates f to an automorphism of X .*

Proposition 2.4 ([BD15] Proposition 2.3) *Let $f \in \text{Cr}_2(\mathbf{K})$ be an elliptic element of infinite order. Then f is conjugate to an automorphism of \mathbb{P}^2 . Furthermore there exists an affine chart with affine coordinates (x, y) on which f acts by automorphism of the following form:*

1. $(x, y) \mapsto (\alpha x, \beta y)$ where $\alpha, \beta \in \mathbf{K}^*$ are such that the kernel of the group homomorphism $\mathbf{Z}^2 \rightarrow \mathbf{K}^*, (i, j) \mapsto \alpha^i \beta^j$ is generated by $(k, 0)$ for some $k \in \mathbf{Z}$;
2. $(x, y) \mapsto (\alpha x, y + 1)$ where $\alpha \in \mathbf{K}^*$ and α is of infinite order if $\text{char}(\mathbf{K}) > 0$.

Remark 2.5 If $\mathbf{K} = \overline{\mathbb{F}_p}$ then every elliptic element is of finite order.

Proof (of Proposition 2.4) By Proposition 2.3 we can suppose that f is an automorphism of \mathbb{P}^2 or of a Hirzebruch surface.

Let's consider first the case when $f \in \text{Aut}(\mathbb{P}^2) = \text{PGL}_3(\mathbf{K})$. By putting the corresponding matrix in Jordan normal form, we can find an affine chart on which f has, up to conjugation, one of the following forms: 1) $(x, y) \mapsto (\alpha x, \beta y)$; 2) $(x, y) \mapsto (\alpha x, y + 1)$; 3) $(x, y) \mapsto (x + y, y + 1)$. If $\text{char}(\mathbf{K}) > 0$ then f can not have the third form since it would have finite order; if $\text{char}(\mathbf{K}) = 0$ then in the third case f is conjugate by $(x, y) \mapsto (x - \frac{1}{2}y(y-1), y)$ to $(x, y) \mapsto (x, y + 1)$. We now show that in the first case α, β can be chosen to verify the condition in the proposition. Let $\phi : (x, y) \mapsto (\alpha x, \beta y)$ be a diagonal automorphism, we denote by $\Delta(\phi)$ the kernel of the group morphism $\mathbf{Z}^2 \rightarrow \mathbf{K}^*, (i, j) \mapsto \alpha^i \beta^j$. For $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{GL}_2(\mathbf{Z})$, we denote by $M(\phi)$ the diagonal automorphism $(x, y) \mapsto (\alpha^a \beta^b x, \alpha^c \beta^d y)$, i.e. the conjugate of ϕ by the monomial map $(x, y) \mapsto (x^a y^b, x^c y^d)$. We have the relation $\Delta(M(\phi)) = (M^T)^{-1}(\Delta(\phi))$. Using the Smith normal form of a matrix with integer entries, up to conjugation by a monomial map we can suppose that our elliptic element f satisfies $\Delta(f) = \langle (k_1, 0), (0, k_1 k_2) \rangle$ where $k_1, k_2 \in \mathbf{Z}$. Since f is of infinite order, $k_1 k_2$ must be 0.

If $f \in \text{Aut}(\mathbb{F}_0) = \text{Aut}(\mathbb{P}^1 \times \mathbb{P}^1)$, then we reduce to the case of \mathbb{P}^2 by blowing up a fixed point and contracting the strict transforms of the two rulings passing through the point. If $f \in \text{Aut}(\mathbb{F}_n)$ for $n \geq 2$ and if f has a fixed point which is not on the exceptional section, then we can reduce to \mathbb{F}_{n-1} by making an elementary transformation at the fixed point.

Suppose now that $f \in \text{Aut}(\mathbb{F}_n)$, $n \geq 2$ and its fixed points are all on the exceptional section. By removing the exceptional section and an invariant fiber of the rational fibration, we get an open subset isomorphic to \mathbb{A}^2 on which f can be written as: $(x, y) \mapsto (\alpha x, \beta y + Q(x))$ or $(x, y) \mapsto (x + 1, \beta y + Q(x))$ where $\alpha, \beta \in \mathbf{K}^*$ and Q is a polynomial of degree $\leq n$.

In the first case, the fact that there is no extra fixed point on the fiber $x = 0$ implies $\beta = 1$ and $Q(0) \neq 0$. The action on the fiber at infinity can be obtained by a change of variables $(x', y') = (1/x, y/x^n)$, so the fact that there is no extra fixed point on it implies $\beta = \alpha^n$ and $\deg(Q) = n$. This forces α to be a primitive r -th root of unity for some $r \in \mathbf{N}$. Conjugating f by $(x, y) \mapsto (x, y + \gamma x^d)$, we replace $Q(x)$ with $Q(x) + \gamma(\alpha^d - 1)x^d$. This allows us to eliminate the term x^d of Q unless $\alpha^d = 1$. So we can assume that f has the form $(x, y) \mapsto (\alpha x, y + \tilde{Q}(x^r))$ where $\alpha^r = 1$ and $\tilde{Q} \in \mathbf{K}[x]$. Then f is conjugate to $(x, y) \mapsto (\alpha x, y + 1)$ by $(x, y) \mapsto (x, y + \tilde{Q}(x^r))$. Remark that this case does not happen in positive characteristic because an automorphism of this form would have finite order. Note that in this paragraph we did not use the fact that f is of infinite order, so that Proposition 2.6 is proved.

Suppose now we are in the second case. There is no extra fixed point if and only if $\beta = 1$ and $\deg(Q) = n$. If $\text{char}(\mathbf{K}) > 0$ and if $\beta = 1$, then f would be of finite order. Therefore we can assume $\text{char}(\mathbf{K}) = 0$. In that case, we can decrease the degree of Q by conjugating f by a well chosen birational transformation of the form $(x, y) \mapsto (x, y + \gamma x^{n+1})$ with $\gamma \in \mathbf{K}^*$. By induction we get $(x, y) \mapsto (x + 1, y)$ at last. \square

Applying the proof of Proposition 2.4 to an automorphism preserving the fibration fiber by fiber, we get the following:

Proposition 2.6 *Let f be an automorphism of a Hirzebruch surface which preserves the rational fibration fiber by fiber (we do not assume that f is of infinite order). Then there exists an affine chart on which f acts as an automorphism of the following form:*

1. $(x, y) \mapsto (x, \beta y)$ where $\beta \in \mathbf{K}^*$;
2. $(x, y) \mapsto (x, y + 1)$.

Here x is the coordinate on the base of the rational fibration.

Once we have the above normal forms, explicit calculations can be done:

Theorem 2.7 ([BD15] Lemmas 2.7 and 2.8) *Let $f \in \text{Cr}_2(\mathbf{K})$ be an elliptic element of infinite order.*

1. *If f is of the form $(x, y) \mapsto (\alpha x, \beta y)$ where $\alpha, \beta \in \mathbf{K}^*$ are such that the kernel of the group homomorphism $\mathbf{Z}^2 \rightarrow \mathbf{K}^*$, $(i, j) \mapsto \alpha^i \beta^j$ is generated by $(k, 0)$ for some $k \in \mathbf{Z}$, then the centralizer of f in $\text{Cr}_2(\mathbf{K})$ is*

$$\text{Cent}(f) = \{(x, y) \mapsto (\eta(x), yR(x^k)) \mid R \in \mathbf{K}(x), \eta \in \text{PGL}_2(\mathbf{K}), \eta(\alpha x) = \alpha\eta(x)\}.$$

2. *If $\text{char}(\mathbf{K}) = 0$ and if f is of the form $(x, y) \mapsto (\alpha x, y + 1)$, then the centralizer of f in $\text{Cr}_2(\mathbf{K})$ is*

$$\text{Cent}(f) = \{(x, y) \mapsto (\eta(x), y + R(x)) \mid \eta \in \text{PGL}_2(\mathbf{K}), \eta(\alpha x) = \alpha\eta(x), R \in \mathbf{K}(x), R(\alpha x) = R(x)\}.$$

If α is not a root of unity then R must be constant and $\eta(x) = \beta x$ for some $\beta \in \mathbf{K}^*$.

3. If $\text{char}(\mathbf{K}) = p > 0$ and if f is of the form $(x, y) \mapsto (\alpha x, y + 1)$ (where α must be of infinite order), then the centralizer of f in $\text{Cr}_2(\mathbf{K})$ is

$$\text{Cent}(f) = \{(x, y) \mapsto (R(y)x, y + t) \mid t \in \mathbf{K}, R(y) = S(y)S(y-1) \cdots S(y-p+1), S \in \mathbf{K}(y)\}.$$

Remark 2.8 Note that the elements η in the above statement can be explicitly described:

$$\{\eta \in \text{PGL}_2(\mathbf{K}) \mid \eta(\alpha x) = \alpha \eta(x)\} = \begin{cases} \text{PGL}_2(\mathbf{K}) & \text{if } \alpha = 1 \\ \{x \mapsto \gamma x^{\pm 1} \mid \gamma \in \mathbf{K}^*\} & \text{if } \alpha = -1 \\ \{x \mapsto \gamma x \mid \gamma \in \mathbf{K}^*\} & \text{if } \alpha \neq \pm 1 \end{cases}$$

Proof First case. We treat first the case where f is of the form $(x, y) \mapsto (\alpha x, \beta y)$. Let $(x, y) \mapsto (\frac{P_1(x, y)}{Q_1(x, y)}, \frac{P_2(x, y)}{Q_2(x, y)})$ be an element of $\text{Cent}(f)$; here $P_1, P_2, Q_1, Q_2 \in \mathbf{K}[x, y]$. The commutation relation gives us

$$\frac{P_1(\alpha x, \beta y)}{Q_1(\alpha x, \beta y)} = \frac{\alpha P_1(x, y)}{Q_1(x, y)}, \quad \frac{P_2(\alpha x, \beta y)}{Q_2(\alpha x, \beta y)} = \frac{\beta P_2(x, y)}{Q_2(x, y)}$$

which imply that P_1, P_2, Q_1, Q_2 are eigenvectors of the \mathbf{K} -linear automorphism $\mathbf{K}[x, y] \rightarrow \mathbf{K}[x, y], g(x, y) \mapsto g(\alpha x, \beta y)$. Therefore each one of the P_1, P_2, Q_1, Q_2 is a product of a monomial in x, y with a polynomial in $\mathbf{K}[x^k]$. Then we must have $\frac{P_1(x, y)}{Q_1(x, y)} = xR_1(x^k)$ and $\frac{P_2(x, y)}{Q_2(x, y)} = yR_2(x^k)$ for some $R_1, R_2 \in \mathbf{K}(x)$. The first factor $\frac{P_1(x, y)}{Q_1(x, y)}$ only depends on x , so for f to be birational it must be an element of $\text{PGL}_2(\mathbf{K})$. The conclusion in this case follows.

Second case. We now treat the case where $\text{char}(\mathbf{K}) = 0$ and where f is of the form $(x, y) \mapsto (\alpha x, y + 1)$. Let $(x, y) \mapsto (\frac{P_1(x, y)}{Q_1(x, y)}, \frac{P_2(x, y)}{Q_2(x, y)})$ be an element of $\text{Cent}(f)$. We have

$$\frac{P_1(\alpha x, y + 1)}{Q_1(\alpha x, y + 1)} = \frac{\alpha P_1(x, y)}{Q_1(x, y)}, \quad \frac{P_2(\alpha x, y + 1)}{Q_2(\alpha x, y + 1)} = \frac{P_2(x, y)}{Q_2(x, y)} + 1. \quad (3)$$

The first equation implies that P_1, Q_1 are eigenvectors of the \mathbf{K} -linear automorphism $\mathbf{K}[x, y] \rightarrow \mathbf{K}[x, y], g(x, y) \mapsto g(\alpha x, y + 1)$. We view an element of $\mathbf{K}[x, y]$ as a polynomial in x with coefficients in $\mathbf{K}[y]$. Since the only eigenvector of the \mathbf{K} -linear automorphism $\mathbf{K}[y] \rightarrow \mathbf{K}[y], g(y) \mapsto g(y + 1)$ is 1 (this is not true if $\text{char}(\mathbf{K}) > 0$), we deduce that P_1, Q_1 depend only on x . Thus, $\frac{P_1(x, y)}{Q_1(x, y)}$ is an element η of $\text{PGL}_2(\mathbf{K})$.

We derive $\psi = \frac{P_2}{Q_2}$ and get

$$\frac{\partial \psi}{\partial y}(\alpha x, y + 1) = \frac{\partial \psi}{\partial y}(x, y), \quad \frac{\partial \psi}{\partial x}(\alpha x, y + 1) = \alpha^{-1} \frac{\partial \psi}{\partial x}(x, y).$$

As before, this means that $\frac{\partial \psi}{\partial y}, \frac{\partial \psi}{\partial x}$ only depend on x (not true if $\text{char}(\mathbf{K}) > 0$). Hence, we can write ψ as $ay + B(x)$ with $a \in \mathbf{K}^*$ and $B \in \mathbf{K}(x)$. Then equation (3) implies

$B(\alpha x) = B(x) + 1 - a$, which implies further $x \frac{\partial B}{\partial x}(x)$ is invariant under $x \mapsto \alpha x$. If α is of infinite order, then $\frac{\partial B}{\partial x}(x) = \frac{c}{x}$ for some constant $c \in \mathbf{K}$. This is only possible if $c = 0$. So B is constant and $a = 1$ in this case. If α is a primitive k -th root of unity, then $(\eta(x), ay + B(x))$ commutes with $f^k : (x, y) \mapsto (x, y + k)$. This yields $a = 1$ and $B(\alpha x) = B(x)$.

Third case. We finally treat the case where $\text{char}(\mathbf{K}) = p > 0$ and where f has the form $(x, y) \mapsto (\alpha x, y + 1)$ with α of infinite order. Let $g \in \text{Cent}(f)$. Then g commutes with $f^p : (x, y) \mapsto (\alpha^p x, y)$ which is in the form of case 1 (the roles of x, y are exchanged). Thus, we know that g can be written as $(A(y)x, \eta(y))$ where $\eta \in \text{PGL}_2(\mathbf{K})$ and $A \in \mathbf{K}(x)$. Then $f \circ g = g \circ f$ implies that η is $y \mapsto y + R$ for some $R \in \mathbf{K}$ and that $A(y + 1) = A(y)$. The last equation implies $A(y) = S(y)S(y - 1) \cdots S(y - p + 1)$ for some $S \in \mathbf{K}(x)$. \square

2.4 Jonquière's twists with trivial action on the base

We follow [CD12b] in this section.

Lemma 2.9 *Let $f \in \text{Jonq}(\mathbf{K})$ be a Jonquière's twist. Let $\text{Cent}(f)$ be the centralizer of f in $\text{Cr}_2(\mathbf{K})$. Then $\text{Cent}(f) \subset \text{Jonq}(\mathbf{K})$.*

Proof The rational fibration preserved by a Jonquière's twist f is unique (cf. [DF01] Lemma 4.5), thus is also preserved by $\text{Cent}(f)$. \square

Let us consider centralizers of Jonquière's twists in $\text{Jonq}_0(\mathbf{K}) = \text{PGL}_2(\mathbf{K}(x))$ which is a linear algebraic group over the function field $\mathbf{K}(x)$. Let $f \in \text{Jonq}_0(\mathbf{K})$ and $M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \text{GL}_2(\mathbf{K}(x))$ be a matrix representing f where $A, B, C, D \in \mathbf{K}[x]$. We introduce the function $\Delta := \frac{\text{Tr}^2}{\det}$ which is well defined in PGL and is invariant by conjugation. This invariant Δ indicates the degree growth:

Lemma 2.10 ([CD12b] Theorem 3.3 [Xie15] Proposition 6.6) *The rational function $\Delta(f)$ is constant if and only if f is an elliptic element.*

Proof Let t_1, t_2 be the two eigenvalues of the matrix M which are elements of the algebraic closure of $\mathbf{K}(x)$. The invariant $\Delta(f)$ equals to $t_1/t_2 + t_2/t_1 + 2$. Since \mathbf{K} is algebraically closed, $\Delta(f) \in \mathbf{K}$ if and only if $t_1/t_2 \in \mathbf{K}$. If $t_1 = t_2$, then by conjugating M to a triangular matrix we can write f in the form $(x, y) \mapsto (x, y + a(x))$ with $a \in \mathbf{K}(x)$ and it follows that f is an elliptic element.

Suppose now that $t_1 \neq t_2$. Let $\zeta : C \rightarrow \mathbb{P}^1$ be the curve corresponding to the finite field extension $\mathbf{K}(x) \hookrightarrow \mathbf{K}(x)(t_1)$, here ζ is the identity map on \mathbb{P}^1 if $t_1, t_2 \in \mathbf{K}(x)$. The birational transformation f induces a birational transformation f_C on $C \times \mathbb{P}^1$ by base change. The induced map f_C is of the form $(x, (t_1/t_2)y)$ where t_1/t_2 is viewed as a function on C . The degree growth of f_C which is the same as f is linear if and only if t_1/t_2 is not a constant, i.e. if and only if $\Delta(f)$ is not a constant. \square

From now on we suppose that f is a Jonquière's twist so that $\Delta(f) \notin \mathbf{K}$. We still denote by t_1, t_2 the two eigenvalues of M as in the above proof, we know that $t_1 \neq t_2$.

We first study the centralizer $\text{Cent}_0(f)$ of f in $\text{Jonq}_0(\mathbf{K}) = \text{PGL}_2(\mathbf{K}(x))$. Let L be the finite extension of $\mathbf{K}(x)$ over which M is diagonalisable; it is $\mathbf{K}(x)$ itself or a quadratic extension of $\mathbf{K}(x)$, depending on whether or not t_1, t_2 are in $\mathbf{K}(x)$. The centralizer $\text{Cent}_0^L(f)$ of f in $\text{PGL}_2(L)$ is isomorphic to the multiplicative group L^* . So $\text{Cent}_0(f)$, being contained in $\text{Cent}_0^L(f)$ and containing all the iterates of f , must be a 1-dimensional torus over $\mathbf{K}(x)$. It is split if $L = \mathbf{K}(x)$, i.e. if $t_1, t_2 \in \mathbf{K}(x)$.

If $L = \mathbf{K}(x)$, then up to conjugation f can be written as $(x, y) \mapsto (x, b(x)y)$ with $b \in \mathbf{K}(x)^*$ and $\text{Cent}_0(f) = \{(x, y) \mapsto (x, \gamma(x)y) \mid \gamma \in \mathbf{K}(x)^*\}$.

If L is a quadratic extension of $\mathbf{K}(x)$ and if $\text{char}(\mathbf{K}) \neq 2$, we can put f in a simpler form and write $\text{Cent}_0(f)$ explicitly as follows. We may assume that the matrix $M = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$ has entry $C = 1$, after conjugation by $\begin{pmatrix} C & 0 \\ 0 & 1 \end{pmatrix}$. Once we have $C = 1$, a

conjugation by $\begin{pmatrix} 2 & D-A \\ 0 & 2 \end{pmatrix}$ allows us to put M in the form $\begin{pmatrix} A & B \\ 1 & A \end{pmatrix}$ with $A, B \in \mathbf{K}[x]$.

Therefore $\text{Cent}_0(f)$ is $\{Id, (x, y) \mapsto (x, \frac{C(x)y+B(x)}{y+C(x)}) \mid C \in \mathbf{K}(x)\}$ as the $(\mathbf{K}(x)$ -points of the) later algebraic group is easily seen to commute with f . Note that B is not a square in $\mathbf{K}(x)$ because M is not diagonalisable over $\mathbf{K}(x)$ and that the transformation $f : (x, y) \mapsto (x, \frac{A(x)y+B(x)}{y+A(x)})$ fixes pointwise the hyperelliptic curve defined by $y^2 = B(x)$.

Now we look at the whole centralizer of f . For $\eta \in \text{PGL}_2(\mathbf{K})$ and $f \in \text{Jonq}_0(\mathbf{K})$ represented by a matrix $\begin{pmatrix} A(x) & B(x) \\ C(x) & D(x) \end{pmatrix}$, we denote by f_η the element of $\text{Jonq}_0(\mathbf{K})$ represented by $\begin{pmatrix} A(\eta(x)) & B(\eta(x)) \\ C(\eta(x)) & D(\eta(x)) \end{pmatrix}$. Let $f \in \text{Jonq}_0(\mathbf{K})$ be a Jonquières twist and $g : (x, y) \mapsto (\eta(x), \frac{a(x)y+b(x)}{c(x)y+d(x)})$ be an element of $\text{Jonq}(\mathbf{K})$. Writing down the commutation equation, we see that g commutes with f if and only if f is conjugate to f_η in $\text{PGL}_2(\mathbf{K}(x))$ by the transformation represented by $\begin{pmatrix} a(x) & b(x) \\ c(x) & d(x) \end{pmatrix}^{-1}$. We have thus $\Delta(f)(x) = \Delta(f_\eta)(x) = \Delta(f)(\eta(x))$. Recall that $\Delta(f) \in \mathbf{K}(x)$ is not in \mathbf{K} . As a consequence the group

$$\{\eta \in \text{PGL}_2(\mathbf{K}), \Delta(f)(x) = \Delta(f)(\eta(x))\}$$

is a finite subgroup of $\text{PGL}_2(\mathbf{K})$. We then obtain:

Theorem 2.11 ([CD12b]) *Let $f \in \text{Jonq}_0(\mathbf{K})$ be a Jonquières twist preserving the rational fibration fiber by fiber. Let $\text{Cent}(f)$ be the centralizer of f in $\text{Cr}_2(\mathbf{K})$. Then $\text{Cent}(f) \subset \text{Jonq}(\mathbf{K})$ and $\text{Cent}_0(f) = \text{Cent}(f) \cap \text{Jonq}_0(\mathbf{K})$ is a finite index normal subgroup of $\text{Cent}(f)$. The group $\text{Cent}_0(f)$ has a structure of a 1-dimensional torus over $\mathbf{K}(x)$. In particular $\text{Cent}(f)$ is virtually abelian.*

Remark 2.12 In [CD12b], the authors give explicit description of the quotient $\text{Cent}(f)/\text{Cent}_0(f)$ when $\mathbf{K} = \mathbf{C}$.

Finite action on the base. If $f \in \text{Jonq}(\mathbf{K})$ is a Jonquières twist which has a finite action on the base, then $f^k \in \text{Jonq}_0(\mathbf{K})$ for some $k \in \mathbf{N}$. As $\text{Cent}(f) \subset \text{Cent}(f^k)$, we can use Theorem 2.11 to describe $\text{Cent}(f)$:

Corollary 2.13 *If $f \in \text{Jonq}(\mathbf{K})$ is a Jonquière's twist which has a finite action on the base, then $\text{Cent}(f)$ is virtually contained in a 1-dimensional torus over $\mathbf{K}(x)$. In particular $\text{Cent}(f)$ is virtually abelian.*

We are contented with this coarse description of $\text{Cent}(f)$ because this causes only a finite index problem as regards the embeddings of \mathbf{Z}^2 to $\text{Cr}_2(\mathbf{K})$. We give an example to show how we expect $\text{Cent}(f)$ to look like:

Example 2.14 *If f is $(x, y) \mapsto (a(x), R(x)y)$ where $R \in \mathbf{K}(x)$ and $a \in \text{PGL}_2(\mathbf{K})$ has order $k < +\infty$. Then all maps of the form $(x, y) \mapsto (x, S(x)S(a(x)) \cdots S(a^{k-1}(x))y)$ with $S \in \mathbf{K}(x)$ commute with f .*

3 Base-wandering Jonquière's twists

We introduce some notations. For a Hirzebruch surface X , let us denote by π the projection of X onto \mathbb{P}^1 , i.e. the rational fibration. When $X = \mathbb{P}^1 \times \mathbb{P}^1$, π is the projection onto the first factor. For $x \in \mathbb{P}^1$, we denote by F_x the fiber $\pi^{-1}(x)$. If f is a birational transformation of a Hirzebruch surface X which preserves the rational fibration, we denote by $\bar{f} \in \text{PGL}_2(\mathbf{K})$ the induced action of f on the base \mathbb{P}^1 and we will consider f as an element of $\text{Jonq}(\mathbf{K})$.

Assume now that f is a Jonquière's twist such that $\bar{f} \in \text{PGL}_2(\mathbf{K})$ if of infinite order, we will call it a *base-wandering Jonquière's twist*. We have an exact sequence:

$$\{1\} \rightarrow \text{Cent}_0(f) \rightarrow \text{Cent}(f) \rightarrow \text{Cent}_b(f) \rightarrow \{1\} \quad (4)$$

where $\text{Cent}_0(f) = \text{Cent}(f) \cap \text{Jonq}_0(\mathbf{K})$ and $\text{Cent}_b(f) \subset \text{Cent}(\bar{f}) \subset \text{PGL}_2(\mathbf{K})$. The action \bar{f} on the base is conjugate to $x \mapsto \alpha x$ with $\alpha \in \mathbf{K}^*$ of infinite order or to $x \mapsto x + 1$. The later case is only possible if $\text{char}(\mathbf{K}) = 0$. Thus $\text{Cent}_b(f)$ is a subgroup of $\{x \mapsto \gamma x, \gamma \in \mathbf{K}^*\}$ or of $\{x \mapsto x + \gamma, \gamma \in \mathbf{K}\}$. In both cases $\text{Cent}_b(f)$ is abelian. We first remark:

Lemma 3.1 *All elements of $\text{Cent}_0(f)$ are elliptic.*

Proof By Theorem 2.11, a Jonquière's twist in $\text{Jonq}_0(\mathbf{K})$ cannot have a base-wandering Jonquière's twist in its centralizer. \square

The rest of the article will essentially be occupied by the proof of the following theorem.

Theorem 3.2 *Let $f \in \text{Jonq}(\mathbf{K})$ be a base-wandering Jonquière's twist. The exact sequence*

$$\{1\} \rightarrow \text{Cent}_0(f) \rightarrow \text{Cent}(f) \rightarrow \text{Cent}_b(f) \rightarrow \{1\}$$

satisfies

- $\text{Cent}_0(f) = \text{Cent}(f) \cap \text{Jonq}_0(\mathbf{K})$, if not trivial, is $\{(x, y) \mapsto (x, ty), t \in \mathbf{K}^*\}$, $\{(x, y) \mapsto (x, y + t), t \in \mathbf{K}\}$ or a torsion group which has order two when $\text{char}(\mathbf{K}) = 0$;

- $\text{Cent}_b(f) \subset \text{PGL}_2(\mathbf{K})$ is isomorphic to the product of a finite cyclic group with \mathbf{Z} . The infinite cyclic subgroup generated by \bar{f} has finite index in $\text{Cent}_b(f)$.

Proof (of Theorem 3.2) The theorem is a consequence of Proposition 3.18, Corollary 3.23 and Proposition 3.35. \square

Corollary 3.3 When $\text{char}(\mathbf{K}) = 0$, the centralizer of a base-wandering Jonquières twist is virtually abelian.

Proof Theorem 3.2 implies that $\text{Cent}_b(f)$ is virtually the cyclic group generated by \bar{f} and that $\text{Cent}_0(f)$ is abelian when $\text{char}(\mathbf{K}) = 0$. The result follows. \square

Remark 3.4 Theorem 3.2 is nearly optimal in the sense that $\text{Cent}_b(f)$ can be \mathbf{Z} (Remark 3.5) or a product of \mathbf{Z} with a non trivial finite cyclic group (Example 3.34) and $\text{Cent}_0(f)$ can be trivial, isomorphic to \mathbf{K} , \mathbf{K}^* or $\mathbf{Z}/2\mathbf{Z}$ (Section 3.3).

Remark 3.5 If a base-wandering Jonquières twist f commutes with an elliptic element of the form $(x, y) \mapsto (x, ty)$ or $(x, y) \mapsto (x, y + t)$ then f can be written as $(\eta(x), yR(x^k))$ or $(\eta(x), y + R(x))$ by Theorem 2.7. A general base-wandering Jonquières twist can not be written as $(\eta(x), yR(x^k))$ or $(\eta(x), y + R(x))$. So the centralizer of a general Jonquières twist f differs from the infinite cyclic group $\langle f \rangle$ only by some finite groups. For example, for a generic choice of $\alpha, \beta \in \mathbf{K}^*$, the centralizer of $f_{\alpha, \beta} : (x, y) \mapsto (\alpha x, \frac{\beta y + x}{y + 1})$ is $\langle f_{\alpha, \beta} \rangle$, this is showed by J. Déserti in [Dés08].

3.1 Algebraically stable maps

If f is a birational transformation of a smooth algebraic surface X over \mathbf{K} , we denote by $\text{Ind}(f)$ the set of indeterminacy points of f . We say that f is *algebraically stable* if there is no curve V on X such that the strict transform $f^k(V) \subset \text{Ind}(f)$ for some integer $k > 0$. There always exists a birational morphism $\hat{X} \rightarrow X$ which lifts f to an algebraically stable birational transformation of \hat{X} ([DF01] Theorem 0.1). The following theorem says that for $f \in \text{Jonq}(\mathbf{K})$, we can get a more precise algebraically stable model:

Theorem 3.6 Let f be a birational transformation of a ruled surface X that preserves the rational fibration. Then there is a ruled surface \hat{X} and a birational map $\varphi : X \dashrightarrow \hat{X}$ such that

- the only singular fibers of \hat{X} are of the form $D_0 + D_1$ where D_0, D_1 are (-1) -curves, i.e. \hat{X} is a conic bundle;
- $f_{\hat{X}} = \varphi \circ f \circ \varphi^{-1}$ is an algebraically stable birational transformation of \hat{X} and it preserves the rational fibration of \hat{X} which is induced by that of X ;
- $f_{\hat{X}}$ sends singular fibers isomorphically to singular fibers and all indeterminacy points of $f_{\hat{X}}$ and its iterates are located on regular fibers.
- φ is a sequence of elementary transformations and blow-ups.

Let $z \in X$ be an indeterminacy point of f . Let $X \xleftarrow{u} Y \xrightarrow{v} X$ be a minimal resolution of the indeterminacy point z , i.e. u, v are birational maps which are regular locally outside the fiber over $\pi(z)$, u^{-1} is a series of n blow-ups at z or at its infinitely near points and n is minimal among possible integers.

Lemma 3.7 *The total transform by u^{-1} in Y of $F_{\pi(z)}$, the fiber containing z , is a chain of $(n+1)$ rational curves $C_0 + C_1 + \dots + C_n$: C_0 is the strict transform of $F_{\pi(z)}$, $C_0^2 = C_n^2 = -1$, $C_i^2 = -2$ for $0 < i < n$ and $C_i \cdot C_{i+1} = 1$ for $0 \leq i < n$.*

Proof Let us write $u : Y \rightarrow X$ as $Y = Y_n \xrightarrow{u_n} Y_{n-1} \dots \xrightarrow{u_2} Y_1 \xrightarrow{u_1} Y_0 = X$ where each u_i is a single contraction of a (-1) -curve. Denote by C_i the (-1) -curve contracted by u_i . By an abuse of notation, we will also use C_i to denote all strict transforms of C_i . The connectedness of the fibers and the preservation of the fibration imply that for each i , the map $f \circ u_1 \circ \dots \circ u_i$ has at most one indeterminacy point on a fiber. To prove the lemma, it suffices to show that the indeterminacy point of $f \circ u_1 \circ \dots \circ u_i$ which by construction lies in C_i is not the intersection point of C_i with C_{i-1} .

Suppose by contradiction that C_{i+1} is obtained by blowing up the intersection point of C_i with C_{i-1} . Then for $j > i$, the auto-intersection of C_i on X_j is less than or equal to -2 . Let us write $v : Y \rightarrow X$ as $Y = Y_n \xrightarrow{v_n} Y_{n-1} \dots \xrightarrow{v_2} Y_1 \xrightarrow{v_1} Y_0 = X$ where each v_i is a single contraction of a (-1) -curve. Since C_i is contracted by v , there must exist an integer k such that $v_{k+1} \circ \dots \circ v_n(C_i)$ is the (-1) -curve on Y_k contracted by v_k . By looking at the auto-intersection number, we see that it is possible only if the $C_j, j > i$ are all contracted by $v_k \circ \dots \circ v_n$. But by the minimality of the integer n , C_n can not be contracted by v . \square

Proof (of Theorem 3.6) Our proof is inspired by the proof of Theorem 0.1 of [DF01]. Let $p_1, \dots, p_k \in X$ be the indeterminacy points of f . By Lemma 3.7, for $1 \leq i \leq k$ the minimal resolution of f at p_i writes as

$$X = X_{i0} \xleftarrow{u_{i1}} X_{i1} \xleftarrow{u_{i2}} \dots \xleftarrow{u_{in_i}} X_{in_i} = Y_{in_i} \xrightarrow{v_{in_i}} \dots \xrightarrow{v_{i2}} Y_{i1} \xrightarrow{v_{i1}} Y_{i0} = X$$


where $u_{i1}, \dots, u_{in_i}, v_{i1}, \dots, v_{in_i}$ are single contractions of (-1) -curves and X_{in_i} has one singular fiber which is a chain of rational curves $C_{i0} + \dots + C_{in_i}$. Let us write the global minimal resolution of indeterminacy of f by keeping in mind the rational fibration:

$$\begin{array}{cccccccccccc} X = X_0 & \xrightarrow{\quad f_0 \quad} & X_1 & \xrightarrow{\quad f_1 \quad} & \dots & \xrightarrow{\quad f_{n-1} \quad} & X_n & \xrightarrow{\quad f_n \quad} & \dots & \xrightarrow{\quad f_{2n-2} \quad} & X_{2n-1} & \xrightarrow{\quad f_{2n-1} \quad} & X_{2n} = X \\ \downarrow \pi & & \downarrow \pi & & & & \downarrow \pi & & & & \downarrow \pi & & \downarrow \pi \\ \mathbb{P}^1 & \xrightarrow{\quad \bar{f}_0 \quad} & \mathbb{P}^1 & \xrightarrow{\quad \bar{f}_1 \quad} & \dots & \xrightarrow{\quad \bar{f}_{n-1} \quad} & \mathbb{P}^1 & \xrightarrow{\quad \bar{f}_n \quad} & \dots & \xrightarrow{\quad \bar{f}_{2n-2} \quad} & \mathbb{P}^1 & \xrightarrow{\quad \bar{f}_{2n-1} \quad} & \mathbb{P}^1 \end{array}$$

where $n = n_1 + \dots + n_k$ and

- f_0, \dots, f_{n-1} are blow-ups which correspond to the inverses of $u_{11}, \dots, u_{1n_1}, \dots, u_{k1}, \dots, u_{kn_k}$;
- f_n, \dots, f_{2n-1} are blow-downs which correspond to $v_{11}, \dots, v_{1n_1}, \dots, v_{k1}, \dots, v_{kn_k}$;
- X_n has k singular fibers which are chains of rational curves $C_{i0} + \dots + C_{in_i}, 1 \leq i \leq k$;

- the abusive notation π is self-explaining; we will also denote by $C_{i'l}$ a strict transform of C_{il} (if it remains a curve) on the surfaces X_j . On $X_0 = X_{2n}$, it is possible that $C_{i'0} = C_{in_i}$ for $1 \leq i, i' \leq k$.



For any $j \in \mathbf{N}$, we denote $j \bmod 2n$ by \bar{j} and we let $X_j = X_{\bar{j}}$, $f_j = f_{\bar{j}}$. If f_j blows up a point $r_j \in X_j$, then we denote by V_{j+1} the exceptional curve on X_{j+1} . If f_j contracts a curve $W_j \subset X_j$ then we denote by s_{j+1} the point $f_j(W_j) \in X_{j+1}$. For each V_j (resp. W_j), there is an i such that V_j (resp. W_j) is among C_{i0}, \dots, C_{in_i} . Suppose that f is not algebraically stable on H . Then there exist integers $1 \leq M < N$ such that f_M contracts W_M and

$$f_{N-1} \circ \dots \circ f_M(W_M) = r_N \in \text{Ind}(f_N).$$

Since f_N is a blow-up and f_M is a blow-down, we have $0 \leq \bar{N} \leq n-1$ and $n \leq \bar{M} \leq 2n-1$. We can assume that the length $(N-M)$ is minimal. Observe first that the minimality of the length implies for all j such that $M \leq j < N-1$, the point $t_{j+1} := f_j \circ \dots \circ f_M(W_M) = f_j \circ \dots \circ f_{M+1}(s_{M+1})$ is neither an indeterminacy point nor a point on a curve contracted by f_{j+1} .

The second observation is that for any j such that $M \leq j < N-1$, t_{j+1} is not on a singular fiber of X_{j+1} . The reason is as follows. Denote by M', N' the two numbers that satisfy $M < M' \leq N' < N$, $\bar{M}' = \bar{N}' = 0$, $M' - M \leq n$ and $N - N' \leq n$. If $M \leq j < M'$ and t_{j+1} was on a singular fiber, then t_{j+1} would be on a component contracted by some f_k with $j+1 \leq k < M'$. This contradicts the minimality of $N-M$ by our first observation. Similarly If $N' \leq j < N$ and t_{j+1} was on a singular fiber, then t_k would be an indeterminacy point of some f_k with $N' \leq k \leq j$, again contradicting the minimality

of $N - M$. If $M' \leq j < N'$ and t_{j+1} was on a singular fiber, then there would exist L such that $M' \leq L \leq j < L + 2n \leq N'$ and at least one of the t_L, \dots, t_{L+2n-1} is not a point where the corresponding map is locally isomorphic, still contradicting the minimality of $N - M$ by our first observation.

The second observation further implies that for j such that $M \leq j < j + 2n < N - 1$, t_{j+1}, t_{j+2n+1} are not on the same fiber of $X_{j+1} = X_{j+2n+1}$. Suppose to the contrary that this is not true for some j . Then there exists j' such that $j < j' \leq j + 2n$ and $\overline{j'} = \overline{M}$. $t_{j'}$ would also be on the same fiber as t_M . However it is the singular fiber containing W_M , contradiction to our second observation that $t_{j'}$ cannot be on a singular fiber.

Since f_{N-1} maps isomorphically the fiber of X_{N-1} containing t_{N-1} (which is regular by the above observation) to the fiber of X_N containing r_N , the fiber containing r_N is just one rational curve. As f_N is a blow-up, the fiber of X_{N+1} containing V_{N+1} is the union of two (-1) -curves, let us say, C_{k0} and $C_{k1} = V_{N+1}$. Then the fiber of X_N containing r_N is just C_{k0} . Similarly the singular fiber of X_M containing W_M is $C_{mm} + C_{m(n_m-1)}$ for some $1 \leq m \leq k$ and $W_M = C_{m(n_m-1)}$.

First case. Suppose that $m = k$ and $n_k = 1$. Let $a \in \mathbb{N}$ be the minimal integer such that $M + 2an > N$. Then for $N < j \leq M + 2an$, the surface X_j has a singular fiber $C_{k0} + C_{k1}$ and the maps $f_N, \dots, f_{M+2an-1}$ are all regular on $C_{k0} + C_{k1}$. Now we blow-up $t_{M+1}, \dots, t_{N-1}, r_N$. For $\overline{j_1} = \overline{j_2}$, we showed that t_{j_1}, t_{j_2} are not on the same fiber of $X_{j_1} = X_{j_2}$. This means that these blow-ups only give rise to singular fibers which are unions of two (-1) -curves. We denote by \hat{X}_j the modified surfaces, and \hat{f}_j the induced maps. Then every \hat{X}_j has singular fibers of the form $C_{k0} + C_{k1}$ and every \hat{f}_j is regular around these singular fibers. Let $\hat{f} = \hat{f}_{2n-1} \circ \dots \circ \hat{f}_0$. The number of indeterminacy points of \hat{f} (it was k for f) has decreased by one. Note that \hat{f} exchanges the two components C_{k0} and C_{k1} . This fact will be used in the proof of Proposition 3.13.

Second case. Suppose that $m = k$ and $n_k > 1$ or simply $m \neq k$. We blow-up r_N and contract the strict transform of the initial fiber containing r_N which is C_{k0} , obtaining a new surface \hat{X}_N whose corresponding fiber is now the single rational curve C_{k1} . We perform elementary transformations at t_{N-1}, \dots, t_{M+1} , i.e. we blow-up X_j at t_j and contract the strict transform of the initial fiber, replacing X_j with \hat{X}_j . This process has no ambiguity: if $\overline{j_1} = \overline{j_2}$, we showed that t_{j_1}, t_{j_2} are not on the same fiber of $X_{j_1} = X_{j_2}$, so the corresponding elementary transformations do not interfere with each other. Let us denote by $\hat{f}_M, \dots, \hat{f}_N$ the maps induced by f_M, \dots, f_N .

We now analyse the effects of $\hat{f}_M, \dots, \hat{f}_N$. First look at f_N , it lifts to a regular isomorphism after blowing up r_N . Thus \hat{f}_N is the blow-up at the point e_N of \hat{X}_N to which C_{k0} is contracted. After this step, the map going from X_{N-1} to \hat{X}_N induced by f_{N-1} is as following: it contracts the fiber containing t_{N-1} to e_N and blows up t_{N-1} . Then we make elementary transformations at t_{N-1}, \dots, t_{M+1} in turn. The maps $\hat{f}_{N-1}, \dots, \hat{f}_{M+1}$ are all regular on the modified fibers, thus they are still single blow-ups or single blow-downs. The behaviour of \hat{f}_M differs from the previous ones: it does not contract $C_{m(n_m-1)}$ any more, but contracts C_{mm} .

The hypothesis $m \neq k$ (or $m = k, n_k > 1$) forbids $C_{k0} \subset X_{N+1}$ to go back into the fiber of $X_{M+2na} = X_M$ containing W_M without being contracted. More precisely this implies the existence of $N' > N$ such that

- $X_{N+1}, \dots, X_{N'}$ all contain C_{k0} and C_{k1} ;

- $f_{N+1}, \dots, f_{N'-1}$ are regular on C_{k0} and $f_{N'}$ contracts C_{k0} ;
- if $a \in \mathbf{N}$ is the minimal integer such that $M + 2na > N$, then $N' < M + 2na$.

On the surfaces $X_{N+1}, \dots, X_{N'}$, C_{k0} is always a (-1) -curve, we contract all these C_{k0} and obtain new surfaces $\hat{X}_{N+1}, \dots, \hat{X}_{N'}$. The second and the third property listed above mean that the new induced maps $\hat{f}_N, \dots, \hat{f}_{N'}$ are all single blow-ups, single blow-downs or simply isomorphisms.

In summary we get a commutative diagram:

$$\begin{array}{cccccccccccccccc}
\hat{X}_0 & \xrightarrow{\hat{f}_0} & \hat{X}_1 & \xrightarrow{\hat{f}_1} & \dots & \xrightarrow{\hat{f}_{n-1}} & \hat{X}_n & \xrightarrow{\hat{f}_n} & \dots & \xrightarrow{\hat{f}_{2n-2}} & \hat{X}_{2n-1} & \xrightarrow{\hat{f}_{2n-1}} & \hat{X}_{2n} = \hat{X}_0 \\
\downarrow & & \downarrow & & & & \downarrow & & & & \downarrow & & \downarrow \\
X_0 & \xrightarrow{f_0} & X_1 & \xrightarrow{f_1} & \dots & \xrightarrow{f_{n-1}} & X_n & \xrightarrow{f_n} & \dots & \xrightarrow{f_{2n-2}} & X_{2n-1} & \xrightarrow{f_{2n-1}} & X_{2n} = X_0
\end{array}$$

where the vertical arrows are composition of elementary transformations and blow-ups. Let us remark that:

- the first vertical arrow $\hat{X}_0 \rightarrow X_0$ is a composition of elementary transformations.
- the blow-ups or the contractions of the \hat{f}_j only concern the k singular fibers and the exceptional curves are always among $C_{10}, \dots, C_{1n_1}, \dots, C_{k0}, \dots, C_{kn_k}$;
- there is no more C_{k0} . We then do a renumbering: C_{k1}, \dots, C_{kn_k} become $C_{k0}, \dots, C_{k(n_k-1)}$.

Let $\hat{f} = \hat{f}_{2n-1} \circ \dots \circ \hat{f}_0$. We repeat the above process. Either we are in the first case and k decreases, or we are in the second case and the total number of $C_{10}, \dots, C_{1n_1}, \dots, C_{k0}, \dots, C_{kn_k}$ decreases. As a consequence, after a finite number of times, either we get an algebraically stable map \hat{f} , or we will get rid of all the $C_{10}, \dots, C_{1n_1}, \dots, C_{k0}, \dots, C_{kn_k}$. In the later case \hat{f} is a regular automorphism, thus automatically algebraically stable. \square

3.2 Elliptic elements in the Jonquières group

Some materials in this section are taken from [Bla11]. We assume $\text{char}(\mathbf{K}) = 0$ in this section.

Lemma 3.8 *Assume $\text{char}(\mathbf{K}) = 0$. Let $f : (x, y) \mapsto (\eta(x), yR(x))$, $\eta \in \text{PGL}_2(\mathbf{K})$, $R \in \mathbf{K}(x)$ be an elliptic element. Then*

1. either $R \in \mathbf{K}$,
2. or $R(x) = \frac{rS(x)}{S(\eta(x))}$ with $r \in \mathbf{K}^*$ and $S \in \mathbf{K}(x) \setminus \mathbf{K}$.

Proof If η is the identity, then we see easily, by looking at the degree growth, that f is elliptic if and only if R is constant. From now on assume that η is not the identity.

We first consider the case where f has infinite order. Then by Proposition 2.4 f is conjugate to an automorphism of a Hirzebruch surface. By Theorem 3.6, the conjugation which turns f into an automorphism of a Hirzebruch surface is a sequence of elementary transformations. After conjugation it preserves the two strict transforms

of the two sections $\{y = 0\}$ and $\{y = \infty\}$. Therefore there exists $g \in \text{Jonq}_0(\mathbf{K})$ of the form $(x, y) \mapsto (x, yS(x))$, $S \in \mathbf{K}(x)$ such that $g \circ f \circ g^{-1}$ is $(x, y) \mapsto (\eta(x), ry)$ with $r \in \mathbf{K}^*$. Hence f is $(x, y) \mapsto (\eta(x), y \frac{rS(x)}{S(\eta(x))})$.

Now assume that f has finite order. Then η has also finite order. Denote by d the order of η and dk the order of f . The expression of f^d is $(x, y) \mapsto (x, T(x)y)$ where

$$T(x) = R(x)R(\eta(x)) \cdots R(\eta^{d-1}(x)).$$

As $f^{dk} = \text{Id}$, we have $T(x)^k = 1$. Therefore $T(x) = \alpha$ is a root of unity. Take a $\beta \in \mathbf{K}^*$ such that $\beta^d = \alpha$. Then $R'(x) = R(x)/\beta$ satisfies

$$R'(x)R'(\eta(x)) \cdots R'(\eta^{d-1}(x)) = 1.$$

In other words R' represents a cocycle of $\mathbf{Z}/d\mathbf{Z}$ with values in $\mathbf{K}^*(X)$. By Hilbert's Theorem 90, the cohomology group $H^1(\mathbf{Z}/d\mathbf{Z}, \mathbf{K}^*(X))$ is trivial. Thus the existence of $S \in \mathbf{K}^*(X)$ such that $R'(x) = \frac{S(x)}{S(\eta(x))}$. The conclusion follows. \square

Remark 3.9 If $\text{char}(\mathbf{K}) > 0$ then we cannot apply Hilbert's Theorem 90 in the above situation because the corresponding field extension may not be separable.

Remark 3.10 In the above lemma S may not be unique. If η has finite order and $T \in \mathbf{K}(x)$ is such that $T(x) = T(\eta(x))$, then $\frac{S(x)}{S(\eta(x))} = \frac{T(x)S(x)}{T(\eta(x))S(\eta(x))}$.

A direct corollary of Lemma 3.8 is:

Corollary 3.11 *If $\text{char}(\mathbf{K}) = 0$ then a diagonalisable elliptic element of $\text{Jonq}_0(\mathbf{K}) = \text{PGL}_2(\mathbf{K}(X))$ is always conjugate to an automorphism of a Hirzebruch surface.*

Lemma 3.12 *Let $f : (x, y) \mapsto (\eta(x), y + R(x))$, $\eta \in \text{PGL}_2(\mathbf{K})$, $R \in \mathbf{K}(x)$ be an elliptic element. Then one of the following holds:*

1. η has finite order;
2. there exists a coordinate x' such that η is $x' \mapsto vx'$ with $v \in \mathbf{K}^*$, and $R(x') = R_1(x') + R_2(\frac{1}{x'})$ where R_1, R_2 are polynomials in x' .
3. there exists a coordinate x' such that η is $x' \mapsto x' + 1$, and R is a polynomial in x' .

Proof Assume that η has infinite order, then for some coordinate x' , η can be written as $\eta(x') = x' \mapsto x' + 1$ or $x' \mapsto vx'$ with $v \in \mathbf{K}^*$. In coordinates (x', y) , write the transformation f as $(x', y) \mapsto (\eta(x'), y + C(x') + \frac{P(x')}{Q(x')})$ where $C, P, Q \in \mathbf{K}[x']$ are such that either $P = 0$ or $\deg P < \deg Q$. There is nothing to prove if $P = 0$. Suppose by contradiction that $\deg P < \deg Q$. For $n \in \mathbf{N}^*$, the iterate f^n is

$$(x', y) \mapsto \left(\eta^n(x'), y + \Sigma(x') + \frac{P(x')}{Q(x')} + \cdots + \frac{P(\eta^{n-1}(x'))}{Q(\eta^{n-1}(x'))} \right)$$

Where $\Sigma(x')$ is a polynomial of degree at most the degree of C . If $\eta(x') = vx'$ and $Q(x') = qx'^d$ for some $q \in \mathbf{K}$ and $d \in \mathbf{N}$, then replacing the coordinate x' with $x^* = 1/x'$ we can write f as $(x^*, y) \mapsto (\frac{x^*}{v}, y + C(\frac{1}{x^*}) + q^{-1}x^{*d}P(1/x^*))$. As $d = \deg Q > \deg P$, the coordinate x^* satisfies the required properties.

In the remaining cases, either $\eta(x') = x + 1$ or Q has a factor $x - a$ with $a \neq 0$. In both cases some factor of Q gives rise to an infinite number of distinct factors in the polynomials $Q(\eta^n(x')), n \in \mathbf{N}$. Then if we write

$$\frac{P(x')}{Q(x')} + \dots + \frac{P(\eta^{n-1}(x'))}{Q(\eta^{n-1}(x'))}$$

as a single fraction, then the degree of its denominator tends to infinity as n tends to infinity. This contradicts the hypothesis that the degrees of the f^n 's are bounded. This means $P = 0$ in these situations. \square

We refer to [Bla11] Section 3 (especially Proposition 3.3 and Lemma 3.9) for results that are stronger than the following proposition.

Proposition 3.13 *Assume that $\text{char}(\mathbf{K}) = 0$. Let $f \in \text{Jonq}(\mathbf{K})$ be an elliptic element that is not conjugate to an automorphism of a Hirzebruch surface. Then f has even finite order $2k$ and is conjugate to an automorphism f' of a conic bundle. Moreover f^k is in $\text{Jonq}_0(\mathbf{K})$ and f^k exchanges the two components of at least one singular fiber of the conic bundle.*

Proof We see by Proposition 2.4 that an elliptic element of infinite order is always conjugate to an automorphism of Hirzebruch surface. Hence our hypothesis implies immediately that f is of finite order. We can assume that f' is an algebraically stable map on a conic bundle X which satisfies the conclusions of Theorem 3.6. We claim that f' is an automorphism of X . Suppose by contradiction that p_0 is an indeterminacy point of f' . It must lie on a regular fiber F of X . Let d be the order of f' . As f' is algebraically stable, p_0 is not an indeterminacy point of f'^{-n} for any n and there exists p_1, \dots, p_d belonging to the fibers $f'^{-1}(F), \dots, f'^{-d}(F)$ such that f' is well defined at each p_j and $f'(p_j) = p_{j-1}$. However $f'^d = \text{Id}$ implies $p_d = p_0$, contradicting that p_0 is an indeterminacy point.

Since by hypothesis X is not a Hirzebruch surface, it must have some singular fibers. As f' has finite order, there exists a minimal $k > 0$ such that f'^k preserves the rational fibration fiber by fiber, i.e. $f'^k \in \text{Jonq}_0(\mathbf{K})$.

Let $M = \begin{pmatrix} A(X) & B(X) \\ C(X) & D(X) \end{pmatrix}$ be a matrix of $\text{GL}_2(\mathbf{K}(X))$ that represents $f'^k \in \text{Jonq}_0(\mathbf{K}) = \text{PGL}_2(\mathbf{K}(X))$.

We claim that M is not diagonalisable. It is proved when $\mathbf{K} = \mathbf{C}$ in [Bla11][page 479 parts (i), (ii) of the proof of Proposition 3.3] that this would imply that f is conjugate to an automorphism of a Hirzebruch surface; the proof works word by word when $\text{char}(\mathbf{K}) = 0$. A consequence is that f'^k exchanges the two components of at least one singular fiber of X because otherwise f'^k would be conjugate to an automorphism of a Hirzebruch surface and would be diagonalisable (see the formula (2)). Thus the order of f^k is even.

Let $2j$ be the order of f^k . Then there exists $\lambda \in \mathbf{K}(x)^*$ such that $M^{2j} = \lambda \text{Id}$ is a scalar matrix. Taking determinants of both side we get $\det M^{2j} = \lambda^2$. Thus there exists $\kappa \in \mathbf{K}(X)$ such that $\kappa^j = \lambda$. Let $P_M \in \mathbf{K}(X)[T]$ be the minimal polynomial of M . Then P_M is a factor of $T^{2j} - \lambda = T^{2j} - \kappa^j$. Since M is not diagonalisable the polynomial P_M is irreducible in $\mathbf{K}(X)[T]$. Any irreducible factor of $T^{2j} - \kappa^j$ has the form $T^2 - \mu \kappa$ for some j -th root of unity μ . Therefore M^2 is a scalar matrix. In other words $j = 1$ and f^k is an involution. \square

Corollary 3.14 *Assume that $\text{char}(\mathbf{K}) = 0$. Let $f \in \text{Jonq}_0(\mathbf{K})$ be an elliptic element which is not conjugate to an automorphism of a Hirzebruch surface. Then f is an involution and is conjugate to an automorphism f' of a conic bundle X . The automorphism f' fixes pointwise a hyperelliptic curve whose projection onto the base \mathbb{P}^1 is a ramified double cover. In some affine chart f can be written as $(x, y) \mapsto (x, \frac{a(x)}{y})$ with $a \in \mathbf{K}[x]$. The hyperelliptic curve is given by the equation $y^2 = a(x)$.*

Such involutions are well known and are called *Jonquières involutions*, see [BB00].

3.3 The group $\text{Cent}_0(f)$

Now we turn back to the study of centralizers. Let f be a base-wandering Jonquières twist. In [CD12b], it is proved by explicit calculations, in the case where $\mathbf{K} = \mathbf{C}$, that $\text{Cent}_0(f)$ is isomorphic to \mathbf{C}^* , $\mathbf{C}^* \rtimes \mathbf{Z}/2\mathbf{Z}$, \mathbf{C} or a finite group (this is not optimal). Their arguments do not work directly when $\text{char}(\mathbf{K}) > 0$. With a more precise description of elements of $\text{Jonq}_0(\mathbf{K})$, we simplify their arguments and improve their results.

Let $g \in \text{Cent}_0(f)$ be non trivial. By Lemma 3.1 g is elliptic. If g is conjugate to an automorphism of a Hirzebruch surface, then by proposition 2.6 by choosing suitable coordinates we can write g as $(x, y) \mapsto (x, \beta y)$ or $(x, y) \mapsto (x, y + 1)$.

Lemma 3.15 *Suppose that there exists a non trivial $g \in \text{Cent}_0(f)$ that can be written as $(x, y) \mapsto (x, \beta y)$ with $\beta \in \mathbf{K}^*$. Either f has the form $(a(x), R(x)y^{-1})$ with $a \in \text{PGL}_2(\mathbf{K})$ and $\text{Cent}_0(f)$ is an order two group generated by the involution $(x, y) \mapsto (x, -y)$, or f has the form $(a(x), R(x)y)$ and $\text{Cent}_0(f)$ is $\{(x, y) \mapsto (x, \gamma y), \gamma \in \mathbf{K}^*\}$.*

Proof The map g preserves $\{y = 0\}$ and $\{y = \infty\}$ and these two curves are the only g -invariant sections. Thus f permutes these two sections and has necessarily the form $(x, y) \mapsto (a(x), R(x)y^{\pm 1})$ where $R \in \mathbf{K}(x)$ and $a \in \text{PGL}_2(\mathbf{K})$ has infinite order. If f is $(a(x), R(x)y^{-1})$, then $\beta = -1$. Since $\text{Cent}(f) \subset \text{Cent}(f^2)$ we replace f by f^2 in what follows so that we can assume f is $(a(x), R(x)y)$.

The only f -invariant sections are $\{y = 0\}$ and $\{y = \infty\}$. Indeed an invariant section s satisfies

$$s(a^n(x)) = R(x) \cdots R(a^{n-1}(x))s(x) \quad \forall n \in \mathbf{N}.$$

If s was not $\{y = 0\}$ nor $\{y = \infty\}$, then the two sides of the equations would be non-zero rational fractions and the degree of the right side would go to infinity with n because f is a Jonquières twist. Thus, an element of $\text{Cent}_0(f)$ permutes the two f -invariant sections and has the form $(x, A(x)y)$ or $(x, \frac{A(x)}{y})$ with $A \in \mathbf{K}(x)$. In the first case the commutation relation implies $A(a(x)) = A(x)$ which further implies that A is a constant. In the second

case the commutation relation gives $A(a(x))^{-1}R(x)^2A(x) = 1$ which further implies that $(a(x), R(x)^2y)$ is conjugate by $(x, A(x)y)$ to an elliptic element $(a(x), y)$. This is not possible because the map $f' : (x, y) \mapsto (a(x), R(x)^2y)$ is a Jonquière's twist. Indeed the iterates f^n, f^m are respectively

$$(a^n(x), R(x) \cdots R(a^{n-1}(x))y) \quad \text{and} \quad (a^n(x), (R(x) \cdots R(a^{n-1}(x)))^2y)$$

and they have the same degree growth.

Reciprocally all elements of the form $(x, y) \mapsto (x, \beta y)$ with $\beta \in \mathbf{K}^*$ commute with $f : (x, y) \mapsto (a(x), R(x)y)$ and we have already observed that $(x, y) \mapsto (x, -y)$ is the only non trivial element of $\text{Jonq}_0(\mathbf{K})$ which commutes with $(a(x), R(x)y^{-1})$. \square

Lemma 3.16 *Suppose that there exists a non trivial $g \in \text{Cent}_0(f)$ that can be written as $(x, y) \mapsto (x, y + 1)$. Then f has the form $(a(x), y + S(x))$ with $S \in \mathbf{K}(x)$ and $\text{Cent}_0(f)$ is $\{(x, y + \gamma), \gamma \in \mathbf{K}\}$.*

Proof The section $\{y = \infty\}$ is the only g -invariant section. Thus f preserves this section and has the form $(x, y) \mapsto (a(x), R(x)y + S(x))$ where $R, S \in \mathbf{K}(x)$ and $a \in \text{PGL}_2(\mathbf{K})$ is of infinite order. Writing down the relation $f \circ g = g \circ f$, we see that $R = 1$. Thus f is $(a(x), y + S(x))$ where S belongs to $\mathbf{K}(x)$ but not to $\mathbf{K}[x]$ since f is a Jonquière's twist. The only f -invariant section is $\{y = \infty\}$. Indeed an invariant section s satisfies

$$s(a^n(x)) = s(x) + S(x) + \cdots + S(a^{n-1}(x)) \quad \forall n \in \mathbf{N}.$$

If s was not $\{y = \infty\}$, then the two sides of the equations are rational fractions. The degree of the right-hand side grows linearly in n while the degree of the left-hand side does not depend on n , contradiction. Thus, an element of $\text{Cent}_0(f)$ fixes $\{y = \infty\}$ and has the form $(x, A(x)y + B(x))$ with $A, B \in \mathbf{K}(x)$. Writing down the commutation relation, we get

$$A(x)y + B(x) + S(x) = A(a(x))y + A(a(x))S(x) + B(a(x)).$$

The fact that a has infinite order implies that A is a constant. Then the equation is reduced to

$$B(x) + (1 - A)S(x) - B(a(x)) = 0.$$

If $A \neq 1$, then $f : (x, y) \mapsto (a(x), y + S(x))$ would be conjugate by $(x, y + \frac{B(x)}{1-A})$ to the elliptic element $(a(x), y)$. Therefore $A = 1$ and B is a constant. Reciprocally we see that all elements of the form $(x, y) \mapsto (x, y + \beta)$ with $\beta \in \mathbf{K}$ commute with $f : (x, y) \mapsto (a(x), y + S(x))$. \square

Lemma 3.17 *Assume that no non-trivial element of $\text{Cent}_0(f)$ is conjugate to an automorphism of a Hirzebruch surface and that $\text{Cent}_0(f)$ has a non-trivial element g . Then $\text{Cent}_0(f)$ is a torsion group. If moreover $\text{char}(\mathbf{K}) = 0$ then g has order two and is the only non-trivial element of $\text{Cent}_0(f)$.*

Proof By Lemma 3.1, g is an elliptic element. Since an elliptic element of infinite order is conjugate to an automorphism of a Hirzebruch surface by Proposition 2.4, g

has finite order. If $\text{char}(\mathbf{K}) = 0$ then g is a Jonquières involution of a conic bundle X by Corollary 3.14.

Assume now that $\text{char}(\mathbf{K}) = 0$. Then by Corollary 3.14 g fixes pointwise a hyperelliptic curve C . The map f induces an action on C , equivariant with respect to the ramified double cover. The action of f on C is infinite, this is possible only if the action of f on the base is up to conjugation $x \mapsto \alpha x$ and if C is a rational curve whose projection on the base \mathbb{P}^1 is ramified over $x = 0, x = \infty$. Then the only singular fibers of X are over $x = 0, x = \infty$. Note that f can have at most one indeterminacy point on each of these two fibers. If f had an indeterminacy point on these two fibers, then it would be a fixed point of g because g commutes with f and g preserves each fiber. But the only fixed point of g on a singular fiber is the intersection point of the two components, which can not be an indeterminacy point by Lemma 3.7. Therefore the Jonquières twist f must have an indeterminacy point over a point whose orbit in the base is infinite. This implies that the indeterminacy points of all the iterates of f form an infinite set. As g commutes with all the iterates of f , it fixes an infinite number of these indeterminacy points. Thus, the hyperelliptic curve C associated to g is the Zariski closure of these indeterminacy points and is uniquely determined by f . However C determines g too. (cf. [BB00] Proposition 2.7). Therefore g is uniquely determined by f and is the only non trivial element of $\text{Cent}_0(f)$. \square

Putting together the three previous lemmas, we obtain the following improvement of [CD12b]:

Proposition 3.18 *Let f be a base-wandering Jonquières twist. If $\text{Cent}_0(f)$ is not trivial, then it is $\{(x, y) \mapsto (x, ty), t \in \mathbf{K}^*\}$, $\{(x, y) \mapsto (x, y + t), t \in \mathbf{K}\}$, $\langle (x, y) \mapsto (x, -y) \rangle$ or a torsion group. If $\text{char}(\mathbf{K}) = 0$ and $\text{Cent}_0(f)$ is a non-trivial torsion group, then it is a group of order two generated by a Jonquières involution.*

3.4 Persistent indeterminacy points

3.4.1 general facts

Let f be a birational transformation of a surface X . An indeterminacy point $x \in X$ of f will be called *persistent* if 1) for every $i > 0$, f^{-i} is regular at x ; and 2) there are infinitely many curves contracted onto x by the iterates f^{-n} , $n \in \mathbf{N}$. This notion of persistence and the following idea appeared first in a non published version of [Can11], and it is also applied to some particular examples in [Dés08]. Note that it is different from the definition of persistent base points given in [BD15].

Proposition 3.19 *Let f be an algebraically stable birational transformation of a surface X . Suppose that there exists at least one persistent indeterminacy point with an infinite backward orbit. Let n denote the number of such indeterminacy points. Then the centralizer $\text{Cent}(f)$ of f admits a morphism $\varphi : \text{Cent}(f) \rightarrow \mathcal{S}_n$ to the symmetric group of order n satisfying the following property: for any $g \in \text{Ker}(\varphi)$, there exists $l \in \mathbf{Z}$ such that $g \circ f^l$ preserves fiber by fiber a pencil of rational curves.*

Proof The algebraic stability of f will be used throughout the proof, we will not recall it each time. Denote by p_1, \dots, p_n the persistent indeterminacy points of f . Let g be

a birational transformation of X which commutes with f . Fix an index $1 \leq n_0 \leq n$. Since $\{f^{-i}(p_{n_0}), i > 0\}$ is infinite, there exists $k_0 > 0$ such that g is regular at $f^{-k}(p_{n_0})$ for all $k \geq k_0$. For infinitely many $j > 0$, f^{-j} contracts a curve onto p_{n_0} , denote these curves by $C_{n_0}^j$. There exists $k_1 > 0$ such that g does not contract $C_{n_0}^k$ for all $k \geq k_1$. We deduce, from the above observations and the fact that f and g commute, that for $k \geq k_0$ the point $g(f^{-k}(p_{n_0}))$ is an indeterminacy point of some f^m with $0 < m \leq k + k_1$. Then there exists $0 \leq m_0 < m$ such that

- for $0 \leq i \leq m_0$, f^i is regular at $g(f^{-k}(p_{n_0}))$;
- $f^{m_0}(g(f^{-k}(p_{n_0}))) = g(f^{m_0-k}(p_{n_0}))$ is an indeterminacy point of f .

By looking at $g(f^{-k}(p_{n_0}))$ and $C_{n_0}^{k'}$ for infinitely many k, k' , we see that the above indeterminacy point does not depend on k and is persistent with an infinite backward orbit. So it is $p_{\sigma_g(n_0)}$ for some $1 \leq \sigma_g(n_0) \leq n$. This gives us a well defined map $\sigma_g : \{1, \dots, n\} \rightarrow \{1, \dots, n\}$.

Now let g, h be two elements of $\text{Cent}(f)$. Then by considering a sufficiently large k for which g is regular at $f^{-k}(p_{n_0})$ and h is regular at $g(f^{-k}(p_{n_0}))$, we see that $\sigma_h \circ \sigma_g = \sigma_{h \circ g}$. By taking $h = g^{-1}$ we see that σ_g is bijective. We have then a group homomorphism φ from $\text{Cent}(f)$ to the symmetric group \mathcal{S}_n which sends g to σ_g .

Assume that n_0 is a fixed point of σ_g , this holds in particular when $g \in \text{Ker}(\varphi)$. We keep the previous notations. Since $g(f^{-k}(p_{n_0}))$ is an indeterminacy point of f^m whose forward orbit meets p_{n_0} , for an appropriate choice of $l \leq k$ we have

$$g \circ f^l(f^{-k}(p_{n_0})) = f^{-k}(p_{n_0})$$

for all $k \geq k_0$. This implies further

$$g \circ f^l(C_{n_0}^{k'}) = C_{n_0}^{k'}$$

for all sufficiently large k' . We conclude by Lemma 3.20 below. \square

The proof of the following lemma in [Can10] is written over \mathbf{C} for rational self-maps. The same proof works in all characteristics for birational self-maps but not for general rational self-maps (this is also observed and used in [Xie15].).

Lemma 3.20 *If a birational transformation f of a smooth algebraic surface preserves infinitely many curves then these curves are members of a pencil of curves and f preserves each member of this pencil of curves.*

Note that Lemma 3.20 is not needed in the proof of Proposition 3.19 if $f \in \text{Jonq}(\mathbf{K})$, and this is the only case in this paper where we apply Proposition 3.19 (see Corollary 3.23).

Remark 3.21 Using the tools introduced in [LU18] and the same idea of Proposition 3.19, Lonjou-Urech generalized Proposition 3.19 and Corollary 3.23 below to higher dimension, see [LU18] Theorem 1.6.

3.4.2 persistent indeterminacy points for Jonquières twists

We examine the notion of persistence in the Jonquières group and give a complement to Theorem 3.6:

Proposition 3.22 *Let f be a Jonquières twist acting algebraically stably on a conic bundle X as in the statement of Theorem 3.6. Then an indeterminacy point p of f is persistent if and only if the orbit of $\pi(p) \in \mathbb{P}^1$ under \bar{f} is infinite. And in that case, every $f^{-i}, i \in \mathbf{N}^*$ contracts a curve onto p .*

Proof If $\pi(p)$ has a finite orbit then p certainly cannot be persistent. Let us assume that the orbit of $\pi(p)$ is infinite. Then \bar{f} is conjugate to $x \mapsto \alpha x$ with $\alpha \in K^*$ of infinite order or to $x \mapsto x + 1$ (only when $\text{char}(\mathbf{K}) = 0$). By the algebraic stability of f , f^{-i} is regular at p for all $i > 0$ and all the points $f^{-i}(p), i > 0$ are on distinct fibers. Denote by x_0, x_1 the points $\pi(p), \bar{f}(\pi(p))$. By Theorem 3.6, we know that the fibers F_{x_0}, F_{x_1} are not singular. Thus f is regular on $F_{x_0} \setminus \{p\}$ and contracts it onto a point $q \in F_{x_1}$; f^{-1} is regular on $F_{x_1} \setminus \{q\}$ and contracts it onto p . Now pick a point x_n in the forward orbit of x_0 by \bar{f} and consider the fiber F_{x_n} . The fiber F_{x_n} cannot be contracted onto q by $f^{-(n-1)}$ because of the algebraic stability of f . As a consequence it is contracted by f^{-n} onto p . \square

Corollary 3.23 *Let f be a Jonquières twist acting algebraically stably on a conic bundle X as in the statement of Theorem 3.6. Suppose that the base action $\bar{f} \in \text{PGL}_2(\mathbf{K})$ has infinite order and there is an indeterminacy point of f located on a fiber $F_x \subset X$ such that $\bar{f}(x) \neq x$.*

1. *If \bar{f} has the form $x \mapsto x + 1$ then $\text{Cent}_b(f)$ is isomorphic to \mathbf{Z} ;*
2. *if \bar{f} has the form $x \mapsto \alpha x$ then $\text{Cent}_b(f)$ is isomorphic to the product of \mathbf{Z} with a finite cyclic group.*

Note that the first case does not occur when $\text{char}(\mathbf{K}) \neq 0$.

Proof Proposition 3.22 shows that the birational transformation f satisfies the hypothesis of Proposition 3.19. Let n denote the number of persistent indeterminacy points of f with infinite backward orbits. Let $g \in \text{Cent}(f)$. It is in $\text{Jonq}(\mathbf{K})$ by Lemma 2.9. Proposition 3.19 says that $g^{n!} \circ f^l$ preserves every member of a pencil of rational curves for some $l \in \mathbf{Z}$. The proof of Proposition 3.19 shows that infinitely many members of this pencil of rational curves are fibers of the initial rational fibration on X . Therefore this pencil of rational curves is the initial rational fibration. This means $\bar{g}^{n!} \circ \bar{f}^l = \text{Id} \in \text{PGL}_2(\mathbf{K})$.

When $\text{char}(\mathbf{K}) = 0$ and \bar{f} is $x \mapsto x + 1$, its centralizer in $\text{PGL}_2(\mathbf{K})$ is isomorphic to the additive group \mathbf{K} and this group is torsion free. Thus, $\text{Cent}_b(f)$ is contained in an infinite cyclic group in which $\langle \bar{f} \rangle$ has index $\leq n!$. The conclusion follows in this case.

When \bar{f} is $x \mapsto \alpha x$ with α of infinite order, its centralizer in $\text{PGL}_2(\mathbf{K})$ is isomorphic to the multiplicative group \mathbf{K}^* . The difference is that, in this case it is possible that \bar{g} has finite order $\leq n!$. Thus, we may have an additional finite cyclic factor in $\text{Cent}_b(f)$. \square

3.5 Local analysis around a fiber

Now we need to study the case where there is no persistent indeterminacy points. In this section we will work in the following setting:

- Let f be a base-wandering Jonquières twist. We can suppose that \bar{f} is $x \mapsto \alpha x$ or $x \mapsto x + 1$.
- Up to taking an algebraically stable model as in Theorem 3.6, we can suppose that f is a birational transformation of a conic bundle X which satisfies the properties in Theorem 3.6. We can assume that X is minimal in the sense that f switches the two components of every singular fiber of X .
- We assume that the only indeterminacy points of f are on the fibers F_0, F_∞ . This is because these two fibers are the only possible invariant fibers and if f has an indeterminacy point elsewhere then the situation is treated in the previous section.

Without loss of generality, let us suppose that f has an indeterminacy point p on the fiber F_∞ . By algebraic stability f^{-1} has an indeterminacy point $q \neq p$ on F_∞ . If $x \in \mathbb{P}^1$ is not 0 nor ∞ , then the orbit of x under \bar{f} is infinite and the fiber F_x is regular. As f has an indeterminacy point on F_∞ , the fiber F_∞ is also regular. Assume that F_0 is singular, then it is the union of two (-1) -curves and f exchanges the two components. Since $\text{Cent}(f) \subset \text{Cent}(f^2)$ and the aim of this section is to prove that $\text{Cent}_b(f)$ is finite by cyclic, it is not harmful to replace f with f^2 so that the two components of F_0 are no more exchanged and we can assume that F_0 is regular. Thus, we can suppose that

- the surface X is a Hirzebruch surface.

If \bar{f} is $x \mapsto \alpha x$, then $\text{Cent}_b(f)$ is contained in $\{(x \mapsto \gamma x), \gamma \in \mathbf{K}^*\}$ and all elements of $\text{Cent}_b(f)$ fix 0 and ∞ . Similarly if \bar{f} is $x \mapsto x + 1$ then all elements of $\text{Cent}_b(f)$ fix ∞ . Thus F_0 or F_∞ is $\text{Cent}(f)$ -invariant (under total transforms), we will study the (semi-)local behaviour of the elements in $\text{Cent}(f)$ around such an invariant fiber.

3.5.1 An infinite chain

We blow up X at p, q the indeterminacy points of f, f^{-1} , obtaining a new surface X_1 . The fiber of X_1 over 0 is a chain of three rational curves $C_{-1} + C_0 + C_1$ where C_1 (resp. C_{-1}) is the exceptional curve corresponding to p (resp. q) and C_0 is the strict transform of $F_\infty \subset X$. Now f induces a birational transformation f_1 of X_1 . As in Lemma 3.7, we know that f_1 (resp. f_1^{-1}) has an indeterminacy point p_2 (resp. q_2) on C_1 (resp. C_{-1}) which is disjoint from C_0 . We then blow up p_2, q_2 and repeat the process. We have:

- for every $n \in \mathbf{N}$, a surface X_n on which f induces a birational transformation f_n ;
- the fiber of X_n over 0 is a chain of rational curves $C_{-n}, \dots, C_0, \dots, C_n$;
- f_n (resp. f_n^{-1}) has an indeterminacy point p_{n+1} (resp. q_{n+1}) on C_n (resp. C_{-n}) disjoint from C_{n-1} (resp. $C_{-(n-1)}$).

Let g be a birational transformation of X which commutes with f . We already observed that F_∞ is an invariant fiber of g . If g is regular on F_∞ , then the commutativity implies that g preserves the set $\{p, q\}$. Suppose that g is not regular on F_∞ . Then g (resp. g^{-1}) has an indeterminacy point p' (resp. q') on F_∞ . Replacing g by g^{-1} or f by f^{-1} , we can suppose that $p' \neq q$. Then for every point $x \in F_\infty$ such that $x \neq p, p'$, we have that $g(q) = g(f(x)) = f(g(x))$ is a point, thus equals q . This further implies $q = q'$. Then we apply the same argument to g, f^{-1} , obtaining $p = p'$. In summary, g is either regular on F_∞ and preserves $\{p, q\}$, or the set of indeterminacy points of g, g^{-1} on F_∞ is exactly $\{p, q\}$.

We lift g to a birational transformation on X_n . By repeating the above arguments, we deduce that for all $n \in \mathbf{N}$ the two indeterminacy points of f_n, f_n^{-1} on the fiber $F_\infty \subset X_n$ coincide with that of g_n, g_n^{-1} if the later exist. This means that for a C_i given, and for sufficiently large n , the rational curve C_i is a component of the fiber of X_n and g_n maps it to another component C_j of the fiber. In other words g acts on the infinite chain of rational curves $\sum_{n \in \mathbf{Z}} C_n$. The dual graph of this infinite chain of rational curves is a chain of vertices indexed by \mathbf{Z} . The action of f on the dual graph is just a non trivial translation. The isomorphism group of the dual graph is isomorphic to $\mathbf{Z} \rtimes \mathbf{Z}/2\mathbf{Z}$. Those isomorphisms which commute with a non trivial translation coincide with the subgroup of translations \mathbf{Z} . The above considerations can be summarized as follows:

Lemma 3.24 *There is a group homomorphism $\Phi : \text{Cent}(f) \rightarrow \mathbf{Z}$ such that $g(C_n) = C_{\Phi(g)+n}$ for $g \in \text{Cent}(f)$. An element $g \in \text{Cent}(f)$ is in the kernel of Φ if and only if $g(C_n) = C_n$ for every $n \in \mathbf{Z}$. In other words an element g of the kernel of Φ is regular on the fiber F_∞ and fixes the indeterminacy points of f, f^{-1} on this fiber.*

Lemma 3.25 *Let g be an element of $\text{Cent}(f)$. Let $x \in \mathbb{P}^1$ be a point not fixed by \bar{f} . Then g can not have any indeterminacy points on the fiber F_x over x .*

Proof By our hypothesis f is regular on all fibers F_{x_n} where $\{x_n, n \in \mathbf{Z}\}$ denote the orbit of x under \bar{f} . If g had an indeterminacy point p on F_x , then $f(p), f^2(p), \dots$ would give us an infinite number of indeterminacy points of g . \square

Corollary 3.26 *Suppose that \bar{f} is conjugate to $x \mapsto x + 1$ (in particular $\text{char}(\mathbf{K}) = 0$). Let $g \in \text{Cent}(f)$ be in the kernel of $\Phi : \text{Cent}(f) \rightarrow \mathbf{Z}$. Then g is an automorphism of X . Furthermore g preserves the rational fibration fiber by fiber.*

Proof Lemma 3.24 says that g does not have any indeterminacy point on the fiber F_∞ . Lemma 3.25 says that g does not have any indeterminacy point elsewhere neither. Thus, g is an automorphism. Since \bar{g} commutes with $\bar{f} : x \mapsto x + 1$, \bar{g} is $x \mapsto x + v$ for some $v \in \mathbf{K}$. Suppose by contradiction that $v \neq 0$. Then g is an elliptic element of infinite order and $f \in \text{Cent}(g)$. We can apply Theorem 2.7 to g, f and put them in normal form. As f is a Jonquières twist, the rational fibration preserved simultaneously by f and g is unique and it must be the rational fibration appeared in the normal form. Hence, Theorem 2.7 forbids \bar{f}, \bar{g} to be both non-trivial and of the form $x \mapsto x + w$ with $w \in \mathbf{K}$. \square

When \bar{f} is of the form $x \mapsto \alpha x$, there are two special fibers F_0, F_∞ and the above easy argument does not work.

3.5.2 Formal considerations along a fiber

In the rest of this section we will assume that \bar{f} is $x \mapsto \alpha x$. There are two invariant fibers F_∞ and F_0 in this case. We assume that f has an indeterminacy point q on F_0 .

The idea of what we do in the sequel is as follows. Let us look at the case where $\mathbf{K} = \mathbf{C}$. The indeterminacy point $q \in F_0$ of f^{-1} is a fixed point of f , at which the differential of f has two eigenvalues 0 and α ; the fiber direction is superattracting and in the transverse direction f is just $x \mapsto \alpha x$. Therefore there is a local invariant manifold at q for f , which is a local holomorphic section of the rational fibration. Likewise, there is a local invariant manifold at $p \in F_0$, the indeterminacy point of f . These two local holomorphic sections allow us to conjugate locally holomorphically f to $(\alpha x, a(x)y)$ where a is a germ of holomorphic function. The structure of Jonquières maps is nice enough to allow us to apply this geometric idea over any field in an elementary way. We need just to work with formal series instead of polynomials.

From now on we fix $f : (x, y) \mapsto (\alpha x, \frac{A(x)y+B(x)}{C(x)y+D(x)})$ where $\alpha \in \mathbf{K}^*$ has infinite order and $A, B, C, D \in \mathbf{K}[x]$. Without loss of generality, we suppose that 1) the point $(0, 0)$ (resp. $(0, \infty)$) is an indeterminacy point of f (resp. f^{-1}); 2) one of the A, B, C, D is not a multiple of x . This implies

$$B(0) = C(0) = D(0) = 0, A(0) \neq 0. \quad (5)$$

We will consider A, B, C, D as elements of the ring of formal series $\mathbf{K}[[x]]$. We will also view f as an element of the formal Jonquières group $\mathrm{PGL}_2(\mathbf{K}((x))) \rtimes \mathbf{K}^*$ whose elements are formal expressions of the form $(\mu x, \frac{a(x)y+b(x)}{c(x)y+d(x)})$ where $\mu \in \mathbf{K}^*$ and a, b, c, d belong to $\mathbf{K}((x))$, the fraction field of $\mathbf{K}[[x]]$.

Normal form. We want to conjugate f to a formal expression of the form $(\alpha x, \beta(x)y)$, $\beta \in \mathbf{K}((x))$ by some formal expression $(x, \frac{E(x)y+F(x)}{G(x)y+H(x)})$ with $E, F, G, H \in \mathbf{K}[[x]]$. This amounts to say that we are looking for $E, F, G, H \in \mathbf{K}[[x]]$ such that $EF - GH \neq 0$ and

$$\begin{pmatrix} E(\alpha x) & F(\alpha x) \\ G(\alpha x) & H(\alpha x) \end{pmatrix}^{-1} \begin{pmatrix} A(x) & B(x) \\ C(x) & D(x) \end{pmatrix} \begin{pmatrix} E(x) & F(x) \\ G(x) & H(x) \end{pmatrix}$$

is a diagonal matrix. By writing out the explicit expressions of the up-right entry and the down-left entry of this matrix product, we obtain two equations to solve:

$$F(x)H(\alpha x)A(x) + H(x)H(\alpha x)B(x) - F(x)F(\alpha x)C(x) - H(x)F(\alpha x)D(x) = 0 \quad (6)$$

$$-E(x)G(\alpha x)A(x) - G(x)G(\alpha x)B(x) + E(x)E(\alpha x)C(x) + G(x)E(\alpha x)D(x) = 0 \quad (7)$$

We will use minuscules to denote the coefficients of the formal series, e.g. $E(x) = \sum_{i \in \mathbf{N}} e_i x^i$. Let us first look at the constant terms of equations (6), (7), they give

$$-e_0 g_0 a_0 - g_0^2 b_0 + e_0^2 c_0 + e_0 g_0 d_0 = 0 = f_0 h_0 a_0 + h_0^2 b_0 - f_0 f_0 c_0 - f_0 h_0 d_0.$$

Since $b_0 = c_0 = d_0 = 0$ and $a_0 \neq 0$ (see Equation (5)), we must have $e_0 g_0 = f_0 h_0 = 0$. We can choose $f_0 = g_0 = 0$ and $e_0 = h_0 = 1$, this guarantees in particular that our solution will satisfy $EH - FG \neq 0$.

Remark that the equations (6) and (7) involve respectively only E, G and F, H , and they have exactly the same form. So it suffices to show the existence of E, G which satisfy equation (6). The constant term is done, let us look at the x term. This leads to a linear equation in e_1, g_1 with coefficients involving $a_0, b_0, c_0, d_0, e_0, g_0$ and α . Therefore there exists at least one solution for e_1, g_1 . Then we turn to the next term and get a linear equation in e_2, g_2 , and so on. Hence, we can find E, F, G, H which satisfy the desired properties. To sum up, we have:

Lemma 3.27 *There exists $E, F, G, H \in \mathbf{K}[[x]]$ such that:*

- $E(0) = H(0) = 1$ and $F(0) = G(0) = 0$, in particular $\begin{pmatrix} E & F \\ G & H \end{pmatrix} \in \mathrm{PGL}_2(\mathbf{K}((x)))$;
- $(x, \frac{E(x)y+F(x)}{G(x)y+H(x)})$ conjugates f to $(\alpha x, \beta(x)y)$ for some $\beta \in \mathbf{K}((x))$;

Projective line over $\mathbf{K}((x))$. We call an element of $\mathbb{P}^1(\mathbf{K}((x))) = \mathbf{K}((x)) \cup \{\infty\}$ a formal section. We say a formal section $\theta(x)$ passes through the origin if $\theta(0) = 0$. An element $u = (\mu x, \frac{a(x)y+b(x)}{c(x)y+d(x)})$ of the formal Jonquières group $\mathrm{PGL}_2(\mathbf{K}((x))) \rtimes \mathbf{K}^*$ acts on $\mathbb{P}^1(\mathbf{K}((x)))$ in the following way:

$$\theta(x) \mapsto u \cdot \theta(x) = \begin{cases} \infty & \text{if } c(\mu^{-1}x)\theta(\mu^{-1}x) + d(\mu^{-1}x) = 0 \\ \frac{a(\mu^{-1}x)\theta(\mu^{-1}x) + b(\mu^{-1}x)}{c(\mu^{-1}x)\theta(\mu^{-1}x) + d(\mu^{-1}x)} & \text{otherwise} \end{cases},$$

$$\infty \mapsto \begin{cases} \infty & \text{if } c = 0 \\ \frac{a(\mu^{-1}x)}{c(\mu^{-1}x)} & \text{if } c \neq 0 \end{cases}.$$

Geometrically this is saying that a formal section of the rational fibration is sent to another by a formal Jonquières transformation. Remark that this action on $\mathbb{P}^1_{\mathbf{K}((x))}$ is not an automorphism of $\mathbf{K}((x))$ -algebraic variety. In scheme theoretic language, we have a commutative diagram:

$$\begin{array}{ccc} \mathbb{P}^1_{\mathbf{K}((x))} & \xrightarrow{\theta \mapsto u \cdot \theta} & \mathbb{P}^1_{\mathbf{K}((x))} \\ \downarrow & & \downarrow \\ \mathrm{Spec}(\mathbf{K}((x))) & \xrightarrow{\mu x \mapsto x} & \mathrm{Spec}(\mathbf{K}((x))). \end{array}$$

Thus, we have a group homomorphism from $\mathrm{PGL}_2(\mathbf{K}((x))) \rtimes \mathbf{K}^*$ to the group of such twisted automorphisms of $\mathbb{P}^1_{\mathbf{K}((x))}$.

Now let $g \in \mathrm{Cent}(f)$ be an element in the kernel of Φ . Recall (see Lemma 3.24) that g is regular on the fiber F_0 and fixes $(0, 0), (0, \infty)$. We showed that f is conjugate by $\begin{pmatrix} E & F \\ G & H \end{pmatrix}$ to a formal expression \hat{f} of the form $(\alpha x, \beta(x)y)$. We conjugate g by $\begin{pmatrix} E & F \\ G & H \end{pmatrix}$ too to get a formal expression \hat{g} . Then \hat{g} commutes with \hat{f} .

Recall that, by Lemma 3.27, we get $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ when we evaluate the formal expression $\begin{pmatrix} E & F \\ G & H \end{pmatrix}$ at $x = 0$. Together with the fact that $g \in \text{Ker}(\Phi)$, this implies that we get $y \mapsto \delta_0 y$ for some $\delta_0 \in \mathbf{K}^*$ when we evaluate \hat{g} at $x = 0$.

Let us consider the actions of \hat{f}, \hat{g} on $\mathbb{P}_{\mathbf{K}((x))}^1$ as described above. Since \hat{f} is in diagonal form, it fixes the points 0 and ∞ of $\mathbb{P}_{\mathbf{K}((x))}^1$.

Lemma 3.28 *If $\theta \in \mathbb{P}_{\mathbf{K}((x))}^1$ satisfies $\theta(0) = 0$ and $\hat{f} \cdot \theta(x) = \theta(x)$, then $\theta = 0$.*

Proof The equation $\hat{f} \cdot \theta(x) = \theta(x)$ can be written as $\beta(\alpha^{-1}x)\theta(\alpha^{-1}x) = \theta(x)$, i.e. $\theta(\alpha x)^{-1}\beta(x)\theta(x) = 1$. Suppose by contradiction that θ is not 0. Then we can write $\theta(x)$ as $x^r \tilde{\theta}(x)$ where $r > 0$ and $\tilde{\theta}(0) \neq 0$. Hence we have $\tilde{\theta}(\alpha x)^{-1}\beta(x)\tilde{\theta}(x) = \alpha^r$. This implies that \hat{f} is conjugate by $(x, \tilde{\theta}(x)y)$ to $(\alpha x, \alpha^r y)$. Since $\tilde{\theta}(0) \neq 0$ and $\begin{pmatrix} E(0) & F(0) \\ G(0) & H(0) \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$, this implies that the initial Jonquieres twist f is regular on the fiber F_0 , contradiction. \square

Since \hat{g} is $y \mapsto \delta_0 y$ at $x = 0$, it sends the formal section $0 \in \mathbb{P}^1(\mathbf{K}((x)))$ to another former section passing through the origin. The fact that \hat{f} and \hat{g} commute and the fact that 0 is the only fixed formal section of \hat{f} which passes through the origin imply that \hat{g} fixes $0 \in \mathbb{P}_{\mathbf{K}((x))}^1$. Likewise \hat{g} fixes ∞ too. Therefore \hat{g} can be written as $(\gamma x, \delta(x)y)$ where $\gamma \in \mathbf{K}^*$ and $\delta \in \mathbf{K}((x))$ satisfies $\delta(0) = \delta_0 \neq 0$.

Normal forms for a pair. Let us assume for the moment that γ is not a root of unity; we are going to prove that this is impossible. We want to, under this hypothesis, conjugate $\hat{g} = (\gamma x, \delta(x)y)$ to $(\gamma x, \delta(0)y)$ by $h = (x, \xi(x)y)$ for some $\xi \in \mathbf{K}[[x]]$. Remark that the conjugate of \hat{f} by h will still be in diagonal form.

We write $\delta = \frac{\omega}{\sigma}$ where $\omega, \sigma \in \mathbf{K}[[x]]$ satisfies $\omega(0) \neq 0, \sigma(0) \neq 0$ and $\frac{\omega(0)}{\sigma(0)} = \delta(0)$.

We will write ξ as $\sum_{i \in \mathbf{N}} \xi_i x^i$, and likewise for σ, ω .

After conjugation by $h = (x, \xi(x)y)$, \hat{g} becomes

$$\tilde{g} = h \circ \hat{g} \circ h^{-1} = (\gamma x, \frac{\xi(\gamma x)}{\xi(x)} \frac{\omega(x)}{\sigma(x)} y).$$

Therefore the equation we want to solve is

$$\xi(\gamma x)\omega(x) = \frac{\omega_0}{\sigma_0} \xi(x)\sigma(x). \quad (8)$$

The constant terms of the two sides are automatically equal, let us just choose $\xi_0 = 1$. Comparing the other terms, we obtain

$$\begin{aligned} \xi_0 \omega_1 + \gamma \xi_1 \omega_0 &= \frac{\omega_0}{\sigma_0} (\xi_0 \sigma_1 + \xi_1 \sigma_0) \\ \xi_0 \omega_2 + \gamma \xi_1 \omega_1 + \gamma^2 \xi_2 \omega_0 &= \frac{\omega_0}{\sigma_0} (\xi_0 \sigma_2 + \xi_1 \sigma_1 + \xi_2 \sigma_0) \\ &\dots \end{aligned}$$

which are equivalent to

$$\begin{aligned}(\gamma - 1)\omega_0\xi_1 &= \frac{\omega_0}{\sigma_0}\xi_0\sigma_1 - \xi_0\omega_1 \\(\gamma^2 - 1)\omega_0\xi_2 &= \frac{\omega_0}{\sigma_0}(\xi_0\sigma_2 + \xi_1\sigma_1) - \xi_0\omega_2 - \gamma\xi_1\omega_1 \\&\dots\end{aligned}$$

For the i -th term, we have a linear equation whose coefficient before ξ_i is $(\gamma^i - 1)\omega_0$. Since $\omega \neq 0$ and we have supposed that γ is not a root of unity, The above equations always have solutions. In summary, we have the following intermediate lemma (we will get from this lemma a contradiction so its hypothesis is in fact absurd):

Lemma 3.29 *Suppose that $g \in \text{Ker}(\Phi)$ and the action of g on the base is of infinite order. Then we can conjugate f and g , simultaneously by an element in $\text{PGL}_2(\mathbf{K}((x)))$ whose evaluation at $x = 0$ is $\text{Id} : y \mapsto y$, to*

$$\tilde{g} = (\gamma x, \delta y), \tilde{f} = (\alpha x, \beta(x)y)$$

where $\alpha, \gamma, \delta \in \mathbf{K}^*$, $\beta \in \mathbf{K}((x))^*$ and α, γ are of infinite order.

Writing down the equation $\tilde{f} \circ \tilde{g} = \tilde{g} \circ \tilde{f}$, we get $\delta\beta(x) = \delta\beta(\gamma x)$. As $\delta \neq 0$, we get $\beta(x) = \beta(\gamma x)$. We write $\beta = \frac{\beta^{num}}{\beta^{den}}$ with $\beta^{num}, \beta^{den} \in \mathbf{K}[[x]]$ such that at least one of the $\beta_0^{num}, \beta_0^{den}$ is not 0. The equation becomes

$$\beta^{num}(x)\beta^{den}(\gamma x) = \beta^{den}(x)\beta^{num}(\gamma x).$$

By comparing the coefficients of two sides, we get

$$\forall k \in \mathbf{N}, \sum_{i+j=k} \beta_i^{num}\beta_j^{den}\gamma^j = \sum_{i+j=k} \beta_i^{den}\beta_j^{num}\gamma^j.$$

Then by induction on k we get from these equations:

1. if $\beta_0^{num} = 0$ then $\beta^{num} = 0$, this is impossible;
2. if $\beta_0^{den} = 0$ then $\beta^{den} = 0$, this is again impossible;
3. if $\beta^{den}\beta^{num} \neq 0$ then $\beta^{num} = \kappa\beta^{den}$ for some $\kappa \in \mathbf{K}^*$. Then $\tilde{f} = (\alpha x, \kappa y)$, this contradicts the fact that the original birational transformation f has an indeterminacy point on the fiber F_0 because to get \tilde{f} we only did conjugations whose evaluation at $x = 0$ are the identity $y \mapsto y$.

Thus, we get

Proposition 3.30 *Suppose that $g \in \text{Ker}(\Phi)$. Then \bar{g} has finite order and g is an elliptic element of $\text{Cr}_2(\mathbf{K})$.*

Proof The previous discussion showed that \bar{g} cannot have infinite order. Then an iterate g^k is in $\text{Jonq}_0(\mathbf{K})$ and $f \in \text{Cent}(g^k)$. By Theorem 2.11, an element which commutes with a Jonquière twist in $\text{Jonq}_0(\mathbf{K})$ cannot have an infinite action on the base. As \bar{f} has infinite order, g^k must be elliptic. So g must be elliptic. \square

3.5.3 Another fiber

The base action $\bar{f} \in \mathrm{PGL}_2(\mathbf{K})$ is $x \mapsto \alpha x$, it has two fixed points 0 and ∞ . Recall that we are always under the hypothesis that the indeterminacy points of f are on the fibers F_0, F_∞ . We have done analysis around the fiber F_0 on which f has an indeterminacy point. We will denote by Φ_0 the homomorphism Φ we considered before. In case f has also an indeterminacy point on F_∞ , we denote the corresponding homomorphism by Φ_∞ .

Lemma 3.31 *The image of $\mathrm{Aut}(X) \cap \mathrm{Ker}(\Phi_0) \subset \mathrm{Cent}(f)$ in $\mathrm{Cent}_b(f) \subset \mathrm{PGL}_2(\mathbf{K})$ is a finite cyclic group.*

Proof We recall first that the automorphism group of a Hirzebruch surface is an algebraic group (see [Mar71]). An element of $\mathrm{Cent}(f)$ which is regular everywhere on H must be in $\mathrm{Ker}(\Phi_0)$. Thus, $\mathrm{Aut}(X) \cap \mathrm{Ker}(\Phi_0) = \mathrm{Aut}(X) \cap \mathrm{Cent}(f)$ is an algebraic subgroup of $\mathrm{Aut}(H)$. An automorphism of a Hirzebruch surface always preserves the rational fibration and there is a morphism of algebraic groups from $\mathrm{Aut}(X)$ to $\mathrm{PGL}_2(\mathbf{K})$ (see [Mar71]). The image of $\mathrm{Aut}(X) \cap \mathrm{Ker}(\Phi_0) \subset \mathrm{Cent}(f)$ in $\mathrm{Cent}_b(f) \subset \mathrm{PGL}_2(\mathbf{K})$ is an algebraic subgroup Λ of $\mathrm{PGL}_2(\mathbf{K})$. By Proposition 3.30, the elements of Λ are all multiplication by roots of unity. If Λ was infinite then it would equal to its Zariski closure in $\mathrm{PGL}_2(\mathbf{K})$ and would be isomorphic to the multiplicative group \mathbf{K}^* . But the existence of a base-wandering Jonquieres twist means that \mathbf{K}^* contains elements of infinite order, for example α . This contradicts the fact that $\Lambda = \mathbf{K}^*$ is torsion. The conclusion follows. \square

We first look at the case where we have two homomorphisms Φ_0, Φ_∞ :

Proposition 3.32 *If f has an indeterminacy point on F_∞ , then $\mathrm{Ker}(\Phi_0) = \mathrm{Ker}(\Phi_\infty)$ is a subgroup of $\mathrm{Aut}(X)$. The image of $\mathrm{Ker}(\Phi_0)$ in $\mathrm{Cent}_b(f) \subset \mathrm{PGL}_2(\mathbf{K})$ is a finite cyclic group.*

Proof Let g be an element of $\mathrm{Ker}(\Phi_0)$. By Proposition 3.30 g is an elliptic element of $\mathrm{Cr}_2(\mathbf{K})$. If $\Phi_\infty(g)$ were not trivial, then g would act by a non trivial translation on the corresponding infinite chain of rational curves and could not be conjugate to any automorphism. This means g must belong to $\mathrm{Ker}(\Phi_\infty)$ and consequently g must be an automorphism of H . The second part of the statement follows from Lemma 3.31. \square

When f is regular on F_∞ , we may need to do a little bit more, but we get more precise information as well:

Proposition 3.33 *If f has no indeterminacy points on F_∞ , then $\mathrm{Ker}(\Phi_0)$ is a finite cyclic group whose elements are automorphisms of $\mathbb{P}^1 \times \mathbb{P}^1$ of the form $(x, y) \mapsto (\gamma x, y)$ with γ a root of unity.*

Proof Assume that f is regular on F_∞ . Let $g \in \mathrm{Ker}(\Phi_0)$ be a non trivial element, it is regular on F_0 . By Lemma 3.25, an indeterminacy point of g can only be located on F_∞ .

Suppose that g has an indeterminacy point p on F_∞ . Then g^{-1} also has an indeterminacy point q on F_∞ . If $p \neq q$, then g would act by translation on the corresponding

infinite chain of rational curves. This means that g would never be conjugate to an automorphism of some surface and contradicts Proposition 3.30 which asserts that g is elliptic. Thus, we have $p = q$. The facts that f commutes with g and that f is regular on F_∞ imply $f(p) = p$. We blow up the Hirzebruch surface X at p to get a new surface X' and induced actions f', g' . The induced action f' is still regular on the fiber F'_∞ and preserves both of the two irreducible components. If g' has an indeterminacy point on F'_∞ , then as before it coincides with the indeterminacy point of g'^{-1} and must be fixed by f' . Then we can keep blowing up indeterminacy points of maps induced from g , or contracting g -invariant (-1) -curves in the fiber, without losing the regularity of the map induced by f . As g is elliptic, we will get at last a surface \hat{X} with induced actions \hat{f}, \hat{g} which are both regular on the fiber over ∞ . We can suppose that \hat{X} is minimal among the surfaces with this property. In particular \hat{g} is an automorphism of \hat{X} . Moreover, the proof of Theorem 3.6 shows that \hat{X} is a conic bundle and the only possible singular fiber is \hat{F}_∞ . We claim that \hat{F}_∞ is in fact regular. Suppose by contradiction that \hat{F}_∞ is singular. Then it is a chain of two (-1) -curves and \hat{g} exchanges the two components. However the conic bundle \hat{X} is obtained from a Hirzebruch surface by a single blow-up, it has a unique section of negative self-intersection which passes through only one of the two components of the singular fiber. As a consequence, the automorphism \hat{g} cannot exchange the two components, contradiction. Thus, replacing X by \hat{X} , we can suppose from the beginning that g is an automorphism of the Hirzebruch surface X .

Suppose by contradiction that g preserves only finitely many sections of the rational fibration. Since f commutes with g , we can assume, after perhaps replacing f by some of its iterates, that f and g preserve simultaneously a section of the rational fibration. Removing this section and the fiber F_0 from H , we get an open set isomorphic to \mathbb{A}^2 restricted to which f can be written as $(x', y') \mapsto (\alpha^{-1}x', A(x')y' + B(x'))$ where $A, B \in \mathbf{K}(x')$. The rational function A must be a constant because f acts as an automorphism on this affine open set. Likewise the rational function B must be a polynomial. But then $(\deg(f^n))_{n \in \mathbf{N}}$ would be a bounded sequence. This contradicts the fact that f is a Jonquières twist.

Hence, if $g \in \text{Ker}(\Phi)$ is non-trivial then it preserves necessarily infinitely many sections. This forces g to preserve each member of a pencil of rational curves on X whose general members are sections (see Lemma 3.20). This is only possible if $X = \mathbb{P}^1 \times \mathbb{P}^1$ and g acts as $(x, y) \mapsto (\gamma x, y)$ with $\gamma \in \mathbf{K}^*$; here the projection of $\mathbb{P}^1 \times \mathbb{P}^1$ onto the first factor is the original rational fibration we were looking at. This allows us to conclude by Lemma 3.31. \square

Example 3.34 Let μ be a k -th root of unity, the pair $f : (x, y) \mapsto (\alpha x, \frac{(1+x^k)y+x^k}{(2+x^k)y+1+x^k})$, $g : (x, y) \mapsto (\mu x, y)$ satisfy the conditions in Proposition 3.33.

Now let f be a base-wandering Jonquières twist which satisfies the hypothesis made at the beginning of Section 3.5; in particular f is regular outside $F_0 \cap F_\infty$ and \bar{f} is $x \mapsto \alpha x$ or $x \mapsto x + 1$. The image $\Phi_\infty(\text{Cent}(f))$ is an infinite cyclic subgroup of \mathbf{Z} and is isomorphic to \mathbf{Z} , it is generated by $\Phi_\infty(g)$ for some $g \in \text{Cent}(f)$. Then for any $h \in \text{Cent}(f)$, there exists $k \in \mathbf{Z}$ such that $g^{-k} \circ h \in \text{Ker}(\Phi_\infty)$. Thus, $\bar{g}^{-k} \circ \bar{h}$ belongs to the image of $\text{Ker}(\Phi_\infty)$ in $\text{Cent}_b(f)$. By Corollary 3.26, Proposition 3.32 and Proposition 3.33, the image of $\text{Ker}(\Phi_\infty)$ in $\text{Cent}_b(f)$ is at worst finite cyclic. Note that $\text{Cent}_b(f)$

is always abelian. Therefore we obtain the last piece of information to prove Theorem 3.2:

Proposition 3.35 *Let f be a base-wandering Jonquières twist which satisfies the hypothesis made at the beginning of Section 3.5. Let g be an element of $\text{Cent}(f)$ such that $\Phi_0(g)$ generates the image of Φ . Then $\text{Cent}_b(f)$ is the product of a finite cyclic group with the infinite cyclic group generated by \bar{g} .*

4 Proofs of the main results

Proof (of Theorem 1.1) The proof is a direct combination of Theorems 2.1, 2.2, 2.7, 2.11 and 3.2. \square

Proof (of Theorem 1.5) Centralizers of loxodromic elements are virtually cyclic by Theorem 2.1 of Blanc-Cantat. It is proved in [Giz80],[Can11] that centralizers of Halphen twists are virtually abelian (see Theorem 2.2). Centralizers of Jonquières twists whose actions on the base are of finite order are contained in tori over the function field $\mathbf{K}(x)$, thus are abelian ([CD12b] see Theorem 2.11). Corollary 3.3 says that centralizers of base-wandering Jonquières twists are virtually abelian when $\text{char}(\mathbf{K}) = 0$. Centralizers of infinite order elliptic elements (due to [BD15]) are described in Theorem 2.7, we use the notations therein. In the first two cases of Theorem 2.7, by using Remark 2.8 we see that $\text{Cent} f$ is abelian if $\alpha \neq \pm 1$, contains an abelian subgroup of index 2 if $\alpha = -1$ and is not virtually abelian if $\alpha = 1$. In the third case of Theorem 2.7, the kernel of the projection onto the y -coordinate is an abelian subgroup of finite index of $\text{Cent} f$. \square

Proof (of Remark 1.3) In the first case the degree function is bounded on Γ . Indeed there exists $j, k \in \mathbf{N}^*$ such that f^j, g^k have degree 1 because f, g are elliptic. Then f^j, g^k generates a subgroup of finite index of Γ all of whose non-trivial elements have degree one.

In the second case, the two Halphen twists f and g are automorphisms of a rational surface X preserving an elliptic fibration $X \rightarrow \mathbb{P}^1$. The elliptic fibration is induced by the linear system corresponding to mK_X for some $m \in \mathbf{N}^*$. For $n \in \mathbf{N}$, the actions of f^n and g^n on $\text{Pic}(X)$ are respectively

$$D \mapsto D - mn(D \cdot K_X)\Delta_i + \left(-\frac{m^2}{2}(D \cdot K_X) \cdot (n\Delta_i)^2 + m(D \cdot (n\Delta_i)) \right) K_X, \quad i = 1, 2$$

where (\cdot) denotes the intersection form and $\Delta_i \in \text{Pic}(X)$ satisfies $\Delta_i \cdot K_X = 0$ (cf. [Giz80], [BD15] Section 5). Therefore the action of $f^i \circ g^j$ on $\text{Pic}(X)$ is

$$D \mapsto D - mi(D \cdot K_X)\Delta_1 - mj(D \cdot K_X)\Delta_2 + \lambda_{ij}K_X \quad \text{where}$$

$$\lambda_{ij} = -\frac{m^2}{2}(D \cdot K_X) \cdot (i^2\Delta_1^2 + j^2\Delta_2^2) + mD \cdot (i\Delta_1 + j\Delta_2) - ijm^2(D \cdot K_X)(\Delta_1 \cdot \Delta_2).$$

Let Λ be an ample class on X . Then the degree of $f^i \circ g^j$ is up to a bounded term (cf. [BD15] Section 5)

$$\Lambda \cdot (f^i \circ g^j)^* \Lambda = \Lambda^2 - \frac{m^2}{2} (\Lambda \cdot K_X)^2 (i^2 \Delta_1^2 + j^2 \Delta_2^2) - ijm^2 (\Lambda \cdot K_X)^2 (\Delta_1 \cdot \Delta_2).$$

Note that Δ_1^2 and Δ_2^2 are negative.

The third case is [BC16] Lemma 5.7.

In the fourth case the description of the degree function follows directly from the explicit expressions. \square

Theorem 4.1 *Suppose that $\text{char}(\mathbf{K}) = 0$. Let $G \subset \text{Cr}_2(\mathbf{K})$ be a maximal abelian subgroup which has at least one element of infinite order. Then up to conjugation one of the following possibilities holds:*

1. G is $\{(x, y) \mapsto (\alpha x, \beta y) \mid \alpha, \beta \in \mathbf{K}^*\}$, $\{(x, y) \mapsto (\alpha x, y + v) \mid \alpha \in \mathbf{K}^*, v \in \mathbf{K}\}$ or $\{(x, y) \mapsto (x + u, y + v) \mid u, v \in \mathbf{K}\}$;
2. G is the product of $\{(x, y) \mapsto (x, \beta y) \mid \beta \in \mathbf{K}^*\}$ with an infinite torsion group G_1 . Each element of G_1 has the form

$$(x, y) \mapsto \left(\eta x, y \frac{S(x)}{S(\eta x)} \right) \quad \eta \text{ is a root of unity, } S \in \mathbf{K}(x)$$

All elements of G are elliptic but G is not conjugate to a group of automorphisms of any rational surface.

3. G is the product of $\{(x, y) \mapsto (x, y + v) \mid v \in \mathbf{K}\}$ with an infinite torsion group G_1 . Each element of G_1 has the form

$$(x, y) \mapsto (\eta x, y + S(x) - S(\eta x)) \quad \eta \text{ is a root of unity, } S \in \mathbf{K}(x)$$

All elements of G are elliptic but G is not conjugate to a group of automorphisms of any rational surface.

4. G has a finite index subgroup contained in $\text{Jonq}_0(\mathbf{K}) = \text{PGL}_2(\mathbf{K}(x))$.
5. A finite index subgroup G' of G is a cyclic group generated by a base-wandering Jonquières twist.
6. A finite index subgroup G' of G is isomorphic to $\mathbf{K}^* \times \mathbf{Z}$ (resp. $\mathbf{K} \times \mathbf{Z}$) where the first factor is $\{(x, y) \mapsto (x, \beta y) \mid \beta \in \mathbf{K}^*\}$ (resp. $\{(x, y) \mapsto (x, y + v) \mid v \in \mathbf{K}\}$) and the second factor is generated by a base-wandering Jonquières twist, as in the fourth case of Theorem 1.1;
7. A finite index subgroup G' of G is isomorphic to \mathbf{Z}^s with $s \leq 8$ and G' preserves fiberwise an elliptic fibration;
8. A finite index subgroup G' of G is a cyclic group generated by a loxodromic element.

The existences of type two and type three maximal abelian groups are less obvious than the others. We give here two examples.

Example 4.2 Let $q \in \mathbf{N}^*$. Let $(\xi_n)_n$ be a sequence of elements of \mathbf{K}^* such that ξ_n is a primitive q^n -th root of unity and $\xi_n^q = \xi_{n-1}$. Let $(R_n)_n$ be a sequence of non-constant rational fractions. For $i \in \mathbf{N}$, put

$$f_{i+1} : (x, y) \mapsto (\xi_{i+1}x, yS_{i+1}(x)) \text{ with } S_{i+1}(x) = \frac{R_i(x^{q^i})}{R_i(\xi_1x^{q^i})} \frac{R_{i-1}(x^{q^{i-1}})}{R_{i-1}(\xi_2x^{q^{i-1}})} \cdots \frac{R_1(x)}{R_1(\xi_ix)}.$$

We have $f_{i+1}^q = f_i$ for all $i \in \mathbf{N}^*$ so that the group G_1 generated by all the f_i is an infinite torsion abelian group. Let $T_i(x) = R_i(x^{q^i}) \cdots R_1(x^q)$. The conjugation by $(x, y) \mapsto (x, yT_i(x))$ sends the group generated by f_1, \dots, f_i into the cyclic group $\{(x, y) \mapsto (\xi_i^j x, y) | j = 0, 1, \dots, q^i - 1\}$ whose elements are elliptic. The product of G_1 with $\{(x, y) \mapsto (x, \beta y) | \beta \in \mathbf{K}^*\}$ is an abelian subgroup of $\text{Cr}_2(\mathbf{K})$.

Suppose by contradiction that up to conjugation G_1 is a group of automorphisms of a projective rational surface X . Since the degree of f_i tends to infinity when i tends to infinity, the f_i s are not contained in the identity component $\text{Aut}^0(X)$ for i sufficiently large. Note that the quotient group $\text{Aut}(X)/\text{Aut}^0(X)$ acts on the Néron-Severi group $\text{NS}(X, \mathbf{Z})$ with finite kernel (cf. [Har87], [CD12a] 1.1) and preserves the intersection form which has signature $(1, \dim \text{NS}(X, \mathbf{Z}) - 1)$ by Hodge Index Theorem. Therefore infinitely many f_i acts non trivially by isometry on $\text{NS}(X, \mathbf{Z})$. Note that $\text{Aut}^0(X)$ is a finite group because $\text{Aut}(X)/\text{Aut}^0(X)$ is infinite (cf. [Har87]). Since the order of the torsion element f_i tends to infinity, this implies that the order of the action of f_i on $\text{NS}(X, \mathbf{Z})$ tends to infinity. Contradiction because an isometry of $\text{NS}(X, \mathbf{Z})$ cannot have arbitrary high torsion order (the quadratic form is positive definite on the orthogonal of an eigenvector).

Example 4.3 We can give an additive version of Example 4.2. Let $(\xi_n)_n$ be as in Example 4.2. Let $(R_n)_n$ be a sequence of rational fractions whose degrees tend to infinity. For $i \in \mathbf{N}$, put

$$f_{i+1} : (x, y) \mapsto (\xi_{i+1}x, y + S_{i+1}(x))$$

with

$$S_{i+1}(x) = R_i(x^{q^i}) - R_i(\xi_1x^{q^i}) + R_{i-1}(x^{q^{i-1}}) - R_{i-1}(\xi_2x^{q^{i-1}}) + \cdots + R_1(x) - R_1(\xi_ix).$$

Let G_1 be the group generated by all the f_i . The product of G_1 with $\{(x, y) \mapsto (x, y + v) | v \in \mathbf{K}\}$ is an abelian subgroup of $\text{Cr}_2(\mathbf{K})$. Again G_1 can not be conjugate to an automorphism group.

Proof (of Theorem 4.1) Let G be a maximal abelian subgroup of $\text{Cr}_2(\mathbf{K})$. Note that if f is a non-trivial element of G , then G is the maximal abelian subgroup of $\text{Cent}(f)$.

If G contains a loxodromic element f , then G is included in $\text{Cent}(f)$ and is virtually the cyclic group generated by f by Theorem 2.1; this corresponds to the last case of the above statement. If G contains a Halphen twist, then by Theorem 2.2 it is virtually

a free abelian group of rank ≤ 8 which preserves fiberwise an elliptic fibration; this corresponds to the seventh case.

Assume that G contains a base-wandering Jonquières twist f . Theorem 3.2 says that $\text{Cent}(f)$ is virtually isomorphic to $\mathbf{K}^* \times \mathbf{Z}$, $\mathbf{K} \times \mathbf{Z}$ or \mathbf{Z} . Thus the same is true for G . This corresponds to the fifth and the sixth case.

Assume that G contains a non-base-wandering Jonquières twist f . Theorem 2.11 says that $\text{Cent}(f)$ is virtually isomorphic to an abelian subgroup of $\text{PGL}_2(\mathbf{K}(x))$, so the same is true for G . This is the fourth case.

In the rest of the proof we assume that G contains only elliptic elements. Note that G is not necessarily conjugate to a group of automorphisms.

Assume that G contains an element $f : (x, y) \mapsto (\alpha x, y + 1)$ with $\alpha \in \mathbf{K}^*$. By Theorem 2.7 we have

$$\text{Cent}(f) = \{(x, y) \mapsto (\eta(x), y + R(x)) \mid \eta \in \text{PGL}_2(\mathbf{K}), \eta(\alpha x) = \alpha \eta(x), R \in \mathbf{K}(x), R(\alpha x) = R(x)\}.$$

In α has infinite order, then $G = \text{Cent}(f) = \{(x, y) \mapsto (\gamma x, y + v) \mid \gamma \in \mathbf{K}^*, v \in \mathbf{K}\}$ and we are in the first case. Therefore we can and will assume that α has finite order d so that f^d is $(x, y) \mapsto (x, y + d)$. Assume in this paragraph that G has an element g with an infinite action on the base of the rational fibration $(x, y) \mapsto x$. If the action of g on the base is conjugate to $x \mapsto \beta x$ with $\beta \in \mathbf{K}^*$, then up to conjugation in $\text{Jonq}(\mathbf{K})$ we can suppose that g is just our initial element $f : (x, y) \mapsto (\alpha x, y + 1)$ (see Proposition 2.4), so that G is isomorphic to $\mathbf{K}^* \times \mathbf{K}$. Now consider the case where the action of g on the base is conjugate to $x' \mapsto x' + 1$. The parabolic element $x' \mapsto x' + 1$ does not commute with $x \mapsto \alpha x$ (note that x, x' are different coordinates) unless $\alpha = 1$. Thus $\alpha = 1$ under the existence of such a g and by choosing an appropriate coordinate x' , the two elements g and f are respectively $(x', y) \mapsto (x' + 1, y + R(x'))$ and $(x', y) \mapsto (x', y + d)$ where R is a polynomial by Lemma 3.12. We can conjugate g and f , simultaneously by $(x', y) \mapsto (x', y + S(x'))$ for some $S \in \mathbf{K}[X]$, to $(x', y) \mapsto (x' + 1, y)$ and $(x', y) \mapsto (x', y + d)$. Then we have

$$G = \text{Cent}(f^d) \cap \text{Cent}(g) = \{(x', y) \mapsto (x' + u, y + v) \mid u, v \in \mathbf{K}\}.$$

We are still under the hypothesis that G contains an element $f : (x, y) \mapsto (\alpha x, y + 1)$ with $\alpha \in \mathbf{K}^*$. Assume now that no element of G has an infinite action on the base of the rational fibration $(x, y) \mapsto x$. Then the description of $\text{Cent}(f)$ implies that G is a subgroup of

$$\{(x, y) \mapsto (\delta x, y + R(x)) \mid \delta \in \mathbf{K}^*, R \in \mathbf{K}(x)\}.$$

Consider the projection $\pi : G \rightarrow \text{PGL}_2(\mathbf{K})$ which records the action on the base. Denote by G_0 the kernel of π and by G_b the image of π . We identify G_b as a subgroup of the multiplicative group of roots of unity of \mathbf{K} . We want to prove that G_b is finite so that G is virtually contained in $\text{Jonq}_0(\mathbf{K}) = \text{PGL}_2(\mathbf{K}(x))$. Assume that G_b is an infinite subgroup of the group of roots of unity. We first claim that G_0 is isomorphic to \mathbf{K} . Let $h : (x, y) \mapsto (x, y + R(x))$, $R \in \mathbf{K}(x)$ be an element of G_0 and $g : (x, y) \mapsto (\beta x, y + S(x))$, $S \in \mathbf{K}(x)$ be an element of G . The commutation relation $f \circ g = g \circ f$ implies $R(x) = R(\beta x)$. Here β can be any element of the infinite group G_b . This

implies that R is constant, which proves the claim. Let $g : (x, y) \mapsto (\gamma x, y + R(x))$ be an element of G . Let d be the order of γ . The transformation g^d is

$$(x, y) \mapsto (x, y + R(x) + R(\gamma x) + \cdots + R(\gamma^{d-1}x))$$

As G_0 is \mathbf{K} , there exists $C \in \mathbf{K}$ such that $R(x) + \cdots + R(\gamma^{d-1}x) = C$. Define $R^* = R - C/d$. Then we have $R^*(x) + \cdots + R^*(\gamma^{d-1}x) = 0$. By additive Hilbert's Theorem 90, there exists $T \in \mathbf{K}(x)$ such that $R^*(x) = S(\gamma x) - S(x)$. Therefore $R(x) = S(\gamma x) - S(x) + C/d$. Note that $g^* : (x, y) \mapsto (\gamma x, y + R^*(x))$ commutes with every element of G once $(x, y) \mapsto (\gamma x, y + R^*(x))$ is so. By maximality g^* is also in G . Remark that g^* has the same order as γ . Therefore such g^* s form a subgroup of G isomorphic to G_b . Thus G is isomorphic to the product $G_0 \times G_b$ and we are in the third case.

Assume that G contains an element $f : (x, y) \mapsto (\alpha x, \beta y)$ where $\alpha, \beta \in \mathbf{K}^*$ and β has infinite order. If α also has infinite order, then Theorem 2.7 implies immediately that $G = \text{Cent}(f)$ is isomorphic to $\mathbf{K}^* \times \mathbf{K}^*$ and we are in the first case. Assume that α has finite order but G contains an element $f_1 : (x, y) \mapsto (\alpha_1 x, yR(x))$ where $R \in \mathbf{K}(x)$ and $\alpha_1 \in \mathbf{K}^*$ has infinite order. We can suppose that $\alpha = 1$ because $\text{Cent}(f) \subset \text{Cent}(f^k)$ for any k . We apply Theorem 3.6 to f_1 to get a surface X on which the conjugate of f_1 acts by automorphism. By Corollary 3.11 X is a Hirzebruch surface. In the proof of Theorem 3.6, to go from $\mathbb{P}^1 \times \mathbb{P}^1$ to X , we perform elementary transformations at the iterates of the indeterminacy points of f_1 . Since f commutes with f_1 and f preserves every fiber, any iterate of an indeterminacy point of f_1 is fixed by f . Hence f remains an automorphism on X . In other words we can simultaneously conjugate f, f_1 to $(x, y) \mapsto (x, \beta y)$ and $(x, y) \mapsto (\alpha_1 x, ry)$ with $r \in \mathbf{K}^*$. Thus, Theorem 2.7, when applied respectively to f and f_1 , shows that $G = \text{Cent}(f) \cap \text{Cent}(f_1)$ is isomorphic to the diagonal group $\mathbf{K}^* \times \mathbf{K}^*$. Hence we are in the first case.

According to the classification of normal forms of elliptic elements of infinite order (see Proposition 2.4), the only remaining case is the following: G contains an element $f : (x, y) \mapsto (\alpha x, \beta y)$ where $\alpha \in \mathbf{K}^*$ has finite order and $\beta \in \mathbf{K}^*$ has infinite order but G contains no elements $(x, y) \mapsto (\alpha_1 x, yR(x))$ with α_1 of infinite order. In this case $\text{Cent}(f)$ is a subgroup of the Jonquières group by Theorem 2.7. Denote by π the projection of G into $\text{PGL}_2(\mathbf{K})$. If $\pi(G)$ is finite then we are in the fourth case of Theorem 4.1. So we assume that $\pi(G)$ is infinite. Then $\pi(G)$ is isomorphic to an infinite subgroup of the group of roots of unity. We want to show that we are in the second case of Theorem 4.1. By Lemma 3.8, each element of G has the form $(x, y) \mapsto (\eta(x), y \frac{rS(x)}{S(\eta(x))})$ with $\eta \in \text{PGL}_2(\mathbf{K}), r \in \mathbf{K}^*, S \in \mathbf{K}(x)$. If $(x, y) \mapsto (\eta(x), y \frac{rS(x)}{S(\eta(x))})$ is an element of G for some r , then $(x, y) \mapsto (\eta(x), y \frac{S(x)}{S(\eta(x))})$ is also an element of G because it commutes with every other element. However the later has the same order in G as η in $\text{PGL}_2(\mathbf{K})$. This means that G has a subgroup isomorphic to $\pi(G)$, so that G is isomorphic to the product of this subgroup with the kernel of π . To finish the proof, it suffices to show that the kernel of π is $\{(x, y) \mapsto (x, \beta y) | \beta \in \mathbf{K}^*\}$. This is because $(x, y) \mapsto (x, \beta y)$ are the only possible elliptic elements by Lemma 3.8. \square

5 References

- [BB00] Lionel Bayle and Arnaud Beauville. Birational involutions of \mathbf{P}^2 . *Asian J. Math.*, 4(1):11–17, 2000. Kodaira’s issue.
- [BC16] Jérémy Blanc and Serge Cantat. Dynamical degrees of birational transformations of projective surfaces. *J. Amer. Math. Soc.*, 29(2):415–471, 2016.
- [BCW09] Christian Bonatti, Sylvain Crovisier, and Amie Wilkinson. The C^1 generic diffeomorphism has trivial centralizer. *Publ. Math. Inst. Hautes Études Sci.*, 109:185–244, 2009.
- [BD05] Eric Bedford and Jeffrey Diller. Energy and invariant measures for birational surface maps. *Duke Math. J.*, 128(2):331–368, 2005.
- [BD15] Jérémy Blanc and Julie Déserti. Degree growth of birational maps of the plane. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, 14(2):507–533, 2015.
- [Bea07] Arnaud Beauville. p -elementary subgroups of the Cremona group. *J. Algebra*, 314(2):553–564, 2007.
- [BF13] Jérémy Blanc and Jean-Philippe Furter. Topologies and structures of the Cremona groups. *Ann. of Math. (2)*, 178(3):1173–1198, 2013.
- [Bis04] Cinzia Bisi. On commuting polynomial automorphisms of \mathbb{C}^2 . *Publ. Mat.*, 48(1):227–239, 2004.
- [Bis08] Cinzia Bisi. On commuting polynomial automorphisms of \mathbb{C}^k , $k \geq 3$. *Math. Z.*, 258(4):875–891, 2008.
- [Bla07] Jérémy Blanc. Finite abelian subgroups of the Cremona group of the plane. *C. R. Math. Acad. Sci. Paris*, 344(1):21–26, 2007.
- [Bla11] Jérémy Blanc. Elements and cyclic subgroups of finite order of the Cremona group. *Comment. Math. Helv.*, 86(2):469–497, 2011.
- [Can10] Serge Cantat. Invariant hypersurfaces in holomorphic dynamics. *Math. Res. Lett.*, 17(5):833–841, 2010.
- [Can11] Serge Cantat. Sur les groupes de transformations birationnelles des surfaces. *Ann. of Math. (2)*, 174(1):299–340, 2011.
- [Can18] Serge Cantat. The Cremona group. In *Algebraic geometry: Salt Lake City 2015*, volume 97 of *Proc. Sympos. Pure Math.*, pages 101–142. Amer. Math. Soc., Providence, RI, 2018.
- [CD12a] Serge Cantat and Igor Dolgachev. Rational surfaces with a large group of automorphisms. *J. Amer. Math. Soc.*, 25(3):863–905, 2012.
- [CD12b] Dominique Cerveau and Julie Déserti. Centralisateurs dans le groupe de Jonquières. *Michigan Math. J.*, 61(4):763–783, 2012.

- [CGL] Serge Cantat, Vincent Guirardel, and Anne Lonjou. Generators of proper normal subgroups of the Cremona group. *preprint*.
- [CX18] Serge Cantat and Junyi Xie. Algebraic actions of discrete groups: the p -adic method. *Acta Math.*, 220(2):239–295, 2018.
- [Dem70] Michel Demazure. Sous-groupes algébriques de rang maximum du groupe de Cremona. *Ann. Sci. École Norm. Sup. (4)*, 3:507–588, 1970.
- [Dés06a] Julie Déserti. Groupe de Cremona et dynamique complexe: une approche de la conjecture de Zimmer. *Int. Math. Res. Not.*, pages Art. ID 71701, 27, 2006.
- [Dés06b] Julie Déserti. Sur les automorphismes du groupe de Cremona. *Compos. Math.*, 142(6):1459–1478, 2006.
- [Dés08] Julie Déserti. Expériences sur certaines transformations birationnelles quadratiques. *Nonlinearity*, 21(6):1367–1383, 2008.
- [DF01] Jeffrey Diller and Charles Favre. Dynamics of bimeromorphic maps of surfaces. *Amer. J. Math.*, 123(6):1135–1169, 2001.
- [DI09] Igor V. Dolgachev and Vasilii A. Iskovskikh. Finite subgroups of the plane Cremona group. In *Algebra, arithmetic, and geometry: in honor of Yu. I. Manin. Vol. I*, volume 269 of *Progr. Math.*, pages 443–548. Birkhäuser Boston, Inc., Boston, MA, 2009.
- [É81] Jean Écalle. *Les fonctions résurgentes. Tome II*, volume 6 of *Publications Mathématiques d’Orsay 81 [Mathematical Publications of Orsay 81]*. Université de Paris-Sud, Département de Mathématique, Orsay, 1981. Les fonctions résurgentes appliquées à l’itération. [Resurgent functions applied to iteration].
- [Enr93] Federigo Enriques. Sui gruppi continui di trasformazioni cremoniane nel piano. *Rom. Acc. L. Rend.*, (5) II₁:468–473, 1893.
- [Fav10] Charles Favre. Le groupe de Cremona et ses sous-groupes de type fini. *Astérisque*, Volume 2008/2009.(332):Exp.No. 998, vii, 11–43, 2010. Séminaire Bourbaki. Exposés 997–1011.
- [FM89] Shmuel Friedland and John Milnor. Dynamical properties of plane polynomial automorphisms. *Ergodic Theory Dynam. Systems*, 9(1):67–99, 1989.
- [Giz80] Marat K. Gizatullin. Rational G -surfaces. *Izv. Akad. Nauk SSSR Ser. Mat.*, 44(1):110–144, 239, 1980.
- [Har87] Brian Harbourne. Rational surfaces with infinite automorphism group and no antipluricanonical curve. *Proc. Amer. Math. Soc.*, 99(3):409–414, 1987.
- [Isk79] Vasilii A. Iskovskikh. Minimal models of rational surfaces over arbitrary fields. *Izv. Akad. Nauk SSSR Ser. Mat.*, 43(1):19–43, 237, 1979.

- [Lam01] Stéphane Lamy. L’alternative de Tits pour $\text{Aut}[\mathbb{C}^2]$. *J. Algebra*, 239(2):413–437, 2001.
- [LU18] Anne Lonjou and Christian Urech. Actions of cremona groups on $\text{CAT}(0)$ cube complexes. *preprint*, <https://arxiv.org/pdf/2001.00783.pdf>, 2018.
- [Man67] Juri I. Manin. Rational surfaces over perfect fields. II. *Mat. Sb. (N.S.)*, 72 (114):161–192, 1967.
- [Mar71] Masaki Maruyama. On automorphism groups of ruled surfaces. *J. Math. Kyoto Univ.*, 11:89–112, 1971.
- [Ser10] Jean-Pierre Serre. Le groupe de Cremona et ses sous-groupes finis. *Astérisque*, Volume 2008/2009.(332):Exp. No. 1000, vii, 75–100, 2010. Séminaire Bourbaki. Exposés 997–1011.
- [Ure] Christian Urech. Subgroups of elliptic elements of the cremona group. *J. Reine Angew. Math.*, to appear.
- [Xie15] Junyi Xie. Periodic points of birational transformations on projective surfaces. *Duke Math. J.*, 164(5):903–932, 2015.

ShengYuan Zhao
 Institut de Recherche Mathématique de Rennes
 Université de Rennes 1
 263 avenue du Général Leclerc, CS 74205
 F-35042 RENNES Cédex

Institute for Mathematical Sciences
 Stony Brook University
 Stony Brook NY 11794-3660, USA
e-mail: shengyuan.zhao@stonybrook.edu