# On instabilities of stationary scalar field configurations supported by reflecting compact stars

Yan Peng<sup>1\*</sup>
<sup>1</sup> School of Mathematical Sciences, Qufu Normal University, Qufu, Shandong 273165, China

### Abstract

We study instabilities of the system composed of stationary scalar fields and asymptotically flat horizonless reflecting compact stars. In the probe limit, we obtain bounds on the scalar field frequency. Below this bound, stationary hairy stars are expected to suffer from nonlinear instabilities under massless field perturbations. In other words, we prove that stationary scalar hairy stars are unstable for scalar fields with small frequency.

PACS numbers: 11.25.Tq, 04.70.Bw, 74.20.-z

<sup>\*</sup> yanpengphy@163.com

### I. INTRODUCTION

There is accumulating evidence that fundamental scalar fields may exist in nature [1]. Theoretically, the scalar field can be either static or stationary. The famous black hole no scalar hair theorems state an interesting property that static scalar hair usually cannot exist in the asymptotically flat black hole backgrounds, see references [2]-[10] and reviews [11, 12]. In contrast, it has recently been shown that rotating black holes allow the existence of stationary massive scalar field hairs [13]-[22]. Moreover, with scalar fields confined in a box, static hairy black hole solutions were constructed in [23–25] and their dynamical formation was studied in [26]. We should mention that the instability properties of hairy black holes were investigated [27, 28].

Interestingly, no static scalar hair behavior also appears in horizonless neutral compact object spacetimes. Hod firstly proved no static scalar hair theorems for asymptotically flat horizonless neutral compact stars with reflective surface boundary conditions [29]. In fact, being filled with matter, it is natural to assume that the compact star surface would have reflective properties [30]. When considering nonminimal couplings between scalar fields and curvature, static scalar hairs also cannot exist outside asymptotically flat horizonless neutral reflecting compact stars [31]. In the asymptotically dS background, no static scalar hair theorems still hold for the horizonless neutral reflecting compact star [32]. So the no static scalar hair behavior is a very general property in the spacetime of horizonless neutral reflecting compact stars.

On the other side, null circular geodesics may exist in the compact object spacetime [33–37]. For horizonless compact objects, if null circular geodesics exist, in general, there will be pairs of the null circular geodesics and the innermost null circular geodesic is stable [38, 39]. So horizonless compact objects with null circular geodesics is expected to be unstable since massless fields can pile up on the innermost stable null geodesic [40, 41]. One known way to evade the no hair theorem for horizonless reflecting compact star is to consider a charged background [42]-[52]. The null circular geodesic was used to study instabilities of charged horizonless static scalar hairy compact stars [53, 54]. Recently, Hod provided another interesting way to evade the no hair theorem for horizonless reflecting compact stars, which is considering stationary scalar field hairs [55]. So it is interesting to disclose the (in)stability of such horizonless stationary scalar hairy reflecting compact stars through properties of null circular geodesics.

In the following, we study the (in)stability of horizonless stationary scalar hairy reflecting compact stars in the asymptotically flat gravity. We analytically obtain bounds for the scalar field frequency. Below this bound, stationary hairy stars are unstable. We give conclusions at the last section.

### II. BOUNDS FOR THE FREQUENCY OF STATIONARY SCALAR FIELDS

We study a gravity model of stationary scalar fields linearly coupled to horizonless reflecting compact stars.

And the matter field Lagrange density is

$$\mathcal{L} = -|\nabla_{\mu}\Psi|^2 - m^2\Psi^2,\tag{1}$$

where  $\Psi$  is the scalar field with mass m.

The line element of the spherically symmetric compact star reads [33]

$$ds^{2} = -fe^{-\chi}dt^{2} + \frac{dr^{2}}{f} + r^{2}(d\theta^{2} + \sin^{2}\theta d\phi^{2}).$$
 (2)

Here f and  $\chi$  are metric functions depending on the radial coordinate r. We define the star radius as  $r_s$ . Since the spacetime is horizonless, there is f(r) > 0 for all  $r \ge r_s$ .  $\theta$  and  $\phi$  are angular coordinates.

The equations of metrics and scalar fields are [53–59]

$$f' = -8\pi r \rho + (1 - f)/r,\tag{3}$$

$$\chi' = -8\pi r(\rho + p)/f,\tag{4}$$

$$(\nabla^{\nu}\nabla_{\nu} - m^2)\Psi = 0 \tag{5}$$

with  $\rho = -T_t^t$ ,  $p = T_r^r$  as the matter field energy density and the radial pressure respectively.

In this work, we neglect scalar fields' backreaction on the background. So there is Schwarzschild type solution  $\chi(r) = 0$  and  $f(r) = 1 - \frac{2M}{r}$  with M as the star mass. We take stationary scalar fields in the form

$$\Psi(t,r) = e^{-i\omega t}\psi(r),\tag{6}$$

where  $\omega$  is the frequency.

And the scalar field equation is

$$\psi'' + (\frac{2}{r} + \frac{f'}{f})\psi' + (\frac{\omega^2}{f^2} - \frac{m^2}{f})\psi = 0$$
 (7)

with  $f = 1 - \frac{2M}{r}$  [60–64].

At the star surface  $r_s$ , we impose the scalar reflecting condition. At the infinity, the general asymptotic behavior is  $\psi \sim A \cdot \frac{1}{r} e^{-\sqrt{m^2 - \omega^2}r} + B \cdot \frac{1}{r} e^{\sqrt{m^2 - \omega^2}r}$ , where A and B are integral constants. Boundness of the scalar field at infinity requires B = 0 [65]. So boundary conditions are

$$\psi(r_s) = 0, \qquad \qquad \psi(\infty) = 0. \tag{8}$$

It was shown that horizonless compact objects with null circular geodesics usually have pairs of null circular geodesics and the innermost null circular geodesic is stable [38]. As massless fields tend to pile up on the stable null circular geodesic, horizonless compact stars with null circular geodesics are unstable to massless field perturbations [40, 41]. So we can study the instability of horizonless stationary hairy configurations by examining whether there is null circular geodesic in the spacetime. In the probe limit, there is exterior null circular geodesic when the would-be null circular geodesic radius is above compact star surface. In the following analysis, we will obtain the instability condition by imposing exterior null circular geodesic radii above upper bounds of hairy star radii.

We introduce a new function  $\tilde{\psi} = \sqrt{r}\psi$ . According to the scalar field equation (7), the equation of the function  $\tilde{\psi}$  can be expressed as

$$r^{2}\tilde{\psi}'' + \left(r + \frac{r^{2}f'}{f}\right)\tilde{\psi}' + \left(-\frac{1}{4} - \frac{rf'}{2f} + \frac{\omega^{2}r^{2}}{f^{2}} - \frac{m^{2}r^{2}}{f}\right)\tilde{\psi} = 0$$

$$(9)$$

with  $f = 1 - \frac{2M}{r}$ .

Boundary conditions of the function  $\tilde{\psi}$  are

$$\tilde{\psi}(r_s) = 0, \qquad \tilde{\psi}(\infty) = 0.$$
 (10)

According to (10), at least one extremum point  $r = r_{peak}$  of the function  $\tilde{\psi}$  exists between the surface  $r_s$  and the infinity. At this extremum point, there are the following relations

$$\{\tilde{\psi}' = 0 \quad and \quad \tilde{\psi}\tilde{\psi}'' \leqslant 0\} \quad for \quad r = r_{peak}.$$
 (11)

Relations (9) and (11) yield the inequality

$$-\frac{1}{4} - \frac{rf'}{2f} + \frac{\omega^2 r^2}{f^2} - \frac{m^2 r^2}{f} \geqslant 0 \quad for \quad r = r_{peak}.$$
 (12)

It can be transformed into

$$m^2 r^2 f(r) \leqslant \omega^2 r^2 - \frac{rff'}{2} - \frac{1}{4} f^2 \quad for \quad r = r_{peak}.$$
 (13)

The regular condition of the spacetime requires  $r_s \ge 2M$ , otherwise there is a horizon at r = 2M above the star surface. With  $r_s \ge 2M$ , there are relations

$$r \geqslant r_s \geqslant 2M,$$
 (14)

$$f = 1 - \frac{2M}{r} = \frac{1}{r}(r - 2M) \geqslant 0,\tag{15}$$

$$rf' = r(\frac{2M}{r^2}) = \frac{2M}{r} \geqslant 0. \tag{16}$$

According to (13-16), the following inequality holds

$$m^2 r^2 f(r) \leqslant \omega^2 r^2 \quad for \quad r = r_{peak}.$$
 (17)

The relation (17) yields the inequality

$$m^2r^2(1-\frac{2M}{r}) \leqslant \omega^2r^2 \quad for \quad r = r_{peak}.$$
 (18)

We can transform (18) into

$$m^2 - \omega^2 \leqslant \frac{2m^2M}{r_{peak}}. (19)$$

From (19), we obtain bounds on the extremum point  $r = r_{peak}$  in the form

$$r_{peak} \leqslant \frac{2m^2M}{m^2 - \omega^2}. (20)$$

It is also the bound on hairy star radii as

$$r_s \leqslant r_{peak} \leqslant \frac{2m^2M}{m^2 - \omega^2}. (21)$$

Following approaches in [66, 67], we derive the characteristic relation for null circular geodesics. The Lagrangian describing geodesics is

$$2\mathcal{L} = -e^{-\chi}f\dot{t}^2 + \frac{1}{f}\dot{r}^2 + r^2\dot{\phi}^2,$$
 (22)

where a dot represents a derivative with respect to the affine parameter along the geodesic.

The Lagrangian is independent of t and  $\phi$ . This implies that the existence of two constants of motion labeled as E and L. According to (22), the generalized momenta can be expressed as

$$p_t = -e^{-\chi} f \dot{t} = -E = const, \tag{23}$$

$$p_{\phi} = r^2 \dot{\phi} = L = const, \tag{24}$$

$$p_r = \frac{1}{f}\dot{r} \ . \tag{25}$$

The Hamiltonian of the system is  $\mathcal{H} = p_t \dot{t} + p_r \dot{r} + p_\phi \dot{\phi} - \mathcal{L}$ , which implies

$$2\mathcal{H} = -E\dot{t} + L\dot{\phi} + \frac{1}{f}\dot{r}^2 = \delta = const. \tag{26}$$

For timelike geodesics, there is  $\delta = 1$  and in this paper with null geodesics, we take  $\delta = 0$ .

From (26), we obtain the expression

$$\dot{r}^2 = f[E\dot{t} - L\dot{\phi}]. \tag{27}$$

With relations (23) and (24), we get expressions for  $\dot{t}$  and  $\dot{\phi}$  in the form

$$\dot{t} = \frac{e^{\chi} E}{f}, \qquad \dot{\phi} = \frac{L}{r^2}. \tag{28}$$

Putting (28) into (27), we arrive at the relation

$$\dot{r}^2 = f\left[\frac{e^{\chi}E^2}{f} - \frac{L^2}{r^2}\right] = e^{\chi}E^2 - \frac{L^2f}{r^2}.$$
 (29)

At the null circular geodesic, there are relations  $\dot{r}^2 = 0$  and  $(\dot{r}^2)' = 0$  [67]. The equation  $\dot{r}^2 = 0$  yields

$$E^2 = \frac{L^2 f}{r^2 e^{\chi}}. (30)$$

The requirement  $(\dot{r}^2)' = 0$  yields

$$(\dot{r}^2)' = \chi' e^{\chi} E^2 + \frac{2L^2 f}{r^3} - \frac{L^2 f'}{r^2} = 0.$$
(31)

According to (30) and (31), the characteristic equation of null circular geodesics is

$$2f(r_{\gamma}) + r_{\gamma}[\chi'(r_{\gamma})f(r_{\gamma}) - f'(r_{\gamma})] = 0.$$
(32)

Without backreaction of scalar fields, the exterior spacetime of the compact star can be described by  $\chi = 0$  and  $f = 1 - \frac{2M}{r}$ . The exterior null circular geodesic radius is

$$r_{\gamma} = 3M. \tag{33}$$

In order to obtain the instability condition, we impose that the radius of the would-be outer null circular geodesics is above the hairy star radius bound (21), which leads to the existence of the exterior null circular geodesic  $r_{\gamma} = 3M$  [53, 54]. So the horizonless hairy star is unstable on condition that

$$r_{\gamma} = 3M \geqslant \frac{2m^2M}{m^2 - \omega^2}.\tag{34}$$

From (34), we obtain bounds for the frequency of stationary scalar fields as

$$\frac{\omega^2}{m^2} \leqslant \frac{1}{3}.\tag{35}$$

It is known that the static ( $\omega = 0$ ) scalar field usually cannot exist outside horizonless neutral reflecting compact stars. In contrast, stationary ( $\omega \neq 0$ ) scalar fields can condense outside the horizonless neutral reflecting compact star. In this work, we further show that horizonless neutral stationary hairy reflecting compact stars are unstable for frequency below the bound (35).

#### III. CONCLUSIONS

We investigated the model of stationary scalar fields linearly coupled to asymptotically flat horizonless neutral compact stars with reflecting boundary conditions. We studied instabilities of scalar hairy reflecting stars through properties of null circular geodesics. We analytically obtained bounds on the frequency of stationary scalar field as  $\frac{\omega^2}{m^2} \leq \frac{1}{3}$ , where  $\omega$  and m are the frequency and mass of the scalar field respectively. Below this bound, stationary scalar hairy configurations supported by horizonless reflecting compact stars are expected to be dynamically unstable under perturbations of massless fields. That is to say the horizonless stationary scalar hairy compact reflecting star is unstable for scalar fields with small frequency.

## Acknowledgments

This work was supported by the Shandong Provincial Natural Science Foundation of China under Grant No. ZR2018QA008. This work was also supported by a grant from Qufu Normal University of China under Grant No. xkjjc201906.

<sup>[1]</sup> Franz E. Schunck, Eckehard W. Mielke, TOPICAL REVIEW: General relativistic boson stars, Class. Quant. Grav. 20(2003)R301-R356.

<sup>[2]</sup> J. D. Bekenstein, Transcendence of the law of baryon-number conservation in black hole physics, Phys. Rev. Lett. 28(1972)452.

<sup>[3]</sup> J. E. Chase, Commun. Math. Phys. 19(1970)276; C. Teitelboim, Lett. Nuovo Cimento 3(1972) 326; J. D. Bekenstein, Physics Today 33(1980)24.

<sup>[4]</sup> R. Ruffini and J. A. Wheeler, Introducing the black hole, Phys. Today 24(1971)30.

 <sup>[5]</sup> Sourav Bhattacharya, Amitabha Lahiri, Black-hole no-hair theorems for a positive cosmological constant, Phys. Rev. Lett. 99(2007)201101.

<sup>[6]</sup> Sourav Bhattacharya, Amitabha Lahiri, No hair theorems for stationary axisymmetric black holes, Phys. Rev. D 83(2011)124017.

<sup>[7]</sup> Norman Gürlebeck, No-hair theorem for Black Holes in Astrophysical Environments, Phys. Rev. Lett. 114(2015)no.15,151102.

<sup>[8]</sup> Robert Benkel, Thomas P. Sotiriou, Helvi Witek, Dynamical scalar hair formation around a Schwarzschild black hole, Phys.Rev. D 94(2016)no.12,121503.

- [9] Joseph Sultana, Demosthenes Kazanas, A no-hair theorem for spherically symmetric black holes in  $\mathbb{R}^2$  gravity, Gen.Rel.Grav. 50(2018)no.11,137.
- [10] Carlos Barceló, Raúl Carballo-Rubio, Stefano Liberati, Generalized no-hair theorems without horizons, Class. Quant.Grav. 36(2019)no.13,13.
- [11] J. D. Bekenstein, Black hole hair: 25-years after, arXiv:gr-qc/9605059.
- [12] Carlos A. R. Herdeiro, Eugen Radu, Asymptotically flat black holes with scalar hair: a review, Int. J. Mod. Phys. D 24(2015)09,1542014.
- [13] S. Hod, Stationary Scalar Clouds Around Rotating Black Holes, Phys. Rev. D 86(2012)104026.
- [14] C. A. R. Herdeiro and E. Radu, Kerr black holes with scalar hair, Phys. Rev. Lett. 112(2014)221101.
- [15] P. V. P. Cunha, C. A. R. Herdeiro, Shadows of Kerr black holes with scalar hair, E. Radu, and H. F. Rúnarsson, Phys. Rev. Lett. 115(2015)211102.
- [16] Y. Brihaye, C. Herdeiro, and E. Radu, Inside black holes with synchronized hair, Phys. Lett. B 760(2016)279.
- [17] Carlos Herdeiro, Eugen Radu, Construction and physical properties of Kerr black holes with scalar hair, Class. Quant. Grav. 32(2015)no.14,144001.
- [18] Shahar Hod, Kerr-Newman black holes with stationary charged scalar clouds, Phys. Rev. D 90(2014)no.2,024051.
- [19] Shahar Hod, The large-mass limit of cloudy black holes, Class. Quant. Grav. 32(2015) no.13,134002.
- [20] Yan Peng, The extreme orbital period in scalar hairy kerr black holes, Phys. Lett. B 792(2019)1-3.
- [21] Jorge F.M. Delgado, Carlos A.R. Herdeiro, Eugen Radu, Kerr black holes with synchronised scalar hair and higher azimuthal harmonic index, Phys. Lett. B 792(2019)436-444.
- [22] Yong-Qiang Wang, Yu-Xiao Liu, Shao-Wen Wei, Excited Kerr black holes with scalar hair, Phys. Rev. D 99(2019)064036.
- [23] S. R. Dolan, S. Ponglertsakul, E. Winstanley, Stability of black holes in Einstein-charged scalar field theory in a cavity, Phys. Rev. D 92(2015)no.12,124047.
- [24] Pallab Basu, Chethan Krishnan, P. N. Bala Subramanian, Hairy Black Holes in a Box, JHEP 1611(2016)041.
- [25] Oscar J. C. Dias, Ramon Masachs, Evading no-hair theorems: hairy black holes in a Minkowski box, Phys. Rev. D 97(2018)no.12,124030.
- [26] N. Sanchis-Gual, J. C. Degollado, P. J. Montero, J. A. Font and C. Herdeiro, Explosion and Final State of an Unstable Reissner-Nordström Black Hole, Phys. Rev. Lett. 116(2016)no.14,141101.
- [27] Bogdan Ganchev, Jorge E. Santos, Scalar Hairy Black Holes in Four Dimensions are Unstable, Phys. Rev. Lett. 120(2018)no.17,171101.
- [28] J. C. Degollado, C. A. R. Herdeiro and E. Radu, Effective stability against superradiance of Kerr black holes with synchronised hair, Phys. Lett. B 781(2018)651-655.
- [29] S.Hod, No-scalar-hair theorem for spherically symmetric reflecting stars, Phys. Rev. D 94(2016)104073.
- [30] Carlos Barceló, Raúl Carballo-Rubio, Luis J. Garay, Gravitational wave echoes from macroscopic quantum gravity effects, JHEP 1705(2017)054.
- [31] S.Hod, No nonminimally coupled massless scalar hair for spherically symmetric neutral reflecting stars, Phys. Rev. D 96(2017)024019.
- [32] Srijit Bhattacharjee, Sudipta Sarkar, No-hair theorems for a static and stationary reflecting star, Phys. Rev. D 95(2017)084027.
- [33] S. Chandrasekhar, The Mathematical Theory of Black Holes, (Oxford University Press, New York, 1983).
- [34] S. L. Shapiro and S. A. Teukolsky, Black Holes, White Dwarfs and Neutron Stars: The Physics of Compact Objects, 1st ed. (Wiley-Interscience, 1983).
- [35] S. Hod, Extremal Kerr-Newman black holes with extremely short charged scalar hair, Phys. Lett. B 751(2015)177.
- [36] V. Cardoso, L. C. B. Crispino, C. F. B. Macedo, H. Okawa, and P. Pani, Light rings as observational evidence for event horizons: long-lived modes, ergoregions and nonlinear instabilities of ultracompact objects, Phys. Rev. D 90(2014)044069.
- [37] P. Grandclément, Light rings and light points of boson stars, Phys. Rev. D 95(2017)084011.
- [38] P. V. P. Cunha, E. Berti, and C. A. R. Herdeiro, Light-Ring Stability for Ultracompact Objects, Phys. Rev. Lett. 119 (2017)251102.
- [39] S. Hod, On the number of light rings in curved spacetimes of ultra-compact objects, Phys. Lett. B 776(2018)1.
- [40] J. Keir, Slowly decaying waves on spherically symmetric spacetimes and ultracompact neutron stars, Classical Quantum Gravity 33(2016)135009.
- [41] S. Hod, Upper bound on the radii of black-hole photonspheres, Phys. Lett. B 727(2013)345.
- [42] S.Hod, Charged massive scalar field configurations supported by a spherically symmetric charged reflecting shell, Phys. Lett. B 763(2016)275.
- [43] S.Hod, Marginally bound resonances of charged massive scalar fields inthebackground of a charged reflecting shell, Phys. Lett. B 768(2017)97-102.
- [44] Yan Peng, Bin Wang, Yunqi Liu, Scalar field condensation behaviors around reflecting shells in Anti-de Sitter spacetimes, Eur. Phys. J. C 78(2018) no.8,680.
- [45] S.Hod, Charged reflecting stars supporting charged massive scalar field configurations, Eur. Phys. J. C 78(2018)no.3,173.
- [46] Shahar Hod, Stationary bound-state scalar configurations supported by rapidly-spinning exotic compact objects, Phys. Lett. B 770(2017)186.
- [47] Shahar Hod, Ultra-spinning exotic compact objects supporting static massless scalar field configurations, Phys.

- Lett. B 774(2017)582.
- [48] Shahar Hod,Onset of superradiant instabilities in rotating spacetimes of exotic compact objects, JHEP 06(2017)132.
- [49] Yan Peng, Scalar field configurations supported by charged compact reflecting stars in a curved spacetime, Phys. Lett. B 780(2018)144-148.
- [50] Yan Peng, Scalar condensation behaviors around regular Neumann reflecting stars, Nucl. Phys. B 934(2018)459-465.
- [51] Yan Peng, Static scalar field condensation in regular asymptotically AdS reflecting star backgrounds, Phys. Lett. B 782(2018)717-722.
- [52] Yan Peng, Hair formation in the background of noncommutative reflecting stars, Nucl. Phys. B 938(2019)143-153.
- [53] S. Hod, Upper bound on the gravitational masses of stable spatially regular charged compact objects, Phys. Rev. D 98(2018)064014.
- [54] Yan Peng, On instabilities of scalar hairy regular compact reflecting stars, JHEP 1810(2018)185.
- [55] Shahar Hod, Stationary bound-state massive scalar field configurations supported by spherically symmetric compact reflecting stars, Eur. Phys. J. C 77(2017)899.
- [56] Darío Núñez, Hernando Quevedo, Daniel Sudarsky, Black holes have no short hair, Phys. Rev. Lett. 76(1996)571-574.
- [57] S. Hod, hairy Black Holes and Null Circular Geodesics, Phys. Rev. D 84(2011)124030.
- [58] A.E. Mayo and J.D. Bekenstein, No hair for spherical black holes: Charged and nonminimally coupled scalar field with selfinteraction, Phys. Rev. D 54(1996)5059-5069.
- [59] H. Bondi, Anisotropic spheres in general relativity, Mon. Not. R. Astron. Soc. 259(1992)365.
- [60] M. Khodaei, H. Mohseni Sadjadi, No skyrmion hair for stationary spherically symmetric reflecting stars, Phys. Lett. B 797(2019)134922.
- [61] Peng Wang, Haitang Yang, Shuxuan Ying, Thermodynamics and phase transition of a Gauss-Bonnet black hole in a cavity, arXiv:1909.01275 [gr-qc].
- [62] Peng Wang, Houwen Wu, Haitang Yang, Thermodynamics and phase transition of a nonlinear electrodynamics black hole in a cavity, J. High Energy Phys. 07 (2019)002
- [63] Yan Peng, Studies of a general flat space/boson star transition model in a box through a language similar to holographic superconductors, JHEP 07(2017)042.
- [64] Yan Peng, Bin Wang, Yunqi Liu, On the thermodynamics of the black hole and hairy black hole transitions in the asymptotically flat spacetime with a box, Eur. Phys. J. C 78(2018)no.3,176.
- [65] Shahar Hod, No hair for spherically symmetric neutral black holes: Nonminimally coupled massive scalar fields, Phys. Rev. D 96(2017)124037.
- [66] V. Cardoso, A. S. Miranda, E. Berti, H. Witek and V. T. Zanchin, Geodesic stability, Lyapunov exponents and quasinormal modes, Phys. Rev. D 79(2009)064016.
- [67] Shahar Hod, Fastest way to circle a black hole, Phys. Rev. D 84(2011)104024.