

Bistability and Synchronization in Coupled Maps Models

Ricardo López-Ruiz

University of Zaragoza
rilopez@unizar.es

Several coupled maps models are sketched and reviewed in this short communication.

First, a discrete logistic type model that was proposed for the symbiotic interaction of two species [1, 2]. The coupling between the species depends on the size of the other species (x_n, y_n) at each time n and of a constant λ that it was called the *mutual benefit*:

$$\begin{aligned} x_{n+1} &= \lambda(3y_n + 1)x_n(1 - x_n) \\ y_{n+1} &= \lambda(3x_n + 1)y_n(1 - y_n). \end{aligned} \quad (1)$$

If the species are isolated, we recover the logistic model for each of them. For λ values where one isolated species would extinguish, the coexistence with the other species allows them to have an alternative state to survive, supporting the idea that the symbiosis between both species makes possible the *bistability* (Fig. 1). For increasing values of λ different dynamical regimes are obtained, from periodic oscillations of the populations, through quasiperiodicity up to a chaotic regime before the final collapse.

Second, if many of these symbiotic species, $i = 1, \dots, N$, evolve together (x_n^i, y_n^i) under the coupling action of a global mean field, X_n^N and Y_n^N , with strength ϵ ,

$$\begin{aligned} x_{n+1}^i &= f_{\lambda, N}(x_n^i, y_n^i) \\ y_{n+1}^i &= f_{\lambda, N}(y_n^i, x_n^i) \end{aligned} \quad (2)$$

with

$$f_{\lambda, N}(x_n^i, y_n^i) = \lambda \left[3 \left(\frac{y_n^i + \epsilon Y_n^N}{1 + \epsilon} \right) + 1 \right] x_n^i (1 - x_n^i)$$

and $X_n^N = \frac{1}{N} \sum_{i=1}^N x_n^i$, $Y_n^N = \frac{1}{N} \sum_{i=1}^N y_n^i$, then the synchronization phenomenon of many oscillators appears, not only under the effect of a strong coupling ϵ if not also under the effect of an increasing size N of the system. This is one of the first models appeared in the literature where the *synchronization* of many chaotic oscillators was reported [1]. Thus, it can be easily seen that when $\epsilon = 0$ the species (maps) are uncoupled and they evolve freely but when $\epsilon \rightarrow \infty$ the maps are totally synchronized and they evolve with the same dynamics all of them, in a periodic, quasiperiodic or chaotic state depending on the mutual benefit λ . But it is still more surprising the effect provoked by the size of the system consisting in that when N increases there is an induced synchronization in this many agents model, such as it was also reported in Ref. [1] and it is sketched in Fig. 2.




Figure 1: *Bistability between the fixed populations P_0 and P_4 and their basins of attraction for the map (1) when $\lambda = 0.8$.*

Third, if the species x_i interact in a symbiotic way with the local mean field X_n^i coming from the nearest neighboring species then we can implement this model in different kind of networks. This implementation was done in Ref. [3] interpreting this model in a neuronal context as a naive brain model where the bistability of the global system mimics the sleeping and the awaking states of a primitive brain. These are the two global synchronized states, the turned off (x_θ) and the turned on (x_+) states of the network in question, that in the context of species would mean the extinction or the coexistence of the species in the ecosystem. This model is

$$x_{n+1}^i = \bar{p}_i x_n^i (1 - x_n^i) \quad (3)$$

where \bar{p}_i is the symbiotic parameter that depends on the local mean value, X_n^i , generated by the neighboring species, $\bar{p}_i = p (3X_n^i + 1)$, with $X_n^i = \frac{1}{N_i} \sum_{j=1}^{N_i} x_n^j$. N_i is the number of neighbors of the i th species (node i of the network), and p is the actual version of the mutual benefit λ of the model (1). The synchronized states of system (3) can be analytically found. The solutions are $x_\theta = 0$ and $x_+ \neq 0$. The first state x_θ is stable for $0 < p < 1$ and the state x_+ is stable for $p > 0.75$. Therefore the bistability between both states, ($x_n^i = x_\theta, x_n^i = x_+$), $\forall i$, is possible for $p > p_0 = 0.75$ in the case of many interacting species on a network. Let us observe at this point that the non-null state x_+ appears in the system as a kind of *explosive synchronization* mediated by a global saddle-node bifurcation in the network that takes place exactly at $p = p_0$, but in this case the appearance of the new synchronized state x_+ coexists with another non-noisy synchronized state x_θ . Also, different routes to enhance or switch on these both states and other similar multi-dimensional models were reported at that time in Refs. [3, 4].

Finally, some new results concerning this last model embedded in different topologies will be presented [5].

References

- [1] R. López-Ruiz, C. Pérez-García, Dynamics of maps with a global multiplicative coupling, *Chaos, Solitons and Fractals* 1 (1991) 511–528.
- [2] R. López-Ruiz, D. Fournier-Prunaret, Complex behaviour in a discrete logistic model for the symbiotic interaction of two species, *Math. Biosci. Eng.* 1 (2004) 307–324.
- [3] R. López-Ruiz, Y. Moreno, A.P. Fernández, S. Boccaletti, D.U. Hwang, Awakening and sleeping of a complex network, arXiv:nlin/0406053 (2004); *Neural Networks* 20 (2007) 102–108.
- [4] R. López-Ruiz, D. Fournier-Prunaret, The bistable brain: A neuronal model with symbiotic interactions, Ch. 10 in *Symbiosis: Evolution, Biology and Ecological Effects*, A.F. Camisao & C.C. Pedroso (Ed.), Nova Books (2012) 235–254.
- [5] R. López-Ruiz, M. Asís-Cánovas, to be published.




Figure 2: (a–c) Iterates and Fourier spectra of the globally coupled map (2) for $\lambda = 1.07$ when $\epsilon = 0.01$ and $N = 100$: (a) Iterates of a single element (the exterior attractor) and of the average (the central spot), (b) temporal Fourier spectrum of a single unit (arbitrary units), (c) temporal Fourier spectrum for the average (mean field). (d–f) The same as (a–c) when $N = 1000$. The synchronization effect mediated by the increasing size N is observed with the appearance of a period-2 oscillation (at $w = 1/2$) in the mean field, and by extension in the single element.