# DISCRETE POWER FUNCTIONS ON A HEXAGONAL LATTICE I: DERIVATION OF DEFINING EQUATIONS FROM THE SYMMETRY OF THE GARNIER SYSTEM IN TWO VARIABLES

NALINI JOSHI, KENJI KAJIWARA, TETSU MASUDA, AND NOBUTAKA NAKAZONO

ABSTRACT. The discrete power function on the hexagonal lattice proposed by Bobenko et al is considered, whose defining equations consist of three cross-ratio equations and a similarity constraint. We show that the defining equations are derived from the discrete symmetry of the Garnier system in two variables.

## 1. INTRODUCTION

The theory of discrete analytic functions based on circle packings and circle patterns was initiated by Thurston's idea of using circle packings to approximate the Riemann mapping [35]. Recently various applications of circle packings and circle patterns have been reported, for example, in probability and statistical physics [24, 31] based on general theory (see for example [32, 33] and references therein).

A theory of discrete analytic functions based on the circle patterns generated by discrete integrable systems has been developed [13]. Nijhoff et al. [26] showed that the cross-ratio equation (the discrete Schwarzian KdV equation) together with a certain similarity constraint can be regarded as a part of the Bäcklund transformations of the Painlevé VI equation. Bobenko found that this system of equations defines a discrete power function on a square lattice [8], and its various properties have been investigated in detail [3–5, 11, 12]. In this context, the cross-ratio equation can be regarded the nonlinear discretization of the Cauchy-Riemann relation. In our previous studies [7, 21], we constructed the explicit formulas for the discrete power functions on a square lattice in terms of the hypergeometric  $\tau$ -functions of the Painlevé VI equation, and clarified the complete characterization of the defining equations in relation to the symmetry of the Painlevé VI equation that is isomorphic to an extension of the affine Weyl group of type  $D_4^{(1)}$ .

Another type of discrete power function defined on a hexagonal lattice has been investigated in [6, 9, 10]. Consider the following system of difference equations consisting of three cross-ratio equations:

$$\frac{(f_{l_1,l_2,l_3} - f_{l_1+1,l_2,l_3})(f_{l_1+1,l_2+1,l_3} - f_{l_1,l_2+1,l_3})}{(f_{l_1+1,l_2,l_3} - f_{l_1,l_2+1,l_3})(f_{l_1,l_2+1,l_3} - f_{l_1,l_2,l_3})} = \frac{1}{x_1},$$

$$\frac{(f_{l_1,l_2,l_3} - f_{l_1,l_2+1,l_3})(f_{l_1,l_2+1,l_3+1} - f_{l_1,l_2,l_3+1})}{(f_{l_1,l_2+1,l_3} - f_{l_1,l_2,l_3+1})(f_{l_1,l_2,l_3+1} - f_{l_1,l_2,l_3})} = \frac{1}{x_2},$$

$$\frac{(f_{l_1,l_2,l_3} - f_{l_1,l_2,l_3+1})(f_{l_1+1,l_2,l_3+1} - f_{l_1+1,l_2,l_3})}{(f_{l_1,l_2,l_3+1} - f_{l_1+1,l_2,l_3} - f_{l_1,l_2,l_3+1})(f_{l_1+1,l_2,l_3+1} - f_{l_1,l_2,l_3})} = \frac{1}{x_3},$$

$$(1.1)$$

<sup>2010</sup> Mathematics Subject Classification. 14H70, 33E17, 34M55, 39A14.

Key words and phrases. discrete power function; Garnier system; affine Weyl group; consistent around a cube;  $\tau$  function.

NJ's ORCID ID is 0000-0001-7504-4444.

and a similarity constraint:

$$\begin{aligned} \alpha_{0}^{0} f_{l_{1},l_{2},l_{3}} &= (l_{1} - \alpha_{1}^{1}) \frac{(f_{l_{1}+1,l_{2},l_{3}} - f_{l_{1},l_{2},l_{3}})(f_{l_{1},l_{2},l_{3}} - f_{l_{1}-1,l_{2},l_{3}})}{f_{l_{1}+1,l_{2},l_{3}} - f_{l_{1}-1,l_{2},l_{3}}} \\ &+ (l_{2} - \alpha_{1}^{2}) \frac{(f_{l_{1},l_{2}+1,l_{3}} - f_{l_{1},l_{2},l_{3}})(f_{l_{1},l_{2},l_{3}} - f_{l_{1},l_{2}-1,l_{3}})}{f_{l_{1},l_{2}+1,l_{3}} - f_{l_{1},l_{2}-1,l_{3}}} \\ &+ (l_{3} - \alpha_{1}^{3}) \frac{(f_{l_{1},l_{2},l_{3}+1} - f_{l_{1},l_{2},l_{3}})(f_{l_{1},l_{2},l_{3}} - f_{l_{1},l_{2},l_{3}-1})}{f_{l_{1},l_{2},l_{3}+1} - f_{l_{1},l_{2},l_{3}-1}}. \end{aligned}$$
(1.2)

Here,  $l_1$ ,  $l_2$ ,  $l_3$  are integers, and  $x_i$ , i = 1, 2, 3,  $\alpha_1^j$ , j = 1, 2, 3, and  $\alpha_0^0$  are complex parameters satisfying  $x_1x_2x_3 = 1$ . A solution  $f_{l_1,l_2,l_3}$  of equations (1.1) and (1.2), which satisfies the initial conditions

$$f_{1,0,0} = 1, \quad f_{0,1,0} = e^{c(a_2 + a_3)\sqrt{-1}}, \quad f_{0,0,-1} = e^{ca_3\sqrt{-1}},$$
 (1.3)

with the identification  $x_j = e^{2a_j\sqrt{-1}} (a_j > 0, a_1 + a_2 + a_3 = \pi), j = 1, 2, 3, and <math>\alpha_0^0 = c/2$ (0 < c < 2),  $\alpha_1^1 = \alpha_1^2 = \alpha_1^3 = 0$ , is associated with a hexagonal circle pattern [6, Definition 1]. For a given set of integers  $(l_1, l_2, l_3)$  with  $l_1 + l_2 + l_3 = 0$ , the six points  $f_{l_1\pm 1, l_2, l_3}, f_{l_1, l_2\pm 1, l_3}$  and  $f_{l_1, l_2, l_3\pm 3}$  lie on a circle with the center at  $f_{l_1, l_2, l_3}$ , which gives rise to a circle pattern. See Figure 1.1 for an example with  $a_i = \pi/3$ , i = 1, 2, 3, and c = 3/2. We refer to such a solution as the discrete power function on a hexagonal lattice. Also, equations (1.1) and (1.2) are referred to as its defining equations.



Figure 1.1. A hexagonal circle pattern with  $a_1 = a_2 = a_3 = \pi/3$  and c = 3/2. Left: circle pattern with the coordinate lines. Right: configuration of the lattice points around  $f_{l_1,l_2,l_3}$  with  $l_1 + l_2 + l_3 = 0$ .

The main purpose of this paper is to show that the defining equations of the discrete power functions on a hexagonal lattice (1.1) and (1.2) arise from the discrete symmetry of the Garnier system in two variables (see [23, 34] for details of the Garnier system). In the next publication, we plan to construct the explicit formulas for the discrete power functions on the hexagonal lattice in terms of the hypergeometric  $\tau$  functions of the Garnier system.

This paper is organized as follows. In Section 2, we formulate the discrete symmetry of the Garnier system in two variables, and derive some bilinear relations for the so called  $\tau$ -variables. In Section 3, we derive a system of partial difference equations with some constraints from the bilinear relations. As a sub-system of the difference equations, we obtain the equations (1.1) and (1.2). Concluding remarks are then given in Section 4. It

is known that integrable difference equations on a hexagonal lattice can be reduced from equations on a multidimensional quadrilateral lattice [17]. In Appendix A, we consider a multi-dimensionally consistent system of partial difference equations, and show that it is reduced to the system of partial difference equations discussed in Section 3.

## 2. DISCRETE SYMMETRY OF THE GARNIER SYSTEM IN TWO VARIABLES

In this section, we first formulate the discrete symmetry of the Garnier system in two variables. Then we introduce  $\tau$ -variables through the Hamiltonians of the Garnier system in two variables and derive several bilinear relations among the  $\tau$ -variables.

2.1. Garnier system in two variables and its symmetry. The Garnier system in two variables [18, 20, 23] is equivalent to the Hamiltonian system

$$\frac{\partial q_j}{\partial t_i} = \frac{\partial H_i}{\partial p_j}, \quad \frac{\partial p_j}{\partial t_i} = -\frac{\partial H_i}{\partial q_j}, \quad (i, j = 1, 2), \tag{2.1}$$

with the Hamiltonians

$$t_{i}(t_{i}-1)H_{i} = q_{i}(q_{1}p_{1}+q_{2}p_{2}+\alpha)(q_{1}p_{1}+q_{2}p_{2}+\alpha+\kappa_{\infty}) + t_{i}p_{i}(q_{i}p_{i}-\theta_{i})$$

$$-\frac{t_{j}(t_{i}-1)}{t_{i}-t_{j}}(q_{j}p_{j}-\theta_{j})q_{i}p_{j} - \frac{t_{i}(t_{i}-1)}{t_{i}-t_{j}}(q_{i}p_{i}-\theta_{i})q_{j}p_{i}$$

$$-\frac{t_{i}(t_{j}-1)}{t_{j}-t_{i}}q_{j}p_{j}(q_{i}p_{i}-\theta_{i}) - \frac{t_{i}(t_{j}-1)}{t_{j}-t_{i}}q_{i}p_{i}(q_{j}p_{j}-\theta_{j})$$

$$-(t_{i}+1)(q_{i}p_{i}-\theta_{i})q_{i}p_{i} + (\kappa_{1}t_{i}+\kappa_{0}-1)q_{i}p_{i}, \quad (i,j) = (1,2), (2,1),$$

$$(2.2)$$

where  $\alpha = -\frac{1}{2}(\theta_1 + \theta_2 + \kappa_0 + \kappa_1 + \kappa_\infty - 1)$ . This system admits the following birational canonical transformations [20, 22]

$$\begin{aligned}
\sigma_{12} &: \theta_{1} \leftrightarrow \theta_{2}, \quad t_{1} \leftrightarrow t_{2}, \quad q_{1} \leftrightarrow q_{2}, \quad p_{1} \leftrightarrow p_{2}, \\
\sigma_{23} &: \theta_{2} \leftrightarrow \kappa_{0}, \quad t_{1} \mapsto \frac{t_{2} - t_{1}}{t_{2} - 1}, \quad t_{2} \mapsto \frac{t_{2}}{t_{2} - 1}, \\
q_{1} \mapsto \frac{t_{2} - t_{1}}{t_{1}(t_{2} - 1)}q_{1}, \quad q_{2} \mapsto \frac{t_{2}}{1 - t_{2}}(g_{t} - 1), \\
p_{1} \mapsto \frac{t_{2} - 1}{t_{2} - t_{1}}(t_{1}p_{1} - t_{2}p_{2}), \quad p_{2} \mapsto (1 - t_{2})p_{2}, \\
\sigma_{34} &: \kappa_{0} \leftrightarrow \kappa_{1}, \quad t_{i} \mapsto \frac{1}{t_{i}}, \quad q_{i} \mapsto \frac{q_{i}}{t_{i}}, \quad p_{i} \mapsto t_{i}p_{i}, \\
\sigma_{45} &: \kappa_{1} \leftrightarrow \kappa_{\infty}, \quad t_{i} \mapsto \frac{t_{i}}{t_{i} - 1}, \\
q_{i} \mapsto \frac{q_{i}}{g_{1} - 1}, \quad p_{i} \mapsto (g_{1} - 1)(p_{i} - \alpha - q_{1}p_{1} - q_{2}p_{2}), \\
g_{t} &= \frac{q_{1}}{t_{1}} + \frac{q_{2}}{t_{2}}, \quad g_{1} &= q_{1} + q_{2}.
\end{aligned}$$
(2.4)

where

Note that these transformations act on the parameters 
$$\theta = (\theta_1, \theta_2, \kappa_0, \kappa_1, \kappa_\infty)$$
 by transposing adjacent pairs, and generate the group isomorphic to the symmetric group  $\mathfrak{S}_5$ . The following mappings

$$r_{i} : \theta_{i} \mapsto -\theta_{i}, \quad p_{i} \mapsto p_{i} - \frac{\theta_{i}}{q_{i}} \quad (i = 1, 2),$$

$$r_{3} : \kappa_{0} \mapsto -\kappa_{0}, \quad p_{i} \mapsto p_{i} - \frac{\kappa_{0}}{t_{i}(g_{t} - 1)},$$

$$r_{4} : \kappa_{1} \mapsto -\kappa_{1}, \quad p_{i} \mapsto p_{i} - \frac{\kappa_{1}}{g_{1} - 1},$$

$$r_{5} : \kappa_{\infty} \mapsto -\kappa_{\infty},$$

$$(2.5)$$

are also birational canonical transformations for the Garnier system in two variables. These satisfy the fundamental relations

$$r_i^2 = 1$$
  $(i = 1, ..., 5),$   $r_i r_j = r_j r_i$   $(i \neq j),$   $\sigma r_i \sigma^{-1} = r_{\sigma(i)}$   $(\sigma \in \mathfrak{S}_5).$  (2.6)

Further there exists the transformation [36]

$$r_{34} : \theta_i \mapsto -\theta_i (i = 1, 2), \quad \kappa_0 \mapsto -\kappa_0 + 1, \quad \kappa_1 \mapsto -\kappa_1 + 1, \quad \kappa_\infty \mapsto -\kappa_\infty,$$

$$q_i \mapsto \frac{t_i p_i (q_i p_i - \theta_i)}{(q_1 p_1 + q_2 p_2 + \alpha)(q_1 p_1 + q_2 p_2 + \alpha + \kappa_\infty)}, \quad q_i p_i \mapsto -q_i p_i.$$

$$(2.7)$$

We have the fundamental relations

$$r_{34}^2 = 1, \qquad r_i r_{34} = r_{34} r_i \quad (i \neq 3, 4),$$
 (2.8)

and

$$r_{34}\sigma_{34} = \sigma_{34}r_{34}, \quad r_{34}\sigma_{ab} = \sigma_{ab}r_{34} (a, b \neq 3, 4), \tag{2.9}$$

where  $\sigma_{ab} = \sigma_{ba} \in \mathfrak{S}_5$  is a transposition for mutually distinct indices  $a, b \in \{1, \dots, 5\}$ .

**Remark 2.1.** The group generated by  $r_i$  (i = 1, ..., 5),  $\sigma_{ab}$  ( $a, b \in \{1, 2, ..., 5\}$ ), and  $r_{34}$  constitute a symmetry group of the Garnier system in two variables. This group includes the affine Weyl group of type  $B_5^{(1)}$  [34]. We note that the full symmetry group of the Garnier system has not yet been completely identified.

Based on  $r_{34}$ , for later convenience, we introduce the transformations  $r_{ab} = r_{ba}$  for other pairs of distinct subscripts by

$$r_{\sigma(a)\sigma(b)} = \sigma r_{ab} \sigma^{-1} \quad (\sigma \in \mathfrak{S}_5), \tag{2.10}$$

where  $a, b \in \{1, ..., 5\}, a \neq b$ . It is easy to see that we have

$$r_{ab}^{2} = 1, \qquad r_{i}r_{ab} = r_{ab}r_{i} \quad (i \neq a, b).$$
 (2.11)

The action of  $r_{ab}$  on the parameters and the dependent variables is given by

$$r_{ab} : \theta_a \mapsto -\theta_a + 1, \quad \theta_b \mapsto -\theta_b + 1, \quad \theta_c \mapsto -\theta_c \ (c \neq a, b),$$
$$q_i \mapsto Q_i^{(ab)}, \quad p_i \mapsto P_i^{(ab)}, \tag{2.12}$$

where  $(\theta_3, \theta_4, \theta_5) = (\kappa_0, \kappa_1, \kappa_\infty)$ , and  $Q_i^{(ab)}, P_i^{(ab)}$  (i = 1, 2) are certain rational functions in the canonical variables  $q_i, p_i$  (i = 1, 2).

We also define the Schlesinger transformations

$$T_{ab} = r_a r_b r_c r_d r_e r_{ab}, \quad \mathcal{T}_{ab} = r_a r_c r_d r_e r_{ab} r_b, \tag{2.13}$$

where  $a, b, c, d, e \in \{1, ..., 5\}$  are mutually distinct. These transformations commute with each other, and act on the parameters by

$$\begin{array}{ll} T_{ab} & : & \theta_a \mapsto \theta_a + 1, & \theta_b \mapsto \theta_b + 1, & \theta_c \mapsto \theta_c & (c \neq a, b), \\ T_{ab} & : & \theta_a \mapsto \theta_a + 1, & \theta_b \mapsto \theta_b - 1, & \theta_c \mapsto \theta_c & (c \neq a, b). \end{array}$$

$$(2.14)$$

2.2. **Transformations of the Hamiltonians.** One can introduce the  $\tau$ -variables through the Hamiltonians [20] in a similar manner to the Painlevé equations (see, for example, [28, 29]). For this purpose, we here describe the action of  $r_i$  (i = 1, ..., 5) and  $r_{34}$  on the Hamiltonians.

Following [34], we first introduce the modified Hamiltonians  $h_i^{(3)}$  (i = 1, 2) by adding correction terms, which do not depend on  $p_i$ ,  $q_i$ , to the Hamiltonian  $H_1$  (2.2) as

$$\begin{split} h_{1}^{(3)} &= H_{1} - \frac{\theta_{1}\theta_{2}}{2} \left( \frac{1}{t_{1} - 1} - \frac{1}{t_{1} - t_{2}} \right) + \frac{\theta_{1}(\kappa_{0} - 1)}{2} \left( \frac{1}{t_{1}} - \frac{1}{t_{1} - 1} \right) - \frac{\theta_{1}\kappa_{1}}{2} \frac{1}{t_{1} - 1} \\ &+ \frac{\theta_{1}^{2}}{8} \left( \frac{1}{t_{1}} - \frac{3}{t_{1} - 1} + \frac{1}{t_{1} - t_{2}} \right) - \frac{\theta_{2}^{2}}{24} \left( \frac{1}{t_{1}} + \frac{1}{t_{1} - 1} - \frac{3}{t_{1} - t_{2}} \right) \\ &+ \frac{(\kappa_{0} - 1)^{2}}{24} \left( \frac{3}{t_{1}} - \frac{1}{t_{1} - 1} - \frac{1}{t_{1} - t_{2}} \right) - \frac{\kappa_{1}^{2}}{24} \left( \frac{1}{t_{1}} + \frac{1}{t_{1} - 1} + \frac{1}{t_{1} - t_{2}} \right) \\ &- \frac{\kappa_{\infty}^{2}}{24} \left( \frac{1}{t_{1}} - \frac{3}{t_{1} - 1} + \frac{1}{t_{1} - t_{2}} \right), \end{split}$$

$$\end{split}$$

$$(2.15)$$

and  $h_2^{(3)} = \sigma_{12}(h_1^{(3)})$ . Then we have

$$r_i(h_j^{(3)}) = h_j^{(3)} \quad (i \neq 3), \qquad r_3(h_j^{(3)}) = h_j^{(3)} - \frac{\kappa_0 q_j}{t_j^2 (g_t - 1)} + \frac{\kappa_0}{6} \left(\frac{3}{t_1} - \frac{1}{t_1 - 1} - \frac{1}{t_1 - t_2}\right). \tag{2.16}$$

We see that the 1-form

$$\omega_3 = \sum_{j=1}^2 h_j^{(3)} dt_j, \qquad (2.17)$$

is invariant with respect to the action of  $\sigma_{ab}$  ( $a, b \neq 3$ ), namely

$$\sigma_{ab}(\omega_3) = \omega_3 \quad (a, b \neq 3). \tag{2.18}$$

Let us introduce the 1-form  $\omega_k$  (k = 1, ..., 5) and the Hamiltonians  $h_j^{(k)}$  (j = 1, 2; k = 1, ..., 5) by

$$\sigma(\omega_k) = \omega_{\sigma(k)} \quad (\sigma \in \mathfrak{S}_5), \tag{2.19}$$

and

$$\omega_k = \sum_{j=1}^2 h_j^{(k)} dt_j, \qquad (2.20)$$

respectively. We also define an auxiliary 1-form  $\omega$  by

$$\omega = \omega_5 + r_5(\omega_5). \tag{2.21}$$

Then it follows that

$$r_i(\omega_k) = \omega_k, \quad (i \neq k),$$
  

$$r_{ab}(\omega_a) = \omega_b, \quad r_{ab}(\omega_b) = \omega_a.$$
(2.22)

It is possible to verify by direct computation the following formulas among the above Hamiltonians

$$r_{1}(h_{1}^{(1)}) + h_{1}^{(1)} - r_{5}(h_{1}^{(5)}) - h_{1}^{(5)} - \frac{\partial}{\partial t_{1}} \log q_{1} = \frac{1}{3} \left( \frac{1}{t_{1} - t_{2}} - \frac{2}{t_{1}} \right),$$

$$r_{2}(h_{2}^{(1)}) + h_{2}^{(1)} - r_{5}(h_{2}^{(5)}) - h_{2}^{(5)} - \frac{\partial}{\partial t_{2}} \log q_{1} = -\frac{1}{3} \left( \frac{1}{t_{1} - t_{2}} + \frac{1}{t_{2} - 1} \right),$$

$$r_{3}(h_{j}^{(3)}) + h_{j}^{(3)} - r_{5}(h_{j}^{(5)}) - h_{j}^{(5)} - \frac{\partial}{\partial t_{j}} \log \left( \frac{q_{1}}{t_{1}} + \frac{q_{2}}{t_{2}} - 1 \right) = -\frac{1}{3} \left( \frac{1}{t_{j} - 1} - \frac{1}{t_{j}} \right),$$

$$r_{4}(h_{j}^{(4)}) + h_{j}^{(4)} - r_{5}(h_{j}^{(5)}) - h_{j}^{(5)} - \frac{\partial}{\partial t_{j}} \log(q_{1} + q_{2} - 1) = -\frac{1}{3} \frac{1}{t_{j} - 1},$$

$$(2.23)$$

$$\begin{aligned} h_1^{(1)} + r_{34}(h_1^{(1)}) - h_1^{(4)} - h_1^{(3)} - \frac{\partial}{\partial t_1} \log(q_1 p_1 - \theta_1) \\ &= -\frac{1}{4} \left( \frac{1}{t_1} - \frac{1}{t_1 - 1} - \frac{1}{t_1 - t_2} \right), \\ h_2^{(1)} + r_{34}(h_2^{(1)}) - h_2^{(4)} - h_2^{(3)} - \frac{\partial}{\partial t_2} \log(q_1 p_1 - \theta_1) \\ &= -\frac{1}{12} \left( \frac{1}{t_2} + \frac{1}{t_2 - 1} + \frac{3}{t_1 - t_2} \right), \end{aligned}$$
(2.24)

and

$$r_{5}(h_{j}^{(5)}) + r_{34}r_{5}(h_{j}^{(5)}) - h_{j}^{(4)} - h_{j}^{(3)} - \frac{\partial}{\partial t_{j}}\log(q_{1}p_{1} + q_{2}p_{2} + \alpha)$$

$$= -\frac{1}{12}\left(\frac{1}{t_{j}} - \frac{3}{t_{j} - 1} + \frac{1}{t_{j} - t_{k}}\right),$$

$$h_{j}^{(5)} + r_{34}(h_{j}^{(5)}) - h_{j}^{(4)} - h_{j}^{(3)} - \frac{\partial}{\partial t_{j}}\log(q_{1}p_{1} + q_{2}p_{2} + \alpha + \kappa_{\infty})$$

$$= -\frac{1}{12}\left(\frac{1}{t_{j}} - \frac{3}{t_{j} - 1} + \frac{1}{t_{j} - t_{k}}\right).$$
(2.25)

From Equations (2.23), (2.24) and (2.25), we can deduce the following relations among the 1-forms  $\omega_k$  (k = 1, ..., 5)

$$d \log q_{1} = r_{1}(\omega_{1}) + \omega_{1} - r_{5}(\omega_{5}) - \omega_{5} + \frac{2}{3}\frac{dt_{1}}{t_{1}} + \frac{1}{3}\frac{dt_{2}}{t_{2}-1} - \frac{1}{3}\frac{dt_{1}-dt_{2}}{t_{1}-t_{2}},$$

$$d \log q_{2} = r_{2}(\omega_{2}) + \omega_{2} - r_{5}(\omega_{5}) - \omega_{5} + \frac{1}{3}\frac{dt_{1}}{t_{1}-1} + \frac{2}{3}\frac{dt_{2}}{t_{2}} - \frac{1}{3}\frac{dt_{1}-dt_{2}}{t_{1}-t_{2}},$$

$$d \log \left(\frac{q_{1}}{t_{1}} + \frac{q_{2}}{t_{2}} - 1\right) = r_{3}(\omega_{1}) + \omega_{3} - r_{5}(\omega_{5}) - \omega_{5} + \sum_{j=1}^{2}\frac{1}{3}\left(\frac{1}{t_{j}-1} - \frac{1}{t_{j}}\right)dt_{j},$$

$$d \log(q_{1}+q_{2}-1) = r_{4}(\omega_{1}) + \omega_{4} - r_{5}(\omega_{5}) - \omega_{5} + \sum_{j=1}^{2}\frac{1}{3}\frac{dt_{j}}{t_{j}-1},$$

$$d \log(q_{1}p_{1}-\theta_{1}) = \omega_{1} + r_{34}(\omega_{1}) - \omega_{4} - \omega_{3} + \frac{1}{4}\left(\frac{1}{t_{1}} - \frac{1}{t_{1}-1}\right)dt_{1} + \frac{1}{12}\left(\frac{1}{t_{2}} + \frac{1}{t_{2}-1}\right)dt_{2} - \frac{1}{4}\frac{dt_{1}-dt_{2}}{t_{1}-t_{2}},$$

 $d \log(q_2 p_2 - \theta_2) = \omega_2 + r_{34}(\omega_2) - \omega_4 - \omega_3$ 

$$+\frac{1}{12}\left(\frac{1}{t_1}+\frac{1}{t_1-1}\right)dt_1+\frac{1}{4}\left(\frac{1}{t_2}-\frac{1}{t_2-1}\right)dt_2-\frac{1}{4}\frac{dt_1-dt_2}{t_1-t_2}$$

and

 $d\log(q_1p_1 + q_2p_2 + \alpha) = \omega_5 + r_{34}r_5(\omega_5) - \omega_4 - \omega_3$ 

$$+\sum_{j=1}^{2}\frac{1}{12}\left(\frac{1}{t_{j}}-\frac{3}{t_{j}-1}\right)+\frac{1}{12}\frac{dt_{1}-dt_{2}}{t_{1}-t_{2}},$$

(2.27)

(2.28)

 $d\log(q_1p_1+q_2p_2+\alpha+\kappa_\infty)=\omega_5+r_{34}(\omega_5)-\omega_4-\omega_3$ 

$$+\sum_{j=1}^{2}\frac{1}{12}\left(\frac{1}{t_{j}}-\frac{3}{t_{j}-1}\right)+\frac{1}{12}\frac{dt_{1}-dt_{2}}{t_{1}-t_{2}},$$

respectively.

2.3.  $\tau$ -variables and bilinear relations. Here we introduce the  $\tau$ -variables for the Garnier system in two variables and formulate the action of the following subgroup

$$G = \langle r_i, r_{ab} \rangle, \qquad i = 1, \dots, 5, \quad a, b \in \{1, \dots, 5\},$$
(2.29)

on the  $\tau$ -variables; see also [34]. Note that the independent variables  $t_i$  (i = 1, 2) are invariant under the action of this subgroup.

It is known that the 1-forms  $\omega_k$  (k = 1, ..., 5) given in Equations (2.20) and (2.21) are closed [20], so that one can define the  $\tau$ -variables  $\tau_k$  (k = 1, ..., 5) and  $\tau$  by

$$\omega_k = d\log \tau_k \quad (k = 1, \dots, 5), \quad \omega = d\log \tau, \tag{2.30}$$

up to constant multiples.

From Equations (2.22), (2.26) and (2.30), one can define the action of  $r_i$  (i = 1, ..., 5) on the variables  $\tau_k$  and the auxiliary variable  $\tau$  by

$$r_{i}(\tau_{k}) = \tau_{k} (i \neq k), \quad r_{i}(\tau) = \tau (i = 1, ..., 5),$$

$$r_{1}(\tau_{1}) = t_{1}^{-2/3}(t_{2} - 1)^{-1/3}(t_{1} - t_{2})^{1/3}q_{1} \frac{\tau}{\tau_{1}},$$

$$r_{2}(\tau_{2}) = t_{2}^{-2/3}(t_{1} - 1)^{-1/3}(t_{1} - t_{2})^{1/3}q_{2} \frac{\tau}{\tau_{2}},$$

$$r_{3}(\tau_{3}) = \prod_{j=1}^{2} t_{j}^{1/3}(t_{j} - 1)^{-1/3} \left(\frac{q_{1}}{t_{1}} + \frac{q_{2}}{t_{2}} - 1\right) \frac{\tau}{\tau_{3}},$$

$$r_{4}(\tau_{4}) = \prod_{j=1}^{2} (t_{j} - 1)^{-1/3} (q_{1} + q_{2} - 1) \frac{\tau}{\tau_{4}}, \quad r_{5}(\tau_{5}) = \frac{\tau}{\tau_{5}}.$$
(2.31)

The actions of  $r_{ab}$  ( $a, b \in \{1, ..., 5\}$ ) on the  $\tau$ -variables can be determined from Equations (2.27) and (2.28). To this end, we define some preliminary notations. For mutually distinct indices  $k, a, b \in \{1, ..., 5\}$ , we define  $\varphi_{ab,k} = \varphi_{ba,k}$  by

$$\varphi_{34,k} = q_k p_k - \theta_k \quad (k = 1, 2), \qquad \varphi_{34,5} = -(q_1 p_1 + q_2 p_2 + \alpha + \kappa_{\infty}), \tag{2.32}$$

and

$$\sigma(\varphi_{ab,k}) = \varphi_{\sigma(a)\sigma(b),\sigma(k)} \quad (\sigma \in \mathfrak{S}_5).$$
(2.33)

Further we introduce  $\varphi_{a5}$  (a = 1, 2, 3, 4) by

$$\varphi_{15} = \frac{\sigma_{13}\sigma_{45}r_{34}(g_1 - 1)}{g_1 - 1}\varphi_{15,4}r_4(\varphi_{15,4}), \quad \varphi_{25} = \frac{\sigma_{23}\sigma_{45}r_{34}(g_1 - 1)}{g_1 - 1}\varphi_{25,4}r_4(\varphi_{25,4}),$$

$$\varphi_{35} = \frac{\sigma_{45}r_{34}(g_1 - 1)}{g_1 - 1}\varphi_{35,4}r_4(\varphi_{35,4}), \qquad \varphi_{45} = \frac{\sigma_{35}r_{34}(g_t - 1)}{g_t - 1}\varphi_{45,3}r_3(\varphi_{45,3}),$$
(2.34)

where  $g_1$  and  $g_t$  are given in Equation (2.4). Note that  $\varphi_{ab,k}$  and  $\varphi_{a5}$  are indeed polynomials of the canonical variables  $q_i$ ,  $p_i$  (i = 1, 2). Then the action of  $r_{ab}$  on the  $\tau$ -variables can be expressed as follows

$$r_{ab} : \tau_{a} \mapsto \tau_{b}, \qquad \tau_{b} \mapsto \tau_{a}, \qquad \tau_{k} \mapsto t_{ab,k}\varphi_{ab,k}\frac{\tau_{a}\tau_{b}}{\tau_{k}} \quad (k \neq a, b),$$
  
$$\tau \mapsto t_{ab,5}{}^{2}\varphi_{ab,5}r_{5}(\varphi_{ab,5})\frac{\tau_{a}{}^{2}\tau_{b}{}^{2}}{\tau} \quad (a, b \neq 5),$$
  
$$\tau \mapsto t_{a5}\varphi_{a5}\frac{\tau_{a}{}^{2}\tau_{5}{}^{2}}{\tau},$$
  
$$(2.35)$$

where  $t_{ab,k} = t_{ba,k}$  and  $t_{ab} = t_{ba}$  are certain power products of  $t_1, t_2, t_1 - 1, t_2 - 1$  and  $t_1 - t_2$ . For readers' convenience we give the explicit form of them. Let introduce the notation

$$\chi(e_1, e_2, e_3, e_4, e_5) = t_1^{e_1} t_2^{e_2} (t_1 - 1)^{e_3} (t_2 - 1)^{e_4} (t_1 - t_2)^{e_5},$$
(2.36)

then  $t_{ab,k}$  and  $t_{ab}$  are given by

$$t_{12,3} = \chi(-\frac{1}{4}, -\frac{1}{4}, -\frac{1}{12}, -\frac{1}{12}, \frac{5}{12}), \quad t_{12,4} = \chi(-\frac{1}{12}, -\frac{1}{12}, -\frac{1}{12}, -\frac{1}{12}, \frac{5}{12}),$$
  

$$t_{12,5} = \chi(-\frac{1}{12}, -\frac{1}{12}, -\frac{1}{4}, -\frac{1}{4}, \frac{5}{12}),$$
(2.37)

$$t_{13,2} = \chi(\frac{5}{12}, -\frac{1}{4}, -\frac{1}{12}, \frac{1}{4}, -\frac{1}{4}), \quad t_{13,4} = \sqrt{-1}\chi(\frac{5}{12}, -\frac{1}{12}, -\frac{1}{12}, -\frac{1}{12}, -\frac{1}{12}),$$

$$t_{13,5} = \chi(\frac{5}{12}, -\frac{1}{12}, -\frac{1}{12}, -\frac{1}{12}, -\frac{1}{12}), \quad (2.38)$$

$$t_{14,2} = -\chi(-\frac{1}{12}, \frac{1}{4}, -\frac{1}{12}, \frac{1}{4}, \frac{1}{4}), \qquad t_{14,3} = -\chi(-\frac{1}{4}, \frac{1}{4}, -\frac{1}{12}, -\frac{1}{12}, -\frac{1}{12}), t_{14,5} = -\chi(-\frac{1}{12}, -\frac{1}{12}, -\frac{1}{4}, \frac{1}{4}, -\frac{1}{12}), \qquad (2.39)$$

$$t_{23,1} = -\chi(-\frac{1}{4}, \frac{5}{12}, \frac{1}{4}, -\frac{1}{12}, -\frac{1}{4}), \qquad t_{23,4} = -\chi(-\frac{1}{12}, \frac{5}{12}, -\frac{1}{12}, -\frac{1}{12}, -\frac{1}{12}), t_{23,5} = \sqrt{-1}\chi(-\frac{1}{12}, \frac{5}{12}, \frac{1}{4}, -\frac{1}{4}, -\frac{1}{12}), \qquad (2.40)$$

$$t_{24,1} = -\sqrt{-1}\chi(\frac{1}{4}, -\frac{1}{12}, \frac{1}{4}, -\frac{1}{12}, -\frac{1}{4}), \quad t_{24,3} = -\chi(\frac{1}{4}, -\frac{1}{4}, -\frac{1}{12}, -\frac{1}{12}, -\frac{1}{12}),$$
  

$$t_{24,5} = -\chi(-\frac{1}{12}, -\frac{1}{12}, \frac{1}{4}, -\frac{1}{4}, -\frac{1}{12}),$$
(2.41)

$$t_{34,1} = \sqrt{-1}\chi(-\frac{1}{12}, -\frac{1}{12}, \frac{1}{4}, -\frac{1}{12}, \frac{1}{4}), \quad t_{34,2} = \sqrt{-1}\chi(-\frac{1}{12}, -\frac{1}{4}, -\frac{1}{12}, \frac{1}{4}, \frac{1}{4}), \quad t_{34,5} = \sqrt{-1}\chi(-\frac{1}{12}, -\frac{1}{12}, \frac{1}{4}, \frac{1}{4}, -\frac{1}{12}), \quad (2.42)$$

$$t_{15,2} = \chi(-\frac{1}{12}, \frac{1}{4}, \frac{5}{12}, -\frac{1}{4}, -\frac{1}{4}), \qquad t_{15,3} = \sqrt{-1}\chi(-\frac{1}{4}, \frac{1}{4}, \frac{5}{12}, -\frac{1}{12}, -\frac{1}{12}), t_{15,4} = \sqrt{-1}\chi(-\frac{1}{12}, -\frac{1}{12}, \frac{5}{12}, -\frac{1}{12}, -\frac{1}{12}), \qquad t_{15} = \chi(-\frac{1}{6}, -\frac{1}{6}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{6}),$$
(2.43)

$$t_{25,1} = -\chi(\frac{1}{4}, -\frac{1}{12}, -\frac{1}{4}, \frac{5}{12}, -\frac{1}{4}), \qquad t_{25,3} = -\chi(\frac{1}{4}, -\frac{1}{4}, -\frac{1}{12}, \frac{5}{12}, -\frac{1}{12}), t_{25,4} = -\chi(-\frac{1}{12}, -\frac{1}{12}, -\frac{1}{12}, \frac{5}{12}, -\frac{1}{12}), \qquad t_{25} = -\chi(-\frac{1}{6}, -\frac{1}{6}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{6}),$$
(2.44)

$$t_{35,1} = \chi(-\frac{1}{4}, -\frac{1}{12}, -\frac{1}{4}, -\frac{1}{12}, \frac{1}{4}), \qquad t_{35,2} = \chi(-\frac{1}{12}, -\frac{1}{4}, -\frac{1}{12}, -\frac{1}{4}, \frac{1}{4}), t_{35,4} = \chi(-\frac{1}{12}, -\frac{1}{12}, -\frac{1}{12}, -\frac{1}{12}, -\frac{1}{12}), \qquad t_{35} = \chi(-\frac{1}{6}, -\frac{1}{6}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{6}),$$
(2.45)

$$t_{45,1} = -\chi(\frac{1}{4}, -\frac{1}{12}, -\frac{1}{4}, -\frac{1}{12}, \frac{1}{4}), \qquad t_{45,2} = -\chi(-\frac{1}{12}, \frac{1}{4}, -\frac{1}{12}, -\frac{1}{4}, \frac{1}{4}), t_{45,3} = -\sqrt{-1}\chi(\frac{1}{4}, \frac{1}{4}, -\frac{1}{12}, -\frac{1}{12}, -\frac{1}{12}), \qquad t_{45} = -\chi(-\frac{1}{6}, -\frac{1}{6}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{6}).$$
(2.46)

On the level of the  $\tau$ -variables, the fundamental relations (2.11) are preserved except for some modifications. First we have

$$r_{ab}^{2}(\tau_{k}) = -\tau_{k} \quad (k \neq a, b).$$
 (2.47)

The commutativity of the Schlesinger transformations  $T_{\mu}, T_{\mu'} \in \{T_{ab}, \mathcal{T}_{ab} \mid a, b \in \{1, ..., 5\}, a \neq b\}$  are modified by

$$T_{\mu}T_{\mu'}(\tau_k) = c_{\mu,\mu'}T_{\mu'}T_{\mu}(\tau_k) \quad (k = 1, \dots, 5), \qquad T_{\mu}T_{\mu'}(\tau) = c_{\mu,\mu'}{}^2T_{\mu'}T_{\mu}(\tau), \tag{2.48}$$

where  $c_{\mu,\mu'}$  are certain constants depending on the pair  $(\mu,\mu')$ .

One can derive bilinear relations for the  $\tau$ -variables from the above formulation. It is easy to verify that we have

$$r_k(\varphi_{ab,k}) - \varphi_{ab,k} = \theta_k. \tag{2.49}$$

Substituting  $t_{ab,k}\varphi_{ab,k} = \frac{\tau_k r_{ab}(\tau_k)}{\tau_a \tau_b}$  into the above formula, we get the bilinear relations

$$\tau_{k,k}r_{ab}(\tau_{k,k}) - \tau_k r_{ab}(\tau_k) = \theta_k t_{ab,k}\tau_a \tau_b, \qquad (2.50)$$

for mutually distinct indices  $a, b, k \in \{1, ..., 5\}$ , where  $\tau_{k,k} = r_k(\tau_k)$ . From the formulas (2.31), we get the bilinear relation

$$\prod_{j=1}^{2} t_{j}^{-1/3} (t_{j} - 1)^{1/3} \tau_{3} \tau_{3,3} = t_{1}^{-1/3} (t_{2} - 1)^{1/3} (t_{1} - t_{2})^{-1/3} \tau_{1} \tau_{1,1} + t_{2}^{-1/3} (t_{1} - 1)^{1/3} (t_{1} - t_{2})^{-1/3} \tau_{2} \tau_{2,2} - \tau_{5} \tau_{5,5}.$$
(2.51)

8

#### 3. DERIVATION OF DIFFERENCE EQUATIONS AND CONSTRAINTS

In this section, we derive a system of partial difference equations with some constraints from the bilinear relations for the  $\tau$ -variables of the Garnier system in two variables. The dependent variable of the system we derive can be regarded as defined on  $\mathbb{Z}^4$ . We obtain the difference equations (1.1) and (1.2) by restricting the domain of the system to  $\mathbb{Z}^3$ .

3.1. *f*-variable and nonlinear relations. Suggested by our preceding works [7, 19], we introduce the *f*-variable in terms of the  $\tau$ -variables by

$$f = \frac{\tau_{5,5}}{\tau_5} = \frac{r_5(\tau_5)}{\tau_5}.$$
(3.1)

We then derive nonlinear relations that yield equations (1.1) and (1.2) in terms of the *f*-variable. Note that the action of the Schlesinger transformations  $T_{ab}$ ,  $\mathcal{T}_{ab}$  ( $a, b \in \{1, ..., 5\}$ ) on *f* commutes with each other, since the *f*-variable is defined by a ratio of  $\tau$ -variables.

**Proposition 3.1.** The *f*-variable satisfies the following relations

$$\frac{f - T_{14}^{-1}(f)}{T_{14}^{-1}(f) - T_{12}^{-1}(f)} \frac{T_{12}^{-1}(f) - T_{24}^{-1}(f)}{T_{24}^{-1}(f) - f} = \frac{1 - t_2}{1 - t_1},$$

$$\frac{f - T_{24}^{-1}(f)}{T_{24}^{-1}(f) - T_{23}^{-1}(f)} \frac{T_{23}^{-1}(f) - T_{34}^{-1}(f)}{T_{34}^{-1}(f) - f} = \frac{1}{1 - t_2},$$

$$\frac{f - T_{34}^{-1}(f)}{T_{34}^{-1}(f) - T_{13}^{-1}(f)} \frac{T_{13}^{-1}(f) - T_{14}^{-1}(f)}{T_{14}^{-1}(f) - f} = 1 - t_1.$$
(3.2)

We remark that each equation in (3.2) corresponds to the cross-ratio equation or discrete Schwarzian KdV equation [25, 30] which is also known as  $Q1_{\delta=0}$  [1, 2, 14–16].

*Proof.* The bilinear relations (2.50) with (a, b, k) = (1, 4, 5) and (2, 4, 5) yield

$$f - T_{14}^{-1}(f) = -\kappa_{\infty} \chi(-\frac{1}{12}, -\frac{1}{12}, -\frac{1}{4}, \frac{1}{4}, -\frac{1}{12}) \frac{\tau_1 \tau_4}{\tau_5 r_{14}(\tau_{5,5})},$$
(3.3)

$$f - T_{24}^{-1}(f) = -\kappa_{\infty} \chi(-\frac{1}{12}, -\frac{1}{12}, \frac{1}{4}, -\frac{1}{4}, -\frac{1}{12}) \frac{\tau_2 \tau_4}{\tau_5 r_{24}(\tau_{5,5})},$$
(3.4)

from which we have

$$\frac{f - T_{14}^{-1}(f)}{f - T_{24}^{-1}(f)} = (t_1 - 1)^{-1/2} (t_2 - 1)^{1/2} \frac{\tau_1 r_{24}(\tau_{5,5})}{\tau_2 r_{14}(\tau_{5,5})}.$$
(3.5)

Applying the transformation  $T_{24}^{-1}r_4$  to (3.3), we get

$$T_{24}^{-1}(f) - T_{12}^{-1}(f) = -\kappa_{\infty} \chi(-\frac{1}{12}, -\frac{1}{12}, -\frac{1}{4}, \frac{1}{4}, -\frac{1}{12}) \frac{r_{24}(\tau_{1,1})\tau_2}{r_{24}(\tau_{5,5})r_{24}r_1r_{14}(\tau_5)}.$$
 (3.6)

Similarly, we obtain

$$T_{14}^{-1}(f) - T_{12}^{-1}(f) = -\kappa_{\infty} \chi(-\frac{1}{12}, -\frac{1}{12}, \frac{1}{4}, -\frac{1}{4}, -\frac{1}{12}) \frac{r_{14}(\tau_{2,2})\tau_1}{r_{14}(\tau_{5,5})r_{14}r_2r_{24}(\tau_5)},$$
(3.7)

by applying  $T_{14}^{-1}r_4$  to (3.4). Due to the relation

$$\frac{r_{24}(\tau_{1,1})}{r_{24}r_{1}r_{14}(\tau_{5})} = \frac{r_{14}(\tau_{2,2})}{r_{14}r_{2}r_{24}(\tau_{5})},$$
(3.8)

we have

$$\frac{T_{24}^{-1}(f) - T_{12}^{-1}(f)}{T_{14}^{-1}(f) - T_{12}^{-1}(f)} = (t_1 - 1)^{-1/2}(t_2 - 1)^{1/2}\frac{\tau_2 r_{14}(\tau_{5,5})}{\tau_1 r_{24}(\tau_{5,5})}.$$
(3.9)

The formulas (3.5) and (3.9) yield the first relation of (3.2).

Similar computations give the other relations of Proposition 3.1. Although they are obtained by repetition of the same process, we exhibit the derivations in detail for readers'

convenience. First, we show the third relation of (3.2). The bilinear relation (2.50) with (a, b, k) = (3, 4, 5) gives

$$f - T_{34}^{-1}(f) = \sqrt{-1} \kappa_{\infty} \chi(-\frac{1}{12}, -\frac{1}{12}, \frac{1}{4}, \frac{1}{4}, -\frac{1}{12}) \frac{\tau_3 \tau_4}{\tau_5 r_{34}(\tau_{5,5})}.$$
(3.10)

Then we have

$$\frac{f - T_{14}^{-1}(f)}{f - T_{34}^{-1}(f)} = \sqrt{-1} \left( t_1 - 1 \right)^{-1/2} \frac{\tau_1 r_{34}(\tau_{5,5})}{\tau_3 r_{14}(\tau_{5,5})},$$
(3.11)

from (3.3) and (3.10). Applying the transformation  $T_{34}^{-1}r_4$  to (3.3), we get

$$T_{34}^{-1}(f) - T_{13}^{-1}(f) = -\kappa_{\infty} \chi(-\frac{1}{12}, -\frac{1}{12}, -\frac{1}{4}, \frac{1}{4}, -\frac{1}{12}) \frac{r_{34}(\tau_{1,1})\tau_3}{r_{34}(\tau_{5,5})r_{34}r_1r_{14}(\tau_5)}.$$
 (3.12)

Similarly we obtain

$$T_{14}^{-1}(f) - T_{13}^{-1}(f) = \sqrt{-1} \kappa_{\infty} \chi(-\frac{1}{12}, -\frac{1}{12}, \frac{1}{4}, \frac{1}{4}, -\frac{1}{12}) \frac{r_{14}(\tau_{3,3})\tau_1}{r_{14}(\tau_{5,5})r_{14}r_3r_{34}(\tau_5)},$$
(3.13)

by applying  $T_{14}^{-1}r_4$  to (3.10). Noticing the relation

$$\frac{r_{34}(\tau_{1,1})}{r_{34}r_{1}r_{14}(\tau_{5})} = \frac{r_{14}(\tau_{3,3})}{r_{14}r_{3}r_{34}(\tau_{5})},$$
(3.14)

we have

$$\frac{T_{34}^{-1}(f) - T_{13}^{-1}(f)}{T_{14}^{-1}(f) - T_{13}^{-1}(f)} = \sqrt{-1} (t_1 - 1)^{-1/2} \frac{\tau_3 r_{14}(\tau_{5,5})}{\tau_1 r_{34}(\tau_{5,5})}.$$
(3.15)

The formulas (3.11) and (3.15) yield the third relation of (3.2).

Finally, we deduce the second relation of (3.2). We have from (3.4) and (3.10)

$$\frac{f - T_{24}^{-1}(f)}{f - T_{34}^{-1}(f)} = \sqrt{-1} \left(t_2 - 1\right)^{-1/2} \frac{\tau_2 r_{34}(\tau_{5,5})}{\tau_3 r_{24}(\tau_{5,5})}.$$
(3.16)

Applying the transformation  $T_{34}^{-1}r_4$  to (3.4), we get

$$T_{34}^{-1}(f) - T_{23}^{-1}(f) = -\kappa_{\infty} \chi(-\frac{1}{12}, -\frac{1}{12}, \frac{1}{4}, -\frac{1}{12}) \frac{r_{34}(\tau_{2,2})\tau_3}{r_{34}(\tau_{5,5})r_{34}r_2r_{24}(\tau_5)}.$$
 (3.17)

Similarly we obtain

$$T_{24}^{-1}(f) - T_{23}^{-1}(f) = \sqrt{-1} \kappa_{\infty} \chi(-\frac{1}{12}, -\frac{1}{12}, \frac{1}{4}, \frac{1}{4}, -\frac{1}{12}) \frac{r_{24}(\tau_{3,3})\tau_2}{r_{24}(\tau_{5,5})r_{24}r_3r_{34}(\tau_5)}$$
(3.18)

by applying  $T_{24}^{-1}r_4$  to (3.10). Then the relation

$$\frac{r_{34}(\tau_{2,2})}{r_{34}r_2r_{24}(\tau_5)} = \frac{r_{24}(\tau_{3,3})}{r_{24}r_3r_{34}(\tau_5)},$$
(3.19)

gives

$$\frac{T_{34}^{-1}(f) - T_{23}^{-1}(f)}{T_{24}^{-1}(f) - T_{23}^{-1}(f)} = \sqrt{-1} (t_2 - 1)^{-1/2} \frac{\tau_3 r_{24}(\tau_{5,5})}{\tau_2 r_{34}(\tau_{5,5})}.$$
(3.20)

The formulas (3.16) and (3.20) yield the second relation of (3.2), which completes the proof of Proposition 3.1.  $\hfill \Box$ 

Proposition 3.2. The *f*-variable satisfies the following relations

$$\left(\frac{1}{f} - \frac{1}{T_{14}^{-1}(f)}\right) (\mathcal{T}_{54}(f) - \mathcal{T}_{51}(f)) = \kappa_{\infty}(\kappa_{\infty} + 1)\chi(-\frac{1}{6}, -\frac{1}{6}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{6}),$$

$$\left(\frac{1}{f} - \frac{1}{T_{24}^{-1}(f)}\right) (\mathcal{T}_{54}(f) - \mathcal{T}_{52}(f)) = \kappa_{\infty}(\kappa_{\infty} + 1)\chi(-\frac{1}{6}, -\frac{1}{6}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{6}),$$

$$\left(\frac{1}{f} - \frac{1}{T_{34}^{-1}(f)}\right) (\mathcal{T}_{54}(f) - \mathcal{T}_{53}(f)) = -\kappa_{\infty}(\kappa_{\infty} + 1)\chi(-\frac{1}{6}, -\frac{1}{6}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{6}),$$

$$(3.21)$$

where  $\chi$  is given by (2.36).

We note that each of (3.21) corresponds  $H1_{\epsilon=0}$  in [1, 2, 14–16].

Proof. We start with the relations

$$f - T_{14}^{-1}(f) = -\kappa_{\infty} \chi(-\frac{1}{12}, -\frac{1}{12}, -\frac{1}{4}, \frac{1}{4}, -\frac{1}{12}) \frac{\tau_1 \tau_4}{\tau_5 r_{14}(\tau_{5,5})},$$

$$\mathcal{T}_{54}(f) - \mathcal{T}_{51}(f) = -(\kappa_{\infty} + 1) \chi(-\frac{1}{12}, -\frac{1}{12}, -\frac{1}{4}, \frac{1}{4}, -\frac{1}{12}) \frac{r_5 r_{45}(\tau_{1,1}) \tau_{5,5}}{\tau_4 r_1 r_5 r_{45} r_{14}(\tau_{5,5})},$$
(3.22)

the first of which is (3.3) and the second can be obtained by applying  $r_4T_{45}$  to the first. Due to the relation

$$\frac{r_5 r_{45}(\tau_{1,1})}{r_1 r_5 r_{45} r_{14}(\tau_{5,5})} = -\frac{r_{14}(\tau_5)}{\tau_1},$$
(3.23)

we have

$$\begin{pmatrix} f - T_{14}^{-1}(f) \end{pmatrix} (\mathcal{T}_{54}(f) - \mathcal{T}_{51}(f)) \\ = -\kappa_{\infty}(\kappa_{\infty} + 1)\chi(-\frac{1}{6}, -\frac{1}{6}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{6}) f T_{14}^{-1}(f),$$

$$(3.24)$$

which is equivalent to the first relation of (3.21).

The second and third relations can be also obtained in a similar way. Let us start with the relations

$$f - T_{24}^{-1}(f) = -\kappa_{\infty} \chi(-\frac{1}{12}, -\frac{1}{12}, \frac{1}{4}, -\frac{1}{4}, -\frac{1}{12}) \frac{\tau_2 \tau_4}{\tau_5 r_{24}(\tau_{5,5})},$$

$$\mathcal{T}_{54}(f) - \mathcal{T}_{52}(f) = -(\kappa_{\infty} + 1) \chi(-\frac{1}{12}, -\frac{1}{12}, \frac{1}{4}, -\frac{1}{4}, -\frac{1}{12}) \frac{r_5 r_{45}(\tau_{2,2}) \tau_{5,5}}{\tau_4 r_2 r_5 r_{45} r_{24}(\tau_{5,5})},$$
(3.25)

the first of which is (3.4) and the second can be obtained by applying  $r_4T_{45}$  to the first. Due to the relation

$$\frac{r_5 r_{45}(\tau_{2,2})}{r_2 r_5 r_{45} r_{24}(\tau_{5,5})} = -\frac{r_{24}(\tau_5)}{\tau_2},$$
(3.26)

we have

$$\begin{pmatrix} f - T_{24}^{-1}(f) \end{pmatrix} (\mathcal{T}_{54}(f) - \mathcal{T}_{52}(f)) \\ = -\kappa_{\infty}(\kappa_{\infty} + 1)\chi(-\frac{1}{6}, -\frac{1}{6}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{6}) f T_{24}^{-1}(f),$$

$$(3.27)$$

which is equivalent to the second relation of (3.21).

Similarly we start with the relations

$$f - T_{34}^{-1}(f) = \sqrt{-1} \kappa_{\infty} \chi(-\frac{1}{12}, -\frac{1}{12}, \frac{1}{4}, \frac{1}{4}, -\frac{1}{12}) \frac{\tau_3 \tau_4}{\tau_5 r_{34}(\tau_{5,5})},$$

$$\mathcal{T}_{54}(f) - \mathcal{T}_{53}(f) = \sqrt{-1} (\kappa_{\infty} + 1) \chi(-\frac{1}{12}, -\frac{1}{12}, \frac{1}{4}, \frac{1}{4}, -\frac{1}{12}) \frac{r_5 r_{45}(\tau_{3,3}) \tau_{5,5}}{\tau_4 r_3 r_5 r_{45} r_{34}(\tau_{5,5})},$$
(3.28)

the first of which is (3.10) and the second can be obtained by applying  $r_4T_{45}$  to to the first. Due to the relation

$$\frac{r_5 r_{45}(\tau_{3,3})}{r_3 r_5 r_{45} r_{34}(\tau_{5,5})} = -\frac{r_{34}(\tau_5)}{\tau_3},$$
(3.29)

we have

$$\begin{pmatrix} (f - T_{34}^{-1}(f)) (\mathcal{T}_{54}(f) - \mathcal{T}_{53}(f)) \\ = \kappa_{\infty}(\kappa_{\infty} + 1) \chi(-\frac{1}{6}, -\frac{1}{6}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{6}) f T_{34}^{-1}(f),$$

$$(3.30)$$

which is equivalent to the third relation of (3.21). This completes the proof of Proposition 3.2.

We next derive some relations that yield the similarity constraint (1.2).

Lemma 3.3. The *f*-variable satisfies the following relations

$$\begin{aligned} \theta_{1} \frac{f - \mathcal{T}_{14}(f)}{T_{14}^{-1}(f) - \mathcal{T}_{14}(f)} + \theta_{2} \frac{f - \mathcal{T}_{24}(f)}{T_{24}^{-1}(f) - \mathcal{T}_{24}(f)} + \kappa_{0} \frac{f - \mathcal{T}_{34}(f)}{T_{34}^{-1}(f) - \mathcal{T}_{34}(f)} \\ &= \frac{\theta_{1} + \theta_{2} + \kappa_{0} + \kappa_{1} + \kappa_{\infty} - 1}{2}, \\ \theta_{1} \frac{f - \mathcal{T}_{14}(f)}{T_{14}^{-1}(f) - \mathcal{T}_{14}(f)} T_{14}^{-1}(f) + \theta_{2} \frac{f - \mathcal{T}_{24}(f)}{T_{24}^{-1}(f) - \mathcal{T}_{24}(f)} T_{24}^{-1}(f) \\ &+ \kappa_{0} \frac{f - \mathcal{T}_{34}(f)}{T_{34}^{-1}(f) - \mathcal{T}_{34}(f)} T_{34}^{-1}(f) = \frac{\theta_{1} + \theta_{2} + \kappa_{0} + \kappa_{1} - \kappa_{\infty} - 1}{2} f. \end{aligned}$$
(3.31)

*Proof.* The second relation can be obtained by applying the transformation  $r_5$  to the first one. Then we demonstrate the proof of the first relation in detail. We start with the relations

$$f - T_{14}^{-1}(f) = -\kappa_{\infty} \chi(-\frac{1}{12}, -\frac{1}{12}, -\frac{1}{4}, \frac{1}{4}, -\frac{1}{12}) \frac{\tau_1 \tau_4}{\tau_5 r_{14}(\tau_{5,5})},$$
  

$$f - \mathcal{T}_{14}(f) = -\kappa_{\infty} \chi(-\frac{1}{12}, -\frac{1}{12}, -\frac{1}{4}, \frac{1}{4}, -\frac{1}{12}) \frac{\tau_{1,1} \tau_4}{\tau_5 r_1 r_{14}(\tau_{5,5})},$$
(3.32)

the first of which is (3.3) and the second can be obtained by applying  $r_1$  to the first. Then we get

$$T_{14}^{-1}(f) - \mathcal{T}_{14}(f) = \theta_1 \kappa_{\infty} \chi(\frac{1}{6}, -\frac{1}{6}, -\frac{1}{2}, \frac{1}{6}, \frac{1}{6}) \frac{\tau_4^2}{r_{14}(\tau_{5,5})r_1 r_{14}(\tau_{5,5})},$$
(3.33)

by using the bilinear relation (2.50) with (a, b, k) = (4, 5, 1). Then we obtain

$$\theta_1 \frac{f - \mathcal{T}_{14}(f)}{T_{14}^{-1}(f) - \mathcal{T}_{14}(f)} = -\chi(-\frac{1}{4}, \frac{1}{12}, \frac{1}{4}, \frac{1}{12}, -\frac{1}{4})\frac{\tau_{1,1}r_{14}(\tau_{5,5})}{\tau_4\tau_5}.$$
(3.34)

In a similar way one can express the other terms of the left-hand side of the first relation in Lemma 3.3 by the  $\tau$ -variables. For the second term we start with the relations

$$f - T_{24}^{-1}(f) = -\kappa_{\infty} \chi(-\frac{1}{12}, -\frac{1}{12}, \frac{1}{4}, -\frac{1}{4}, -\frac{1}{12}) \frac{\tau_2 \tau_4}{\tau_5 r_{24}(\tau_{5,5})},$$
  

$$f - \mathcal{T}_{24}(f) = -\kappa_{\infty} \chi(-\frac{1}{12}, -\frac{1}{12}, \frac{1}{4}, -\frac{1}{4}, -\frac{1}{12}) \frac{\tau_{2,2} \tau_4}{\tau_5 r_2 r_{24}(\tau_{5,5})},$$
(3.35)

the first of which is (3.4) and the second can be obtained by applying  $r_2$  to the first. We have

$$T_{24}^{-1}(f) - \mathcal{T}_{24}(f) = \theta_2 \kappa_{\infty} \chi(-\frac{1}{6}, \frac{1}{6}, \frac{1}{6}, -\frac{1}{2}, \frac{1}{6}) \frac{\tau_4^2}{r_{24}(\tau_{5,5})r_2r_{24}(\tau_{5,5})}$$
(3.36)

by using the bilinear relation (2.50) with (a, b, k) = (4, 5, 2). Then we obtain

$$\theta_2 \frac{f - \mathcal{T}_{24}(f)}{T_{24}^{-1}(f) - \mathcal{T}_{24}(f)} = -\chi(\frac{1}{12}, -\frac{1}{4}, \frac{1}{12}, \frac{1}{4}, -\frac{1}{4})\frac{\tau_{2,2}r_{24}(\tau_{5,5})}{\tau_4\tau_5}.$$
(3.37)

Similarly, starting with the relations

$$f - T_{34}^{-1}(f) = \sqrt{-1} \kappa_{\infty} \chi(-\frac{1}{12}, -\frac{1}{12}, \frac{1}{4}, \frac{1}{4}, -\frac{1}{12}) \frac{\tau_3 \tau_4}{\tau_5 r_{34}(\tau_{5,5})},$$
  

$$f - \mathcal{T}_{34}(f) = \sqrt{-1} \kappa_{\infty} \chi(-\frac{1}{12}, -\frac{1}{12}, \frac{1}{4}, \frac{1}{4}, -\frac{1}{12}) \frac{\tau_{3,3} \tau_4}{\tau_5 r_3 r_{34}(\tau_{5,5})},$$
(3.38)

the first of which is (3.10) and the second can be obtained by applying  $r_3$  to the first, we get

$$T_{34}^{-1}(f) - \mathcal{T}_{34}(f) = \kappa_0 \kappa_\infty \chi(\frac{1}{6}, \frac{1}{6}, \frac{1}{6}, \frac{1}{6}, -\frac{1}{6}) \frac{\tau_4^2}{r_{34}(\tau_{5,5})r_3r_{34}(\tau_{5,5})}.$$
 (3.39)

Then the bilinear relation (2.50) with (a, b, k) = (4, 5, 3) gives

$$\kappa_0 \frac{f - \mathcal{T}_{34}(f)}{T_{34}^{-1}(f) - \mathcal{T}_{34}(f)} = \sqrt{-1}\chi(-\frac{1}{4}, -\frac{1}{4}, \frac{1}{12}, \frac{1}{12}, \frac{1}{12})\frac{\tau_{3,3}r_{34}(\tau_{5,5})}{\tau_4\tau_5}.$$
(3.40)

Noticing that we have

$$r_{14}(\tau_{5,5}) = r_{45}(\tau_{1,1}), \quad r_{24}(\tau_{5,5}) = r_{45}(\tau_{2,2}), \quad r_{34}(\tau_{5,5}) = r_{45}(\tau_{3,3}),$$
 (3.41)

we find that the left-hand side of the first relation in Lemma 3.3 can be expressed by

$$\frac{1}{t_{45,1}} \frac{\tau_{1,1} r_{45}(\tau_{1,1})}{\tau_4 \tau_5} + \frac{1}{t_{45,2}} \frac{\tau_{2,2} r_{45}(\tau_{2,2})}{\tau_4 \tau_5} + \frac{1}{t_{45,3}} \frac{\tau_{3,3} r_{45}(\tau_{3,3})}{\tau_4 \tau_5}$$

$$= r_1(\varphi_{45,1}) + r_2(\varphi_{45,2}) + r_3(\varphi_{45,3}).$$
(3.42)

From the definition (2.32) and (2.33) of  $\varphi_{ab,k}$  we have

$$r_1(\varphi_{45,1}) + r_2(\varphi_{45,2}) + r_3(\varphi_{45,3}) = -\alpha, \tag{3.43}$$

where  $\alpha = -\frac{1}{2}(\theta_1 + \theta_2 + \kappa_o + \kappa_1 + \kappa_\infty - 1)$ . Then we obtain the first relation in Lemma 3.3.

We have the following relation from Lemma 3.3, which can be verified by a direct computation.

**Proposition 3.4.** The *f*-variable satisfies the following relation

$$\kappa_{\infty}f = \theta_{1} \frac{(\mathcal{T}_{14}(f) - f)(f - T_{14}^{-1}(f))}{\mathcal{T}_{14}(f) - T_{14}^{-1}(f)} + \theta_{2} \frac{(\mathcal{T}_{24}(f) - f)(f - T_{24}^{-1}(f))}{\mathcal{T}_{24}(f) - T_{24}^{-1}(f)} + \kappa_{0} \frac{(\mathcal{T}_{34}(f) - f)(f - T_{34}^{-1}(f))}{\mathcal{T}_{34}(f) - T_{34}^{-1}(f)}.$$
(3.44)

We finally derive additional relations that will be used to derive the similarity constraint (1.2).

Proposition 3.5. The *f*-variable satisfies the following relations

$$\begin{split} \widetilde{\chi} \,\kappa_{\infty} & \left( \frac{\theta_{1}(t_{2}-1)}{(t_{1}-1)(T_{14}^{-1}(f)-\mathcal{T}_{14}(f))} + \frac{\theta_{2}}{T_{24}^{-1}(f)-\mathcal{T}_{24}(f)} \right. \\ & \left. - \frac{\kappa_{0}(t_{2}-1)}{T_{34}^{-1}(f)-\mathcal{T}_{34}(f)} \right) = \frac{1}{T_{54}^{-1}(f)}, \end{split}$$

$$\begin{split} \widetilde{\chi} \,\kappa_{\infty} & \left( \frac{\theta_{1}(t_{2}-1)}{(t_{1}-1)(1/T_{14}^{-1}(f)-1/\mathcal{T}_{14}(f))} \right. \\ & \left. + \frac{\theta_{2}}{1/T_{24}^{-1}(f)-1/\mathcal{T}_{24}(f)} - \frac{\kappa_{0}(t_{2}-1)}{1/T_{34}^{-1}(f)-1/\mathcal{T}_{34}(f)} \right) = \mathcal{T}_{54}(f), \end{split}$$

$$\end{split}$$

$$\end{split}$$

$$\end{split}$$

where

$$\widetilde{\chi} = \chi(-\frac{1}{6}, -\frac{1}{6}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{6}).$$
(3.46)

*Proof.* The second relation can be obtained by applying the transformation  $r_5$  to the first. Then we explain the proof on the first relation in detail. From (3.33) and (3.41) we have

$$\frac{\widetilde{\chi}\kappa_{\infty}\theta_1}{T_{14}^{-1}(f) - \mathcal{T}_{14}(f)} = t_1^{-1/3}(t_1 - 1)(t_2 - 1)^{-2/3}(t_1 - t_2)^{-1/3}r_{45}\left(\frac{\tau_1\tau_{1,1}}{\tau_5^2}\right).$$
 (3.47)

Similarly, we obtain from (3.36), (3.39) and (3.41)

$$\frac{\widetilde{\chi}\kappa_{\infty}\theta_{2}}{T_{24}^{-1}(f) - \mathcal{T}_{24}(f)} = t_{2}^{-1/3}(t_{1} - 1)^{1/3}(t_{1} - t_{2})^{-1/3}r_{45}\left(\frac{\tau_{2}\tau_{2,2}}{\tau_{5}^{2}}\right),$$

$$\frac{\widetilde{\chi}\kappa_{\infty}\kappa_{0}}{T_{34}^{-1}(f) - \mathcal{T}_{34}(f)} = t_{1}^{-1/3}t_{2}^{-1/3}(t_{1} - 1)^{1/3}(t_{2} - 1)^{-2/3}r_{45}\left(\frac{\tau_{3}\tau_{3,3}}{\tau_{5}^{2}}\right).$$
(3.48)

Then the bilinear relation (2.51) gives

$$\begin{aligned} \widetilde{\chi}_{K_{\infty}} \left( \frac{\theta_1(t_2 - 1)}{(T_{14}^{-1}(f) - \mathcal{T}_{14}(f))(t_1 - 1)} + \frac{\theta_2}{T_{24}^{-1}(f) - \mathcal{T}_{24}(f)} - \frac{\kappa_0(t_2 - 1)}{T_{34}^{-1}(f) - \mathcal{T}_{34}(f)} \right) \\ &= r_{45} \left( \frac{\tau_5 \tau_{5,5}}{\tau_5^2} \right) = r_{45}(f) = \frac{1}{T_{54}^{-1}(f)}, \end{aligned}$$
(3.49)

which is the first relation of (3.45).

3.2. **Restriction to a subsystem.** Here we introduce a subgroup of G (defined in (2.29)) and focus on the action of the subgroup. Considering some elements of infinite order of the subgroup we see that the relations for the f-variable obtained in the previous subsection yield a system of partial difference equations together with some constraints.

Let us define the transformations  $s_i^k$  (i = 0, 1; k = 0, 1, 2, 3) and  $\pi_k$  (k = 0, 1, 2, 3) by

$$s_{1}^{k} = r_{k} \quad (k = 1, 2, 3), \qquad s_{0}^{0} = r_{5}, \pi_{1} = r_{2}r_{3}r_{5}r_{14}, \quad \pi_{2} = r_{1}r_{3}r_{5}r_{24}, \quad \pi_{3} = r_{1}r_{2}r_{5}r_{34}, \quad \pi_{0} = r_{1}r_{2}r_{3}r_{54}, s_{0}^{k} = \pi_{k}s_{1}^{k}\pi_{k} \quad (k = 1, 2, 3), \qquad s_{1}^{0} = \pi_{0}s_{0}^{0}\pi_{0}.$$

$$(3.50)$$

Then these elements satisfy the following relations

$$(s_i^k)^2 = 1, \qquad s_i^k s_j^l = s_j^l s_i^k \quad (k \neq l), (\pi_k)^2 = 1, \qquad \pi_k \pi_l = \pi_l \pi_k \quad (k \neq l), \pi_k s_i^l = s_i^l \pi_k \quad (k \neq l).$$
 (3.51)

This means that for each  $k \in \{0, 1, 2, 3\}$  the group generated by the transformations  $s_0^k$  and  $s_1^k$  is isomorphic to the affine Weyl group of type  $A_1^{(1)}$  and  $\pi_k$  plays the role of the automorphism of the Dynkin diagram;

$$\langle s_0^k, s_1^k \rangle \cong W(A_1^{(1)}) \quad (k = 0, 1, 2, 3), \qquad \langle s_0^k, s_1^k, \pi_k \rangle \cong \widetilde{W}(A_1^{(1)}) \quad (k = 0, 1, 2, 3).$$
(3.52)

Also, these four groups  $\langle s_0^k, s_1^k, \pi_k \rangle$  (k = 0, 1, 2, 3) commute with each other. Let us introduce the parameters  $\alpha_i^k$  (i = 0, 1; k = 0, 1, 2, 3, 4) by

$$\alpha_1^1 = \theta_1, \quad \alpha_1^2 = \theta_2, \quad \alpha_1^3 = \kappa_0, \quad \alpha_1^4 = \kappa_1, \quad \alpha_0^0 = \kappa_\infty, \\
\alpha_0^k = 1 - \alpha_1^k \quad (k = 1, 2, 3, 4), \quad \alpha_1^0 = 1 - \alpha_0^0.$$
(3.53)

Then the action of the transformations  $s_i^k$  and  $\pi_k$  on these parameters is given by

$$s_0^k : \alpha_0^k \mapsto -\alpha_0^k, \quad \alpha_1^k \mapsto \alpha_1^k + 2\alpha_0^k, \\ s_1^k : \alpha_1^k \mapsto -\alpha_1^k, \quad \alpha_0^k \mapsto \alpha_0^k + 2\alpha_1^k$$

$$(3.54)$$

and

$$\pi_k : (\alpha_0^k, \alpha_1^k, \alpha_0^4, \alpha_1^4) \mapsto (\alpha_1^k, \alpha_0^k, \alpha_1^4, \alpha_0^4) \quad (k = 0, 1, 2, 3).$$
(3.55)

Let us consider the elements

$$\rho_k = \pi_k s_1^k \quad (k = 0, 1, 2, 3), \tag{3.56}$$

of the group  $\langle s_0^k, s_1^k, \pi_k \rangle$ . Each of them acts on the parameters  $\alpha_0^k$  and  $\alpha_1^k$  as a parallel translation:

$$\rho_k : (\alpha_0^k, \alpha_1^k) \mapsto (\alpha_0^k + 1, \alpha_1^k - 1).$$
(3.57)

Note that we have

$$\rho_k(f) = T_{k4}^{-1}(f), \quad \rho_k^{-1}(f) = \mathcal{T}_{k4}(f) \qquad (k = 1, 2, 3),$$
  

$$\rho_0(f) = \mathcal{T}_{54}(f), \qquad \rho_0^{-1}(f) = T_{54}^{-1}(f).$$
(3.58)

Then the relations given in Proposition 3.1 can be expressed by

$$\frac{f - \rho_1(f)}{\rho_1(f) - \rho_1\rho_2(f)} \frac{\rho_1\rho_2(f) - \rho_2(f)}{\rho_2(f) - f} = \frac{1 - t_2}{1 - t_1},$$

$$\frac{f - \rho_2(f)}{\rho_2(f) - \rho_2\rho_3(f)} \frac{\rho_2\rho_3(f) - \rho_3(f)}{\rho_3(f) - f} = \frac{1}{1 - t_2},$$

$$\frac{f - \rho_3(f)}{\rho_3(f) - \rho_3\rho_1(f)} \frac{\rho_3\rho_1(f) - \rho_1(f)}{\rho_1(f) - f} = 1 - t_1.$$
(3.59)

Let us introduce an infinite family of variables  $f_{l_1,l_2,l_3,l_0}$   $(l_1,l_2,l_3,l_0 \in \mathbb{Z})$  by

$$f_{l_1,l_2,l_3,l_0} = \rho_1^{l_1} \rho_2^{l_2} \rho_3^{l_3} \rho_0^{l_0}(f) \quad (l_1,l_2,l_3,l_0 \in \mathbb{Z}).$$
(3.60)

By using the results in the previous subsection, we obtain several difference equations for the variables  $f_{l_1,l_2,l_3,l_0}$ . First, we have the following statement from (3.59) and (3.21).

**Theorem 3.6.** The variables  $f_{l_1,l_2,l_3,l_0}$  satisfy the partial difference equations

$$\frac{f_{l_1,l_2,l_3,l_0} - f_{l_1+1,l_2,l_3,l_0}}{f_{l_1+1,l_2,l_3,l_0} - f_{l_1,l_2+1,l_3,l_0} - f_{l_1,l_2+1,l_3,l_0}} = \frac{1 - t_2}{1 - t_1},$$

$$\frac{f_{l_1,l_2,l_3,l_0} - f_{l_1,l_2+1,l_3,l_0}}{f_{l_1,l_2+1,l_3,l_0} - f_{l_1,l_2+1,l_3+1,l_0} - f_{l_1,l_2,l_3,l_0}} = \frac{1}{1 - t_2},$$

$$\frac{f_{l_1,l_2,l_3,l_0} - f_{l_1,l_2+1,l_3,l_0}}{f_{l_1,l_2+1,l_3,l_0} - f_{l_1,l_2,l_3+1,l_0} - f_{l_1,l_2,l_3+1,l_0}} = \frac{1}{1 - t_2},$$

$$\frac{f_{l_1,l_2,l_3,l_0} - f_{l_1,l_2,l_3+1,l_0}}{f_{l_1,l_2,l_3+1,l_0} - f_{l_1,l_2,l_3+1,l_0} - f_{l_1,l_2,l_3,l_0}} = 1 - t_1,$$
(3.61)

and

$$\begin{pmatrix} \frac{1}{f_{l_1,l_2,l_3,l_0}} - \frac{1}{f_{l_1+1,l_2,l_3,l_0}} \end{pmatrix} (f_{l_1,l_2,l_3,l_0+1} - f_{l_1+1,l_2,l_3,l_0+1}) \\ = (\alpha_0^0 + l_0)(\alpha_0^0 + l_0 + 1)\chi(-\frac{1}{6}, -\frac{1}{2}, \frac{1}{2}, -\frac{1}{6}), \\ \begin{pmatrix} \frac{1}{f_{l_1,l_2,l_3,l_0}} - \frac{1}{f_{l_1,l_2+1,l_3,l_0}} \end{pmatrix} (f_{l_1,l_2,l_3,l_0+1} - f_{l_1,l_2+1,l_3,l_0+1}) \\ = (\alpha_0^0 + l_0)(\alpha_0^0 + l_0 + 1)\chi(-\frac{1}{6}, -\frac{1}{6}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{6}), \\ \begin{pmatrix} \frac{1}{f_{l_1,l_2,l_3,l_0}} - \frac{1}{f_{l_1,l_2,l_3+1,l_0}} \end{pmatrix} (f_{l_1,l_2,l_3,l_0+1} - f_{l_1,l_2,l_3+1,l_0+1}) \\ = -(\alpha_0^0 + l_0)(\alpha_0^0 + l_0 + 1)\chi(-\frac{1}{6}, -\frac{1}{6}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{6}). \end{cases}$$
(3.62)

We also get the following from (3.44) and (3.45).

**Theorem 3.7.** The variables  $f_{l_1,l_2,l_3,l_0}$  satisfy the difference equations

$$\begin{aligned} (\alpha_{0}^{0}+l_{0}) f_{l_{1},l_{2},l_{3},l_{0}} &= (\alpha_{1}^{1}-l_{1}) \frac{(f_{l_{1}-l_{2},l_{3},l_{0}}-f_{l_{1},l_{2},l_{3},l_{0}})(f_{l_{1},l_{2},l_{3},l_{0}}-f_{l_{1}+1,l_{2},l_{3},l_{0}})}{f_{l_{1}-l_{1},l_{2},l_{3},l_{0}}-f_{l_{1}+1,l_{2},l_{3},l_{0}}} \\ &+ (\alpha_{1}^{2}-l_{2}) \frac{(f_{l_{1},l_{2}-1,l_{3},l_{0}}-f_{l_{1},l_{2},l_{3},l_{0}})(f_{l_{1},l_{2},l_{3},l_{0}}-f_{l_{1},l_{2}+1,l_{3},l_{0}})}{f_{l_{1},l_{2}-1,l_{3},l_{0}}-f_{l_{1},l_{2}+1,l_{3},l_{0}}} \\ &+ (\alpha_{1}^{3}-l_{3}) \frac{(f_{l_{1},l_{2},l_{3}-1,l_{0}}-f_{l_{1},l_{2},l_{3},l_{0}})(f_{l_{1},l_{2},l_{3}-1,l_{0}}-f_{l_{1},l_{2},l_{3}+1,l_{0}})}{f_{l_{1},l_{2},l_{3}-1,l_{0}}-f_{l_{1},l_{2},l_{3}+1,l_{0}}}, \end{aligned}$$
(3.63)

and

$$\frac{1}{f_{l_1,l_2,l_3,l_0-1}} = \chi \left( \alpha_0^0 + l_0 \right) \left( \frac{(\alpha_1^1 - l_1)(t_2 - 1)}{(t_1 - 1)(f_{l_1+1,l_2,l_3,l_0} - f_{l_1-1,l_2,l_3,l_0})} + \frac{(\alpha_1^2 - l_2)}{f_{l_1,l_2+1,l_3,l_0} - f_{l_1,l_2-1,l_3,l_0}} - \frac{(\alpha_1^3 - l_3)(t_2 - 1)}{f_{l_1,l_2,l_3+1,l_0} - f_{l_1,l_2,l_3-1,l_0}} \right),$$

$$f_{l_1,l_2,l_3,l_0+1} = \chi \left( \alpha_0^0 + l_0 \right) \left( \frac{(\alpha_1^1 - l_1)(t_2 - 1)}{(t_1 - 1)(1/f_{l_1+1,l_2,l_3,l_0} - 1/f_{l_1-1,l_2,l_3,l_0})} + \frac{(\alpha_1^2 - l_2)}{1/f_{l_1,l_2+1,l_3,l_0} - 1/f_{l_1,l_2-1,l_3,l_0}} - \frac{(\alpha_1^3 - l_3)(t_2 - 1)}{1/f_{l_1,l_2-1,l_3,l_0}} - \frac{(\alpha_1^3 - l_3)(t_2 - 1)}{1/f_{l_1,l_2,l_3-1,l_0}} \right).$$
(3.64)

Equations (3.63) and (3.64) can be regarded as some constraints for the system of partial difference equations (3.61) and (3.62). One can determine all the values of  $f_{l_1,l_2,l_3,l_0}$  from a finite number of initial values, as we will show in Appendix A.

To conclude this section, we explicitly relate the above result to the defining equations of the discrete power function on a hexagonal lattice.

**Theorem 3.8.** We introduce  $f_{l_1,l_2,l_3}$  by

$$f_{l_1,l_2,l_3} = f_{l_1,l_2,l_3,0}.$$
(3.65)

Then it satisfies the system of partial difference equations (1.1) with

$$x_1 = \frac{1 - t_1}{1 - t_2}, \quad x_2 = 1 - t_2, \quad x_3 = \frac{1}{1 - t_1}$$
 (3.66)

together with the similarity constraint (1.2).

When we consider the discrete power function on a hexagonal lattice, we have in equation (1.2)  $\alpha_1^1 = \alpha_1^2 = \alpha_1^3 = 0$ . In this case, by putting  $l_1 = l_2 = l_3 = 0$ , we see from (1.2) that  $f_{0,0,0}$  cannot be freely chosen but instead must be 0. Also, by giving the initial values at the six points  $f_{\pm 1,0,0}$ ,  $f_{0,0,\pm 1}$ ,  $f_{0,0,\pm 1}$  one can determine all the values of  $f_{l_1,l_2,l_3}$ . However, the way to determination of values is different from the generic case. If we start with the initial data at the three points  $f_{1,0,0}$ ,  $f_{0,0,-1}$ , as given by (1.3), we can determine the values of  $f_{l_1,l_2,l_3}$  consistently only for  $l_1 \ge 0$ ,  $l_2 \ge 0$ ,  $l_3 \le 0$ . In this sense, the initial value problem for the discrete power function is consistent, but different from the generic case. The specialization  $\alpha_1^1 = \alpha_1^2 = \alpha_1^3 = 0$  for the equation (1.2) suggests that the discrete power function on a hexagonal lattice relates to the special solution of the hypergeometric type to the Garnier system in two variables.

## 4. CONCLUDING REMARKS

In this paper, we constructed a system of partial difference equations together with certain constraints from the discrete symmetry of the Garnier system in two variables. This contains the difference equations (1.1) and (1.2), as a sub-system, which are associated with the discrete power functions on a hexagonal lattice.

It is known that the Garnier system in two variables admits special solutions expressed by Appel's hypergeometric functions. Consequently, we expect to deduce an explicit formula of the discrete power function in terms of the hypergeometric  $\tau$ -functions of the Garnier system in two variables. This will be reported in detail in a separate paper.

**Acknowledgment.** This research was supported by an Australian Research Council grant # DP160101728 and JSPS KAKENHI Grant Numbers JP16H03941, JP19K14559 and JP17J00092.

16

APPENDIX A. A MULTI-DIMENSIONALLY CONSISTENT SYSTEM OF PARTIAL DIFFERENCE EQUATIONS

The system of six multi-affine linear quad-equations is said to have a consistent-arounda-cube (CAC) property [1,27], if they are placed on six faces of a cube in such a way that, given initial values on 4 vertices of the cube, the solutions of the equations are uniquely defined at each remaining vertex. Integrable partial difference equations arise by iterating such consistent quad-equations on neighbouring cubes that tile the lattice  $\mathbb{Z}^3$ .

The concept of the CAC property can be extended to a hypercube. The system of equations on a hypercube is said to be multi-dimensionally consistent, if the equations on every sub-cube of the hypercube have the CAC property. Extension to *n*-dimensional hypercubes that are space-filling polytopes in the lattice  $\mathbb{Z}^n$  leads to a system of integrable partial difference equations in *n*-directions.

In this section, we consider a system of partial difference equations (A.1) and (A.2), which is an extension of the system discussed in [21] to four dimensions. This system is reduced to the system of difference equations (3.61) and (3.62) under some conditions. The symmetry of the equations (A.1) and (A.2) is taken over to the symmetry of the reduced equations. Furthermore we describe a birational action of the four Weyl groups  $\langle s_0^k, s_1^k, \pi_k \rangle$  (k = 0, 1, 2, 3), which recovers the system of difference equations (1.1) and (1.2), through the system of equations (3.61) and (3.62) together with the constraints (3.63) and (3.64).

A.1. A multi-dimensionally consistent system of partial difference equations. Here, we consider a multi-dimensionally consistent system of partial difference equations and its symmetry.

Let  $u : \mathbb{Z}^4 \mapsto \mathbb{C}$  be a function assigning a value to each vertex of the 4-dimensional integer lattice  $\mathbb{Z}^4$ . We denote the value of u at the vertex  $(l_1, l_2, l_3, l_0) \in \mathbb{Z}^4$  by  $u_{l_1, l_2, l_3, l_0}$ . Suppose  $\mu_l^{(i)} \in \mathbb{C}$   $(l \in \mathbb{Z}; i = 0, 1, 2, 3)$  are parameters assigned to each edge of the hypercube.

We assume that the function u satisfies the relations

$$\frac{(u_{l_1,l_2,l_3,l_0} + u_{l_1+1,l_2,l_3,l_0})(u_{l_1,l_2+1,l_3,l_0} + u_{l_1+1,l_2+1,l_3,l_0})}{(u_{l_1,l_2,l_3,l_0} + u_{l_1,l_2+1,l_3,l_0})(u_{l_1+1,l_2+1,l_3,l_0} + u_{l_1+1,l_2,l_3,l_0})}{(u_{l_1,l_2,l_3,l_0} + u_{l_1,l_2+1,l_3,l_0})(u_{l_1,l_2,l_3+1,l_0} + u_{l_1,l_2+1,l_3+1,l_0})}{(u_{l_1,l_2,l_3,l_0} + u_{l_1,l_2,l_3+1,l_0})(u_{l_1,l_2+1,l_3+1,l_0} + u_{l_1,l_2+1,l_3,l_0})} = \frac{\mu_{l_1}^{(2)}}{\mu_{l_3}^{(3)}},$$
(A.1)
$$\frac{(u_{l_1,l_2,l_3,l_0} + u_{l_1,l_2,l_3+1,l_0})(u_{l_1+1,l_2,l_3+1,l_0} + u_{l_1,l_2+1,l_3,l_0})}{(u_{l_1,l_2,l_3,l_0} + u_{l_1+1,l_2,l_3+1,l_0})(u_{l_1+1,l_2,l_3+1,l_0} + u_{l_1,l_2,l_3+1,l_0})} = \frac{\mu_{l_3}^{(3)}}{\mu_{l_3}^{(1)}},$$

 $\alpha$ 

and

$$\begin{pmatrix} \frac{1}{u_{l_1,l_2,l_3,l_0}} + \frac{1}{u_{l_1+1,l_2,l_3,l_0}} \end{pmatrix} (u_{l_1,l_2,l_3,l_0+1} + u_{l_1+1,l_2,l_3,l_0+1}) = -\mu_{l_1}^{(1)} \mu_{l_0}^{(0)},$$

$$\begin{pmatrix} \frac{1}{u_{l_1,l_2,l_3,l_0}} + \frac{1}{u_{l_1,l_2+1,l_3,l_0}} \end{pmatrix} (u_{l_1,l_2,l_3,l_0+1} + u_{l_1,l_2+1,l_3,l_0+1}) = -\mu_{l_2}^{(2)} \mu_{l_0}^{(0)},$$

$$\begin{pmatrix} \frac{1}{u_{l_1,l_2,l_3,l_0}} + \frac{1}{u_{l_1,l_2,l_3+1,l_0}} \end{pmatrix} (u_{l_1,l_2,l_3,l_0+1} + u_{l_1,l_2,l_3+1,l_0+1}) = -\mu_{l_3}^{(3)} \mu_{l_0}^{(0)}.$$

$$(A.2)$$

This list of six equations above corresponds to twenty four 2-dimensional faces of a 4dimensional hypercube which forms a fundamental cell of  $\mathbb{Z}^4$ . See Figure A.1 for the corresponding labelling assigning coordinates to the vertices of a hypercube and parameters to each direction.

Adler et al. [1,2] and Boll [14–16] classified the six face equations of 3D cube by using the CAC property and additional property called tetrahedron property. Every 3D sub-cube for the system of equations (A.1) and (A.2) is fit with one in [14–16]. In these papers, each of equations (A.1) is called  $Q1_{\delta=0}$  while each of equations (A.2) is called  $H1_{\epsilon=0}$ . Note



Figure A.1. A 4-dimensional hypercube. The sixteen variables  $u_{l_1,l_2,l_3,l_0}, \ldots, u_{l_1+1,l_2+1,l_3+1,l_0+1}$  are assigned to the vertices of the hypercube. Note that  $u = u_{l_1,l_2,l_3,l_0}$  and the subscript *i* means +1 shift in the  $l_i$ -direction, e.g.  $u_1 = u_{l_1+1,l_2,l_3,l_0}$  and  $u_{10} = u_{l_1+1,l_2,l_3,l_0+1}$ . Moreover, each of the lattice parameters  $\mu_{l_i}^{(i)}$ , i = 0, 1, 2, 3, is assigned to the edge connecting *u* and  $u_i$  and to its parallel edges.

that each of partial difference equations (A.1) is known as the discrete Schwarzian KdV equation [25, 30].

It is easy to see that the system of equations (A.1) and (A.2) is invariant under the transformations  $w_i^k$  (i = 0, 1; k = 0, 1, 2, 3) and  $\varpi_k$  (k = 0, 1, 2, 3) given by

$$\begin{split} w_{0}^{1} &: u_{l_{1},l_{2},l_{3},l_{0}} \mapsto u_{2-l_{1},l_{2},l_{3},l_{0}}, \qquad \mu_{l}^{(1)} \mapsto \mu_{1-l}^{(1)}, \\ w_{1}^{1} &: u_{l_{1},l_{2},l_{3},l_{0}} \mapsto u_{-l_{1},l_{2},l_{3},l_{0}}, \qquad \mu_{l}^{(1)} \mapsto \mu_{-1-l}^{(1)}, \\ \overline{\varpi}_{1} &: u_{l_{1},l_{2},l_{3},l_{0}} \mapsto -u_{1-l_{1},l_{2},l_{3},l_{0}}, \qquad \mu_{l}^{(1)} \mapsto \mu_{-l}^{(1)}, \\ w_{0}^{2} &: u_{l_{1},l_{2},l_{3},l_{0}} \mapsto u_{l_{1},2-l_{2},l_{3},l_{0}}, \qquad \mu_{l}^{(2)} \mapsto \mu_{1-l}^{(2)}, \\ w_{1}^{2} &: u_{l_{1},l_{2},l_{3},l_{0}} \mapsto u_{l_{1},-l_{2},l_{3},l_{0}}, \qquad \mu_{l}^{(2)} \mapsto \mu_{-l}^{(2)}, \\ \overline{\varpi}_{2} &: u_{l_{1},l_{2},l_{3},l_{0}} \mapsto -u_{l_{1},1-l_{2},l_{3},l_{0}}, \qquad \mu_{l}^{(2)} \mapsto \mu_{-l}^{(2)}, \\ w_{0}^{3} &: u_{l_{1},l_{2},l_{3},l_{0}} \mapsto u_{l_{1},l_{2},2-l_{3},l_{0}}, \qquad \mu_{l}^{(3)} \mapsto \mu_{1-l}^{(3)}, \\ w_{1}^{3} &: u_{l_{1},l_{2},l_{3},l_{0}} \mapsto u_{l_{1},l_{2},-l_{3},l_{0}}, \qquad \mu_{l}^{(3)} \mapsto \mu_{-l}^{(3)}, \\ w_{0}^{3} &: u_{l_{1},l_{2},l_{3},l_{0}} \mapsto -u_{l_{1},l_{2},1-l_{3},l_{0}}, \qquad \mu_{l}^{(0)} \mapsto \mu_{-l}^{(0)}, \\ w_{0}^{0} &: u_{l_{1},l_{2},l_{3},l_{0}} \mapsto \frac{1}{u_{l_{1},l_{2},l_{3},-l_{0}}, \qquad \mu_{l}^{(0)} \mapsto \mu_{-l-l}^{(0)}, \\ w_{1}^{0} &: u_{l_{1},l_{2},l_{3},l_{0}} \mapsto \frac{1}{u_{l_{1},l_{2},l_{3},-l_{0}}, \qquad \mu_{l}^{(0)} \mapsto \mu_{-l-l}^{(0)}, \\ \overline{\varpi}_{0} &: u_{l_{1},l_{2},l_{3},l_{0}} \mapsto -\frac{1}{u_{l_{1},l_{2},l_{3},-l_{0}}, \qquad \mu_{l}^{(0)} \mapsto \mu_{-l-l}^{(0)}, \\ \overline{\varpi}_{0} &: u_{l_{1},l_{2},l_{3},l_{0}} \mapsto -\frac{1}{u_{l_{1},l_{2},l_{3},-l_{0}}, \qquad \mu_{l}^{(0)} \mapsto \mu_{-l-l}^{(0)}, \\ \overline{\varpi}_{0} &: u_{l_{1},l_{2},l_{3},l_{0}} \mapsto -\frac{1}{u_{l_{1},l_{2},l_{3},-l_{0}}, \qquad \mu_{l}^{(0)} \mapsto \mu_{-2-l}^{(0)}. \end{split}$$

These transformations can be read from the setting that the variables  $u_{l_1,l_2,l_3,l_0}$  and the parameters  $\mu_l^{(i)}$  are respectively assigned to the vetices and edges of the lattice. For each of  $k \in \{0, 1, 2, 3\}$  the transformations  $w_0^k, w_1^k$  and  $\varpi_k$  give a realization of the extended affine

Weyl group  $\widetilde{W}(A_1^{(1)})$ , namely they satisfy the fundamental relations

$$(w_0^k)^2 = (w_1^1)^2 = (\overline{\omega}_k)^2 = 1, \quad \overline{\omega}_k w_0^k = w_1^k \overline{\omega}_k.$$
 (A.4)

Furthermore, the four group  $\langle w_0^k, w_1^k, \varpi_k \rangle$ , k = 0, 1, 2, 3, commute with each other.

A.2. Reduction of the system of equations (A.1) and (A.2). Let us consider the reduction of (A.1) and (A.2) to the system derived from the discrete symmetry of the Garnier system in two variables. To this end, we specialize  $\mu_l^{(i)}$  (*i* = 0, 1, 2, 3) as

$$\mu_l^{(1)} = \chi(-\frac{1}{6}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{6}), \quad \mu_l^{(2)} = \chi(-\frac{1}{6}, -\frac{1}{6}, \frac{1}{2}, -\frac{1}{2}, -\frac{1}{6}),$$

$$\mu_l^{(3)} = -\chi(-\frac{1}{6}, -\frac{1}{6}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{6}), \quad \mu_l^{(0)} = (\alpha_0^0 + l) (\alpha_0^0 + l + 1),$$
(A.5)

where  $\chi$  is given by (2.36) and  $\alpha_0^0$  is a complex parameter. Then, the equations (A.1) and (A.2) become the equations (3.61) and (3.62), respectively, by the change of variables

$$f_{l_1,l_2,l_3,l_0} = (-1)^{l_1+l_2+l_3} u_{l_1,l_2,l_3,l_0}.$$
(A.6)

Recall that the difference equations (3.61) and (3.62) are parts of the system derived from the discrete symmetry of the Garnier system in two variables. This system also contains the constraints (3.64) and

$$\frac{\alpha_{1}^{1} + \alpha_{1}^{2} + \alpha_{1}^{3} - \alpha_{0}^{4} + \alpha_{0}^{0} - l_{1} - l_{2} - l_{3} + l_{0}}{2} = (\alpha_{1}^{1} - l_{1}) \frac{f_{l_{1},l_{2},l_{3},l_{0}} - f_{l_{1}-1,l_{2},l_{3},l_{0}}{f_{l_{1}+1,l_{2},l_{3},l_{0}} - f_{l_{1}-1,l_{2},l_{3},l_{0}}} + (\alpha_{1}^{2} - l_{2}) \frac{f_{l_{1},l_{2},l_{3},l_{0}} - f_{l_{1},l_{2}-1,l_{3},l_{0}}{f_{l_{1},l_{2}+1,l_{3},l_{0}} - f_{l_{1},l_{2}-1,l_{3},l_{0}}} + (\alpha_{1}^{3} - l_{3}) \frac{f_{l_{1},l_{2},l_{3},l_{0}} - f_{l_{1},l_{2},l_{3}-1,l_{0}}}{f_{l_{1},l_{2},l_{3}+1,l_{0}} - f_{l_{1},l_{2},l_{3}-1,l_{0}}}, \qquad (A.7)$$

$$\frac{\alpha_{1}^{1} + \alpha_{1}^{2} + \alpha_{1}^{3} - \alpha_{0}^{4} - \alpha_{0}^{0} - l_{1} - l_{2} - l_{3} - l_{0}}{f} f_{l_{1},l_{2},l_{3},l_{0}} + (\alpha_{1}^{2} - l_{2}) \frac{f_{l_{1},l_{2},l_{3},l_{0}} - f_{l_{1}-1,l_{2},l_{3},l_{0}}}{f_{l_{1}+1,l_{2},l_{3},l_{0}} - f_{l_{1}-1,l_{2},l_{3},l_{0}}} f_{l_{1}+1,l_{2},l_{3},l_{0}} + (\alpha_{1}^{2} - l_{2}) \frac{f_{l_{1},l_{2},l_{3},l_{0}} - f_{l_{1}-1,l_{2},l_{3},l_{0}}}{f_{l_{1},l_{2}+1,l_{3},l_{0}} - f_{l_{1},l_{2}-1,l_{3},l_{0}}} f_{l_{1},l_{2}+1,l_{3},l_{0}} + (\alpha_{1}^{3} - l_{3}) \frac{f_{l_{1},l_{2},l_{3},l_{0}} - f_{l_{1},l_{2}-1,l_{3},l_{0}}}{f_{l_{1},l_{2},l_{3}+1,l_{0}} - f_{l_{1},l_{2}-1,l_{3},l_{0}}} f_{l_{1},l_{2}+1,l_{3},l_{0}} + (\alpha_{1}^{3} - l_{3}) \frac{f_{l_{1},l_{2},l_{3},l_{0}} - f_{l_{1},l_{2}-1,l_{3},l_{0}}}{f_{l_{1},l_{2},l_{3}-1,l_{0}}} f_{l_{1},l_{2},l_{3}+1,l_{0}},$$

where  $\alpha_1^i$  (*i* = 1, 2, 3) and  $\alpha_0^j$  (*j* = 0, 4) are also complex parameters. Note that the constraints (A.7) directly come from Lemma 3.3. By using the partial difference equations (A.1) and (A.2) together with the constraints (3.64) and (A.7), one can determine all the values of the variables  $f_{l_1,l_2,l_3,l_0}$  from a finite number of initial values. For instance, one can choose

 $f_0 = f_{0,0,0,0}, \quad f_{\pm 1} = f_{\pm 1,0,0,0}, \quad f_{\pm 2} = f_{0,\pm 1,0,0}, \quad f_{\pm 3} = f_{0,0,\pm 1,0}$ (A.8)

as the initial values, where these seven values satisfy the relations

$$\frac{\alpha_1^1 + \alpha_1^2 + \alpha_1^3 - \alpha_0^4 + \alpha_0^0}{2} = \alpha_1^1 \frac{f_0 - f_{-1}}{f_1 - f_{-1}} + \alpha_1^2 \frac{f_0 - f_{-2}}{f_2 - f_{l-2}} + \alpha_1^3 \frac{f_0 - f_{-3}}{f_3 - f_{-3}},$$

$$\frac{\alpha_1^1 + \alpha_1^2 + \alpha_1^3 - \alpha_0^4 - \alpha_0^0}{2} f_0 = \alpha_1^1 \frac{f_0 - f_{-1}}{f_1 - f_{-1}} f_1 + \alpha_1^2 \frac{f_0 - f_{-2}}{f_2 - f_{-2}} f_2 + \alpha_1^3 \frac{f_0 - f_{-3}}{f_3 - f_{-3}} f_3$$
(A.9)

obtained from the constraints (A.7).

The symmetry of the system of equations (A.1) and (A.2) described by  $\langle w_0^k, w_1^k, \overline{\omega}_k \rangle$ , k = 0, 1, 2, 3, is taken over to that of the reduced system discussed in §3.2 by the following isomorphism:

$$w_1^k \mapsto s_1^k, \quad w_0^k \mapsto s_0^k, \quad \overline{\omega}_k \mapsto \pi_k \qquad (k = 0, 1, 2, 3).$$
 (A.10)

From the actions (A.3) we obtain

$$\begin{aligned}
 s_{0}^{1} &: & f_{l_{1},l_{2},l_{3},l_{0}} \mapsto f_{2-l_{1},l_{2},l_{3},l_{0}}, & s_{1}^{1} &: & f_{l_{1},l_{2},l_{3},l_{0}} \mapsto f_{-l_{1},l_{2},l_{3},l_{0}}, \\
 \pi_{1} &: & f_{l_{1},l_{2},l_{3},l_{0}} \mapsto f_{1-l_{1},l_{2},l_{3},l_{0}}, \\
 s_{0}^{2} &: & f_{l_{1},l_{2},l_{3},l_{0}} \mapsto f_{l_{1},2-l_{2},l_{3},l_{0}}, & s_{1}^{2} &: & f_{l_{1},l_{2},l_{3},l_{0}} \mapsto f_{l_{1},-l_{2},l_{3},l_{0}}, \\
 \pi_{2} &: & f_{l_{1},l_{2},l_{3},l_{0}} \mapsto f_{l_{1},1-l_{2},l_{3},l_{0}}, & s_{1}^{3} &: & f_{l_{1},l_{2},l_{3},l_{0}} \mapsto f_{l_{1},l_{2},-l_{3},l_{0}}, \\
 s_{0}^{3} &: & f_{l_{1},l_{2},l_{3},l_{0}} \mapsto f_{l_{1},l_{2},2-l_{3},l_{0}}, & s_{1}^{3} &: & f_{l_{1},l_{2},l_{3},l_{0}} \mapsto f_{l_{1},l_{2},-l_{3},l_{0}}, \\
 \pi_{3} &: & f_{l_{1},l_{2},l_{3},l_{0}} \mapsto f_{l_{1},l_{2},1-l_{3},l_{0}},
 \end{aligned}$$
(A.11)

and

$$s_{0}^{0} : f_{l_{1},l_{2},l_{3},l_{0}} \mapsto \frac{1}{f_{l_{1},l_{2},l_{3},-l_{0}}}, \quad \alpha_{0}^{0} \mapsto -\alpha_{0}^{0},$$

$$s_{1}^{0} : f_{l_{1},l_{2},l_{3},l_{0}} \mapsto \frac{1}{f_{l_{1},l_{2},l_{3},-2-l_{0}}}, \quad \alpha_{0}^{0} \mapsto 2 - \alpha_{0}^{0},$$

$$\pi_{0} : f_{l_{1},l_{2},l_{3},l_{0}} \mapsto \frac{1}{f_{l_{1},l_{2},l_{3},-1-l_{0}}}, \quad \alpha_{0}^{0} \mapsto 1 - \alpha_{0}^{0},$$
(A.12)

Let us introduce the auxiliary parameters  $\alpha_0^i$  (*i* = 1, 2, 3) and  $\alpha_1^j$  (*j* = 0, 4) by

$$\alpha_0^k = 1 - \alpha_1^k, \quad k = 0, 1, 2, 3, 4.$$
 (A.13)

Since we impose the constraints (3.64) and (A.7), one can formulate the birational action of  $\langle s_0^k, s_1^k, \pi_k \rangle$ , k = 0, 1, 2, 3, on the field  $\mathbb{K}(f_0, f_{\pm 1}, f_{\pm 2}, f_{\pm 3})$  of rational functions with the coefficient field  $\mathbb{K} = \mathbb{C}(\{\alpha_i^j\})$ . Recall that the action on the coefficient field  $\mathbb{K}$  is given by (3.54) and (3.55). It is easy to see that the transformations  $s_1^k$  (k = 1, 2, 3) and  $s_0^0$  are given by

$$s_{1}^{k}(f_{k}) = f_{-k}, \quad s_{1}^{k}(f_{-k}) = f_{k} \quad (k = 1, 2, 3),$$
  

$$s_{0}^{0}(f_{k}) = \frac{1}{f_{k}} \qquad (k = 0, \pm 1, \pm 2, \pm 3).$$
(A.14)

The description of the transformations  $\pi_k$  (k = 1, 2, 3, 0) is a little bit complicated. Regarding the transformation  $\pi_1$ , we have  $\pi_1(f_0) = f_1, \pi_1(f_1) = f_0$  and

$$\frac{f_0 - f_{-3}}{f_{-3} - \pi_1(f_{-3})} \frac{\pi_1(f_{-3}) - f_1}{f_1 - f_0} = 1 - t_1, \quad \frac{f_0 - f_3}{f_3 - \pi_1(f_3)} \frac{\pi_1(f_3) - f_1}{f_1 - f_0} = 1 - t_1,$$

$$\frac{f_0 - f_{-1}}{f_{-1} - \pi_1(f_{-2})} \frac{\pi_1(f_{-2}) - f_{-2}}{f_{-2} - f_0} = \frac{1 - t_2}{1 - t_1}, \quad \frac{f_0 - f_1}{f_1 - \pi_1(f_2)} \frac{\pi_1(f_2) - f_2}{f_2 - f_0} = \frac{1 - t_2}{1 - t_1},$$
(A.15)

which come from the equations (A.1). By solving each equations of (A.15) for  $\pi_1(f_{-3})$ ,  $\pi_1(f_3)$ ,  $\pi_1(f_{-2})$  and  $\pi_1(f_2)$ , we see that these are expressed by the rational functions of  $f_0, f_{\pm 1}, f_{\pm 2}, f_{\pm 3}$ . From the first relation of (A.9) we also get

$$\frac{\alpha_0^1 + \alpha_1^2 + \alpha_1^3 - \alpha_0^4 + \alpha_0^0}{2} = \alpha_0^1 \frac{f_1 - \pi_1(f_{-1})}{f_0 - \pi_1(f_{-1})} + \alpha_1^2 \frac{f_1 - \pi_1(f_{-2})}{f_2 - \pi_1(f_{-2})} + \alpha_1^3 \frac{f_1 - \pi_1(f_{-3})}{f_3 - \pi_1(f_{-3})}, \quad (A.16)$$

which gives the expression of  $\pi_1(f_{-1})$  by the rational functions of  $f_0, f_{\pm 1}, f_{\pm 2}, f_{\pm 3}$ . In a similar way,  $\pi_k(f_i)$  (k = 2, 3;  $i \in \{0, \pm 1, \pm 2, \pm 3\}$ ) can be expressed in terms of the rational functions of  $f_0, f_{\pm 1}, f_{\pm 2}, f_{\pm 3}$ . Let us consider the transformation  $\pi_0$ . From the first relation of (3.64), we have

$$\pi_0(f_0) = \widetilde{\chi} \, \alpha_0^0 \left( \frac{\alpha_1^1(t_2 - 1)}{(t_1 - 1)(f_1 - f_{-1})} + \frac{\alpha_1^2}{f_2 - f_{-2}} - \frac{\alpha_1^3(t_2 - 1)}{f_3 - f_{-3}} \right), \tag{A.17}$$

where  $\tilde{\chi}$  is given by (3.46). One can compute  $\pi_0(f_k)$  and  $\pi_0(f_{-k})$  for k = 1, 2, 3 by  $\pi_0(f_k) = \pi_0\pi_k(f_0) = \pi_k\pi_0(f_0)$  and  $\pi_0(f_{-k}) = \pi_0s_1^k(f_k) = s_1^k\pi_0(f_k)$ , respectively. Then, we obtain the expression of  $\pi_0(f_i)$  ( $i \in \{0, \pm 1, \pm 2, \pm 3\}$ ) by the rational functions of  $f_0, f_{\pm 1}, f_{\pm 2}, f_{\pm 3}$ . The transformations  $s_0^k$  (k = 1, 2, 3) and  $s_1^0$  are compositions of the above transformations, namely we have

$$s_0^k = \pi_k s_1^k \pi_k \quad (k = 1, 2, 3), \qquad s_1^0 = \pi_0 s_0^0 \pi_0.$$
 (A.18)

We can verify that the action of  $\langle s_0^k, s_1^k, \pi_k \rangle$ , k = 0, 1, 2, 3, constructed here is equivalent to that in Section 3.2.

#### References

- V. E. Adler, A. I. Bobenko, and Y. B. Suris. Classification of integrable equations on quad-graphs. The consistency approach. *Comm. Math. Phys.*, 233(3):513–543, 2003.
- [2] V. E. Adler, A. I. Bobenko, and Y. B. Suris. Discrete nonlinear hyperbolic equations: classification of integrable cases. *Funktsional. Anal. i Prilozhen.*, 43(1):3–21, 2009.
- [3] S. I. Agafonov. Imbedded circle patterns with the combinatorics of the square grid and discrete Painlevé equations, *Discrete Comput. Geom.*, 29 (2003), 305–319.
- [4] S. I. Agafonov. Discrete Riccati equation, hypergeometric functions and circle patterns of Schramm type, Glasg. Math. J., 47 (2005), 1–16.
- [5] S. I. Agafonov and A. I. Bobenko. Discrete Z<sup>y</sup> and Painlevé equations. *Internat. Math. Res. Notices*, no. 4:165–193, 2000.
- [6] S. I. Agafonov and A. I. Bobenko. Hexagonal circle patterns with constant intersection angles and discrete Painlevé and Riccati equations. J. Math. Phys., 44(8):3455–3469, 2003.
- [7] H. Ando, M. Hay, K. Kajiwara, and T. Masuda. An explicit formula for the discrete power function associated with circle patterns of Schramm type. *Funkcial. Ekvac.* 57(1):1–41, 2014.
- [8] A. I. Bobenko. Discrete conformal maps and surfaces. Symmetries and integrability of difference equations (Canterbury, 1996), 97–108, London Math. Soc. Lecture Note Ser. 255 (Cambridge Univ. Press, Cambridge, 1999).
- [9] A. I.Bobenko and T. Hoffmann. Hexagonal circle patterns and integrable systems: patterns with constant angles. *Duke Math. Journal* 116:525–566, 2003.
- [10] A. I. Bobenko and T. Hoffmann. Conformally symmetric circle packings. A generalization of Doyle spirals. *Exp. Math.* 10:141–150, 2001.
- [11] A. I. Bobenko and A. Its, The asymptotic behavior of the discrete holomorphic map  $Z^a$  via the Riemann-Hilbert method, *Duke Math. J.* 165(14): 2607-2682, 2016.
- [12] A. I. Bobenko and U. Pinkall. Discretization of surfaces and integrable systems. In *Discrete integrable geometry and physics (Vienna, 1996)*, volume 16 of *Oxford Lecture Ser. Math. Appl.*, pages 3–58. Oxford Univ. Press, New York, 1999.
- [13] A. I. Bobenko and Y. B. Suris. Discrete differential geometry, volume 98 of Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, 2008. Integrable structure.
- [14] R. Boll. Classification of 3D consistent quad-equations. J. Nonlinear Math. Phys., 18(3):337-365, 2011.
- [15] R. Boll. Corrigendum: Classification of 3D consistent quad-equations. J. Nonlinear Math. Phys., 19(4):1292001, 3, 2012.
- [16] R. Boll. Classification and Lagrangian Structure of 3D Consistent Quad-Equations. Doctoral Thesis, Technische Universität Berlin, 2012.
- [17] A. Doliwa, M. Nieszporski, and P. M. Santini. Integrable lattices and their sublattices. II. From the Bquadrilateral lattice to the self-adjoint schemes on the triangular and the honeycomb lattices. J. of Math. Phys., 48(11):113506, 2007.
- [18] R. Garnier. Sur des équations différentielles du troisième ordre dont líntégrale générale est uniforme et sur un classe déquations nouvelles dórder supérieur dont líntégrale générale a ses points critiques fixes, Ann. Sci. École Norm. Sup. (3) 29, 1-126, 1912 (French).
- [19] M. Hay, K. Kajiwara, and T. Masuda. Bilinearization and special solutions to the discrete Schwarzian KdV equation. J. Math-for-Ind. 3A:53–62, 2011.
- [20] K. Iwasaki, H. Kimura, S. Shimomura, and M. Yoshida. From Gauss to Painlevé: A modern theory of special functions, Vieweg, Braunschweig, 1991.
- [21] N. Joshi, K. Kajiwara, T. Masuda, N. Nakazono, and Y. Shi. Geometric description of a discrete power function associated with the sixth Painlevé equation. Proc. R. Soc. A., 473(2207):20170312, 19 pages, 2017.
- [22] H. Kimura. Symmetries of the Garnier system and of the associated polynomial Hamiltonian system. Proc. Japan Acad. Ser. A Math. Sci., 66(7):176–178, 1990.
- [23] H. Kimura and K. Okamoto. On the polynomial Hamiltonian structure of the Garnier systems. J. Math. Pures Appl. (9), 63(1):129–146, 1984.
- [24] M. Lis. Circle Patterns and Critical Ising Models. Commun. Math. Phys., 370(2): 507-530, 2019.

- [25] F. W. Nijhoff, H. W. Capel, G. L. Wiersma, and G. R. W. Quispel. Bäcklund transformations and threedimensional lattice equations. *Phys. Lett. A*, 105(6):267–272, 1984.
- [26] F. W. Nijhoff, A. Ramani, B. Grammaticos, and Y. Ohta. On discrete Painlevé equations associated with the lattice KdV systems and the Painlevé VI equation. *Stud. Appl. Math.*, 106(3):261–314, 2001.
- [27] F. W. Nijhoff and A. J. Walker. The discrete and continuous Painlevé VI hierarchy and the Garnier systems. *Glasg. Math. J.*, 43A:109–123, 2001. Integrable systems: linear and nonlinear dynamics (Islay, 1999).
- [28] M. Noumi. Painlevé equations through symmetry, volume 223 of Translations of Mathematical Monographs. American Mathematical Society, Providence, RI, 2004. Translated from the 2000 Japanese original by the author.
- [29] K. Okamoto. Studies on the Painlevé equations. I. Sixth Painlevé equation P<sub>VI</sub>. Ann. Mat. Pura Appl. Serie Quarta, 146(1):337–381, 1987.
- [30] G. R. W. Quispel, F. W. Nijhoff, H. W. Capel, and J. van der Linden. Linear integral equations and nonlinear difference-difference equations. *Phys. A*, 125(2-3):344–380, 1984.
- [31] S. Smirnov, Discrete Complex Analysis and Probability. *Proceedings of the International Congress of Mathematicians*, Hyderabad, India, 2010.
- [32] K. Stephenson. Circle packing: a mathematical tale. Notices of The AMS, 50(11): 1376–1388, 2003.
- [33] K. Stephenson. Introduction to circle packing. Cambridge University Press, Cambridge, 2005.
- [34] T. Suzuki. Affine Weyl group symmetry of the Garnier system. Funkcial. Ekvac., 48(2):203–230, 2005.
- [35] W. P. Thurston. The finite Riemann mapping theorem. Invited Address. International Symposium in Celebration of the Proof of the Bieberbach Conjecture. Purdue University (1985).
- [36] T. Tsuda. Rational solutions of the Garnier system in terms of Schur polynomials. Int. Math. Res. Not. IMRN, (43):2341–2358, 2003.

School of Mathematics and Statistics F07, The University of Sydney, NSW 2006, Australia. *Email address*: nalini.joshi@sydney.edu.au

Institute of Mathematics for Industry, Kyushu University, 744 Motooka, Fukuoka 819-0395, Japan. *Email address*: kaji@imi.kyushu-u.ac.jp

Department of Mathematical Sciences, Aoyama Gakuin University, Sagamihara, Kanagawa 252-5258, Japan. *Email address*: masuda@gem.aoyama.ac.jp

Institute of Engineering, Tokyo University of Agriculture and Technology, 2-24-16 Nakacho Koganei, Tokyo 184-8588, Japan.

Email address: nakazono@go.tuat.ac.jp

22