

# REMARKS ON TORIC MANIFOLDS WHOSE CHERN CHARACTERS ARE POSITIVE

HIROSHI SATO AND YUSUKE SUYAMA

ABSTRACT. We show that the second Chern character of any projective toric manifold of Picard number three is not positive. In connection with this result, we give various examples of the positivity of higher Chern characters of projective toric manifolds.

## CONTENTS

1. Introduction	1
2. Preliminaries	2
3. Toric manifolds of Picard number three	4
4. Positivities of higher Chern characters	7
References	10

## 1. INTRODUCTION

For a smooth projective variety  $X$  over an algebraically closed field and a positive integer  $k$ , the  $k$ -th Chern character  $\mathrm{ch}_k(X)$  is said to be *positive* (resp. *nef*) if the intersection number  $(\mathrm{ch}_k(X) \cdot z)$  is positive (resp. non-negative) for any  $z \in \overline{\mathrm{NE}}_k(X) \setminus \{0\}$ , where  $\mathrm{NE}_k(X)$  is the cone of numerical effective  $k$ -cycles on  $X$ . A variety with  $\mathrm{ch}_1(X)$  positive is nothing but a Fano variety. Fano varieties with  $\mathrm{ch}_2(X)$  nef are called *2-Fano manifolds* and they are first studied by de Jong–Starr [JS] in order to investigate the existence of rational surfaces on a Fano variety. Fano  $d$ -folds with  $\mathrm{ch}_2(X)$  positive and index  $\geq d - 2$  are classified in [AC].

In this paper, we focus on the toric case. A *toric manifold* is a smooth complete toric variety. For  $d \geq 2$ , the  $d$ -dimensional projective space  $\mathbb{P}^d$  is a toric manifold and one can easily see that the second Chern character  $\mathrm{ch}_2(\mathbb{P}^d)$  is positive. On the other hand, we do not know toric manifolds whose second Chern character is positive other than projective spaces at the present moment. Therefore, we study projective toric manifolds of Picard number three in order to find a new example of such toric manifolds. However, on the contrary, we show the following:

**Theorem 1.1** (Theorem 3.6). *The second Chern character of any projective toric manifold of Picard number three which does not have a Fano contraction is not nef.*

Moreover, by combining Theorem 1.1 with the known result in [S2] (see Theorem 2.6), we obtain the following:

**Corollary 1.2** (Corollary 3.7). *Let  $X$  be a smooth projective toric  $d$ -fold of Picard number at most three. If the second Chern character  $\mathrm{ch}_2(X)$  of  $X$  is positive, then  $X$  is isomorphic to the  $d$ -dimensional projective space  $\mathbb{P}^d$ .*

---

Date: 2019/12/6, version 0.12.

2010 *Mathematics Subject Classification*. Primary 14M25; Secondary 14C17, 14J45.

*Key words and phrases*. toric manifolds, Chern character, 2-Fano manifolds.

Also, we deal with the positivity of higher Chern characters of projective toric manifolds. In contrast to the second Chern character, we show that there exists a projective toric manifold with  $\text{ch}_k(X)$  positive other than the projective space for each  $k \geq 3$ . We also give some phenomena which do not appear in the case of the positivity of the second Chern character.

This paper is organized as follows: In Section 2, we collect basic results in the toric geometry. We define the positivity of Chern characters for a projective toric manifold. Section 3 is devoted to the calculation of intersection numbers on projective toric manifolds of Picard number three which admit no Fano contraction. As a result, we can show that the second Chern characters of such manifolds are not nef. In Section 4, we introduce various phenomena about the positivity of higher Chern characters.

**Acknowledgments.** The authors would like to thank Professor Osamu Fujino for his supports. The first author was partially supported by JSPS KAKENHI Grant Number JP18K03262. The second author was partially supported by JSPS KAKENHI Grant Number JP18J00022.

## 2. PRELIMINARIES

In this section, we introduce the fundamental notation and concepts in the toric geometry. For details, please see [CLS], [F] and [O] for the toric geometry. Also see [FS], [M] and [R] for the toric Mori theory. We will work over an algebraically closed field  $K = \overline{K}$ .

Let  $N := \mathbb{Z}^d$ ,  $M := \text{Hom}_{\mathbb{Z}}(N, \mathbb{Z})$ ,  $N_{\mathbb{R}} := N \otimes_{\mathbb{Z}} \mathbb{R}$  and  $M_{\mathbb{R}} := M \otimes_{\mathbb{Z}} \mathbb{R}$ .  $\text{Cone}(S)$  stands for the cone generated by  $S \subset N_{\mathbb{R}}$ . For a fan  $\Sigma$  in  $N$ , we denote by  $X = X_{\Sigma}$  the toric  $d$ -fold associated to  $\Sigma$ . We denote by  $T = T_N$  the algebraic torus for  $N$ . For a smooth complete fan  $\Sigma$ , put

$$G(\Sigma) := \{\text{the primitive generators of 1-dimensional cones in } \Sigma\} \subset N.$$

For a cone  $\sigma \in \Sigma$ , we put  $G(\sigma) := \sigma \cap G(\Sigma)$ , that is, the set of primitive generators of  $\sigma$ . For any  $x \in G(\Sigma)$ , there is the corresponding torus invariant divisor  $D_x$  on  $X$ . It is well-known that there exists the following isomorphism of abelian groups:

$$A_1(X) \cong \left\{ (a_x)_{x \in G(\Sigma)} \in \mathbb{Z}^{G(\Sigma)} \left| \sum_{x \in G(\Sigma)} a_x x = 0 \right. \right\},$$

where  $A_1(X)$  is the group of numerical 1-cycles on  $X$ . Thus, we can regard a linear relation among elements in  $G(\Sigma)$  as a numerical 1-cycle on  $X$ . For a  $(d-1)$ -dimensional cone  $\tau \in \Sigma$ , there exists a linear relation

$$y_1 + y_2 + a_1 x_1 + \cdots + a_{d-1} x_{d-1} = 0,$$

where  $G(\tau) = \{x_1, \dots, x_{d-1}\}$ ,  $a_1, \dots, a_{d-1} \in \mathbb{Z}$  and for  $\{y_1, y_2\} \subset G(\Sigma)$ ,  $G(\tau) \cup \{y_1\}$  and  $G(\tau) \cup \{y_2\}$  generate maximal cones in  $\Sigma$ . This linear relation corresponds to the torus invariant curve  $C_{\tau}$  associated to  $\tau$ . We remark that

$$2 + a_1 + \cdots + a_{d-1} = (-K_X \cdot C_{\tau}).$$

The following relation is convenient to describe a complete smooth fan:

**Definition 2.1** ([B1, Definitions 2.6, 2.7 and 2.8]). Let  $X = X_{\Sigma}$  be a toric manifold. We call a nonempty subset  $P \subset G(\Sigma)$  a *primitive collection* in  $\Sigma$  if  $P$  does not generate a cone in  $\Sigma$ , while any proper subset of  $P$  generates a cone in  $\Sigma$ .

For a primitive collection  $P = \{x_1, \dots, x_r\}$ , there exists the unique cone  $\sigma(P) \in \Sigma$  such that  $x_1 + \dots + x_r$  is contained in the relative interior of  $\sigma(P)$ . Put  $G(\sigma) = \{y_1, \dots, y_s\}$ . Then, we have a linear relation

$$x_1 + \dots + x_r = a_1 y_1 + \dots + a_s y_s \quad (a_1, \dots, a_s \in \mathbb{Z}_{>0}).$$

We call this relation the *primitive relation* of  $P$ . Thus, by the above argument, we obtain the numerical 1-cycle  $r(P) \in A_1(X)$  for any primitive collection  $P \subset G(\Sigma)$ . We remark that

$$r - (a_1 + \dots + a_s) = (-K_X \cdot r(P)).$$

We can characterize toric Fano manifolds using the notion of primitive relations as follows:

**Theorem 2.2** ([B2, Proposition 2.3.6]). *Let  $X = X_\Sigma$  be a toric manifold. Then  $X$  is Fano if and only if  $(-K_X \cdot r(P)) > 0$  for any primitive collection  $P$  in  $\Sigma$ .*

In this paper, we study the positivity of the  $k$ -th Chern character  $\text{ch}_k(X)$  for a smooth projective toric  $d$ -fold  $X = X_\Sigma$ . Let  $D_1, \dots, D_n$  be the torus invariant prime divisors on  $X$ . Then,  $\text{ch}_k(X)$  is described as

$$\text{ch}_k(X) = \frac{1}{k!} (D_1^k + \dots + D_n^k).$$

The following proposition is fundamental for our theory. The proof is completely similar to the one for [O, Proposition 2.26] in the Japanese version. We describe a sketch of the proof for the reader's convenience.

**Proposition 2.3.** *Let  $X = X_\Sigma$  be a toric manifold of  $\dim X = d$  and  $1 \leq k \leq d$ . For any  $k$ -dimensional irreducible closed subvariety  $Y \subset X$ , there exist torus invariant  $k$ -dimensional irreducible closed subvarieties  $Y_1, \dots, Y_l$  such that  $Y$  is numerically equivalent to*

$$a_1 Y_1 + \dots + a_l Y_l$$

for positive integers  $a_1, \dots, a_l$ .

*Sketch of the proof.* Let  $V(\sigma)$  be the smallest torus invariant irreducible closed subvariety associated to a cone  $\sigma \in \Sigma$  which contains  $Y$ . Obviously,  $\dim \sigma \leq d - k$ . So, there exists a  $(d - k)$ -dimensional cone  $\tau \in \Sigma$  such that  $\sigma$  is a face of  $\tau$ . For  $n \in (\text{Relint}(\tau)) \cap N$ , where  $\text{Relint}(\tau)$  stands for the relative interior of  $\tau$ , let  $\gamma_n : K^\times \rightarrow T$  be the corresponding one-parameter subgroup. For a morphism

$$\begin{array}{ccc} \Phi : & V(\sigma) \times K^\times & \rightarrow & V(\sigma) \times K^\times, \\ & \downarrow & & \downarrow \\ & (u, \lambda) & \mapsto & (\gamma_n(\lambda)u, \lambda) \end{array}$$

let  $Z := \overline{\Phi(Y \times K^\times)}$  be the closure of  $\Phi(Y \times K^\times)$  in  $V(\sigma) \times K$  and  $\text{pr}_2 : Z \rightarrow K$  the second projection. Then, there exist  $(d - k)$ -dimensional cones  $\tau_1, \dots, \tau_l$  which contain  $\sigma$  as a face such that

$$\text{pr}_2^{-1}(1) = Y, \text{ while } \text{pr}_2^{-1}(0) = V(\tau_1) \cup \dots \cup V(\tau_l)$$

as sets, where  $V(\tau_1), \dots, V(\tau_l)$  are the torus invariant irreducible closed subvarieties associated to  $\tau_1, \dots, \tau_l$ , respectively (see [CLS, Proposition 3.2.2] and [O, Proposition 1.6 (v)]). Therefore,  $Y$  is rationally equivalent to

$$a_1 V(\tau_1) + \dots + a_l V(\tau_l)$$

for positive integers  $a_1, \dots, a_l$ . In particular, they are numerically equivalent.  $\square$

Proposition 2.3 tells us that for a toric manifold  $X$ , we may define the positivity (resp. non-negativity) of  $\text{ch}_k(X)$  as follows:

**Definition 2.4.** Let  $X$  be a toric manifold of  $\dim X = d$ . For  $1 \leq k \leq d$ , we say  $\text{ch}_k(X)$  is *positive* (resp. *nef*) if  $(\text{ch}_k(X) \cdot Y) > 0$  (resp.  $\geq 0$ ) for any  $k$ -dimensional closed torus invariant subvariety  $Y \subset X$ . When  $\text{ch}_k(X)$  is positive (resp. nef), we say  $X$  is  *$\text{ch}_k$ -positive* (resp.  *$\text{ch}_k$ -nef*).

**Remark 2.5.** When  $k = 1$ , we have  $\text{ch}_1(X) = c_1(X) = -K_X$ . So, a  $\text{ch}_1$ -positive (resp.  $\text{ch}_1$ -nef) toric manifold is nothing but a toric Fano (resp. weak Fano) manifold.

It is not difficult to see that the  $d$ -dimensional projective space  $\mathbb{P}^d$  is  $\text{ch}_k$ -positive for  $1 \leq k \leq d$ . Moreover, for  $k = 2$ , the following holds:

**Theorem 2.6** ([S2, Theorem 1.3]). *Let  $X = X_\Sigma$  be a smooth projective toric  $d$ -fold. If  $X = X_\Sigma$  is  $\text{ch}_2$ -positive and  $X$  has a Fano contraction, then  $X \cong \mathbb{P}^d$ . In particular, if  $\rho(X) = 2$ , then  $X$  is not  $\text{ch}_2$ -positive.*

### 3. TORIC MANIFOLDS OF PICARD NUMBER THREE

In this section, we determine whether a projective toric manifold  $X$  of  $\rho(X) = 3$  is  $\text{ch}_2$ -positive or not. By Theorem 2.6, we may assume  $X$  has no Fano contraction. We use the notation of [B1] throughout this section:

**Theorem 3.1** ([B1, Theorem 6.6]). *Let  $X = X_\Sigma$  be a projective toric manifold of Picard number three. If  $X$  has no Fano contraction, then the fan  $\Sigma$  is explicitly described as follows: Let*

$$G(\Sigma) = \{v_1, \dots, v_{p_0}, y_1, \dots, y_{p_1}, z_1, \dots, z_{p_2}, t_1, \dots, t_{p_3}, u_1, \dots, u_{p_4}\},$$

where  $p_0, p_1, p_2, p_3, p_4$  are positive integers and  $p_0 + p_1 + p_2 + p_3 + p_4 - 3 = d := \dim X$ . Then, the primitive relations of  $\Sigma$  are

$$v_1 + \dots + v_{p_0} + y_1 + \dots + y_{p_1} = c_2 z_2 + \dots + c_{p_2} z_{p_2} + (b_1 + 1)t_1 + \dots + (b_{p_3} + 1)t_{p_3},$$

$$y_1 + \dots + y_{p_1} + z_1 + \dots + z_{p_2} = u_1 + \dots + u_{p_4},$$

$$z_1 + \dots + z_{p_2} + t_1 + \dots + t_{p_3} = 0,$$

$$t_1 + \dots + t_{p_3} + u_1 + \dots + u_{p_4} = y_1 + \dots + y_{p_1} \text{ and}$$

$$u_1 + \dots + u_{p_4} + v_1 + \dots + v_{p_0} = c_2 z_2 + \dots + c_{p_2} z_{p_2} + b_1 t_1 + \dots + b_{p_3} t_{p_3}$$

for  $b_1, \dots, b_{p_3}, c_2, \dots, c_{p_2} \in \mathbb{Z}_{\geq 0}$ .

We may assume  $c_2 = \min\{c_2, \dots, c_{p_2}\}$  and  $b_1 = \min\{b_1, \dots, b_{p_3}\}$ .

Let

$$V_1, \dots, V_{p_0}, Y_1, \dots, Y_{p_1}, Z_1, \dots, Z_{p_2}, T_1, \dots, T_{p_3}, U_1, \dots, U_{p_4}$$

be the torus invariant divisors in  $\text{Pic}(X)$  associated to

$$v_1, \dots, v_{p_0}, y_1, \dots, y_{p_1}, z_1, \dots, z_{p_2}, t_1, \dots, t_{p_3}, u_1, \dots, u_{p_4},$$

respectively. By calculating the rational functions for the dual basis of  $G(\Sigma) \setminus \{v_1, z_1, u_1\}$ , we have the relations

$$V_2 - V_1 = 0, \dots, V_{p_0} - V_1 = 0,$$

$$Y_1 + U_1 - V_1 = 0, \dots, Y_{p_1} + U_1 - V_1 = 0,$$

$$Z_2 - Z_1 + c_2 V_1 = 0, \dots, Z_{p_2} - Z_1 + c_{p_2} V_1 = 0,$$

$$T_1 - Z_1 - U_1 + (b_1 + 1)V_1 = 0, \dots, T_{p_3} - Z_1 - U_1 + (b_{p_3} + 1)V_1 = 0,$$

$$U_2 - U_1 = 0, \dots, U_{p_4} - U_1 = 0$$

in  $\text{Pic}(X)$ . In particular, we obtain

$$V_1 = \dots = V_{p_0}, \quad Y_1 = \dots = Y_{p_1}, \quad U_1 = \dots = U_{p_4}, \quad T_i = T_1 + (b_1 - b_i)V_1 \quad (2 \leq i \leq p_3).$$

On the other hand, the primitive collections of  $\Sigma$  tell us that

$$V_1 \cdots V_{p_0} \cdot Y_1 \cdots Y_{p_1} = 0, \quad Y_1 \cdots Y_{p_1} \cdot Z_1 \cdots Z_{p_2} = 0, \quad Z_1 \cdots Z_{p_2} \cdot T_1 \cdots T_{p_3} = 0, \\ T_1 \cdots T_{p_3} \cdot U_1 \cdots U_{p_4} = 0 \text{ and } U_1 \cdots U_{p_4} \cdot V_1 \cdots V_{p_0} = 0.$$

Finally, we should remark that

$$2\text{ch}_2(X) = V_1^2 + \cdots + V_{p_0}^2 + Y_1^2 + \cdots + Y_{p_1}^2 + Z_1^2 + \cdots + Z_{p_2}^2 + T_1^2 + \cdots + T_{p_3}^2 + U_1^2 + \cdots + U_{p_4}^2$$

for our calculation below.

First of all, we calculate  $(\text{ch}_2(X) \cdot S_1)$  for the torus invariant subsurface  $S_1$  associated to the  $(d-2)$ -dimensional cone  $\tau \in \Sigma$  such that  $\underline{G(\tau)} = \underline{G(\Sigma) \setminus \{v_1, y_1, z_1, t_1, u_1\}}$ . In this case,  $\rho(S_1) = 3$ . Since

$$S_1 = V_2 \cdots V_{p_0} \cdot Y_2 \cdots Y_{p_1} \cdot Z_2 \cdots Z_{p_2} \cdot T_2 \cdots T_{p_3} \cdot U_2 \cdots U_{p_4},$$

we have

$$V_1 \cdot Y_1 \cdot S_1 = Y_1 \cdot Z_1 \cdot S_1 = Z_1 \cdot T_1 \cdot S_1 = T_1 \cdot U_1 \cdot S_1 = U_1 \cdot V_1 \cdot S_1 = 0.$$

We can calculate the intersection numbers as follows:

$$\begin{aligned} (V_1^2 \cdot S_1) &= (V_1 \cdot (Y_1 + U_1) \cdot S_1) = 0, \\ (Y_1^2 \cdot S_1) &= (Y_1 \cdot (V_1 - U_1) \cdot S_1) = -1, \\ (U_1^2 \cdot S_1) &= (U_1 \cdot (V_1 - Y_1) \cdot S_1) = -1, \\ (Z_1^2 \cdot S_1) &= (Z_1 \cdot (T_1 - U_1 + (b_1 + 1)V_1) \cdot S_1) = -1 + b_1 + 1 = b_1, \\ (Z_2^2 \cdot S_1) &= ((Z_1 - c_2 V_1) \cdot (Z_1 - c_2 V_1) \cdot S_1) = (Z_1^2 \cdot S_1) - 2c_2(Z_1 \cdot V_1 \cdot S_1) + c_2^2(V_1^2 \cdot S_1) = b_1 - 2c_2, \\ &\vdots \\ (Z_{p_2}^2 \cdot S_1) &= b_1 - 2c_{p_2}, \\ (T_1^2 \cdot S_1) &= (T_1 \cdot (Z_1 + U_1 - (b_1 + 1)V_1) \cdot S_1) = -(b_1 + 1), \\ (T_2^2 \cdot S_1) &= ((T_1 + (b_1 - b_2)V_1) \cdot (Z_1 + U_1 - (b_2 + 1)V_1) \cdot S_1) \\ &= (T_1 \cdot (-(b_2 + 1)V_1) \cdot S_1) + ((b_1 - b_2)V_1 \cdot Z_1 \cdot S_1) + ((b_1 - b_2)V_1 \cdot (-(b_2 + 1)V_1) \cdot S_1) \\ &= -(b_2 + 1) + (b_1 - b_2) = b_1 - 2b_2 - 1, \\ &\vdots \\ (T_{p_3}^2 \cdot S_1) &= -(b_{p_3} + 1) + (b_1 - b_{p_3}) = b_1 - 2b_{p_3} - 1. \end{aligned}$$

Therefore,

$$\begin{aligned} 2(\text{ch}_2(X) \cdot S_1) &= -p_1 - p_4 + b_1 p_2 - 2(c_2 + \cdots + c_{p_2}) \\ &\quad - (b_1 + 1) + (b_1 - b_2 - (b_2 + 1)) + \cdots + (b_1 - b_{p_3} - (b_{p_3} + 1)). \end{aligned}$$

So, we have the following:

**Lemma 3.2.** *If  $\text{ch}_2(X)$  is nef, then  $b_1 \geq 1$  and  $p_2 > p_3$ . In particular,  $p_2 \geq 2$ .*

*Proof.* By the above calculation,

$$2(\text{ch}_2(X) \cdot S_1) \leq -p_1 - p_4 + b_1 p_2 - (b_1 + 1)p_3 = -p_1 - p_3 - p_4 + b_1(p_2 - p_3).$$

Thus,  $b_1$  and  $p_2 - p_3$  have to be positive.  $\square$

Next, we treat the case where the Picard number of the torus invariant subsurface  $S = S_\tau \subset X_\Sigma$  associated to a  $(d-2)$ -dimensional cone  $\tau \in \Sigma$  is two. In this case, we can apply the following result:

**Proposition 3.3** ([S1, Theorem 3.5 and Proposition 4.4]). *Let  $X = X_\Sigma$  be a projective toric manifold and  $S = S_\tau$  the torus invariant subsurface associated to a  $(d-2)$ -dimensional cone  $\tau \in \Sigma$  in  $X$  such that  $\rho(S) = 2$ . In this case,  $S$  is isomorphic to the Hirzebruch surface  $F_\alpha$  of degree  $\alpha \geq 0$ . Let  $G(\tau) = \{x_1, \dots, x_{d-2}\}$ . We have exactly four maximal cones*

$$\text{Cone}(G(\tau) \cup \{w_1, w_2\}), \text{Cone}(G(\tau) \cup \{w_2, w_3\}),$$

$$\text{Cone}(G(\tau) \cup \{w_3, w_4\}) \text{ and } \text{Cone}(G(\tau) \cup \{w_4, w_1\})$$

which contain  $\tau$  for  $w_1, w_2, w_3, w_4 \in G(\Sigma)$  such that

$$w_1 + w_3 + a_1 x_1 + \dots + a_{d-2} x_{d-2} = 0 \text{ and } w_2 + w_4 - \alpha w_1 + e_1 x_1 + \dots + e_{d-2} x_{d-2} = 0.$$

Then,

$$(\text{ch}_2(X) \cdot S) = \frac{1}{2} (\alpha(2 + a_1^2 + \dots + a_{d-2}^2) + 2(-\alpha + a_1 e_1 + \dots + a_{d-2} e_{d-2})).$$

We calculate the intersection number  $(\text{ch}_2(X) \cdot S)$  for two cases: (1)  $G(\tau) = G(\Sigma) \setminus \{v_1, z_1, z_2, t_1, t_2\}$  and (2)  $G(\tau) = G(\Sigma) \setminus \{v_1, y_1, z_1, z_2, u_1\}$ .

(1)  $G(\tau) = G(\Sigma) \setminus \{v_1, z_1, z_2, t_1, t_2\}$ . We have two linear relations

$$z_1 + z_2 + z_3 + \dots + z_{p_2} + y_1 + \dots + y_{p_1} - (u_1 + \dots + u_{p_4}) = 0 \text{ and}$$

$$t_1 + t_2 + t_3 + \dots + t_{p_3} + u_1 + \dots + u_{p_4} - (y_1 + \dots + y_{p_1}) = 0$$

for  $\text{Cone}(G(\tau) \cup \{t_1\})$  and  $\text{Cone}(G(\tau) \cup \{z_1\})$ , respectively. One can easily see that  $S \cong \mathbb{P}^1 \times \mathbb{P}^1$ , and we obtain

$$2(\text{ch}_2(X) \cdot S) = 2(-p_1 - p_4) < 0$$

by Proposition 3.3. So, we have the following by Lemma 3.2:

**Lemma 3.4.** *If  $\text{ch}_2(X)$  is nef, then  $p_3 = 1$ .*

*Proof.* If  $p_2 \geq 2$  and  $p_3 \geq 2$ , then  $(\text{ch}_2(X) \cdot S) < 0$ . Lemma 3.2 says that  $p_2 \geq 2$ . So, we have  $p_3 = 1$ .  $\square$

Therefore, we may assume  $p_3 = 1$  in the following calculation.

(2)  $G(\tau) = G(\Sigma) \setminus \{v_1, y_1, z_1, z_2, u_1\}$ . First, we remark that this subsurface always exists by Lemma 3.2. We have two linear relations

$$z_1 + z_2 + \dots + z_{p_2} + t_1 = 0 \text{ and}$$

$$v_1 + y_1 + v_2 + \dots + v_{p_0} + y_2 + \dots + y_{p_1} - (c_2 z_2 + \dots + c_{p_2} z_{p_2} + (b_1 + 1)t_1) = 0$$

for  $\text{Cone}(G(\tau) \cup \{v_1\})$  and  $\text{Cone}(G(\tau) \cup \{z_2\})$ , respectively. One can easily see that  $S \cong F_{c_2}$ , and we obtain

$$2(\text{ch}_2(X) \cdot S) = c_2(p_2 + 1) + 2(-(c_2 + \dots + c_{p_2} + b_1 + 1))$$

by Proposition 3.3. If  $p_2 \geq 3$ , we have

$$2(\text{ch}_2(X) \cdot S) = c_2 + (c_2 - c_2) + \dots + (c_2 - c_{p_2}) + c_2 - (c_2 + \dots + c_{p_2}) - 2(b_1 + 1) < 0.$$

Thus, we obtain the following lemma (see Lemma 3.2):

**Lemma 3.5.** *If  $\text{ch}_2(X)$  is nef, then  $p_2 = 2$  and  $c_2 - 2(b_1 + 1) \geq 0$ .*

Thus, we conclude the following:

**Theorem 3.6.** *Let  $X$  be a projective toric manifold. If  $\rho(X) = 3$  and  $X$  has no Fano contraction, then  $\text{ch}_2(X)$  is not nef.*

*Proof.* Suppose that  $\text{ch}_2(X)$  is nef. Then, we may assume that  $p_2 = 2$ ,  $p_3 = 1$ ,  $b_1 \geq 1$  and  $c_2 - 2(b_1 + 1) \geq 0$  by Lemmas 3.2, 3.4 and 3.5. However, the last inequality obviously contradicts the another inequality

$$\begin{aligned} 2(\text{ch}_2(X) \cdot S_1) &= -p_1 - p_4 + 2b_1 - 2c_2 - (b_1 + 1) \geq 0 \\ \iff b_1 &\geq 2c_2 + p_1 + p_4 + 1. \end{aligned}$$

Thus,  $\text{ch}_2(X)$  is not nef.  $\square$

By combining Theorems 2.6 and 3.6, we have the following:

**Corollary 3.7.** *If  $X = X_\Sigma$  is a  $\text{ch}_2$ -positive smooth projective toric  $d$ -fold and  $\rho(X) \leq 3$ , then  $X \cong \mathbb{P}^d$ .*

#### 4. POSITIVITIES OF HIGHER CHERN CHARACTERS

In this section we study toric manifolds  $X$  with  $\text{ch}_k(X)$  nef for  $k \geq 3$ . In the case  $d = k$ , it is not difficult to construct toric  $d$ -folds  $X$  with  $\text{ch}_d(X)$  positive.

**Example 4.1.** Let  $d \geq 3$  and  $a \geq 1$ . Let  $X = X_\Sigma$  be a toric manifold of  $\rho(X) = 2$  such that  $G(\Sigma) = \{x_1, x_2, x_3, y_1, \dots, y_{d-1}\}$  and the primitive relations of  $\Sigma$  are

$$x_1 + x_2 + x_3 = ay_1 \text{ and } y_1 + \dots + y_{d-1} = 0.$$

Then  $X$  is isomorphic to the  $\mathbb{P}^{d-2}$ -bundle

$$\mathbb{P}_{\mathbb{P}^2}(\mathcal{O}(a) \oplus \mathcal{O}^{\oplus d-2})$$

over  $\mathbb{P}^2$ . We denote by  $D_1, D_2, D_3, E_1, \dots, E_{d-1}$  the torus invariant divisors in  $\text{Pic}(X)$  associated to  $x_1, x_2, x_3, y_1, \dots, y_{d-1}$ , respectively. Then we have the relations

$$D_1 = D_2 = D_3 \text{ and } aD_3 + E_1 = E_2 = \dots = E_{d-1}$$

in  $\text{Pic}(X)$ . We can calculate the intersection numbers as follows:

$$D_1^d = D_2^d = D_3^d = 0,$$

$$E_1^d = E_1^2(E_2 - aD_3)^{d-2}$$

$$= E_1^2 \left( E_2^{d-2} - a(d-2)D_3 \cdot E_2^{d-3} + a^2 \frac{(d-2)(d-3)}{2} D_3^2 \cdot E_2^{d-4} \right)$$

$$= E_1^2 \cdot E_2 \cdots E_{d-1} - a(d-2)D_3 \cdot E_1^2 \cdot E_2 \cdots E_{d-2}$$

$$+ a^2 \frac{(d-2)(d-3)}{2} D_2 \cdot D_3 \cdot E_1^2 \cdot E_2 \cdots E_{d-3}$$

$$= 0 - a(d-2) \cdot (-a) + a^2 \frac{(d-2)(d-3)}{2}$$

$$= a^2(d-2) + a^2 \frac{(d-2)(d-3)}{2} > 0,$$

$$E_2^d = \dots = E_{d-1}^d = (aD_3 + E_1) \cdot E_2 \cdot E_3 \cdots E_{d-1}^2 = aD_3 \cdot E_2 \cdot E_3 \cdots E_{d-1}^2 = a \cdot a = a^2 > 0.$$

Thus,  $\text{ch}_d(X)$  is positive. Note that  $X$  is Fano if  $a \leq 2$  (this follows from Theorem 2.2).

Next we consider the case  $d > k$ . For odd  $k$ , there is a toric  $d$ -fold  $X$  with  $\text{ch}_k(X)$  positive. However, our example does not provide an example of a toric manifold with  $\text{ch}_k(X)$  positive for even  $k$ .

**Proposition 4.2.** *Let  $d > k \geq 3$  and  $a \geq 1$  with  $d - a^k \geq 1$ . Let  $X = X_\Sigma$  be a toric manifold of  $\rho(X) = 2$  such that  $G(\Sigma) = \{x_1, \dots, x_d, y_1, y_2\}$  and the primitive relations of  $\Sigma$  are*

$$x_1 + \dots + x_d = ay_1 \text{ and } y_1 + y_2 = 0.$$

*Then the following hold:*

- (1) If  $k$  is odd, then  $\text{ch}_k(X)$  is positive.  
(2) If  $k$  is even, then  $\text{ch}_k(X)$  is nef, but not positive.

**Remark 4.3.** In Proposition 4.2,  $X$  is isomorphic to the  $\mathbb{P}^1$ -bundle  $\mathbb{P}_{\mathbb{P}^{d-1}}(\mathcal{O}(a) \oplus \mathcal{O})$  over  $\mathbb{P}^{d-1}$ , and it is Fano by Theorem 2.2.

*Proof of Proposition 4.2.* We denote by  $D_1, \dots, D_d, E_1, E_2$  the torus invariant divisors in  $\text{Pic}(X)$  associated to  $x_1, \dots, x_d, y_1, y_2$ , respectively. Then we have the relations

$$D_1 = \dots = D_d \text{ and } aD_d + E_1 = E_2$$

in  $\text{Pic}(X)$ . Put  $\tau_1$  (resp.  $\tau_2$ ) as the  $(d-k)$ -dimensional cone in  $\Sigma$  whose generators are  $x_1, \dots, x_{d-k}$  (resp.  $x_1, \dots, x_{d-k-1}, y_1$ ). We denote by  $V_i$  the  $k$ -dimensional torus invariant closed subvariety of  $X$  associated to  $\tau_i$  for  $i = 1, 2$ . It follows from Proposition 2.3 and the relations in  $\text{Pic}(X)$  that  $\text{ch}_k(X)$  is positive (resp. nef) if and only if  $(\text{ch}_k(X) \cdot V_i)$  is positive (resp. non-negative) for each  $i = 1, 2$ . Since

$$\begin{aligned} D_1^k \cdot D_1 \cdots D_{d-k} &= D_2^k \cdot D_1 \cdots D_{d-k} = \dots = D_d^k \cdot D_1 \cdots D_{d-k} = 0, \\ E_1^k \cdot D_1 \cdots D_{d-k} &= D_1 \cdots D_{d-k} \cdot (E_2 - aD_d)^{k-2} \cdot E_1^2 = D_1 \cdots D_{d-k} \cdot (-aD_d)^{k-2} \cdot E_1^2 \\ &= (-a)^{k-2} D_1 \cdots D_{d-2} \cdot E_1^2 = (-a)^{k-1}, \\ E_2^k \cdot D_1 \cdots D_{d-k} &= D_1 \cdots D_{d-k} \cdot (aD_d + E_1)^{k-2} \cdot E_2^2 = D_1 \cdots D_{d-k} \cdot (aD_d)^{k-2} \cdot E_2^2 \\ &= a^{k-2} D_1 \cdots D_{d-2} \cdot E_2^2 = a^{k-1}, \end{aligned}$$

we have

$$(\text{ch}_k(X) \cdot V_1) = \begin{cases} 2a^{k-1} & (k \text{ is odd}), \\ 0 & (k \text{ is even}). \end{cases}$$

Since

$$\begin{aligned} D_1^k \cdot D_1 \cdots D_{d-k-1} \cdot E_1 &= D_2^k \cdot D_1 \cdots D_{d-k-1} \cdot E_1 = \dots = D_d^k \cdot D_1 \cdots D_{d-k-1} \cdot E_1 = 1, \\ E_1^k \cdot D_1 \cdots D_{d-k-1} \cdot E_1 &= D_1 \cdots D_{d-k-1} \cdot (E_2 - aD_d)^{k-1} \cdot E_1^2 \\ &= (-a)^{k-1} D_1 \cdots D_{d-2} \cdot E_1^2 = (-a)^k, \\ E_2^k \cdot D_1 \cdots D_{d-k-1} \cdot E_1 &= 0, \end{aligned}$$

we have

$$(\text{ch}_k(X) \cdot V_2) = \begin{cases} d - a^k & (k \text{ is odd}), \\ d + a^k & (k \text{ is even}). \end{cases}$$

Thus,  $\text{ch}_k(X)$  is nef for any  $k \geq 3$ , and it is positive if and only if  $k$  is odd.  $\square$

At present, the only known examples of toric  $d$ -folds with  $\text{ch}_4(X)$  positive for  $d \geq 5$  are projective spaces. Finally, we prove the following theorem:

**Theorem 4.4.** *Any smooth complete toric  $d$ -fold  $X$  of  $\rho(X) = 2$  with  $d \geq 5$  is not  $\text{ch}_4$ -positive.*

*Proof.* Any toric manifold of Picard number two is a projective space bundle over a projective space. Let  $s-1$  be the dimension of a fiber. If  $s = 2$ , then  $\text{ch}_4(X)$  is not positive by Proposition 4.2. Thus, it suffices to prove the assertion in the case  $s \geq 3$ .

(1) *The case where  $s = 3$ .* Let  $X = X_\Sigma$  be a toric manifold of  $\rho(X) = 2$  such that  $G(\Sigma) = \{x_1, \dots, x_{d-1}, y_1, y_2, y_3\}$  and the primitive relations of  $\Sigma$  are

$$x_1 + \dots + x_{d-1} = a_1 y_1 + a_2 y_2 \text{ and } y_1 + y_2 + y_3 = 0,$$

where  $a_1 \geq a_2 \geq 0$ . Then  $X$  is the  $\mathbb{P}^2$ -bundle  $\mathbb{P}_{\mathbb{P}^{d-2}}(\mathcal{O}(a_1) \oplus \mathcal{O}(a_2) \oplus \mathcal{O})$  over  $\mathbb{P}^{d-2}$ . We denote by  $D_1, \dots, D_{d-1}, E_1, E_2, E_3$  the torus invariant divisors in  $\text{Pic}(X)$  associated to  $x_1, \dots, x_{d-1}, y_1, y_2, y_3$ , respectively. Then we have the relations

$$D_1 = \dots = D_{d-1} \text{ and } a_i D_{d-1} + E_i - E_3 = 0$$



for  $i = 1, 2$  in  $\text{Pic}(X)$ . Put  $\tau$  as the  $(d-4)$ -dimensional cone in  $\Sigma$  whose generators are  $x_1, \dots, x_{d-5}, y_1$ . We denote by  $V$  the 4-dimensional torus invariant closed subvariety of  $X$  associated to  $\tau$ . We show  $(\text{ch}_4(X) \cdot V) \leq 0$ . Since  $D_i^4 \cdot D_1 \cdots D_{d-5} \cdot E_1 = 0$  for every  $i = 1, \dots, d-1$ ,  $E_2^4 \cdot D_1 \cdots D_{d-5} \cdot E_1 = -a_2^3$  and  $E_3^4 \cdot D_1 \cdots D_{d-5} \cdot E_1 = a_2^3$ , it suffices to show  $E_1^4 \cdot D_1 \cdots D_{d-5} \cdot E_1 \leq 0$ . We calculate the intersection number as follows:

$$\begin{aligned}
& E_1^4 \cdot D_1 \cdots D_{d-5} \cdot E_1 \\
&= D_1 \cdots D_{d-5} \cdot E_1^3 \cdot (E_2 - (a_1 - a_2)D_{d-1}) \cdot (E_3 - a_1 D_{d-1}) \\
&= D_1 \cdots D_{d-5} \cdot E_1^3 \cdot (-a_1 D_{d-1} \cdot E_2 - (a_1 - a_2)D_{d-1} \cdot E_3 + a_1(a_1 - a_2)D_{d-1}^2) \\
&= D_1 \cdots D_{d-5} \cdot E_1^3 \cdot (-a_1 D_{d-1} \cdot E_2 - (a_1 - a_2)D_{d-1} \cdot (a_2 D_{d-1} + E_2) + a_1(a_1 - a_2)D_{d-1}^2) \\
&= D_1 \cdots D_{d-5} \cdot E_1^3 \cdot ((a_2 - 2a_1)D_{d-1} \cdot E_2 + (a_1 - a_2)^2 D_{d-1}^2) \\
&= D_1 \cdots D_{d-5} \cdot E_1^2 \cdot (E_3 - a_1 D_{d-1}) \cdot ((a_2 - 2a_1)D_{d-1} \cdot E_2 + (a_1 - a_2)^2 D_{d-1}^2) \\
&= (a_1 - a_2)^2 D_1 \cdots D_{d-3} \cdot E_1^2 \cdot E_3 - a_1(a_2 - 2a_1)D_1 \cdots D_{d-3} \cdot E_1^2 \cdot E_2 \\
&\quad - a_1(a_1 - a_2)^2 D_1 \cdots D_{d-2} \cdot E_1^2 \\
&= -(a_1 - a_2)^3 + a_1^2(a_2 - 2a_1) - a_1(a_1 - a_2)^2 \leq 0.
\end{aligned}$$

Thus,  $(\text{ch}_4(X) \cdot V) \leq 0$ .

(2) *The case where  $s \geq 4$ .* Let  $X = X_\Sigma$  be a toric manifold of  $\rho(X) = 2$  such that  $G(\Sigma) = \{x_1, \dots, x_{d-s+2}, y_1, \dots, y_s\}$  and the primitive relations of  $\Sigma$  are

$$x_1 + \cdots + x_{d-s+2} = a_1 y_1 + \cdots + a_{s-1} y_{s-1} \text{ and } y_1 + \cdots + y_s = 0,$$

where  $a_1 \geq \cdots \geq a_{s-1} \geq 0$ . Then  $X$  is the  $\mathbb{P}^{s-1}$ -bundle  $\mathbb{P}_{\mathbb{P}^{d-s+1}}(\mathcal{O}(a_1) \oplus \cdots \oplus \mathcal{O}(a_{s-1}) \oplus \mathcal{O})$  over  $\mathbb{P}^{d-s+1}$ . We denote by  $D_1, \dots, D_{d-s+2}, E_1, \dots, E_s$  the torus invariant divisors in  $\text{Pic}(X)$  associated to  $x_1, \dots, x_{d-s+2}, y_1, \dots, y_s$ , respectively. Then we have the relations

$$D_1 = \cdots = D_{d-s+2} \text{ and } a_i D_{d-s+2} + E_i - E_s = 0$$

for  $i = 1, \dots, s-1$  in  $\text{Pic}(X)$ . Put  $\tau$  as the  $(d-4)$ -dimensional cone in  $\Sigma$  whose generators are  $x_1, \dots, x_{d-s}, y_1, \dots, y_{s-4}$ . We denote by  $V$  the 4-dimensional torus invariant closed subvariety of  $X$  associated to  $\tau$ . We show  $(\text{ch}_4(X) \cdot V) \leq 0$ . For every  $i = 1, \dots, d-s+2$ , we have  $D_i^4 \cdot D_1 \cdots D_{d-s} \cdot E_1 \cdots E_{s-4} = 0$ . We calculate

$$\begin{aligned}
& E_1^4 \cdot D_1 \cdots D_{d-s} \cdot E_1 \cdots E_{s-4} \\
&= D_1 \cdots D_{d-s} \cdot E_1^2 \cdot E_2 \cdots E_{s-4} \cdot (E_{s-3} + (a_{s-3} - a_1)D_{d-s+2}) \\
&\quad \cdot (E_{s-2} + (a_{s-2} - a_1)D_{d-s+2}) \cdot (E_{s-1} + (a_{s-1} - a_1)D_{d-s+2}) \\
&= D_1 \cdots D_{d-s} \cdot E_1^2 \cdot E_2 \cdots E_{s-4} \cdot (E_{s-3} \cdot E_{s-2} \cdot E_{s-1} + (a_{s-3} - a_1)D_{d-s+2} \cdot E_{s-2} \cdot E_{s-1} \\
&\quad + (a_{s-2} - a_1)D_{d-s+2} \cdot E_{s-3} \cdot E_{s-1} + (a_{s-1} - a_1)D_{d-s+2} \cdot E_{s-3} \cdot E_{s-2}) \\
&= -a_1 + (a_{s-3} - a_1) + (a_{s-2} - a_1) + (a_{s-1} - a_1) \leq 0.
\end{aligned}$$

Similarly,

$$E_i^4 \cdot D_1 \cdots D_{d-s} \cdot E_1 \cdots E_{s-4} = -a_i + (a_{s-3} - a_i) + (a_{s-2} - a_i) + (a_{s-1} - a_i) \leq 0$$

for every  $i = 2, \dots, s-4$ . Hence it suffices to show

$$(E_{s-3}^4 + E_{s-2}^4 + E_{s-1}^4 + E_s^4) \cdot D_1 \cdots D_{d-s} \cdot E_1 \cdots E_{s-4} \leq 0.$$

We calculate

$$\begin{aligned}
& E_{s-3}^4 \cdot D_1 \cdots D_{d-s} \cdot E_1 \cdots E_{s-4} \\
&= D_1 \cdots D_{d-s} \cdot E_1 \cdots E_{s-4} \cdot E_{s-3}^2 \\
&\cdot (E_{s-2} + (a_{s-2} - a_{s-3})D_{d-s+2}) \cdot (E_{s-1} + (a_{s-1} - a_{s-3})D_{d-s+2}) \\
&= D_1 \cdots D_{d-s} \cdot E_1 \cdots E_{s-4} \cdot E_{s-3}^2 \\
&\cdot (E_{s-2} \cdot E_{s-1} + (a_{s-2} - a_{s-3})D_{d-s+2} \cdot E_{s-1} + (a_{s-1} - a_{s-3})D_{d-s+2} \cdot E_{s-2}) \\
&= -a_{s-3} + (a_{s-2} - a_{s-3}) + (a_{s-1} - a_{s-3}) \\
&= (a_{s-3} + a_{s-2} + a_{s-1}) - 4a_{s-3}.
\end{aligned}$$

Similarly,

$$\begin{aligned}
E_{s-2}^4 \cdot D_1 \cdots D_{d-s} \cdot E_1 \cdots E_{s-4} &= (a_{s-3} + a_{s-2} + a_{s-1}) - 4a_{s-2}, \\
E_{s-1}^4 \cdot D_1 \cdots D_{d-s} \cdot E_1 \cdots E_{s-4} &= (a_{s-3} + a_{s-2} + a_{s-1}) - 4a_{s-1}.
\end{aligned}$$

Furthermore, we calculate

$$\begin{aligned}
& E_s^4 \cdot D_1 \cdots D_{d-s} \cdot E_1 \cdots E_{s-4} \\
&= D_1 \cdots D_{d-s} \cdot E_1 \cdots E_{s-4} \cdot E_s^2 \cdot (E_{s-2} + a_{s-2}D_{d-s+2}) \cdot (E_{s-1} + a_{s-1}D_{d-s+2}) \\
&= D_1 \cdots D_{d-s} \cdot E_1 \cdots E_{s-4} \cdot E_s^2 \cdot (E_{s-2} \cdot E_{s-1} + a_{s-2}D_{d-s+2} \cdot E_{s-1} + a_{s-1}D_{d-s+2} \cdot E_{s-2}) \\
&= a_{s-3} + a_{s-2} + a_{s-1}.
\end{aligned}$$

Hence we have

$$\begin{aligned}
& (E_{s-3}^4 + E_{s-2}^4 + E_{s-1}^4 + E_s^4) \cdot D_1 \cdots D_{d-s} \cdot E_1 \cdots E_{s-4} \\
&= 4(a_{s-3} + a_{s-2} + a_{s-1}) - 4a_{s-3} - 4a_{s-2} - 4a_{s-1} = 0.
\end{aligned}$$

Thus,  $(\text{ch}_4(X) \cdot V) \leq 0$ .

In every case,  $\text{ch}_4(X)$  is not positive. □

## REFERENCES

- [AC] C. Araujo and A-M. Castravet, Classification of 2-Fano manifolds with high index, A Celebration of Algebraic Geometry, Clay Math. Proc., vol. 18, Amer. Math. Soc., Providence, RI, 2013, 1–36.
- [B1] V. Batyrev, On the classification of smooth projective toric varieties, Tohoku Math. J. **43** (1991), no. 4, 569–585.
- [B2] V. Batyrev, On the classification of toric Fano 4-folds, Algebraic geometry, 9. J. Math. Sci. (New York) **94** (1999), 1021–1050.
- [CLS] D. A. Cox, J. B. Little, H. K. Schenck, *Toric varieties*, Graduate Studies in Mathematics, **124**. American Mathematical Society, Providence, RI, 2011.
- [JS] A. J. de Jong and J. Starr, Higher Fano manifolds and rational surfaces, Duke Math. J. **139** (2007), no. 1, 173–183.
- [FS] O. Fujino and H. Sato, Introduction to the toric Mori theory, Michigan Math. J. **52** (2004), no. 3, 649–665.
- [F] W. Fulton, *Introduction to toric varieties*, Annals of Mathematics Studies, **131**. The William H. Roever Lectures in Geometry. Princeton University Press, Princeton, NJ, 1993.
- [M] K. Matsuki, *Introduction to the Mori program*, Universitext, Springer-Verlag, New York, 2002.
- [O] T. Oda, *Convex bodies and algebraic geometry, An introduction to the theory of toric varieties*, Translated from the Japanese, Results in Mathematics and Related Areas (3) **15**, Springer-Verlag, Berlin, 1988.
- [R] M. Reid, Decomposition of toric morphisms, *Arithmetic and geometry, Vol. II*, 395–418, Progr. Math. **36**, Birkhäuser Boston, Boston, MA, 1983.
- [S1] H. Sato, The numerical class of a surface on a toric manifold, Int. J. Math. Math. Sci. 2012, 9 pages.
- [S2] H. Sato, Toric 2-Fano manifolds and extremal contractions, Proc. Japan Acad. Ser. A Math. Sci. **92** (2016), no. 10, 121–124.

DEPARTMENT OF APPLIED MATHEMATICS, FACULTY OF SCIENCES, FUKUOKA UNIVERSITY, 8-19-1,  
NANAKUMA, JONAN-KU, FUKUOKA 814-0180, JAPAN

*E-mail address:* `hirosato@fukuoka-u.ac.jp`

DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, OSAKA UNIVERSITY, TOYON-  
AKA, OSAKA 560-0043, JAPAN

*E-mail address:* `y-suyama@cr.math.sci.osaka-u.ac.jp`