

# One-dimensional optimal system for 2D Rotating Ideal Gas

Andronikos Paliathanasis\*

*Institute of Systems Science, Durban University of Technology*

*PO Box 1334, Durban 4000, Republic of South Africa*

September 4, 2019

## Abstract

We derive the one-dimensional optimal system for a system of three partial differential equations which describe the two-dimensional rotating ideal gas with polytropic parameter  $\gamma > 2$ . The Lie symmetries and the one-dimensional optimal system are determined for the nonrotating and rotating systems. We compare the results and we found that when there is no Coriolis force the system admits eight Lie point symmetries, while the rotating system admits seven Lie point symmetries. Consequently the two systems are not algebraic equivalent as in the case of  $\gamma = 2$  which was found by previous studies. For the one-dimensional optimal system we determine all the Lie invariants, while we demonstrate our results by reducing the system of partial differential equations into a system of first-order ordinary differential equations which can be solved by quadratures.

Keywords: Lie symmetries; invariants; shallow water; similarity solutions

## 1 Introduction

A powerful mathematical treatment for the determination of exact solutions for nonlinear differential equations is the Lie symmetry analysis [1–3]. Specifically, Lie point symmetries help us in the simplification of differential equations by means of similarity transformations which reduce the differential equation. The reduction process is based on the existence of functions which are invariant under a specific group of point transformations. When someone uses these invariants as new dependent and independent variables the differential equation is reduced. The reduction process differs between ordinary differential equations (ODEs) and partial differential equations (PDEs). For ODEs Lie point symmetries are applied to reduce the order of ODE by one; while on PDEs Lie point symmetries are applied to reduce by one the number of independent variables, while the order of the PDE remain the same. The solutions which are found with the application of those invariant functions are called similarity solutions. Some applications on the determination of similarity solutions for nonlinear differential equations can be found in [6–11] and references therein.

A common characteristic in the reduction process is that the Lie point symmetries are not preserved during the reduction, hence we can say that the symmetries can be lost. That is not an accurate statement, because

---

\*Email: anpaliat@phys.uoa.gr

symmetries are not “destroyed” or “created” under point transformations but the “nature” of the symmetry change. In addition, Lie symmetries can be used to construct new similarity solutions for a given differential equation by applying the adjoint representation of the Lie group [5].

It is possible that a given differential equation to admit more than one similarity solution when the given differential equation admits a “large” number of Lie point symmetries. Hence, in order for someone to classify a differential equation according to the admitted similarity solutions, all the inequivalent Lie subalgebras of the admitted Lie symmetries should be determined.

The first group classification problem was carried out by Ovsianikov [4] who demonstrated the construction of the one-dimensional optimal system for the Lie algebra. Since then, the classification of the one-dimensional optimal system has become a main tool for the study of nonlinear differential equations [12–15].

In this work, we focus on the classification of the one-dimensional optimal system for the two-dimensional rotating ideal gas system for a described by the following system of PDEs [16–18]

$$h_t + (hu)_x + (hv)_y = 0, \quad (1)$$

$$u_t + uu_x + vu_y + h^{\gamma-2}h_x - fv = 0, \quad (2)$$

$$v_t + uv_x + vv_y + h^{\gamma-2}h_y + fu = 0. \quad (3)$$

where  $u$  and  $v$  are the velocity components on the  $x$  and  $y$  directions respectively,  $h$  is the density of the ideal gas,  $f$  is the Coriolis parameter and  $\gamma$  is the polytropic parameter of the fluid. Usually  $\gamma$  is assumed to be  $\gamma = 2$  where equations (1)-(3) reduce to the shallow water system. However in this work we consider that  $\gamma > 2$ . In this work, polytropic index  $\gamma$  is defined as  $\frac{C_p}{C_v} = \gamma - 1$ .

Shallow-water equations describe the flow of a fluid under a pressure surface. There are various physical phenomena which are described by the Shallow-water system with emphasis on atmospheric and oceanic phenomena [19–21]. Hence, the existence of the Coriolis force it becomes critical in the description of the physical phenomena.

In the case of  $\gamma = 2$ , the complete symmetry analysis of the system (1)-(3) is presented in [22]. It was found that for  $\gamma = 2$  the given system of PDEs is invariant under a nine-dimensional Lie algebra. The same Lie algebra but in a different representation is also admitted by the nonrotating system, i.e.  $f = 0$ . One of the main results of [22] is that the transformation which relates the two representations of the admitted Lie algebras for the rotating and nonrotating system, transform the rotating system (1)-(3) into the nonrotating one. For other applications of Lie symmetries on Shallow-water equation we refer the reader in [23–28].

For the case of an ideal gas [17], i.e. parameter  $\gamma > 1$  from our analysis it follows that this property is lost. The nonrotating system and the rotating one are invariant under a different number of Lie symmetries and consequently under different Lie algebras. For each of the Lie algebras we æther the one-dimensional optimal system and all the Lie invariants. The results are presented in tables. We demonstrate the application of the Lie invariants by determining some similarity solutions for the system (1)-(3) for  $\gamma > 2$ . The paper is structured as follows.

In Section 2. we briefly discuss the theory of Lie symmetries for differential equations and the adjoint representation. The nonrotating system (1)-(3) is studied in Section 3. Specifically we determine the Lie points symmetries which form an eight-dimensional Lie algebra. The commutators and the adjoint representation are presented. We make use of these results and we perform a classification of the one-dimensional optimal system. We found that in total there are twenty-three one-dimensional independent Lie symmetries and possible

reductions, the corresponding invariants are determined and presented in tables. In Section 4 we perform the same analysis for the rotating system. There we find that the admitted Lie symmetries form a seven-dimensional Lie algebra while there are twenty independent one-dimensional Lie algebras. We demonstrate the results by reducing the system of PDEs (1)-(3) into an integrable system of three first-order ODEs which solution is given by quadratures. In Section 5 we discuss our results and draw our conclusions. Finally, in Appendix A we present the tables which includes the results of our analysis.

## 2 Lie symmetry analysis

Let  $H^A(x^i, \Phi^A, \Phi_i^A, \dots) = 0$  be a system of partial differential equations (PDEs) where  $\Phi^A$  denotes the dependent variables and  $x^i$  are the independent variables, at this point it is important to mention that we make use of the Einstein summation convention. By definition under the action of the infinitesimal one-parameter point transformation (1PPT)

$$\bar{x}^i = x^i(x^j, \Phi^B; \varepsilon), \quad \bar{\Phi}^A = \Phi^A(x^j, \Phi^B; \varepsilon), \quad (4)$$

which connects two different points  $P(x^j, \Phi^B) \rightarrow Q(\bar{x}^j, \bar{\Phi}^B, \varepsilon)$ , the differential equation  $H^A = 0$  remains invariant if and only if  $\bar{H}^A = H^A$ , that is [2]

$$\lim_{\varepsilon \rightarrow 0} \frac{\bar{H}^A(\bar{y}^i, \bar{u}^A, \dots; \varepsilon) - H^A(y^i, u^A, \dots)}{\varepsilon} = 0. \quad (5)$$

The latter condition means that the  $\Phi^A(P)$  and  $\Phi^A(Q)$  are connected through the transformation.

The lhs of expression (5) defines the Lie derivative of  $H^A$  along the vector field  $X$  of the one-parameter point transformation (4), in which  $X$  is defined as

$$X = \frac{\partial \bar{x}^i}{\partial \varepsilon} \partial_i + \frac{\partial \bar{\Phi}^A}{\partial \varepsilon} \partial_A.$$

Thus, condition (5) is equivalent with the following expression [2]

$$\mathcal{L}_X(H^A) = 0, \quad (6)$$

where  $\mathcal{L}$  denotes the Lie derivative with respect to the vector field  $X^{[n]}$  which is the  $n$ th-extension of generator  $X$  of the transformation (4) in the jet space  $\{x^i, \Phi^A, \Phi_{,i}^A, \Phi_{,ij}^A, \dots\}$  given by the expression [2]

$$X^{[n]} = X + \eta^{[1]} \partial_{\Phi_i^A} + \dots + \eta^{[n]} \partial_{\Phi_{i_1 i_2 \dots i_n}^A}, \quad (7)$$

in which

$$\eta^{[n]} = D_i \eta^{[n-1]} - u_{i_1 i_2 \dots i_{n-1}} D_i \left( \frac{\partial \bar{x}^j}{\partial \varepsilon} \right), \quad i \succeq 1, \quad \eta^{[0]} = \left( \frac{\partial \bar{\Phi}^A}{\partial \varepsilon} \right). \quad (8)$$

Conditions (6) provides a system of PDEs whose solution determine the components of the  $X$ , consequently the infinitesimal transformation. The vector fields  $X$  which satisfy condition (6) are called Lie symmetries for the differential equation  $H^A = 0$ . The Lie symmetries for a given differential equation form a Lie algebra.

Lie symmetries can be used by different ways [2] in order to study a differential equation. However, their direct application is on the determination of the so-called similarity solutions. The steps which we follow to determine a similarity solution is based on the determination and application of the Lie invariant functions.

Let  $X$  be a Lie symmetry for a given differential equation  $H^A = 0$ , then the differential equation  $X(F) = 0$ , where  $F$  is a function, provides the Lie invariants where by replacing in the differential equation  $H^A = 0$ , we reduce the number of the independent variables (in the case of PDEs) or the order of the differential equation (in the case of ordinary differential equations (ODEs)).

## 2.1 Optimal system

Consider the  $n$ -dimensional Lie algebra  $G_n$  with elements  $X_1, X_2, \dots, X_n$ . Then we shall say that the two vector fields [2]

$$Z = \sum_{i=1}^n a_i X_i, \quad W = \sum_{i=1}^n b_i X_i, \quad a_i, b_i \text{ are constants.} \quad (9)$$

are equivalent iff there

$$\mathbf{W} = \lim_{\varepsilon \rightarrow 0} \text{Ad}(\exp(\varepsilon X_i)) \mathbf{Z} \quad (10)$$

or

$$W = cZ, \quad c = \text{const.} \quad (11)$$

where the operator [2]

$$\text{Ad}(\exp(\varepsilon X_i)) X_j = X_j - \varepsilon [X_i, X_j] + \frac{1}{2} \varepsilon^2 [X_i, [X_i, X_j]] + \dots \quad (12)$$

is called the adjoint representation.

Therefore, in order to perform a complete classification for the similarity solutions of a given differential equation we should determine all the one-dimensional independent symmetry vectors of the Lie algebra  $G_n$ .

We continue our analysis by calculating the Lie point symmetries for the system (1)-(3) for the case where the system is rotating ( $f \neq 0$ ) and nonrotating ( $f = 0$ ).

## 3 Symmetries and optimal system for nonrotating shallow water

We start our analysis by applying the symmetry condition (6) for the Coriolis free system (1)-(3) with  $f = 0$ . We found that the system of PDEs admit eight Lie point symmetries as they are presented in the following [4]

$$\begin{aligned} X_1 &= \partial_t, \quad X_2 = \partial_x, \quad X_3 = \partial_y, \\ X_4 &= t\partial_x + \partial_u, \quad X_5 = t\partial_y + \partial_v, \\ X_6 &= y\partial_x - x\partial_y + v\partial_u - u\partial_v, \\ X_7 &= t\partial_t + x\partial_x + y\partial_y, \\ X_8 &= (\gamma - 1)(x\partial_x + y\partial_y + u\partial_u + v\partial_v) + 2h\partial_h. \end{aligned}$$

The commutators of the Lie symmetries and the adjoint representation are presented in tables 1 and 3 respectively.

We continue by determine the one-dimensional optimal system. Let us consider the generic symmetry vector

$$Z^8 = a_1 X_1 + a_2 X_2 + a_3 X_3 + a_4 X_4 + a_5 X_5 + a_6 X_6 + a_7 X_7 + a_8 X_8$$

From table 3 we see that by applying the following adjoint representations

$$Z'^8 = \text{Ad}(\exp(\varepsilon_5 X_5)) \text{Ad}(\exp(\varepsilon_4 X_4)) \text{Ad}(\exp(\varepsilon_3 X_3)) \text{Ad}(\exp(\varepsilon_2 X_2)) \text{Ad}(\exp(\varepsilon_1 X_1)) Z^8$$

parameters  $\varepsilon_1, \varepsilon_2, \varepsilon_3, \varepsilon_4$  and  $\varepsilon_5$  can be determined such that

$$Z'^8 = a'_6 X_6 + a'_7 X_7 + a'_8 X_8$$

Table 1: Commutators of the admitted Lie point symmetries for the nonrotating 2D Shallow water

$[ , ]$	$\mathbf{X}_1$	$\mathbf{X}_2$	$\mathbf{X}_3$	$\mathbf{X}_4$	$\mathbf{X}_5$	$\mathbf{X}_6$	$\mathbf{X}_7$	$\mathbf{X}_8$
$\mathbf{X}_1$	0	0	0	$X_2$	$X_3$	0	$-(\gamma - 1)X_1$	0
$\mathbf{X}_2$	0	0	0	0	0	$-X_3$	0	$(\gamma - 1)X_2$
$\mathbf{X}_3$	0	0	0	0	0	$X_2$	0	$(\gamma - 1)X_3$
$\mathbf{X}_4$	$-X_2$	0	0	0	0	$-X_5$	$(\gamma - 1)X_4$	$(\gamma - 1)X_4$
$\mathbf{X}_5$	$-X_3$	0	0	0	0	$X_4$	$(\gamma - 1)X_5$	$(\gamma - 1)X_5$
$\mathbf{X}_6$	0	$X_3$	$-X_2$	$X_5$	$-X_4$	0	0	0
$\mathbf{X}_7$	$(\gamma - 1)X_1$	0	0	$-(\gamma - 1)X_4$	$-(\gamma - 1)X_5$	0	0	0
$\mathbf{X}_8$	0	$-(\gamma - 1)X_2$	$-(\gamma - 1)X_3$	$-(\gamma - 1)X_4$	$-(\gamma - 1)X_5$	0	0	0

Parameters  $a_6$ ,  $a_7$ , and  $a_8$  are the relative invariants of the full adjoint action. Indeed in order to determine the relative invariants we solve the following system of partial differential equation [1]

$$\Delta(\phi(a_i)) = C_{ij}^k a^i \frac{\partial}{\partial a_j}$$

where  $C_{ij}^k$  are the structure constants of the admitted Lie algebra as they presented in Table 1. Consequently, in order to derive all the possible one-dimensional Lie symmetries we should study various cases where non of the invariants are zero, one of the invariants are zero, two of the invariants are zero or all the invariants are zero.

Hence, for the first three cases infer the following one-dimensional independent Lie algebras

$$X_6, X_7, X_8, \xi_{(67)} = X_6 + \alpha X_7, \xi_{(68)} = X_6 + \alpha X_8$$

$$\xi_{(78)} = X_7 + \alpha X_8, \xi_{(678)} = X_6 + \alpha X_7 + \beta X_8.$$

We apply the same procedure for the rest of the possible linear combinations of the symmetry vectors and we find the one-dimensional dependent Lie algebras

$$X_1, X_2, X_3, X_4, X_5, \xi_{(12)} = X_1 + \alpha X_2, \xi_{(13)} = X_1 + \alpha X_3, \xi_{(23)} = X_2 + \alpha X_3, \xi_{(14)} = X_1 + \alpha X_4,$$

$$\xi_{(15)} = X_1 + \alpha X_5, \xi_{(16)} = X_1 + \alpha X_6, \xi_{(34)} = X_3 + \alpha X_4, \xi_{(25)} = X_2 + \alpha X_5, \xi_{(45)} = X_4 + \alpha X_5,$$

$$\xi_{(123)} = X_1 + \alpha X_2 + \beta X_3, \xi_{(145)} = X_1 + \alpha X_4 + \beta X_5, \xi_{(125)} = X_1 + \alpha X_2 + \beta X_5, \xi_{(134)} = X_1 + \alpha X_3 + \beta X_4,$$

in which  $\alpha$  and  $\beta$  are constants.

Therefore by applying one of the above Lie symmetry vectors we find all the possible reductions from a system of 1 + 2 PDEs to a system of 1 + 1 PDEs. The reduced system will not admit all the remaining Lie symmetries. The Lie symmetries which survive under a reduction process are given as described in the following example.

Let a PDE admits the Lie point symmetries  $\Gamma_1, \Gamma_2$  which are such that  $[\Gamma_1, \Gamma_2] = C_{12}^1 X_1$ , with  $C_{12}^1 \neq 0$ . Reduction with the symmetry vector  $\Gamma_1$  leads to a reduced differential equation which admits  $\Gamma_2$  as Lie symmetry. On the other hand, reduction of the mother equation with respect to the Lie symmetry  $\Gamma_2$  leads to a different reduced differential equation which does not admit as a Lie point symmetry the vector field  $\Gamma_1$ .

In case the two Lie symmetries form an Abelian Lie algebra, i.e.  $C_{12}^1 = 0$ , then under any reduction process symmetries are preserved by any reduction.

We found that the optimal system admit twenty-three one-dimensional Lie symmetries, and possible independent reductions. All the possible twenty-three Lie invariants are presented in tables 4 and 5.

An application of the Lie invariants is presented below.

### 3.1 Application of $\xi_{145}$

Let us now demonstrate the results of tables 4 and 5 by the Lie invariants of the symmetry vector  $\xi_{145}$  and construct the similarity solution for the system.

The application of  $\xi_{145}$  in the nonrotating system (1)-(3) reduce the PDEs in the following system

$$(hu)_z + (hv)_w = 0 \quad (13)$$

$$\alpha + uu_z + vu_w + h^{\gamma-2}h_z = 0 \quad (14)$$

$$\beta + uv_z + vv_w + h^{\gamma-2}h_w = 0 \quad (15)$$

where  $z = x - \frac{\alpha}{2}t^2$  and  $w = y - \frac{\beta}{2}t^2$ .

System (13)-(15) admits the Lie point symmetries

$$\partial_z, \partial_w, z\partial_z + w\partial_w + \frac{2}{\gamma-1}h\partial_h + u\partial_u + v\partial_v \quad (16)$$

Reduction with the symmetry vector  $\partial_z + c\partial_w$  provides the following system of first-order ODEs

$$Fh_\sigma = (c\alpha - \beta)h^2, \quad (17)$$

$$Fv_\sigma = \frac{(\alpha - c\beta)ch^\gamma - \alpha h(v - cu)^2}{v - cu}, \quad (18)$$

$$Fh_\sigma = \frac{(\alpha - c\beta)cu^\gamma - \beta h(v - cu)^2}{v - cu}. \quad (19)$$

where  $F = (1 + c^2)h^\gamma - h(v - cu)^2$  and  $\sigma = z + cw$ .

By performing the change of variable  $d\sigma = f d\tau$  function  $f$  can be removed from the above system. For  $h(\tau) = 0$ , system (17), (18), (19) admits a solution  $u = u_0, v = v_0$  which is a critical point. The latter special solutions are always unstable when  $\alpha c > \beta$ .

We proceed our analysis by considering the rotating system.

## 4 Symmetries and optimal system for rotating shallow water

For the rotating system ( $f \neq 0$ ), the Lie symmetries are

$$Y_1 = \partial_t, Y_2 = \partial_x, Y_3 = \partial_y,$$

$$Y_4 = y\partial_x - x\partial_y + v\partial_u - u\partial_v,$$

$$Y_5 = \sin(ft)\partial_x + \cos(ft)\partial_y + f(\cos(ft)\partial_u - \sin(ft)\partial_v)$$

$$Y_6 = \cos(ft)\partial_x - \sin(ft)\partial_y - f(\sin(ft)\partial_u + \cos(ft)\partial_v)$$

$$Y_7 = (\gamma - 1)(x\partial_x + y\partial_y + u\partial_u + v\partial_v) + 2h\partial_h$$

Table 2: Commutators of the admitted Lie point symmetries for the rotating 2D Shallow water

$[ , ]$	$\mathbf{Y}_1$	$\mathbf{Y}_2$	$\mathbf{Y}_3$	$\mathbf{Y}_4$	$\mathbf{Y}_5$	$\mathbf{Y}_6$	$\mathbf{Y}_7$
$\mathbf{Y}_1$	0	0	0	0	$fY_6$	$-fY_5$	0
$\mathbf{Y}_2$	0	0	0	$-Y_3$	0	0	$(\gamma - 1)Y_2$
$\mathbf{Y}_3$	0	0	0	$Y_2$	0	0	$(\gamma - 1)Y_3$
$\mathbf{Y}_4$	0	$Y_3$	$-Y_2$	0	$-Y_6$	$Y_5$	0
$\mathbf{Y}_5$	$-fY_6$	0	0	$Y_6$	0	0	$(\gamma - 1)Y_5$
$\mathbf{Y}_6$	$fY_5$	0	0	$-Y_5$	0	0	$(\gamma - 1)Y_6$
$\mathbf{Y}_7$	0	$-(\gamma - 1)Y_2$	$-(\gamma - 1)Y_3$	0	$-(\gamma - 1)Y_5$	$-(\gamma - 1)Y_6$	0

The commutators and the adjoint representation are given in tables 2 and 6. The Lie symmetries for the rotating system form a smaller dimension Lie algebra than the non-rotating system. That is not the case when  $\gamma = 2$ , where the two Lie algebras have the same dimensional and are equivalent under point transformation [22]. Therefore, for  $\gamma > 2$  the Coriolis force cannot be eliminated by a point transformation as in the  $\gamma = 2$  case.

As far as the admitted Lie symmetries admitted by the given system of PDEs with or without the Coriolis terms for  $\gamma > 2$ , we remark that the rotating and the nonrotating system have a common Lie subalgebra of one-parameter point transformations consists by the symmetry vectors  $Y_1, Y_2, Y_3, Y_4$  and  $Y_7$ , or for the nonrotating system  $X_1, X_2, X_3, X_6$  and  $X_8$ .

We proceed with the determination of the one-dimensional optimal system and the invariant functions. Specifically, the relative invariants for the adjoint representation are calculated to be  $a_1, a_7$  and  $a_8$ . From tables 2 and 6 we can find the one-dimensional optimal system, which is

$$\begin{aligned}
 & Y_1, Y_2, Y_3, Y_4, Y_5, Y_6, Y_7, \chi_{12} = Y_1 + \alpha Y_2, \chi_{13} = Y_1 + \alpha Y_3, \\
 & \chi_{14} = Y_1 + \alpha Y_4, \chi_{15} = Y_1 + \alpha Y_5, \chi_{16} = Y_1 + \alpha Y_6, \chi_{17} = Y_1 + \alpha Y_7, \\
 & \chi_{23} = Y_2 + \alpha Y_3, \chi_{45} = Y_4 + \alpha Y_5, \chi_{46} = Y_4 + \alpha Y_6, \chi_{56} = Y_5 + \alpha Y_6 \\
 & \chi_{47} = Y_4 + \alpha Y_6, \chi_{123} = Y_1 + \alpha Y_2 + \beta Y_3, \chi_{147} = Y_1 + \alpha Y_4 + \beta Y_7.
 \end{aligned}$$

The Lie invariants which correspond to all the above one-dimensional Lie algebras are presented in tables 7 and 8.

Let us demonstrate the application of the Lie invariants by the following applications, from where we can see that the Lie invariants reduce the nonlinear field equations into a system of integrable first-order ODE which can be solved with quadratures.

#### 4.1 Application of $\chi_{12}$

We consider the travel-wave similarity solution in the  $x$ -plane provided by the symmetry vector  $\chi_{12}$ , and the vector field  $Y_3$ . The resulting equations is described by the following system of first order ODEs

$$v_z = f \frac{u}{\alpha - u} \quad (20)$$

$$\bar{F}u_z = f(\alpha - u)vh \quad (21)$$

$$\bar{F}h_z = fvh^2 \quad (22)$$

where  $\bar{F} = h^\gamma - (a - u)^2 h$  and  $z = t - \alpha x$ . Because we performed reduction with a subalgebra admitted by the nonrotating system, by setting  $f = 0$  in (20), (21), (22) we get the similarity solution for the nonrotating system where in this case is found to be  $h(z) = h_0$ ,  $u(z) = u_0$  and  $v(z) = v_0$ .

We perform the substitution  $dz = \frac{\bar{F}}{f v} d\tau$  and the latter system is simplified as follows

$$\frac{v}{\bar{F}} v_\tau = \frac{u}{\alpha - u} \quad (23)$$

$$u_\tau = (\alpha - u) h \quad (24)$$

$$h_\tau = h^2 \quad (25)$$

from where we get the solution

$$h(\tau) = (h_0 - \tau)^{-1}, \quad u(\tau) = \alpha + u_0 - \frac{u_0}{h_0} \tau \quad (26)$$

and

$$v(t)^2 = 2 \int \frac{\left(a + u_0 - \frac{u_0}{h_0} \tau\right)}{\frac{u_0}{h_0} (h_0 - \tau)} \left( (h_0 - \tau)^{-\gamma} + \left(\frac{u_0}{h_0}\right)^2 \tau - \frac{(u_0)^2}{h_0} \right) d\tau. \quad (27)$$

## 4.2 Application of $\chi_{23}$

Consider now the reduction with the symmetry vector fields  $\chi_{23}$ . The resulting system of 1+1 differential equations admit five Lie point symmetries, they are

$$\begin{aligned} & \partial_t, \partial_w, (\sin(ft) + \alpha \cos(ft)) \partial_w + f(\sin(ft) \partial_u + \cos(ft) \partial_v) \\ & (\alpha \sin(ft) - \cos(ft)) \partial_w - f(\cos(ft) \partial_u - \sin(ft) \partial_v), (\gamma - 1)(\partial_w + u \partial_u + v \partial_v) + 2h \partial_h. \end{aligned}$$

where  $w = y - \alpha x$ . For simplicity on our calculations let us assume  $\gamma = 3$ .

Reduction with the scaling symmetry provides the following system of first order ODEs

$$H_t = 2H(\alpha U - V), \quad (28)$$

$$U_t = \alpha H^2 + u(\alpha U - V) + fV, \quad (29)$$

$$V_t = -H^2 - v(\alpha U - V) - fU, \quad (30)$$

where  $h = wH$ ,  $u = wU$  and  $v = wV$ . The latter system is integrable and can be solved with quadratures.

Reduction with respect the symmetry vector  $(\alpha \sin(ft) - \cos(ft)) \partial_w - f(\cos(ft) \partial_u - \sin(ft) \partial_v)$  we find the reduced system

$$\frac{H_t}{H} = -\frac{\alpha \cos(ft) + \sin(ft)}{\cos(ft) - \alpha \sin(ft)}, \quad (31)$$

$$U_t = -\alpha f \frac{\sin(ft) V - \cos(ft) U}{\cos(ft) - \alpha \sin(ft)}, \quad (32)$$

$$V_t = -f \frac{\sin(ft) V - \cos(ft) U}{\cos(ft) - \alpha \sin(ft)}, \quad (33)$$

where now

$$h = H(t), \quad (34)$$

$$u = \frac{\cos(ft)}{\cos(ft) - \alpha \sin(ft)} f w + U(t), \quad (35)$$

$$v = -\frac{\sin(ft)}{\cos(ft) - \alpha \sin(ft)} f w + V(t). \quad (36)$$

System (31), (32), (33) is integrable and the solution is expressed in terms of quadratures.

## 5 Conclusions

In this work, we determined the one-dimensional optimal system for the two-dimensional ideal gas equations. The nonrotating system it was found that is invariant under an eight-dimensional group of one-parameter point transformations. and there are twenty-three independent one-dimensional Lie algebras. One the other hand, when the Coriolis force is introduced, the dynamical admits seven Lie point symmetries and twenty one-dimensional Lie algebras.

For all the independent Lie algebras we determined all the invariant functions which correspond to all the independent similarity solutions.

In a future work we plan to classify all the independent one-dimensional Lie algebras which lead to analytic forms for the similarity solutions.

### **Acknowledgements**

The author wants to thank Professor Kevin Duffy for gracious hospitality and Professor PGL Leach for a fruitful discussion on the subject.

## A Tables

In this appendix we present the tables 3, 4, 5, 6, 7 and 8 which are refereed in the main article.

Table 3: Adjoint representation of the admitted Lie point symmetries for the nonrotating 2D Shallow water

$Ad(e^{\varepsilon \mathbf{X}_i}) \mathbf{X}_j$	$\mathbf{X}_1$	$\mathbf{X}_2$	$\mathbf{X}_3$	$\mathbf{X}_4$	$\mathbf{X}_5$	$\mathbf{X}_6$	$\mathbf{X}_7$	$\mathbf{X}_8$
$\mathbf{X}_1$	$X_1$	$X_2$	$X_3$	$X_4 - \varepsilon X_2$	$X_5 - \varepsilon X_3$	$X_6$	$X_7 + \varepsilon(\gamma - 1)X_1$	$X_8$
$\mathbf{X}_2$	$X_1$	$X_2$	$X_3$	$X_4$	$X_5$	$X_6 + \varepsilon X_3$	$X_7$	$X_8 - \varepsilon(\gamma - 1)X_2$
$\mathbf{X}_3$	$X_1$	$X_2$	$X_3$	$X_4$	$X_5$	$X_6 - \varepsilon X_2$	$X_7$	$X_8 - \varepsilon(\gamma - 1)X_3$
$\mathbf{X}_4$	$X_1 + \varepsilon X_2$	$X_2$	$X_3$	$X_4$	$X_5$	$X_6 + \varepsilon X_5$	$X_7 - \varepsilon(\gamma - 1)X_4$	$X_8 - \varepsilon(\gamma - 1)X_4$
$\mathbf{X}_5$	$X_1 + \varepsilon X_3$	$X_2$	$X_3$	$X_4$	$X_5$	$X_6 - \varepsilon X_4$	$X_7 - \varepsilon(\gamma - 1)X_5$	$X_8 - \varepsilon(\gamma - 1)X_5$
$\mathbf{X}_6$	$X_1$	$X_2 \cos \varepsilon - X_3 \sin \varepsilon$	$X_2 \sin \varepsilon + X_3 \cos \varepsilon$	$X_4 \cos \varepsilon - X_5 \sin \varepsilon$	$X_4 \sin \varepsilon + X_5 \cos \varepsilon$	$X_6$	$X_7$	$X_8$
$\mathbf{X}_7$	$e^{-(\gamma-1)\varepsilon} X_1$	$X_2$	$X_3$	$e^{-(\gamma-1)\varepsilon} X_4$	$e^{-(\gamma-1)\varepsilon} X_5$	$X_6$	$X_7$	$X_8$
$\mathbf{X}_8$	$X_1$	$e^{(\gamma-1)\varepsilon} X_3$	$e^{(\gamma-1)\varepsilon} X_4$	$e^{-(\gamma-1)\varepsilon} X_4$	$e^{-(\gamma-1)\varepsilon} X_5$	$X_6$	$X_7$	$X_8$

Table 4: Lie invariants for the optimal system of the nonrotating system

Symmetry	Invariants
$\mathbf{X}_1$	$x, y, h(x, y), u(x, y), v(x, y)$
$\mathbf{X}_2$	$t, y, h(t, y), u(t, y), v(t, y)$
$\mathbf{X}_3$	$t, x, h(t, x), u(t, x), v(t, x)$
$\mathbf{X}_4$	$t, y, h(t, y), \frac{x}{t} + U(t, y), v(t, y)$
$\mathbf{X}_5$	$t, x, h(t, x), u(t, x), \frac{y}{t} + V(t, x)$
$\mathbf{X}_6$	$t, x^2 + y^2, h(t, x^2 + y^2), \frac{xU(t, x^2 + y^2) + yV(t, x^2 + y^2)}{\sqrt{x^2 + y^2}}, \frac{yU(t, x^2 + y^2) - xV(t, x^2 + y^2)}{\sqrt{x^2 + y^2}}$
$\mathbf{X}_7$	$\frac{x}{t}, \frac{y}{t}, h(\frac{x}{t}, \frac{y}{t}), u(\frac{x}{t}, \frac{y}{t}), v(\frac{x}{t}, \frac{y}{t})$
$\mathbf{X}_8$	$H(x, y)t^{\frac{2}{1-\gamma}}, U(x, y)t^{-1}, V(x, y)t^{-1}$
$\xi_{(12)}$	$x - \alpha t, y, h(x - \alpha t, y), u(x - \alpha t, y), v(x - \alpha t, y)$
$\xi_{(13)}$	$x, y - \alpha t, h(x, y - \alpha t), u(x, y - \alpha t), v(x, y - \alpha t)$
$\xi_{(14)}$	$x - \frac{\alpha}{2}t^2, y, h(x - \frac{\alpha}{2}t^2, y), u(x - \frac{\alpha}{2}t^2, y), v(x - \frac{\alpha}{2}t^2, y)$
$\xi_{(15)}$	$x, y - \frac{\alpha}{2}t^2, h(x, y - \frac{\alpha}{2}t^2), u(x, y - \frac{\alpha}{2}t^2), v(x, y - \frac{\alpha}{2}t^2)$

Table 5: Lie invariants for the optimal system of the nonrotating system

Symmetry	Invariants
$\xi_{(16)}$	$t, e^{-\alpha t} (x^2 + y^2), u(t, e^{-\alpha t} x^2 + y^2) \cos(\alpha t) + v(t, e^{-\alpha t} x^2 + y^2) \sin(\alpha t)$ $\frac{y}{x}, h(e^{-\alpha t} x^2 + y^2, \frac{y}{x}), u(t, e^{-\alpha t} x^2 + y^2) \sin(\alpha t) - v(t, e^{-\alpha t} x^2 + y^2) \cos(\alpha t)$
$\xi_{(23)}$	$t, x - \alpha y, h(t, x - \alpha y), u(t, x - \alpha y), v(t, x - \alpha y)$
$\xi_{(34)}$	$t, y - \frac{x}{\alpha t}, h(t, y - \frac{x}{\alpha t}), u(t, y - \frac{x}{\alpha t}), v(t, y - \frac{x}{\alpha t})$
$\xi_{(25)}$	$t, y - \alpha t x, h(t, y - \alpha t x), u(t, y - \alpha t x), v(t, y - \alpha t x)$
$\xi_{(45)}$	$t, y - \alpha x, h(t, y - \alpha x), \alpha \frac{x}{t} + U(t, y - \alpha x), \alpha \frac{x}{t} + V(t, y - \alpha x)$
$\xi_{(123)}$	$t - \alpha x, t - \beta y, h(t - \alpha x, t - \beta y), u(t - \alpha x, t - \beta y), v(t - \alpha x, t - \beta y)$
$\xi_{(145)}$	$x - \frac{\alpha}{2} t^2, y - \frac{\beta}{2} t^2, h(x - \frac{\alpha}{2} t^2, y - \frac{\beta}{2} t^2), \alpha t + U(x - \frac{\alpha}{2} t^2, y - \frac{\beta}{2} t^2), \beta t + V(x - \frac{\alpha}{2} t^2, y - \frac{\beta}{2} t^2)$
$\xi_{(125)}$	$x - \alpha t, y - \frac{\beta}{2} t^2, h(x - \alpha t, y - \frac{\beta}{2} t^2), u(x - \alpha t, y - \frac{\beta}{2} t^2), \beta t + V(x - \alpha t, y - \frac{\beta}{2} t^2)$
$\xi_{(134)}$	$x - \frac{\beta}{2} t^2, y - \alpha t, h(x - \frac{\beta}{2} t^2, y - \alpha t), \beta t + U(x - \frac{\beta}{2} t^2, y - \alpha t), V(x - \frac{\beta}{2} t^2, y - \alpha t)$
$\xi_{(67)}$	$\frac{\ln t}{\alpha}, w = \frac{t^{-\frac{\alpha + \sqrt{\alpha(\alpha-4)-4}}{2\alpha}}}{2\sqrt{\alpha(\alpha-4)-4}} \left( x - \left( \alpha + \sqrt{\alpha(\alpha-4)-4} \right) y \right), z = \frac{t^{-\frac{\alpha + \sqrt{\alpha(\alpha-4)-4}}{2\alpha}}}{2\sqrt{\alpha(\alpha-4)-4}} \left( x + \left( \alpha + \sqrt{\alpha(\alpha-4)-4} \right) y \right)$ $h(w, z), U(w, z) \sin\left(\frac{\ln t}{\alpha}\right) + V(w, z) \sin\left(\frac{\ln t}{\alpha}\right), U(w, z) \cos\left(\frac{\ln t}{\alpha}\right) - V(w, z) \sin\left(\frac{\ln t}{\alpha}\right)$
$\xi_{(68)}$	$t, x^2 + y^2, x^{-\frac{2}{\gamma-1}} h(t, x^2 + y^2), \frac{U(t, x^2 + y^2) \cos\left(\frac{\ln x}{\alpha}\right) + V(t, x^2 + y^2) \sin\left(\frac{\ln x}{\alpha}\right)}{x}, \frac{U(t, x^2 + y^2) \sin\left(\frac{\ln x}{\alpha}\right) - V(t, x^2 + y^2) \cos\left(\frac{\ln x}{\alpha}\right)}{x}$
$\xi_{(78)}$	$w = x t^{-\frac{(\gamma-1)}{\alpha(\gamma-1)-2}}, z = y t^{-\frac{(\gamma-1)}{\alpha(\gamma-1)-2}}, t^{-\frac{2\alpha}{\alpha(\gamma-1)-2}} h(w, z), t^{-\frac{(\gamma-1)\alpha}{\alpha(\gamma-1)-2}} u(w, z), t^{-\frac{(\gamma-1)\alpha}{\alpha(\gamma-1)-2}} v(w, z)$
$\xi_{(678)}$	$t, t^{-1-\beta} (x^2 + y^2), t^{-\beta} (U(t, x^2 + y^2) \sin(\alpha t) + V(t, x^2 + y^2) \cos(\alpha t))$ $t^{-\frac{2\beta}{\gamma-1}} H(t, x^2 + y^2), t^{-\beta} (U(t, x^2 + y^2) \cos(\alpha t) - V(t, x^2 + y^2) \sin(\alpha t))$

Table 6: Adjoint representation of the admitted Lie point symmetries for the rotating 2D Shallow water

$Ad(e^{(\varepsilon \mathbf{Y}_i)}) \mathbf{Y}_j$	$\mathbf{Y}_1$	$\mathbf{Y}_2$	$\mathbf{Y}_3$	$\mathbf{Y}_4$	$\mathbf{Y}_5$	$\mathbf{Y}_6$	$\mathbf{Y}_7$
$\mathbf{Y}_1$	$Y_1$	$Y_2$	$Y_3$	$Y_4$	$Y_5 \cos(f\varepsilon) - Y_6 \sin(f\varepsilon)$	$Y_5 \sin(f\varepsilon) + Y_6 \cos(f\varepsilon)$	$Y_7$
$\mathbf{Y}_2$	$Y_1$	$Y_2$	$Y_3$	$Y_4 + \varepsilon Y_3$	$Y_5$	$Y_6$	$Y_7 - \varepsilon(\gamma - 1)Y_2$
$\mathbf{Y}_3$	$Y_1$	$Y_2$	$Y_3$	$Y_4 - \varepsilon Y_2$	$Y_5$	$Y_6$	$Y_7 - \varepsilon(\gamma - 1)Y_3$
$\mathbf{Y}_4$	$Y_1$	$Y_2 \cos \varepsilon - Y_3 \sin \varepsilon$	$Y_2 \sin \varepsilon + Y_3 \cos \varepsilon$	$Y_4$	$Y_5 \cos \varepsilon + Y_6 \sin \varepsilon$	$Y_6 \cos \varepsilon - Y_5 \sin \varepsilon$	$Y_7$
$\mathbf{Y}_5$	$Y_1 + f\varepsilon Y_6$	$Y_2$	$Y_3$	$Y_4 - \varepsilon Y_6$	$Y_5$	$Y_6$	$Y_7 - \varepsilon(\gamma - 1)Y_5$
$\mathbf{Y}_6$	$Y_1 - f\varepsilon Y_5$	$Y_2$	$Y_3$	$Y_4 + \varepsilon Y_5$	$Y_5$	$Y_6$	$Y_7 - \varepsilon(\gamma - 1)Y_6$
$\mathbf{Y}_7$	$Y_1$	$e^{(\gamma-1)\varepsilon} Y_2$	$e^{(\gamma-1)\varepsilon} Y_3$	$Y_4$	$e^{(\gamma-1)\varepsilon} Y_5$	$e^{(\gamma-1)\varepsilon} Y_6$	$Y_7$

Table 7: Lie invariants for the optimal system of the rotating system

Symmetry	Invariants
$\mathbf{Y}_1$	$x, y, h(x, y), u(x, y), v(x, y)$
$\mathbf{Y}_2$	$t, y, h(t, y), u(t, y), v(t, y)$
$\mathbf{Y}_3$	$t, x, h(t, x), u(t, x), v(t, x)$
$\mathbf{Y}_4$	$t, x^2 + y^2, h(t, x^2 + y^2), \frac{xU(t, x^2 + y^2) + yV(t, x^2 + y^2)}{\sqrt{x^2 + y^2}}, \frac{yU(t, x^2 + y^2) - xV(t, x^2 + y^2)}{\sqrt{x^2 + y^2}}$
$\mathbf{Y}_5$	$t, x \cot(ft) - y, h(t, x \cot(ft) - y), fx \cot(ft) + U(t, x \cot(ft) - y), -fx + V(t, x \cot(ft) - y)$
$\mathbf{Y}_6$	$t, x \tan(ft) + y, h(t, x \tan(ft) + y), -fx \tan(ft) + U(t, x \tan(ft) + y), -fx + V(t, x \tan(ft) + y)$
$\mathbf{Y}_7$	$\frac{x}{t}, \frac{y}{t}, h\left(\frac{x}{t}, \frac{y}{t}\right), u\left(\frac{x}{t}, \frac{y}{t}\right), v\left(\frac{x}{t}, \frac{y}{t}\right)$
$\chi(12)$	$x - \alpha t, y, h(x - \alpha t, y), u(x - \alpha t, y), v(x - \alpha t, y)$
$\chi(13)$	$x, y - \alpha t, h(x, y - \alpha t), u(x, y - \alpha t), v(x, y - \alpha t)$
$\chi(14)$	$t, e^{-\alpha t}(x^2 + y^2), u(t, e^{-\alpha t}x^2 + y^2) \cos(\alpha t) + v(t, e^{-\alpha t}x^2 + y^2) \sin(\alpha t)$ $\frac{y}{x}, h(e^{-\alpha t}x^2 + y^2, \frac{y}{x}), u(t, e^{-\alpha t}x^2 + y^2) \sin(\alpha t) - v(t, e^{-\alpha t}x^2 + y^2) \cos(\alpha t)$

Table 8: Lie invariants for the optimal system of the rotating system

Symmetry	Invariants
$\chi_{(15)}$	$x + \frac{\alpha}{f} \cos(ft), y - \frac{\alpha}{f} \sin(ft), h\left(x + \frac{\alpha}{f} \cos(ft), y - \frac{\alpha}{f} \sin(ft)\right),$ $\alpha \sin(ft) + U\left(x + \frac{\alpha}{f} \cos(ft), y - \frac{\alpha}{f} \sin(ft)\right), \alpha \cos(ft) + V\left(x + \frac{\alpha}{f} \cos(ft), y - \frac{\alpha}{f} \sin(ft)\right)$
$\chi_{(16)}$	$x - \frac{\alpha}{f} \sin(ft), y - \frac{\alpha}{f} \cos(ft), h\left(x - \frac{\alpha}{f} \sin(ft), y - \frac{\alpha}{f} \cos(ft)\right),$ $\alpha \cos(ft) + U\left(x - \frac{\alpha}{f} \sin(ft), y - \frac{\alpha}{f} \cos(ft)\right), -\alpha \sin(ft) + V\left(x - \frac{\alpha}{f} \sin(ft), y - \frac{\alpha}{f} \cos(ft)\right)$
$\chi_{(17)}$	$xe^{-\alpha t}, ye^{-\alpha t}, e^{\frac{2\alpha}{\gamma-1}t} h(xe^{-\alpha t}, ye^{-\alpha t}), e^{\alpha t} u(xe^{-\alpha t}, ye^{-\alpha t}), e^{\alpha t} v(xe^{-\alpha t}, ye^{-\alpha t})$
$\chi_{(23)}$	$t, x - \alpha y, h(t, x - \alpha y), u(t, x - \alpha y), v(t, x - \alpha y)$
$\chi_{(45)}$	$t, w = (x^2 + y^2 - 2x \cos(ft) + 2y \sin(ft)), \frac{U(t,w)+f \sin(ft)}{V(t,w)+f \cos(ft)}, \frac{U(t,w)^2+V(t,w)^2}{2} + f(U(t,w) \sin(ft) + V(t,w) \cos(ft))$
$\chi_{(46)}$	$t, w = (x^2 + y^2 - 2x \sin(ft) - 2y \cos(ft)), \frac{U(t,w)-f \cos(ft)}{V(t,w)+f \sin(ft)}, \frac{U(t,w)^2+V(t,w)^2}{2} + f(V(t,w) \sin(ft) - U(t,w) \cos(ft))$
$\chi_{(56)}$	$t, z = y - \frac{x(\cos(ft)-\alpha \sin(ft))}{\sin(ft)+\alpha \cos(ft)}, h(t, z), f \frac{x(\cos(ft)-\alpha \sin(ft))}{\sin(ft)+\alpha \cos(ft)} + U(t, z), -x + V(t, z)$
$\chi_{(47)}$	$t, x^2 + y^2, x^{-\frac{2}{\gamma-1}} h(t, x^2 + y^2), \frac{U(t,x^2+y^2) \cos(\frac{\ln x}{\alpha}) + V(t,x^2+y^2) \sin(\frac{\ln x}{\alpha})}{x}, \frac{U(t,x^2+y^2) \sin(\frac{\ln x}{\alpha}) - V(t,x^2+y^2) \cos(\frac{\ln x}{\alpha})}{x}$
$\chi_{(123)}$	$t - \alpha x, t - \beta y, h(t - \alpha x, t - \beta y), u(t - \alpha x, t - \beta y), v(t - \alpha x, t - \beta y)$
$\chi_{(147)}$	$z = e^{-t(\gamma-1)} (x \cos t - y \sin t), w = e^{-t(\gamma-1)} (y \cos t + x \sin t),$ $e^{-t(\gamma-1)} h(z, w), e^{-t(\gamma-1)} (U(z, w) \cos t - V(z, w) \sin t), e^{-t(\gamma-1)} (U(z, w) \sin t + V(z, w) \cos t)$

## References

- [1] P.J. Olver, Applications of Lie Groups to Differential Equations, Springer-Verlag, New York, (1993)
- [2] G.W. Bluman and S. Kumei, Symmetries and Differential Equations, Springer-Verlag, New York, (1989)
- [3] N.H. Ibragimov, CRC Handbook of Lie Group Analysis of Differential Equations, Volume I: Symmetries, Exact Solutions, and Conservation Laws, CRS Press LLC, Florida (2000)
- [4] L. V. Ovsiannikov, Group analysis of differential equations, Academic Press, New York, (1982)
- [5] S Jamal, P.G.L. Leach and A. Paliathanasis, Quaestiones Mathematicae 42, 125 (2019)
- [6] G.M. Webb, J. Phys A: Math. Gen. 23, 3885 (1990)
- [7] M. Tsamparlis, A. Paliathanasis and L. Karpathopoulos, J. Phys. A: Math. Theor. 45, 275201 (2012)
- [8] H. Azad and M.T. Mustafa, Commun. Nonlinear. Sci. Numer. Simul. 15, 1132 (2010)
- [9] M. Tsamparlis, A. Paliathanasis and A. Qadir, Int. J. Geom. Methods Mod. Phys. 12, 155003 (2005)
- [10] S.V. Meleshko and V.P. Shapeev, J. Nonl. Math. Phys. 18, 195 (2011)
- [11] A. Halder, A. Paliathanasis and P.G.L. Leach, Symmetry 10, 744 (2018)
- [12] K. S. Chou and C. Z. Qu, Acta Applicandae Mathematicae, 83, 257 (2004).
- [13] F. Galas and E. W. Richter, Phys. D, (50) 297 (1991).
- [14] S. V. Coggeshalla and J.Meyer-ter-Vehn, J.Math. Phys. 33, 3585 (1992).
- [15] X. Hu, Y. Li and Y. Chen, J. Math. Phys. 56, 053504 (2015)
- [16] G. K. Vallis, Atmospheric and Oceanic Fluid Dynamics: Fundamentals and Large-Scale Circulation, Cambridge University Press, Cambridge (2006)
- [17] R. Courant and K. O. Friedrichs, Supersonic Flow and Shock Waves, Interscience Publ. (1948)
- [18] J. Kevorkian, Partial Differential Equations: Analytical Solutions Techniques, Chapman and Hall, New York (1990)
- [19] V. Caleffi, A. Valiani and A. Zanni, J. Hydraulic Research 41, 167 (2003)
- [20] R.A.D. Akkermans, L.P.J. Kamp, H.J.H. Clercx and G.J.F. van Heijst, Phys. Rev. E 82, 026314 (2010)
- [21] D.H. Kim, Y.S. Cho and Y.K. Yi, Ocean Engineering 34, 1164 (2007)
- [22] A.A. Chesnokov, Eur. J. Appl. Math. 20, 461 (2009)
- [23] X. Xin, L. Zhang, Y. Xia and H. Liu, Appl. Math. Lett. 94, 112 (2019)
- [24] S. Szatmari and A. Bihlo, Comm. Nonl. Sci. Num. Sim. 19, 530 (2014)
- [25] A.A. Chesnokov, J. Appl. Mech. Techn. Phys. 49, 737 (2008)

- [26] J.-G. Liu, Z.-F. Zeng, Y. He and G.-P. Ai, *Int. J. Nonl. Sci. Num. Sim.* 16, 114 (2013)
- [27] M. Pandey, *Int. J. Nonl. Sci. Num. Sim.* 16, 93 (2015)
- [28] A. Paliathanasis, to appear in *Zeitschrift für Naturforschung A* DOI:10.1515/zna-2019-0063  
[arXiv:1906.00689]