

Counting curves in a linear system with upto eight singular points

Somnath Basu and Ritwik Mukherjee

Abstract

In this paper, we develop a systematic approach to enumerate curves with a certain number of nodes and one further singularity which maybe more degenerate. As a result, we obtain an explicit formula for the number of curves in a sufficiently ample linear system, passing through the right number of generic points, that have δ nodes and one singularity of codimension k , for all $\delta + k \leq 8$. In particular, we recover the formulas for curves with upto six nodal points obtained by Vainsencher. Moreover, all the codimension seven numbers we have obtained agree with the formulas obtained by Kazarian. Finally, in codimension eight, we recover the formula of A.Weber, M.Mikosz and P.Pragacz for curves with one singular point and we also recover the formula of Kleiman and Piene for eight nodal curves. All the other codimension eight numbers we have obtained are new.

Contents

1	Introduction	1
2	A survey of related results	3
3	A brief overview of the method we use	5
4	Necessary and sufficient criteria for a singularity	7
5	Notation	9
6	Recursive Formulas	11
7	Proof of the Recursive Formulas via Euler class computation	15
8	Low degree checks in \mathbb{P}^2	42
9	Low degree checks in $\mathbb{P}^1 \times \mathbb{P}^1$	46

1 Introduction

Enumerative geometry is a branch of mathematics concerned with the following question:

How many geometric objects are there which satisfy prescribed constraints?

It is a classical subject that dates back to over 150 years ago. Hilbert's fifteenth problem was to lay a rigorous foundation for enumerative Schubert calculus. While the problems in this field are typically easy to state, solutions to almost all of them require various deep concepts from various branches of

mathematics.

In order to motivate the results of this paper, let us first state a very general problem in enumerative geometry:

Question 1.1. *Let $L \rightarrow X$ be a holomorphic line bundle over a compact complex surface and $\mathcal{D} := \mathbb{P}H^0(X, L) \approx \mathbb{P}^{\delta_L}$ be the space of non zero holomorphic sections upto scaling. What is $N(A_1^\delta \mathfrak{X})$, the number of curves in X , that belong to the linear system $H^0(X, L)$, passing through $\delta_L - (k + \delta)$ generic points and having δ distinct nodes and one singularity of type \mathfrak{X} whose codimension is k .*

The above question has been studied by several mathematicians over the last thirty years; it becomes increasingly hard when the codimension $\delta + k$ increases and/or when the singularity \mathfrak{X} becomes more degenerate. In this paper, we give develop a systematic method to approach this question and obtain a complete answer to the above question till codimension eight. Before stating the main result of this paper, let us make a couple of definitions:

Definition 1.2. Let $L \rightarrow X$ be a holomorphic line bundle over a complex surface and and $f \in H^0(X, L)$ a holomorphic section. A point $q \in f^{-1}(0)$ is of singularity type A_k, D_k, E_6, E_7, E_8 or X_9 if there exists a coordinate system $(x, y) : (\mathcal{U}, q) \rightarrow (\mathbb{C}^2, 0)$ such that $f^{-1}(0) \cap \mathcal{U}$ is given by

$$\begin{aligned} A_k : y^2 + x^{k+1} = 0 \quad k \geq 0, \quad D_k : y^2x + x^{k-1} = 0 \quad k \geq 4, \\ E_6 : y^3 + x^4 = 0, \quad E_7 : y^3 + yx^3 = 0, \quad E_8 : y^3 + x^5 = 0, \\ X_9 : x^4 + y^4 = 0. \end{aligned}$$

In more common terminology, q is a *smooth* point of $f^{-1}(0)$ if it is a singularity of type A_0 ; a *simple node* (or just node) if its singularity type is A_1 ; a *cuspid* if its type is A_2 ; a *tacnode* if its type is A_3 ; a *an ordinary triple point* if its type is D_4 ; and an *ordinary quadruple point* if its type is X_9 .

Definition 1.3. A holomorphic line bundle $L \rightarrow X$ over a compact complex manifold X is sufficiently k -ample if $L \approx L_1^{\otimes n} \otimes \xi \rightarrow X$ for some $n \geq k$, where $L_1 \rightarrow X$ is a very ample line bundle and $\xi \rightarrow X$ is a line bundle such that the linear system $H^0(X, \xi)$ is base point free.

Remark 1.4. We will frequently use the phrase “a singularity of codimension k ”. Roughly speaking, this refers to the number of conditions having that singularity imposes on the space of curves. More precisely, it is the expected codimension of the equisingular strata. Hence, a singularity of type A_k, D_k or $E_{k \leq 8}$ is a singularity of codimension k . However, a singularity of type X_9 is of codimension 8, not 9. The reason that it is standard to denote an ordinary quadruple point by X_9 and not X_8 is because singularities are traditionally indexed by their Milnor number and not by the codimension of the equisingular strata. For A_k, D_k and $E_{k \leq 8}$ singularities, these two numbers happen to be the same. In general, these two numbers need not be the same.

The main result of this paper is as follows:

Main Result 1.5. *Let $L \rightarrow X$ be a line bundle and \mathfrak{X} be one of the singularities defined in 1.2 whose codimension is k . We obtain an explicit formula for $N(A_1^\delta \mathfrak{X})$ if $\delta + k \leq 8$, provided the line bundle is sufficiently $(2\delta + C_{\mathfrak{X}})$ -ample, where*

$$C_{A_k} = k, \quad C_{D_4} = 3, \quad C_{D_k} = k - 2 \quad \text{if } k > 4, \quad C_{E_k} = k - 3 \quad \text{and} \quad C_{X_9} = 4.$$

Remark 1.6. The ampleness requirement is there to prove that the sections we encounter are transverse to the zero set. However, the bound we impose is not the optimal bound. In particular, our bound is coarser than the bound given by Kleiman and Piene in [17]. One of the reasons our bound is coarse, is because while proving transversality, we restrict ourselves exclusively to the tangent space of the curves; we do not utilize the fact that we can also move the marked point. This makes the proof of transversality much easier, but it requires a higher bound on the ampleness of the line bundle. We believe that with some further effort we should be able to improve our bound (namely by moving both the curve and the point); we intend to investigate that question in the future.

Remark 1.7. In order to keep this paper of a reasonable length, we have decided to split the proof of our main result into two papers. In this paper, we focus exclusively on the Euler class computation. We precisely state the relevant transversality, closure and multiplicity claims and then go on to compute the enumerative numbers. Our second paper [1] is devoted exclusively to proving the technical results on closure and multiplicities that we use in this paper. We also prove in [1] the necessary transversality results that we use in this paper. The transversality results actually follow without too much effort. In [6] and [3] we prove the relevant transversality results for \mathbb{P}^2 when there are one or two singular points. It is easy to see from the proof presented in [3], how the proof of transversality generalizes to more than one singular point. In [27] we show how to generalize the proof of transversality to the case of general linear system that is sufficiently ample. In particular, the proof of the relevant transversality results that we use in this paper is very similar to the proof presented in our earlier three papers ([6], [3] and [27]).

2 A survey of related results

2.1 Counting singular curves in a linear system

We now give a brief survey of related results in this area of mathematics. As early as 1873, Zeuthen computed the number of nodal cubics and the number of tri nodal quartics in \mathbb{P}^2 , passing through the appropriate number of generic points (cf [46]). A lot of progress has since been made in enumerating plane nodal curves; the question is now very well understood. Ziv Ran and then Caporaso and Harris completely solved the problem of enumerating δ -nodal curves in \mathbb{P}^2 (cf [34], [33] and [11]). Caporaso and Harris gave a recursive formula for δ -nodal curves, in terms of the number of curves with lesser number of nodes, but which have contacts of arbitrary order with a given generic line. Later on their result for \mathbb{P}^2 was extended by Ravi Vakil to certain del-Pezzo surfaces (cf. [44]). An alternative proof of the Caporasso-Harris formula was also given by Ionel-Parker (cf. [14]) and Tehrani-Zinger (cf. [40]), using the Symplectic Sum formula.¹

Let us now look at the more general question of enumerating curves in an algebraic surface (such as $\mathbb{P}^1 \times \mathbb{P}^1$). In [43], Vainsencher obtained an explicit formula for curves in a sufficiently ample linear system that have upto six nodes.

Next, Kazarian computed *all possible* codimension seven numbers in [15]. More precisely, he obtained a formula for the number of curves in a sufficiently ample linear system (passing through the right number of generic points) and having a singularity of type $\chi_{k_1}, \chi_{k_2}, \dots, \chi_{k_n}$, provided $\kappa := k_1 + k_2 + \dots + k_n \leq 7$. Here χ_{k_i} is a singularity of codimension k_i .

We now mention the only two results that we are aware of in codimension eight. In [31], [32] and [25], A.Weber, M.Mikosz and P.Pragacz extended the method of Kazarian to enumerate curves

¹It would be interesting to see if the result of Ravi Vakil can also be obtained using the Symplectic Sum Formula.

with one singular point. In his slides [45, page 52], Weber has explicitly written down the formula for $N(A_8)$ and $N(E_8)$; our formulas agree with theirs.

Finally, we look at the results of S.Kleiman and R.Piene. In [17, Theorem 1.2], Kleiman and Piene obtained an explicit formula for *eight* nodal curves in a suitably ample linear system! They also obtained a formula for $N(A_1^\delta D_4)$, $N(A_1^\delta D_6)$ and $N(A_1^\delta E_7)$, with δ at most 3, 1 and 0 respectively (basically till codimension seven).² They also give a very precise condition for the ampleness requirement of their line bundle. They consider a line bundle $L \rightarrow X$, that is of the form $L := M^{\otimes m} \otimes N$, where m is at least three times the expected codimension of the equisingular strata and where N is spanned by global holomorphic sections. Using this ampleness condition, they prove that the numbers they compute are indeed enumerative, i.e. each curve appears with a multiplicity of one in this linear system.

Finally we note that a great deal of progress has been made in proving that universal formulas exist in terms of Chern classes for the number of curves in a sufficiently ample linear system passing through the right number of generic points and having singularities of type $\chi_{k_1}, \chi_{k_2}, \dots, \chi_{k_n}$. The fact that a universal formula exists was conjectured by Göttsche when the singularities $\chi_{k_1}, \dots, \chi_{k_n}$ are all simple nodes. Two independent proofs of this conjecture have now been given using methods of BPS calculus and using degeneration methods by Kool, Shende and Thomas ([21]) and Tzeng ([41]) respectively. There is also an earlier approach by Liu in [22] and [23]. Recently, Li and Tzeng gave a proof for the existence of universal formulas for any collection of singularities $\chi_{k_1}, \chi_{k_2}, \dots, \chi_{k_n}$ in [42], thereby generalizing the Göttsche conjecture. A further generalization of the Göttsche conjecture was proved by Rennemo, where he shows that there is a universal polynomial to count hypersurfaces in a sufficiently ample linear system, with any collection of isolated singularities ([35, Proposition 7.8]).

It should be noted that even for δ -nodal curves in \mathbb{P}^2 , it is nontrivial question as to when the polynomials obtained by the Göttsche conjecture are actually enumerative. They are enumerative if d is large. However, Göttsche conjectured that these polynomials are actually enumerative for all d roughly greater than $\frac{d}{2}$. This has now been proven by Kleiman and Shende in [18].

For a more detailed overview of this subject, we direct the reader to the comprehensive survey article [16] by Kleiman.

2.2 Counting curves of a fixed genus: Gromov-Witten theory

Let us now consider another classical enumerative geometry problem; namely counting curves of a fixed genus. To begin with, let us look at genus zero curves (i.e. rational curves). Let X be a compact complex surface and $\beta \in H_2(X, \mathbb{Z})$ a given homology class. A classical question is to ask what is n_β , the number of genus zero degree β curves in X passing through $\langle c_1(TX), \beta \rangle - 1$ generic points? For the case of \mathbb{P}^2 , the numbers n_1, n_2, n_3 and n_4 were computed by Zeuthen in the late nineteenth century ([46]). There was essentially no progress in this question, until in the early 1990's a formula for n_d was announced by Kontsevich-Manin and Ruan-Tian. More precisely, in [20] and [39], the authors obtain a formula for n_β for any complex del-Pezzo surface.³ Both the approaches involve Symplectic Geometry; it involves using a gluing theorem which can be algebraically reformulated as the associativity of Quantum Cohomology. A lot of progress has since been made in this question

²The paper by Kleiman and Piene is earlier than the paper by Kazarian; so their formulas for the codimension seven numbers were all new.

³Strictly speaking, Kontsevich-Manin compute the genus zero Gromov-Witten invariants of del-Pezzo surfaces. The fact that the Gromov-Witten invariants are actually enumerative for all del-Pezzo surfaces was later proved by Pandharipande and Göttsche in [30].

over the last thirty years using Symplectic and Algebraic geometry and more recently using Tropical geometry (cf [24]).

The question of enumerating genus zero curves with higher singularities is a much harder question. Pandahripande and later Kock computed the number of rational cuspidal curves in \mathbb{P}^2 and $\mathbb{P}^1 \times \mathbb{P}^1$ using algebro geometric methods (cf [29] and [19]). Later on, Zinger used methods from Symplectic Geometry to approach the question of enumerating rational curves with cusps and higher singularities (cf [49], [52], [50] and [51]). It should also be noted that although Zinger obtains his formulas for curves in \mathbb{P}^2 , his methods are applicable in a much more general setup. In fact, his methods apply to surfaces where the Gromov-Witten Theory and its relationship with Enumerative geometry is well understood. In particular, his methods apply very nicely to del-Pezzo surfaces. A few of Zinger's results have been extended by the second author to del-Pezzo surfaces (cf. [7], [9] and [8]).

Let us now explain how this question is relevant to the problem we are studying in this paper. The problem of counting curves of a fixed genus enables us to subject our formulas to several low degree checks. For example, let us look at n_d , the number of degree d rational curves in \mathbb{P}^2 passing through $3d - 1$ generic points. A little bit of thought shows that n_3 is same as the number of nodal cubics through 8 points; hence, this number helps us verify the formula for $N(A_1)$. Next, we consider n_4 . A little bit of thought shows that this is same as the number of irreducible quartics with three (unordered) nodes through 11 points. Hence, the number n_4 helps us verify the the formula for $N(A_1^3)$. And so on.

Let us now consider higher singularities. For example, in [49], Zinger has computed the number of rational degree d curves with one E_6 -singularity. For $d = 3$, this ought to be zero (which it is) and for $d = 4$, his answer ought to be the same as the number of quartics through 8 points having an E_6 -singularity (again, that is indeed the case). Hence, these results for higher singularities also help us verify many of our formulas.

In section 8, we subject our formulas to several low degree checks. In section 9, we correct some of the low degree checks made by Vainsencher in his paper [43] and show how the numbers are consistent with the values predicted by the formula of Kontsevich and Manin in [20].

3 A brief overview of the method we use

We now give a brief description of the method we use to compute the enumerative numbers. Our starting point is the following fact from differential topology (cf. [10, Proposition 12.8]):

Theorem 3.1. *Let $V \rightarrow X$ be an oriented vector bundle over a compact oriented manifold X and suppose $s : X \rightarrow V$ is a smooth section that is transverse to the zero set. Then the Poincaré dual of $[s^{-1}(0)]$ is the Euler class of V . In particular, if the rank of V is same as the dimension of X then the signed cardinality of $s^{-1}(0)$ is the Euler class of V , evaluated on the fundamental class of X , i.e.,*

$$|\pm s^{-1}(0)| = \langle e(V), [X] \rangle.$$

Our goal is to enumerate curves with certain singularities. The fact that a curve has a specific singularity means that certain derivatives vanish (an example of this fact is the Implicit Function Theorem and the Morse Lemma). We interpret these derivatives as the section of some bundle. Hence, our enumerative numbers are the zeros of a section of some bundle restricted to the *open* part of a manifold (or variety). One of the reasons we typically have to restrict ourselves to the open part of a manifold is because we are usually enumerating curves with more than one singular point;

hence we have to consider the space of curves with a collection of marked points, where the marked points are all distinct.

Next, we observe that if the line bundle is sufficiently ample, then this section (induced by taking derivatives) will be transverse to the zero set restricted to the open part.⁴ We then evaluate the Euler class of this bundle on the fundamental class of the manifold/variety⁵ and hope that this number is what we want. As one might expect, that hope is almost always too naive. This is because, the section will typically vanish on the boundary and hence give an excess (degenerate) contribution to the Euler class. The real challenge is therefore to compute this degenerate contribution to the Euler class.

The most famous and well known method of computing degenerate contributions to the Euler class is global excess intersection theory, which is developed in Fulton's book [12]. This approach is global and rigid in nature and involves blowing up the degenerate loci to smooth them out. This method has been successfully applied to solve a large class of enumerative problems; in particular it has been applied by Vainsencher and Kleiman and Piene to obtain their formula for δ -nodal curves.

In his PhD thesis [48], Aleksey Zinger developed a completely different approach to address the problem of computing degenerate contributions to the Euler class; he developed the method of *local intersection theory*. This approach is soft and local in nature and does not involve taking any blowups. The method involves perturbing the relevant section *smoothly* and counting (with a sign) the number of zeros of the section near the degenerate locus. He applied this method successfully to solve a very large class of enumerative problems; these include counting rational curves in \mathbb{P}^2 (and even \mathbb{P}^n) with various types of singularities (which include cusp, tacnode, triple point and E_6 -singularities). He also used local intersection theory to compute the degenerate contribution to the Symplectic Invariant of \mathbb{P}^n and computed the genus g Enumerative Invariant of \mathbb{P}^n for $g = 2$ and 3 , thereby extending the result of Pandharipande and Ionel for genus one (cf. [28] and [13]).

Zinger further applied the machinery of local intersection theory in his paper [47] to enumerate plane degree d curves with singularities till codimension three. In particular, he computed all the possible codimension three numbers $N(A_1^\delta \mathfrak{X})$ in his paper by using local intersection theory.

The second author's PhD thesis (cf [26]) comprises of extending the approach developed in [47] and applying *local* intersection theory to compute $N(A_1^\delta \mathfrak{X})$ for higher codimension. In his thesis, the second author obtained a formula for $N(A_1^\delta \mathfrak{X})$ for \mathbb{P}^2 when $\delta + k \leq 7$; part of that thesis has been published in [6] and [5] (namely the cases $\delta = 0$ and 1 have been published). In this paper we finally obtain a formula for all the numbers $N(A_1^\delta \mathfrak{X})$ for any sufficiently ample linear system till codimension 8, thereby going beyond the results of the second author's thesis. Furthermore, we give *alternative* proofs for the results of Vainsencher, Kleiman-Piene and Kazarian⁶ and we also obtain new results in the end (basically all the codimension eight numbers are new except for the formula for eight nodal curves and the formulas for one singular point).

Finally, let us give a brief description of the method used by Kazarian. His method works on the principle that there exists a universal formula (in terms of Chern classes) for the Thom polynomial associated to a given singularity. He then goes on to consider enough special cases to find out what that exact combination is. As an example, suppose there is a polynomial of degree m . To find out what the polynomial is, we simply have to find the value of the polynomial at enough points

⁴The open part of the variety we consider is always going to be smooth in our case.

⁵Any algebraic variety defines a homology class since the singularities have at least real codimension two. This follows from the main result of [53], namely that any singular space whose singularities are of real codimension two or more (i.e. a pseudocycle) defines a homology class.

⁶However, we should mention that we only compute numbers of the form $N(A_1^\delta \mathfrak{X})$; Kazarian computes all numbers till codimension seven; such as $N(A_3 A_4)$ for example.

([15, page 667]). One of the challenges of this method is to prove the existence of such a universal formula. This fact is shown by R.Rimanyi and A. Szucs in [36], [37] and [38]. This method has been successfully applied to compute all the codimension seven numbers. It has also been extended by A.Weber, M.Mikosz and P.Pragacz to enumerate curves with one singular point in codimension eight (cf. [31], [32] and [25]).

4 Necessary and sufficient criteria for a singularity

For the convenience of the reader, let us recapitulate from the arXiv version of our paper [2, section 5], a necessary and sufficient criterion for a curve $f^{-1}(0)$ to have a singularity of type \mathfrak{X} at the point q .⁷ Let $f = f(x, y)$ be a holomorphic function defined on a neighborhood of the origin in \mathbb{C}^2 and i, j be non-negative integers. We define

$$f_{ij} := \left. \frac{\partial^{i+j} f}{\partial^i x \partial^j y} \right|_{(x,y)=(0,0)}.$$

Let us now define the following directional derivatives, which are functions of f_{ij} :

$$\begin{aligned} A_3^f &:= f_{30}, & A_4^f &:= f_{40} - \frac{3f_{21}^2}{f_{02}}, & A_5^f &:= f_{50} - \frac{10f_{21}f_{31}}{f_{02}} + \frac{15f_{12}f_{21}^2}{f_{02}^2}, \\ A_6^f &:= f_{60} - \frac{15f_{21}f_{41}}{f_{02}} - \frac{10f_{31}^2}{f_{02}} + \frac{60f_{12}f_{21}f_{31}}{f_{02}^2} + \frac{45f_{21}^2f_{22}}{f_{02}^2} - \frac{15f_{03}f_{21}^3}{f_{02}^3} - \frac{90f_{12}^2f_{21}^2}{f_{02}^3}, \\ A_7^f &:= f_{70} - \frac{21f_{21}f_{51}}{f_{02}} - \frac{35f_{31}f_{41}}{f_{02}} + \frac{105f_{12}f_{21}f_{41}}{f_{02}^2} + \frac{105f_{21}^2f_{32}}{f_{02}^2} + \frac{70f_{12}f_{31}^2}{f_{02}^2} + \frac{210f_{21}f_{22}f_{31}}{f_{02}^2} \\ &\quad - \frac{105f_{03}f_{21}^2f_{31}}{f_{02}^3} - \frac{420f_{12}^2f_{21}f_{31}}{f_{02}^3} - \frac{630f_{12}f_{21}^2f_{22}}{f_{02}^3} - \frac{105f_{13}f_{21}^3}{f_{02}^3} + \frac{315f_{03}f_{12}f_{21}^3}{f_{02}^4} + \frac{630f_{12}^3f_{21}^2}{f_{02}^4}, \\ A_8^f &:= f_{80} - \frac{28f_{21}f_{61}}{f_{02}} - \frac{56f_{31}f_{51}}{f_{02}} + \frac{210f_{21}^2f_{42}}{f_{02}^2} + \frac{420f_{21}f_{22}f_{41}}{f_{02}^2} - \frac{210f_{03}f_{21}^2f_{41}}{f_{02}^3} + \frac{560f_{21}f_{31}f_{32}}{f_{02}^3} \\ &\quad - \frac{840f_{13}f_{21}^2f_{31}}{f_{02}^3} - \frac{420f_{21}^3f_{23}}{f_{02}^3} + \frac{1260f_{03}f_{21}^3f_{22}}{f_{02}^4} - \frac{35f_{41}^2}{f_{02}} + \frac{280f_{22}f_{31}^2}{f_{02}^2} - \frac{280f_{03}f_{21}f_{31}^2}{f_{02}^3} - \frac{1260f_{21}^2f_{22}^2}{f_{02}^3} + \\ &\quad - \frac{105f_{04}f_{21}^4}{f_{02}^4} - \frac{315f_{03}^2f_{21}^4}{f_{02}^5} + \frac{168f_{21}f_{51}f_{12}}{f_{02}^2} + \frac{280f_{31}f_{41}f_{12}}{f_{02}^2} - \frac{1680f_{21}^2f_{32}f_{12}}{f_{02}^3} - \frac{3360f_{21}f_{22}f_{31}f_{12}}{f_{02}^3} + \\ &\quad - \frac{2520f_{03}f_{21}^2f_{31}f_{12}}{f_{02}^4} + \frac{2520f_{13}f_{21}^3f_{12}}{f_{02}^4} - \frac{840f_{21}f_{41}f_{12}^2}{f_{02}^3} + \frac{7560f_{21}^2f_{22}f_{12}^2}{f_{02}^4} - \frac{560f_{31}^2f_{12}^2}{f_{02}^3} - \frac{5040f_{03}f_{21}^3f_{12}^2}{f_{02}^5} \\ &\quad + \frac{3360f_{21}f_{31}f_{12}^3}{f_{02}^4} - \frac{5040f_{21}^2f_{12}^4}{f_{02}^5} \end{aligned} \tag{1}$$

and

$$D_6^f := f_{40}, \quad D_7^f := f_{50} - \frac{5f_{31}^2}{3f_{12}}, \quad D_8^f := f_{60} + \frac{5f_{03}f_{31}f_{50}}{3f_{12}^2} - \frac{5f_{31}f_{41}}{f_{12}} - \frac{10f_{03}f_{31}^3}{3f_{12}^3} + \frac{5f_{22}f_{31}^2}{f_{12}^2}. \tag{2}$$

We will now state a necessary and sufficient criteria for a curve to have a specific singularity.

⁷The arXiv version contains a few details that were omitted from our published paper [6]; hence we will often refer to the arXiv version of our paper.

Lemma 4.1. *Let $f = f(x, y)$ be a holomorphic function defined on a neighborhood of the origin in \mathbb{C}^2 such that $f_{00} = 0$ and $\nabla f|_{(0,0)} \neq 0$. Then the curve has an A_0 singularity at the origin (i.e. a smooth point).*

Lemma 4.2. *Let $f = f(x, y)$ be a holomorphic function defined on a neighbourhood of the origin in \mathbb{C} such that $f_{00}, \nabla f|_{(0,0)} = 0$ and $\nabla^2 f|_{(0,0)}$ is non-degenerate. Then the curve has an A_1 singularity at the origin.*

Remark 4.3. Lemma 4.1 is also known as the **Implicit Function Theorem** and Lemma 4.2 is also known as the **Morse Lemma**.

We now state the remaining Lemmas, which can be thought of as a continuation of Lemma 4.2.

Lemma 4.4. *Let $f = f(r, s)$ be a holomorphic function defined on a neighbourhood of the origin in \mathbb{C} such that $f(0, 0), \nabla f|_{(0,0)} = 0$ and there exists a non-zero vector $w = (w_1, w_2)$ such that at the origin $\nabla^2 f(w, \cdot) = 0$, i.e., the Hessian is degenerate. Let $x = w_1 r + w_2 s, y = -\bar{w}_2 r + \bar{w}_1 s$ and f_{ij} be the partial derivatives with respect to the new variables x and y . Then, the curve $f^{-1}(0)$ has an A_k singularity at the origin (for $2 \leq k \leq 7$) if $f_{02} \neq 0$ and the directional derivatives A_i^f defined in (1) are zero for all $i \leq k$ and $A_{k+1}^f \neq 0$.*

Lemma 4.5. *Let $f = f(x, y)$ be a holomorphic function defined on a neighbourhood of the origin in \mathbb{C} such that $f_{00}, \nabla f|_{(0,0)}, \nabla^2 f|_{(0,0)} = 0$ and there does not exist a non-zero vector $w = (w_1, w_2)$ such that at the origin $\nabla^3 f(w, w, \cdot) = 0$. Then the curve $f^{-1}(0)$ has a D_4 singularity at the origin.*

Lemma 4.6. *Let $f = f(r, s)$ be a holomorphic function defined on a neighbourhood of the origin in \mathbb{C} such that $f_{00}, \nabla f|_{(0,0)}, \nabla^2 f|_{(0,0)} = 0$ and there exists a non-zero vector $w = (w_1, w_2)$ such that at the origin $\nabla^3 f(w, w, \cdot) = 0$. Let $x = w_1 r + w_2 s, y = -\bar{w}_2 r + \bar{w}_1 s$ and f_{ij} be the partial derivatives with respect to the new variables x and y . Then, the curve $f^{-1}(0)$ has a D_k -node at the origin (for $5 \leq k \leq 7$) if $f_{12} \neq 0$ and the directional derivatives D_i^f defined in (2) are zero for all $i \leq k$ and $D_{k+1}^f \neq 0$.*

Lemma 4.7. *Let $f = f(r, s)$ be a holomorphic function defined on a neighbourhood of the origin in \mathbb{C} such that $f_{00} = \nabla f|_{(0,0)} = \nabla^2 f|_{(0,0)} = 0$ and there exists a non-zero vector $w = (w_1, w_2)$ such that at the origin $\nabla^3 f(w, w, \cdot) = 0$. Let $x = w_1 r + w_2 s, y = -\bar{w}_2 r + \bar{w}_1 s$ and f_{ij} be partial derivatives with respect to the new coordinates, x and y . Then, the curve $f^{-1}(0)$ has an E_6 singularity at the origin if $f_{12} = 0$ and $f_{03} \neq 0, f_{40} \neq 0$.*

Lemma 4.8. *Let $f = f(r, s)$ be a holomorphic function defined on a neighbourhood of the origin in \mathbb{C} such that $f_{00}, \nabla f|_{(0,0)}, \nabla^2 f|_{(0,0)} = 0$ and there exists a non-zero vector $w = (w_1, w_2)$ such that at the origin $\nabla^3 f(w, w, \cdot) = 0$. Let $x = w_1 r + w_2 s, y = -\bar{w}_2 r + \bar{w}_1 s$. Let f_{ij} be the partial derivatives with respect to the new variables x and y . Then, the curve $f^{-1}(0)$ has an E_7 singularity at the origin if $f_{12} = 0$ and $f_{03} \neq 0, f_{31} \neq 0$.*

Lemma 4.9. *Let $f = f(r, s)$ be a holomorphic function defined on a neighbourhood of the origin in \mathbb{C} such that $f_{00}, \nabla f|_{(0,0)}, \nabla^2 f|_{(0,0)} = 0$ and there exists a non-zero vector $w = (w_1, w_2)$ such that at the origin $\nabla^3 f(w, w, \cdot) = 0$. Let $x = w_1 r + w_2 s, y = -\bar{w}_2 r + \bar{w}_1 s$. Let f_{ij} be the partial derivatives with respect to the new variables x and y . Then, the curve $f^{-1}(0)$ has an E_8 singularity at the origin if $f_{12} = 0, f_{31} = 0$ and $f_{03} \neq 0, f_{50} \neq 0$.*

Proofs of Lemmas 4.4 to 4.8: This is proved in section 3 of our paper [6]. The proof of Lemma 4.9 is very similar to the proof of the other Lemmas; it involves looking at the Taylor expansion and making an appropriate change of coordinates. \square

5 Notation

We will now introduce some basic notation that will be used for stating our recursive formulas. Our setup is as follows; we consider a sufficiently ample line bundle $L \rightarrow X$ over a compact complex surface X . Let

$$\mathcal{D} \approx \mathbb{P}H^0(X, L) \approx \mathbb{P}^{\delta_L},$$

denote the space of holomorphic sections of the line bundle $L \rightarrow X$, upto scaling. Let \mathfrak{X} be a singularity of a given type. We will also denote \mathfrak{X} to be the space of curves and a marked point such that the curve has a singularity of type \mathfrak{X} at the marked point. More precisely,

$$\mathfrak{X} := \{([f], q) \in \mathcal{D} \times X : f \text{ has a singularity of type } \mathfrak{X} \text{ at } q\}.$$

For example,

$$A_2 := \{([f], q) \in \mathcal{D} \times X : f \text{ has a singularity of type } A_2 \text{ at } q\}.$$

Next, given n subsets S_1, S_2, \dots, S_n of $\mathcal{D} \times \mathbb{P}^2$, we define

$$S_1 \circ S_2 \circ \dots \circ S_n := \{([f], q_1, \dots, q_n) \in \mathcal{D} \times (\mathbb{P}^2)^n : ([f], q_i) \in S_i, \dots, ([f], q_n) \in S_n, \\ \text{and } q_1, \dots, q_n \text{ are all distinct}\}.$$

For example, $A_1^2 \circ A_2$ is the space of curves with three ordered points, where the curve has a simple node at the first two points and a cusp at the last point and all the three points are distinct. Similarly, $A_1^2 \circ \overline{A}_2$ is the space of curves with three distinct ordered points, where the curve has a simple node at the first two points and a singularity at least as degenerate as a cusp at the last point; the curve could have a tacnode at the last marked point.

It will also be useful to consider the space of curves with a singularity and a specific direction along which certain directional derivatives vanish. More precisely, let $\mathbb{P}TX \rightarrow X$ denote the projectivization of the tangent bundle X and let \mathfrak{X} be a singularity of type $A_{k \geq 2}$ or $D_{k \geq 5}$ or E_6, E_7 or E_8 . Then we define

$$\begin{aligned} \mathcal{P}A_k &:= \{([f], l_q) \in \mathcal{D} \times \mathbb{P}TX : ([f], q) \in A_k, \quad \nabla^2 f(v, \cdot) = 0 \quad \forall v \in l_q\} & \text{if } k \geq 2, \\ \mathcal{P}D_k &:= \{([f], l_q) \in \mathcal{D} \times \mathbb{P}TX : ([f], q) \in D_k, \quad \nabla^3 f(v, v, \cdot) = 0 \quad \forall v \in l_q\} & \text{if } k \geq 5, \\ \mathcal{P}E_k &:= \{([f], l_q) \in \mathcal{D} \times \mathbb{P}TX : ([f], q) \in E_k, \quad \nabla^3 f(v, v, \cdot) = 0 \quad \forall v \in l_q\} & \text{if } k = 6, 7, 8. \end{aligned}$$

For example, $\mathcal{P}A_2$ is the space of curves with a marked point and a marked direction, such that the curve has a cusp at the marked point and the marked direction belongs to the kernel of the Hessian. Note that in all the above cases, the projection map $\pi : \mathcal{P}\mathfrak{X} \rightarrow \mathfrak{X}$ is one to one. Next, let us define

$$\begin{aligned} \mathcal{P}A_1 &:= \{([f], l_q) \in \mathcal{D} \times \mathbb{P}TX : ([f], q) \in A_1, \quad \nabla^2 f(v, v) = 0 \quad \forall v \in l_q\}, \\ \mathcal{P}D_4 &:= \{([f], l_q) \in \mathcal{D} \times \mathbb{P}TX : ([f], q) \in D_4, \quad \nabla^3 f(v, v, v) = 0 \quad \forall v \in l_q\}, \\ \mathcal{P}X_9 &:= \{([f], l_q) \in \mathcal{D} \times \mathbb{P}TX : ([f], q) \in X_9, \quad \nabla^4 f(v, v, v, v) = 0 \quad \forall v \in l_q\}. \end{aligned}$$

For example, $\mathcal{P}A_1$ is the space of curves with a marked point and a marked direction, such that the curve has a node at the marked point and the second derivative along the marked direction vanishes. Note that there are two such distinguished directions. Hence, the projection map $\pi : \mathcal{P}A_1 \rightarrow A_1$ is two to one. Similarly, the maps

$$\pi : \mathcal{P}D_4 \rightarrow D_4 \quad \text{and} \quad \pi : \mathcal{P}X_9 \rightarrow X_9$$

are three to one and four to one respectively. Let us also define

$$\hat{\mathfrak{X}} := \{([f], l_q) \in \mathcal{D} \times \mathbb{P}TX : ([f], q) \in \mathfrak{X}\} = \pi^{-1}(\mathfrak{X}).$$

Finally, if S_1, \dots, S_n are subsets of $\mathcal{D} \times X$ and T is a subset of $\mathcal{D} \times \mathbb{P}TX$, then we define

$$S_1 \circ S_2 \circ \dots \circ S_n \circ T := \{([f], q_1, \dots, q_n, l_{q_{n+1}}) \in \mathcal{D} \times X^n \times \mathbb{P}TX : ([f], q_1) \in S_1, \dots, ([f], q_n) \in S_n, \\ ([f], l_{q_{n+1}}) \in T \quad \text{and} \\ q_1, \dots, q_n, q_{n+1} \quad \text{are all distinct}\}.$$

For example, $A_1^2 \circ \mathcal{P}A_2$ is the space of curves with three distinct ordered points, where the curve has a simple node at the first two points and a cusp at the last point and a distinguished direction at the last marked point, such that the Hessian vanishes along that direction. We are now in a position to define a few numbers. Let us define

$$N(A_1^\delta \mathfrak{X}, n_1, m_1, m_2) := \langle y^{\delta L - k - n_1 - m_1 - 2m_2 - \delta} c_1^{m_1} x_1^{m_1} x_2^{m_2}, \overline{[A_1^\delta \circ \mathfrak{X}]} \rangle, \quad (3)$$

$$N(A_1^\delta \mathcal{P}\mathfrak{X}, n_1, m_1, m_2, \theta) := \langle y^{\delta L - k - n_1 - m_1 - 2m_2 - \delta - \theta} c_1^{m_1} x_1^{m_1} x_2^{m_2} \lambda^\theta, \overline{[A_1^\delta \circ \mathcal{P}\mathfrak{X}]} \rangle \quad \text{and} \quad (4)$$

$$N(A_1^\delta \hat{\mathfrak{X}}, n_1, m_1, m_2, \theta) := \langle y^{\delta L - k - n_1 - m_1 - 2m_2 - \delta - \theta} c_1^{m_1} x_1^{m_1} x_2^{m_2} \lambda^\theta, \overline{[A_1^\delta \circ \hat{\mathfrak{X}}]} \rangle, \quad (5)$$

where

$$c_1 := c_1(L), \quad x_i := c_i(T^*X), \quad \lambda := c_1(\hat{\gamma}^*), \quad y := c_1(\gamma_D^*)$$

and $\gamma_D \rightarrow D$ and $\hat{\gamma} \rightarrow \mathbb{P}TX$ are the tautological line bundles.

Let us now make a few observations. First, let us abbreviate $N(A_1^\delta \mathfrak{X}, 0, 0, 0)$ as $N(A_1^\delta \mathfrak{X})$; this quantity is the number of curves passing through the right number of points that have δ distinct nodes and one singularity of type \mathfrak{X} . Next, we note that

$$N(A_1^\delta \mathfrak{X}, n_1, m_1, m_2) = \frac{1}{\deg(\pi)} N(A_1^\delta \mathcal{P}\mathfrak{X}, n_1, m_1, m_2, 0), \quad (6)$$

where $\deg(\pi)$ is the degree of the projection map $\pi : \mathcal{P}\mathfrak{X} \rightarrow \mathfrak{X}$. We remind the reader that the degree is always one, except in the cases when $\mathfrak{X} = A_1, D_4$ or X_9 , in which case the degree is 2, 3 and 4 respectively.

Next, we note that using the ring structure of $H^*(\mathbb{P}TX, \mathbb{Z})$ (cf [10], pp. 270), we conclude that

$$N(A_1^\delta \mathcal{P}\mathfrak{X}, n_1, m_1, m_2, \theta) = N(A_1^\delta \mathcal{P}\mathfrak{X}, n_1, m_1 + 1, m_2, \theta - 1) - N(A_1^\delta \mathcal{P}\mathfrak{X}, n_1, m_1, m_2 + 1, \theta - 2) \\ \text{if } \theta > 1. \quad (7)$$

Finally, we note that

$$N(A_1^\delta \hat{\mathfrak{X}}, n_1, m_1, m_2, \theta) = 0 \quad \text{if } \theta = 0, \\ N(A_1^\delta \hat{\mathfrak{X}}, n_1, m_1, m_2, \theta) = N(A_1^\delta \mathfrak{X}, n_1, m_1, m_2) \quad \text{if } \theta = 1 \quad \text{and} \\ N(A_1^\delta \hat{\mathfrak{X}}, n_1, m_1, m_2, \theta) = N(A_1^\delta \hat{\mathfrak{X}}, n_1, m_1 + 1, m_2, \theta - 1) - N(A_1^\delta \hat{\mathfrak{X}}, n_1, m_1, m_2 + 1, \theta - 2) \\ \text{if } \theta > 1. \quad (8)$$

This again follows from [10], pp. 270.

6 Recursive Formulas

We are now ready to state the recursive formulas. We are going to give a recursive formula for curves with δ -nodes and one singularity of type \mathfrak{X} in terms of number of curves with $\delta - 1$ nodes and one singularity more degenerate than \mathfrak{X} . Combined with the base case of the recursion (Theorem 6.1), equations (6), (7) and (8), Theorems 6.3 to 6.22 enable us to obtain an explicit formula for $N(A_1^\delta \mathfrak{X})$ if $\delta + k \leq 8$.

We have written a mathematica program to implement these recursive formulas and obtain the final formulas. The program is available on the second author's homepage

<https://www.sites.google.com/site/ritwik371/home>

For the convenience of the reader, we have generated an output of all the numbers $N(A_1^\delta \mathfrak{X})$ and have appended it at the end of this paper (after the Bibliography).

We remind the reader that in the formulas for $N(A_1^\delta \mathfrak{X})$, the nodes are always *ordered*. To obtain the corresponding number of curves where the nodes are unordered, we have to divide by $(\delta + 1)!$ or $\delta!$, depending on whether the last singularity (\mathfrak{X}) is A_1 or something different, respectively.

Theorem 6.1. *The number $N(A_1, n_1, m_1, m_2)$ is given by*

$$N(A_1, n_1, m_1, m_2) = \begin{cases} 3c_1^2 + 2c_1x_1 + x_2 & \text{if } (n_1, m_1, m_2) = (0, 0, 0), \\ 3c_1^2 + c_1x_1 & \text{if } (n_1, m_1, m_2) = (1, 0, 0), \\ c_1^2 & \text{if } (n_1, m_1, m_2) = (2, 0, 0), \\ 3c_1x_1 + x_1^2 & \text{if } (n_1, m_1, m_2) = (0, 1, 0), \\ c_1x_1 & \text{if } (n_1, m_1, m_2) = (1, 1, 0), \\ x_1^2 & \text{if } (n_1, m_1, m_2) = (0, 2, 0), \\ x_2 & \text{if } (n_1, m_1, m_2) = (0, 0, 1), \\ 0 & \text{otherwise.} \end{cases} \quad (9)$$

Remark 6.2. Theorem 6.1 will serve as the base case of our recursion.

Theorem 6.3. *If $1 \leq \delta \leq 7$, then*

$$\begin{aligned} N(A_1^\delta A_1, n_1, m_1, m_2) &= N(A_1^\delta, 0, 0, 0) \times N(A_1, n_1, m_1, m_2) \\ &\quad - \binom{\delta}{1} \left(N(A_1^{\delta-1}, n_1, m_1, m_2) + N(A_1^{\delta-1}, n_1 + 1, m_1, m_2) + 3N(A_1^{\delta-1} A_2, n_1, m_1, m_2) \right) \\ &\quad + 4 \binom{\delta}{2} N(A_1^{\delta-2} A_3, n_1, m_1, m_2) + 18 \binom{\delta}{3} N(A_1^{\delta-3} D_4, n_1, m_1, m_2), \end{aligned}$$

provided the line bundle is sufficiently $(2\delta + 1)$ -ample.

Theorem 6.4. *If $0 \leq \delta \leq 7$, then $N(A_1^\delta \mathcal{P}A_1, n_1, m_1, m_2, \theta)$ is as follows:*

$$\begin{aligned} \theta = 0: & \quad 2N(A_1^\delta A_1, n_1, m_1, m_2) \\ \theta = 1: & \quad 2N(A_1^\delta A_1, n_1, m_1 + 1, m_2) + N(A_1^\delta A_1, n_1, m_1, m_2) + N(A_1^\delta A_1, n_1 + 1, m_1, m_2) \\ & \quad - 2 \binom{\delta}{2} N(A_1^{\delta-2} \mathcal{P}D_4, n_1, m_1, m_2, \theta - 1), \end{aligned}$$

provided the line bundle is sufficiently $(2\delta + 1)$ -ample.

Theorem 6.5. *If $0 \leq \delta \leq 6$, then*

$$\begin{aligned}
N(A_1^\delta \mathcal{P}A_2, n_1, m_1, m_2, \theta) &= N(A_1^\delta \mathcal{P}A_1, n_1, m_1, m_2, \theta) + N(A_1^\delta \mathcal{P}A_1, n_1, m_1 + 1, m_2, \theta) \\
&\quad + N(A_1^\delta \mathcal{P}A_1, n_1 + 1, m_1, m_2, \theta) - 2 \binom{\delta}{1} N(A_1^{\delta-1} \mathcal{P}A_3, n_1, m_1, m_2, \theta) \\
&\quad - 3 \binom{\delta}{1} N(A_1^{\delta-1} \hat{D}_4, n_1, m_1, m_2, \theta) - 4 \binom{\delta}{2} N(A_1^{\delta-2} \mathcal{P}D_4, n_1, m_1, m_2, \theta) \\
&\quad - 4 \binom{\delta}{2} N(A_1^{\delta-2} \hat{D}_5, n_1, m_1, m_2, \theta),
\end{aligned}$$

provided the line bundle is sufficiently $(2\delta + 2)$ -ample.

Theorem 6.6. *If $0 \leq \delta \leq 5$, then*

$$\begin{aligned}
N(A_1^\delta \mathcal{P}A_3, n_1, m_1, m_2, \theta) &= 3N(A_1^\delta \mathcal{P}A_2, n_1, m_1, m_2, \theta + 1) + N(A_1^\delta \mathcal{P}A_2, n_1 + 1, m_1, m_2, \theta) \\
&\quad + N(A_1^\delta \mathcal{P}A_2, n_1, m_1, m_2, \theta) - 2 \binom{\delta}{1} N(A_1^{\delta-1} \mathcal{P}A_4, n_1, m_1, m_2, \theta) \\
&\quad - 2 \binom{\delta}{2} N(A_1^{\delta-2} \mathcal{P}D_5, n_1, m_1, m_2, \theta),
\end{aligned}$$

provided the line bundle is sufficiently $(2\delta + 3)$ -ample.

Theorem 6.7. *If $0 \leq \delta \leq 4$, then*

$$\begin{aligned}
N(A_1^\delta \mathcal{P}A_4, n_1, m_1, m_2, \theta) &= 2N(A_1^\delta \mathcal{P}A_3, n_1, m_1, m_2, \theta + 1) + 2N(A_1^\delta \mathcal{P}A_3, n_1, m_1 + 1, m_2, \theta) \\
&\quad + 2N(A_1^\delta \mathcal{P}A_3, n_1, m_1, m_2, \theta) + 2N(A_1^\delta \mathcal{P}A_3, n_1 + 1, m_1, m_2, \theta) \\
&\quad - 2 \binom{\delta}{1} N(A_1^{\delta-1} \mathcal{P}A_5, n_1, m_1, m_2, \theta) - 4 \binom{\delta}{2} N(A_1^{\delta-2} \mathcal{P}D_6, n_1, m_1, m_2, \theta),
\end{aligned}$$

provided the line bundle is sufficiently $(2\delta + 4)$ -ample.

Theorem 6.8. *If $0 \leq \delta \leq 4$, then $N(A_1^\delta \mathcal{P}D_4, n_1, m_1, m_2, \theta)$ is as follows:*

$$\begin{aligned}
\theta = 0: \quad & 2N(A_1^\delta \mathcal{P}A_3, n_1, m_1 + 1, m_2, \theta) - 2N(A_1^\delta \mathcal{P}A_3, n_1, m_1, m_2, \theta + 1) \\
& + N(A_1^\delta \mathcal{P}A_3, n_1, m_1, m_2, \theta) + N(A_1^\delta \mathcal{P}A_3, n_1 + 1, m_1, m_2, \theta) \\
& - 2 \binom{\delta}{1} N(A_1^{\delta-1} D_5, n_1, m_1, m_2) - 2 \binom{\delta}{2} N(A_1^{\delta-2} \mathcal{P}D_6, n_1, m_1, m_2, 0) \\
\theta = 1: \quad & \frac{1}{3}(N(A_1^\delta \mathcal{P}D_4, n_1, m_1, m_2, \theta - 1) + 3N(A_1^\delta \mathcal{P}D_4, n_1, m_1 + 1, m_2, \theta - 1) \\
& + N(A_1^\delta \mathcal{P}D_4, n_1 + 1, m_1, m_2, \theta - 1)) - 24 \binom{\delta}{3} N(X_9, n_1, m_1, m_2) \\
\theta \geq 2: \quad & N(A_1^\delta \mathcal{P}D_4, n_1, m_1 + 1, m_2, \theta - 1) - N(A_1^\delta \mathcal{P}D_4, n_1, m_1, m_2 + 1, \theta - 2),
\end{aligned}$$

provided the line bundle is sufficiently $(2\delta + 4)$ -ample.

Remark 6.9. Note that the structure of this formula is a little bit different; we have something different when $\theta = 0$ and when $\theta = 1$. The reason for this will be clear when we prove the Theorem (subsection 7.7).

Theorem 6.10. *If $0 \leq \delta \leq 3$, then $N(A_1^\delta \mathcal{P}D_5, n_1, m_1, m_2, \theta)$ is as follows:*

$$\begin{aligned} N(A_1^\delta \mathcal{P}D_5, n_1, m_1, m_2, \theta) = & N(A_1^\delta \mathcal{P}D_4, n_1, m_1, m_2, \theta + 1) + N(A_1^\delta \mathcal{P}D_4, n_1, m_1 + 1, m_2, \theta) \\ & + N(A_1^\delta \mathcal{P}D_4, n_1, m_1, m_2, \theta) + N(A_1^\delta \mathcal{P}D_4, n_1 + 1, m_1, m_2, \theta) \\ & - 2 \binom{\delta}{1} N(A_1^{\delta-1} \mathcal{P}D_6, n_1, m_1, m_2, \theta) - 12 \binom{\delta}{1} N(A_1^{\delta-1} \hat{X}_9, n_1, m_1, m_2, \theta) \\ & - 18 \binom{\delta}{3} N(\mathcal{P}X_9, n_1, m_1, m_2, \theta), \end{aligned}$$

provided the line bundle is sufficiently $(2\delta + 5)$ -ample.

Theorem 6.11. *If $0 \leq \delta \leq 3$, then*

$$\begin{aligned} N(A_1^\delta \mathcal{P}A_5, n_1, m_1, m_2, \theta) = & 3N(A_1^\delta \mathcal{P}A_4, n_1, m_1, m_2, \theta + 1) + 2N(A_1^\delta \mathcal{P}A_4, n_1, m_1 + 1, m_2, \theta) \\ & + 2N(A_1^\delta \mathcal{P}A_4, n_1, m_1, m_2, \theta) + 2N(A_1^\delta \mathcal{P}A_4, n_1 + 1, m_1, m_2, \theta) \\ & - 2 \binom{\delta}{1} N(A_1^{\delta-1} \mathcal{P}A_6, n_1, m_1, m_2, \theta) - \binom{\delta}{1} N(A_1^{\delta-1} \mathcal{P}E_6, n_1, m_1, m_2, \theta) \\ & - 4 \binom{\delta}{2} N(A_1^{\delta-2} \mathcal{P}D_7, n_1, m_1, m_2, \theta), \end{aligned}$$

provided the line bundle is sufficiently $(2\delta + 5)$ -ample.

Theorem 6.12. *If $0 \leq \delta \leq 2$, then*

$$\begin{aligned} N(A_1^\delta \mathcal{P}D_6, n_1, m_1, m_2, \theta) = & 4N(A_1^\delta \mathcal{P}D_5, n_1, m_1, m_2, \theta + 1) + N(A_1^\delta \mathcal{P}D_5, n_1, m_1, m_2, \theta) \\ & + N(A_1^\delta \mathcal{P}D_5, n_1 + 1, m_1, m_2, \theta) - 2 \binom{\delta}{1} N(A_1^{\delta-1} \mathcal{P}D_7, n_1, m_1, m_2, \theta) \\ & - \binom{\delta}{1} N(A_1^{\delta-1} \mathcal{P}E_7, n_1, m_1, m_2, \theta), \end{aligned}$$

provided the line bundle is sufficiently $(2\delta + 6)$ -ample.

Theorem 6.13. *If $0 \leq \delta \leq 2$, then $N(A_1^\delta \mathcal{P}E_6, n_1, m_1, m_2, \theta)$ is as follows:*

$$\begin{aligned} N(A_1^\delta \mathcal{P}E_6, n_1, m_1, m_2, \theta) = & 2N(A_1^\delta \mathcal{P}D_5, n_1, m_1 + 1, m_2, \theta) - N(A_1^\delta \mathcal{P}D_5, n_1, m_1, m_2, \theta + 1) \\ & + N(A_1^\delta \mathcal{P}D_5, n_1, m_1, m_2, \theta) + N(A_1^\delta \mathcal{P}D_5, n_1 + 1, m_1, m_2, \theta) \\ & - \binom{\delta}{1} N(A_1^{\delta-1} \mathcal{P}E_7, n_1, m_1, m_2, \theta) - 4 \binom{\delta}{1} N(\hat{X}_9, n_1, m_1, m_2, \theta), \end{aligned}$$

provided the line bundle is sufficiently $(2\delta + 6)$ -ample.

Theorem 6.14. *If $0 \leq \delta \leq 2$, then $N(A_1^\delta \mathcal{P}A_6, n_1, m_1, m_2, \theta)$ is as follows:*

$$\begin{aligned} N(A_1^\delta \mathcal{P}A_6, n_1, m_1, m_2, \theta) = & 2N(A_1^\delta \mathcal{P}A_5, n_1, m_1, m_2, \theta + 1) + 4N(A_1^\delta \mathcal{P}A_5, n_1, m_1 + 1, m_2, \theta) \\ & + 3N(A_1^\delta \mathcal{P}A_5, n_1, m_1, m_2, \theta) + 3N(A_1^\delta \mathcal{P}A_5, n_1 + 1, m_1, m_2, \theta) \\ & - 2N(A_1^\delta \mathcal{P}D_6, n_1, m_1, m_2, \theta) - N(A_1^\delta \mathcal{P}E_6, n_1, m_1, m_2, \theta) \\ & - 2 \binom{\delta}{1} N(A_1^{\delta-1} \mathcal{P}A_7, n_1, m_1, m_2, \theta) - 3 \binom{\delta}{1} N(A_1^{\delta-1} \mathcal{P}E_7, n_1, m_1, m_2, \theta) \\ & - 12 \binom{\delta}{1} N(A_1^{\delta-1} \hat{X}_9, n_1, m_1, m_2, \theta) - 6 \binom{\delta}{2} N(\mathcal{P}D_8, n_1, m_1, m_2), \end{aligned}$$

provided the line bundle is sufficiently $(2\delta + 6)$ -ample.

Theorem 6.15. *If $0 \leq \delta \leq 1$, then*

$$\begin{aligned} N(A_1^\delta \mathcal{P}D_7, n_1, m_1, m_2, \theta) &= 4N(A_1^\delta \mathcal{P}D_6, n_1, m_1, m_2, \theta + 1) + 2N(A_1^\delta \mathcal{P}D_6, n_1, m_1 + 1, m_2, \theta) \\ &\quad + 2N(A_1^\delta \mathcal{P}D_6, n_1, m_1, m_2, \theta) + 2N(A_1^\delta \mathcal{P}D_6, n_1 + 1, m_1, m_2, \theta) \\ &\quad - 2 \binom{\delta}{1} N(\mathcal{P}D_8, n_1, m_1, m_2, \theta), \end{aligned}$$

provided the line bundle is sufficiently $(2\delta + 7)$ -ample.

Theorem 6.16. *If $0 \leq \delta \leq 1$, then*

$$\begin{aligned} N(A_1^\delta \mathcal{P}E_7, n_1, m_1, m_2, \theta) &= N(A_1^\delta \mathcal{P}D_6, n_1, m_1, m_2, \theta) - N(A_1^\delta \mathcal{P}D_6, n_1, m_1, m_2, \theta + 1) \\ &\quad 2N(A_1^\delta \mathcal{P}D_6, n_1, m_1 + 1, m_2, \theta) + N(A_1^\delta \mathcal{P}D_6, n_1 + 1, m_1, m_2, \theta) \\ &\quad - 3 \binom{\delta}{1} N(\mathcal{P}X_9, n_1, m_1, m_2, \theta), \end{aligned}$$

provided the line bundle is sufficiently $(2\delta + 3)$ -ample.

Theorem 6.17. *If $0 \leq \delta \leq 1$, then*

$$\begin{aligned} N(A_1^\delta \mathcal{P}A_7, n_1, m_1, m_2, \theta) &= -N(A_1^\delta \mathcal{P}A_6, n_1, m_1, m_2, \theta + 1) + 8N(A_1^\delta \mathcal{P}A_6, n_1, m_1 + 1, m_2, \theta) \\ &\quad + 5N(A_1^\delta \mathcal{P}A_6, n_1, m_1, m_2, \theta) + 5N(A_1^\delta \mathcal{P}A_6, n_1 + 1, m_1, m_2, \theta) \\ &\quad - 5N(A_1^\delta \hat{X}_9, n_1, m_1, m_2, \theta) \\ &\quad - 6N(A_1^\delta \mathcal{P}D_7, n_1, m_1, m_2, \theta) - 7N(A_1^\delta \mathcal{P}E_7, n_1, m_1, m_2, \theta) \\ &\quad - 2 \binom{\delta}{1} N(\mathcal{P}A_8, n_1, m_1, m_2, \theta) - 14 \binom{\delta}{1} N(\mathcal{P}E_8, n_1, m_1, m_2, \theta), \end{aligned}$$

provided the line bundle is sufficiently $(2\delta + 7)$ -ample.

Theorem 6.18. *The number $N(\mathcal{P}E_8, n_1, m_1, m_2, \theta)$ is given by*

$$\begin{aligned} N(\mathcal{P}E_8, n_1, m_1, m_2, \theta) &= N(\mathcal{P}E_7, n_1, m_1, m_2, \theta) + N(\mathcal{P}E_7, n_1, m_1 + 1, m_2, \theta) + 3N(\mathcal{P}E_7, n_1 + 1, m_1, m_2, \theta) \\ &\quad + 2N(\mathcal{P}E_7, n_1, m_1, m_2, \theta + 1) \end{aligned}$$

provided the line bundle is sufficiently 8-ample.

Theorem 6.19. *The number $N(\mathcal{P}D_8, n_1, m_1, m_2, \theta)$ is given by*

$$\begin{aligned} N(\mathcal{P}D_8, n_1, m_1, m_2, \theta) &= 4N(\mathcal{P}D_7, n_1, m_1, m_2, \theta) + 6N(\mathcal{P}D_7, n_1, m_1 + 1, m_2, \theta) + 3N(\mathcal{P}A_7, n_1, m_1, m_2, \theta + 1) \\ &\quad 4N(\mathcal{P}D_7, n_1 + 1, m_1, m_2, \theta) - 3N(\mathcal{P}E_8, n_1, m_1, m_2, \theta) \end{aligned}$$

provided the line bundle is sufficiently 8-ample.

Theorem 6.20. *The number $N(\mathcal{P}X_9, n_1, m_1, m_2, \theta)$ is given by*

$$N(\mathcal{P}X_9, n_1, m_1, m_2, \theta) = N(\mathcal{P}E_7, n_1, m_1, m_2, \theta) + 4N(\mathcal{P}E_7, n_1, m_1, m_2, \theta + 1) + N(\mathcal{P}E_7, n_1 + 1, m_1, m_2, \theta)$$

provided the line bundle is sufficiently 4-ample.

It remains to give a formula for $N(\mathcal{P}A_8, n_1, m_1, m_2, \theta)$. Before we do that, we need to define one more space.

$$\mathcal{P}X_9^\vee := \{([f], l_q) \in \mathcal{D} \times \mathbb{P}TX : ([f], l_q) \in \hat{X}_9, \left(-\frac{f_{31}^3}{8} + \frac{3f_{22}f_{31}f_{40}}{16} - \frac{f_{13}f_{40}^2}{16}\right) = 0\}.$$

Theorem 6.21. *The number $N(\mathcal{P}X_9^\vee, n_1, m_1, m_2, \theta)$ is given by*

$$\begin{aligned} N(\mathcal{P}X_9^\vee, n_1, m_1, m_2, \theta) &= 3N(\hat{X}_9, n_1, m_1, m_2, \theta) + 3N(\hat{X}_9, n_1, m_1 + 1, m_2, \theta) \\ &\quad + 6N(\hat{X}_9, n_1, m_1, m_2, \theta + 1) + 3N(\hat{X}_9, n_1 + 1, m_1, m_2, \theta), \end{aligned}$$

provided the line bundle is sufficiently 4-ample.

Theorem 6.22. *The number $N(\mathcal{P}A_8, n_1, m_1, m_2)$ is given by*

$$\begin{aligned} N(\mathcal{P}A_8, n_1, m_1, m_2, \theta) &= 6N(\mathcal{P}A_7, n_1, m_1, m_2) - 2N(\mathcal{P}A_7, n_1, m_1, m_2, 1) + 10N(\mathcal{P}A_7, n_1, m_1 + 1, m_2) \\ &\quad + 6N(\mathcal{P}A_7, n_1 + 1, m_1, m_2) - 8N(\mathcal{P}D_8, n_1, m_1, m_2) - 16N(\mathcal{P}E_8, n_1, m_1, m_2) \\ &\quad - 6N(\mathcal{P}X_9^\vee, n_1, m_1, m_2, \theta), \end{aligned}$$

provided the line bundle is sufficiently 8-ample.

7 Proof of the Recursive Formulas via Euler class computation

We will now give a proof of the recursive formulas we stated in section 6. The relevant claims on closure, multiplicity and transversality are proved in our earlier papers [6], [5], [27] and [4], when $\delta = 0$ or 1 and $\delta + k \leq 7$. For the remaining cases (namely $\delta > 1$ or $\delta + k > 7$), we will prove the relevant closure, multiplicity and transversality claims in [1].

7.1 Proof of Theorem 6.1: computing $N(A_1, n_1, m_1, m_2)$

This is proved in [4], Proposition 5.2. □

7.2 Proof of Theorem 6.3: computing $N(A_1^\delta A_1, n_1, m_1, m_2)$

Recall that in section 5 we have defined the space

$$\begin{aligned} A_1^\delta \circ (\mathcal{D} \times X) &:= \{([f], q_1, \dots, q_\delta, q_{\delta+1}) \in \mathcal{D} \times (X)^{\delta+1} : f \text{ has a singularity of type } A_1 \text{ at } q_1, \dots, q_\delta, \\ &\quad q_1, \dots, q_{\delta+1} \text{ all distinct}\}. \end{aligned}$$

Let μ be a generic pseudocycle representing the homology class Poincaré dual to

$$c_1^{n_1} x_1^{m_1} x_2^{m_2} y^{\delta_L - (n_1 + m_1 + 2m_2 + \delta + 1)}.$$

We now consider sections of the following two bundles that are induced by the evaluation map and the vertical derivative, namely:

$$\begin{aligned} \psi_{A_0} : A_1^\delta \circ (\mathcal{D} \times X) \cap \mu &\longrightarrow \mathcal{L}_{A_0} := \gamma_{\mathcal{D}}^* \otimes L, & \{\psi_{A_0}([f], q_1, \dots, q_{\delta+1})\}(f) &:= f(q_{\delta+1}) \\ \psi_{A_1} : \psi_{A_0}^{-1}(0) &\longrightarrow \mathcal{V}_{A_1} := \gamma_{\mathcal{D}}^* \otimes T^*X \otimes L, & \{\psi_{A_1}([f], q_1, \dots, q_{\delta+1})\}(f) &:= \nabla f|_{q_{\delta+1}}. \end{aligned}$$

In [1], we show that if L is sufficiently $(2\delta + 1)$ -ample, then these sections are transverse to the zero set. Next, let us define

$$\mathcal{B} := \overline{A_1^\delta \circ (\mathcal{D} \times X)} - A_1^\delta \circ (\mathcal{D} \times X).$$

Hence

$$\langle e(\mathcal{L}_{A_0})e(\mathcal{V}_{A_1}), \overline{[A_1^\delta \circ (\mathcal{D} \times X)]} \cap [\mu] \rangle = N(A_1^\delta A_1, n_1, m_1, m_2) + \mathcal{C}_{\mathcal{B} \cap \mu},$$

where $\mathcal{C}_{\mathcal{B} \cap \mu}$ denotes the contribution of the section to the Euler class from the points of $\mathcal{B} \cap \mu$. We now give an explicit description of the boundary $\mathcal{B} \cap \mu$; the proof will be given in [1]. First of all we note that the whole of \mathcal{B} is not relevant while computing the contribution to the Euler class; only the points at which the section vanishes is relevant. Hence, we only need to consider the component of $\mathcal{B} \cap \mu$ where one (or more) of the q_i become equal to the last point $q_{\delta+1}$. Let us define

$$\mathcal{B}(q_1, q_{\delta+1}) := \{([f], q_1, \dots, q_{\delta+1}) \in \mathcal{B} : q_1 = q_{\delta+1}\}.$$

The spaces $\mathcal{B}(q_2, q_{\delta+1}), \dots, \mathcal{B}(q_\delta, q_{\delta+1})$ are defined similarly. It is shown in [5, Corollary 6.6] and [4, Proposition 6.1] that the contribution from $\mathcal{B}(q_1, q_{\delta+1}) \cap \mu$ is

$$N(A_1^{\delta-1}, n_1, m_1, m_2) + N(A_1^{\delta-1}, n_1 + 1, m_1, m_2) + 3N(A_1^{\delta-1} A_2, n_1, m_1, m_2).$$

The same contribution arises from $\mathcal{B}(q_i, q_{\delta+1}) \cap \mu$ for all $i = 1$ to δ . Hence the total contribution from $\mathcal{B}(q_1, q_{\delta+1}) \cap \mu, \dots, \mathcal{B}(q_\delta, q_{\delta+1}) \cap \mu$ is

$$\delta \left(N(A_1^{\delta-1}, n_1, m_1, m_2) + N(A_1^{\delta-1}, n_1 + 1, m_1, m_2) + 3N(A_1^{\delta-1} A_2, n_1, m_1, m_2) \right).$$

Next, let us define

$$\mathcal{B}(q_1, q_2, q_{\delta+1}) := \{([f], q_1, \dots, q_{\delta+1}) \in \mathcal{B} : q_1 = q_2 = q_{\delta+1}\}.$$

The spaces $\mathcal{B}(q_{i_1}, q_{i_2}, q_{\delta+1})$, and more generally $\mathcal{B}(q_{i_1}, q_{i_2}, \dots, q_{i_j}, q_{\delta+1})$, are defined similarly. Let us now focus on $\mathcal{B}(q_1, q_2, q_{\delta+1})$. We claim that

$$\mathcal{B}(q_1, q_2, q_{\delta+1}) \approx \overline{A_1^{\delta-2} \circ A_3}. \tag{10}$$

Let us give an intuitive justification as to why this is a believable claim. Geometrically, $\mathcal{B}(q_1, q_2, q_{\delta+1})$ denotes the component of the boundary when the nodal points q_1 and q_2 collide with each other. Geometrically, we expect the following thing to happen:




Figure 1: Two nodes colliding into a tacnode

That is indeed the case. We give a rigorous proof of (10) in [5, Lemma 6.3(2)]. We will show in [1] that the contribution from each of the tacnodal points, namely $\mathcal{B}(q_1, q_2, q_{\delta+1}) \cap \mu$ is 4. Hence the total contribution from all the components of type $\mathcal{B}(q_{i_1}, q_{i_2}, q_{\delta+1})$ is

$$4 \binom{\delta}{2} N(A_1^\delta A_3, n_1, m_1, m_2).$$

Next, let us focus on $\mathcal{B}(q_1, q_2, q_3, q_{\delta+1})$. We claim that

$$\mathcal{B}(q_1, q_2, q_3, q_{\delta+1}) \cap \mu \approx \overline{A_1^{\delta-3} \circ D_4} \cap \mu. \tag{11}$$

Again, let us give an intuitive justification as to why this is a believable claim. Geometrically, $\mathcal{B}(q_1, q_2, q_3, q_{\delta+1})$ denotes the component of the boundary when three of the nodal points q_1, q_2 and q_3 collide with each other. Geometrically, we expect the following thing to happen:




Figure 2: Three nodes colliding into a triple point

That is indeed the case. We will give a rigorous proof of (11) in [1]. We will also show that the contribution from each of the points of $\mathcal{B}(q_1, q_2, q_3, q_{\delta+1}) \cap \mu$ is 18. Hence the total contribution from all the components of type $\mathcal{B}(q_{i_1}, q_{i_2}, q_{i_3}, q_{\delta+1})$ is

$$18 \binom{\delta}{3} N(A_1^\delta D_4, n_1, m_1, m_2).$$

Finally, we will also show in [1] that if we consider a one dimensional family of r -nodal curves, then all the r -nodes can not collide together if $4 \leq r \leq 7$. In other words,

$$\mathcal{B}(q_1, q_2, \dots, q_r, q_{\delta+1}) \cap \mu = \emptyset \quad \forall 4 \leq r \leq 7. \quad (12)$$

Hence, the remaining stratum of \mathcal{B} does not contribute to the Euler class, giving us Theorem 6.3. \square

Remark 7.1. Note that (12) is true because we intersected the left hand side with μ . It is *not* true that

$$\mathcal{B}(q_1, q_2, \dots, q_r, q_{\delta+1}) = \emptyset \quad \forall 4 \leq r \leq 7.$$

Equation (12) is saying that when more than three nodes come together, the resulting singularity will have such a high codimension that it will not intersect a generic codimension one class (i.e. if we have a one dimensional family of curves, then more than three nodes can not come together). Similarly, (11) is true only with the μ present; it is *not* true that

$$\mathcal{B}(q_1, q_2, q_3, q_{\delta+1}) \cap \mu \approx \overline{A_1^{\delta-3} \circ D_4}. \quad (13)$$

Geometrically, the following thing can also occur:




Figure 3: Three nodes colliding into an A_5 -singularity

Hence, three nodes can collide to form an A_5 -singularity as well. However, since this is a codimension 5 singularity, its intersection with μ will be empty; only codimension four singularities will survive. All this will be proven rigorously in [1].

7.3 Proof of Theorem 6.4: computing $N(A_1^\delta \mathcal{P}A_1, n_1, m_1, m_2, \theta)$

When $\theta = 0$, the formula follows from (6). Let us now assume $\theta > 0$.

Recall that in section 5 we have defined the space

$$A_1^\delta \circ \overline{\hat{A}}_1 := \{([f], q_1, \dots, q_\delta, l_{q_{\delta+1}}) \in \mathcal{D} \times (X)^\delta \times \mathbb{P}TX : f \text{ has a singularity of type } A_1 \text{ at } q_1, \dots, q_\delta, \\ ([f], l_{q_{\delta+1}}) \in \overline{\hat{A}}_1, \quad q_1, \dots, q_{\delta+1} \text{ all distinct}\}.$$

Let μ be a generic pseudocycle representing the homology class Poincaré dual to

$$c_1^{n_1} x_1^{m_1} x_2^{m_2} \lambda^\theta y^{\delta L - (n_1 + m_1 + 2m_2 + \theta + \delta + 2)}.$$

We now define a section of the following bundle

$$\psi_{\mathcal{P}A_1} : A_1^\delta \circ \overline{\hat{A}}_1 \longrightarrow \mathcal{L}_{\mathcal{P}A_1} := \gamma_D^* \otimes \hat{\gamma}^{*2} \otimes L, \quad \text{given by} \\ \{\psi_{\mathcal{P}A_1}([f], q_1, \dots, q_\delta, l_{q_{\delta+1}})\}(f \otimes v^{\otimes 2}) := \nabla^2 f|_{q_{\delta+1}}(v, v).$$

Here $\hat{\gamma} \longrightarrow \mathbb{P}TX$ is the tautological line bundle over $\mathbb{P}TX$. We will show in [1] that if L is sufficiently $(2\delta + 2)$ -ample, then this section is transverse to the zero set. Next, let us define

$$\mathcal{B} := \overline{A_1^\delta \circ \overline{\hat{A}}_1} - A_1^\delta \circ \overline{\hat{A}}_1.$$

Hence

$$\langle e(\mathcal{L}_{\mathcal{P}A_1}), \overline{[A_1^\delta \circ \overline{\hat{A}}_1]} \cap [\mu] \rangle = N(A_1^\delta \mathcal{P}A_1, n_1, m_1, m_2, \theta) + \mathcal{C}_{\mathcal{B} \cap \mu}, \quad (14)$$

where as before, $\mathcal{C}_{\mathcal{B} \cap \mu}$ denotes the contribution of the section to the Euler class from $\mathcal{B} \cap \mu$. We now give an explicit description of \mathcal{B} . As before, we only need to consider the component of \mathcal{B} where one (or more) of the q_i become equal to the last point $q_{\delta+1}$. Let us define

$$\mathcal{B}(q_1, l_{q_{\delta+1}}) := \{([f], q_1, \dots, q_\delta, l_{q_{\delta+1}}) \in \mathcal{B} : q_1 = q_{\delta+1}\}.$$

The spaces $\mathcal{B}(q_2, q_{\delta+1}), \dots, \mathcal{B}(q_\delta, q_{\delta+1})$ are defined similarly. Following the same argument as in (10), we conclude that

$$\mathcal{B}(q_1, l_{q_{\delta+1}}) \approx \overline{A_1^{\delta-1} \circ \hat{A}_3}.$$

Let us now intersect $\overline{A_1^{\delta-1} \circ \hat{A}_3}$ with μ . This will be an isolated set of finite points. Hence, the section $\psi_{\mathcal{P}A_1}$ will not vanish on $\overline{A_1^{\delta-1} \circ \hat{A}_3} \cap \mu$. Hence it does not contribute to the Euler class. Next, let us define

$$\mathcal{B}(q_1, q_2, l_{q_{\delta+1}}) := \{([f], q_1, \dots, q_\delta, l_{q_{\delta+1}}) \in \mathcal{B} : q_1 = q_2 = l_{q_{\delta+1}}\}.$$

The spaces $\mathcal{B}(q_{i_1}, q_{i_2}, \dots, q_{i_j}, l_{q_{\delta+1}})$ are defined similarly. Following the same argument as in (11), we conclude that

$$\mathcal{B}(q_1, q_2, l_{q_{\delta+1}}) \approx \overline{A_1^{\delta-2} \circ \hat{D}_4}.$$

The section $\psi_{\mathcal{P}A_1}$ vanishes everywhere on $\overline{A_1^{\delta-2} \circ \hat{D}_4}$; hence it also vanishes on $\overline{A_1^{\delta-2} \circ \hat{D}_4} \cap \mu$. We will show in [1] that the contribution from each of the points of $\mathcal{B}(q_1, q_2, l_{q_{\delta+1}}) \cap \mu$ is 6. Hence the total contribution from all the components of type $\mathcal{B}(q_{i_1}, q_{i_2}, l_{q_{\delta+1}})$ is

$$6 \binom{\delta}{2} N(A_1^{\delta-2} \hat{D}_4, n_1, m_1, m_2, \theta).$$

Finally, using (12) we conclude that

$$\mathcal{B}(q_1, q_2, \dots, q_r, l_{q_{\delta+1}}) \cap \mu = \emptyset \quad \forall 3 \leq r \leq 7.$$

Hence the remaining stratum of $\mathcal{B} \cap \mu$ does not contribute to the Euler class, giving Theorem (6.4). \square

Remark 7.2. The reason we need to compute $N(A_1^\delta \mathcal{P}A_1, n_1, m_1, m_2, \theta)$ is because it arises in the recursive formula for $N(A_1^\delta \mathcal{P}A_2, n_1, m_1, m_2, \theta)$, as will be clear shortly.

7.4 Proof of Theorem 6.5: computing $N(A_1^\delta \mathcal{P}A_2, n_1, m_1, m_2, \theta)$

Recall that in section 5 we have defined the space

$$A_1^\delta \circ \overline{\mathcal{P}A_1} := \{([f], q_1, \dots, q_\delta, l_{q_{\delta+1}}) \in \mathcal{D} \times (X)^\delta \times \mathbb{P}TX : f \text{ has a singularity of type } A_1 \text{ at } q_1, \dots, q_\delta, \\ ([f], l_{q_{\delta+1}}) \in \overline{\mathcal{P}A_1}, \quad q_1, \dots, q_{\delta+1} \text{ all distinct}\}.$$

Let μ be a generic pseudocycle representing the homology class Poincaré dual to

$$c_1^{n_1} x_1^{m_1} x_2^{m_2} \lambda^\theta y^{\delta_L - (n_1 + m_1 + 2m_2 + \theta + \delta + 2)}.$$

We now define a section of the following line bundle

$$\Psi_{\mathcal{P}A_2} : A_1^\delta \circ \overline{\mathcal{P}A_1} \longrightarrow \mathbb{L}_{\mathcal{P}A_2} := \gamma_D^* \otimes \hat{\gamma}^* \otimes (\pi^* TX / \hat{\gamma})^* \otimes L, \quad \text{given by} \\ \{\Psi_{\mathcal{P}A_2}([f], q_1, \dots, q_\delta, l_{q_{\delta+1}})\}(f \otimes v \otimes w) := \nabla^2 f|_{q_{\delta+1}}(v, w).$$

If L is sufficiently $(2\delta + 2)$ -ample, then this section is transverse to zero. Next, let us define

$$\mathcal{B} := \overline{A_1^\delta \circ \overline{\mathcal{P}A_1}} - A_1^\delta \circ \overline{\mathcal{P}A_1}.$$

Hence

$$\langle e(\mathbb{L}_{\mathcal{P}A_2}), [\overline{A_1^\delta \circ \overline{\mathcal{P}A_1}}] \cap [\mu] \rangle = N(A_1^\delta \mathcal{P}A_2, n_1, m_1, m_2, \theta) + \mathcal{C}_{\mathcal{B} \cap \mu}.$$

We now give an explicit description of \mathcal{B} . As before, we only need to consider the component of \mathcal{B} where one (or more) of the q_i become equal to the last point $q_{\delta+1}$. Define $\mathcal{B}(q_{i_1}, \dots, q_{i_k}, l_{q_\delta})$ as before. We will show in [1] that

$$\mathcal{B}(q_1, l_{q_{\delta+1}}) \approx \overline{A_1^{\delta-1} \circ \mathcal{P}A_3} \cup \overline{A_1^{\delta-1} \circ \hat{D}_4}.$$

Furthermore, we show that the contribution from $\overline{A_1^{\delta-1} \circ \mathcal{P}A_3} \cap \mu$ is 2, while the contribution from $\overline{A_1^{\delta-1} \circ \hat{D}_4} \cap \mu$ is 3. Hence the total contribution from all the components of type $\mathcal{B}(q_{i_1}, l_{q_{\delta+1}})$ equals

$$2 \binom{\delta}{1} N(A_1^{\delta-1} \mathcal{P}A_3, n_1, m_1, m_2, \theta) + 3 \binom{\delta}{1} N(A_1^{\delta-1} \hat{D}_4, n_1, m_1, m_2, \theta).$$

Next, we show in [1] that

$$\mathcal{B}(q_1, q_2, l_{q_{\delta+1}}) \approx \overline{A_1^{\delta-2} \circ \mathcal{P}D_4} \cup \overline{A_1^{\delta-2} \circ \hat{D}_5}.$$

We also show that the contribution from $\overline{A_1^{\delta-2} \circ \hat{D}_4} \cap \mu$ is 2, while the contribution from $\overline{A_1^{\delta-2} \circ \mathcal{P}D_5}$ is 3. Hence the total contribution from all the components of type $\mathcal{B}(q_{i_1}, q_{i_2}, l_{q_{\delta+1}})$ equals

$$4 \binom{\delta}{2} N(A_1^{\delta-2} \mathcal{P}D_4, n_1, m_1, m_2, \theta) + 4 \binom{\delta}{2} N(A_1^{\delta-2} \hat{D}_5, n_1, m_1, m_2, \theta).$$

Finally, we show in [1] that

$$\mathcal{B}(q_1, q_2, \dots, q_r, l_{q_{\delta+1}}) \cap \mu = \emptyset \quad \forall 3 \leq r \leq 5. \quad (15)$$

Hence the remaining stratum of \mathcal{B} does not contribute to the Euler class, giving us Theorem 6.5. \square

Remark 7.3. Although we have devoted [1] for the rigorous proofs of our closure claims, there is very subtle point about equation (15) that we should mention here. Let us look at the claim for $r = 3$. This is essentially saying that four nodes can not come together, once we intersect the variety with μ . Let us see if this claim is believable; consider the following picture:




Figure 4: Four nodes collapsing to a D6-singularity

Now it seems naively that one of the components of $\mathcal{B}(q_1, q_2, q_3, l_{q_4})$ is \hat{D}_6 . If that were indeed the case then the intersection with μ would not be zero when $\theta = 1$ (it would be zero when $\theta = 0$). Hence, what we are claiming here is that one of the components of $\mathcal{B}(q_1, q_2, \dots, q_3, l_{q_4})$ is *not* \hat{D}_6 , but is in fact \mathcal{PD}_6 . Let us look at it a bit more carefully. We are *not* considering here the space of curves with four distinct nodes. We are in fact considering the space of curves with three nodes and one node with a distinguished branch of the node; namely the space $A_1^3 \circ \mathcal{PA}_1$ as opposed to A_1^4 . Hence, any random curve with a D_6 -singularity will not be in the closure; i.e. all elements of \hat{D}_6 are not in the closure. Only those elements will be in the closure where a specific directional derivative vanishes, namely only elements of \mathcal{PD}_6 will be in the closure. Hence, (15) is consistent with the picture we have drawn.

Next, let us look at the claim for $r = 5$. This is essentially saying that six nodes can not come together, once we intersect the variety with μ . Again, let us see if this claim is believable by looking at the following picture:




Figure 5: Six nodes collapsing to an ordinary quadruple point

Now it seems naively that one of the components of $\mathcal{B}(q_1, q_2, q_3, q_4, q_5, l_{q_6})$ is \hat{X}_9 . If that were indeed the case then the intersection with μ would not be zero when $\theta = 1$ (it would be zero when $\theta = 0$). Hence, what we are claiming here is that one of the components of $\mathcal{B}(q_1, q_2, \dots, q_5, l_{q_6})$ is *not* \hat{X}_9 , but is in fact $\mathcal{P}X_9$. Let us look at it a bit more carefully. Again, we are *not* considering here the space of curves with six distinct nodes. We are in fact considering the space of curve with five nodes and one node with a distinguished branch of the node; namely the space $A_1^5 \circ \mathcal{P}A_1$ as opposed to A_1^6 . Hence, any random curve with an X_9 singularity will not be in the closure; i.e. all elements of \hat{X}_9 are not in the closure. Only those elements will be in the closure where a specific directional derivative vanishes, namely only elements of $\mathcal{P}X_9$ will be in the closure. Hence, (15) is consistent with the picture we have drawn.

7.5 Proof of Theorem 6.6: computing $N(A_1^\delta \mathcal{P}A_3, n_1, m_1, m_2, \theta)$

Recall that in section 5 we have defined the space

$$A_1^\delta \circ \overline{\mathcal{P}A_2} := \{([f], q_1, \dots, q_\delta, l_{q_{\delta+1}}) \in \mathcal{D} \times (X)^\delta \times \mathbb{P}TX : f \text{ has a singularity of type } A_1 \text{ at } q_1, \dots, q_\delta, \\ ([f], l_{q_{\delta+1}}) \in \overline{\mathcal{P}A_2}, q_1, \dots, q_{\delta+1} \text{ all distinct}\}.$$

Let μ be a generic pseudocycle representing the homology class Poincaré dual to

$$c_1^{n_1} x_1^{m_1} x_2^{m_2} \lambda^\theta y^{\delta L - (n_1 + m_1 + 2m_2 + \theta + \delta + 3)}.$$

We now define a section of the following bundle

$$\Psi_{\mathcal{P}A_3} : A_1^\delta \circ \overline{\mathcal{P}A_2} \longrightarrow \mathbb{L}_{\mathcal{P}A_3} := \gamma_D^* \otimes \hat{\gamma}^{*3} \otimes L, \quad \text{given by} \\ \{\Psi_{\mathcal{P}A_3}([f], q_1, \dots, q_\delta, l_{q_{\delta+1}})\}(f \otimes v^{\otimes 3}) := \nabla^3 f|_{q_{\delta+1}}(v, v, v).$$

By [2, Lemma 7.1, statement 8], we have that

$$\overline{\mathcal{P}A_2} = \mathcal{P}A_2 \cup \overline{\mathcal{P}A_3} \cup \overline{D_4}. \quad (16)$$

Hence, let us define

$$\mathcal{B} := \overline{A_1^\delta \circ \overline{\mathcal{P}A_2}} - A_1^\delta \circ (\mathcal{P}A_2 \cup \overline{\mathcal{P}A_3}).$$

If L is sufficiently $(2\delta + 3)$ -ample, then the section $\Psi_{\mathcal{P}A_3}$ vanishes on the points of $A_1^\delta \circ \mathcal{P}A_3$ transversally (all the points of $A_1^\delta \circ \mathcal{P}A_3$ are smooth points of the variety $A_1^\delta \circ \overline{\mathcal{P}A_2}$). Hence,

$$\langle e(\mathbb{L}_{\mathcal{P}A_3}), \overline{[A_1^\delta \circ \overline{\mathcal{P}A_2}] \cap [\mu]} \rangle = N(A_1^\delta \mathcal{P}A_3, n_1, m_1, m_2, \theta) + \mathcal{C}_{\mathcal{B} \cap \mu}.$$

We now give an explicit description of \mathcal{B} . Let us first define

$$\mathcal{B}_0 := \{([f], q_1, \dots, q_\delta, l_{q_{\delta+1}}) \in \mathcal{B} : q_1, q_2, \dots, q_{\delta+1} \text{ are all distinct}\}.$$

In other words, \mathcal{B}_0 is that component of the boundary, where all the points are still distinct. By (16), we conclude that

$$\mathcal{B}_0 = \overline{A_1^\delta} \circ \overline{\hat{D}_4}.$$

If we intersect \mathcal{B}_0 with μ then we will get a finite set of points. Since the representative μ is generic, we conclude that the third derivative along v will not vanish, i.e. the section $\Psi_{\mathcal{P}A_3}$ will not vanish on those points. Hence, $\mathcal{B}_0 \cap \mu$ does not contribute to the Euler class.

Next, let us consider the components of \mathcal{B} where one (or more) of the q_i become equal to the last point $q_{\delta+1}$. Define $\mathcal{B}(q_{i_1}, \dots, q_{i_k}, l_{q_\delta})$ as before. We show in [5, Lemma 3.1] that

$$\mathcal{B}(q_1, l_{q_{\delta+1}}) \approx \overline{A_1^{\delta-1} \circ \mathcal{P}A_4} \cup \overline{A_1^{\delta-1} \circ \hat{D}_5}.$$

We also show in [5] (Corollary 6.13) that the contribution to the Euler class from each of the points of $\overline{A_1^{\delta-1} \circ \mathcal{P}A_4} \cap \mu$ is 2. Furthermore, the section $\Psi_{\mathcal{P}A_3}$ does not vanish on $\overline{A_1^{\delta-1} \circ \hat{D}_5} \cap \mu$, since μ is generic. Hence, the total contribution from all the components of type $\mathcal{B}(q_{i_1}, l_{q_{\delta+1}})$ equals

$$2 \binom{\delta}{1} N(A_1^{\delta-1} \mathcal{P}A_4, n_1, m_1, m_2, \theta).$$

Next, we analyze what happens when two nodes and one cusp come together. We show in [1] that

$$\mathcal{B}(q_1, q_2, l_{q_{\delta+1}}) \approx \overline{A_1^{\delta-2} \circ \mathcal{P}D_5}.$$

Geometrically this is a believable statement due to the following picture:




Figure 6: Two nodes and one cusp collapsing to a D_5 -singularity

We also show that the contribution from each of the points of $\overline{A_1^{\delta-2} \circ \mathcal{P}D_5} \cap \mu$ is 2. Hence the total contribution from all the components of type $\mathcal{B}(q_{i_1}, q_{i_2}, l_{q_{\delta+1}})$ equals

$$2 \binom{\delta}{2} N(A_1^{\delta-2} \mathcal{P}D_5, n_1, m_1, m_2, \theta).$$

Finally, we claim that

$$\mathcal{B}(q_1, q_2, \dots, q_r, l_{q_{\delta+1}}) \cap \mu = \emptyset \quad \forall 3 \leq r \leq 5.$$

In other words, a one dimensional family of curves with r -nodes and one cusp can not come together if $3 \leq r \leq 5$. Hence the remaining stratum of \mathcal{B} does not contribute to the Euler class, giving us Theorem 6.6. \square

7.6 Proof of Theorem 6.7: computing $N(A_1^\delta \mathcal{P}A_4, n_1, m_1, m_2, \theta)$

Recall that in section 5 we have defined the space

$$A_1^\delta \circ \overline{\mathcal{P}A_3} := \{([f], q_1, \dots, q_\delta, l_{q_{\delta+1}}) \in \mathcal{D} \times (X)^\delta \times \mathbb{P}TX : f \text{ has a singularity of type } A_1 \text{ at } q_1, \dots, q_\delta, \\ ([f], l_{q_{\delta+1}}) \in \overline{\mathcal{P}A_3}, q_1, \dots, q_{\delta+1} \text{ all distinct}\}.$$

Let μ be a generic pseudocycle representing the homology class Poincaré dual to

$$c_1^{n_1} x_1^{m_1} x_2^{m_2} \lambda^\theta y^{\delta L - (n_1 + m_1 + 2m_2 + \theta + \delta + 4)}.$$

Let $v \in \hat{\gamma}$ and $w \in \pi^*TX/\hat{\gamma}$ be two fixed non zero vectors. Let us introduce the following abbreviation:

$$f_{ij} := \nabla^{i+j} f|_q \left(\underbrace{v, \dots, v}_i, \underbrace{w, \dots, w}_j \right).$$

We now define a section of the following bundle

$$\psi_{\mathcal{P}A_4} : A_1^\delta \circ \overline{\mathcal{P}A_3} \longrightarrow \mathbb{L}_{\mathcal{P}A_4} := \gamma_D^{*2} \otimes \hat{\gamma}^{*4} \otimes (\pi^*TX/\hat{\gamma})^{*2} \otimes L^{*2}, \\ \{\psi_{\mathcal{P}A_4}([f], q_1, \dots, q_\delta, l_{q_{\delta+1}})\} (f^{\otimes 2} \otimes v^{\otimes 4} \otimes w^{\otimes 2}) := f_{02} A_4^f, \quad \text{where} \quad A_4^f := f_{40} - \frac{3f_{21}^2}{f_{02}}.$$

By [2, Lemma 7.1, statement 9], we have that

$$\overline{\mathcal{P}A_3} = \mathcal{P}A_3 \cup \overline{\mathcal{P}A_4} \cup \overline{\mathcal{P}D_4}. \quad (17)$$

Hence, let us define

$$\mathcal{B} := \overline{A_1^\delta \circ \overline{\mathcal{P}A_3}} - A_1^\delta \circ (\mathcal{P}A_3 \cup \overline{\mathcal{P}A_4}).$$

If L is sufficiently $(2\delta + 4)$ -ample, then the section $\Psi_{\mathcal{P}A_4}$ vanishes on the points of $A_1^\delta \circ \mathcal{P}A_4$ transversally (all the points of $A_1^\delta \circ \mathcal{P}A_4$ are smooth points of the variety $A_1^\delta \circ \overline{\mathcal{P}A_3}$). Next, let us define

$$\mathcal{B} := \overline{A_1^\delta \circ \mathcal{P}A_3} - A_1^\delta \circ \mathcal{P}A_3 \cup \overline{\mathcal{P}A_4}.$$

Hence,

$$\langle e(\mathbb{L}_{\mathcal{P}A_4}), [\overline{A_1^\delta \circ \overline{\mathcal{P}A_3}}] \cap [\mu] \rangle = N(A_1^\delta \mathcal{P}A_4, n_1, m_1, m_2, \theta) + \mathcal{C}_{\mathcal{B} \cap \mu}.$$

We now give an explicit description of \mathcal{B} . As before, let us first define \mathcal{B}_0 as

$$\mathcal{B}_0 := \{([f], q_1, \dots, q_\delta, l_{q_{\delta+1}}) \in \mathcal{B} : q_1, q_2, \dots, q_{\delta+1} \text{ are all distinct}\}.$$

Hence, \mathcal{B}_0 is that component of the boundary, where all the points are still distinct. By (17), we conclude that

$$\mathcal{B}_0 = \overline{A_1^\delta \circ \overline{\mathcal{P}D_4}}.$$

If we intersect \mathcal{B}_0 with μ then we will get a finite set of points. Since the representative μ is generic, we conclude that the directional derivative f_{21} will not vanish on those points. Since $f_{02} = 0$ on \mathcal{B}_0 we conclude that

$$f_{02}A_4^f = f_{02}f_{40} - 3f_{21}^2 \neq 0$$

if $f_{21} \neq 0$. Hence the section $\Psi_{\mathcal{P}A_4}$ will not vanish on $\mathcal{B}_0 \cap \mu$.

Next, let us consider the components of \mathcal{B} where one (or more) of the q_i become equal to the last point $q_{\delta+1}$. Define $\mathcal{B}(q_{i_1}, \dots, q_{i_k}, l_{q_\delta})$ as before. We show in [5, Lemma 6.3(4)], that

$$\mathcal{B}(q_1, l_{q_{\delta+1}}) \approx \overline{A_1^{\delta-1} \circ \mathcal{P}A_5} \cup \overline{A_1^{\delta-1} \circ \mathcal{P}D_5^\vee},$$

where we define $\mathcal{P}D_k^\vee$ to be the space

$$\mathcal{P}D_k^\vee := \{([f], l_q) \in \mathcal{D} \times \mathbb{P}TX : ([f], q) \in \mathcal{D}_k, f_{30} = 0 \text{ and } f_{21} \neq 0\}. \quad (18)$$

The section $\Psi_{\mathcal{P}A_4}$ will not vanish on $\mathcal{P}D_5^\vee$, since by definition on $\mathcal{P}D_k^\vee$ we have $f_{02} = 0$ but $f_{21} \neq 0$. Hence, the section will not vanish on $\overline{A_1^{\delta-1} \circ \mathcal{P}D_5^\vee} \cap \mu$. We also show in [5, Corollary 6.13], that the contribution of the section $\Psi_{\mathcal{P}A_4}$ to the Euler class from each of the points of $\overline{A_1^{\delta-1} \circ \mathcal{P}A_5} \cap \mu$ is 2. Hence the total contribution from all the components of type $\mathcal{B}(q_{i_1}, l_{q_{\delta+1}})$ equals

$$2 \binom{\delta}{1} N(A_1^{\delta-1} \mathcal{P}A_5, n_1, m_1, m_2, \theta).$$

Next, we analyze what happens when two nodes and one tacnode come together. We show in [1] that

$$\mathcal{B}(q_1, q_2, l_{q_{\delta+1}}) \cap \mu \approx \overline{A_1^{\delta-2} \circ \mathcal{P}D_6} \cap \mu.$$

Geometrically this is a believable statement due to the following picture:




Figure 7: Two nodes and one tacnode collapsing to a D_6 -singularity

We also show that the contribution from each of the points of $\overline{A_1^{\delta-2} \circ \mathcal{P}D_6} \cap \mu$ is 4. Hence the total contribution from all the components of type $\mathcal{B}(q_{i_1}, q_{i_2}, l_{q_{\delta+1}})$ equals

$$4 \binom{\delta}{2} N(A_1^{\delta-2} \mathcal{P}D_6, n_1, m_1, m_2, \theta).$$

Finally, we show in [1] that

$$\mathcal{B}(q_1, q_2, \dots, q_r, l_{q_{\delta+1}}) \cap \mu = \emptyset \quad \forall 3 \leq r \leq 4.$$

In other words, a one dimensional family of curves with r -nodes and one tacnode can not come together if $3 \leq r \leq 4$. Hence the remaining stratum of \mathcal{B} does not contribute to the Euler class, giving us Theorem 6.7. \square

7.7 Proof of Theorem 6.8: computing $N(A_1^\delta \mathcal{P}D_4, n_1, m_1, m_2, \theta)$

Let us first do the case $\theta = 0$. Define a section of the following bundle

$$\begin{aligned} \Psi_{\mathcal{P}D_4} : A_1^\delta \circ \overline{\mathcal{P}A_3} &\longrightarrow \mathbb{L}_{\mathcal{P}D_4} := \gamma_D^* \otimes (\pi^* TX / \hat{\gamma})^{*2} \otimes L, & \text{given by} \\ \{\Psi_{\mathcal{P}D_4}([f], q_1, \dots, q_\delta, l_{q_{\delta+1}})\} & (f \otimes w^{\otimes 2}) := \nabla^2 f|_q(w, w). \end{aligned}$$

By [2, Lemma 7.1, statement 9], we conclude that

$$\overline{\mathcal{P}A_3} = \mathcal{P}A_3 \cup \overline{\mathcal{P}A_4} \cup \overline{\mathcal{P}D_4}. \quad (19)$$

Hence, let us define

$$\mathcal{B} := \overline{A_1^\delta \circ \overline{\mathcal{P}A_3}} - A_1^\delta \circ (\mathcal{P}A_3 \cup \overline{\mathcal{P}D_4}).$$

If L is sufficiently $(2\delta + 4)$ -ample, then the section $\Psi_{\mathcal{P}D_4}$ vanishes on the points of $A_1^\delta \circ \mathcal{P}D_4$ transversally (all the points of $A_1^\delta \circ \mathcal{P}D_4$ are smooth points of the variety $A_1^\delta \circ \overline{\mathcal{P}A_3}$). Hence,

$$\langle e(\mathbb{L}_{\mathcal{P}D_4}), [\overline{A_1^\delta \circ \mathcal{P}A_3}] \cap [\mu] \rangle = N(A_1^\delta \mathcal{P}D_4, n_1, m_1, m_2, \theta) + \mathcal{C}_{\mathcal{B} \cap \mu}.$$

We now give an explicit description of \mathcal{B} . Let us define \mathcal{B}_0 as before; it denotes the component of the boundary where all the points are distinct. The section $\Psi_{\mathcal{P}D_4}$ does not vanish on $\overline{A_1^\delta \circ \mathcal{P}A_4}$; hence (19), the section does not vanish on \mathcal{B}_0 .

Next, let us analyze the component of \mathcal{B} where one (or more) of the q_i become equal to the last point $q_{\delta+1}$. Define $\mathcal{B}(q_{i_1}, \dots, q_{i_k}, l_{q_\delta})$ as before. We show in [5, Lemma 6.3(4)] that

$$\mathcal{B}(q_1, l_{q_{\delta+1}}) \approx \overline{A_1^{\delta-1} \circ \mathcal{P}A_5} \cup \overline{A_1^{\delta-1} \circ \mathcal{P}D_5^\vee},$$

where $\mathcal{P}D_5^\vee$ is as defined in (18). We also show in [5, Corollary 6.17] that the section $\Psi_{\mathcal{P}D_4}$ vanishes on $A_1^{\delta-1} \circ \mathcal{P}D_5^\vee \cap \mu$ with a multiplicity of 2. Furthermore, we notice that the section does not vanish on $A_1^{\delta-1} \circ \mathcal{P}A_5 \cap \mu$. Hence the total contribution from all the components of type $\mathcal{B}(q_{i_1}, l_{q_{\delta+1}})$ equals

$$2 \binom{\delta}{1} N(A_1^{\delta-1} \mathcal{P}D_5^\vee, n_1, m_1, m_2, 0).$$

Next, we note that the projection map $\pi : A_1^\delta \circ \mathcal{P}D_5^\vee \rightarrow A_1^\delta \circ D_5$ is one to one. Hence,

$$N(A_1^{\delta-1} \mathcal{P}D_5^\vee, n_1, m_1, m_2, 0) = N(A_1^{\delta-1} D_5, n_1, m_1, m_2).$$

Hence the total contribution from all the components of type $\mathcal{B}(q_{i_1}, l_{q_{\delta+1}})$ equals

$$2 \binom{\delta}{1} N(A_1^{\delta-1} D_5, n_1, m_1, m_2).$$

Next, let us describe the boundary when three of the points come together. As mentioned earlier in the Proof of Theorem 6.7, we show in [1] that

$$\mathcal{B}(q_1, q_2, l_{q_{\delta+1}}) \approx \overline{A_1^{\delta-2} \circ \mathcal{P}D_6}.$$

We also show that the contribution from each of the points of $\overline{A_1^{\delta-2} \circ \mathcal{P}D_6} \cap \mu$ is 2. Hence the total contribution from all the components of type $\mathcal{B}(q_{i_1}, q_{i_2}, l_{q_{\delta+1}})$ equals

$$2 \binom{\delta}{2} N(A_1^{\delta-2} \mathcal{P}D_6, n_1, m_1, m_2, \theta).$$

Finally, we show in [1] that

$$\mathcal{B}(q_1, q_2, \dots, q_r, l_{q_{\delta+1}}) \cap \mu = \emptyset \quad \forall 3 \leq r \leq 4.$$

In other words, if we have a one dimensional family of curves with r -nodes and one tacnode, then all the points can not come together, if r is 3 or 4. Hence the remaining stratum of \mathcal{B} does not contribute to the Euler class, giving us Theorem 6.8 for $\theta = 0$.

Next, we consider the case $\theta = 1$. Let us consider the following space in section 5, namely

$$A_1^\delta \circ \overline{\widehat{D}}_4 := \{([f], q_1, \dots, q_\delta, l_{q_{\delta+1}}) \in \mathcal{D} \times (X)^\delta \times \mathbb{P}TX : f \text{ has a singularity of type } A_1 \text{ at } q_1, \dots, q_\delta, \\ ([f], l_{q_{\delta+1}}) \in \overline{\widehat{D}}_4, \quad q_1, \dots, q_{\delta+1} \text{ all distinct}\}.$$

Let μ be a generic pseudocycle representing the homology class Poincaré dual to

$$c_1^{n_1} x_1^{m_1} x_2^{m_2} \lambda^\theta y^{\delta_L - (n_1 + m_1 + 2m_2 + \theta + \delta + 4)}.$$

We again consider the section of the following bundle

$$\Psi_{\mathcal{P}A_3} : A_1^\delta \circ \overline{\widehat{D}}_4 \rightarrow \mathbb{L}_{\mathcal{P}A_3} := \gamma_D^* \otimes \hat{\gamma}^{*3} \otimes L, \\ \{\Psi_{\mathcal{P}A_3}([f], q_1, \dots, q_\delta, l_{q_{\delta+1}})\} (f \otimes v^{\otimes 3}) := \nabla^3 f|_q(v, v, v).$$

If L is sufficiently $(2\delta + 4)$ -ample, then this section is transverse to the zero set. Let,

$$\mathcal{B} := \overline{A_1^\delta \circ \widehat{D}_4} - A_1^\delta \circ \widehat{D}_4.$$

Hence

$$\langle e(\mathbb{L}_{\mathcal{P}A_3}), \overline{[A_1^\delta \circ \widehat{D}_4]} \cap [\mu] \rangle = N(A_1^\delta \mathcal{P}D_4, n_1, m_1, m_2, \theta) + \mathcal{C}_{\mathcal{B} \cap \mu}.$$

Let us denote \mathcal{B}_0 to be the component of the boundary where all the points are distinct. The section $\Psi_{\mathcal{P}A_3}$ will not vanish on $\mathcal{B}_0 \cap \mu$, since the last marked point will comprise of a generic D_5 -singularity.

Next, let us consider the component when one or more points come together. Define $\mathcal{B}(q_1, \dots, q_r, l_{q_{\delta+1}})$ as before. We show in [1] that

$$\mathcal{B}(q_1, l_{q_{\delta+1}}) = \widehat{D}_6.$$

Again, when we intersect \widehat{D}_6 , the section $\Psi_{\mathcal{P}A_3}$ will not vanish. Hence, it does not contribute to the Euler class. Next, we show in [1] that

$$\mathcal{B}(q_1, q_2, l_{q_{\delta+1}}) \cap \mu = \emptyset.$$

Hence, $\mathcal{B}(q_1, q_2, l_{q_{\delta+1}})$ does not contribute to the Euler class. Finally, we show that when three nodes and one triple point come together, we get a triple point. In other words, we show in [1] that:

$$\mathcal{B}(q_1, q_2, q_3, l_{q_4}) = \widehat{X}_9. \tag{20}$$

Geometrically this is a believable statement due to the following picture:




Figure 8: Three nodes and one D_4 -singularity collapsing to a quadruple point

We also show in [1] that the contribution of the section from each of the points of $\overline{X}_9 \cap \mu$ is 24. That gives us Theorem 6.8 for $\theta = 1$. \square

7.8 Proof of Theorem 6.10: computing $N(A_1^\delta \mathcal{P}D_5, n_1, m_1, m_2, \theta)$

Recall that in section 5, we have defined the following space

$$A_1^\delta \circ \overline{\mathcal{P}D}_4 := \{([f], q_1, \dots, q_\delta, l_{q_{\delta+1}}) \in \mathcal{D} \times (X)^\delta \times \mathbb{P}TX : f \text{ has a singularity of type } A_1 \text{ at } q_1, \dots, q_\delta, \\ ([f], l_{q_{\delta+1}}) \in \overline{\mathcal{P}D}_4, q_1, \dots, q_{\delta+1} \text{ all distinct}\}.$$

Let μ be a generic pseudocycle representing the homology class Poincaré dual to

$$c_1^{n_1} x_1^{m_1} x_2^{m_2} \lambda^\theta y^{\delta L - (n_1 + m_1 + 2m_2 + \theta + \delta + 5)}.$$

As before, we let $v \in \hat{\gamma}$ and $w \in \pi^*TX/\hat{\gamma}$ be two fixed non zero vectors and let f_{ij} be defined as before. We now define a section of the following bundle

$$\Psi_{\mathcal{P}D_5} : A_1^\delta \circ \overline{\mathcal{P}D}_4 \longrightarrow \mathbb{L}_{\mathcal{P}D_5} := \gamma_D^* \otimes \hat{\gamma}^{*2} \otimes (\pi^*TX/\hat{\gamma})^* \otimes L^*, \quad \text{given by} \\ \{\Psi_{\mathcal{P}D_5}([f], q_1, \dots, q_\delta, l_{q_{\delta+1}})\}(f \otimes v^{\otimes 2} \otimes w) := f_{21}.$$

In [2, Lemma 7.1, statement 4], we prove that

$$\overline{\mathcal{P}D}_4 = \mathcal{P}D_4 \cup \overline{\mathcal{P}D}_5 \cup \overline{\mathcal{P}D}_5^\vee. \quad (21)$$

Hence, let us define

$$\mathcal{B} := \overline{A_1^\delta \circ \overline{\mathcal{P}D}_4} - A_1^\delta \circ (\mathcal{P}D_4 \cup \overline{\mathcal{P}D}_5).$$

If L is sufficiently $(2\delta + 4)$ -ample, then the section $\Psi_{\mathcal{P}D_5}$ vanishes on the points of $A_1^\delta \circ \mathcal{P}D_5$ transversally (all the points of $A_1^\delta \circ \mathcal{P}D_5$ are smooth points of the variety $A_1^\delta \circ \overline{\mathcal{P}D}_4$). Hence,

$$\langle e(\mathbb{L}_{\mathcal{P}D_5}), [\overline{A_1^\delta \circ \overline{\mathcal{P}D}_4}] \cap [\mu] \rangle = N(A_1^\delta \mathcal{P}D_5, n_1, m_1, m_2, \theta) + \mathcal{C}_{\mathcal{B} \cap \mu}.$$

Let us define \mathcal{B}_0 as before, namely the component of the boundary where all the points are distinct. The section $\Psi_{\mathcal{P}D_5}$ does not vanish on $\mathcal{P}D_5^\vee$ by definition (recall that by definition of the space $\mathcal{P}D_5^\vee$, the directional derivative $f_{21} \neq 0$). Hence, the section $\Psi_{\mathcal{P}D_5}$ does not vanish on $\overline{A_1^\delta \circ \overline{\mathcal{P}D}_5^\vee} \cap \mu$ for a generic μ . Therefore, \mathcal{B}_0 does not contribute to the Euler class.

Next, let us analyze the boundary when one or more points come together; define $\mathcal{B}(q_{i_1}, \dots, q_{i_k}, l_{q_\delta})$ as before. In [1], we show that

$$\mathcal{B}(q_1, l_{q_{\delta+1}}) \approx \overline{A_1^{\delta-1} \circ \mathcal{P}D_6} \cup \overline{A_1^{\delta-1} \circ \mathcal{P}D_6^\vee}.$$

Again, by the definition of $\mathcal{P}D_k^\vee$, the directional derivative f_{21} does not vanish on $\mathcal{P}D_6^\vee$; hence the section $\overline{\Psi_{\mathcal{P}D_5}}$ does not vanish on $\overline{A_1^{\delta-1} \circ \mathcal{P}D_6^\vee} \cap \mu$. We also show in [1] that the section vanishes on $\overline{A_1^{\delta-1} \circ \mathcal{P}D_6} \cap \mu$ with a multiplicity of 2. Hence the total contribution from all the components of type $\mathcal{B}(q_{i_1}, l_{q_{\delta+1}})$ equals

$$2 \binom{\delta}{1} N(A_1^{\delta-1} \mathcal{P}D_6, n_1, m_1, m_2, \theta).$$

Next, let us consider the boundary when two points come together. We show in [1] that

$$\mathcal{B}(q_1, q_2, l_{q_{\delta+1}}) \approx \overline{A_1^{\delta-2} \circ \hat{X}_9}.$$

Furthermore, we show that the section vanishes on $\overline{A_1^{\delta-2} \circ \hat{X}_9}$. Hence, the contribution from all points of type $\mathcal{B}(q_{i_1}, q_{i_2}, l_{q_{\delta+1}})$ equals

$$12 \binom{\delta}{1} N(A_1^{\delta-2} \hat{X}_9, n_1, m_1, m_2, \theta).$$

Finally, we show in [1] that

$$\mathcal{B}(q_1, q_2, q_3, l_{q_4}) \cap = \overline{\mathcal{P}X_9}. \tag{22}$$

This is geometrically believable due to the following picture:




Figure 9: Three nodes and one triple point collapsing to a quadruple point

We also show in [1] that the contribution from each of the point of $\overline{\mathcal{P}X}_9$ is 18, thereby giving us Theorem 6.10. \square

Remark 7.4. We remind the reader again that there is a subtle difference between (20) and (22). Both the statements happen to be true, but they are not exactly the same statements. In the first statement we are looking at the space of curves with three nodes and one triple point; in the closure we get a quadruple point. In other words we were looking at the space $A_1^3 \circ \hat{D}_4$ and getting \hat{X}_9 in the closure.

In the second case, we are looking at the space of curves with three nodes and a triple point with a distinguished branch along which the third derivative vanishes. In other words, we are looking at the space $A_1 \circ \mathcal{P}D_4$ as opposed to $A_1^3 \circ \hat{D}_4$. What we get in the closure is not any random quadruple point. We get a quadruple point with a distinguished branch such that the fourth derivative along that direction vanishes. In other words, we do not get \hat{X}_9 in the closure, but rather $\mathcal{P}X_9$.

7.9 Proof of Theorem 6.11: computing $N(A_1^\delta \mathcal{P}A_5, n_1, m_1, m_2, \theta)$

Recall that in section 5 we have defined the space

$$A_1^\delta \circ \overline{\mathcal{P}A}_4 := \{([f], q_1, \dots, q_\delta, l_{q_{\delta+1}}) \in \mathcal{D} \times (X)^\delta \times \mathbb{P}TX : f \text{ has a singularity of type } A_1 \text{ at } q_1, \dots, q_\delta, \\ ([f], l_{q_{\delta+1}}) \in \overline{\mathcal{P}A}_4, \quad q_1, \dots, q_{\delta+1} \text{ all distinct}\}.$$

Let μ be a generic pseudocycle representing the homology class Poincaré dual to

$$c_1^{n_1} x_1^{m_1} x_2^{m_2} \lambda^\theta y^{\delta_L - (n_1 + m_1 + 2m_2 + \theta + \delta + 5)}.$$

As before, we let $v \in \hat{\gamma}$ and $w \in \pi^*TX/\hat{\gamma}$ be two fixed non zero vectors and define f_{ij} as before. We now define a section of the following bundle

$$\Psi_{\mathcal{P}A_5} : A_1^\delta \circ \overline{\mathcal{P}A}_4 \longrightarrow \mathbb{L}_{\mathcal{P}A_5} := \hat{\gamma}^{*5} \otimes \gamma_D^{*3} \otimes (\pi^*TX/\hat{\gamma})^{*4} \otimes L^{*3} \quad \text{given by} \\ \{\Psi_{\mathcal{P}A_5}([f], q_1, \dots, q_\delta, l_{q_{\delta+1}})\}(f^{\otimes 2} \otimes v^{\otimes 5} \otimes w^{\otimes 4}) := f_{02}^2 A_5^f,$$

where

$$A_5^f := f_{50} - \frac{10f_{21}f_{31}}{f_{02}} + \frac{15f_{12}f_{21}^2}{f_{02}^2}.$$

In [2, Lemma 7.1, statement 10], we prove that

$$\overline{\mathcal{P}A}_4 = \mathcal{P}A_4 \cup \overline{\mathcal{P}A}_5 \cup \overline{\mathcal{P}D}_5. \quad (23)$$

Hence, let us define

$$\mathcal{B} := \overline{A_1^\delta \circ \overline{\mathcal{P}A}_4} - A_1^\delta \circ (\mathcal{P}A_4 \cup \overline{\mathcal{P}A}_5).$$

If L is sufficiently $(2\delta + 5)$ -ample, then the section $\Psi_{\mathcal{P}A_5}$ vanishes on the points of $A_1^\delta \circ \mathcal{P}A_5$ transversally (all the points of $A_1^\delta \circ \mathcal{P}A_5$ are smooth points of the variety $A_1^\delta \circ \overline{\mathcal{P}A_4}$). Hence,

$$\langle e(\mathbb{L}_{\mathcal{P}A_5}), [\overline{A_1^\delta \circ \mathcal{P}A_4}] \cap [\mu] \rangle = N(A_1^\delta \mathcal{P}A_5, n_1, m_1, m_2, \theta) + \mathcal{C}_{\mathcal{B} \cap \mu}.$$

Let us define \mathcal{B}_0 as before, namely the component of the boundary where none of the (q_1, \dots, q_δ) points are equal to the last marked point $q_{\delta+1}$. We show in [2, Corollary 7.4], that the section $\Psi_{\mathcal{P}A_5}$ vanishes on all points of $\overline{A_1^\delta \circ \mathcal{P}D_5}$ with a multiplicity of 2. Hence, the contribution of \mathcal{B}_0 to the Euler class is

$$2N(A_1^\delta \mathcal{P}D_5, n_1, m_1, m_2, \theta).$$

Next, let us analyze the boundary when one (or more) of the q_i become equal to the last point $q_{\delta+1}$. Define $\mathcal{B}(q_{i_1}, \dots, q_{i_k}, l_{q_\delta})$ as before. We show in [5, Lemma 6.3, statement 5] that,

$$\mathcal{B}(q_1, l_{q_{\delta+1}}) \approx \overline{A_1^{\delta-1} \circ \mathcal{P}A_6} \cup \overline{A_1^{\delta-1} \circ \mathcal{P}E_6} \cup \overline{A_1^{\delta-1} \circ \mathcal{P}D_7}.$$

The contribution from $\overline{A_1^{\delta-1} \circ \mathcal{P}A_6} \cap \mu$ and $\overline{A_1^{\delta-1} \circ \mathcal{P}E_6} \cap \mu$ are 2 and 5 respectively. Since the pseudocycle μ is generic, it will not intersect $\overline{A_1^{\delta-1} \circ \mathcal{P}D_7} \cap \mu$. Hence, the total contribution from all the components of type $\mathcal{B}(q_{i_1}, l_{q_{\delta+1}})$ equals

$$2 \binom{\delta}{1} N(A_1^{\delta-1} \mathcal{P}A_6, n_1, m_1, m_2, \theta) + 5 \binom{\delta}{1} N(A_1^{\delta-1} \mathcal{P}E_6, n_1, m_1, m_2, \theta).$$

Next, we show in [1] that

$$\mathcal{B}(q_1, q_2, l_{q_{\delta+1}}) \approx \overline{A_1^{\delta-2} \circ \mathcal{P}D_7}.$$

This is geometrically believable due to the following picture:




Figure 10: Two nodes and nodes and one A_4 -singularity collapsing to a D_7 -singularity

We also show that the contribution from each of the points of $\overline{A_1^{\delta-2} \circ \mathcal{P}D_7} \cap \mu$ is 6. Hence the total contribution from all the components of type $\mathcal{B}(q_{i_1}, q_{i_2}, l_{q_{\delta+1}})$ equals

$$6 \binom{\delta}{2} N(A_1^{\delta-2} \mathcal{P}D_7, n_1, m_1, m_2, \theta).$$

Finally, we show in [1] that

$$\mathcal{B}(q_1, q_2, q_3, l_{q_4}) \cap \mu = \emptyset.$$

Hence the remaining stratum of \mathcal{B} does not contribute to the Euler class, giving us Theorem 6.11. \square

7.10 Proof of Theorem 6.14: computing $N(A_1^\delta \mathcal{P}A_6, n_1, m_1, m_2, \theta)$

Recall that in section 5, we have defined the space

$$A_1^\delta \circ \overline{\mathcal{P}A_5} := \{([f], q_1, \dots, q_\delta, l_{q_{\delta+1}}) \in \mathcal{D} \times (X)^\delta \times \mathbb{P}TX : f \text{ has a singularity of type } A_1 \text{ at } q_1, \dots, q_\delta, \\ ([f], l_{q_{\delta+1}}) \in \mathcal{P}A_5, q_1, \dots, q_{\delta+1} \text{ all distinct}\}.$$

Let μ be a generic pseudocycle representing the homology class Poincaré dual to

$$c_1^{n_1} x_1^{m_1} x_2^{m_2} \lambda^\theta y^{\delta_L - (n_1 + m_1 + 2m_2 + \theta + \delta + 6)}.$$

As before, we let $v \in \hat{\gamma}$ and $w \in \pi^*TX/\hat{\gamma}$ be two fixed non zero vectors and define f_{ij} as before. We now define a section of the following bundle

$$\Psi_{\mathcal{P}A_6} : A_1^\delta \circ \mathcal{P}A_5 \longrightarrow \mathbb{L}_{\mathcal{P}A_6} := \hat{\gamma}^{*6} \otimes \gamma_D^{*4} \otimes (\pi^*TX/\hat{\gamma})^{*6} \otimes L^{*4} \quad \text{given by} \\ \{\Psi_{\mathcal{P}A_6}([f], q_1, \dots, q_\delta, l_{q_{\delta+1}})\}(f^{\otimes 2} \otimes v^{\otimes 6} \otimes w^{\otimes 6}) := f_{02}^3 A_6^f,$$

where

$$A_6^f := f_{60} - \frac{15f_{21}f_{41}}{f_{02}} - \frac{10f_{31}^2}{f_{02}} + \frac{60f_{12}f_{21}f_{31}}{f_{02}^2} + \frac{45f_{21}^2f_{22}}{f_{02}^2} - \frac{15f_{03}f_{21}^3}{f_{02}^3} - \frac{90f_{12}^2f_{21}^2}{f_{02}^3}.$$

In [2, Lemma 7.1, statement 11], we prove that

$$\overline{\mathcal{P}A_5} = \mathcal{P}A_5 \cup \overline{\mathcal{P}A_6} \cup \overline{\mathcal{P}D_6} \cup \overline{\mathcal{P}E_6}. \quad (24)$$

Hence, let us define

$$\mathcal{B} := \overline{A_1^\delta \circ \overline{\mathcal{P}A_5}} - A_1^\delta \circ (\mathcal{P}A_5 \cup \overline{\mathcal{P}A_6}).$$

If L is sufficiently $(2\delta + 5)$ -ample, then the section $\Psi_{\mathcal{P}A_6}$ vanishes on the points of $A_1^\delta \circ \mathcal{P}A_6$ transversally (all the points of $A_1^\delta \circ \mathcal{P}A_6$ are smooth points of the variety $A_1^\delta \circ \overline{\mathcal{P}A_5}$). Hence,

$$\langle e(\mathbb{L}_{\mathcal{P}A_6}), [\overline{A_1^\delta \circ \mathcal{P}A_5}] \cap [\mu] \rangle = N(A_1^\delta \mathcal{P}A_6, n_1, m_1, m_2, \theta) + \mathcal{C}_{\mathcal{B} \cap \mu}.$$

Let us define \mathcal{B}_0 as before, namely the component of the boundary where all the points are distinct. We show in [2, Corollary 7.7 and Corollary 7.9] that the section $\Psi_{\mathcal{P}A_6}$ vanishes on all points of $\overline{A_1^\delta \circ \mathcal{P}D_6} \cap \mu$ and $\overline{A_1^\delta \circ \mathcal{P}E_6} \cap \mu$ with a multiplicity of 4 and 3 respectively. Hence, the contribution of \mathcal{B}_0 to the Euler class is

$$4N(A_1^\delta \mathcal{P}D_6, n_1, m_1, m_2, \theta) + 3N(A_1^\delta \mathcal{P}E_6, n_1, m_1, m_2, \theta).$$

Next, let us analyze the boundary when one (or more) of the q_i become equal to the last point $q_{\delta+1}$. Define $\mathcal{B}(q_{i_1}, \dots, q_{i_k}, l_{q_\delta})$ as before. We show in [5, Lemma 6.3, statement 6] and [1] that

$$\mathcal{B}(q_1, l_{q_{\delta+1}}) \approx \overline{A_1^{\delta-1} \circ \mathcal{P}A_7} \cup \overline{A_1^{\delta-1} \circ \mathcal{P}E_7} \cup \overline{A_1^{\delta-1} \circ \mathcal{P}D_8} \cup \overline{A_1^{\delta-1} \circ \hat{X}_9}.$$

We also show in [5, Corollary 6.13 and 6.23] that the contribution from the points of $\overline{A_1^{\delta-1} \circ \mathcal{P}A_7} \cap \mu$ and $\overline{A_1^{\delta-1} \circ \mathcal{P}E_7} \cap \mu$ are 2 and 6 respectively. We also show in [1] that the contribution from the points of $\overline{A_1^{\delta-1} \circ \hat{X}_9} \cap \mu$ is 12. Since the pseudocycle μ is generic, it will not intersect $\overline{A_1^{\delta-1} \circ \mathcal{P}D_8}$. Hence the total contribution from all the components of type $\mathcal{B}(q_{i_1}, l_{q_{\delta+1}})$ equals

$$2 \binom{\delta}{1} N(A_1^{\delta-1} \mathcal{P}A_7, n_1, m_1, m_2, \theta) + 6 \binom{\delta}{1} N(A_1^{\delta-1} \mathcal{P}E_7, n_1, m_1, m_2, \theta) + 12 \binom{\delta}{1} N(A_1^{\delta-1} \hat{X}_9, n_1, m_1, m_2, \theta).$$

Finally, we show in [1] that

$$\mathcal{B}(q_1, q_2, l_{q_3}) \cap \mu = \emptyset.$$

Hence the remaining stratum of \mathcal{B} does not contribute to the Euler class, giving us Theorem 6.14. \square

7.11 Proof of Theorem 6.12 and 6.13: computing $N(A_1^\delta \mathcal{P}D_6, n_1, m_1, m_2, \theta)$ and $N(A_1^\delta \mathcal{P}E_6, n_1, m_1, m_2, \theta)$

The computation of $N(A_1^\delta \mathcal{P}D_6, n_1, m_1, m_2, \theta)$ and $N(A_1^\delta \mathcal{P}E_6, n_1, m_1, m_2, \theta)$ are over the same spaces; hence it is more efficient to do these to computations simultaneously. Recall that in section 5, we have defined the following space

$$A_1^\delta \circ \overline{\mathcal{P}D_5} := \{([f], q_1, \dots, q_\delta, l_{q_{\delta+1}}) \in \mathcal{D} \times (X)^\delta \times \mathbb{P}TX : f \text{ has a singularity of type } A_1 \text{ at } q_1, \dots, q_\delta, \\ ([f], l_{q_{\delta+1}}) \in \overline{\mathcal{P}D_5}, q_1, \dots, q_{\delta+1} \text{ all distinct}\}.$$

Let μ be a generic pseudocycle representing the homology class Poincaré dual to

$$c_1^{n_1} x_1^{m_1} x_2^{m_2} \lambda^\theta y^{\delta L - (n_1 + m_1 + 2m_2 + \theta + \delta + 6)}.$$

As before, we let $v \in \hat{\gamma}$ and $w \in \pi^*TX/\hat{\gamma}$ be two fixed non zero vectors and define f_{ij} as before. We now define sections of the following two bundles:

$$\begin{aligned} \Psi_{\mathcal{P}D_6} : A_1^\delta \circ \overline{\mathcal{P}D_5} &\longrightarrow \mathbb{L}_{\mathcal{P}D_6} := \gamma_D^* \otimes \hat{\gamma}^{*4} \otimes L, && \text{given by} \\ \{\Psi_{\mathcal{P}D_6}([f], q_1, \dots, q_\delta, l_{q_{\delta+1}})\} &:= f_{40} && \text{and} \\ \Psi_{\mathcal{P}E_6} : A_1^\delta \circ \overline{\mathcal{P}D_5} &\longrightarrow \mathbb{L}_{\mathcal{P}E_6} := \hat{\gamma}^* \otimes \gamma_D^* \otimes (TX/\hat{\gamma})^{*2} \otimes L^*, && \text{given by} \\ \{\Psi_{\mathcal{P}E_6}([f], q_1, \dots, q_\delta, l_{q_{\delta+1}})\} &:= f_{12}. \end{aligned}$$

In [2, Lemma 7.1, statement 6], we prove that

$$\overline{\mathcal{P}D_5} = \mathcal{P}D_5 \cup \overline{\mathcal{P}D_6} \cup \overline{\mathcal{P}E_6}. \quad (25)$$

Hence, let us define

$$\begin{aligned} \mathcal{B} &:= \overline{A_1^\delta \circ \overline{\mathcal{P}D_5}} - A_1^\delta \circ (\mathcal{P}D_5 \cup \overline{\mathcal{P}D_6}) \quad \text{and} \\ \mathcal{B}' &:= \overline{A_1^\delta \circ \overline{\mathcal{P}D_5}} - A_1^\delta \circ (\mathcal{P}D_5 \cup \overline{\mathcal{P}E_6}) \end{aligned}$$

If L is sufficiently $(2\delta+4)$ -ample and $(2\delta+3)$ -ample, then the sections $\Psi_{\mathcal{P}D_6}$ and $\Psi_{\mathcal{P}E_6}$ vanish on the points of $A_1^\delta \circ \mathcal{P}D_6$ and $A_1^\delta \circ \mathcal{P}E_6$ transversally, respectively (all the points of $A_1^\delta \circ \mathcal{P}D_6$ and $A_1^\delta \circ \mathcal{P}E_6$ are smooth points of the variety $A_1^\delta \circ \overline{\mathcal{P}D_5}$). Hence

$$\begin{aligned} \langle e(\mathbb{L}_{\mathcal{P}D_6}), \overline{[A_1^\delta \circ \mathcal{P}D_5]} \cap [\mu] \rangle &= N(A_1^\delta \mathcal{P}D_6, n_1, m_1, m_2, \theta) + \mathcal{C}_{\mathcal{B} \cap \mu}(\Psi_{\mathcal{P}D_6}) \quad \text{and} \\ \langle e(\mathbb{L}_{\mathcal{P}E_6}), \overline{[A_1^\delta \circ \mathcal{P}D_5]} \cap [\mu] \rangle &= N(A_1^\delta \mathcal{P}E_6, n_1, m_1, m_2, \theta) + \mathcal{C}_{\mathcal{B}' \cap \mu}(\Psi_{\mathcal{P}E_6}). \end{aligned}$$

Here $\mathcal{C}_{\mathcal{B} \cap \mu}(\Psi_{\mathcal{P}D_6})$ and $\mathcal{C}_{\mathcal{B}' \cap \mu}(\Psi_{\mathcal{P}E_6})$ denote the contribution from the boundary to the Euler class from the sections $\Psi_{\mathcal{P}D_6}$ and $\Psi_{\mathcal{P}E_6}$ respectively.

Let us define \mathcal{B}_0 and \mathcal{B}'_0 as before, namely the component of the boundary where all the points are distinct. It is easy to see that the section $\Psi_{\mathcal{P}D_6}$ does not vanish on $\mathcal{B}'_0 \cap \mu$ and the section $\Psi_{\mathcal{P}E_6}$ does not vanish on $\mathcal{B}_0 \cap \mu$. Hence, \mathcal{B}_0 and \mathcal{B}'_0 do not contribute to the Euler class.

Next, let us analyze the boundary when one (or more) of the q_i become equal to the last point $q_{\delta+1}$. Define $\mathcal{B}(q_{i_1}, \dots, q_{i_k}, l_{q_\delta})$ and $\mathcal{B}'(q_{i_1}, \dots, q_{i_k}, l_{q_\delta})$ as before. We show in [5, Lemma 6.3, statement 9] and [1] that

$$\mathcal{B}(q_1, l_{q_{\delta+1}}) \approx \mathcal{B}'(q_1, l_{q_{\delta+1}}) \approx \overline{A_1^{\delta-1} \circ \mathcal{P}D_7} \cup \overline{A_1^{\delta-1} \circ \mathcal{P}E_7} \cup \overline{A_1^{\delta-1} \circ \hat{X}_9}.$$

Let us first focus on the section $\Psi_{\mathcal{P}D_6}$. We show in [5, Corollary 6.28 and Corollary 6.30], that the contribution of the section $\Psi_{\mathcal{P}D_6}$ from the points of $\overline{A_1^{\delta-1} \circ \mathcal{P}D_7} \cap \mu$ and $\overline{A_1^{\delta-1} \circ \mathcal{P}E_7} \cap \mu$ are 2 and 1 respectively. Let us now consider $\overline{A_1^{\delta-1} \circ \hat{X}_9} \cap \mu$. Suppose

$$([f], q_1, \dots, q_\delta, l_{q_{\delta+1}}) \in \overline{A_1^{\delta-1} \circ \hat{X}_9} \cap \mu.$$

Since μ is a generic pseudocycle, we conclude that $f_{40} \neq 0$. Hence, the section $\Psi_{\mathcal{P}D_6}$ does not vanish on $\overline{A_1^{\delta-1} \circ \hat{X}_9} \cap \mu$. Hence the total contribution of the section $\Psi_{\mathcal{P}D_6}$ from all the components of type $\mathcal{B}(q_{i_1}, l_{q_{\delta+1}})$ equals

$$2 \binom{\delta}{1} N(A_1^{\delta-1} \mathcal{P}D_7, n_1, m_1, m_2, \theta) + \binom{\delta}{1} N(A_1^{\delta-1} \mathcal{P}E_7, n_1, m_1, m_2, \theta).$$

Next, we show in [1] that

$$\mathcal{B}(q_1, q_2, l_{q_3}) \cap \mu = \emptyset.$$

Hence the remaining stratum of \mathcal{B} does not contribute to the Euler class, giving us Theorem 6.12.

Next, let us consider the section $\Psi_{\mathcal{P}E_6}$. We show in [5, Corollary 6.30], and in [1] that the contribution of the section $\Psi_{\mathcal{P}E_6}$ from the points of $\overline{A_1^{\delta-1} \circ \mathcal{P}E_7} \cap \mu$ and $\overline{A_1^{\delta-1} \circ \hat{X}_9} \cap \mu$ are 1 and

4 respectively. The section $\Psi_{\mathcal{P}E_6}$ does not vanish on $\overline{A_1^{\delta-1} \circ \mathcal{P}D_7}$; hence it does not vanish on $\overline{A_1^{\delta-1} \circ \overline{\mathcal{P}D_7}} \cap \mu$. Hence the total contribution from all the components of type $\mathcal{B}(q_{i_1}, l_{q_{\delta+1}})$ equals

$$1 \times \binom{\delta}{1} N(A_1^{\delta-1} \mathcal{P}E_7, n_1, m_1, m_2, \theta) + 4 \binom{\delta}{1} N(A_1^{\delta-1} \hat{X}_9, n_1, m_1, m_2, \theta).$$

Finally, as stated earlier, we prove in [1] that

$$\mathcal{B}(q_1, q_2, l_{q_3}) \cap \mu = \emptyset.$$

Hence, the remaining stratum of \mathcal{B} does not contribute to Euler class, giving us Theorem 6.13. \square

7.12 Proof of Theorem 6.15 and 6.16: computing $N(A_1^\delta \mathcal{P}D_7, n_1, m_1, m_2, \theta)$ and $N(A_1^\delta \mathcal{P}E_7, n_1, m_1, m_2, \theta)$

The computation of $N(A_1^\delta \mathcal{P}D_7, n_1, m_1, m_2, \theta)$ and $N(A_1^\delta \mathcal{P}E_7, n_1, m_1, m_2, \theta)$ are over the same spaces; hence it is more efficient to prove these two computations simultaneously. Recall that in section 5, we have defined the following space

$$A_1^\delta \circ \overline{\mathcal{P}D_6} := \{([f], q_1, \dots, q_\delta, l_{q_{\delta+1}}) \in \mathcal{D} \times (X)^\delta \times \mathbb{P}TX : f \text{ has a singularity of type } A_1 \text{ at } q_1, \dots, q_\delta, \\ ([f], l_{q_{\delta+1}}) \in \overline{\mathcal{P}D_6}, q_1, \dots, q_{\delta+1} \text{ all distinct}\}.$$

Let μ be a generic pseudocycle representing the homology class Poincaré dual to

$$c_1^{n_1} x_1^{m_1} x_2^{m_2} \lambda^\theta y^{\delta L - (n_1 + m_1 + 2m_2 + \theta + \delta + 7)}.$$

As before, we let $v \in \hat{\gamma}$ and $w \in \pi^*TX/\hat{\gamma}$ be two fixed non zero vectors and define f_{ij} as before. We now define sections of the following two bundles

$$\begin{aligned} \Psi_{\mathcal{P}D_7} : A_1^\delta \circ \overline{\mathcal{P}D_6} &\longrightarrow \mathbb{L}_{\mathcal{P}D_7} := \hat{\gamma}^{*6} \otimes \gamma_D^{*2} \otimes (\pi^*TX/\hat{\gamma})^{*2} \otimes L^{*2}, & \text{given by} \\ \{\Psi_{\mathcal{P}D_7}([f], q_1, \dots, q_\delta, l_{q_{\delta+1}})\} & (f^{\otimes 2} \otimes v^{\otimes 6} \otimes w^{\otimes 2}) := f_{12} D_7^f & \text{and} \\ \Psi_{\mathcal{P}E_7} : A_1^\delta \circ \overline{\mathcal{P}D_6} &\longrightarrow \mathbb{L}_{\mathcal{P}E_7} := \gamma_D^* \otimes \hat{\gamma}^* \otimes (\pi^*TX/\hat{\gamma})^{*2} \otimes L, & \text{given by} \\ \{\Psi_{\mathcal{P}E_7}([f], q_1, \dots, q_\delta, l_{q_{\delta+1}})\} & (f \otimes v \otimes w^{\otimes 2}) := f_{12}, \end{aligned}$$

where $D_7^f := f_{50} - \frac{5f_{31}^2}{3f_{12}}$.

In [2, Lemma 7.1, statement 7], we prove that

$$\overline{\mathcal{P}D_6} = \mathcal{P}D_6 \cup \overline{\mathcal{P}D_7} \cup \overline{\mathcal{P}E_7}. \quad (26)$$

Hence, let us define

$$\begin{aligned} \mathcal{B} &:= \overline{A_1^\delta \circ \overline{\mathcal{P}D_6}} - A_1^\delta \circ (\mathcal{P}D_6 \cup \overline{\mathcal{P}D_7}) & \text{and} \\ \mathcal{B}' &:= \overline{A_1^\delta \circ \overline{\mathcal{P}D_5}} - A_1^\delta \circ (\mathcal{P}D_6 \cup \overline{\mathcal{P}E_7}) \end{aligned}$$

If L is sufficiently $(2\delta + 5)$ -ample and $(2\delta + 4)$ -ample, then the sections $\Psi_{\mathcal{P}D_7}$ and $\Psi_{\mathcal{P}E_7}$ vanish on the points of $A_1^\delta \circ \mathcal{P}D_7$ and $A_1^\delta \circ \mathcal{P}E_7$ transversally, respectively (all the points of $A_1^\delta \circ \mathcal{P}D_7$ and $A_1^\delta \circ \mathcal{P}E_7$ are smooth points of the variety $A_1^\delta \circ \overline{\mathcal{P}D_6}$). Hence

$$\begin{aligned} \langle e(\mathbb{L}_{\mathcal{P}D_7}), \overline{[A_1^\delta \circ \mathcal{P}D_6]} \cap [\mu] \rangle &= N(A_1^\delta \mathcal{P}D_7, n_1, m_1, m_2, \theta) + \mathcal{C}_{\mathcal{B} \cap \mu}(\Psi_{\mathcal{P}D_7}) & \text{and} \\ \langle e(\mathbb{L}_{\mathcal{P}E_7}), \overline{[A_1^\delta \circ \mathcal{P}D_6]} \cap [\mu] \rangle &= N(A_1^\delta \mathcal{P}E_7, n_1, m_1, m_2, \theta) + \mathcal{C}_{\mathcal{B}' \cap \mu}(\Psi_{\mathcal{P}E_7}). \end{aligned}$$

Let us define \mathcal{B}_0 and \mathcal{B}'_0 as before, namely the component of the boundary where all the points are distinct. Note that the section $\Psi_{\mathcal{P}D_7}$ does not vanish on $\overline{A_1^\delta} \circ \mathcal{P}E_7$. To see why that is so, we note that on $\mathcal{P}E_7$, the directional derivative $f_{12} = 0$ but f_{03} and $f_{31} \neq 0$; this follows from Lemma 4.8. Hence,

$$f_{12}D_7^f = f_{12}f_{50} - \frac{5}{3}f_{31}^2 = -\frac{5}{3}f_{31}^2 \neq 0.$$

Hence, the section $\Psi_{\mathcal{P}D_7}$ does not vanish on $\mathcal{B}_0 \cap \mu$. It is also immediate that the section $\Psi_{\mathcal{P}E_7}$ does not vanish on $\overline{A_1^\delta} \circ \mathcal{P}D_7$; hence it does not vanish on $\mathcal{B}'_0 \cap \mu$. Hence, \mathcal{B}_0 and \mathcal{B}'_0 do not contribute to the Euler class.

Next, let us analyze the boundary when one (or more) of the q_i become equal to the last point $q_{\delta+1}$. Define $\mathcal{B}(q_{i_1}, \dots, q_{i_k}, l_{q_\delta})$ and $\mathcal{B}'(q_{i_1}, \dots, q_{i_k}, l_{q_\delta})$ as before. We show in [1], that

$$\mathcal{B}(q_1, l_{q_{\delta+1}}) \approx \mathcal{B}'(q_1, l_{q_{\delta+1}}) \approx \overline{A_1^{\delta-1} \circ \mathcal{P}D_8} \cup \overline{A_1^{\delta-1} \circ \mathcal{P}X_9}.$$

Let us first focus on the section $\Psi_{\mathcal{P}D_7}$. We show that the contribution of the section $\Psi_{\mathcal{P}D_7}$ from the points of $\overline{A_1^{\delta-1} \circ \mathcal{P}D_8} \cap \mu$ is 2. Next, let us consider $\overline{A_1^{\delta-1} \circ \mathcal{P}X_9} \cap \mu$. Suppose

$$([f], q_1, \dots, q_\delta, l_{q_{\delta+1}}) \in \overline{A_1^{\delta-1} \circ \mathcal{P}X_9} \cap \mu.$$

Since μ is a generic pseudocycle, we conclude that $f_{31} \neq 0$. Hence, the section $\Psi_{\mathcal{P}D_7}$ does not vanish on $\overline{A_1^{\delta-1} \circ \mathcal{P}X_9} \cap \mu$. Hence the total contribution of the section $\Psi_{\mathcal{P}D_7}$ from $\mathcal{B} \cap \mu$ is

$$2 \binom{\delta}{1} N(A_1^{\delta-1} \mathcal{P}D_8, n_1, m_1, m_2, \theta).$$

That gives us Theorem 6.15.

Next, let us look at the section $\Psi_{\mathcal{P}E_7}$. We show that the contribution of the section $\Psi_{\mathcal{P}E_7}$ from the points of $\overline{A_1^{\delta-1} \circ \mathcal{P}X_9} \cap \mu$ is 3. Furthermore, the section does not vanish on $A_1^{\delta-1} \circ \mathcal{P}D_8$ (and hence does not vanish on $A_1^{\delta-1} \circ \mathcal{P}D_8 \cap \mu$ since the cycle μ is generic). Hence, the total contribution of the section from $\mathcal{B} \cap \mu$ is

$$3 \binom{\delta}{1} N(A_1^{\delta-1} \mathcal{P}X_9, n_1, m_1, m_2, \theta).$$

That proves Theorem 6.15. □

7.13 Proof of Theorem 6.17: computing $N(A_1^\delta \mathcal{P}A_7, n_1, m_1, m_2, \theta)$

Recall that in section 5, we have defined the space

$$A_1^\delta \circ \overline{\mathcal{P}A_6} := \{([f], q_1, \dots, q_\delta, l_{q_{\delta+1}}) \in \mathcal{D} \times (X)^\delta \times \mathbb{P}TX : f \text{ has a singularity of type } A_1 \text{ at } q_1, \dots, q_\delta, \\ ([f], l_{q_{\delta+1}}) \in \overline{\mathcal{P}A_6}, q_1, \dots, q_{\delta+1} \text{ all distinct}\}.$$

Let μ be a generic pseudocycle representing the homology class Poincaré dual to

$$c_1^{n_1} x_1^{m_1} x_2^{m_2} \lambda^\theta y^{\delta_L - (n_1 + m_1 + 2m_2 + \theta + \delta + 7)}.$$

As before, we let $v \in \hat{\gamma}$ and $w \in \pi^*TX/\hat{\gamma}$ be two fixed non zero vectors and define f_{ij} as before. We now define section of the following bundle

$$\Psi_{\mathcal{P}A_7} : A_1^\delta \circ \overline{\mathcal{P}A_6} \longrightarrow \mathbb{L}_{\mathcal{P}A_7} := \hat{\gamma}^{*7} \otimes \gamma_D^{*5} \otimes (\pi^*TX/\hat{\gamma})^{*8} \otimes L^{*5}, \quad \text{given by} \\ \{\Psi_{\mathcal{P}A_7}([f], q_1, \dots, q_\delta, l_{q_{\delta+1}})\} (f^{\otimes 5} \otimes v^{\otimes 7} \otimes w^{\otimes 8}) := f_{02}^4 A_7^f.$$

In [2, Lemma 7.1, statement 12], we prove that

$$\overline{\mathcal{P}A_6} = \mathcal{P}A_6 \cup \overline{\mathcal{P}A_7} \cup \overline{\mathcal{P}D_7} \cup \overline{\mathcal{P}E_7} \cup \overline{\hat{X}_9}. \quad (27)$$

Hence, let us define

$$\mathcal{B} := \overline{A_1^\delta \circ \overline{\mathcal{P}A_6}} - A_1^\delta \circ (\mathcal{P}A_6 \cup \overline{\mathcal{P}A_7}).$$

If L is sufficiently $(2\delta + 7)$ -ample then the section vanishes on the points of $A_1^\delta \circ \mathcal{P}A_7$ transversally (all the points of $A_1^\delta \circ \mathcal{P}A_7$ are smooth points of the variety $A_1^\delta \circ \overline{\mathcal{P}A_6}$). Hence

$$\langle e(\mathbb{L}_{\mathcal{P}A_7}), \overline{[A_1^\delta \circ \mathcal{P}A_6]} \cap [\mu] \rangle = N(A_1^\delta \mathcal{P}A_7, n_1, m_1, m_2, \theta) + \mathcal{C}_{\mathcal{B} \cap \mu}. \quad (28)$$

Let us define \mathcal{B}_0 as before, namely the component of the boundary all the points are distinct. In [2, Corollary 7.11, Corollary 7.13 and Equation 7.62], we prove that the section $\Psi_{\mathcal{P}A_7}$ vanishes on $\overline{A_1^\delta \circ \mathcal{P}D_7} \cap \mu$, $\overline{A_1^\delta \circ \mathcal{P}E_7} \cap \mu$ and $\overline{A_1^\delta \circ \hat{X}_9} \cap \mu$ with a multiplicity of 6, 7 and 5 respectively. Thus, the contribution of $\mathcal{B}_0 \cap \mu$ to the Euler class is

$$6N(A_1^\delta \mathcal{P}D_7, n_1, m_1, m_2, \theta) + 7N(A_1^\delta \mathcal{P}E_7, n_1, m_1, m_2, \theta) + 5N(A_1^\delta \hat{X}_9, n_1, m_1, m_2, \theta).$$

Next, let us analyze the boundary when one (or more) of the q_i become equal to the last point $q_{\delta+1}$. We show in [1] that

$$\mathcal{B}(q_1, l_{q_{\delta+1}}) \cap \mu = \left(\overline{\mathcal{P}A_8} \cap \mu \right) \cup \left(\overline{\mathcal{P}E_8} \cap \mu \right).$$

We also show that the contribution to the Euler class from $\overline{\mathcal{P}A_8} \cap \mu$ and $\overline{\mathcal{P}E_8} \cap \mu$ is 2 and 14 respectively. Hence the total contribution from points of type $\mathcal{B}(q_{i_1}, l_{q_{\delta+1}})$ is

$$2 \binom{\delta}{1} N(A_1^\delta \mathcal{P}A_8, n_1, m_1, m_2, \theta) + 14 \binom{\delta}{1} N(A_1^\delta \mathcal{P}E_8, n_1, m_1, m_2, \theta).$$

This gives us Theorem 6.17. \square

The rest of the computations involve enumerating curves with just one codimension eight singular point. Hence, there will be no reference to a δ , since δ is going to be zero.

7.14 Proof of Theorem 6.19: computing $N(\mathcal{P}D_8, n_1, m_1, m_2, \theta)$

Let μ be a generic pseudocycle representing the homology class Poincaré dual to

$$c_1^{n_1} x_1^{m_1} x_2^{m_2} \lambda^\theta y^{\delta_L - (n_1 + m_1 + 2m_2 + \theta + 8)}.$$

As before, we let $v \in \hat{\gamma}$ and $w \in \pi^*TX/\hat{\gamma}$ be two fixed non zero vectors and define f_{ij} as before. We now define a section of the following bundles

$$\begin{aligned} \Psi_{\mathcal{P}D_8} : \overline{\mathcal{P}D_7} &\longrightarrow \mathbb{L}_{\mathcal{P}D_8} := \gamma_D^{*3} \otimes \hat{\gamma}^{*9} \otimes (\pi^*TX/\hat{\gamma})^{*6} \otimes L^{*4}, & \text{given by} \\ \{\Psi_{\mathcal{P}D_8}([f], q_1, \dots, q_\delta, l_{q_{\delta+1}})\} &(f^{\otimes 2} \otimes v^{\otimes 6} \otimes w^{\otimes 2}) := f_{12}^3 D_8^f, \end{aligned}$$

where

$$D_8^f := f_{60} + \frac{5f_{03}f_{31}f_{50}}{3f_{12}^2} - \frac{5f_{31}f_{41}}{f_{12}} - \frac{10f_{03}f_{31}^3}{3f_{12}^3} + \frac{5f_{22}f_{31}^2}{f_{12}^2}.$$

If L is sufficiently $(2\delta + 6)$ -ample, then this section is transverse to the zero set. We show in [1] that

$$\overline{\mathcal{P}D}_7 = \mathcal{P}D_7 \cup \overline{\mathcal{P}D}_8 \cup \overline{\mathcal{P}E}_8.$$

Hence, let us define

$$\mathcal{B} := \overline{\mathcal{P}D}_7 - \mathcal{P}D_7.$$

Hence

$$\langle e(\mathbb{L}_{\mathcal{P}D_8}), \overline{[A_1^\delta \circ \mathcal{P}D_7]} \cap [\mu] \rangle = N(A_1^\delta \mathcal{P}D_8, n_1, m_1, m_2, \theta) + \mathcal{C}_{\mathcal{B} \cap \mu}.$$

We also show in [1] that the contribution of $\overline{\mathcal{P}E}_8 \cap \mu$ to the Euler class is 3. That gives us Theorem 6.19. \square

7.15 Proof of Theorem 6.18 and 6.20: computing $N(\mathcal{P}E_8, n_1, m_1, m_2, \theta)$ and $N(\mathcal{P}X_9, n_1, m_1, m_2, \theta)$

Since the computation of both these numbers is done on top of the same space, it is more efficient to compute them together. Let μ be a generic pseudocycle representing the homology class Poincaré dual to

$$c_1^{n_1} x_1^{m_1} x_2^{m_2} \lambda^\theta y^{\delta_L - (n_1 + m_1 + 2m_2 + \theta + 8)}.$$

As before, we let $v \in \hat{\gamma}$ and $w \in \pi^*TX/\hat{\gamma}$ be two fixed non zero vectors and define f_{ij} as before. We now define sections of the following bundles

$$\begin{aligned} \Psi_{\mathcal{P}E_8} : \overline{\mathcal{P}E}_7 &\longrightarrow \mathbb{L}_{\mathcal{P}E_8} := \gamma_D^* \otimes \hat{\gamma}^{*3} \otimes (\pi^*TX/\hat{\gamma})^* \otimes L^* && \text{given by} \\ \{\Psi_{\mathcal{P}E_8}([f], q_1, \dots, q_\delta, l_{q_{\delta+1}})\} & (f \otimes v^{\otimes 3} \otimes w) := f_{31} && \text{and} \\ \Psi_{\mathcal{P}X_9} : \overline{\mathcal{P}E}_7 &\longrightarrow \mathbb{L}_{\mathcal{P}X_9} := \gamma_D^* \otimes (\pi^*TX/\hat{\gamma})^{*3} \otimes L^* && \text{given by} \\ \{\Psi_{\mathcal{P}X_9}([f], q_1, \dots, q_\delta, l_{q_{\delta+1}})\} & (f \otimes w^{\otimes 3}) := f_{03}. \end{aligned}$$

If L is sufficiently $(2\delta + 4)$ -ample, then these sections are transverse to the zero set. We show in [1] that

$$\overline{\mathcal{P}E}_7 = \mathcal{P}E_7 \cup \overline{\mathcal{P}E}_8 \cup \overline{\mathcal{P}X}_9. \quad (29)$$

The directional derivative f_{31} does not vanish on $\mathcal{P}X_9$ after we intersect it with μ . Similarly, the directional derivative f_{03} does not vanish on $\mathcal{P}E_8$. Hence,

$$\begin{aligned} \langle e(\mathbb{L}_{\mathcal{P}E_8}), \overline{[\mathcal{P}E}_7]} \cap [\mu] \rangle &= N(\mathcal{P}E_8, n_1, m_1, m_2, \theta) && \text{and} \\ \langle e(\mathbb{L}_{\mathcal{P}X_9}), \overline{[\mathcal{P}E}_7]} \cap [\mu] \rangle &= N(\mathcal{P}X_9, n_1, m_1, m_2, \theta). \end{aligned}$$

That gives us Theorem 6.18 and 6.20. \square

7.16 Proof of Theorem 6.21: computing $N(\mathcal{P}X_9^\vee, n_1, m_1, m_2, \theta)$

Let μ be a generic pseudocycle representing the homology class Poincaré dual to

$$c_1^{n_1} x_1^{m_1} x_2^{m_2} \lambda^\theta y^{\delta_L - (n_1 + m_1 + 2m_2 + \theta + 8)}.$$

As before, we let $v \in \hat{\gamma}$ and $w \in \pi^*TX/\hat{\gamma}$ be two fixed non zero vectors and define f_{ij} as before. We now define sections of the following bundle

$$\begin{aligned} \Psi_{\mathcal{P}X_9^\vee} : \overline{\hat{X}}_9 &\longrightarrow \mathbb{L}_{\mathcal{P}X_9^\vee} := \gamma_D^{*3} \otimes \hat{\gamma}^{*9} \otimes (\pi^*TX/\hat{\gamma})^{*3} \otimes L^{*3}, & \text{given by} \\ \{\Psi_{\mathcal{P}X_9^\vee}\}(v^{\otimes 9} \otimes w^{\otimes 3}) &:= \left(-\frac{f_{31}^3}{8} + \frac{3f_{22}f_{31}f_{40}}{16} - \frac{f_{13}f_{40}^2}{16} \right). \end{aligned}$$

We show in [1], that there is no extra boundary contribution while computing $N(\mathcal{P}X_9^\vee, n_1, m_1, m_2, \theta)$. Hence,

$$\langle e(\mathbb{L}_{\mathcal{P}X_9^\vee}), [\overline{\hat{X}}_9] \cap [\mu] \rangle = N(\mathcal{P}X_9^\vee, n_1, m_1, m_2, \theta).$$

This gives us Theorem 6.21. □

7.17 Proof of Theorem 6.22: computing $N(\mathcal{P}A_8, n_1, m_1, m_2, \theta)$

Finally, we are ready to compute $N(\mathcal{P}A_8, n_1, m_1, m_2, \theta)$. Let μ be a generic pseudocycle representing the homology class Poincaré dual to

$$c_1^{n_1} x_1^{m_1} x_2^{m_2} \lambda^\theta y^{\delta_L - (n_1 + m_1 + 2m_2 + \theta + 8)}.$$

As before, we let $v \in \hat{\gamma}$ and $w \in \pi^*TX/\hat{\gamma}$ be two fixed non zero vectors and define f_{ij} as before. We now define sections of the following bundle

$$\begin{aligned} \Psi_{\mathcal{P}A_8} : \overline{\mathcal{P}A}_7 &\longrightarrow \mathbb{L}_{\mathcal{P}A_8} := \gamma_D^{*6} \otimes \hat{\gamma}^{*8} \otimes (\pi^*TX/\hat{\gamma})^{*10} \otimes L^{*6}, & \text{given by} \\ \{\Psi_{\mathcal{P}A_8}([f], l_q)\}(f^{\otimes 6} \otimes v^8 \otimes w^{\otimes 10}) &:= f_{02}^5 A_8^f, \end{aligned}$$

where

$$\begin{aligned} A_8^f &:= f_{80} - \frac{28f_{21}f_{61}}{f_{02}} - \frac{56f_{31}f_{51}}{f_{02}} + \frac{210f_{21}^2f_{42}}{f_{02}^2} + \frac{420f_{21}f_{22}f_{41}}{f_{02}^2} - \frac{210f_{03}f_{21}^2f_{41}}{f_{02}^3} + \frac{560f_{21}f_{31}f_{32}}{f_{02}^2} - \\ &\frac{840f_{13}f_{21}^2f_{31}}{f_{02}^3} - \frac{420f_{21}^3f_{23}}{f_{02}^3} + \frac{1260f_{03}f_{21}^3f_{22}}{f_{04}^4} - \frac{35f_{41}^2}{f_{02}} + \frac{280f_{22}f_{31}^2}{f_{02}^2} - \frac{280f_{03}f_{21}f_{31}^2}{f_{02}^3} - \frac{1260f_{21}^2f_{22}^2}{f_{02}^3} + \\ &\frac{105f_{04}f_{21}^4}{f_{02}^4} - \frac{315f_{03}^2f_{21}^4}{f_{02}^5} + \frac{168f_{21}f_{51}f_{12}}{f_{02}^2} + \frac{280f_{31}f_{41}f_{12}}{f_{02}^2} - \frac{1680f_{21}^2f_{32}f_{12}}{f_{02}^3} - \frac{3360f_{21}f_{22}f_{31}f_{12}}{f_{02}^3} + \\ &\frac{2520f_{03}f_{21}^2f_{31}f_{12}}{f_{02}^4} + \frac{2520f_{13}f_{21}^3f_{12}}{f_{02}^4} - \frac{840f_{21}f_{41}f_{12}^2}{f_{02}^3} + \frac{7560f_{21}^2f_{22}f_{12}^2}{f_{02}^4} - \frac{560f_{31}^2f_{12}^2}{f_{02}^3} - \frac{5040f_{03}f_{21}^3f_{12}^2}{f_{02}^5} \\ &+ \frac{3360f_{21}f_{31}f_{12}^3}{f_{02}^4} - \frac{5040f_{21}^2f_{12}^4}{f_{02}^5}. \end{aligned}$$

We show in [1] that

$$\overline{\mathcal{P}A}_7 = \mathcal{P}A_7 \cup \overline{\mathcal{P}A}_8 \cup \overline{\mathcal{P}D}_8 \cup \overline{\mathcal{P}E}_8 \cup \overline{\mathcal{P}X}_9^\vee.$$

We also show in [1] that the contributions from $\mathcal{P}D_8 \cap \mu$, $\mathcal{P}E_8 \cap \mu$ and $\mathcal{P}X_9^\vee \cap \mu$ are 8, 16 and 6. Hence,

$$\begin{aligned} \langle e(\mathbb{L}_{\mathcal{P}A_8}), [\overline{\mathcal{P}A}_7] \cap [\mu] \rangle &= N(\mathcal{P}A_8, n_1, m_1, m_2, \theta) \\ &+ 8N(\mathcal{P}D_8, n_1, m_1, m_2, \theta) + 16N(\mathcal{P}E_8, n_1, m_1, m_2, \theta) + 6N(\mathcal{P}X_9^\vee, n_1, m_1, m_2, \theta). \end{aligned}$$

That gives us Theorem 6.22. □

8 Low degree checks in \mathbb{P}^2

8.1 Verifying the number $N(A_1^8)$

We will now subject our formulas to some low degree checks. Let start by verifying Kleiman-Piense's formula for 8-nodal curves for quintics in \mathbb{P}^2 . We need to find out the number for degree 5 curves in \mathbb{P}^2 , through 12 generic points that have 8 nodes (let us say unordered). The genus of a smooth quintic is 6; hence the curve has to break into at least three components. The possibilities are as follows: it could break into a nodal cubic through 8 points and a pair of lines through the remaining four points. The total number of such configurations are

$$\binom{12}{8} \times 12 \times \binom{4}{2} \times \binom{2}{2} \times \frac{1}{2}.$$

Note that the lines are indistinguishable; hence we divide by two. Next, the quintic could break into two conics and a line. The total number of such configurations is

$$\binom{12}{5} \times \binom{7}{5} \times \frac{1}{2}.$$

Again, we divide by two since the conics are indistinguishable. Adding up, we get that the total number is

$$\binom{12}{8} \times 12 \times \binom{4}{2} \times \binom{2}{2} \times \frac{1}{2} + \binom{12}{5} \times \binom{7}{5} \times \frac{1}{2} = 26136.$$

This is precisely the value predicted by the formula of Kleiman and Piense!

8.2 Verifying the number $N(A_1^6 A_2)$

Let us directly compute the number of quintics through 12 points in \mathbb{P}^2 that have 6 nodes and one cusp. This has to break into at least two components. The possibilities are: it could break as a cuspidal cubic going through 7 points and a conic. The total number of such configurations is

$$\binom{12}{7} \times 24 \times \binom{5}{5}.$$

Next, it could also break into a rational quartic with two nodes and one cusp through 10 points and a line through the remaining two points. The total number of such configurations is

$$\binom{12}{10} \times 2304 \times \binom{2}{2}.$$

Adding up, we get that the total number is

$$\binom{12}{7} \times 24 \times \binom{5}{5} + \binom{12}{10} \times 2304 \times \binom{2}{2} = 171072.$$

That is precisely the value predicted by our formula!

8.3 Verifying the number $N(A_1^4 D_4)$

Let us directly compute the number of quintics in \mathbb{P}^2 through 12 points that have 4 nodes and one triple point. This has to break into at least two components. The possibilities are as follows: it could break into a quartic through 10 points having a triple point and a line through the remaining two points. The total number of such configurations are

$$\binom{12}{10} \times 60 \times \binom{2}{2}.$$

Next, it could be an irreducible quartic through 11 points, with three nodes and a line through the remaining point also going through any of the three nodes. The total number of such configurations is

$$\binom{12}{11} \times 620 \times 3.$$

Next, we could put a line through any of the two points. And we could put an irreducible quartic through 10 points, that has a node on the given line. First, let us see how we can compute that last number. Using our formula for $N(A_1, n_1, m_1, m_2)$ we can compute that the number of quartics through 10 points that have a three nodes and one of them on a given line; that number is 843 (the free nodes are unordered in the final answer). Of these, we have to figure out how many are irreducible. There could be a cubic through 9 points and a line through the last point and any of the three points the cubic intersects. Or, we could place a line through two points and a cubic through the remaining eight points and the point of intersection of the two lines. The total number of reducible configurations are

$$\binom{10}{9} \times 3 + \binom{10}{2} = 75.$$

Hence, the number of irreducible cubics through 10 points with three nodes, one of them on a line is 768. Hence, the number of ways we could put a line through two points and put an irreducible quartic through 10 points, that has a node on the given line is

$$\binom{12}{2} \times 768.$$

Next, the curve could split as $3+2$. We could put a nodal cubic through 8 points and a conic through the remaining 4 points and the nodal point. That gives us

$$\binom{12}{8} \times 12.$$

We could also put a conic through 5 points and a cubic through 7 points with a node on the given conic. The total number of such configurations is

$$\binom{12}{5} \times 6 \times 2.$$

Note that the number of cubics through 7 points with a node on a given line is 6; hence the corresponding number with a node on a given conic is 12.

Finally, the curve can split as $3+1+1$. We could put a cubic through 9 points, a line through

two points and then a line through the remaining point and any one of the three points of intersection. The total such configurations is

$$\binom{12}{9} \times \binom{3}{2} \times 3.$$

Note that in this case the lines are distinguishable (one of them goes through two points, the other one goes through one point). Hence, we do not divide by two. Finally, we could put a line through two point, another line through two point and a cubic through the remaining eight points and the point of intersection. The total number of such configurations is

$$\binom{12}{2} \times \binom{10}{2} \times \frac{1}{2}.$$

Note that in this case, the two lines are indistinguishable, hence, we divide by two. Adding up all these numbers, we get that the desired number is

$$\begin{aligned} & \binom{12}{10} \times 60 \times \binom{2}{2} + \binom{12}{11} \times 620 \times 3 \\ & + \binom{12}{2} \times 768 + \binom{12}{8} \times 12 + \binom{12}{5} \times 6 \times 2 \\ & + \binom{12}{9} \times \binom{3}{2} \times 3 + \binom{12}{2} \times \binom{10}{2} \times \frac{1}{2} = 95877. \end{aligned}$$

This is precisely the number predicted by our formula!

8.4 Verifying the number $N(A_1^4 D_4, [L])$

Let us denote $N(A_1^4 D_4, [L])$ to be the number of degree d curves in \mathbb{P}^2 through the appropriate number of generic points that have 4 nodes and one triple point lying on a line. Let us verify this number directly for quintics. We will directly compute the number of quintics through 11 points that have four nodes and one triple point lying on a line.

The curve could break as $4 + 1$. It could break as quartic through 9 points with a triple point on a line and a line through the remaining two points. Total number of such configurations is

$$\binom{11}{9} \times 12.$$

Note that the number of quartics through 9 points with a triple point on a line is 12. Next, we could put an irreducible quartic through ten point with a node on the given line and put a line through the remaining eleventh point and the nodal point of the quartic that lies on the line. The total number of such configurations is

$$\binom{11}{10} \times 768.$$

Note that 768 is the number of irreducible quartics through 10 points with three nodes, one of them on a line. Next, let us compute the number of irreducible quartics through 9 points, with three nodes one of them on a given point. The total number of such quartics is 105 (which we can obtain by using our formula for $N(A_1^\delta, n_1, m_1, m_2)$). The reducible configuration is a cubic through any of the 8 points and the given special point and a line through the remaining ninth point and the given

special point. The total number of irreducible quartics through 9 point, with three nodes, one of them lying on a point is

$$105 - \binom{9}{8} = 96.$$

Hence, one final configuration that can occur is a line through two points and an irreducible quartic through nine points with a node lying on the given point of intersection of the two lines. That is

$$\binom{11}{2} \times 96.$$

Similarly, the $3 + 2$ configurations are as follows: put a conic through five points and then a nodal cubic through any one of the two points of intersection of the conic and the line. Or, put a nodal cubic through 7 points, with a node on a given line and put a conic through the remaining five points. The sum total is

$$\binom{11}{7} \times 6 + \binom{11}{5} \times 2.$$

Note that the number of cubic through seven points with a node on a given line is 6.

Finally, the quintic could break as $3 + 1 + 1$. We could put a cubic through 9 points and a line each through the remaining two connecting the the point of intersection of the cubic with the given line. The number of such configurations is

$$\binom{11}{9} \times 3 \times \binom{2}{1} \times \frac{1}{2}.$$

Note that the lines are indistinguishable here. Finally, we can also put a line through two points, a line through one more of the remaining points and connecting the point of intersection of the previous line with the given line and a cubic through the remaining eight points and the nodal point. The number of such configurations is

$$\binom{11}{2} \times \binom{9}{1}.$$

Note that the lines are distinguishable here. Adding up, we get that the desired number is

$$\begin{aligned} & \binom{11}{9} \times 12 + \binom{11}{10} \times 768 + \binom{11}{2} \times 96 + \binom{11}{7} \times 6 + \binom{11}{5} \times 2 \\ & + \binom{11}{9} \times 3 \times \binom{2}{1} \times \frac{1}{2} + \binom{11}{2} \times \binom{9}{1} = 17952. \end{aligned}$$

This is precisely the number predicted by our formula!

8.5 Verifying the number $N(A_1^4 D_4, [\text{pt}])$

Let us denote $N(A_1^4 D_4, [\text{pt}])$ to be the number of degree d curves in \mathbb{P}^2 through the appropriate number of generic points that have 4 nodes and one triple point lying on a point. Let us verify this number directly for quintics. We will directly compute the number of quintics through 10 points that have four nodes and one triple point lying on a point

The curve could break as $4 + 1$. It could break as quartic through 8 points with a triple point on a point and a line through the remaining two points. Total number of such configurations is

$$\binom{10}{8} \times 1.$$

Next, we could put an irreducible quartic through 9 points with three nodes, one of them on a given point. The total number of such configurations is

$$\binom{10}{9} \times 96.$$

Note that 96 is the number of irreducible quartics through 9 points with three nodes, one on a given point (as we calculated earlier).

Next, the curve could break as $3 + 2$. We could put a conic through four points and the given point and then put a cubic through six points with node on the given point. The total number of such configurations are

$$\binom{10}{6}.$$

Finally, the curve could break as $3 + 1 + 1$. Put a cubic through any of the 8 points and the given special point. Put a line through any one of the remaining point and the given special point. And put a line through the last point and the given special point. Total number of such configurations is

$$\binom{10}{8} \times \binom{2}{1} \times \binom{1}{1} \times \frac{1}{2}.$$

Note that the lines are indistinguishable here, hence we divide by two. Adding up we get that the desired number is

$$\binom{10}{8} + \binom{10}{9} \times 96 + \binom{10}{4} + \binom{10}{8} \times \binom{2}{1} \times \frac{1}{2} = 1260.$$

That is precisely the number predicted by our formula!

9 Low degree checks in $\mathbb{P}^1 \times \mathbb{P}^1$

In this section, we correct a few minor oversights made by Vainsencher in [43], while he makes low degree checks for his formulas for curves in $\mathbb{P}^1 \times \mathbb{P}^1$. First, let us recall that that in [20], Kontsevich and Manin obtain a formula for the number of rational degree β curves in any del-Pezzo surface through $\langle c_1(TX), [\beta] \rangle - 1$ generic points. In particular, Kontsevich-Manin's formula can be applied to $\mathbb{P}^1 \times \mathbb{P}^1$. With this in mind, let us look at a couple of low degree checks made by Vainsencher in his paper [43].

In [43, Page 17], Vainsencher, directly computes the number of irreducible curves in $\mathbb{P}^1 \times \mathbb{P}^1$ of bi-degree $(2, 5)$ passing through 13 points that have 4-nodes. The answer he obtains is 3684. However, the number of rational curves of bi-degree $(2, 5)$ through 13 points is predicted to be 3840 by Kontsevich-Manin's formula. The reason for this mismatch is a simple combinatorial oversight made by the author. The total number of curves of bi-degree $(2, 5)$ through 13-points, with 4-nodes is indeed 7038. Of these curves, we have to subtract off all the reducible curves. One could have binodal curves of type $(2, 4)$ through 12 points and a curves of type $(0, 1)$ through the remaining point. The number of bi-nodal curves of type $(2, 4)$ through 12 points is indeed 252. However, not all of them are irreducible! There is the following configuration which occurs; a curve of type $(2, 3)$ through 11 points and a curve of type $(0, 1)$ through one point. There are a total of $\binom{12}{1}$ such configurations. Hence, the total number of irreducible binodal curves of type $(2, 4)$ is $252 - 12 = 240$. The total number of such configurations occurring for the 6-nodal curves is

$$\binom{13}{1} \times 240.$$

Finally, the curve can break as $(2, 3) + (0, 1) + (0, 1)$. The two classes $(0, 1)$ are *indistinguishable*! Hence, the total number of such configurations is

$$\frac{1}{2} \times \binom{13}{11} \times \binom{2}{1} \times \binom{1}{1}.$$

Hence, the final answer is

$$7038 - \binom{13}{1} \times 240 - \frac{1}{2} \binom{13}{11} \times \binom{2}{1} \times \binom{1}{1} = 3840.$$

This is precisely the value predicted by Kontsevich-Manin's formula!

Next, we look at another low degree check made by Vainsencher. In [43, Page 18], he directly compute the number of irreducible curves of type $(3, 4)$ through 13 points that have 6 nodes. The answer he obtains is 90508. However, the number of rational curves of bi-degree $(3, 4)$ passing through 13 generic points is predicted by Kontsevich-Manin's formula to be 87544. The reason for this mismatch is because the author omitted to consider one more configuration that occurs.

The number of 6-nodal curves of class $(3, 4)$ through 13 points is indeed 122865. To get the desired number, we simply have to subtract off all the reducible configurations. First, there will be nodal curves of class $(2, 3)$ through 10 points and a curve of class $(1, 1)$ through 3 points. The total number of such configurations is

$$\binom{13}{3} \times 20.$$

Next, there can be curves of class $(2, 2)$ through 8 points and a curve of class $(1, 2)$ through 5 points. The total number of such configurations is

$$\binom{13}{8}.$$

Next, the curve can break as $(3, 2) + (0, 1) + (0, 1)$. The total number of such configurations is

$$\frac{1}{2} \times \binom{13}{11} \times \binom{2}{1}.$$

Note that we divide by 2 since the two $(0, 1)$ curves are indistinguishable! Now we warn the reader, that we will not separately consider the configuration $(2, 3) + (1, 0) + (0, 1)$; this will be part of a configuration we will soon consider. We don't consider this separately, since all the classes are different.

Next, we consider trinodal curves of class $(3, 3)$ and one curve of class $(0, 1)$. The number of trinodal curves of class $(3, 3)$ through 12 points is 1944. This can break in two possible ways; either as $(3, 2) + (0, 1)$ or $(2, 3) + (1, 0)$. Let us only subtract off those configurations, where the curve breaks as $(3, 2) + (0, 1)$. The number of total configurations with 6-nodal curves, excluding the ones of type $(3, 2) + (0, 1) + (0, 1)$ is

$$\binom{13}{12} \times (1944 - \binom{12}{11}).$$

Note that the above number does include configurations of type $(2, 3) + (1, 0) + (0, 1)$. This configuration doesn't require any special treatment, since $(1, 0)$ and $(0, 1)$ are two distinct classes.

Finally, we come to the configuration that was overlooked by Vainsencher. We can have a bi-nodal curve of type $(2, 4)$ through 12 points and a curve of class $(1, 0)$ through one point. We

have already calculated the number of irreducible bi-nodal curves of type $(2, 4)$; it is $252 - 12 = 240$. Hence, the number of configuration of type $(2, 4) + (1, 0)$ where the $(2, 4)$ is irreducible and binodal is

$$\binom{13}{12} \times 240.$$

Hence, the final answer is

$$122865 - \left(\binom{13}{8} \times 20 + \binom{13}{8} + \frac{1}{2} \times \binom{13}{11} \times \binom{2}{1} + \binom{13}{12} \times (1944 - 12) + 240 \times \binom{13}{12} \right) = 87544.$$

This is precisely the value predicted by Kontsevich-Manin's formula!

References

- [1] S. BASU AND R. MUKHERJEE, *Collision of upto seven singular points*. in preparation.
- [2] ———, *Enumeration of curves with one singular point*. available at <http://arxiv.org/abs/1308.2902>.
- [3] ———, *Enumeration of curves with singularities: Further details*. available at <https://www.sites.google.com/site/ritwik371/home>.
- [4] ———, *Enumeration of curves with upto two singular points in a general linear system*. available at <http://arxiv.org/abs/1501.01557>.
- [5] ———, *Enumeration of curves with two singular points*, Bull. Sci. Math, (2014), pp. 667–735.
- [6] ———, *Enumeration of curves with one singular point*, Journ of Geom and Phys, 104 (2016), pp. 175–203.
- [7] I. BISWAS, S. D’MELLO, R. MUKHERJEE, AND V. PINGALI, *Rational cuspidal curves on del-pezzo surfaces*. available at <http://arxiv.org/abs/1509.06300>.
- [8] I. BISWAS, R. MUKHERJEE, AND V. THAKRE, *Genus two enumerative invariants in del-pezzo surfaces with a fixed complex structure*. available at <http://arxiv.org/abs/1511.04900>.
- [9] ———, *Genus one enumerative invariants in del-pezzo surfaces with a fixed complex structure*, Comptes Rendus Mathematique, (2016).
- [10] R. BOTT AND L. W. TU, *Differential forms in algebraic topology*, vol. 82 of Graduate Texts in Mathematics, Springer-Verlag, New York, 1982.
- [11] L. CAPORASO AND J. HARRIS, *Counting plane curves of any genus*, Invent. Math., 131 (1998), pp. 345–392.
- [12] W. FULTON, *Intersection theory*, vol. 2 of Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], Springer-Verlag, Berlin, second ed., 1998.
- [13] E. IONEL, *Genus one enumerative invariants in \mathbb{P}^2* , Duke Math Jour, 94 (1998), pp. 279–324.
- [14] E. IONEL AND T. PARKER, *The Symplectic Sum Formula for Gromov-Witten invariants*, Annals of Mathematics, 159 (2004), pp. 935–1025.
- [15] M. È. KAZARIAN, *Multisingularities, cobordisms, and enumerative geometry*, Uspekhi Mat. Nauk, 58 (2003), pp. 29–88.
- [16] S. KLEIMAN, *Counting curves à la Göttsche: A celebration of algebraic geometry*, Clay Mathematics Proceedings, Amer. Math. Soc., Providence, RI, 2013, pp. 451–456.
- [17] S. KLEIMAN AND R. PIENE, *Enumerating singular curves on surfaces*, in Algebraic geometry: Hirzebruch 70 (Warsaw, 1998), vol. 241 of Contemp. Math., Amer. Math. Soc., Providence, RI, 1999, pp. 209–238.

- [18] S. KLEIMAN AND V. SHENDE, *On the Göttsche threshold: A celebration of algebraic geometry*, Clay Mathematics Proceedings, Amer. Math. Soc., Providence, RI, 2013, pp. 429–449.
- [19] J. KOCK, *Characteristic number of rational curves with cusp or prescribed triple contact*, Math. Scand., 92 (2003), pp. 223–245.
- [20] M. KONTSEVICH AND Y. MANIN, *Gromov-Witten classes, quantum cohomology, and enumerative geometry* [MR1291244 (95i:14049)], in Mirror symmetry, II, vol. 1 of AMS/IP Stud. Adv. Math., Amer. Math. Soc., Providence, RI, 1997, pp. 607–653.
- [21] M. KOOL, V. SHENDE, AND R. P. THOMAS, *A short proof of the Göttsche conjecture*, Geom. Topol., 15 (2011), pp. 397–406.
- [22] A.-K. LIU, *The algebraic proof of the universality theorem*. available at <http://arxiv.org/abs/math/0402045>.
- [23] ———, *Family blowup formula, admissible graphs and the enumeration of singular curves*, J. Differential Geom., 56 (2000), pp. 381–579.
- [24] G. MIKHALIKIN, *Enumerative tropical algebraic geometry in*, J. Amer. Math. Soc., 18 (2005), pp. 313–377.
- [25] M. MIKOSZ, P. PRAGACZ, AND A. WEBER, *Positivity of Thom polynomials ii; the lagrange singularities*, Fund. Math., 202 (2009).
- [26] R. MUKHERJEE, *Enumerative geometry via topological computations*. PhD Thesis, Stony Brook University, 2011 (available on request).
- [27] ———, *Enumeration of singular hypersurfaces on arbitrary complex manifolds*, International Journal of Mathematics, (2016). In press.
- [28] R. PANDHARIPANDE, *Counting elliptic plane curves with fixed j -invariant*, Proceedings of AMS, 125 (1997), pp. 3471–3479.
- [29] R. PANDHARIPANDE, *Intersections of \mathbb{Q} -divisors on Kontsevichs moduli space and Enumerative Geometry*, Trans. Amer. Math. Soc., 351 (1999), pp. 1481–1505.
- [30] R. PANDHARIPANDE AND L. GÖTTSCHE, *The quantum cohomology of blowup of \mathbb{P}^2 and enumerative geometry*, Jour of Diff Geometry, 48 (1998), pp. 61–90.
- [31] P. PRAGACZ AND A. WEBER, *Positivity of Schur function expansions of Thom polynomials*, Fund. Math., 195 (2007), pp. 85–95.
- [32] ———, *Thom polynomials of invariant cones, Schur functions and positivity*, Algebraic Cycles, Sheaves, Shtukas, and Moduli, Trends in Mathematics, Birkhauser, (2007), pp. 117–129.
- [33] Z. RAN, *On nodal plane curves*, Invent. Math., 86 (1986), pp. 529–534.
- [34] Z. RAN, *Enumerative geometry of singular plane curves*, Invent. Math., 97 (1989), pp. 447–465.
- [35] J. RENNEMO, *Universal polynomials for tautological integrals on Hilbert schemes*, Journal of Geometry and Topology, (2016). In press.

- [36] R. RIMANYI, *Thom polynomials, symmetries and incidences of singularities*, *Inventiones*, 143 (2001), pp. 499–521.
- [37] ———, *Multiple point formulas a new point of view*, *Pacific Jour of Mathematics*, 202 (2002), pp. 475–490.
- [38] R. RIMANYI AND A. SZUCS, *Pontryagin Thom type construction for maps with singularities*, *Topology*, 37 (1998), pp. 1177–1191.
- [39] Y. RUAN AND G. TIAN, *A mathematical theory of quantum cohomology*, *J. Differential Geom.*, 42 (1995), pp. 259–367.
- [40] M. TEHRANI AND A. ZINGER, *On Symplectic Sum Formulas in Gromov-Witten theory*. available at <http://arxiv.org/abs/1404.1898>.
- [41] Y.-J. TZENG, *A proof of the Göttsche-Yau-Zaslow formula*, *J. Differential Geom.*, 90 (2012), pp. 439–472.
- [42] Y.-J. TZENG AND J. LI, *Universal polynomials for singular curves on surfaces*, *Compos. Math.*, 150 (2014), pp. 1169–1182.
- [43] I. VAINSENCHER, *Enumeration of n -fold tangent hyperplanes to a surface*, *J. Algebraic Geom.*, 4 (1995), pp. 503–526.
- [44] R. VAKIL, *Counting curves on rational surfaces*, *Manuscripta Mathematica*, 102 (2000), pp. 53–84.
- [45] A. WEBER, *Characteristic classes in singularity theory*. Slides, available on <http://homepages.math.uic.edu/~jaca2009/notes/Weber.pdf>.
- [46] H. ZEUTHEN, *Almindelige egenskaber ved systemer af plane kurver*, *Kongelige Danske Videnskaberne Selskabs Skrifter*, 10 (1873), pp. 285–393.
- [47] A. ZINGER, *Counting plane rational curves: old and new approaches*. available at <http://arxiv.org/abs/math/0507105>.
- [48] ———, *Enumerative algebraic geometry via techniques of symplectic topology and analysis of local obstructions*. PhD Thesis, MIT, 2002 (available from the author on request).
- [49] A. ZINGER, *Enumeration of genus-two curves with a fixed complex structure in \mathbb{P}^2 and \mathbb{P}^3* , *J. Differential Geom.*, 65 (2003), pp. 341–467.
- [50] ———, *Enumeration of one-nodal rational curves in projective spaces*, *Topology*, 43 (2004), pp. 793–829.
- [51] ———, *Counting rational curves of arbitrary shape in projective spaces*, *Geom. Topol.*, 9 (2005), pp. 571–697 (electronic).
- [52] ———, *Enumeration of genus-three plane curves with a fixed complex structure*, *J. Algebraic Geom.*, 14 (2005), pp. 35–81.
- [53] ———, *Pseudocycles and integral homology*, *Transactions of AMS*, 360 (2008), pp. 2741–2765.

DEPARTMENT OF MATHEMATICS, IISER KOLKATA, WB 741246, INDIA

E-mail address : `basu.somnath@gmail.com`, `somnath.basu@iiserkol.ac.in`

SCHOOL OF MATHEMATICS, NISER, BHUBANESWAR (HBNI), ODISHA 752050, INDIA

E-mail address : `ritwik371@gmail.com`, `ritwikm@niser.ac.in`