# A Generalised Construction of Multiple Complete Complementary Codes and Asymptotically Optimal Aperiodic Quasi-Complementary Sequence Sets

Zhengchun Zhou, Member, IEEE, Fangrui Liu, Avik Ranjan Adhikary, and Pingzhi Fan, Fellow, IEEE.

Abstract—In recent years, complementary sequence sets have found many important applications in multi-carrier code-division multiple-access (MC-CDMA) systems for their good correlation properties. In this paper, we propose a construction, which can generate multiple sets of (N,N,N)- complete complementary codes (CCCs) over  $\mathbb{Z}_N$ , where N ( $N \geq 3$ ) is a positive integer of the form N=pq (p is the least prime factor of N and q is an odd integer). Interestingly, the maximum inter-set aperiodic cross-correlation magnitude of the proposed CCCs is bounded by N. When N is odd, the combination of the proposed CCCs results to a new set of sequences to obtain asymptotically optimal aperiodic quasi-complementary sequence sets (QCSSs) with parameters  $(N(p-1)^2,N,N,N)$ , having much greater set size than the set size of the asymptotically optimal and near-optimal QCSSs reported till date, with more flexible parameters.

Index Terms—Complete complementary codes (CCCs), asymptotically optimal quasi-complementary sequence set (QCSSs), maximum aperiodic cross-correlation magnitude, multi-carrier code-division multiple-access (MC-CDMA).

#### I. INTRODUCTION

PERFECT complementary sequence set (PCSS) refers to a set of two-dimensional matrices with non-trivial autocorrelations and cross-correlations sum to zero for any non-zero time-shift. Such sequence sets have wide applications due to their ideal correlation properties, such as reducing the peak-to-average power ratio (PAPR) [1], channel estimation [2], [3], RADAR waveform design [4], etc. However, PCSSs can support only a limited number of users in multi-carrier code division multiple access (MC-CDMA) systems since the set size (i.e. the number of users) must not be larger than the flock size (i.e. the number of channels) [5].

In search of sequence sets which can support a large number of users in MC-CDMA systems, Liu *et al.* [16] initially proposed low correlation zone complementary sequence set (LCZ-CSS). Further Liu *et al.* [7] generalised the concept and proposed quasi complementary sequence sets (QCSSs) in 2013, which also includes Z-complementary sequence sets (ZCSSs) [8]–[14]. A  $(K, M, N, \delta_{\max})$ - QCSS is a set of K  $M \times N$  sequence sets, where M denotes the flock size, and N denotes the sequence length.  $\delta_{\max}$  is the maximum periodic or aperiodic correlation tolerance, depending on what we are considering. When  $\delta_{\max} = 0$  and  $K \leq M$ , QCSS becomes

TABLE I: Parameters of asymptotically optimal aperiodic OCSSs.

Ref.	K	M	N	Alphabet	Parameter constraint	
[25]	r(r-1)	r	r	$\mathbb{Z}_r$	r is an odd prime.	
Th. 1 [24]	s(s+1)	s	s	$\mathbb{Z}_s$	s is power of prime.	
Th. 3 [24]	$s^2$	s	s-1	$\mathbb{Z}_s$	s is power of prime and $s \geq 3$ .	
Proposed	$N(p-1)^2$	N	N	$\mathbb{Z}_N$	N = pq, $p$ is the least prime factor of $N$ . and $q$ is odd	

PCSS. A PCSS is called a complete complementary code (CCC), when K=M [7].

In 1974, Welch [15] derived a lower bound for the correlation of sequences. Later in 2011, Liu *et al.* [16], [17] tightened the lower bound for aperiodic QCSSs when the set size  $K \geq 3M$ . Liu *et al.* [18] further tightened the lower bound for the cases of aperiodic QCSSs, when  $\frac{\lfloor \frac{\pi^2 M}{4} \rfloor}{M} < \frac{K}{M} < 3 + \frac{1}{M}$  for sufficiently large N.

In [19], Liu et al. stated the construction of aperiodic QCSSs, achieving the bounds proposed in [17], as a open problem. Since then a lot of research have been going on, both for periodic and aperiodic cases, for the construction of QCSSs with various parameters and correlation properties. However, most of the works, including [19], are related to periodic OCSS. Constructions of asymptotically optimal and nearoptimal periodic QCSS based on difference sets and almost difference sets were reported in [20] and [21], respectively. In [22] and [23], Li et al. gave a construction of periodic QCSSs based on characters over finite fields. To the best of the authors knowledge, very few works are available in the literature on the constructions of optimal (asymptotically) aperiodic QCSSs. Working on this problem, recently in 2019 Li et al. [24] proposed asymptotically optimal aperiodic QCSSs based on low-correlation CSSs. In another work of Li et al. [25], the authors proposed near-optimal QCSS based on multiple sets

The works on the construction of aperiodic QCSS reported in [24] and [25] are all based on prime or power of prime integers. The parameters of the QCSS are given in details in Table I. In search of aperiodic QCSS with more generalised parameters, we propose a new set of permutations on  $\mathbb{Z}_N$ , where N=pq (p is the least prime factor of N and q is an odd integer). Based on this new set of permutations, we generate multiple sets of (N, N, N)- CCCs. Combining the proposed CCCs, we construct  $(N(p-1)^2, N, N, N)$ - QCSS. Since, our proposed QCSS satisfies the condition  $K \geq 3M, M \geq 2$ 

Z. Zhou, F. Liu, and A. Adhikary are with School of Mathematics, Southwest Jiaotong University, Chengdu, 610031, China. E-mail: zzc@swjtu.edu.cn, 754521389@qq.com, Avik.Adhikary@ieee.org.

P. Fan is with Institute of Mobile Communications, Southwest Jiaotong University, Chengdu, 610031, China. E-mail: p.Fan@ieee.org

and  $N \geq 2$ , we consider the correlation bound proposed by Liu et~al.~[17] to check the optimality condition. Interestingly, all the proposed CCCs have an inter-set cross correlation upper bounded by N. Hence, all the proposed QCSSs are asymptotically optimal, having the optimality factor  $\rho \approx 1$  for  $p \geq 5$ . The proposed QCSS is near-optimal when p=3, with very low value of optimality factor  $\rho \ (\approx 1.53)$ . Also, our construction is more generalised as compared to [24], since N>2 can be a prime number also. Till date, there is no systematic construction of asymptotically optimal aperiodic QCSSs over  $\mathbb{Z}_N$ , when N is not a prime or a power of prime integer.

The rest of this paper is organized as follows. In Section II, we introduce some correlation bounds of QCSS. In Section III, we propose a new set of permutations, a new construction of multiple sets of CCCs based on the new set of permutations, also prove that all the sequence sets are cyclically distinct. In Section IV, we obtain asymptotically optimal aperiodic QCSS by combining these CCCs into a new set. In section V, we make a comparison our work with the existing works in the literature. We conclude our work in Section VI.

### II. BOUNDS OF QCSS

In this section we recall some correlation bounds of QCSS. Before that let us define some notations and definitions, which will be used throughout the paper. N  $(N \geq 3)$  be a positive integer such that N=pq, where p is the least prime factor of N and q is an odd integer.  $\mathbb{Z}_N$  denotes the ring of integers modulo N.  $\omega_N=e^{\frac{2\pi i}{N}}$  be a primitive N-th root of unity.  $\mathbb{Z}_N^*=\{r:1\leq r< N,\ \gcd(r,N)=1\}.$   $\bigcup_{i=0}^{n-1}C_i=\{C_0,\ldots,C_{n-1}\}$  denotes the combination of n set of matrices  $C_i$ .

Definition 1: Let  $u=(u_0,u_1,\cdots,u_{N-1})$  and  $v=(v_0,v_1,\cdots,v_{N-1})$  denote two complex-valued sequences with length N. The aperiodic correlation function between u and v is defined as

$$\tilde{R}_{u,v}(\tau) = \begin{cases} \sum_{\substack{t=0 \\ t=0}}^{N-1-\tau} u_t v_{t+\tau}^*, & 0 \le \tau \le N-1 \\ \sum_{t=0}^{N-1-\tau} u_{t-\tau} v_{\tau}^*, & -N+1 \le \tau \le 0 \end{cases}$$
 (1)

where  $x^*$  denotes the complex conjugation of a complex-valued sequence x, and the addition  $t+\tau$  is performed modulo N.

Definition 2: Let  $C = \{C^0, C^1, \dots, C^{K-1}\}$  be a set of K two-dimensional matrices, each of size  $M \times N$ , i.e.,

$$C^{k} = \begin{bmatrix} C_{0}^{k} \\ C_{1}^{k} \\ \vdots \\ C_{M-1}^{k} \end{bmatrix}, 0 \le k \le K - 1, \tag{2}$$

where the m-th row  $C_m^k = \left(c_{m,0}^k, c_{m,1}^k, \cdots, c_{m,N-1}^k\right)$  denotes the m-th length-N constituent sequence,  $0 \leq m \leq M-1$ . The sequence set C is called a quasi-complementary sequence set (QCSS) if for any  $C^{k_1}, C^{k_2} \in C, \ 0 \leq k_1, k_2 \leq K-1, \ 0 \leq \tau \leq N-1, k_1 \neq k_2$  or  $0 < \tau \leq N-1, k_1 = k_2$ ,

$$|\tilde{R}_{C^{k_1},C^{k_2}}(\tau)| = \left| \sum_{m=0}^{M-1} \tilde{R}_{C_m^{k_1},C_m^{k_2}}(\tau) \right| \le \delta_{\max}, \quad (3)$$

where K is the set size, M is the size of each sequence set, N is the length of constituent sequences, and  $\delta_{\max}$  is the maximum aperiodic cross-correlation magnitude. The set C can be denoted by  $(K, M, N, \delta_{\max})$ -QCSS. Specially, we call C a complete complementary code (CCC) if K=M and  $\delta_{\max}=0$ .

The following lemma gives a lower bound of  $\delta_{\rm max}$ .

Lemma 1: [15] For an aperiodic quasi-complementary sequence set  $(K, M, N, \delta_{\text{max}})$ -QCSS, the parameters satisfy:

$$\delta_{\max} \ge MN \cdot \sqrt{\frac{\left(\frac{K}{M} - 1\right)}{K(2N - 1) - 1}}.$$
 (4)

For  $K \ge 3M$ ,  $M \ge 2$  and  $N \ge 2$ , Liu *et al.* [17] proposed a tighter correlation bound for aperiodic QCSS, which is given in the following lemma.

Lemma 2: [17] For an aperiodic quasi-complementary sequence set  $(K, M, N, \delta_{\text{max}})$ -QCSS with  $K \geq 3M, M \geq 2$  and  $N \geq 2$ , we have

$$\delta_{\text{max}} \ge \sqrt{MN \left(1 - 2\sqrt{\frac{M}{3K}}\right)}$$
(5)

In this paper, a QCSS is optimal if  $\delta_{max}$  achieves the lower bound in (5). Define the optimality factor  $\rho$  as follows

$$\rho = \frac{\delta_{\text{max}}}{\sqrt{MN\left(1 - 2\sqrt{\frac{M}{3K}}\right)}}.$$
 (6)

Note that  $\rho \geq 1$ . A QCSS is optimal when  $\rho = 1$  and near-optimal if  $1 < \rho \leq 2$ .

### III. MULTIPLE CCCs WITH LOW MAXIMUM INTER-SET APERIODIC CROSS-CORRELATION MAGNITUDE

In this section, first we derive a set of conditions for a set of permutations over  $\mathbb{Z}_N$ . Then we propose a construction of multiple CCCs with low maximum inter-set aperiodic cross-correlation magnitude, based on that permutations.

Construction 1: Let N = pq as defined. Also, let  $\Pi = \{\pi_a | 1 \le a < p\}$  be a set of permutations of  $\mathbb{Z}_N$ , such that the equation  $\pi_{a_1}(x+b) = c\pi_{a_2}(x) \pmod{N}, \ 1 \le a_1 \ne a_2 < p$  has only one solution for any given  $b \in \mathbb{Z}_N$  and  $c \in \mathbb{Z}_N^*$ . Then for each  $0 \le m, s < N$  and  $k \in \mathbb{Z}_N^*$ , let

$$f_s^{(a,k,m)}(t) = ks \cdot \pi_a(t) + mt \pmod{N}, \text{ for } 0 \le t < N.$$
 (7)

For each pair of (a, k), define a sequence set

$$C^{(a,k)} = \{C^{(a,k,0)}, C^{(a,k,1)}, \cdots, C^{(a,k,N-1)}\},$$
(8)

where

$$C^{(a,k,m)} = \begin{bmatrix} C_1^{(a,k,m)} \\ C_2^{(a,k,m)} \\ \vdots \\ C_N^{(a,k,m)} \end{bmatrix} = \begin{bmatrix} C_{(1,1)}^{(a,k,m)}, C_{(1,2)}^{(a,k,m)}, \cdots, C_{(1,N)}^{(a,k,m)} \\ C_{(2,1)}^{(a,k,m)}, C_{(2,2)}^{(a,k,m)}, \cdots, C_{(2,N)}^{(a,k,m)} \\ \vdots \\ C_{(N,1)}^{(k,m)}, C_{(N,2)}^{(a,k,m)}, \cdots, C_{(N,N)}^{(a,k,m)} \end{bmatrix}$$

(3) and 
$$C_{(s,t)}^{(a,k,m)} = \omega_N^{f_s^{(a,k,m)}(t)}$$

Let  $t = t_1 + t_2 \cdot q$  where  $0 \le t_1 < q$  and  $0 \le t_2 < p$  and  $\tau = \tau_1 + \tau_2 \cdot q$  where  $0 \le \tau_1 < q$  and  $0 \le \tau_2 < p$ . Define

$$\delta_{t_1,\tau_1} = \begin{cases} 0, & t_1 + \tau_1 < q, \\ 1, & t_1 + \tau_1 \ge q. \end{cases}$$
(9)

This notations will be used in the rest of the paper.

In the following lemma, we will check if the sequences in the sequence set  $C^{(a,k)}$ , defined in (8) are cyclically distinct or not.

Lemma 3: For  $0 \le s < N$ , all sequences in  $C^{(a,k,m)}$  are cyclically distinct.

*Proof:* There are two parts to prove. Let us prove the first part.

Assume that for  $0 \le s < N$ , all sequences in  $C^{(a,k,m)}$  are not cyclically distinct. Then from (7), we have for  $s_1 \ne s_2$  and  $0 \le \tau < N$ 

$$ks_1 \cdot \pi_a(t) + mt \equiv ks_2 \cdot \pi_a(t+\tau) + m(t+\tau) \pmod{N}$$
. (10)

Since  $\pi_a(t)$  is a permutation on  $\mathbb{Z}_N$ , so  $\pi_a(t) \neq \pi_a(t+\tau)$ . Hence, the above equation is impossible, and the proof follows.

Remark 1: For two different sequence sets  $C^{(a_1,k_1,m)}$  and  $C^{(a_2,k_2,m)}$  the sequences are not always cyclically distinct.

Therefore, to make the sequences cyclically distinct, we need to modify (7). Let  $\mathcal{A}$  be the set of all possible permutations of a and k. Also, let  $\mathcal{B}$  be the set of all possible combinations of a and k, excluding the cases when a=k,  $\mathcal{C}=\{(a,k): a=k=i, 1\leq i\leq (p-1)/2\}$  and  $\mathcal{D}=\mathcal{B}\cup\mathcal{C}$ . Modify the family of functions  $f_s^{(a,k,m)}: \mathbb{Z}_N \to \mathbb{Z}_N$  defined in (7), as follows

$$f_s^{(a,k,m)}(t)$$

$$= \begin{cases} ks \cdot \pi_a(t) + mt \pmod{N} & \text{if } (a,k) \in \mathcal{D}, \\ N - [ks \cdot \pi_a(t) + mt] \pmod{N} & \text{if } (a,k) \in \mathcal{A} \setminus \mathcal{D}. \end{cases}$$
(11)

Theorem 1: Let  $C^{(a,k)}$ ,  $1 \leq a,k < p$ , be the multiple sequence sets generated by Construction 1 using the modified function  $f_s^{(a,k,m)}: \mathbb{Z}_N \to \mathbb{Z}_N$ , as defined in (11). Then,

- 1) Each sequence set  $C^{(a,k)}$  is a (N, N, N)-CCC.
- 2) For any  $a_1, \ a_2$  (where  $1 \le a_1, a_2 < p$ ),  $|\sum_{s=1}^N \tilde{R}_{C_s^{(a_1,k_1,m_1)}, C_s^{(a_2,k_2,m_2)}}(\tau)| \le N \text{ for all } 0 \le \tau \le N-1, k_1 \ne k_2 \in \mathbb{Z}_p^* \text{ and } 1 \le m_1, m_2 \le N.$

*Proof:* Please see Appendix A.

Lemma 5 presents a class of permutations which satisfy the condition of  $\Pi$  in construction 1. But we need the following lemma to prove Lemma 5.

Lemma 4: [26] Let  $\xi(x) = x^e$ , where e is a positive integer with  $\gcd(r-1,e) = 1$ . Then  $\xi$  is a permutation on  $\mathbb{Z}_r$ , where r is any odd prime, such that the equation  $\xi(x+a) \equiv c\xi(x) \pmod{r}$  has only one solution in  $\mathbb{Z}_r$  for any  $a \in \mathbb{Z}_r$  and any  $c \in \mathbb{Z}_r \setminus \{0,1\}$ .

Lemma 5: Let us define a set of functions on  $\mathbb{Z}_N$  as

$$\Pi = \{ \pi_a | \pi_a(t) = qh(at_2) + g(at_1), 1 \le a \le p - 1 \}$$
 (12)

where g is a permutation over  $\mathbb{Z}_q$  and h is a permutation over  $\mathbb{Z}_p$  given by g(x) = x and h(x) = x. Then,  $\Pi$  is a set of

permutations such that  $\pi_{a_1}(t+\tau) \equiv c\pi_{a_2}(t) \pmod{N}$  has only one solution for any given  $\tau \in \mathbb{Z}_N$  and  $1 \leq a_1 \neq a_2 \leq p-1, c \in \mathbb{Z}_N \setminus \{0\}$ .

*Proof:* For  $1 \le a_1 \ne a_2 \le p-1$ ,

$$\pi_{a_1}(t+\tau) - c\pi_{a_2}(t)$$

$$= q \cdot h(a_1(t_2 + \tau_2 + \delta_{t_1,\tau_1})) + g(a_1(t_1 + \tau_1))$$

$$- qc \cdot h(a_2t_2) - cg(a_2t_1) \qquad (13)$$

$$= q \cdot (h(a_1(t_2 + \tau_2 + \delta_{t_1,\tau_1})) - ch(a_2t_2))$$

$$+ (g(a_1(t_1 + \tau_1)) - cg(a_2t_1))$$

It obvious that  $\pi_{a_1}(t+\tau)-c\pi_{a_2}(t)=0$  if and only if  $h(a_1(t_2+\tau_2+\delta_{t_1,\tau_1}))-ch(a_2t_2)=0$  and  $g(a_1(t_1+\tau_1))-cg(a_2t_1)=0$ . For  $a_1=a_2,\ c=1,\ \pi_{a_1}(t+\tau)-\pi_{a_2}(t)=0$  if and only if  $\tau_1=0$  and  $\tau_2=0$ , since g and h are permutations over  $\mathbb{Z}_q$  and  $\mathbb{Z}_p$ , respectively. Therefore,  $\pi_a(t)$  is a permutation over  $\mathbb{Z}_N$  for each  $1\leq a\leq p-1$ . For  $a_1\neq a_2,\ g(a_1(t_1+\tau_1))-cg(a_2t_1)=0$  has only one solution for  $\tau_1\in\mathbb{Z}_q$  according to Lemma 4. Similarly,  $h(a_1(t_2+\tau_2+\delta_{t_2,\tau_2}))-h(a_2t_2)=0$  has only one solution for  $\tau_2\in\mathbb{Z}_p$ . Therefore,  $\pi_{a_1}(t+\tau)\equiv\pi_{a_2}(t)$  (mod N) has only one solution for any given  $\tau\in\mathbb{Z}_N$  and  $1\leq a_1\neq a_2\leq p-1$ .

From Theorem 1 and Lemma 5 we have the following corollary.

Corollary 1: Let  $\Pi$  be a family of permutations of  $\mathbb{Z}_N$  defined by

$$\Pi = \{ \pi_a | \pi_a(t) = qh(at_2) + g(at_1), 1 \le a \le p - 1 \}$$
 (14)

where g is a permutation over  $\mathbb{Z}_q$  and h is a permutation over  $\mathbb{Z}_p$ . Then, the multiple sequence sets  $C^{(a,k)}$ ,  $1 \leq a,k < p$ , generated by Construction 1 using the modified function  $f_s^{(a,k,m)}: \mathbb{Z}_N \to \mathbb{Z}_N$ , as defined in (11) have the following properties

- 1) Each sequence set  $C^{(a,k)}$  is a (N, N, N)-CCC.
- 2) For any  $a_1$ ,  $a_2$  (where  $1 \le a_1, a_2 < p$ ),  $|\sum_{s=1}^{N} \tilde{R}_{C_s^{(a_1,k_1,m_1)}, C_s^{(a_2,k_2,m_2)}}(\tau)| \le N$  for all  $0 \le \tau \le N-1, k_1 \ne k_2 \in \mathbb{Z}_p^*$  and  $1 \le m_1, m_2 \le N$ .

Example 1: Let N=11 with p=11 and q=1. Let h(t)=t and g(t)=t be a permutation over  $\mathbb{Z}_3$ . Then  $\Pi=\{\pi_a|\pi_a(t)=3at_2+at_1,1\leq a\leq 2\}$ . By Construction 1, we get hundred (11,11,11)-CCCs,  $C^{(1,1)},C^{(1,2)},\ldots,C^{(10,9)},C^{(10,10)}$ . The first three complementary sets of first two CCCs are shown in Table II.

In the following section, we propose a construction of asymptotically optimal aperiodic QCSSs.

## IV. PROPOSED CONSTRUCTION OF ASYMPTOTICALLY OPTIMAL QCSSs

In this section, from multiple CCCs with low maximum aperiodic cross-correlation magnitude, we will obtain an aperiodic near-optimal QCSS by combining these complementary sequence sets together, as shown in the following theorem.

Theorem 2: Let  $C = C^{(1,1)} \cup C^{(1,2)} \cup \cdots \cup C^{(p-1,p-1)}$ , then the sequence set C is an asymptotically optimal aperiodic  $(N(p-1)^2, N, N, N)$ -QCSS.

TABLE II: The first three complementary sets of first two (11, 11, 11)- CCCs over  $\mathbb{Z}_{11}$  in Example 1.

For $a = 1, k = 1$						
$C^{(1,1,0)}$	$C^{(1,1,1)}$	$C^{(1,1,2)}$				
000000000000	0 1 2 3 4 5 6 7 8 9 10	0 2 4 6 8 10 1 3 5 7 9				
0 1 2 3 4 5 6 7 8 9 10	0 2 4 6 8 10 1 3 5 7 9	0 3 6 9 1 4 7 10 2 5 8				
0 2 4 6 8 10 1 3 5 7 9	0 3 6 9 1 4 7 10 2 5 8	0 4 8 1 5 9 2 6 10 3 7				
0 3 6 9 1 4 7 10 2 5 8	0 4 8 1 5 9 2 6 10 3 7	0 5 10 4 9 3 8 2 7 1 6				
0 4 8 1 5 9 2 6 10 3 7	0 5 10 4 9 3 8 2 7 1 6	0 6 1 7 2 8 3 9 4 10 5				
051049382716	0 6 1 7 2 8 3 9 4 10 5	073106295184				
061728394105	073106295184	0 8 5 2 10 7 4 1 9 6 3				
073106295184	0 8 5 2 10 7 4 1 9 6 3	0 9 7 5 3 1 10 8 6 4 2				
0 8 5 2 10 7 4 1 9 6 3	0 9 7 5 3 1 10 8 6 4 2	0 10 9 8 7 6 5 4 3 2 1				
0 9 7 5 3 1 10 8 6 4 2	0 10 9 8 7 6 5 4 3 2 1	000000000000				
0 10 9 8 7 6 5 4 3 2 1	$0\; 0\; 0\; 0\; 0\; 0\; 0\; 0\; 0\; 0\; 0$	0 1 2 3 4 5 6 7 8 9 10				
	For $a = 1, k = 2$					
$C^{(1,2,0)}$	$C^{(1,2,1)}$	$C^{(1,2,2)}$				
000000000000	0 1 2 3 4 5 6 7 8 9 10	0 2 4 6 8 10 1 3 5 7 9				
0 2 4 6 8 10 1 3 5 7 9	0 3 6 9 1 4 7 10 2 5 8	0 4 8 1 5 9 2 6 10 3 7				
0 4 8 1 5 9 2 6 10 3 7	0 5 10 4 9 3 8 2 7 1 6	0 6 1 7 2 8 3 9 4 10 5				
061728394105	073106295184	0 8 5 2 10 7 4 1 9 6 3				
0 8 5 2 10 7 4 1 9 6 3	0 9 7 5 3 1 10 8 6 4 2	0 10 9 8 7 6 5 4 3 2 1				
0 10 9 8 7 6 5 4 3 2 1	$0\; 0\; 0\; 0\; 0\; 0\; 0\; 0\; 0\; 0\; 0$	0 1 2 3 4 5 6 7 8 9 10				
0 1 2 3 4 5 6 7 8 9 10	0 2 4 6 8 10 1 3 5 7 9	0 3 6 9 1 4 7 10 2 5 8				
0 3 6 9 1 4 7 10 2 5 8	0 4 8 1 5 9 2 6 10 3 7	0 5 10 4 9 3 8 2 7 1 6				
051049382716	0 6 1 7 2 8 3 9 4 10 5	073106295184				
073106295184	0 8 5 2 10 7 4 1 9 6 3	0 9 7 5 3 1 10 8 6 4 2				
0 9 7 5 3 1 10 8 6 4 2	0 10 9 8 7 6 5 4 3 2 1	000000000000				

*Proof:* According to Theorem 1, each sequence set  $C^{(a,k)}$ , is a (N,N,N)-CCC for each  $k\in S$ , and the maximum aperiodic cross-correlation amplitude is N. Then C is an aperiodic  $(K,M,N,\delta_{\max})$ -QCSS, where  $K=N(p-1)^2,\ M=N,\ N=N,\ \delta_{\max}=max\{0,N\}=N.$ 

The optimality factor of  $(N(p-1)^2, N, N, N)$ -QCSS is

$$\rho = \frac{N}{\sqrt{N^2(1 - 2\sqrt{\frac{N}{3N(p-1)^2}})}}$$

$$= \frac{1}{\sqrt{1 - \frac{2}{\sqrt{3}(p-1)}}}.$$
(15)

Obviously,  $\rho \approx 1$  for  $p \geq 5$ . Then C is an asymptotically optimal aperiodic QCSS. Hence, this theorem follows.

Example 2: The hundred (11,11,11)-CCCs generated in Example 1, has a inter-set aperiodic cross-correlation magnitude, upper-bounded by 11. Therefore,  $\bigcup\limits_{i,j=1}^{10}C^{(i,j)}$  forms an asymptotically optimal (1100,11,11,11)- QCSS, with optimality factor  $\rho=1.0633$ .

Based on Theorem 2, a list of parameters of asymptotically oprimal aperiodic QCSSs are given in Table III. In the following section, we compare our proposed constructions with the previous works.

#### V. COMPARISON WITH PREVIOUS WORKS

We compare the optimality factor for the asymptotically optimal QCSS with the results reported in [24]. We have




Fig. 1: A glimpse of correlation magnitudes of the QCSS in *Example 2*.

taken the minimum of all the optimality factors we can get from the previous works reported in [24], for each alphabet size, whenever it is possible, and denoted it by  $\rho_{\min}$  for the comparison. Our construction is different from [cite] in following ways

- 1) The constructions of asymptotically optimal QCSSs reported in [24] are based on the construction of low correlation CSS, whereas our construction is based on multiple sets of CCCs.
- 2) Our proposed construction can generate asymptotically optimal QCSS with parameters which are not covered by the results proposed in [24]. For example, the asymptotically optimal QCSS, given in Table III, can only be constructed by our construction.
- 3) Although for the case when N is a prime number, previous constructions have reported asymptotically optimal aperiodic QCSS, compared to the previous constructions, our proposed construction can generate QCSS with larger set size and lower optimality factor. In Table IV we have listed down the parameters of the asymptotically optimal QCSS, where the alphabet size  $11 \leq p < 100$  is odd prime, as example, to show that our proposed QCSS, have a lower optimality factor compared to the previously reported results.
- 4) To have an unbiased comparison, we have also compared the cases when N is a power of prime. Our proposed results can generate QCSS having almost same optimality factor to that of the QCSS generated by the previous result reported in [24]. In Table V we have listed down some cases, as example.

#### VI. CONCLUSION

In this paper, we have proposed a new set of permutations on  $\mathbb{Z}_N$ , where N=pq, p is the least prime factor of N and q is

TABLE III: The parameters of asymptotically optimal aperiodic QCSSs.

Alphabet	K	M	N	ρ
$\mathbb{Z}_{11*13}$	14300	143	143	1.0633
$\mathbb{Z}_{13*17}$	31824	221	221	1.0519
$\mathbb{Z}_{17*19}$	82688	323	323	1.0382
$\mathbb{Z}_{19*23}$	141588	437	437	1.0337
$\mathbb{Z}_{23*31}$	345092	713	713	1.0273

TABLE IV: The comparison when N=r, r is prime.

Alphabet	K	M	N	ρ	$\rho_{min}$
$\mathbb{Z}_{11}$	1100	11	11	1.0633	1.2247
$\mathbb{Z}_{13}$	1872	13	13	1.0519	1.2026
$\mathbb{Z}_{17}$	4352	17	17	1.0382	1.1722
$\mathbb{Z}_{19}$	6156	19	19	1.0337	1.1611
$\mathbb{Z}_{23}$	11132	23	23	1.0273	1.1438
$\mathbb{Z}_{29}$	22736	29	29	1.0213	1.1257
$\mathbb{Z}_{31}$	27900	31	31	1.0198	1.1209
$\mathbb{Z}_{37}$	47952	37	37	1.0164	1.1093
$\mathbb{Z}_{41}$	65600	41	41	1.0148	1.1031
$\mathbb{Z}_{43}$	75852	43	43	1.0140	1.1003
$\mathbb{Z}_{47}$	99452	47	47	1.0128	1.0954
$\mathbb{Z}_{53}$	143312	53	53	1.0113	1.0892
$\mathbb{Z}_{59}$	198476	59	59	1.0101	1.0841
$\mathbb{Z}_{61}$	219600	61	61	1.0098	1.0825
$\mathbb{Z}_{67}$	291852	67	67	1.0089	1.0783
$\mathbb{Z}_{71}$	347900	71	71	1.0084	1.0759
$\mathbb{Z}_{73}$	378432	73	73	1.0081	1.0747
$\mathbb{Z}_{79}$	480636	79	79	1.0075	1.0716
$\mathbb{Z}_{83}$	558092	83	83	1.0071	1.0696
$\mathbb{Z}_{89}$	689216	89	89	1.0066	1.0670
$\mathbb{Z}_{97}$	893952	97	97	1.0061	1.0640

TABLE V: The comparison when  $N = r^2$ , r is prime.

Alphabet	K	M	N	ρ	$ ho_{min}$
$\mathbb{Z}_{23*23}$	256036	529	529	1.0273	1.0271
$\mathbb{Z}_{29*29}$	659344	841	841	1.0213	1.0211
$\mathbb{Z}_{31*31}$	864900	961	961	1.0198	1.0197
$\mathbb{Z}_{37*37}$	1774224	1369	1369	1.0164	1.0164
$\mathbb{Z}_{41*41}$	2689600	1681	1681	1.0148	1.0147
$\mathbb{Z}_{43*43}$	3261636	1849	1849	1.0140	1.0140
$\mathbb{Z}_{47*47}$	4674244	2209	2209	1.0128	1.0127

an odd integer. Using this permutation we first proposed a construction of  $(p-1)^2$  sets of (N,N,N)-CCCs. Then combined these  $(p-1)^2$  sets of (N,N,N)-CCCs to construct aperiodic  $(N(p-1)^2,N,N,N)$ -QCSS. All the proposed QCSS are asymptotically optimal for  $p \geq 5$ , since the optimality factor  $\rho \approx 1$ . The proposed asymptotically optimal QCSSs have paprameters which have not been reported before. Also for the alphabets over which asymptotically optimal QCSS have been reported in previous works, our construction generates QCSS having much larger set size, more flexible parameters and lower optimality factor as compared to the previous works. An interesting future work will be to design QCSSs with flexible parameters having optimality factor equal to one.

#### APPENDIX A

#### PROOF OF THEOREM 1

Let us prove the first part. Let  $C^{(a,k,m_1)},C^{(a,k,m_2)}\in C^{(a,k)}$ , where  $0\leq m_1,m_2\leq N-1$  and  $(a,k)\in \mathcal{D}$ . Then the aperiodic correlation of  $C^{(a,k,m_1)}$  and  $C^{(a,k,m_2)}$  is

$$\begin{split} \sum_{s=0}^{N-1} \tilde{R}_{C_s^{(a,k,m_1)},C_s^{(a,k,m_2)}}(\tau) \\ &= \sum_{s=0}^{N-1} \sum_{t=0}^{N-1-\tau} c_{s,t}^{(a,k,m_1)} \cdot \left( c_{s,t+\tau}^{(a,k,m_2)} \right)^* \\ &= \sum_{s=0}^{N-1} \sum_{t=0}^{N-1-\tau} \omega_N^{f_s^{(a,k,m_1)}(t)} \cdot \omega_N^{-f_s^{(a,k,m_2)}(t+\tau)} \\ &= \sum_{s=0}^{N-1} \sum_{t=0}^{N-1-\tau} \omega_N^{ks(\pi_a(t)-\pi_a(t+\tau))+t(m_1-m_2)-m_2\tau}. \end{split}$$

We have the following four cases to consider.

Case 1: When  $m_1 = m_2, \tau = 0$ . We have

$$\sum_{s=0}^{N-1} \tilde{R}_{C_s^{(a,k,m_1)},C_s^{(a,k,m_2)}}(0) = N^2.$$
 (16)

Case 2: When  $m_1 = m_2, 1 \le \tau \le N - 1$ .

$$\begin{split} \sum_{s=0}^{N-1} \tilde{R}_{C_s^{(a,k,m_1)},C_s^{(a,k,m_2)}}(\tau) \\ &= \sum_{s=0}^{N-1} \sum_{t=0}^{N-1-\tau} \omega_N^{-m_2\tau} \cdot \omega_N^{ks(\pi_a(t)-\pi_a(t+\tau))} \\ &= \omega_N^{-m_2\tau} \cdot \sum_{t=0}^{N-1-\tau} \sum_{s=0}^{N-1} \omega_N^{ks(\pi_a(t)-\pi_a(t+\tau))} = 0. \end{split}$$

Since  $\pi_a(t)$  is a permutation on  $\mathbb{Z}_N$  and  $\tau \neq 0$ ,  $\pi_a(t) \neq \pi_a(t+\tau)$ , therefore the last identity holds.

Case 3: When  $m_1 \neq m_2, \tau = 0$ .

$$\sum_{s=0}^{N-1} \tilde{R}_{C_s^{(a,k,m_1)},C_s^{(a,k,m_2)}}(0) = \sum_{s=0}^{N-1} \sum_{t=0}^{N-1} \omega_N^{t(m_1-m_2)} = 0.$$
(17)

Case 4: When  $m_1 \neq m_2, 1 \leq \tau \leq N-1$ .

$$\sum_{s=0}^{N-1} \tilde{R}_{C_s^{(a,k,m_1)},C_s^{(a,k,m_2)}}(\tau)$$

$$= \sum_{t=0}^{N-1-\tau} \omega_N^{t\cdot(m_1-m_2)-m_2\tau} \cdot \sum_{s=0}^{N-1} \omega_N^{k(\pi_a(t)-\pi_a(t+\tau))\cdot s} = 0.$$
(18)

The four cases above illustrate that each sequence set  $C^{(a,k)}$  is a (N,N,N)-CCC. Similarly we can prove it when  $(a,k) \in \mathcal{A} \setminus \mathcal{D}$ .

We now prove the second part. Let us consider  $C^{(a_1,k_1,m_1)}\in C^{(a_1,k_1)}$ , when  $(a_1,k_1)\in \mathcal{D}$  and  $C^{(a_2,k_2,m_2)}\in \mathcal{D}$ 

 $C^{(a_2,k_2)}$ , when  $(a_2,k_2)\in\mathcal{A}\setminus\mathcal{D}$ . Then the aperiodic correlation of  $C^{(a_1,k_1,m_1)}$  and  $C^{(a_2,k_2,m_2)}$  is

$$\begin{split} \sum_{s=0}^{N-1} \tilde{R}_{C_s^{(a_1,k_1,m_1)},C_s^{(a_2,k_2,m_2)}}(\tau) \\ &= \sum_{s=0}^{N-1} \sum_{t=0}^{N-1-\tau} \omega_N^{f_s^{(a_1,k_1,m_1)}(t)} \cdot \omega_N^{-f_s^{(a_2,k_2,m_2)}(t+\tau)} \\ &= \sum_{s=0}^{N-1} \sum_{t=0}^{N-1-\tau} \omega_N^{t(m_1+m_2)+m_2\tau+s\left(k_1\pi_{a_1}(t)+k_2\pi_{a_2}(t+\tau)\right)-N} \\ &= \sum_{t=0}^{N-1-\tau} \omega_N^{t(m_1+m_2)+m_2\tau} \cdot \sum_{s=0}^{N-1} \omega_N^{s\left(k_1\pi_{a_1}(t)+k_2\pi_{a_2}(t+\tau)\right)}. \end{split}$$

Recall from our assumption in Construction 1,  $k_1\pi_{a_1}(t)+k_2\pi_{a_2}(t+\tau)=0\pmod{N}$  for any  $0\leq\tau\leq N-1, k_1\neq k_2$  has only one solution. Therefore, for  $0\leq t'\leq N-1$  and  $k_1\pi_{a_1}(t')+k_2\pi_{a_2}(t'+\tau)=0\pmod{N}$ . If  $N-1\geq t'\geq N-\tau$ , then  $\sum_{s=0}^{N-1}\tilde{R}_{C_s^{(a_1,k_1,m_1)},C_s^{(a_2,k_2,m_2)}(\tau)}=0 \text{ due to } \sum_{s=0}^{N-1}\omega_N^{s(k_1\pi_{a_1}(t)+k_2\pi_{a_2}(t+\tau))}=0.$  For  $0\leq t'\leq N-\tau$ , we have

$$\begin{split} \sum_{s=0}^{N-1} & \tilde{R}_{C_s^{(a_1,k_1,m_1)},C_s^{(a_2,k_2,m_2)}}(\tau) \\ &= \omega_N^{m_2\tau} \cdot [\omega_N^{(m_1+m_2)\cdot t'} \cdot N + \\ &\sum_{0 \leq t \leq N-1-\tau, \qquad 0 \leq s \leq N-1} \omega_N^{(k_1\pi a_1(t)+k_2\pi a_2(t+\tau))\cdot s}] \\ &= \omega_N^{m_2\tau+(m_1+m_2)t'} \cdot N. \end{split}$$

Hence,  $|\sum\limits_{s=0}^{N-1} \tilde{R}_{C_s^{(a_1,k_1,m_1)},C_s^{(a_2,k_2,m_2)}}(\tau)| \leq N$  for all  $0 \leq \tau \leq N-1, k_1 \neq k_2$  and  $1 \leq m_1, m_2 \leq N-1$ .

Let us consider  $C^{(a_1,k_1,m_1)} \in C^{(a_1,k_1)}$  and  $C^{(a_2,k_2,m_2)} \in C^{(a_2,k_2)}$ , when  $(a_1,k_1), (a_2,k_2) \in \mathcal{A} \setminus \mathcal{D}$ . Then the aperiodic correlation of  $C^{(a_1,k_1,m_1)}$  and  $C^{(a_2,k_2,m_2)}$  is

$$\begin{split} &\sum_{s=0}^{N-1} \tilde{R}_{C_s^{(a_1,k_1,m_1)},C_s^{(a_2,k_2,m_2)}}(\tau) \\ &= \sum_{s=0}^{N-1} \sum_{t=0}^{N-1-\tau} \omega_N^{f_{s}^{(a_1,k_1,m_1)}(t)} \cdot \omega_N^{-f_{s}^{(a_2,k_2,m_2)}(t+\tau)} \\ &= \sum_{s=0}^{N-1} \sum_{t=0}^{N-1-\tau} \omega_N^{t(m_2-m_1)+m_2\tau+s\left(k_2\pi_{a_2}(t+\tau)-k_1\pi_{a_1}(t)\right)} \\ &= \sum_{t=0}^{N-1-\tau} \omega_N^{t(m_2-m_1)+m_2\tau} \cdot \sum_{s=0}^{N-1} \omega_N^{s\left(k_2\pi_{a_2}(t+\tau)-k_1\pi_{a_1}(t)\right)}. \end{split}$$

Recall from our assumption in Construction 1,  $k_1\pi_{a_1}(t)-k_2\pi_{a_2}(t+\tau)=0\pmod{N}$  for any  $0\leq\tau\leq N-1, k_1\neq k_2$  has only one solution. Therefore, for  $0\leq t'\leq N-1$  and  $k_1\pi_{a_1}(t')-k_2\pi_{a_2}(t'+\tau)=0\pmod{N}$ . If  $N-1\geq t'\geq N-\tau$ , then  $\sum_{s=0}^{N-1}\tilde{R}_{C_s^{(a_1,k_1,m_1)},C_s^{(a_2,k_2,m_2)}(\tau)}=0$  due to

$$\sum_{\substack{s=0\\\text{have}}}^{N-1} \omega_N^{s(k_2\pi_{a_2}(t+\tau)-k_1\pi_{a_1}(t))} = 0. \text{ For } 0 \le t' \le N-\tau, \text{ we}$$

$$\begin{split} \sum_{s=0}^{N-1} \tilde{R}_{C_s^{(a_1,k_1,m_1)},C_s^{(a_2,k_2,m_2)}}(\tau) \\ &= \omega_N^{m_2\tau} \cdot [\omega_N^{(m_2-m_1)\cdot t'} \cdot N + \\ &\sum_{0 \leq t \leq N-1-\tau, \qquad 0 \leq s \leq N-1} \omega_N^{(k_2\pi a_2(t+\tau)-k_1\pi a_1(t))\cdot s}] \\ &= \omega_N^{m_2\tau + (m_1+m_2)t'} \cdot N. \end{split}$$

Hence, 
$$|\sum_{s=0}^{N-1} \tilde{R}_{C_s^{(a_1,k_1,m_1)},C_s^{(a_2,k_2,m_2)}}(\tau)| \leq N$$
 for all  $0 \leq \tau \leq N-1, k_1 \neq k_2$  and  $1 \leq m_1, m_2 \leq N-1$ .

Similarly, we can prove the theorem for other cases. Hence, the theorem follows.

#### REFERENCES

- J. A. Davis and J. Jedwab, "Peak-to-mean power control in OFDM, Golay complementary sequences, and Reed-Muller codes," *IEEE Trans. Inf. Theory*, vol. 45, no. 7, pp. 2397-2417, 1999.
- [2] P. Spasojevic and C. N. Georghiades, "Complementary sequences for ISI channel estimation," *IEEE Trans. Inf. Theory*, vol. 47, no. 3, pp. 1145-1152, 2001.
- [3] S. Wang and A. Abdi, "MIMO ISI channel estimation using uncorrelated Golay complementary sets of polyphase sequences," *IEEE Trans. Inf. Theory*, vol. 56, no. 5, pp. 3024-3039, 2007.
- [4] A. Pezeshki, A. R. Calderbank, W. Moran, and S. D. Howard, "Doppler resilient Golay complementary waveforms," *IEEE Trans. Inf. Theory*, vol. 54, no.9, pp. 4254-4266, 2008.
- [5] H. H. Chen, "The Next Generation CDMA Techonolgies," 1st edition. John Wiley & Sons, 2007.
- [6] Z. Liu, Y. L. Guan, B. C. Ng and H. Chen, "Correlation and Set Size Bounds of Complementary Sequences with Low Correlation Zone," *IEEE Trans. Commun.*, vol. 59, no. 12, pp. 3285-3289, 2011.
- [7] Z. Liu, U. Parampalli, Y. L. Guan, and S. Bozetas, "Constructions of optimal and near-optimal quasi-complementary sequence sets from singer difference sets," *IEEE Wire. Commun. Lett.*, vol. 2, no.5, pp. 487-490, 2013.
- [8] P. Ke and Z. Zhou, "A Generic Construction of Z-Periodic Complementary Sequence Sets with Flexible Flock Size and Zero Correlation Zone Length," *IEEE Signal Process. Lett.*, vol. 22, no. 9, pp. 1462-1466, 2015.
- [9] Z. Liu, U. Parampalli and Y. L. Guan, "Optimal odd-length binary Z-complementary pairs," *IEEE Trans. Inf. Theory*, vol. 60, no. 9, pp. 5768-5781, 2014.
- [10] Y. Li, C. Xu, N. Jing and K. Liu, "Constructions of Z-periodic complementary sequence set with flexible flock size?," *IEEE Commun. Lett.*, vol. 18, no. 2, pp. 201-204, 2014.
- [11] A. R. Adhikary, S. Majhi, "New construction of optimal aperiodic Z-complementary sequence sets of odd-lengths," *Electron. Lett.*, 2019.
- [12] P. Sarkar, S. Majhi, and Z. Liu, "Optimal Z-complementary code set from generalized Reed-Muller codes," *IEEE Trans. Commun.*, vol. 67, no. 3, pp. 1783-1796, 2018.
- [13] C. Chen, "A novel construction of Z-complementary pairs based on generalized Boolean functions", *IEEE Signal Process. Lett.* vol. 24, no. 7, pp. 987-990, 2017.
- [14] A. R. Adhikary, S. Majhi, Z. Liu, and Y. L. Guan, "New sets of even length binary Z-complementary pairs with asymptotic ZCZ ratio of 3/4," *IEEE Sign. Process. Lett.*, vol. 25, no. 7, pp. 970-973, 2018.
- [15] L. R. Welch, "Lower bounds on the maximum cross-correlation of signals," *IEEE Trans. Inf. Theory*, vol. IT-20, no. 3, pp. 397C399, 1974.
- [16] Z. Liu, Y. L. Guan and W. H. Mow, "Improved lower bound for quasi-complementary sequence set," *Proc. IEEE Int. Symp. Inf. Theory*, St. Petersburg, 2011, pp. 489-493.
- [17] Z. Liu, Y. L. Guan and W. H. Mow, "A tighter correlation lower bound for quasi-complementary sequence sets," *IEEE Trans. Inf. Theory*, vol. 60, no. 1, pp. 388-396, 2014.

- [18] Z. Liu, Y. L. Guan and W. H. Mow, "Asymptotically locally optimal weight vector design for a tighter correlation lower Bound of Quasi-Complementary Sequence Sets," *IEEE Trans. Signal Process.*, vol. 65, no. 12, pp. 3107-3119, 2017.
- [19] Z. Liu, U. Parampalli, Y. L. Guan and S. Boztas, "Constructions of optimal and near-optimal quasi-complementary sequence sets from singer difference sets," *IEEE Wireless Commun. Lett.*, vol. 2, no. 5, pp. 487-490, 2013.
- [20] Y. Li, T. Liu and C. Q. Xu, "Constructions of asymptotically optimal quasi-complementary sequence sets," *IEEE Commun. Lett.* vol. 22, no. 8, pp. 1516-1519, 2018.
- [21] Y. Li, L. Y. Tian, T. Liu and C. Q. Xu, "Constructions of quasi-complementary sequence sets associated with characters," *IEEE Trans. Inf. Theory*, vol. 65, no. 7, pp. 4597-4608, 2019.
- [22] Y. Li, T. Yan and C. Lv, "Construction of a near-optimal quasi-complementary sequence set from almost difference set," *Cryptogr. Commun.*, pp. 815-824, Jul. 2019.
- [23] Y. Li, L. Tian, T. Liu and C. Xu, "Two constructions of asymptotically optimal quasi-complementary sequence sets," *IEEE Trans. Commun.*, vol. 67, no. 3, pp. 1910-1924, 2019.
- [24] Y. Li, L. Tian and C. Xu, "Constructions of asymptotically optimal aperiodic quasi-complementary sequence sets," *IEEE Trans. Commun.*, Early access.
- [25] T. Liu, C. Xu and Y. Li, "Multiple complementary sequence sets with low inter-set cross-correlation property," *IEEE Signal Process. Lett.*, vol. 26, no. 6, pp. 913-917, 2019.
- [26] Z. Zhou, D. Zhang, T. Helleseth and J. Wen, "A construction of multiple optimal zcz sequence sets with good cross correlation," *IEEE Trans. Inf. Theory*, vol. 64, no. 2, pp. 1340-1346, 2018.