# A geometrically based criterion to avoid infimum-gaps in Optimal Control

Michele Palladino and Franco Rampazzo

September 16, 2019

#### Abstract

In optimal control theory the expression *infimum qap* means a stricly negative difference between the infimum value of a given minimum problem and the infimum value of a new problem obtained by the former by *extending* the original family  $\mathcal{V}$  of controls to a larger family  $\mathcal{W}$ . Now, for some classes of domain-extensions –like convex relaxation or impulsive embedding of unbounded control problems- the *normality* of an extended minimizer has been shown to be sufficient for the avoidance of an infimum gaps. A natural issue is then the search of a general hypothesis under which the criterium 'normality *implies no gap'* holds true. We prove that, far from being a peculiarity of those specific extensions and from requiring the convexity of the extended dynamics, this criterium is valid provided the original family  $\mathcal{V}$  of controls is *abundant* in the extended family  $\mathcal{W}$ . Abundance, which is stronger than the mere  $C^0$ -density of the original trajectories in the set of extended trajectories, is a dynamical-topological notion introduced by J. Warga, and is here utilized in a 'non-convex' version which, moreover, is adapted to differential manifolds. To get the main result, which is based on set separation arguments, we prove an open mapping result valid for Quasi-Differential-Quotient (QDQ) approximating cones, a notion of 'tangent cone' resulted as a peculiar specification of H. Sussmann's Approximate-Generalized-Differential-Quotients (AGDQ) approximating cone.

### 1 Introduction

One of the main reason for enlarging the domain of a minimum problem relies on the aim of establishing the existence of at least one solution. Actually, domain extension is a quite common and variously motivated practice, in particular in the Calculus of Variations and in Optimal Control. Of course, a crucial requisite of such a domain enlargement consists in the *density* of the original problem in the new one: the extended minimum should be approximable by processes of the original problem. However, because of the presence of a final target, even a dense extension of the domain may result in the occurrence of an *infimum gap*: namely, it can happen that the infimum value of the original problem is strictly greater than the infimum value of the extended problem. This might be undesirable in many respects, for instance in the convergence of numerical schemes as well as in the identification of the value function via Hamilton-Jacobi equations. This raises a natural question: how can one avoid this gap phenomenon? A sufficient condition for gap avoidance seems to emerge from investigations by J. Warga [42, 43, 44, 45] and from some other more recent papers [28, 31, 32, 33, 34], dealing with some particular cases: this criterion is the so-called *normality* of minimizers. Therefore, the mentioned question can be turned into the following one:

**Q.** Under which hypotheses on a general optimal control problem normality is sufficient for gap-avoidance?

In order be more precise, let us briefly sketch the abstract setting of our optimal control problem. The *state* variable y will range on a Riemannian manifold  $\mathcal{M}$ , while the control maps  $v(\cdot)$  will belong to an *original* family  $\mathcal{V} \subset \mathcal{W} := L^1([0, S], \mathfrak{W})$  —where  $\mathfrak{W}$  is a subset of a metric space– or to a larger set  $\mathcal{W}$ , which will be called the *extended* family of controls. Given an initial state  $\bar{y} \in \mathcal{M}$  and a time interval [0, S], we will consider the control system

(E) 
$$\begin{cases} \frac{dy}{ds}(s) = f(s, y(s), w(s))\\ y(0) = \bar{y}, \end{cases}$$

and, for every  $w \in \mathcal{W}$ , we will use  $y[w] : [0, S] \to \mathcal{M}$  the corresponding (supposedly unique) solution. The *original optimal control problem* is defined as

$$(P)_{\mathcal{V}} \qquad Minimize\bigg\{ h\big(y[v](S)\big) \quad | \quad v \in \mathcal{V}, \quad y[v](S) \in \mathfrak{T} \bigg\},$$

where the cost function  $h: \mathcal{M} \to \mathbb{R}$  is continuous, and  $\mathfrak{T} \subset \mathcal{M}$  is a closed set called *target*.

Replacing the family of controls  $\mathcal{V}$  by the larger set  $\mathcal{W}$ , one obtains the *extended optimal* control problem:

$$(P)_{\mathcal{W}} \qquad Minimize \Big\{ h \big( y[w](S) \big) \mid w \in \mathcal{W}, \quad y[w](S) \in \mathfrak{T} \Big\}.$$

We will assume the existence of a local minimum for the extended problem, namely a control  $\hat{w} \in \mathcal{W}$  such that, for some  $C^0$  neighbourhood  $\mathcal{O}$  of  $y[\hat{w}]$ ,  $h(y[\hat{w}](S)) \leq h(y[w](S))$  for all  $w \in \mathcal{W}$  such that  $y[w](S) \in \mathfrak{T}$  and  $y[w] \in \mathcal{O}$ . The non-occurrence of infimum gaps means that the original infimum value is unaffected by the introduction of the extended controls, namely

$$h(y[\hat{w}](S)) = \inf \left\{ h(y[w](S)) \quad | \quad v \in \mathcal{V}, \quad y[v](S) \in \mathfrak{T}, \ y[v] \in \mathcal{O} \right\}$$

for all sufficiently small neighbourhoods  $\mathcal{O}$  of  $y[\hat{w}]$ 

If, on the contrary, there exists a neighbourhood  $\mathcal O$  such that

$$h(y[\hat{w}](S)) < \inf \left\{ h(y[w](S)) \mid v \in \mathcal{V}, \quad y[v](S) \in \mathfrak{T}, \ y[v] \in \mathcal{O} \right\},$$

one says that the optimal control problem has an infimum gap at  $y[\hat{w}]$ . Obviously, via the usual reductions, one can formulate a notion of infimum gap for a general Bolza problem –where an integral cost is involved as well–.

For problems defined on Euclidean spaces and such that the extended dynamics is convex, an insightfull investigation of the gap question and its relation with normality was carried out by J.Warga (see e.g. [43]). More recently two specific classes of domain extensions –still assuming the convexity of the extended dynamics– have been studied in [28, 32, 33, 34]. As mentioned above, these investigations share the fact that a certain condition turns out to be necessary for the gap occurrence:

(A) There is an infimum-gap only if the minimum of the extended problem is an abnormal extremal.  $^{1}$ 

Since any version of the PMP states that 'an optimal process  $(\hat{y}, \hat{w}) := (y[\hat{w}], \hat{w})$  for the extended problem  $(P)_{W}$  is an extremal', in order for **(A)** to have a precise meaning one has to specify which version of the Pontryagin Maximum Principle (PMP) one is considering. In turn, this is equivalent to specify which kind of *approximating cones* we are going to utilize for both the *reachable set* and the target  $\mathfrak{T}$ . Actually, for this goal we shall introduce a generalized differential called *Quasi Differential Quotient* (QDQ) (Def. 2.3)<sup>2</sup> and the

<sup>&</sup>lt;sup>1</sup>Equivalently: if the minimum is normal (=not abnormal) there is no gap.

<sup>&</sup>lt;sup>2</sup>A QDQ is a special case of Sussmann's Approximate Generalized Differential Quotient [40].

associated notion of QDQ approximating cone (Def. 2.5). Let us say immediately that this choice is perhaps the most important step for the validity of the main result. And while it is impossible at this stage to give an exhaustive description of what QDQ approximating cones are, let us point out that, on the one hand, they are sufficiently small for a fixed point theorem to hold true and, on the other hand, they are enough large to allow the utilization of the notion of abundance, which, as we shall see, proves crucial for normality to imply no gap.

This said, let us give the notion of *extremal*. For simplicity, we consider here only the case when the state ranges on a Euclidean space. Moreover, if  $C \subset \mathbb{R}^n$  is a cone, we use  $C^{\perp}$  to denote the *polar cone of* C, namely the set of linear forms  $\lambda \in (\mathbb{R}^n)^*$  such that  $\lambda \cdot c \leq 0$  for all  $c \in C$ .

**Definition 1.1** (Extremal). Consider a control  $\hat{w} \in W$  and the corresponding trajectory  $\hat{y} := y[\hat{w}]$ . Assume that  $\hat{y}(S) \in \mathfrak{T}$ , and let C be a QDQ approximating cone of the target  $\mathfrak{T}$  at  $\hat{y}(S)$ . We say that the process  $(\hat{y}, \hat{w})$  is an extremal (with respect to h and C) if there exist an absolutely continuous (adjoint) path  $\lambda \in W^{1,1}([0, S]; (\mathbb{R}^n)^*)$  and a cost multiplier  $\lambda_c \in \{0, 1\}$  such that  $(\lambda, \lambda_c) \neq 0$  and

- $(i) \quad \frac{d\lambda}{ds} = -\lambda \cdot \frac{\partial f}{\partial x}(s,\lambda(s),\hat{w}(s))$
- (*ii*)  $\max_{\mathfrak{w}\in\mathfrak{W}}\lambda(s)\cdot f(s,\hat{y}(s),\mathfrak{w}) = \lambda(s)\cdot f(s,\hat{y}(s),\hat{w}(s)) \quad a.e. \ s \in [0,S];$

$$(iii) \quad \lambda(S) \in -\lambda_c \nabla h(\hat{y}(S)) - C^{\perp}$$

Furthermore, we say that an extremal  $(\hat{y}, \hat{w})$  is normal if for every choice of the pair  $(\lambda, \lambda_c)$  one has  $\lambda_c = 1$ . We say that an extremal  $(\hat{y}, \hat{w})$  is abnormal if it is not normal, namely, if exists a choice of  $(\lambda, \lambda_c)$  with  $\lambda_c = 0$ .

As mentioned above, in [32, 33, 34], where the original set of controls  $\mathcal{V}$  was embedded in the set  $\mathcal{W}$  of relaxed controls, it has been shown the validity of criterion (**A**), that is: *if* the optimal process  $(\hat{y}, \hat{w})$  is a normal extremal, then an infimum gap cannot occur at  $\hat{y}$ . An akin result has been achieved in [28], where the system is control-affine and the original set  $\mathcal{V}$  comprises unbounded controls ranging in a convex cone. In that case a space-time, impulsive, extension is considered, namely the larger set of trajectories corresponding to  $\mathcal{W}$  comprises space-time paths which are allowed to evolve along fixed time directions.<sup>3</sup>

It is worth noticing that in both the investigated cases the original set of trajectories is *dense* in the set of extended trajectories, when the latter is endowed with  $C^0$  topology. So, one might conjecture that criterion (A) is still valid as soon as the trajectories corresponding to  $\mathcal{V}$  are dense in the set of trajectories corresponding to  $\mathcal{W}$ . In fact, *this is not the case*, as shown by the simple example in Appendix A.1.

Hence, a condition stronger than density is needed. For this goal we introduce Kaskoz' version of J. Warga's notion of  $\mathcal{V}$  being *abundant* in  $\mathcal{W}$  (Def. 4.1). This condition strengthens density by requiring that the trajectories of the extended system's convexification are uniformly approached by trajectories of the original system.

We will further extend the notion of abundant subfamily  $\mathcal{V} \subset \mathcal{W}$  to systems defined on manifolds and to fairly general classes of controls (which are merely required to belong to a metric space). Then, aiming to express normality of extended trajectories in geometric terms, we invoke local set separation of the target from the original reachable set.

A crucial result for the achievement of the main theorem consists in showing that, with this notion of abundance, every needle-variational cone  $\mathbf{C}$  at  $\hat{y}$  corresponding to the enlarged domain  $\mathcal{W}$  is also a QDQ approximating cone to the original reachable set (Theorem 4.1).

The next step consists in showing that the local set separation of the target from the original reachable set implies the linear separability between a QDQ approximating cone

 $<sup>^{3}</sup>$ It is well-known that under commutativity hypotheses these paths could be regarded as measure, while the measure-theoretical approach is unfit for non-commutative problems, see e.g. [8],[24].

to the target and the above mentioned needle-variational cone  $\mathbf{C}$  (Theorem 5.1). This is exactly the point where the choice of QDQ approximating cones –rather then other more classical cones, e.g. Boltiansky cones– plays essential. By expressing this linear separation in terms of adjoint paths, one finally gets the main result of the paper (Corollary 5.1), where, under the abundance hypothesis, statement (A) is turned into an actual theorem.

In Section 7 we apply the main theorem to nonlinear systems whose dynamics are neither bounded nor convex. Finally, since normality cannot be verified *a priori*, it is important to find sufficient conditions on the data guaranteeing that all minimizers are normal. This is what is provided by Theorem 5.2, where a directly verifiable criterion for normality is proved to hold true in the general setting.

#### **1.1** Basic notions and notation

#### 1.1.1 Linear spaces, manifolds

Let *E* be a real linear space, and let us use  $E^*$  to denote the algebraic dual of *E*. If  $\langle \cdot, \cdot \rangle$  is a given scalar product on E,<sup>4</sup> we will use  $|\cdot|$  to denote the norm associated with  $\langle \cdot, \cdot \rangle$ , namely, for every  $e \in E$  we set  $|e| = \sqrt{\langle e, e \rangle}$ . For every  $e \in E$  and every real number  $r \ge 0$  let us use  $e + B_r$  to denote the closed ball of center *e* and radius *r*, namely  $e + B_r = \{e + f \mid |f| \le r\}$ . When e = 0 we will write  $B_r$  instead of  $0 + B_r$ 

If  $E_1, E_2$  are real linear spaces and  $L \in Lin(E_1, E_2)$ , we shall use  $L \cdot e$  to denote the element of  $E_2$  coinciding with the image of  $e \in E_1$ . We will use the symbol  $\cdot$  also to mean duality. Furthermore, if  $\lambda \in E_2^*$  and and  $L \in Lin(E_1, E_2)$ , sometimes we will use the notation  $\lambda \cdot L$  to mean "the element of  $E_1^*$  coinciding the image of  $\lambda$  through the dual map of L." While it doesn't generate any confusion, this promiscuous use of the notation " $\cdot$ " makes the writing  $\lambda \cdot L \cdot e$  unambiguous, for one has  $(\lambda \cdot L) \cdot e = \lambda \cdot (L \cdot e)$  for all  $(e, \lambda) \in E_1 \times E_2^*$ .

By saying that  $(\mathcal{M}, \langle \cdot, \cdot \rangle)$  is a Riemannian differentiable manifold we will mean that  $\mathcal{M}$  is a differential manifold and  $\langle \cdot, \cdot \rangle$  is a Riemannan metric. For every  $x \in \mathcal{M}$  and  $e, f \in T_x \mathcal{M}, \langle e, f \rangle_x$  will denote the corresponding scalar product of e, f, and  $|e|_x := \sqrt{\langle e, e \rangle_x}$  will be called the *norm* of e. We will often omit the subscript and we will write  $\langle e, f \rangle$  and  $|e|_x$ .

We will use d to denote the distance induced on  $\mathcal{M}$  by  $\langle \cdot, \cdot \rangle$ . We recall that, if  $x_1, x_2 \in \mathcal{M}$ , the distance  $d(x_1, x_2)$  is defined as the minimum among the  $\langle \cdot, \cdot \rangle$ -lenghts of the absolutely continuous curves having  $x_1, x_2$  as end-points. For any  $x \in \mathcal{M}$  and any  $r \geq 0$ , we will use  $\mathcal{B}[x, r]$  to denote the closed ball of radius r and center x, i.e.  $\mathcal{B}[x, r] := \{y \in \mathcal{M} \mid d(x, y) \leq r\}.$ 

### **1.1.2** Cones

Let *E* be a real linear space. A subset  $K \subset E$  is a *cone* if  $\alpha k \in K$  for all  $(\alpha, k) \in [0, +\infty[\times K. \text{ If } A \subset E \text{ is any subset, we use <math>span^+A$  to denote the smallest convex cone containing *A*. Let us introduce a notion of *transversality* for cones.

The idea of a non-trivial intersection between cones, which plays essential in setseparation results like Theorem 2.3 below, is made formal is made formal by the following notion of *tranversality*:

**Definition 1.2.** Let E be a linear space and let  $K_1, K_2 \subseteq E$  be convex cones. We say that 1.  $K_1$  and  $K_2$  are transverse, if  $K_1 - K_2 := \{k_1 - k_2, (k_1, k_2) \in K_1 \times K_2\} = E$ ;

2.  $K_1$  and  $K_2$  are strongly transverse, if they are transverse and  $K_1 \cap K_2 \supseteq \{0\}$ .

<sup>&</sup>lt;sup>4</sup>By *scalar product* we mean a positive definite, symmetric, bilinear form.

Transversality differs from strong transversality only when  $K_1$  and  $K_2$  are complementary subspaces:

**Proposition 1.1.** Let E be a linear space, and let  $K_1, K_2 \subseteq E$  be convex cones. Then  $K_1, K_2$  are transverse if and only if either  $K_1, K_2$  are strongly transverse or  $K_1, K_2$  are linear subspaces such that  $K_1 \oplus K_2 = E$ .

**Definition 1.3.** Let E be a finite-dimensional linear space, and let  $E^*$  be its dual space. For any subset  $A \subset E$ , the (convex) cone

$$A^{\perp} \doteq \{ p \in E^* : p \cdot w \le 0 \quad \forall \ w \in A \} \subseteq E^*$$

(where the symbol  $\cdot$  denotes duality) will be called the polar cone of A.

The transversality of two cones is equivalent to their linearly separability. More precisely:

**Proposition 1.2.** Two convex cones  $K_1$  and  $K_2$  are not transverse if and only if  $K_1^{\perp} \cap K_2^{\perp} \setminus \{0\} \neq \emptyset$ , namely there exists a linear form  $\lambda \neq 0$  such that

 $\lambda \cdot k_1 \ge 0 \ \forall k_1 \in K_1 \quad and \quad \lambda \cdot k_2 \le 0 \ \forall k_2 \in K_2.$ 

In this case one also says that  $K_1$  and  $K_2$  are linearly separable.

### 1.1.3 Scorza-Dragoni points

**Definition 1.4** (Scorza-Dragoni point). Given a compact set  $X \subset \mathcal{M}$  and an interval  $[a,b] \subseteq \mathbb{R}, a < b$ , let us consider a function  $\varphi : [a,b] \times X \to \mathbb{R}^n$  verifying

- i)  $[a,b] \ni s \mapsto \varphi(s,y) \in \mathbb{R}^n$  is measurable for each  $y \in X$ ;
- ii)  $X \ni y \mapsto \varphi(s, y)$  is continuous for each  $s \in [a, b]$ ,

We say that  $\bar{s} \in [a, b]$  is a Scorza-Dragoni point for  $\varphi$  if, for all  $y \in X$ ,

$$\lim_{r \to 0} \lim_{\delta \to 0} \frac{1}{\delta} \int_{\bar{s}}^{\bar{s}+\delta} \Lambda_r(s, y) \, ds = 0 \tag{1.1}$$

where

$$\Lambda_r(s,y) := \sup_{x \in X, \ |x-y| \le r} d\left(\varphi(s,x), \varphi(\bar{s},y)\right) \tag{1.2}$$

We shall use SD  $\{\varphi\}$  to denote the set of all the Scorza-Dragoni points for the function  $\varphi$ .

Notice in particular that, if  $s \in SD \{\varphi\}$ , one has

$$\lim_{s \to y, \delta \searrow 0} \varphi(s + \delta, x) = \varphi(s, y), \tag{1.3}$$

for any  $y \in X$ . The importance of Scorza-Dragoni points relies on the fact they they form a full measure set [36]:

**Theorem 1.1** (Scorza-Dragoni). The set of all the Scorza-Dragoni points of a Caratheodory function  $\varphi : [a, b] \times X \to \mathbb{R}^n$  has measure equal to b - a.

### 2 Set separation and open mappings

### 2.1 Quasi Differential Quotients

In order to state the set-separation theorem (Th. 2.3) we need the notion of *Quasi Differential Quotients approximating cone* to a set  $\mathcal{E}$  at a point of its boundary. For this purpose let us introduce the notion of Quasi Differential Quotient, which in turn is a particular case of Sussmann's Approximate Generalized Differential Quotient [40]. The corresponding set-separation theorem is based on an Open Mapping we prove below.

Let us recall the notion of Cellina continuously approximable set-valued function, which is the building block in the definition of Approximate Generalized Differential Quotient.

**Definition 2.1** (CCA). Let  $F : \mathbb{R}^N \to \mathbb{R}^n$  be a set-valued map. We say that F is a Cellina continuously approximable (CCA) set-valued map *if*, for any compact set  $K \subset \mathbb{R}^N$ ,

- the restriction of F on K has compact graph, that is, the set  $\operatorname{Gr}(F_{|_K}) := \{(x, y) \in K \times \mathbb{R}^n : y \in F(x)\}$  is compact, and
- there exists a sequence of single-valued, continuous maps  $f_k : K \to \mathbb{R}^n$ ,  $k \in \mathbb{N}$ , such that the following condition holds: for every open set  $\Omega$  satisfying  $\operatorname{Gr}(F_{|_K}) \subset \Omega$ , there exists  $k_{\Omega}$  such that  $\operatorname{Gr}(f_k) := \{(x, y) \in K \times \mathbb{R}^n : y \in f_k(x)\} \subset \Omega$  for every  $k \ge k_{\Omega}$ .

We will say that a function  $\rho : [0; +\infty[\rightarrow [0; +\infty]]$  is a *a pseudo-modulus* if it is monotonically nondecreasing and  $\lim_{s\to 0^+} \rho(s) = \rho(0) = 0$ . We call *modulus* a pseudo-modulus taking values in  $[0, +\infty[$ .

**Definition 2.2** (AGDQ). Assume that  $F : \mathbb{R}^N \to \mathbb{R}^n$  is a set-valued map,  $(\bar{\gamma}, \bar{y}) \in \mathbb{R}^N \times \mathbb{R}^n$ ,  $\Lambda \subset Lin\{\mathbb{R}^N, \mathbb{R}^n\}$  is a compact set, and  $\Gamma \subset \mathbb{R}^N$  is any subset. We say that  $\Lambda$  is an Approximate Generalized Differential Quotient (AGDQ) of F at  $(\bar{\gamma}, \bar{y})$  in the direction of  $\Gamma$  if there exists a pseudo-modulus  $\rho$  having the property that

(\*) for every  $\delta > 0$  such that  $\rho(\delta) < +\infty$ , there exists CCA set-valued map  $A^{\delta}: (\bar{\gamma} + B_{\delta} \cap \Gamma) \to Lin\{\mathbb{R}^{N}, \mathbb{R}^{n}\} \times \mathbb{R}^{n}$  such that

$$\inf_{L' \in \Lambda} |L - L'| \le \rho(\delta), \quad |h(\gamma)| \le \delta \rho(\delta), \quad and \ \bar{y} + L \cdot (\gamma - \bar{\gamma}) + h \in F(\gamma)^{-5}$$

whenever  $\gamma \in \overline{\gamma} + B(\delta) \cap \Gamma$  and  $(L, h) \in A^{\delta}(\gamma)$ .

We will use a subclass of AGDQs, which we call the *Quasi Differential Quotients*. Their main property consists in the validity of an actual, *not punctured*, open mapping theorem (see Theorem 2.2below).

**Definition 2.3** (QDQ). Assume that  $F : \mathbb{R}^N \to \mathbb{R}^n$  is a set-valued map,  $(\bar{\gamma}, \bar{y}) \in \mathbb{R}^N \times \mathbb{R}^n$ ,  $\Lambda \subset Lin\{\mathbb{R}^N, \mathbb{R}^n\}$  is a compact set, and  $\Gamma \subset \mathbb{R}^N$  is any subset. We say that  $\Lambda$  is a Quasi Differential Quotient (QDQ) of F at  $(\bar{\gamma}, \bar{y})$  in the direction of  $\Gamma$  if there exists modulus  $\rho : [0, +\infty[ \to [0, +\infty[$  having the property that

(\*) for every  $\delta > 0$  there exists a continuous map  $(L_{\delta}, h_{\delta}) : (\bar{\gamma} + B_{\delta} \cap \Gamma) \to Lin\{\mathbb{R}^{N}, \mathbb{R}^{n}\} \times \mathbb{R}^{n}$  such that

$$\min_{L' \in \Lambda} |L_{\delta}(\gamma) - L'| \le \rho(\delta), \quad |h_{\delta}(\gamma)| \le \delta \rho(\delta), \quad and \ \bar{y} + L_{\delta} \cdot (\gamma - \bar{\gamma}) + h_{\delta}(\gamma) \in F(\gamma),$$

whenever  $\gamma \in \overline{\gamma} + B_{\delta} \cap \Gamma$ .

**Definition 2.4** (AGDQ and QDQ on manifolds). Let  $\mathcal{N}$ ,  $\mathcal{M}$  be  $C^1$  Riemannian manifolds. Assume that  $\tilde{F} : \mathcal{N} \rightsquigarrow \mathcal{M}$  is a set-valued map,  $(\bar{\gamma}, \bar{y}) \in \mathcal{N} \times \mathcal{M}$ ,  $\tilde{\Lambda} \subset Lin\{T_{\gamma}\mathcal{N}, T_{y}\mathcal{M}\}$ is a compact set, and  $\Gamma \subset \mathcal{N}$  is any subset. Moreover, let  $\phi : U \to \mathbb{R}^N$  and  $\psi : V \to \mathbb{R}^n$ be charts defined on neighbourhoods U and V of  $\bar{\gamma}$  and  $\bar{y}$ , respectively, and assume that  $\phi(\bar{\gamma}) = 0, \psi(\bar{y}) = 0$ . Consider the map  $\psi \circ \tilde{F} \circ \phi^{-1} : \psi(U) \to \mathbb{R}^n$  and extend it arbitrarily to a map  $F : \mathbb{R}^N \to \mathbb{R}^n$ . We say that  $\tilde{\Lambda}$  is an Approximate Generalized Differential Quotient (AGDQ) [resp. a Quasi Differential Quotient (QDQ)] of  $\tilde{F}$  at  $(\bar{\gamma}, \bar{y})$  in the direction of  $\tilde{\Gamma}$ if  $\Lambda := D\psi(\bar{y}) \circ \tilde{\Lambda} \circ D\phi^{-1}(0)$  is an Approximate Generalized Differential Quotient [resp. a Quasi Differential Quotient] of F at (0, 0) in the direction of  $\Gamma := \phi(\tilde{\Gamma} \cap U)$ .

As pointed out in [40], this definition is intrinsic, that is, it is independent of the choice of the charts  $\phi$  and  $\psi$ .

<sup>&</sup>lt;sup>5</sup> Here  $|\cdot|$  denotes the operator norm, namely  $|M| = \sup_{|v|=1} |M \cdot v|$ , for every linear operator  $M \in Lin(\mathbb{R}^N, \mathbb{R}^n)$ .

### 2.2 Open Mapping results

**Theorem 2.1** (Directional Open Mapping). Let N, n be positive integers, and let  $\Gamma$  be a convex cone in  $\mathbb{R}^N$ . Let  $F : \mathbb{R}^N \rightsquigarrow \mathbb{R}^n$  be a set-valued map, and let  $\Lambda$  be a AGDQ of F at  $(\bar{\gamma}, \bar{y})$  in the direction of  $\Gamma$ . Let us assume that there is an element  $\bar{w} \in \mathbb{R}^n$  such that  $\bar{w} \in Int(L \cdot \Gamma)$  for every  $L \in \Lambda$ . Then there exist a closed convex cone  $D \subseteq \mathbb{R}^n$  and positive constants  $\alpha, \beta$  verifying  $\bar{w} \in Int(D)$  and

$$\bar{y} + (B_a \setminus \{0\} \cap D) \subset F(\bar{\gamma} + B_{a\beta} \cap \Gamma) \quad for \ all \ a \in ]0, \alpha].$$
 (2.4)

If one takes  $\bar{w} = 0$  in the statement of Theorem 2.1, the cone *D* necessarily coincides with the whole  $\mathbb{R}^n$ . As a consequence, one obtains the following 'punctured' Open Mapping Theorem.

**Corollary 2.1** ('Punctured' Open Mapping). <sup>6</sup> Let N, n be positive integers, and let  $\Gamma$  be a convex cone in  $\mathbb{R}^N$ . Let  $F : \mathbb{R}^N \rightsquigarrow \mathbb{R}^n$  be a set-valued map, and let  $\Lambda$  be an QDQ of Fat  $(\bar{\gamma}, \bar{y})$  in the direction of  $\Gamma$ . Let us assume that  $\Lambda$  is surjective, by which we mean that  $L \cdot \Gamma = \mathbb{R}^n$  for every  $L \in \Lambda$ . Then there are positive constants  $\alpha, \beta$  verifying

$$\bar{y} + (B_a \setminus \{0\}) \subset F(\bar{\gamma} + B_{a\beta} \cap \Gamma) \quad for \ all \ a \in ]0, \alpha].$$
 (2.5)

If we replace AGDQ's with QDQ we get a real, non-punctured, open mapping result:

**Theorem 2.2** (Open Mapping). Let N, n be positive integers, and let  $\Gamma$  be a convex cone in  $\mathbb{R}^N$ . Let  $F : \mathbb{R}^N \rightsquigarrow \mathbb{R}^n$  be a set-valued map, and let  $\Lambda$  be a GDQ of F at  $(\bar{\gamma}, \bar{y})$  in the direction of  $\Gamma$ . As above, let us assume that  $\Lambda$  is surjective, by which we mean that  $L \cdot \Gamma = \mathbb{R}^n$  for every  $L \in \Lambda$ . Then the following statements (i), (ii) hold true:

(i) there are positive constants  $\alpha, \beta$  having the property that

$$\bar{y} + (B_a \setminus \{0\}) \subset F(\bar{\gamma} + B_{a\beta} \cap \Gamma) \quad for \ all \ a \in ]0, \alpha];$$
 (2.6)

(ii) there exists  $\check{\delta} > 0$  such that, for every  $\delta \leq \check{\delta}$  and every  $(L_{\delta}, h_{\delta})$  as in Definition 2.3, there exists  $\gamma_{\delta} \in \bar{\gamma} + \Gamma \cap B_{\delta}$  such that

$$\bar{y} = L_{\delta}(\gamma_{\delta}) \cdot (\gamma_{\delta} - \bar{\gamma}) + h_{\delta}(\gamma_{\delta}) \quad \Big[ \in F(\gamma_{\delta}) \Big].$$
(2.7)

In particular, by possibly reducing the size of  $\alpha$ , one gets the open-mapping inclusions

$$\bar{y} + B_a \subset F(\bar{\gamma} + B_{a\beta} \cap \Gamma)$$
 for all  $a \in ]0, \alpha]$ .

*Proof.* Without loss of generality, we can assume  $(\bar{\gamma}, \bar{y}) = (0, 0)$  and, since a QDQ is an AGDQ, it is sufficient to prove only statement (*ii*). Namely, for every  $\delta > 0$  sufficiently small, we have to establish the existence of a  $\gamma_{\delta} \in B_{\delta} \cap \Gamma$  such that

$$0 = L_{\delta}(\gamma_{\delta}) \cdot \gamma_{\delta} + h_{\delta}(\gamma_{\delta}). \tag{2.8}$$

For every  $\delta > 0$ , let us define the set-valued map  $L_{\delta}^{-1_r} : B_{\delta} \cap \Gamma \rightsquigarrow Lin(\mathbb{R}^n, \mathbb{R}^N)$  by setting, for every  $\gamma \in B_{\delta} \cap \Gamma$ ,

$$L_{\delta}^{-1_r}(\gamma) := \Big\{ M \in Lin(\mathbb{R}^n, \mathbb{R}^N), \ L_{\delta}(\gamma) \circ M = Id_{\mathbb{R}^n} \Big\}.$$

Namely,  $L_{\delta}^{-1_r}(\gamma)$  is the set of right inverse of  $L_{\delta}(\gamma)$ . Let us first observe that, for every  $\gamma \in B_{\delta} \cap \Gamma$ , and  $\delta$  sufficiently small,  $L_{\delta}^{-1_r}(\gamma)$  is non-empty. Indeed, it contains the Moore-Penrose pseudo-inverse

$$M^{\sharp}_{\delta}(\gamma) := L^{tr}_{\delta}(\gamma) \circ \left( L_{\delta}(\gamma) \circ L^{tr}_{\delta}(\gamma) \right)^{-1},$$

<sup>&</sup>lt;sup>6</sup> The adjective *punctured* here refers to the fact that  $\bar{y}$  does not belong the image  $F(\bar{\gamma} + B_{a\beta} \cap \Gamma)$ .

where  $t^r$  denotes transposition. Furthermore, it is trivial to verify that the set-valued map  $L_{\delta}^{-1_r}$  is convex-valued. Finally, by possibly reducing the size of  $\check{\delta}$ , for every  $\delta \in [0, \check{\delta}]$ , the set-valued map  $L_{\delta}^{-1_r}$  has compact graph. Indeed, there exist a constant K > 0 such that  $\Lambda^{\rho(\delta)}$  is a compact subset made of linear operators whose right inverse are bounded (in the operator norm) by K. Moreover, let us consider a sequence  $(\gamma_m)_{m \in \mathbb{N}} \subset B_{\delta} \cap \Gamma$  converging to  $\tilde{\gamma} \in B_{\delta} \cap \Gamma$ , and, for every  $m \in \mathbb{N}$ , let us choose  $M_m \in L_{\delta}^{-1_r}(\gamma_m)$ . Hence, one has that  $L_{\delta}(\gamma_m) \circ M_m = Id_{\mathbb{R}^n}$  and, since the sequence  $(M_m)$  ranges in a compact set, there exists a subsequence  $(M_{m_k})$  converging to a linear operator  $\tilde{M}$ . In particular,

$$L_{\delta}(\tilde{\gamma}) \circ \tilde{M} = \lim_{k \to \infty} \left( L_{\delta}(\gamma_{m_k}) \circ M_{m_k} \right) = Id_{\mathbb{R}^n},$$

so that  $\tilde{M} \in L^{-1_r}_{\delta}(\tilde{\gamma})$ . This proves that the set-valued map  $\gamma \mapsto L^{-1_r}_{\delta}(\gamma)$  has compact graph.

Now consider the set-valued map  $\Psi_{\delta}: B_{\delta} \cap \Gamma \rightsquigarrow \mathbb{R}^N$  defined by setting

$$\Psi_{\delta}(\gamma) := \left\{ -M \cdot h_{\delta}(\gamma) \mid M \in L_{\delta}^{-1_{r}}(\gamma) \right\} \bigcap \Gamma, \quad \gamma \in B_{\delta} \cap \Gamma.$$

To prove that this map has non-empty values for every  $\gamma \in B_{\delta} \cap \Gamma$ , it is sufficient to determine a linear mapping  $M^{\flat} : \mathbb{R}^n \to \mathbb{R}^N$  and an element  $v \in \Gamma$  such that

$$(L_{\delta}(\gamma) \circ M^{\flat}) \cdot w = w \quad \forall w \in \mathbb{R}^{n} \ \left( \iff M^{\flat} \in L_{\delta}^{-1_{r}}(\gamma) \right), \qquad -M^{\flat} \cdot h_{\delta}(\gamma) = v \quad (2.9)$$

Fix  $\gamma \in B_{\delta} \cap \Gamma$  and choose  $v \in \Gamma$  verifying  $L_{\delta}(\gamma) \cdot v = -h_{\delta}(\gamma)$ . Such a v exists, since  $L_{\delta}(\gamma)$  is surjective. Now, a geometrical intuition suggests that  $M^{\flat}$  might be obtained by adding a suitable linear operator to an element of  $L_{\delta}^{-1_{r}}(\gamma)$ , for instance the linear operator  $M^{\sharp}$ . Actually, following [40], if  $\langle \cdot, \cdot \rangle$  is any scalar product on  $\mathbb{R}^{n}$ , we define the linear map  $M^{\flat} : \mathbb{R}^{n} \to \mathbb{R}^{N}$  by setting, for every  $w \in \mathbb{R}^{n}$ ,

$$M^{\flat} \cdot w := M^{\sharp} \cdot w - \frac{\langle w, h_{\delta}(\gamma) \rangle}{\langle h_{\delta}(\gamma), h_{\delta}(\gamma) \rangle} \left( v - M^{\sharp} \cdot h_{\delta}(\gamma) \right).$$

It is straightforward to verify that  $M^{\flat}$  verifies conditions (2.9), so that  $\Psi_{\delta}(\gamma)$  is not empty.

Since for every  $\delta$  the map  $h_{\delta}$  is continuous and  $||h_{\delta}|| \leq \delta\rho(\delta)$ , by possibly reducing the size of  $\check{\delta}$  we conclude that, for every  $\delta \in [0, \check{\delta}]$ , the set-valued map  $\Psi_{\delta}$  verifies  $\Psi_{\delta}(B_{\delta} \cap \Gamma) \subset \overline{B}_{\delta} \cap \Gamma$  and has non-empty, convex values, and a closed graph. Since the domain of  $\Psi_{\delta}$  is compact and convex, the set-valued map  $\Psi_{\delta}$  verifies the hypotheses of the Kakutani fixed point theorem, so that there exists  $\gamma_{\delta} \in B_{\delta} \cap \Gamma$  such that  $\gamma_{\delta} \in \Psi_{\delta}(\gamma_{\delta})$ . It follows that there is a matrix  $M \in L_{\delta}^{-1_{r}}(\gamma_{\delta})$  such that  $0 = \gamma_{\delta} + M \cdot h_{\delta}(\gamma_{\delta})$ . Therefore, one gets

$$0 = L_{\delta}(\gamma_{\delta}) \cdot \left(\gamma_{\delta} + M \cdot h_{\delta}(\gamma_{\delta})\right) = L_{\delta}(\gamma_{\delta}) \cdot \gamma_{\delta} + h_{\delta}(\gamma_{\delta}),$$

which concludes the proof.

### 2.3 QDQ approximating cones and set separation

Assume that  $\mathcal{M}$  is a  $C^1$  differentiable manifold,  $\mathcal{E} \subset \mathcal{M}$ , and  $z \in \mathcal{E}$ . If X is a linear space, let us call *convex multicone* in X any family of convex cones of X.

**Definition 2.5.** An AGDQ approximating multicone [resp. a QDQ approximating multicone] to  $\mathcal{E}$  at z is a convex multicone  $\mathcal{C} \subseteq T_z \mathcal{M}$  such that there exist a non-negative integer N, a set-valued map  $F : \mathbb{R}^N \rightsquigarrow \mathcal{M}$ , a convex cone  $\Gamma \subset \mathbb{R}^N$ , and an AGDQ [resp. a QDQ]  $\Lambda$  of F at (0, z) in the direction of  $\Gamma$  such that  $F(\Gamma) \subset \mathcal{E}$  and  $\mathcal{C} = \{L \cdot \Gamma : L \in \Lambda\}$ .

In the particular case when an AGDQ approximating multicone [resp. a QDQ approximating multicone] is a singleton, namely  $\Lambda = \{L\}$  for some  $L \in Lin(\mathbb{R}^N, \mathbb{R}^n)$ , we say that  $C := L \cdot \Gamma$  is an AGDQ approximating cone [resp. a QDQ approximating cone] to  $\mathcal{E}$ at z. Let us introduce the notion of local set-separation:

**Definition 2.6.** Let  $\mathcal{X}$  be a topological space, and let us consider two subsets  $\mathcal{A}_1, \mathcal{A}_2 \subset \mathcal{X}$ and a point  $z \in \mathcal{A}_1 \cap \mathcal{A}_2$ . We say that  $\mathcal{A}_1$  and  $\mathcal{A}_2$  are locally separated at z provided there exists a neighborhood V of z such that

$$\mathcal{A}_1 \cap \mathcal{A}_2 \cap V = \{z\}.$$

We are now ready to state our set-separation result, which connects set separation with the linear separability of QDQ approximating cones. Furthermore the result includes a special property in the case when the approximating cones are complementary linear subspaces.

**Theorem 2.3** (Set separation). Let  $\mathcal{E}_1, \mathcal{E}_2$  be subsets of  $\mathcal{M}$ , and let  $z \in \mathcal{E}_1 \cap \mathcal{E}_2$  Assume that  $C_1, C_2$  are AGDQ approximating cones of  $\mathcal{E}_1$  and  $\mathcal{E}_2$ , respectively, at z.

- i) If  $C_1$  and  $C_2$  are strongly transverse, then the sets  $\mathcal{E}_1$  and  $\mathcal{E}_2$  are not locally separated.
- ii) If moreover:
  - 1.  $C_1$ ,  $C_2$  are QDQ cones,
  - 2.  $C_1$  and  $C_2$  are complementary linear subspaces, i.e.  $C_1 \oplus C_2 = T_z \mathcal{M}$ ,
  - 3. for each  $i = 1, 2, \Gamma_i \subset \mathbb{R}^{N_i}$  is a convex cone,  $F_i : \mathbb{R}^{N_i} \rightsquigarrow \mathcal{M}$  is a set-valued map, and  $\Lambda_i = \{L^i\} \in Lin(\mathbb{R}^{N_i}, T_z\mathcal{M})$  is a QDQ of  $F_i$  at (0, z) in the direction of  $\Gamma_i$ ,  $F_i(\Gamma_i) \subseteq \mathcal{E}_i$ , and  $C_i = L^i \cdot \Gamma_i$ ,

then there exists a sequence  $(\gamma_{1_k}, \gamma_{2_k}) \in \Gamma_1 \times \Gamma_2$  such that  $z_k \in F_1(\gamma_{1_k}) \cap F_2(\gamma_{2_k})$ and  $z_k \to z$ .

**Remark 2.1.** Property *ii*), whose proof is based on the Open Mapping result stated in Theorem 2.2, is not true if we replace QDQ approximating cones with AGDQ approximating cones. Of course, this is connected with the non validity of a non-punctured open mapping result for AGDQ's.

Proof of Theorem 2.3. Statement i) of Theorem 2.3 is direct consequence of [40], Theorem 4.37, where an analogous result concerning the non-separation of multicones is provided

Let us prove statement *ii*). Because of the local character of the statement, there is not loss of generality in considering only the Euclidean case when  $\mathcal{M} = \mathbb{R}^n$ . For every i = 1, 2, let  $n_i \ge 0$  be the dimensions of the subspace  $C_i$  (so that  $n_1 + n_2 = n$ ), and let  $N_i \ge 0$  an integer such that  $\Gamma_i \subset \mathbb{R}^{N_i}$ . By hypothesis, for every i = 1, 2 there exists a modulus  $\rho_i : [0, +\infty[ \to [0, +\infty[$  having the property that, for every  $\delta > 0$ , there exists a continuous map  $(L_{\delta}^i, h_{\delta}^i) : B_{\delta} \cap \Gamma_i \to Lin\{\mathbb{R}^{N_i}, \mathbb{R}^{n_i}\} \times \mathbb{R}^{n_i}$ , such that

$$|L^i_{\delta}(\gamma_i) - L^i| \le \rho_i(\delta), \quad |h^i_{\delta}| \le \delta \cdot \rho_i(\delta), \text{ and } z + L^i_{\delta}(\gamma_i) \cdot \gamma_i + h^i_{\delta}(\gamma_i) \in F_i(\gamma_i)$$

whenever  $\gamma_i \in B_{\delta} \cap \Gamma_i$ . Let us consider the cone  $\Gamma := \Gamma_1 \times \Gamma_2 \subset \mathbb{R}^{N_1 + N_2}$  and the set-valued map  $F : \Gamma \rightsquigarrow \mathbb{R}^n$  defined by setting

$$F(\gamma_1, \gamma_2) := F_2(\gamma_2) - F_1(\gamma_1) = \left\{ z_2 - z_1 \mid (z_1, z_2) \in F_1(\gamma_1) \times F_1(\gamma_2) \right\} \quad \forall (\gamma_1, \gamma_2) \in \Gamma_1 \times \Gamma_2,$$

and observe that

if 
$$(\bar{\gamma}_1, \bar{\gamma}_2)$$
 is such that  $0 \in F(\bar{\gamma}_1, \bar{\gamma}_2)$  then  $\emptyset \neq F_2(\bar{\gamma}_2) \cap F_1(\bar{\gamma}_1) \subseteq \mathcal{E}_2 \cap \mathcal{E}_1$ .

Furthermore, let us set  $\rho(\delta) := \rho_1(\delta) + \rho_2(\delta)$  and let us define the continuous map

$$(L_{\delta}, h_{\delta})(\gamma_1, \gamma_2) := \left( \left( L_{\delta}^2(\gamma_2), -L_{\delta}^1(\gamma_1) \right) , h_{\delta}^2(\gamma_2) - h_{\delta}^1(\gamma_1) \right), \qquad (\gamma_1, \gamma_2) \in B_{\delta} \cap \Gamma.$$

Defining the linear map  $L \in Lin\{\mathbb{R}^{N_1+N_2}, \mathbb{R}^n\}$  by setting  $L(v_1, v_2) := L^2 \cdot v_2 - L^1 \cdot v_1$ , one obtains  $|L_{\delta}(\gamma_1, \gamma_2) - L| \leq \rho(\delta), |h_{\delta}| \leq \delta \cdot \rho(\delta)$ , and

$$L_{\delta}(\gamma_1, \gamma_2) \cdot (\gamma_1, \gamma_2) + h_{\delta}(\gamma_1, \gamma_2) \in F(\gamma_1, \gamma_2).$$

whenever  $(\gamma_1, \gamma_2) \in B_{\delta} \cap \Gamma$ . Hence,  $\Lambda$  is a QDQ of F at (0,0). Moreover, one has  $L \cdot \Gamma = C_1 \oplus C_2 = \mathbb{R}^n$  so that, by the Open Mapping result stated in Theorem 2.2 for  $k \in \mathbb{N}$  sufficiently large, we get the existence of  $(\bar{\gamma}_1^{\frac{1}{k}}, \bar{\gamma}_2^{\frac{1}{k}}) \in \Gamma \cap B_{\frac{1}{k}} \subset \Gamma_1 \times \Gamma_2$  such that, setting  $(\gamma_{1_k}, \gamma_{2_k}) := (\bar{\gamma}_1^{\frac{1}{k}}, \bar{\gamma}_2^{\frac{2}{k}})$ , one has

$$z_k := z + L^1_{\frac{1}{k}}(\gamma_{1_k}) \cdot \gamma_{1_k} + h^1_{\frac{1}{k}}(\gamma_{1_k}) = z + L^2_{\frac{1}{k}}(\gamma_{2_k}) \cdot \gamma_{2_k} + h^2_{\frac{1}{k}}(\gamma_{2_k}) \in F_1(\gamma_{1_k}) \cap F_2(\gamma_{2_k}).$$

Notice that, by  $h_{\frac{1}{k}}^1(\gamma_{1_k}) \leq \rho_1(\frac{1}{k}) \cdot \frac{1}{k}$ , and  $|\gamma_{1_k}| \leq \frac{1}{k}$  one has  $\lim_{k \to \infty} z_k = z$ , which concludes the proof.

### **3** Gaps and set-separation

#### **3.1** Original and extended controls

Let  $(\mathcal{M}, \langle \cdot, \cdot \rangle)$  be a Riemannian  $C^2$ -differentiable manifold, let [0, S] be a *time*-interval and let  $\mathfrak{W}$  be a metric space which we call the set of *control values*. For every  $(s, \mathfrak{w}) \in [0, S] \times \mathfrak{W}$ , let  $\mathcal{M} \ni y \mapsto (y, f(s, y, \mathfrak{w})) \in T\mathcal{M}$  be a vector field. We will consider *two families of controls*  $\mathcal{V}, \mathcal{W} := L^1([0, S], \mathfrak{W})$ , with  $\mathcal{V} \subset \mathcal{W}$ . We will call  $\mathcal{V}$  and  $\mathcal{W}$  the *original family of controls* and the *extended family of controls*, respectively.

Let us choose an *initial point*  $\bar{y} \in \mathcal{M}$ , and, for any *control* map  $w \in \mathcal{W}$ , let us consider the Cauchy problem

(E) 
$$\begin{cases} \frac{dy}{ds}(s) = f(s, y(s), w(s)) & \text{a.e. } s \in [0, S] \\ y(0) = \bar{y} \end{cases}$$

We shall assume the following regularity hypothesis: **Hypothesis (SH) :** 

- (i) for each  $(s, \mathfrak{w}) \in [0, S] \times \mathfrak{W}$ , the vector field  $y \mapsto f(s, y, \mathfrak{w})$  is of class  $C^1$  on  $\mathcal{M}$ ;
- (ii) there exists an integrable function  $c \in L^1([0, S]; \mathbb{R})$  such that, for a.e.  $s \in [0, S]$ ,

$$|f(s, y, \mathfrak{w})| \le c(s), \qquad |Df(s, y, \mathfrak{w})| \le c(s)$$
(3.10)

for every  $(y, \mathfrak{w}) \in \mathcal{M} \times \mathfrak{W}$ .

- (iii) for every  $(y, \mathfrak{w}) \in \mathcal{M} \times \mathfrak{W}$ , the map  $s \mapsto f(s, y, \mathfrak{w})$  is measurable;
- (iv) for every  $s \in [0, S]$ , the map  $(y, \mathfrak{w}) \mapsto f(s, y, \mathfrak{w})$  is continuous.

In particular, for every  $w \in \mathcal{W}$  there exists a unique trajectory y[w] of (E).

Let us fix a closed set  $\mathfrak{T} \subseteq \mathcal{M}$ , which we will refer to as *target*.

**Remark 3.1.** Of course, through standard cut-off arguments, in many situations one can replace (ii) in hypothesis (SH) with a weaker assumption concerning a neighbourhood of  $\hat{y}([0, S])$  istead of the whole state-space  $\mathcal{M}$ .

**Definition 3.1.** For any control  $v \in \mathcal{V}$  [resp.  $w \in \mathcal{W}$ ], the pair (y, v) := (y[v], v) [resp. (y, w) := (y[w], w)] will be called original process [resp. extended processes]. An extended process – in particular, an original process – (y, w) is called feasible if  $y(S) \in \mathfrak{T}$ .

### 3.2 Infimum gaps

Let us endow the set of controls  $\mathcal{W}$  with the pseudo-distance  $d_f$  defined as

$$d_f(w_1, w_2) := d_{\infty}(y[w_1], y[w_2]) \left( := \max_{s \in [0, S]} d(y[w_1](s), y[w_2](s)) \right),$$
(3.11)

for all controls  $w_1, w_2 \in \mathcal{W}$ .

The set

$$\mathcal{R}_{\mathcal{V}} := \left\{ y[v](S) : \quad v \in \mathcal{V} \right\} \subset \mathcal{M}$$
(3.12)

will be called the original reachable set, and the set

$$\mathcal{R}_{\mathcal{W}} := \left\{ y[w](S) : \quad w \in \mathcal{W} \right\} \subset \mathcal{M}$$
(3.13)

will be called the *extended reachable set*.

We will also consider local versions of the above reachable sets. Precisely, for a given extended process  $(\hat{y}, \hat{w})$  and  $r \ge 0$ , we set

$$\mathcal{R}_{\mathcal{V}}^{\hat{w},r} := \begin{cases} y[v](S) : & v \in \mathcal{V}, \quad d_f(\hat{w},v) < r \end{cases} \\ \mathcal{R}_{\mathcal{W}}^{\hat{w},r} := \begin{cases} y[w](S) : & w \in \mathcal{W}, \quad d_f(\hat{w},w) < r \end{cases}.$$

Clearly  $\mathcal{R}_{\mathcal{W}} \supseteq \mathcal{R}_{\mathcal{V}}$  and  $\mathcal{R}_{\mathcal{W}}^{\hat{w},r} \supseteq \mathcal{R}_{\mathcal{V}}^{\hat{w},r}$ , for all  $r \ge 0$ .

The occurrence of a local infimum gap is captured by the following definition:

**Definition 3.2.** Let  $(\hat{y}, \hat{w})$  be a feasible extended process such that  $\hat{y}(S) \in \mathcal{R}_{\mathcal{W}} \setminus \mathcal{R}_{\mathcal{V}}$ . We say that  $(\hat{y}, \hat{w})$  satisfies the infimum gap condition *if*, for any continuous cost function  $h : \mathbb{R}^n \to \mathbb{R}$ , there exists r > 0 such that one has

$$h(\hat{y}(S)) < \inf \left\{ h(y) : \quad y \in \mathcal{R}_{\mathcal{V}}^{\hat{w},r} \cap \mathfrak{T} \right\}$$
(3.14)

Despite the name, the infimum gap condition (3.14) is clearly a fully dynamical property. Actually, it could be as well rephrased in terms of 'supremum gap' or even independently of any optimization procedure as shown in Lemma 3.1 below.

**Definition 3.3.** Let  $(\hat{y}, \hat{w})$  be a feasible extended process such that  $\hat{y}(S) \in \mathcal{R}_{\mathcal{W}} \setminus \mathcal{R}_{\mathcal{V}}$ . We say that  $(\hat{y}, \hat{w})$  is isolated from  $\mathcal{V}$  if, for some r > 0 the sets  $\left(\mathcal{R}_{\mathcal{V}}^{\hat{w}, r} \cup \{\hat{y}(S)\}\right)$  and  $\mathfrak{T}$  are locally separated at  $\hat{y}(S)$ , namely, there exists a neighborhood  $\mathcal{N} \subset \mathcal{M}$  of  $\hat{y}(S)$  such that  $\left(\mathcal{R}_{\mathcal{V}}^{\hat{w}, r} \cup \{\hat{y}(S)\}\right) \cap \mathfrak{T} \cap \mathcal{N} = \{\hat{y}(S)\}.$ 

**Lemma 3.1.** Let  $(\hat{y}, \hat{w})$  be an extended feasible process such that  $\hat{y}(S) \in \mathcal{R}_{\mathcal{W}} \setminus \mathcal{R}_{\mathcal{V}}$ . Then the following conditions are equivalent:

- i)  $(\hat{y}, \hat{w})$  satisfies (3.14) for a given continuous cost function h and  $\hat{r} > 0$ ;
- *ii)* the process  $(\hat{y}, \hat{w})$  is isolated from  $\mathcal{V}$ ;
- iii) the process  $(\hat{y}, \hat{w})$  satisfies the infimum gap condition. Furthermore the right handside of (3.14) is equal to  $+\infty$ .

*Proof.* We give a proof just for the sake of completeness, all arguments being trivial.

Let us start proving that i) implies ii). This means that one has to show that there exists  $\hat{r} > 0$  such that

$$\mathcal{R}_{\mathcal{V}}^{w,r} \cap \mathfrak{T} = \emptyset \quad \forall r < \hat{r}.$$

$$(3.15)$$

Assume that (3.15) is false, which means that there exists a sequence  $r_n \downarrow 0$  such that  $\mathcal{R}_{\mathcal{V}}^{\hat{w},r_n} \cap \mathfrak{T} \neq \emptyset$  for all natural *n*. This implies that there exists a sequence  $(y_k)_{k \in \mathbb{N}}$  verifying

 $y_k \in \left(\mathcal{R}_{\mathcal{V}}^{\hat{w}, r_k} \cap \mathfrak{T}\right)$  for every  $k \in \mathbb{N}$ , so that  $y_n \to \hat{y}(S)$ , which, in view of the continuity of h, contradicts i). Hence, (3.15) holds true, from which we get ii).

Let us now prove that ii)  $\Rightarrow iii$ ). By hypothesis, there exists a neighborhood  $\mathcal{N}$  of  $\hat{y}(S)$  such that  $\left(\mathcal{R}_{\mathcal{V}}^{\hat{w},r} \cup \{\hat{y}(S)\}\right) \cap \mathfrak{T} \cap \mathcal{N} = \{\hat{y}(S)\}$ . Since  $\hat{y}(S) \in \mathcal{R}_{\mathcal{W}} \setminus \mathcal{R}_{\mathcal{V}}$ , by possibly reducing the size of r > 0 one obtains that  $\mathcal{R}_{\mathcal{V}}^{\hat{w},r} \cap \mathfrak{T} = \emptyset$ , which obviously implies iii), with the right hand-side of (3.14) equal to  $+\infty$ . Finally, the relation iii)  $\Rightarrow i$ ) is trivial.

4 Abundance

Our main results –namely Theorem 5.1 and Corollaries 5.1 and 5.2– strongly rely on a property introduced by J. Warga and called "abundance". It consists in a particular pervasiveness of  $\mathcal{V}$  in  $\mathcal{W}$ , which happens to be stronger than density. In fact, because of the presence of a closed final constraint, the mere density of  $\mathcal{R}_{\mathcal{V}}$  into  $\mathcal{R}_{\mathcal{W}}$  is not enough in order to normality to be a sufficient condition for gaps' avoidance (see Subsection (A.1)). We will make use of a generalization of abundance provided in [21] and we will extend it to manifolds.

For every positive integer N, let  $\Gamma_N$  be the convex hull of the union of the origin with the N-simplex, namely

$$\Gamma_N := \Big\{ \gamma = (\gamma^1, ..., \gamma^N) \in \mathbb{R}^N : \sum_{j=1}^N \gamma^j \le 1, \ \gamma^j \ge 0, \ j = 1, ..., N \Big\}.$$

For any  $\gamma \in \Gamma_N$ , let us consider the control system on  $\mathcal{M}$ 

$$\begin{cases} \frac{dy}{ds}(s) = f_{\gamma}\left(s, y(s), w(s), w_{1}(s), ..., w_{N}(s)\right) \\ y(0) = \bar{y}, \end{cases}$$
(4.16)

where: i) the control  $(w, w_1, ..., w_N)$  belongs to  $\mathcal{W}^{1+N}$ , and ii) the vector field  $f_{\gamma}$  is defined by setting, for every  $(s, y) \in [0, S] \times \mathcal{M}$  and  $(\mathfrak{w}, \mathfrak{w}_1, ..., \mathfrak{w}_N) \in \mathfrak{W}^{1+N}$ ,

$$f_{\gamma}(s, y, \mathfrak{w}, (\mathfrak{w}_1, ..., \mathfrak{w}_N)) := f(s, y, \mathfrak{w}) + \sum_{i=1}^N \gamma^i \Big( f(s, y, \mathfrak{w}_i) - f(s, y, \mathfrak{w}) \Big).$$

For every value of the parameter  $\gamma \in \Gamma_N$  and every control  $(w, w_1, ..., w_N) \in \mathcal{W}^{1+N}$ , let us use  $y_{\gamma} \left[ w, w_1, ..., w_N \right]$  to denote the corresponding solution of (4.16).<sup>7</sup> Notice, in particular, that  $y[w] = y_{\gamma} \left[ w, w, ..., w \right]$  for all  $w \in \mathcal{W}$  and for all  $\gamma \in \Gamma_N$ .

**Definition 4.1.** [21] We say that a subclass of controls  $\mathcal{V} \subset \mathcal{W}$  is abundant in  $\mathcal{W}$  if, for every integer N, every (1+N)-tuple of controls  $(w, w_1, ..., w_N) \in \mathcal{W}^{1+N}$ , and every  $\delta > 0$ , there exists a continuous mapping  $\theta_{w,w_1,...,w_N}^{\delta} : \Gamma_N \to \mathcal{V}$  such that

$$d\Big(y_{\gamma}\Big[w, w_1, ..., w_N\Big](S), y\Big[\theta_{w, w_1, ..., w_N}^{\delta}(\gamma)\Big](S)\Big) < \delta, \qquad \forall \gamma \in \Gamma_N.$$

$$(4.17)$$

A sufficient condition for abundance, based on *concatenation*, is given in Proposition 4.1 below.

<sup>&</sup>lt;sup>7</sup> Under hipothesis (SH) such a solution exists and is unique.

**Definition 4.2.** We say that a set of controls  $\mathcal{V}$  satisfies the concatenation property if, for every  $\bar{s} \in ]0, S[$  and for any  $v_1, v_2 \in \mathcal{V}$ , one has  $v_1\chi_{[0,\bar{s}[} + v_2\chi_{[\bar{s},S]} \in \mathcal{V}, {}^8$  where we have used  $\chi_E$  to denote the indicator function of a subset  $E \subseteq [0, S]$ 

**Proposition 4.1.** [21] Assume that the subfamily  $\mathcal{V} \subset \mathcal{W}$  satisfies the concatenation property and is dense in  $\mathcal{W}$  with respect to the pseudo-metric  $d_f$ . Then  $\mathcal{V}$  is an abundant subset of  $\mathcal{W}$ .

The proof of this result for the special case when  $\mathcal{M} = \mathbb{R}^n$  was given in ([21], Theorem IV.3.9) by developing some arguments in [18]. The required, obvious, changes to prove the result on a Riemannian manifold reduce to a reformulation of estimate (4.17) in local coordinates, so we omit them.

### 4.1 Approximating the original reachable set by extended cones

Let us fix a a feasible extended process  $(\hat{y}, \hat{w})$ , and, for any  $s, \check{s} \in [0, S]$ ,  $s > \check{s}$ , let  $M(s, \check{s}) : T_{\hat{y}(\check{s})}\mathcal{M} \to T_{\hat{y}(s)}\mathcal{M}$  denote the differential of the diffeomorphism established by the differential equation  $\dot{y} = f(s, y, \hat{w})$  from a neighborhood of  $\hat{y}(\check{s})$  to a neighborhood of  $\hat{y}(s)$ . As is known,  $s \to M(s, \check{s})$  is the solution of the *variational* Cauchy problem having the following coordinate representation:

$$\frac{dM}{ds}(s) = \frac{\partial f}{\partial y}(s, \hat{y}(s), \hat{w}(s)) \circ M(s), \qquad M(\check{s}, \check{s}) = id_{T_{\hat{y}(\check{s})}\mathcal{M}}.$$
(4.18)

**Definition 4.3.** Consider a positive integer N, N control values  $\mathfrak{w}_1, ..., \mathfrak{w}_N \in \mathfrak{W}$ , and N instants  $s_1, ..., s_N \in \mathrm{SD}\{f(\cdot, \cdot, \hat{w}(\cdot))\} \cap \mathrm{SD}\{f(\cdot, \cdot, \mathfrak{w}_1)\} \cap ... \cap \mathrm{SD}\{f(\cdot, \cdot, \mathfrak{w}_N)\}, 9 \ 0 < s_1 < ..., < s_N \leq S$ . The convex cone

$$\mathbf{C}_{\mathfrak{w}_{1},\ldots,\mathfrak{w}_{N}}^{s_{1},\ldots,s_{N}} = span^{+} \left\{ M(S,s_{i}) \cdot \left( f(s_{i},\hat{y}(s_{i}),\mathfrak{w}_{i}) - f(s_{i},\hat{y}(s_{i}),\hat{w}(s_{i})) \right) : i = 1,\ldots,N \right\} \subset T_{\hat{y}(S)} \mathcal{M}_{i}$$

will be called extended variational cone corresponding to the feasible extended process  $(\hat{y}, \hat{w})$ .

The following result can be regarded as claiming a sort of *infinitesimal thickness* of  $\mathcal{V}$  in  $\mathcal{W}$ .

**Theorem 4.1.** Let the original family of controls  $\mathcal{V} \subset \mathcal{W}$  be abundant in  $\mathcal{W}$ , and let a feasible extended process  $(\hat{y}, \hat{w})$  be given. Consider a positive integer N, N control values  $\mathfrak{w}_1, ..., \mathfrak{w}_N \in \mathfrak{W}$ , and N instants  $s_1, ..., s_N \in \mathrm{SD}\{f(\cdot, \cdot, \hat{w}(\cdot))\} \cap \mathrm{SD}\{f(\cdot, \cdot, \mathfrak{w}_1)\} \cap ... \cap \mathrm{SD}\{f(\cdot, \cdot, \mathfrak{w}_N)\}, 0 < s_1 < ..., < s_N \leq S$ . Then, for any r > 0, the extended variational cones  $\mathbf{C}^{s_1, ..., s_N}_{\mathfrak{W}_1, ..., \mathfrak{W}_N}$  is a QDQ approximating cone to  $\mathcal{R}^{\hat{w}, r}_{\mathcal{V}} \cup \{\hat{y}(S)\}$  at  $\hat{y}(S)$ .

**Remark 4.1.** While the fact that  $\mathbf{C}_{\mathfrak{w}_1,\ldots,\mathfrak{w}_N}^{s_1,\ldots,s_N}$  is a QDQ approximating cone to the extended reachable set  $\mathcal{R}_{\mathcal{W}}^{\hat{w},r}$  at  $\hat{y}(S)$  (for any r > 0) is a classical argument, utilized in the proof of the Maximum Principle,<sup>10</sup> the fact that  $\mathbf{C}_{\mathfrak{w}_1,\ldots,\mathfrak{w}_N}^{s_1,\ldots,s_N}$  is a first order approximation for the small reachable set  $\mathcal{R}_{\mathcal{V}}^{\hat{w},r}$  is anything but obvious: it means, in a sense, that this cone is not too large.

$$v_1\chi_E + v_2\chi_{([0,S]\setminus E)} \in \mathcal{S}.$$

<sup>&</sup>lt;sup>8</sup>Concatenation is weaker than *decomposability* of a set S of paths on an interval [a, b] [16, 23, 30], which prescribes that for any pair of paths  $v_1, v_2 \in S$  and any measurable set  $E \subset [0, S]$ , one has

<sup>&</sup>lt;sup>9</sup>We have used the notation  $SD(\phi)(\cdot, \cdot)$  –introduced in Subsection 1.1– to mean the (full measure) Scorza-Dragoni set of a function  $\phi = \phi(s, y)$ 

<sup>&</sup>lt;sup>10</sup>Actually the same is true for other, more classical, cones, e.g. the Boltyanski cone and the regular tangent cone.

Proof of Theorem 4.1. We will prove this theorem assuming that  $\mathcal{M}$  is an open subset of  $\mathbb{R}^n$ , so that we can identify  $T_{\hat{y}(S)}\mathcal{M}$  with  $\mathbb{R}^n$ . Clearly, this is not restrictive because of the local character of the result.

Let  $\mathbf{C}_{\mathfrak{w}_1,...,\mathfrak{w}_N}^{s_1,...,s_N}$  be an extended variational cone. Let us set  $s_0 = 0$  and, for every i = 1, ..., N, consider a number  $\delta_i \leq s_i - s_{i-1}$  and the control

$$w_i^{\delta^i}(s) := \begin{cases} \hat{w}(s), & \forall s \in [0, S] \setminus [s_i - \delta^i, s_i] \\ \mathfrak{w}_i & \forall s \in [s_i - \delta^i, s_i]. \end{cases}$$
(4.19)

Let us set  $\overline{\delta} := \frac{N}{2} \min\{s_i - s_{i-1}, i = 1, \dots, N\}$ . Let us define the set-valued map  $F : \mathbb{R}^N \rightsquigarrow \mathbb{R}^n$  as

$$F(\epsilon) = \left\{ y \Big[ \theta_{\hat{w}, w_1^{\delta/N}, \dots, w_N^{\delta/N}}^{\delta^2} \left( \pi \left( \frac{\epsilon}{\delta} \right) \right) \Big](S) : \qquad 0 < \delta \le \bar{\delta} \right\} \quad \forall \epsilon \in \mathbb{R}^N, \tag{4.20}$$

where  $\pi : \mathbb{R}^N \to \Gamma_N$  denotes the orthogonal projection on  $\Gamma_N$  (which, because of the convexity of  $\Gamma_N$ , is a continuous, single-valued, map). Notice that, by construction  $F(\epsilon) \subseteq \mathcal{R}_{\mathcal{V}}$ , for every  $\epsilon \in \mathbb{R}^N$ .

For each  $\delta \in [0, \overline{\delta}]$  and  $\epsilon \in \Gamma_N \cap B_{\delta}$ , let us choose  $\gamma = (\gamma^1, \dots, \gamma^N) := (\frac{\epsilon^1}{\delta}, \dots, \frac{\epsilon^N}{\delta}) \in \Gamma_N$ . From (4.24) in Lemma 4.1 below it follows that

$$y_{\gamma}[\hat{w}, w_{1}^{\delta/N}, \dots, w_{N}^{\delta/N}](S) = y_{\epsilon/\delta}[\hat{w}, w_{1}^{\delta/N}, \dots, w_{N}^{\delta/N}](S) =$$
  
=  $\hat{y}(S) + \frac{1}{N} \sum_{i=1}^{N} \epsilon^{i} M(S, s_{i}) \cdot \left( f(s_{i}, \hat{y}(s_{i}), \mathfrak{w}_{i}) - f(s_{i}, \hat{y}(s_{i}), \hat{w}(s_{i})) + \phi(\epsilon, \delta), \right)$   
(4.21)

for every  $\epsilon \in \Gamma_N \cap B_{\delta}$ , where  $\phi : \bigcup_{\substack{0 < \delta \le \overline{\delta} \\ 0 < \delta \le \overline{\delta}}} ((\Gamma_N \cap B_{\delta}) \times \{\delta\}) \to \mathbb{R}^n$  is a continuous function

which verifies  $\max \{ |\phi(\epsilon, \delta)|, \ \epsilon \in \Gamma_N \cap B_{\delta} \} = o(\delta)$ . In view of the abundance property, for each  $\delta \in ]0, \bar{\delta}]$  and  $\epsilon \in \Gamma_N \cap B_{\delta}$ , there exists  $\tilde{\phi} : \bigcup_{0 < \delta \leq \bar{\delta}} ((\Gamma_N \cap B_{\delta}) \times \{\delta\}) \to \mathbb{R}^n$  such that

$$y\left[\theta_{\hat{w},w_1^{\delta/N},\ldots,w_N^{\delta/N}}^{\delta^2}\left(\frac{\epsilon}{\delta}\right)\right](S) - y_{\epsilon/\delta}[\hat{w},w_1^{\delta/N},\ldots,w_N^{\delta/N}](S) = \tilde{\phi}(\epsilon,\delta)$$

for all  $\epsilon \in \Gamma_N \cap B_{\delta}$ , with  $\max \{ |\tilde{\phi}(\epsilon, \delta)|, \epsilon \in \Gamma_N \cap B_{\delta} \} \leq \delta^2$ . Therefore

$$y \left[ \theta_{\hat{w}, w_1^{\delta/N}, \dots, w_N^{\delta/N}}^{\delta^2} \left( \frac{\epsilon}{\delta} \right) \right] (S) =$$
  
=  $\hat{y}(S) + \frac{1}{N} \sum_{i=1}^{N} \epsilon^i M(S, s_i) \cdot \left( f(s_i, \hat{y}(s_i), \mathfrak{w}_i) - f(s_i, \hat{y}(s_i), \hat{w}(s_i)) + h^{\delta}(\epsilon), \right)$ (4.22)

where  $h^{\delta}(\epsilon) := \phi(\epsilon, \delta) + \tilde{\phi}(\epsilon, \delta)$ . Observe that

$$|h^{\delta}(\epsilon)| \le \delta\rho(\delta), \quad \forall \epsilon \in B_{\delta}, \tag{4.23}$$

where we have set  $\rho(\delta) := \frac{\max\{|\phi(\epsilon, \delta)|, \epsilon \in \Gamma_N \cap B_\delta\}}{\delta} + \delta.$ 

For every  $\delta \in ]0, \bar{\delta}]$ , let us define the map

$$A^{\delta}: \Gamma_N \cap B_{\delta} \to Lin(\mathbb{R}^N, \mathbb{R}^n) \times \mathbb{R}^n \\ \epsilon \mapsto A^{\delta}(\epsilon) := (L, h^{\delta}(\epsilon)),$$

where L is the linear map defined as

$$L \cdot b = \frac{1}{N} \sum_{i=1}^{N} b^{i} M(S, s_{i}) \cdot \left( f(s_{i}, \hat{y}(s_{i}), \mathfrak{w}_{i}) - f(s_{i}, \hat{y}(s_{i}), \hat{w}(s_{i})) \right), \quad \forall b \in \mathbb{R}^{N}.$$

Notice that, because of the continuity w.r.t.  $\epsilon$  of the left-hand side of (4.22), for every  $\delta > 0$ , the map  $\epsilon \mapsto A^{\delta}(\epsilon)$  is continuous. By rewriting relation (4.22) as

$$y\left[\theta_{\hat{w},w_1^{\delta/N},\ldots,w_N^{\delta/N}}^{\delta^2}\left(\frac{\epsilon}{\delta}\right)\right](S) = \hat{y}(S) + L \cdot \epsilon + h^{\delta}(\epsilon),$$

we get

$$\hat{y}(S) + L \cdot \epsilon + h^{\delta}(\epsilon) \in F(\epsilon),$$

which means that L is a QDQ of F at  $(0, \hat{y}(S))$  in the direction of the set  $\Gamma_N$ . Therefore,  $\Lambda$  is also a QDQ of F at  $(0, \hat{y}(S))$  in the direction of the cone  $\Gamma = [0, +\infty[^N$ . Since  $\mathbf{C}_{\mathfrak{w}_1,\ldots,\mathfrak{w}_N}^{s_1,\ldots,s_N} = L \cdot \Gamma$ , one concludes that  $\mathbf{C}_{\mathfrak{w}_1,\ldots,\mathfrak{w}_N}^{s_1,\ldots,s_N}$  is a QDQ approximating cone to  $\mathcal{R}_{\mathcal{V}} \cup \{\hat{y}(S)\}$  at  $\hat{y}(S)$ .

**Lemma 4.1.** Fix  $\gamma \in \Gamma_N$ . Then, the map  $\epsilon \to y_{\gamma} \Big[ \hat{w}, w_1^{\epsilon^1}, ..., w_N^{\epsilon^N} \Big](S)$  verifies

$$y_{\gamma} \left[ \hat{w}, w_{1}^{\epsilon^{i}}, ..., w_{N}^{\epsilon^{N}} \right] (S) - \hat{y}(S) = \sum_{i=1}^{N} \gamma^{i} \epsilon^{i} M(S, s_{i}) \cdot \left( f(s_{i}, \hat{y}(s_{i}), \mathfrak{w}_{i}) - f(s_{i}, \hat{y}(s_{i}), \hat{w}(s_{i})) \right) + \phi(\epsilon, \gamma),$$
(4.24)

where  $\phi$  is a continuous function verifying  $\max \{\phi(\epsilon, \gamma) : \gamma \in \Gamma_N\} = o(\epsilon)$ .

*Proof.* Let us begin proving the lemma in the case when N = 1 and  $0 \le \gamma \le 1$ . One has

$$y_{\gamma} \left[ \hat{w}, w_{1}^{\epsilon^{1}} \right] (s_{1}) - \hat{y}(s_{1}) = y_{\gamma} \left[ \hat{w}, w_{1}^{\epsilon^{1}} \right] (s_{1} - \epsilon^{1}) - \hat{y}(s_{1} - \epsilon^{1}) + \int_{s_{1} - \epsilon^{1}}^{s_{1}} \left( f_{\gamma} \left( s, y_{\gamma} [\hat{w}, w_{1}^{\epsilon^{1}}](s), \hat{w}(s), \mathfrak{w}_{1} \right) - f(s, \hat{y}(s), \hat{w}(s)) \right) ds = \int_{s_{1} - \epsilon^{1}}^{s_{1} - \epsilon^{1}} \left( f_{\gamma} \left( s, y_{\gamma} [\hat{w}, w_{1}^{\epsilon^{1}}](s), \hat{w}(s), \mathfrak{w}_{1} \right) - f(s, \hat{y}(s), \hat{w}(s)) \right) ds = \epsilon^{1} \left( f_{\gamma} \left( s_{1}, \hat{y}(s_{1}), \hat{w}(s_{1}), \mathfrak{w}_{1} \right) - f(s_{1}, \hat{y}(s_{1}), \hat{w}(s_{1})) \right) + \Phi_{1}(\epsilon^{1}, \gamma) + \Phi_{2}(\epsilon^{1}) = \gamma \epsilon^{1} \cdot \left( f \left( s_{1}, \hat{y}(s_{1}), \hat{\mathfrak{w}}_{1} \right) - f \left( s_{1}, \hat{y}(s_{1}), \hat{w}(s_{1}) \right) \right) + \Phi_{1}(\epsilon^{1}, \gamma) + \Phi_{2}(\epsilon^{1}),$$

$$(4.25)$$

where

$$\begin{split} \Phi_1(\epsilon^1,\gamma) &:= \int_{s_1-\epsilon^1}^{s_1} \left( f_\gamma\left(s, y_\gamma[\hat{w}, w_1^{\epsilon^1}](s), \hat{w}(s), \mathfrak{w}_1\right) - f_\gamma\left(s_1, \hat{y}(s_1), \hat{w}(s_1), \mathfrak{w}_1\right) \right) ds, \\ \Phi_2(\epsilon^1) &:= \int_{s_1-\epsilon^1}^{s_1} \left( f(s_1, \hat{y}(s_1), \hat{w}(s_1)) - f(s, \hat{y}(s), \hat{w}(s)) \right) ds. \end{split}$$

To simplify the notation, in what follows we will write  $y_{\gamma}(s)$  in place of  $y_{\gamma}[\hat{w}, w_1^{\epsilon^1}](s)$ . Using hypothesis **(SH)**-(ii), one obtains the following estimate:

$$\begin{aligned} |y_{\gamma}(s_{1}) - \hat{y}(s_{1})| &\leq \int_{s_{1}-\epsilon^{1}}^{s_{1}} |f(s, y_{\gamma}(s), \hat{w}(s)) - f(s, \hat{y}(s), \hat{w}(s))| \, ds + \\ \gamma \int_{s_{1}-\epsilon^{1}}^{s_{1}} |f(s, y_{\gamma}(s), \mathfrak{w}_{1}) - f(s, y_{\gamma}(s), \hat{w}(s)) + f(s, \hat{y}(s), \hat{w}(s)) - f(s, \hat{y}(s), \mathfrak{w}_{1})| \, ds + \\ \gamma \int_{s_{1}-\epsilon^{1}}^{s_{1}-\epsilon^{1}} |f(s, \hat{y}(s), \mathfrak{w}_{1}) - f(s, \hat{y}(s), \hat{w}(s))| \, ds \leq \\ (1+2\gamma) \int_{s_{1}-\epsilon^{1}}^{s_{1}} c(s) \, |y_{\gamma}(s) - \hat{y}(s)| \, ds + \gamma \int_{s_{1}-\epsilon^{1}}^{s_{1}} |f(s, \hat{y}(s), \mathfrak{w}_{1}) - f(s, \hat{y}(s), \hat{w}(s))| \, ds. \end{aligned}$$

$$(4.26)$$

Setting, for  $s \in [s_1 - \epsilon^1, s_1]$ ,

$$\alpha(s) = \gamma \int_{s_1 - \epsilon^1}^s |f(s, \hat{y}(s), \mathfrak{w}_1) - f(s, \hat{y}(s), \hat{w}(s))| \, ds, \tag{4.27}$$

it follows from the Gronwall's Lemma that

$$|y_{\gamma}(s_{1}) - \hat{y}(s_{1})| \leq \alpha(s_{1}) + \int_{s_{1}-\epsilon^{1}}^{s_{1}} (1+2\gamma)c(s)\exp\{(1+2\gamma)c(s)(s_{1}-s)\}\alpha(s)ds \leq \\ \leq 2\gamma \int_{s_{1}-\epsilon^{1}}^{s_{1}} c(s)ds + o(\epsilon^{1}) \to 0$$
(4.28)

when  $\epsilon^1 \to 0$ . Therefore,

$$|y_{\gamma}(s) - \hat{y}(s_{1})| \le |y_{\gamma*}(s) - y_{\gamma*}(s_{1})| + |y_{\gamma*}(s_{1}) - \hat{y}(s_{1})| \le (1 + 4\gamma) \int_{s_{1} - \epsilon^{1}}^{s_{1}} c(s)ds + o(\epsilon^{1}).$$
(4.29)

Since

- $s_1$  is a Scorza-Dragoni point of  $f(\cdot, \cdot, \hat{w}(\cdot))$  and  $f(\cdot, \cdot, \mathfrak{w}_1)$ , and
- the maps  $y \mapsto f(s, y, \hat{w}(s))$ ,  $y \mapsto f(s, y, \mathfrak{w}_1)$  are Lipschitz continuous in a neighbourhood of  $\hat{y}([0, S])$ ,

in view of (4.28), (4.29), one easily gets

$$\max\left\{\Phi_1(\epsilon^1, \gamma): \ 0 \le \gamma \le 1\right\} = o(\epsilon^1), \qquad \Phi_2(\epsilon^1) = o(\epsilon^1). \tag{4.30}$$

If we set  $\phi(\epsilon^1, \gamma) := \Phi_1(\epsilon^1, \gamma) + \Phi_2(\epsilon^1)$ , it follows by estimates (4.25) and (4.30) that

$$y_{\gamma} \left[ \hat{w}, w_{1}^{\epsilon^{1}} \right] (s_{1}) - \hat{y}(s_{1}) = \gamma \epsilon^{1} \left( f \left( s_{1}, \hat{y}(s_{1}), \hat{\mathfrak{w}}_{1} \right) - f \left( s_{1}, \hat{y}(s_{1}), \hat{w}(s_{1}) \right) \right) + \phi(\epsilon^{1}, \gamma).$$

Hence, one has

$$\frac{d}{d\epsilon^{1}}y_{\gamma}\left[\hat{w}, w_{1}^{\epsilon^{1}}\right](s_{1})_{|\epsilon^{1}=0} = \gamma\left(f\left(s_{1}, \hat{y}(s_{1}), \hat{\mathfrak{w}}_{1}\right) - f\left(s_{1}, \hat{y}(s_{1}), \hat{w}(s_{1})\right)\right),$$

which, by the basic theory of linear ODE's, implies that

$$\frac{d}{d\epsilon^1} y_{\gamma} \left[ \hat{w}, w_1^{\epsilon^1} \right] (S)_{|\epsilon^1 = 0} = M(S, s_1) \cdot \frac{d}{d\epsilon^1} y_{\gamma} \left[ \hat{w}, w_1^{\epsilon^1} \right] (s_1)_{|\epsilon^1 = 0} = M(S, s_1) \cdot \gamma \Big( f\left(s_1, \hat{y}(s_1), \hat{\mathfrak{w}}_1\right) - f\left(s_1, \hat{y}(s_1), \hat{w}(s_1)\right) \Big).$$

Therefore, the lemma is proved for N = 1 and for any  $0 \le \gamma \le 1$ . The general case  $N \ge 2$  is easily obtained by a finite induction argument. The latter doesn't present any new difficulty with respect to the proof of the case N = 1. Moreover is almost verbatim the one utilized in the proof of the Pontryagin Maximum Principle when passing from single to multiple, finitely many needle variations (see e.g. [35], Theorem 4.2.1). For this reason we omit it.

The reason why we have adopted QDQ approximating cones as tangential objects relies on the validity of the following result.  $^{11}$ 

**Theorem 4.2.** Let the original family of controls  $\mathcal{V} \subset \mathcal{W}$  be abundant in  $\mathcal{W}$ , and let a feasible extended process  $(\hat{y}, \hat{w})$  be given. Consider a positive integer N, N control values  $\mathfrak{w}_1, ..., \mathfrak{w}_N \in \mathfrak{W}$ , and N instants  $s_1, ..., s_N \in \mathrm{SD}\{f(\cdot, \cdot, \hat{w}(\cdot))\} \cap \mathrm{SD}\{f(\cdot, \cdot, \mathfrak{w}_1)\} \cap \ldots \cap \mathrm{SD}\{f(\cdot, \cdot, \mathfrak{w}_N)\}, 0 < s_1 < \ldots, < s_N \leq S$ . Moreover, let C be a QDQ approximating

<sup>&</sup>lt;sup>11</sup>Such a result would be not true if we chose to utilize AGDQ approximating cones

cone to the target  $\mathfrak{T}$  at  $\hat{y}(S)$ . If  $\mathbf{C}_{\mathfrak{w}_1,\ldots,\mathfrak{w}_N}^{s_1,\ldots,s_N}$  and C, are complementary subspaces, i.e.  $\mathbf{C}_{\mathfrak{w}_1,\ldots,\mathfrak{w}_N}^{s_1,\ldots,s_N} \oplus C = T_{\hat{y}(S)}\mathcal{M}$ , then there exists a sequence  $(z_k)_{k\in\mathbb{N}} \subset \mathcal{R}_{\mathcal{V}} \cap \mathfrak{T}$  such that

$$\lim_{k \to \infty} z_k = \hat{y}(S)$$

Proof. In view of Theorem 4.1,  $\mathbf{C}_{\mathfrak{w}_1,\ldots,\mathfrak{w}_N}^{s_1,\ldots,s_N}$  is a QDQ approximating cone to  $\mathcal{R}_{\mathcal{V}} \cup \{\hat{y}(S)\}$  at  $\hat{y}(S)$ . Furthermore, since C is a QDQ approximating cone to the target  $\mathfrak{T}$  at  $\hat{y}(S)$ , there exist a positive integer M, a set-valued map  $G : \mathbb{R}^M \to \mathcal{M}$ , a convex cone  $\Gamma \subset \mathbb{R}^M$ , and a Quasi Differential Quotient L of G at  $(0, \hat{y}(S))$  in the direction of  $\Gamma$  such that  $G(\Gamma) \subseteq \mathfrak{T}$  and  $C = L \cdot \Gamma$ . In order to conclude the proof, it is enough to apply Theorem 2.3, ii), with  $C_1 = \mathbf{C}_{\mathfrak{w}_1,\ldots,\mathfrak{w}_N}^{s_1,\ldots,s_N}$ ,  $C_2 = C$ ,  $N_1 = N$ ,  $N_2 = M$ ,  $\Gamma_1 = [0,\infty)^N$ ,  $\Gamma_2 = \Gamma$ ,  $F_1$  defined as in (4.20), and  $F_2 = G$ . This concludes the proof.

### 5 The main result

**Theorem 5.1** (A GEOMETRIC PRINCIPLE FOR GAPS). Let us assume that the family of controls  $\mathcal{V}$  is abundant in  $\mathcal{W}$ . Let  $(\hat{y}, \hat{w})$  be a feasible extended process satisfying the infimum gap condition. Then any QDQ approximating cone C to  $\mathfrak{T}$  at  $\hat{y}(S)$  is linearly separable from any extended variational cone  $\mathbf{C}_{\mathfrak{w}_1,\ldots,\mathfrak{w}_N}^{s_1,\ldots,s_N}$ , i.e. there exists a non-zero linear form  $\xi \in T^*_{\hat{u}(S)}\mathcal{M}$  such that

$$\xi \cdot c_1 \le 0 \le \xi \cdot c_2, \qquad \forall (c_1, c_2) \in \mathbf{C}^{s_1, \dots, s_N}_{\mathfrak{w}_1, \dots, \mathfrak{w}_N} \times C.$$

Let us give the definition of abnormal extremal, normal h-extremal, and h-abnormal extremal.

**Definition 5.1** (Abnormal extremal). Let  $(\hat{y}, \hat{w})$  be a feasible extended process, and let C be a QDQ approximating cone to the target  $\mathfrak{T}$  at  $\hat{y}(S)$ . We say that the process  $(\hat{y}, \hat{w})$  is an abnormal extremal (with respect to C) if there exists a lift  $(\hat{y}, \lambda) \in W^{1,1}([0, S]; T^*\mathcal{M})$  of  $\hat{y}$  verifying the following conditions:

- (i)  $\frac{d\lambda}{ds} = -\lambda \cdot \frac{\partial f}{\partial y}(s, \hat{y}(s), \hat{w}(s));$
- (*ii*)  $\max_{\mathfrak{w}\in\mathfrak{M}}\lambda(s)\cdot f(s,\hat{y}(s),\mathfrak{w}) = \lambda(s)\cdot f(s,\hat{y}(s),\hat{w}(s)) \quad a.e. \ s \in [0,S];$
- (*iii*)  $\lambda(S) \in -C^{\perp};$
- (*iv*)  $\lambda \neq 0$ .

**Definition 5.2** (*h*-extremal). Let  $(\hat{y}, \hat{w})$  be a feasible extended process, let a (cost) function  $h : \mathcal{M} \to \mathbb{R}$  be differentiable at  $\hat{y}(S)$ , and let C be a QDQ approximating cone of the target  $\mathfrak{T}$  at  $\hat{y}(S)$ . We say that the process  $(\hat{y}, \hat{w})$  is an *h*-extremal (with respect to *h* and *C*) if there exist a lift  $(\hat{y}, \lambda) \in W^{1,1}([0, S]; T^*\mathcal{M})$  of  $\hat{y}(\cdot)$  and a cost multiplier  $\lambda_c \in \{0, 1\}$  such that:

- (i)  $\frac{d\lambda}{ds} = -\lambda \cdot \frac{\partial f}{\partial y}(s, \hat{y}(s), \hat{w}(s));$
- (ii)  $\max_{\mathfrak{m}\in\mathfrak{M}}\lambda(s)\cdot f(s,\hat{y}(s),\mathfrak{m}) = \lambda(s)\cdot f(s,\hat{y}(s),\hat{w}(s)) \quad a.e. \ s \in [0,S];$
- (*iii*)  $\lambda(S) \in -\lambda_c \nabla h(\hat{y}(S)) C^{\perp};$
- $(iv) \quad (\lambda, \lambda_c) \neq 0.$

Furthermore, we say that an h-extremal  $(\hat{y}, \hat{w})$  is normal if for every choice of the pair  $(\lambda, \lambda_c)$ , one has  $\lambda_c = 1$ . We say that an h-extremal  $(\hat{y}, \hat{w})$  is abnormal if it is not normal, namely if exists a choice of  $(\lambda, \lambda_c)$  with  $\lambda_c = 0$ .

**Remark 5.1.** Though these definitions have intrinsic meanings, we have chosen to adopt a notation reminiscent of coordinates. Of course, the adjoint equation (i) might be expressed –when coupled with the dynamics– as the Hamiltonian system

$$\frac{d}{dt}(y,\lambda) = X_H(s,x,\lambda) := J \cdot DH(s,x,\lambda),$$

where  $H: T^*\mathcal{M} \to \mathbb{R}$  is the maximized Hamiltonian defined by setting

$$H(s, x, \lambda) := \max_{\mathfrak{m} \in \mathfrak{M}} \lambda(s) \cdot f(s, x, \mathfrak{m}) \qquad \forall (s, x, \lambda) \in [0, S] \times T^* \mathcal{M}$$

and  $X_H$  is the Hamiltonian vector field, namely  $X_H := J \cdot DH$ , J being the symplectic matrix and D the differential operator with respect to x and  $\lambda$ .

Let us also point out that the dot  $\cdot$  has obvious different meanings according to the context: for instance, in (i) of the definitions above, it denotes a linear operator on the cotangent space, while in (ii) it denotes the duality product.

Observe that every abnormal extremal is an abnormal *h*-extremal for any cost *h* differentiable at  $\hat{y}(S)$ , while every abnormal *h*-extremal is an abnormal extremal. We are now ready to state our main result on infimum gaps.

**Corollary 5.1** (NORMALITY NO-GAP CRITERION). Let us assume that the family of controls  $\mathcal{V}$  is abundant in  $\mathcal{W}$ . If a feasible extended process  $(\hat{y}, \hat{w})$  satisfies the infimum gap condition, then, for every QDQ approximating cone C to  $\mathfrak{T}$  at  $\hat{y}(S)$ ,  $(\hat{y}, \hat{w})$  is an abnormal extremal with respect to C.

When referred to a specific cost h, the contrapositive version of this theorem provides a sufficient condition for the absence of local infimum gaps. Precisely:

**Corollary 5.2** (A SUFFICIENT CONDITION FOR AVOIDING INFIMUM GAPS). Let us assume that the family of controls  $\mathcal{V}$  is abundant in  $\mathcal{W}$ , and let  $(\hat{y}, \hat{w})$  be a feasible extended process. Let  $h : \mathcal{M} \to \mathbb{R}$  be a cost function, differentiable at  $\hat{y}(S)$ , and let  $(\hat{y}, \hat{w})$  be a normal hextremal for some QDQ approximating cone C to  $\mathfrak{T}$  at  $\hat{y}(S)$ . Then there is no infimum gap at  $(\hat{y}, \hat{w})$ .

As we have mentioned in the Introduction, the relation between gap phenomena and abnormality has been quite investigated in two cases of embeddings: the embedding of bounded optimal control problems into their *convex relaxation* [32, 33, 34] and the embedding of unbounded (convex) control systems into their *impulsive, space-time closure* [28]. Since the original control families in such embeddings turn out to be abundant in their extensions, these kinds of results can be also obtained by Theorem 5.1.<sup>12</sup> In Section 7, we are going to present a new application to a dynamics which is neither convex nor bounded.

### 5.1 A verifiable sufficient condition for normality

In practical situations, it may be difficult or even impossible to directly verify the normality of an extremal, which, in view of Corollary 5.2, would guarantee the absence of gaps. This motivates Theorem 5.2 below, which provides a sufficient condition on the data of the problem in order for the extremals to be normal.

In the following definition we assume that a control system

$$\frac{dy}{ds}(s) = f(s, y(s), w(s)), \qquad w \in \mathcal{W}, \qquad \text{a.e.} \quad s \in [0, S], \tag{5.31}$$

as above is given, with an initial condition

$$y(0) = \bar{y},\tag{5.32}$$

and, still, we use  $\mathcal{R}_{\mathcal{W}}$  to denote the reachable set from  $\bar{y}$ .

 $<sup>^{12}</sup>$ Although the use of different types of cones describing the non-transversality condition makes Theorem 5.1 and the results in [28, 32, 33, 34] distinct (see [31] for the details).

**Definition 5.3.** Consider a point  $\tilde{y} \in \mathfrak{T}$  and let  $C \subset T_{\tilde{y}}\mathcal{M}$  be a QDQ approximating cone of  $\mathfrak{T}$  at  $\tilde{y}$ . If S > 0, we say that the point  $\tilde{y}$  is C-needle-controllable (with respect to (5.31)-(5.32)) at S, if, for every  $\boldsymbol{\xi} \in C^{\perp} \setminus \{0\}$ , there exist  $\delta_1 > 0$  and  $\delta_2 \in (0, S]$  such that

$$\inf_{\check{\mathfrak{w}}\in\mathfrak{W}}\sup_{\mathfrak{w}\in\mathfrak{W}}\boldsymbol{\xi}\cdot(f(s,\tilde{y},\mathfrak{w})-f(s,\tilde{y},\check{\mathfrak{w}}))\geq\delta_1\qquad a.e.\ s\in[S-\delta_2,S].$$
(5.33)

**Theorem 5.2.** Consider a feasible process  $(\hat{y}, \hat{w}) : [0, S] \to \mathcal{M} \times \mathfrak{W}$  of (5.31)-(5.32). Let C be a QDQ approximating cone to  $\mathfrak{T}$  at  $\hat{y}(S)$ , and let  $\hat{y}(S)$  be C-needle-controllable at S. Then the process  $(\hat{y}, \hat{w})$  is not an abnormal extremal, so, in particular, it does not satisfy the infimum-gap condition.

*Proof.* Assume by contradiction that the extremal  $(\hat{y}, \hat{w})$  is abnormal, namely that there exists an absolutely continuous lift  $(\hat{y}, \lambda) : [0, S] \to T^*\mathcal{M}$  of  $\hat{y}$  such that  $\lambda \neq 0, \lambda(S) \in -C^{\perp}$ , and the inequality

$$\lambda(s) \cdot \left( f(s, \hat{y}(s), \mathfrak{w}) - f(s, \hat{y}(s), \hat{w}(s)) \right) \le 0 \tag{5.34}$$

holds true for almost every  $s \in [0, S] \setminus I_0$  and every  $\mathfrak{w} \in \mathfrak{W}$ ,  $I_0$  having Lebesgue measure equal to zero. Taking  $\boldsymbol{\xi} := \lambda(S)$  in (5.33), we deduce that there exist  $\delta_1, \delta_2 > 0$  and a neighbourhood  $U \subset T^* \mathcal{M}$  of  $(\hat{y}(S), \boldsymbol{\xi})$  such that, for all  $(y, p) \in U$ ,

$$\sup_{\mathfrak{w}\in\mathfrak{W}} p \cdot (f(s, y, \mathfrak{w}) - f(s, y, \check{\mathfrak{w}})) > \frac{\delta_1}{2} \quad \forall \check{\mathfrak{w}}\in\mathfrak{W}, \quad a.e. \ s \in [S - \delta_2, S].$$

Now, by choosing  $\varepsilon \in [0, \delta_2]$  sufficiently small, for every  $s \in [S - \varepsilon, S]$  one has  $(\hat{y}(s), \lambda(s)) \in U$ , so that

$$\sup_{\mathfrak{w}\in\mathfrak{W}}\lambda(s)\cdot(f(s,\hat{y}(s),\mathfrak{w})-f(s,\hat{y}(s),\check{\mathfrak{w}}))>\frac{\delta_1}{2}\quad\forall\check{\mathfrak{w}}\in\mathfrak{W},\quad\text{a.e.}\ s\in[S-\varepsilon,S].$$

In particular, for all  $s \in [S - \varepsilon, S] \setminus I_0$ ,

$$\sup_{\mathfrak{w}\in\mathfrak{W}}\lambda(s)\cdot(f(s,\hat{y}(s),\mathfrak{w})-f(s,\hat{y}(s),\hat{w}(s)))>0,$$

which contradicts the maximization relation (5.34).

### 6 Proofs of the main results

#### 6.1 Proof of the Geometric Principle (Theorem 5.1)

By a basic result on control system (see e.g. [35], [9]),  $\mathbf{C}_{\mathfrak{w}_1,\ldots,\mathfrak{w}_N}^{s_1,\ldots,\mathfrak{w}_N}$  turns out to be a QDQ approximating cone to the (local) extended reachable set  $\mathcal{R}_{\mathcal{W}}^{\hat{w},r}$  at  $\hat{y}(S)$ .<sup>13</sup> More importantly, Theorem 4.1 states that  $\mathbf{C}_{\mathfrak{w}_1,\ldots,\mathfrak{w}_N}^{s_1,\ldots,s_N}$  is also a QDQ approximating cone to the (local) original reachable set  $\mathcal{R}_{\mathcal{V}}^{\hat{w},r} \cup \{\hat{y}(S)\}$  at  $\hat{y}(S)$ . Therefore, by Lemma 3.1, the sets  $\left(\mathcal{R}_{\mathcal{V}}^{\hat{w},r} \cup \{\hat{y}(S)\}\right)$  and  $\mathfrak{T}$  are locally separated at  $\hat{y}(S)$ , which by Theorem 2.3, i), implies that the cones  $\mathbf{C}_{\mathfrak{w}_1,\ldots,\mathfrak{w}_N}^{s_1,\ldots,s_N}$  and C are not strongly transverse. Since linear separability is equivalent to non-transversality (Proposition 1.2) we have to prove that *these cones are not transverse as well*. Indeed, in view of Proposition 1.1 the only case in which the cones are complementary subspaces of  $T_{\hat{y}(S)}\mathcal{M}$ . However, such an instance is excluded by Theorem 4.2 and the occurrence of an infimum gap. In fact, if  $\mathbf{C}_{\mathfrak{w}_1,\ldots,\mathfrak{w}_N}^{s_1,\ldots,s_N}$  and C satisfy  $\mathbf{C}_{\mathfrak{w}_1,\ldots,\mathfrak{w}_N}^{s_1,\ldots,s_N} \oplus C = T_{\hat{y}(S)}\mathcal{M}$ , then Theorem 4.2 assures the existence of a sequence  $(y_k)_{k\in\mathbb{N}} \subset \mathcal{R}_{\mathcal{V}} \cap \mathfrak{T}$  such that  $y_k \to \hat{y}(S)$ , which contradicts the fact that  $(\hat{y}, \hat{w})$  verifies the infimum gap condition. This concludes the proof.

<sup>&</sup>lt;sup>13</sup>For instance: it is well-known that  $\mathbf{C}_{\mathfrak{w}_1,\ldots,\mathfrak{w}_N}^{s_1,\ldots,s_N}$  is a Boltyanski approximating cone to  $\mathcal{R}_{\mathcal{W}}^{\hat{w},r}$  at  $\hat{y}(S)$  (see e.g. [39]). Furthermore, a Boltyanski approximating cone is clearly a QDQ approximating cone.

### 6.2 Proof of the Normality No-Gap Criterion (Corollary 5.1)

By Theorem 5.1, the cones  $\mathbf{C}_{\mathfrak{w}_1,\ldots,\mathfrak{w}_N}^{s_1,\ldots,s_N}$  and C are linearly separable. This means that there exists  $\xi \in (T_{\hat{y}(S)}\mathcal{M})^* \setminus \{0\}$  such that  $\xi \in -C^{\perp} \cap (\mathbf{C}_{\mathfrak{w}_1,\ldots,\mathfrak{w}_N}^{s_1,\ldots,s_N})^{\perp}$ . Now let us set  $\lambda(s) := \xi \cdot M(S,s)$ , where M(S,s) is the fundamental matrix defined in (4.18), so that

$$\lambda \neq 0, \qquad \lambda(S) \in -C^{\perp}, \quad \frac{d\lambda}{ds}(s) = -\lambda(s) \cdot \frac{\partial f}{\partial y}(s, \hat{y}(s), \hat{w}(s)), \quad \text{for a.e. } s \in [0, S].$$

By  $\xi \in \left(\mathbf{C}_{\mathfrak{w}_1,\ldots,\mathfrak{w}_N}^{s_1,\ldots,s_N}\right)^{\perp}$ , it follows that, for every  $i = 1,\ldots,N$ ,

$$0 \geq \xi \cdot \left( M(S, s_i) \cdot \left( f(s_i, \hat{y}(s_i), \mathfrak{w}_i) - f(s_i, \hat{y}(s_i), \hat{w}(s_i)) \right) \right) \\ = \left( \xi \cdot M(S, s_i) \right) \cdot \left( f(s_i, \hat{y}(s_i), \mathfrak{w}_i) - f(s_i, \hat{y}(s_i), \hat{w}(s_i)) \right) \\ = \lambda(s_i) \cdot \left( f(s_i, \hat{y}(s_i), \mathfrak{w}_i) - f(s_i, \hat{y}(s_i), \hat{w}(s_i)) \right).$$

$$(6.35)$$

Therefore the lift  $(\hat{y}, \lambda)$  verifies (i)-(iv) of Definition 5.1, except that (iii) is verified only for every finite set of pairs  $(s_i, \mathfrak{w}_i) \in [0, S] \times \mathfrak{W}, i = 1, \ldots, N$ , such that  $s_1, \ldots, s_N \in$  $SD\{f(\cdot, \cdot, \hat{w}(\cdot))\} \cap SD\{f(\cdot, \cdot, \mathfrak{w}_1)\} \cap \ldots \cap SD\{f(\cdot, \cdot, \mathfrak{w}_N)\}, 0 < s_1 < \ldots, < s_N \leq S$ . To conclude the proof we have to show the validity of (iii) in the whole control value set  $\mathfrak{W}$ and almost all times. This is achieved through non-empty intersection arguments borrowed from those utilized in [39] to prove the Maximum Principle.

#### 6.2.1 The case of a finite subset of controls

Let us consider a *finite subset* of control values  $\bar{\mathfrak{W}} \subseteq \mathfrak{W}$  and let us set

$$E(\bar{\mathfrak{W}}) := \bigcap_{w \in \bar{\mathfrak{W}}} \operatorname{SD}\{f(\cdot, \hat{y}(\cdot), \mathfrak{w})\} \bigcap \operatorname{SD}\{f(\cdot, \hat{y}(\cdot), \hat{w}(\cdot))\} \quad \Big( \subset [0, S] \Big).$$

Since  $\overline{\mathfrak{W}}$  is finite,  $E(\overline{\mathfrak{W}})$  has measure equal to S. Therefore, by Lusin's theorem we can write

$$E(\bar{\mathfrak{W}}) = \bigcup_{j=0}^{\infty} E_j, \tag{6.36}$$

where  $E_0$  has zero measure and, for every j, the set  $E_j$  is compact, and, for every  $w \in \mathfrak{W}$ , the restrictions to  $E_j$  of the map

$$s \mapsto r^{\mathfrak{w}}(s) := f(s, \hat{y}(s), \mathfrak{w}) - f(s, \hat{y}(s), \hat{w}(s)),$$

is continuous.

For every j let  $D_j$  be the set of *density points* <sup>14</sup> of  $E_j$ . Since, for every natural number j,  $E_j$  and  $D_j$  have the same measure, one obtains that <sup>15</sup>

$$meas(E(\bar{\mathfrak{W}})) = meas(D)$$

where we have set  $D := \bigcup_{j=0}^{\infty} D_j$ .

Now let F be an arbitrary, non-empty, subset of  $D \times \overline{\mathfrak{W}}$ , and let us define the subset  $\Lambda(F, \overline{\mathfrak{W}}) \subseteq (T_{\hat{\mathfrak{Y}}(S)}\mathcal{M})^*$  by setting

$$\Lambda(F,\bar{\mathfrak{W}}) := \left\{ \bar{\lambda} \in (T_{\hat{y}(S)}\mathcal{M})^*, \quad |\bar{\lambda}| = 1, \quad \bar{\lambda} \text{ verifies } (\mathbf{P})_F \right\},\$$

where property  $(\mathbf{P})_F$  is as follows:

 $^{14}\text{We}$  recall that an element  $t\in B\subset \mathbb{R}$  is a density point for B if

$$\lim_{\delta \to 0+} \frac{meas([t-\delta, t+\delta])}{2\delta} = .1$$

<sup>15</sup>For every measurable subset  $A \subseteq [0, S]$ , meas(A) denotes the Lebesge measurable of A.

**Property**  $(\mathbf{P})_F$ . The pair  $(\hat{y}, \lambda) \in W^{1,1}([0, S]; T^*\mathcal{M})$  is a lift of  $\hat{y}(\cdot)$  such that:

- (1)  $\lambda(S) = \bar{\lambda} \in -C^{\perp}$ (2)  $d\lambda$   $\partial f_{(\alpha, \hat{\alpha}(\alpha), \hat{\alpha}(\alpha))}$
- (2)  $\begin{aligned} \frac{d\lambda}{ds} &= -\lambda \cdot \frac{\partial f}{\partial y}(s, \hat{y}(s), \hat{w}(s)), \qquad \text{a.e } s \in [0, S]; \\ (3) \quad \lambda(s) \cdot f(s, \hat{y}(s), \hat{w}(s)) \geq \lambda(s) \cdot f(s, \hat{y}(s), \mathfrak{w}) \\ \text{for every } (s, \mathfrak{w}) \in F. \end{aligned}$

Notice that, for every subset  $F \in D \times \overline{\mathfrak{W}}$ ,  $\Lambda(F, \overline{\mathfrak{W}})$  is compact and, moreover,

$$\Lambda(F_1 \cup F_2, \bar{\mathfrak{W}}) = \Lambda(F_1, \bar{\mathfrak{W}}) \cap \Lambda(F_2, \bar{\mathfrak{W}})$$

for all  $F_1, F_2 \in D \times \overline{\mathfrak{V}}$ .

and

By Theorem 5.1,  $\Lambda(F, \overline{\mathfrak{W}}) \neq \emptyset$  as soon as F is *finite* and can be written as

$$F = \{(s_1, \mathfrak{w}_1), ..., (s_m, \mathfrak{w}_m)\} \qquad 0 \le s_1 < ... < s_i < ... < s_m < S, \ \mathfrak{w}_i \in \mathfrak{W}.$$

**Claim:** One has  $\Lambda(F, \overline{\mathfrak{W}}) \neq \emptyset$  even when F is an arbitrary finite subset of  $D \times \overline{\mathfrak{W}}$ , namely F can be written as

$$F = \{(s_1, \mathfrak{w}_1), ..., (s_m, \mathfrak{w}_m)\} \qquad 0 \le s_1 \le ... \le s_i \le ... \le s_m, \ \mathfrak{w}_i \in \mathfrak{W}.$$

Indeed, every  $s_i$  belongs to a suitable  $D_h$ , which can be labelled as  $D_{h(i)}$ . Since  $D_{h(i)}$  is made of density points, there exist sequences  $(s_{i,j})$  such that

$$s_{i,j} \in D_{h(i)} \quad \forall j, \qquad s_i = \lim_{j \to \infty} s_{i,j},$$
  
 $s_{1,j} < \dots < s_{m,j} \quad \forall j \in \mathbb{N}.$ 

Set  $F_j = \{(s_{i,j}, \mathfrak{w}_1), ..., (s_{m,j}, \mathfrak{w}_m)\}$  -so that  $\Lambda(F_j, \mathfrak{W}) \neq \emptyset$ - and choose  $\bar{\lambda}_j \in \Lambda(F_j, \mathfrak{W})$ . Since  $|\bar{\lambda}_j| = 1$  for all j, by possibly taking a subsequence we can assume that  $(\bar{\lambda}_j)_{j \in \mathbb{N}}$  converges to some  $\bar{\lambda}$ . For every  $s \in [s_1, S]$ , define the lifts  $(\hat{y}, \bar{\lambda}), (\hat{y}, \bar{\lambda}_j) \in W^{1,1}([s_1, S]; T\mathcal{M})$  of  $\hat{y}$  such that  $\bar{\lambda}(S) = \bar{\lambda}, \bar{\lambda}_j(S) = \bar{\lambda}_j$  and both satisfying the equation

$$\frac{d\lambda}{ds} = -\lambda \cdot \frac{\partial f}{\partial y}(s, \hat{y}(s), \hat{w}(s)), \quad \text{a.e } s \in [s_1, S].$$

The mapping  $s \mapsto \overline{\lambda}_j(s)$  satisfies the inequality

$$\lambda_j(s_{i,j}) \cdot f(s_{i,j}, \hat{y}(s_{i,j}), \hat{w}(s_{i,j})) \ge \lambda_j(s_{i,j}) \cdot f(s_{i,j}, \hat{y}(s_{i,j}), \mathfrak{w})$$

for all  $j \in \mathbb{N}$ , every  $i = 1, \ldots, m$  and  $\mathfrak{w} \in \mathfrak{W}$ . Since, for every  $i = 1, \ldots, m$ , the map  $s \mapsto r^{\mathfrak{w}_i}(s) := f(s, \hat{y}(s), \mathfrak{w}_i) - f(s, \hat{y}(s), \hat{w}(s))$  is continuous on  $D_{h(i)}$ , the function  $s \mapsto \overline{\lambda}_j(s) \cdot r^{\mathfrak{w}_i}(s)$  is also continuous on  $D_{h(i)}$ , so passing to the limit we can conclude that

$$\overline{\lambda}(s_i) \cdot f(s_i, \hat{y}(s_i), \hat{w}(s_i)) \ge \overline{\lambda}(s_i) \cdot f(s_i, \hat{y}(s_i), \mathfrak{w})$$

for every i = 1, ..., m and  $\mathfrak{w} \in \overline{\mathfrak{W}}$ . Since one also has  $0 \neq \overline{\lambda} = \overline{\lambda}(S) \in -C^{\perp}$ , the claim is proved.

#### 6.2.2 The general case of an infinite control set

Up to now we have shown that, if  $\overline{\mathfrak{W}}$  is finite, and  $F \subset D \times \overline{\mathfrak{W}}$  is finite —and we write  $card(F) < \infty$ —, then  $\Lambda(F, \overline{\mathfrak{W}})$  is a nonempty compact set. We now conclude the proof

through a standard non-empty intersection argument (see e.g. [39]). If we take a finite family  $F^1, \ldots, F^r \subset D \times \overline{\mathfrak{W}}$  such that  $card(F_i) < \infty$  for every  $i = 1, \ldots, r$ , one has

$$\Lambda(F^1,\bar{\mathfrak{W}})\cap\cdots\cap\Lambda(F^r,\bar{\mathfrak{W}})=\Lambda(F^1\cup\cdots\cup F^r,\bar{\mathfrak{W}})\neq\emptyset,$$

(for  $card(F^1 \cup \cdots \cup F^r) < \infty$ ). Hence,

$$\{\Lambda(F,\bar{\mathfrak{W}}) \mid F \subset D \times \bar{\mathfrak{W}}, \quad \text{card} F < \infty\}$$

is a family of compact subsets such that each finite intersection is nonempty. This implies that the (infinite) intersection of all  $\Lambda(F, \overline{\mathfrak{V}})$  such that  $cardF < \infty$  is nonempty. Therefore

$$\Lambda(D\times\bar{\mathfrak{W}},\mathfrak{W})=\Lambda\left(\bigcup_{card(F)<\infty}F,\mathfrak{W}\right)=\bigcap_{card(F)<\infty}\Lambda(F,\bar{\mathfrak{W}})\neq\emptyset$$

To end the proof in the general case when  $card(\mathfrak{W})$  is infinite, for any arbitrary subset  $\hat{\mathfrak{W}} \subseteq \mathfrak{W}$  define

$$\Lambda(\hat{\mathfrak{W}}) := \left\{ \bar{\lambda} \in (T_{\hat{y}(S)}\mathcal{M})^*, \quad |\bar{\lambda}| = 1, \quad \bar{\lambda} \text{ verifies } (\mathbf{PP})_F \right\},\$$

where property  $(\mathbf{PP})_{\hat{F}}$  is as follows:

- $(\mathbf{PP})_{\hat{F}}$ : The pair  $(\hat{y}, \lambda) \in W^{1,1}([0, S]; T^*\mathcal{M})$  is a lift of  $\hat{y}(\cdot)$  such that:
- (1)  $\lambda(S) = \bar{\lambda} \in -C^{\perp}$
- (2)  $\frac{d\lambda}{ds} = -\lambda \cdot \frac{\partial f}{\partial y}(s, \hat{y}(s), \hat{w}(s)),$  a.e  $s \in [0, S];$

(3) For each  $\mathfrak{w} \in \hat{\mathfrak{W}}$ , there exists a subset of full measure  $I_{\mathfrak{w}} \subseteq [0, S]$  such that

$$\lambda(s) \cdot f(s, \hat{y}(s), \hat{w}(s)) \ge \lambda(s) \cdot f(s, \hat{y}(s), \mathfrak{w})$$

for every  $s \in I_{\mathfrak{w}}, \mathfrak{w} \in \hat{\mathfrak{W}}$ .

So, proving Theorem 5.1 is equivalent to showing that

$$\Lambda(\mathfrak{W}) \neq \emptyset. \tag{6.37}$$

Since

$$\Lambda(\mathfrak{W}) = \bigcap_{\operatorname{card}(\hat{\mathfrak{W}}) < \infty} \Lambda(\hat{\mathfrak{W}}), \tag{6.38}$$

once again we have to show that the (possibly infinite) family

$$\Big\{\Lambda(\hat{\mathfrak{W}}), \quad card(\hat{\mathfrak{W}}) < \infty \Big\}$$

has non-empty intersection. This can easily achieved by the same arguments as above. Indeed,  $\Lambda(\hat{\mathfrak{W}})$  is not empty and compact as soon as  $\hat{\mathfrak{W}}$  is finite. Furthermore, for every  $\mathfrak{W}_1, \mathfrak{W}_2 \subseteq \mathfrak{W}$  one has

$$\Lambda(\mathfrak{W}_1\cup\mathfrak{W}_2)=\Lambda(\mathfrak{W}_1)\cap\Lambda(\mathfrak{W}_2).$$

In particular, the family  $\left\{\Lambda(\hat{\mathfrak{W}}): card(\hat{\mathfrak{W}}) < \infty\right\}$  is made of compact subsets and satisfies the finite intersection property, that is, the intersection of any finite finite subfamily  $\left\{\Lambda(\hat{\mathfrak{W}}): card(\hat{\mathfrak{W}}) < \infty\right\}$  is not empty. Therefore, it has non-empty intersection, namely

$$\Lambda(\mathfrak{W}) = \bigcap_{\operatorname{card}(\hat{\mathfrak{W}}) < \infty} \Lambda(\hat{\mathfrak{W}}) \neq \emptyset \; .$$

This concludes the proof of Theorem 5.1.

### 7 An application to non-convex, unbounded, problems

Impulsive optimal control problems –where the dynamics is *unbounded*– have been extensively studied together with their applications [3, 4, 5, 8, 10, 11, 13, 14, 19, 20, 22, 24, 25, 37, 46]. The space-time representation (see (7.40) below) can be regarded as an *extension* of unbounded control systems. An important case is the one of a minimum problem with of control-affine dynamics:

$$\begin{cases} \text{Minimize } h(t_2, x(t_2), \eta(t_2)) \\ \text{over } t_2 \in \mathbb{R}, \ t_2 > t_1, \ (x, \eta, u) \in AC([t_1, t_2], \mathcal{M} \times \mathbb{R}) \times L^1([t_1, t_2], U) \\ \text{such that} \end{cases}$$

$$(P) \begin{cases} \frac{dx}{dt}(t) = f(t, x(t)) + \sum_{j=1}^{m} g_j(t, x(t)) u^j(t) \quad \text{a.e. } t \in [t_1, t_2] \\ \frac{d\eta}{dt}(t) = |u(t)| \quad \text{a.e. } t \in [t_1, t_2] \\ (x(t_1), \eta(t_1)) = (\bar{x}, 0), \quad (t_2, x(t_2), \eta(t_2)) \in \bar{\mathfrak{T}} \times [0, K] \end{cases}$$

Here the set U where the controls u take values is *unbounded*. Furthermore, the state x range over a n-dimensional Riemannian manifold  $\mathcal{M}$  of class  $C^2$ , and the time-dependent vector fields  $f, g_1, \ldots, g_m$  are of class  $C^1$  in x, measurable in t, and uniformly bounded by a  $L^1$  map. Moreover, the cost  $h: \mathbb{R} \times \mathcal{M} \times \mathbb{R} \to \mathbb{R}$  is a continuous function,  $(t_1, \bar{x}) \in \mathbb{R} \times \mathcal{M}$  is a fixed initial condition, K is a non negative fixed constant, possibly equal to  $+\infty$ , and the end-point constraint  $\overline{\mathfrak{T}} \subseteq \mathbb{R} \times \mathbb{R}^n$  is a closed subset. Notice incidentally that the function  $\eta(t)$  coincides with the  $L^1$ -norm of the control function  $u := (u^1, u^2, ..., u^m)$  on the interval  $[t_1, t]$ .

The gap-abnormality criterion for this kind of systems (where one considers the spacetime extension (7.40) below) has been already investigated in the case when the set of controls U is a convex cone [28]. Actually, thanks to Corollary 5.1 (see also [31]), the main result in [28] can be extended to the case in which the state ranges on a Riemannian manifold. However the generalization made possible by Corollary 5.1 allows one to go much further. Indeed, in what follows we are able to deduce from Corollary 5.1 that the gap-abnormality criterium holds true also in the situation when the control set U is unbounded but is *neither convex nor a cone*.

More precisely, we will consider the following two cases:

- CASE (I) (Space-time convex extension) The controls take values on a (necessarily unbounded) subset  $U \subseteq \mathbb{R}^m$  such that  $\mathbf{co} U$  is a (convex) cone of  $\mathbb{R}^m$ , where we have used  $\mathbf{co} E$  to denote the convex hull of a subset  $E \subseteq \mathbb{R}^m$ ; For instance, one could consider the set  $U = \mathbb{N}^m$ , so that  $\mathbf{co}(U) = [0, +\infty]^m$ .
- CASE (II) (Space-time non-convex extension) The controls take values on a (necessarily unbounded) subset  $U \subseteq \mathbb{R}^m$  such that

$$(r, u) \in [0, +\infty[\times U \Longrightarrow \exists \rho > r \ s.t. \ \rho u \in U \tag{7.39}$$

Notice that, if for a given set E we consider the  $\operatorname{conic}(E) := \{re \mid (r, e) \in [0, +\infty[\times E] \\ -a \text{ cone which we call the conic envelope of } E-, hypothesis (7.39) implies that} \}$ 

$$\inf_{u \in U} d(u, \operatorname{conic}(U)) = 0$$

For instance, one could consider the set  $U = \{(n^2, 0), (0, -m^3) \mid m, n \in \mathbb{N}\}$ , so that **conic**  $(U) = [0, +\infty[\times\{0\} \cup \{0\} \times ] - \infty, 0].$ 

**Remark 7.1.** We will treat CASE (I) in detail, describing the extension to the convex space-time system obtained by both convexification of the dynamics and the closure of

suitably reparameterized processes. Instead, we will only suggest the needed changes to deal with CASE (II), where the only extension comes from reparameterization. However, CASE (II) is somehow more significative, in that it marks the most important improvement with respect to the former literature initiated by Warga's work. Indeed, in this case not only the original dynamics but *also the extended dynamics is non-convex*. This can be of interest in those application where the convexification of the dynamics is not needed (for instance because one gets existence of minima without invoking convexification).

### 7.1 CASE (I) (Space-time convex extension)

In order to formulate this problem by means of the terminology adopted in Theorem 5.1, we need to embed our system into a suitably extended one. To this aim we need to perform both a 'compactification' (to manage unboundedness) and a 'convexification'. Let us begin by setting

$$\begin{aligned} A &:= \left\{ a = (a^1, \dots a^{n+3}) \in [0,1]^{n+3}, \ \sum_{i=1}^{n+3} a^i = 1 \right\} & \mathcal{A} &:= L^1 \left( [0,S], A \right) \\ \mathbf{W} &:= \left\{ (w^0, w) \in [0, \infty) \times \mathbf{co} \ U : \ w^0 + |w| = 1 \right\} & \hat{\mathcal{W}} &:= L^1 \left( [0,S], \mathbf{W} \right) \\ \mathbf{V} &:= \left\{ (v^0, v) \in (0, \infty) \times U : \ v^0 + |v| = 1 \right\} & \hat{\mathcal{V}} &:= L^1 \left( [0,S], \mathbf{V} \right) \\ D &:= [-0.5, 0.5] & \mathcal{D} &:= L^1 \left( [0,S], \mathbf{V} \right) \end{aligned}$$

$$\mathcal{W} := \mathcal{A} \times (\hat{\mathcal{W}})^{n+3} \times \mathcal{D}, \qquad \mathcal{V} := \{(1, 0, \dots, 0)\} \times (\hat{\mathcal{V}})^{n+3} \times \mathcal{D},$$

and let us consider the optimal control problem

$$(P)_{\mathcal{W}}^{h} \begin{cases} \text{Minimize } h\left(z^{0}(\hat{S}), z(\hat{S}), \nu(\hat{S})\right) \\ \text{over } (z^{0}, z, \nu, a, (w_{1}^{0}, w_{1}), \dots, (w_{n+3}^{0}, w_{n+3}), d)(\cdot) \in AC([0, \hat{S}], \mathbb{R} \times \mathcal{M} \times \mathbb{R}) \times \mathcal{W} \times \mathcal{D} \\ \text{s. t., for a.e. } s \in [0, \hat{S}], \\ \begin{cases} \frac{dz^{0}}{ds}(s) = (1 + d(s)) \sum_{i=1}^{n+3} a^{i}(s) w_{i}^{0}(s) \\ \frac{dz}{ds}(s) = (1 + d(s)) \sum_{i=1}^{n+3} a^{i}(s) \left(f(z^{0}(s), z(s)) w_{i}^{0}(s) + \sum_{j=1}^{m} g_{j}(z^{0}(s), z(s)) w_{i}^{j}(s)\right) \\ \frac{d\nu}{ds}(s) = (1 + d(s)) \sum_{i=1}^{n+3} a^{i}(s) |w_{i}(s)| \\ (z^{0}(0), z(0), \nu(0)) = (0, \bar{x}, 0), \qquad (z^{0}(\hat{S}), z(\hat{S}), \nu(\hat{S})) \in \bar{\mathfrak{T}} \times [0, K] \end{cases}$$

$$(7.40)$$

Accordingly, a pair

$$\left(\left((z^0, z, \nu)\right), (a, (w_1^0, w_1), \dots, (w_{n+3}^0, w_{n+3}), d\right)\right)$$

such that  $(z^0, z, \nu)$  is the solution of the above control system corresponding to the *control*  $(a, (w_1^0, w_1), \ldots, (w_{n+3}^0, w_{n+3}), d)$  is called *a process of*  $(P)_{\mathcal{W}}^h$ . The embedding of the problem (P) into  $(P)_{\mathcal{W}}^h$  is as follows: fix  $\hat{S} > 0$ , and, for every control  $u : [t_1, t_2^u] \to U$ , consider the function  $\sigma_u : [t_1, t_2^u] \to [0, \hat{S}]$  defined by

$$\sigma_u(t) := \frac{\hat{S}}{t_2^u + \|u\|_1} \int_{t_1}^t \left(1 + |u(\tau)|\right) d\tau = \frac{\hat{S}}{t_2^u + \|u\|_1} (t + \eta(t)).$$
(7.41)

Then define  $\mathcal{I}: \mathbb{R} \times AC([t_1, t_2], \mathcal{M} \times \mathbb{R} \times \mathbb{R}^m) \to \mathbb{R} \times AC([0, S], \mathbb{R} \times \mathcal{M} \times \mathbb{R} \times \mathbb{R} \times \mathbb{R}^m)$  by setting

$$\mathcal{I}(x,\eta,u) := \left( \left( (z^0, z, \nu) , (a, (w_1^0, w_1), \dots, (w_{n+3}^0, w_{n+3}), d) \right) \right)$$

where, for all  $s \in [0, \hat{S}]$  and all  $i = 1, \ldots, m$ ,

$$(z^{0}, z, \nu)(s) := (id, x, \eta) \circ \sigma_{u}^{-1}(s), \qquad \forall s \in [0, \hat{S}].$$
$$a := (1, 0, \dots, 0), \quad d := \frac{t_{2}^{u} + ||u||_{1}}{\hat{S}} - 1, \quad (w_{i}^{0}, w_{i}) := \left(\frac{1}{1 + |u|}(1, u)\right) \circ \sigma_{u}^{-1}(s).$$

By a trivial use of the chain rule one gets the following result (see e.g. [3] for a similar embedding):

**Lemma 7.1.** The embedding  $\mathcal{I}$  is injective <sup>16</sup>. Moreover, the image space of the embedding  $\mathcal{I}$  coincides with the set of all processes  $\left(\left((z^0, z, \nu), (a, (w_1^0, w_1), \ldots, (w_{n+3}^0, w_{n+3}), d)\right)$  such that

$$(a, (w_1^0, w_1), \dots, (w_{n+3}^0, w_{n+3}), d) \in \mathcal{V}$$

Thanks to Lemma 7.1 we can identify the original problem (P) with the problem

$$(P)_{\mathcal{V}}^{h} \begin{cases} \text{Minimize } h\left(z^{0}(\hat{S}), z(\hat{S}), \nu(\hat{S})\right) \\ \text{over } (z^{0}, z, \nu, a, (w_{1}^{0}, w_{1}), \dots, (w_{n+3}^{0}, w_{n+3}), d)(\cdot) \in AC([0, \hat{S}], \mathbb{R} \times \mathcal{M} \times \mathbb{R}) \times \mathcal{V} \times \mathcal{D} \\ \text{s. t., for a.e. } s \in [0, \hat{S}], \\ \begin{cases} \frac{dz^{0}}{ds}(s) = (1 + d(s)) \sum_{i=1}^{n+3} a^{i}(s) w_{i}^{0}(s) \\ \frac{dz}{ds}(s) = (1 + d(s)) \sum_{i=1}^{n+3} a^{i}(s) \left(f(z^{0}(s), z(s)) w_{i}^{0}(s) + \sum_{j=1}^{m} g_{j}(z^{0}(s), z(s)) w_{i}^{j}(s)\right) \\ \frac{d\nu}{ds}(s) = (1 + d(s)) \sum_{i=1}^{n+3} a^{i}(s) |w_{i}(s)| \\ (z^{0}(0), z(0), \nu(0)) = (0, \bar{x}, 0), \qquad (z^{0}(\hat{S}), z(\hat{S}), \nu(\hat{S})) \in \bar{\mathfrak{T}} \times [0, K] \end{cases}$$

We can now apply the theory developed in the previous sections. In view of the sufficient condition provided by Theorem 4.1, it is trivial to verify that the family of controls  $\mathcal{V}$  is abundant in  $\mathcal{W}$  w.r.t. the dynamics of problem  $(P)^h_{\mathcal{W}}$ . Therefore, by Corollary 5.1, one obtains the following infimum-gap result:

**Theorem 7.1.** Consider a feasible extended process

$$((y^0, y, \nu), (\hat{a}, (\hat{w}^0_1, \hat{w}_1), \dots, (\hat{w}^0_{n+3}, \hat{w}_{n+3}), \hat{d})), \quad \hat{d} \equiv 0,$$

and assume that it satisfies the infimum gap condition. Then, for all approximating cones C to  $\mathfrak{T} := \overline{\mathfrak{T}} \times [0, K]$  at  $(\hat{y}^0, \hat{y}, \hat{\nu})(\hat{S})$ , there exist a number  $\beta \leq 0$ , an absolutely continuous path  $(\lambda^0, \lambda, \lambda^{\nu}) \in W^{1,1}([0, \hat{S}]; \mathbb{R}^{(1+n+1)})$  and a zero-measure subset  $I_0$  such that the following conditions hold true:

(i) 
$$(\lambda^0, \lambda, \lambda^\nu) \neq 0$$
  
(ii.1)  $\frac{d\lambda^0}{ds}(s) = -\lambda(s) \cdot \left[\sum_{i=1}^{n+3} \hat{a}^i(s) \left(\frac{\partial f}{\partial y^0}(\hat{y}^0(s), \hat{y}(s))\hat{w}_i^0(s) + \sum_{j=1}^m \frac{\partial g_j}{\partial y^0}(\hat{y}^0(s), \hat{y}(s))\hat{w}_i^j(s)\right)\right]$ 

<sup>&</sup>lt;sup>16</sup>Notice that the injectivity is a consequence of the fact that  $w_i^0(s) + |w_i(s)| = 1$  for a.e.  $s \in [0, S]$  and for  $i = 1, \ldots, m$ .

$$\begin{aligned} (ii.2) \quad & \frac{d\lambda}{ds}(s) = -\lambda(s) \cdot \left[ \sum_{i=1}^{n+3} \hat{a}^{i}(s) \left( \frac{\partial f}{\partial y} (\hat{y}^{0}(s), \hat{y}(s)) \hat{w}_{i}^{0}(s) + \sum_{j=1}^{m} \frac{\partial g_{j}}{\partial y} (\hat{y}^{0}(s), \hat{y}(s)) \hat{w}_{i}^{j}(s) \right) \right] \\ & a.e. \ s \in [0, \hat{S}] \\ (iii) \quad & (1+d) \sum_{i=1}^{n+3} a^{i} \Big[ \lambda^{0}(s) w_{i}^{0} + \lambda(s) \cdot \Big( \hat{f}(s) w_{i}^{0} + \sum_{j=1}^{m} \hat{g}_{j}(s) w_{i}^{j} \Big) + \beta |w_{i}| \Big] \\ & \leq \sum_{i=1}^{n+3} \hat{a}^{i}(s) \Big[ \lambda^{0}(s) \hat{w}_{i}^{0}(s) + \lambda(s) \cdot \Big( \hat{f}(s) \hat{w}_{i}^{0}(s) + \sum_{j=1}^{m} \hat{g}_{j}(s) \hat{w}_{i}^{j}(s) \Big) + \beta |\hat{w}_{i}(s)| \Big] \\ & for \ every \ (w^{0}, w, d) \in \mathbf{W} \times [-0.5, 0.5] \ and \ s \in [0, \hat{S}] \backslash I_{0}^{17} \\ (iv) \quad & (\lambda^{0}(\hat{S}), \lambda(\hat{S}), \beta) \in -C^{\perp}. \end{aligned}$$

### 7.2 CASE (II) (Space-time non-convex extension)

Let us recall that we are assuming that (7.39), namely

$$(r, u) \in [0, +\infty] \times U \Longrightarrow \exists \rho > r \text{ s.t. } \rho u \in U.$$

Unlike the previous case, we are not going to convexify the dynamics, while we will consider just the impulsive extension. Without repeating all steps, we just observe that the sought extension is obtained by neglecting the sets A and A, and by replacing  $\mathbf{W}$  with the (generally non-convex) set  $\mathbf{W}^{nc} := \{(w^0, w) \in [0, \infty) \times U : w^0 + |w| = 1\}$ , respectively. In turn, problem  $(P)^h_{\mathcal{W}}$  simplifies into the following non-convex problem  $(P^{nc})^h_{\mathcal{W}}$ :

$$(P^{nc})_{\mathcal{W}}^{h} \begin{cases} \text{Minimize } h\left(z^{0}(\hat{S}), z(\hat{S}), \nu(\hat{S})\right) \\ \text{over } (z^{0}, z, \nu, (w^{0}, w), d) \in AC([0, \hat{S}], \mathbb{R} \times \mathcal{M} \times \mathbb{R}) \times \mathcal{W} \times \mathcal{D} \\ \text{s. t., for a.e. } s \in [0, \hat{S}], \\ \begin{cases} \frac{dz^{0}}{ds}(s) = (1 + d(s))w^{0}(s) \\ \frac{dz}{ds}(s) = (1 + d(s))\left(f(z^{0}(s), z(s))w^{0}(s) + \sum_{j=1}^{m} g_{j}(z^{0}(s), z(s))w^{j}(s)\right) \\ \frac{d\nu}{ds}(s) = (1 + d(s))|w(s)| \\ (z^{0}(0), z(0), \nu(0)) = (0, \bar{x}, 0), \qquad (z^{0}(\hat{S}), z(\hat{S}), \nu(\hat{S})) \in \bar{\mathfrak{T}} \times [0, K] \end{cases}$$

$$(7.42)$$

The other objects simplify accordingly, and, still because of the concatenation property, the resulting family  $\mathcal{V}$  is abundant in the resulting  $\mathcal{W}$ : Therefore, by applying the infimum-gap result stated in Corollary 5.1 we get:

**Theorem 7.2.** Consider a feasible extended process

$$\left(\left(y^0, y, \nu\right), \left(\left(\hat{w}^0, \hat{w}\right), \hat{d}\right)\right), \quad \hat{d} \equiv 0,$$

and assume that it satisfies the infimum gap condition. Then, for all QDQ approximating cones C to  $\mathfrak{T} := \overline{\mathfrak{T}} \times [0, K]$  at  $(\hat{y}^0, \hat{y}, \hat{\nu})(\hat{S})$ , there exist a number  $\beta \leq 0$ , an absolutely continuous path  $(\lambda^0, \lambda, \lambda^{\nu}) \in W^{1,1}([0, \hat{S}]; \mathbb{R}^{(1+n+1)})$  and a zero-measure subset  $I_0$  such that the following conditions hold true:

<sup>&</sup>lt;sup>17</sup>We have set  $\hat{f}(s) := f(\hat{y}^0(s), \hat{y}(s)), \hat{g}_j(s) := g_j(\hat{y}^0(s), \hat{y}(s))$ , for all  $s \in [0, \hat{S}]$  and i = 1, ..., m

$$\begin{aligned} (i) \quad & (\lambda^0, \lambda, \lambda^\nu) \neq 0; \\ (ii) \quad & \frac{d\lambda^0}{ds}(s) = -\lambda(s) \cdot \left(\frac{\partial f}{\partial y^0}(\hat{y}^0(s), \hat{y}(s))\hat{w}^0(s) + \sum_{j=1}^m \frac{\partial g_j}{\partial y^0}(\hat{y}^0(s), \hat{y}(s))\hat{w}^j(s)\right) \\ & \frac{d\lambda}{ds}(s) = -\lambda(s) \cdot \left(\frac{\partial f}{\partial y}(\hat{y}^0(s), \hat{y}(s))\hat{w}^0_i(s) + \sum_{j=1}^m \frac{\partial g_j}{\partial y}(\hat{y}^0(s), \hat{y}(s))\hat{w}^j_i(s)\right), \end{aligned}$$

for a.e.  $s \in [0, \hat{S}];$ 

$$\begin{aligned} (iii) \quad (1+d) \left[ \lambda^0(s) w^0 + \lambda(s) \cdot \left( \hat{f}(s) w^0 + \sum_{j=1}^m \hat{g}_j(s) w^j \right) + \beta |w| \right] \\ \leq \left[ \lambda^0(s) \hat{w}^0(s) + \lambda(s) \cdot \left( \hat{f}(s) \hat{w}^0(s) + \sum_{j=1}^m \hat{g}_j(s) \hat{w}^j(s) \right) + \beta |\hat{w}(s)| \right] \\ for \ every \ (w^0, w, d) \in \mathbf{W}^{nc} \times [-0.5, 0.5] \ and \ s \in [0, \hat{S}] \backslash I_0 \end{aligned}$$

(*iv*)  $(\lambda^0(\hat{S}), \lambda(\hat{S}), \beta) \in -C^{\perp}$ .

## A Appendix

### A.1 An example on why abundance is crucial

The following example, which is due to H.J. Sussmann,<sup>18</sup> shows how the abundance hypothesis plays crucial for the validity of Theorem 5.2.

Consider the families of controls  $\mathcal{V}\subset\mathcal{W}$  defined as

$$\mathcal{W} := L^1([0,1],[0,5]) \qquad \mathcal{V} := \left\{ v \in \mathcal{W} : \int_0^1 v(s) ds \neq 1 \right\},$$

and the optimal control problems

$$(P)_{\mathcal{V}} \begin{cases} \text{Minimize } y(1) \\ \text{over processes } (y, v)(\cdot) \in W^{1,1}([0, 1], \mathbb{R}) \times \mathcal{V} \\ \frac{dy}{ds}(s) = v(s), \quad \text{a.e. } s \in [0, 1] \\ y(0) = 0, \quad y(1) = 1. \end{cases}$$
$$(P)_{\mathcal{W}} \begin{cases} \text{Minimize } y(1) \\ \text{over processes } (y, w)(\cdot) \in W^{1,1}([0, 1], \mathbb{R}) \times \mathcal{W} \\ \frac{dy}{ds}(s) = w(s), \quad \text{a.e. } s \in [0, 1] \\ y(0) = 0, \quad y(1) = 1. \end{cases}$$

The process  $(\hat{y}, \hat{w})(s) := (s, 1)$  is a minimizer of the extended problem  $(P)_{\mathcal{W}}$ , with cost equal to 1. If we restrict the controls to the *original* family of controls  $\mathcal{V}$ , the cost of the problem raises to  $+\infty$ , since every solution y[v] with  $v \in \mathcal{V}$  fails to be feasible. In other words the process  $(\hat{y}, \hat{w})$  satisfies the infimum gap condition.

,

By applying the Pontryagin's Maximum Principle to the minimizer  $(\hat{y}, \hat{w})$  of  $(P)_{\mathcal{W}}$ , we get that there exist multipliers  $(\lambda(\cdot), \lambda_c) \neq (0, 0)$  such that

$$\frac{d\lambda}{ds}(s) \equiv 0, \qquad \lambda(s)\mathfrak{w} \leq \lambda(s) \qquad \forall \mathfrak{w} \in [0,5], \quad s \in [0,1].$$

In particular this implies  $\lambda(s) \equiv 0$  and  $\lambda_c > 0$ . Therefore, if we set h(y) := y for every  $y \in \mathbb{R}$ , the process  $(\hat{y}, \hat{w})$  turns out to be a normal h-extremal. Therefore, in view of

<sup>&</sup>lt;sup>18</sup>Personal communication.

Corollary 5.1 the set  $\mathcal{V}$ , though being dense in  $\mathcal{W}$ , cannot be abundant in  $\mathcal{W}$ . As a matter of fact, one can easily find a positive integer  $N, \delta > 0$  and N + 1 controls  $w, w_1, \ldots, w_N$ for which  $\theta_{w,w_1,\ldots,w_N}^{\delta} : \Gamma_N \to \mathcal{V}$  verifying the properties of Definition 4.1 does not exist. Indeed, consider  $\Gamma_1(=[0,1]), w(s) := 0, w_1(s) := 2, \forall s \in [0,1], \delta > 0$ , and take any mapping  $\theta^{\delta} : [0,1] \to \mathcal{V}$ . In view of Definition 4.1, one has

$$\lim_{\delta \to 0} \int_0^1 \theta^{\delta}(\gamma)(s) ds = \int_0^1 w(s) + \gamma \big( w_1(s) - w(s) \big) = 2\gamma \qquad \forall \gamma \in \Gamma_1.$$

Then, for every  $\delta$  sufficiently small, either there exists a  $\gamma_{\delta} \in [0, 1]$  such that

$$\int_0^1 \theta^\delta(\gamma_\delta)(s) ds = 1,$$

or the map  $\gamma \mapsto \int_0^1 \theta^{\delta}(\gamma)(s) ds$  is not continuous. Since the former case is ruled out by the fact that the map  $\theta^{\delta}(\cdot)(s)$  has to take values in  $\mathcal{V}$ , the map  $\gamma \mapsto \int_0^1 \theta^{\delta}(\gamma)(s) ds$  is not continuous, so providing a contradiction.

### References

- M. S. Aronna, J. F. Bonnans, P. Martinon, A well-posed shooting algorithm for optimal control problems with singular arcs, J. Optim. Theory Applic. 158, 2, 2013, pp. 419-459.
- [2] M.S. Aronna, M. Motta, F. Rampazzo, Infimum gaps for limit solutions. Set-Valued Var. Anal. 23, no. 1, 3–22, 2015.
- [3] M.S. Aronna, F. Rampazzo, L<sup>1</sup> limit solutions for control systems. J. Differential Equations 258, no. 3, 954–979, 2015.
- [4] A. Arutyunov, D. Karamzin and F. Pereira, Pontryagin's maximum principle for constrained impulsive control problems. Nonlinear Anal. 75, no. 3, 1045–1057, 2012.
- [5] A. Arutyunov, D. Karamzin and F. Pereira, A nondegenerate maximum principle for the impulse control problem with state constraints. SIAM J. Control Optim. 43, no. 5, 1812–1843, 2005.
- [6] D. Azimov, R. Bishop (2005) New trends in astrodynamics and applications: optimal trajectories for space guidance. Ann. New York Acad. Sci. 1065(1), 189–209.
- [7] Bonnans, J. F. and Shapiro, A. Perturbation Analysis of Optimization Problems, Springer, New York, 2000
- [8] A.Bressan, F. Rampazzo, On differential systems with vector-valued impulsive controls. Boll. Un. Mat. Ital. B (7) 2, no. 3, 641–656, 1988.
- [9] A. Bressan, B. Piccoli, Introduction to the mathematical theory of control. AIMS Series on Applied Mathematics, 2. American Institute of Mathematical Sciences (AIMS), Springfield, MO, 2007.
- [10] Aldo Bressan, Hyper-impulsive motions and controllizable coordinates for Lagrangean systems, Atti Accad. Naz. Lincei, Memorie, Serie VIII, Vol. XIX, 197–246, 1990.
- [11] Aldo Bressan, On some control problems concerning the ski or swing, Atti Accad. Naz. Lincei, Memorie, Serie IX, Vol. I, 147–196, 1991.
- [12] A. Catllá, D. Schaeffer, T. Witelski, E. Monson, A. Lin, (2008) On spiking models for synaptic activity and impulsive differential equations, SIAM Rev. 50(3), 553–569
- [13] V. A. Dykhta, The variational maximum principle and quadratic conditions for the optimality of impulse and singular processes. (Russian) Sibirsk. Mat. Zh. 35 (1994), no. 1, 70–82, ii; translation in Siberian Math. J. 35, no. 1, 65–76, 1994.

- [14] V. A. Dykhta Second order necessary optimality conditions for impulse control problem and multiprocesses. Singular solutions and perturbations in control systems (Pereslavl-Zalessky, 1997), 97–101, IFAC Proc. Ser., IFAC, Laxenburg, 1997.
- [15] A.L. Dontchev, T. Zolezzi, Well-posed optimization Springer-Verlag, New York, 1993.
- [16] A. Fryszowsky Continuous selections for a class of non-convex multivalued maps, Studia Math., vol. 76, 163 - 174, 1983
- [17] P. Gajardo, H. Ramirez C., A. Rapaport, Minimal time sequential batch reactors with bounded and impulse controls for one or more species, SIAM J. Control Optim. 47 (6) 2827–2856, 2008.
- [18] R. V. Gamkrelidze, On Some Extremal Problems in the Theory of Differential Equations with Applications to Optimal Control, SIAM Journal on Control, vol. 3, pp. 106-128, 1965.
- [19] M. Guerra, A. Sarychev, Fréchet generalized trajectories and minimizers for variational problems of low coercivity, J. Dyn. Control Syst. 21, no. 3, 351–377, 2015.
- [20] D.Y. Karamzin; V.A. de Oliveira, F.L. Pereira, G.N. Silva, On the properness of an impulsive control extension of dynamic optimization problems, ESAIM Control Optim. Calc. Var. 21, no. 3, 857–875, 2015.
- [21] B. Kaskosz, Extremality, controllability and abundant subsets of generalized control systems, Journal of Optimization Theory and Applications, 101, (1999), pp. 73-108.
- [22] A. C. Lai, M. Motta and F. Rampazzo, Minimum restraint functions for unbounded dynamics: general and control-polynomial systems, Pure and Applied Functional Analysis, Volume 1, Number 4, pp. 583-612.
- [23] A. Mariconda, Contractibility and Fixed Point Property: the case of Decomposable Sets, Nonlinear Analysis, T.M.A., vol. 17, n. 7, pp. 689-695, 1992.
- [24] M. Miller, E. Y. Rubinovich, Impulsive control in continuous and discrete-continuous systems. Kluwer Academic/Plenum Publishers, New York, 2003.
- [25] M. Motta, F. Rampazzo, Space-time trajectories of nonlinear systems driven by ordinary and impulsive controls. Differential Integral Equations 8, no. 2, 269–288, 1995.
- [26] M. Motta, F. Rampazzo, Dynamic programming for nonlinear systems driven by ordinary and impulsive controls. SIAM J. Control Optim. 34, no. 1, 199–225, 1996.
- [27] M. Motta, F. Rampazzo, State-constrained control problems with neither coercivity nor L<sup>1</sup> bounds on the controls. Ann. Mat. Pura Appl. (4) 177, 117–142, 1999.
- [28] M. Motta, F. Rampazzo, R. B. Vinter, Normality and Gap Phenomena in Optimal Unbounded Control, to appear on ESAIM-COCV.
- [29] M. Motta, C. Sartori, On asymptotic exit-time control problems lacking coercivity. ESAIM Control Optim. Calc. Var. 20, no. 4, 957–982, 2014.
- [30] C. Olech, Decomposability as a substitute of convexity, Lecture Notes in Mathematics "Multifunctions and Integrands", vol. 1091, 193-2015, 1984.
- [31] M. Palladino, F. Rampazzo A No Infimum-Gap Criterion, to appear on Annual Conference on Conference on Decision and Control (CDC) 2019, 58th.
- [32] M. Palladino, R. B. Vinter, *Minimizers that are not also Relaxed Minimizers*, SIAM J. Control and Optim., 52 (2014), no. 4, pp. 2164 2179.
- [33] M. Palladino and R. B. Vinter, When are Minimizing Controls also Minimizing Relaxed Controls?, Discrete Contin. Dyn. Syst. Series A, 35 (2015), no. 9, pp. 4573 -4592.
- [34] M. Palladino and R. B. Vinter, When does relaxation reduce the minimum cost of an optimal control problem?, Annual Conference on Decision and Control (CDC), 2013 IEEE 52nd.

- [35] H. Schättler, U. Ledzewicz, Geometric Optimal Control: Theory, Methods and Examples, Springer, 2012.
- [36] G. Scorza Dragoni, Un teorema sulle funzioni continue rispetto ad una e misurabili rispetto ad un'altra variabile, (Italian) Rend. Sem. Mat. Univ. Padova 17 (1948), 102-106.
- [37] G. Silva, R. Vinter, Measure driven differential inclusions, J. Math. Anal. Appl., 202 no. 3, 727–746, 1996.
- [38] G. Silva, R. Vinter, Necessary conditions for optimal impulsive control problems. SIAM J. Control Optim. 35, no. 6, 1829–1846, 1997.
- [39] H.J. Sussmann, Geometry and Optimal Control, Mathematical Control Theory, J. Baillieul and J. C. Willems, Eds., Springer-Verlag, New York, 1998, pp.140-198.
- [40] H.J. Sussmann, Generalized differentials, variational generators, and the maximum principle with state constraints, Mathematical Control Theory, J. Baillieul and J. C. Willems, Eds., Springer-Verlag, New York, 1998, pp.140-198.
- [41] R. B. Vinter, Optimal Control, Birkhäuser, Boston, 2000.
- [42] J. Warga, Normal Control Problems have no Minimizing Strictly Original Solutions, Bulletin of the Amer. Math. Soc., 77, 4, 1971, pp. 625-628.
- [43] J. Warga, Optimal Control of Differential and Functional Equations, Academic Press, New York, 1972.
- [44] J. Warga, Controllability, extremality, and abnormality in nonsmooth optimal control, J. Optim. Theory and Applic., 41, 1, 1983, pp. 239-260.
- [45] J. Warga, Optimization and Controllability Without Differentiability Assumptions, SIAM J. Control and Optim., 21, 6, 1983, pp. 837-855.
- [46] P. Wolenski, S. Zabić, A differential solution concept for impulsive systems, Dyn. Contin. Discrete Impuls. Syst. Ser. A Math. Anal., 13B, 199–210, 2006.