

On multivariate polynomials with many roots over a finite grid

Olav Geil
Department of Mathematical Sciences
Aalborg University

Abstract

In this note we consider roots of multivariate polynomials over a finite grid. When given information on the leading monomial with respect to a fixed monomial ordering, the footprint bound [8, 5] provides us with an upper bound on the number of roots, and this bound is sharp in that it can always be attained by trivial polynomials being a constant times a product of an appropriate combination of terms consisting of a variable minus a constant. In contrast to the one variable case, there are multivariate polynomials attaining the footprint bound being not of the above form. This even includes irreducible polynomials. The purpose of the note is to determine a large class of polynomials for which only the mentioned trivial polynomials can attain the bound, implying that to search for other polynomials with the maximal number of roots one must look outside this class.

Keywords: finite field, finite grid, footprint bound, multivariate polynomial, root, variety

MSC: 11C08; 11G25; 11R09; 11T06;

1 Introduction

As is well-known, the number of roots of a univariate non-zero polynomial $F(X)$ over a field \mathbb{F} is upper bounded by its degree, and if $a \in \mathbb{F}$ is a root then $X - a$ divides F . For multivariate polynomials the situation is very different in that such polynomials often possess infinitely many roots when the field is not finite, and that (a_1, \dots, a_m) can be a root of $F(X_1, \dots, X_m)$ without F being divisible by any $X_\ell - a_\ell$, $\ell \in \{1, \dots, m\}$.

In many applications, however, the point set under consideration is finite, e.g. it corresponds to a grid $S_1 \times \dots \times S_m$ where S_ℓ , $\ell = 1, \dots, m$ are finite subsets of \mathbb{F} , one particular case being the grid \mathbb{F}_q^m where \mathbb{F}_q denotes the finite field with q elements. For finite grids, of course, the number of roots of a polynomial F becomes finite and the Schwartz-Zippel lemma provides us with an upper bound using information on the total degree of F . Having knowledge of the leading monomial with respect to some fixed monomial ordering, the footprint bound produces even more precise information, as we recall in a moment.

One is often interested in determining polynomials which attain the maximal number of roots according to the above bounds. For instance this is the case when one wants to produce good algebraic geometric codes [19, 9] or when trying to determine minimal weight code words of generalized Reed-Muller codes and their relatives, e.g. [3, 12, 17, 6, 15, 14, 7, 11]. As it turns out a way to produce polynomials with the

maximal number of roots according to the footprint bound is to take a non-zero constant times a product of terms of the form $X_\ell - a$, but as already hinted this trivial construction will typically not give us all the desired polynomials. For instance it does not produce any irreducible polynomials of degree more than one. In the present note we determine a large class of polynomials for which the above type of trivial polynomials are exactly those attaining the maximal number of roots, implying that to search for other polynomials with the maximal number of roots one must look outside this class.

The note is organized as follows. In Section 2 we recall the footprint bound for a single polynomial and demonstrate its sharpness. Then in Section 3 we give necessary conditions for non-trivial polynomials to attain the footprint bound.

2 The footprint bound

The footprint bound [8, 5] is a general method to upper bound the size of varieties [2]. For a single polynomial $F(X_1, \dots, X_m)$ and the point set being a finite grid one obtains a simple closed formula expression. This formula assumes that $\deg_{X_\ell} \text{lm}(F) < s_\ell$, $\ell = 1, \dots, m$, where s_ℓ denotes the size of S_ℓ and where $\text{lm}(F)$ denotes the leading monomial of F . Of course this condition in particular is satisfied when not only the leading monomial, but all monomials in the support satisfy the conditions on the degree which we will often assume throughout the note. Observe that even the last assumption is no real restriction as $F(X_1, \dots, X_m)$ has exactly the same roots over the finite grid [1] as does

$$F(X_1, \dots, X_m) \text{ rem } \left\{ \prod_{\alpha \in S_1} (X_1 - \alpha), \dots, \prod_{\alpha \in S_m} (X_m - \alpha) \right\}, \quad (1)$$

where the latter notation means the remainder modulo the polynomials in the curly brackets. This remainder is produced using the multivariate division algorithm [2]. It is clear that (1) satisfies the requirement on the degree for each monomial in the support of it and we therefore introduce the following notation for the set of remainders:

$$\mathbb{F}[X_1, \dots, X_m]_{<(s_1, \dots, s_m)} = \{F(X_1, \dots, X_m) \in \mathbb{F}[X_1, \dots, X_m] \mid \deg_{X_\ell} F < s_\ell, \text{ for } \ell = 1, \dots, m\}.$$

The footprint bound for a single polynomial now is [8, 5]:

Theorem 1 *Let the notation be as above and consider an arbitrary, but fixed, monomial ordering on the set of monomials in the variables X_1, \dots, X_m . For a non-zero polynomial $F(X_1, \dots, X_m) \in \mathbb{F}[X_1, \dots, X_m]$ write its leading monomial as $X_1^{i_1} \dots X_m^{i_m}$ and assume $i_1 < s_1, \dots, i_m < s_m$ (e.g. $F(X_1, \dots, X_m) \in \mathbb{F}[X_1, \dots, X_m]_{<(s_1, \dots, s_m)}$). Then F possesses at most $D(X_1^{i_1} \dots X_m^{i_m}) = s_1 \dots s_m - (s_1 - i_1) \dots (s_m - i_m)$ roots over $S_1 \times \dots \times S_m$.*

We remark that the Schwartz-Zippel lemma [13, 4, 16, 21] can be obtained as a corollary. This bound states that over a finite grid $S \times \dots \times S$, a polynomial of total degree $t < s$ can at most have ts^{m-1} roots where $s = \#S$.

It is not difficult to see that Theorem 1 is sharp in that for any prescribed leading monomial corresponding polynomials exist having as many roots as the upper bound. Namely, consider any subsets

$$S'_\ell \subseteq S_\ell, i_\ell = \#S'_\ell, \ell = 1, \dots, m \quad (2)$$

and $k \in \mathbb{F} \setminus \{0\}$. Then the trivial polynomial

$$k \prod_{\ell=1}^m \prod_{\alpha \in S'_\ell} (X_\ell - \alpha) \quad (3)$$

has exactly $(s_1 - i_1) \cdots (s_m - i_m)$ non-roots in the finite grid and therefore the number of roots as predicted in Theorem 1, as obviously for any monomial ordering the leading monomial of (3) equals $X_1^{i_1} \cdots X_m^{i_m}$. Observe that the polynomial in (3) is an example of a polynomial satisfying that if $(a_1, \dots, a_m) \in S_1 \times \cdots \times S_m$ is a root then some $X_\ell - a_\ell$ divides it.

The polynomials described in (3) are by no means the only ones producing equality in the footprint bound. We illustrate this observation with a classic example.

Example 1 *Let q be a prime power and consider the point set $\mathbb{F}_{q^2} \times \mathbb{F}_{q^2}$. The Hermitian polynomial $F(X_1, X_2) = X_1^{q+1} - X_2^q - X_2$ has q^3 roots which is exactly the upper bound from Theorem 1 when choosing a monomial ordering such that X_2^q becomes the leading monomial. The roots are established by using the fact that X_1^{q+1} is the norm function related to the field extension $\mathbb{F}_{q^2}/\mathbb{F}_q$ and that $X_2^q + X_2$ is the similar trace function. Employing the properties of these functions one determines [18, 20] the roots. Clearly, for no $(a_1, a_2) \in \mathbb{F}_{q^2} \times \mathbb{F}_{q^2}$ it holds that $X_1 - a_1$, nor that $X_2 - a_2$, divides $F(X_1, X_2)$. Actually the Hermitian polynomial is absolutely irreducible.*

In the next section we derive information on when a polynomial can possibly meet the footprint bound without being of the form (3).

3 Necessary conditions for attaining the footprint bound

We start our investigations with a simple, yet crucial lemma.

Lemma 2 *Let $F(X_1, \dots, X_m) \in \mathbb{F}[X_1, \dots, X_m]$ and $a \in S_\ell$. Then the following bi-implication holds true:*

$$\begin{aligned} & X_\ell - a \text{ divides } F(X_1, \dots, X_m) \\ \Leftrightarrow & \\ & (a_1, \dots, a_{\ell-1}, a, a_{\ell+1}, \dots, a_m) \text{ is a root of } F(X_1, \dots, X_m) \\ & \text{for all } (a_1, \dots, a_{\ell-1}, a_{\ell+1}, \dots, a_m) \in S_1 \times \cdots \times S_{\ell-1} \times S_{\ell+1} \times \cdots \times S_m \end{aligned}$$

Proof: Without loss of generality we assume that $F(X_1, \dots, X_m) \in \mathbb{F}[X_1, \dots, X_m]_{<(s_1, \dots, s_m)}$ (if this is not the case, we first perform reduction modulo $\{\prod_{\alpha \in S_1} (X_1 - \alpha), \dots, \prod_{\alpha \in S_m} (X_m - \alpha)\}$). When F is the zero polynomial then the bi-implication clearly holds. Hence,

assume that F is not the zero polynomial. The "only if" part is easily verified. To see the "if" part we assume that F has all the requested roots and apply the division algorithm to obtain $F(X_1, \dots, X_m) = Q(X_1, \dots, X_{\ell-1}, X_{\ell+1}, \dots, X_m)(X_\ell - a) + R(X_1, \dots, X_{\ell-1}, X_{\ell+1}, \dots, X_m)$. Aiming for a contradiction assume the R is not the zero-polynomial. Observe, that all monomials M in the support of R satisfy $\deg_{X_i} M < s_i$, for all $i \in \{1, \dots, \ell-1, \ell+1, \dots, m\}$ and in particular this holds for the leading monomial with respect to any monomial ordering (the leading monomial exists by assumption). According to the footprint bound applied to the point set $S_1 \times \dots \times S_{\ell-1} \times S_{\ell+1} \times \dots \times S_m$, R therefore cannot have all elements of this point set as roots. But $Q(X_1, \dots, X_{\ell-1}, X_{\ell+1}, \dots, X_m)(X_\ell - a)$ has all the requested roots of the lemma and therefore also this holds for R , which is a contradiction. \square

We shall need two more lemmas.

Lemma 3 Consider $F(X_1, \dots, X_m) \in \mathbb{F}[X_1, \dots, X_m]$ with $\deg_{X_\ell} \text{lm}(F) < s_\ell$, $\ell = 1, \dots, m$ with respect to some fixed monomial ordering (e.g. $F(X_1, \dots, X_m) \in \mathbb{F}[X_1, \dots, X_m]_{<(s_1, \dots, s_m)}$). Write

$$F(X_1, \dots, X_m) = G(X_1, \dots, X_m)H(X_1, \dots, X_m)$$

where

$$G(X_1, \dots, X_m) = \prod_{i=1}^m \prod_{\alpha \in S'_i} (X_i - \alpha)$$

for some $S'_\ell \subseteq S_\ell$, $\ell = 1, \dots, m$ and write $T_\ell = S_\ell \setminus S'_\ell$ and $t_\ell = \#T_\ell$. Then F attains the footprint bound over $S_1 \times \dots \times S_m$ if and only if H attains the footprint bound over $T_1 \times \dots \times T_m$.

Proof:

The set of non-roots of G over $S_1 \times \dots \times S_m$ equals $T_1 \times \dots \times T_m$. Hence, the set of non-roots of F over $S_1 \times \dots \times S_m$ equals the set of non-roots of H over $T_1 \times \dots \times T_m$. Let $\text{lm}(F) = X_1^{i_1} \dots X_m^{i_m}$, then $\text{lm}(H) = X_1^{i_1 - s'_1} \dots X_m^{i_m - s'_m}$ and according to the footprint bound therefore the number of non-roots of H over $T_1 \times \dots \times T_m$ is at least

$$(t_1 - (i_1 - s'_1)) \cdots (t_m - (i_m - s'_m)) = (s_1 - i_1) \cdots (s_m - i_m)$$

and exactly when equality holds the number of roots of F over $S_1 \times \dots \times S_m$ becomes $s_1 \cdots s_m - (s_1 - i_1) \cdots (s_m - i_m) = D(\text{lm}(F))$. \square

Example 2 Recall [10], that the trace map $\text{Tr} : \mathbb{F}_{q^2} \rightarrow \mathbb{F}_q$ is given by $\text{Tr}(\alpha) = \alpha^q + \alpha$ and that the preimage of any element in \mathbb{F}_q is of size exactly q . Now let $F(X_1, X_2) = G(X_1, X_2)H(X_1, X_2) \in \mathbb{F}_{q^2}[X_1, X_2]$ where

$$G(X_1, X_2) = \prod_{\text{Tr}(\alpha)=0} (X_1 - \alpha) \prod_{\text{Tr}(\alpha)=0} (X_2 - \alpha)$$

and

$$H(X_1, X_2) = \text{Tr}(X_1) - \text{Tr}(X_2) = X_1^q - X_2^q + X_1 - X_2.$$

We apply Lemma 3 to show that this polynomial attains the footprint bound over $S_1 \times S_2$, where $S_1 = S_2 = \mathbb{F}_{q^2}$. The leading monomial of F clearly is $X_1^{2q}X_2^q$ or $X_1^qX_2^{2q}$, respectively, depending on the choice of monomial ordering. Hence, according to the footprint bound F potentially has $q^4 - (q^2 - 2q)(q^2 - q) = 3q^3 - 2q^2$ roots. The non-roots of $G(X_1, X_2)$ are $T_1 \times T_2$ where $T_1 = T_2 = \{\alpha \mid \text{Tr}(\alpha) \neq 0\}$. Therefore, according to Lemma 3, F attains the footprint bound over $S_1 \times S_2$ if and only if H attains the footprint bound over $T_1 \times T_2$ meaning that H has $(q^2 - q)^2 - (q^2 - 2q)(q^2 - q) = q^3 - q^2$ roots in this set. This is exactly the case, the roots being

$$\{(\alpha, \beta) \mid \text{Tr}(\alpha) = \text{Tr}(\beta) \neq 0\}.$$

Hence, $F(X_1, X_2)$ does attain the footprint bound possessing $3q^3 - 2q^2$ roots over $S_1 \times S_2 = \mathbb{F}_{q^2} \times \mathbb{F}_{q^2}$.

Lemma 4 Given the point set $S_1 \times \dots \times S_m$ let D be the already introduced map and let D' be the map from $\{X_1^{i_1} \dots X_m^{i_m} \mid i_\ell < s_\ell, \ell = 1, \dots, m\}$ to \mathbb{N}_0 defined by

$$D'(X_1^{i_1} \dots X_m^{i_m}) = s_1 \dots s_{m-1} - (s_1 - i_1) \dots (s_{m-1} - i_{m-1}).$$

Then for all monomials $M, N \in \{X_1^{i_1} \dots X_m^{i_m} \mid i_\ell < s_\ell, \ell = 1, \dots, m\}$ we have:

1. if $\deg_{X_m} M \geq 1$ then $D(M) > s_m D'(M)$
2. if N divides M and $N \neq M$ then $D(N) < D(M)$.

Proof:

By inspection. □

We are now ready for our first result on which polynomials can possibly attain the footprint bound.

Theorem 5 Consider a non-constant polynomial $H(X_1, \dots, X_m) \in \mathbb{F}[X_1, \dots, X_m]_{<(s_1, \dots, s_m)}$ with no factor $X_\ell - a$, $a \in S_\ell$ for any ℓ . Define

$$\Omega = \max\{D(M) \mid M \in \text{Supp}(H)\}$$

where $\text{Supp}(H)$ denotes the support of H , and let $\{M_1, \dots, M_\mu\}$ be the monomials in the support with D -value equal to Ω . Then necessary conditions for H to have Ω roots over $S_1 \times \dots \times S_m$ are that:

1. for any monomial ordering the leading monomial of H belongs to $\{M_1, \dots, M_\mu\}$
2. $\gcd(M_1, \dots, M_\mu) = 1$

Proof:

We only prove that condition 2 is needed. If $\gcd(M_1, \dots, M_\mu) \neq 1$ then there exists some X_ℓ which divides M_1, \dots, M_μ . Without loss of generality assume $\ell = m$. For each $a \in S_m$ we then consider

$$H'(X_1, \dots, X_{m-1}) = H(X_1, \dots, X_{m-1}, a)$$

which according to Lemma 2 is non-zero by the assumption that H has no factor of the form $X_\ell - a$. Let N be the leading monomial of H' with respect to some monomial ordering on the set of monomials in the variable X_1, \dots, X_{m-1} . We now claim that $D'(N) < \Omega/s_m$. If N divides M for some $M \in \{M_1, \dots, M_\mu\}$ then by 1. in Lemma 4 we obtain $\Omega > s_m D'(M)$ which in turn is larger than or equal to $s_m D'(N)$ by 2. in the same lemma. If N does not divide any monomial in $\{M_1, \dots, M_\mu\}$ then for any M in the support of H such that N divides M it holds that $s_m D'(N) \leq s_m D'(M) \leq D(M)$ which by assumption is strictly smaller than Ω . As N necessarily divides some monomial in the support of H this proves the claim. By Theorem 1 for each a the corresponding leading monomial N of H' gives us the upper bound $D'(N)$ on the number of roots of H' over $S_1 \times \dots \times S_{m-1}$. Summing up the contribution from each $a \in S_m$ we obtain fewer than Ω roots in total of H over $S_1 \times \dots \times S_m$. \square

Example 3 This is a continuation of Example 2 where we considered the polynomial $H(X_1, X_2) = X_1^q - X_2^q + X_1 - X_2$ attaining the footprint bound over $T_1 \times T_2$, with $T_1 = T_2 = \{\alpha \in \mathbb{F}_{q^2} \mid \text{Tr}(\alpha) \neq 0\}$. Clearly, H has no factor $X_\ell - a$, and therefore by 2. in Theorem 5 we need to have $\gcd(X_1^q, X_2^q) = 1$ which indeed is the case. Furthermore, in accordance with 1. in Theorem 5 we have $\Omega = D(X_1^q) = D(X_2^q)$.

Remark 6 If the polynomial H in Theorem 5 contains exactly one monomial with the highest D -value then condition 2. is never satisfied (here, we used the assumption from Theorem 5 that H is not a constant).

Theorem 7 Let $F(X_1, \dots, X_m) \in \mathbb{F}[X_1, \dots, X_m]_{<(s_1, \dots, s_m)}$ be a polynomial having a unique monomial M in its support of highest D -value and assume that for some monomial ordering this is the leading monomial. Then it is either of the form (3) or it possesses fewer than $D(M)$ roots.

Proof:

Assume that F has $D(M)$ roots. Then F is square-free and we may write $F = GH$ as in Lemma 3. We will show that H is a constant. The crucial observation is that $\text{lm}(H)$ is the unique monomial in the support of H such that the D -value with respect to the point set $T_1 \times \dots \times T_m$ is maximal. But then the assumption that F has $D(M)$ roots by Lemma 3, 2. in Theorem 5, and Remark 6 implies that H is a constant. \square

We present two corollaries to Theorem 7. The first concerns a family of polynomials which we call *monomial ordering invariant*.

Definition 8 A polynomial is said to be *monomial ordering invariant* if it in its support has a monomial M which is divisible by any (other) monomial in the support.

Clearly, for a monomial ordering invariant polynomial F and any choice of monomial ordering the leading monomial of F equals the M in Definition 8 and by 2. in Lemma 4 the D -value of M is strictly larger than the D -value of any other monomial in the support of F . As the name indicates there are no other polynomials besides the monomial ordering invariant ones having a unique leading monomial. Let namely

$M = X_1^{i_1} \cdots X_m^{i_m}$ be the leading monomial of F with respect to some monomial ordering and aiming for a contradiction assume that F in its support has a polynomial $N = X_1^{i'_1} \cdots X_m^{i'_m}$ with $i_t < i'_t$ for some $t \in \{1, \dots, m\}$. But then for a lexicographic ordering with X_t larger than the other variables it holds that N is larger than M and therefore M is not the leading monomial with respect to this ordering. Observe that univariate non-zero polynomials as well as the polynomials in (3) satisfy the condition for being monomial ordering invariant. As an immediate corollary to Theorem 7 we obtain:

Corollary 9 *Let $F(X_1, \dots, X_m) \in \mathbb{F}[X_1, \dots, X_m]_{<(s_1, \dots, s_m)}$ be a monomial ordering invariant polynomial. Then over $S_1 \times \cdots \times S_m$ the footprint bound is the same for all choices of monomial ordering and when attained, $F(X_1, \dots, X_m)$ is necessarily of the form (3).*

Remark 10 *Recall from the above discussion that a non-zero polynomial which is not monomial ordering invariant must have in its support at least two different monomials each being the leading monomial with respect to some monomial ordering. But then according to the footprint bound if $F(X_1, \dots, X_m) \in \mathbb{F}[X_1, \dots, X_m]_{<(s_1, \dots, s_m)}$ is not monomial ordering invariant then the number of roots cannot attain the maximal value of $D(M)$ for M in the support of it when the polynomial has only one monomial in its support with this D -value. Hence, the non-trivial information given in Theorem 7 is actually that of Corollary 9.*

The last corollary to Theorem 7 concerns irreducible polynomials.

Corollary 11 *Let $F(X_1, \dots, X_m) \in \mathbb{F}_q[X_1, \dots, X_m]_{<(q, \dots, q)}$ be an irreducible polynomial different from $k(X_\ell - a)$, where $k \in \mathbb{F}_q \setminus \{0\}$, and with s roots from the point set $S_1 \times \cdots \times S_m = \mathbb{F}_q^m$. Then at least one of the following two conditions are satisfied:*

1. F contains in its support at least two monomials with D -value equal to s
2. F contains in its support a monomial of D -value strictly larger than s .

Example 4 *This is a continuation of Example 1 where we considered the Hermitian polynomial $X_1^{q+1} - X_2^q - X_2$ which attains the footprint bound, i.e. it possesses $D(X_2^q) = q^3$ roots over \mathbb{F}_{q^2} (here we use a monomial ordering with X_2^q larger than X_1^{q+1}). As already noted the Hermitian polynomial is (absolutely) irreducible, and indeed it satisfies condition 2. of Corollary 11 as $D(X_1^{q+1}) = q^3 + q^2 > q^3$.*

Remark 12 *One can interpret Theorem 5, Theorem 7 and Corollary 9 as results on which polynomials different from (3) can possibly attain the footprint bound over a finite grid.*

Acknowledgments

The author would like to thank René Bødker Christensen for comments on a previous version of the paper.

References

- [1] N. Alon. Combinatorial nullstellensatz. *Comb. Prob. Comput.*, 8(1-2):7–29, 1999.
- [2] D. A. Cox, J. Little, and D. O’Shea. *Ideals, varieties, and algorithms: an introduction to computational algebraic geometry and commutative algebra*, volume 10. Springer, 1997.
- [3] P. Delsarte, J. M. Goethals, and F. J. Mac Williams. On generalized Reed-Muller codes and their relatives. *Information and Control*, 16(5):403–442, 1970.
- [4] R. A. DeMillo and R. J. Lipton. A probabilistic remark on algebraic program testing. Technical report, Georgia Institute of Technology, Atlanta School of Information and Computer Science, 1977.
- [5] O. Geil and T. Høholdt. Footprints or generalized Bezout’s theorem. *IEEE Trans. Inform. Theory*, 46(2):635–641, 2000.
- [6] O. Geil and T. Høholdt. On hyperbolic codes. In S. Boztaş and I. E. Shparlinski, editors, *Applied Algebra, Algebraic Algorithms and Error-Correcting Codes*, volume 2227 of *Lecture Notes in Comput. Sci.*, pages 159–171. Springer, 2001.
- [7] O. Geil and C. Thomsen. Weighted Reed-Muller codes revisited. *Des. Codes Cryptogr.*, 66(1-3):195–220, 2013.
- [8] T. Høholdt. On (or in) Dick Blahut’s ‘footprint’. *Codes, Curves and Signals*, pages 3–9, 1998.
- [9] T. Høholdt, J. H. van Lint, and R. Pellikaan. Algebraic geometry codes. In V. S. Pless and W. C. Huffman, editors, *Handbook of Coding Theory*, volume 1, pages 871–961. Elsevier, Amsterdam, 1998.
- [10] R. Lidl and H. Niederreiter. *Introduction to finite fields and their applications*. Cambridge University Press, Cambridge, 1986.
- [11] H. H. López, C. Rentería-Márquez, and R. H. Villarreal. Affine Cartesian codes. *Des. Codes Cryptogr.*, 71(1):5–19, 2014.
- [12] J. L. Massey, D. J. Costello, and J. Justesen. Polynomial weights and code constructions. *IEEE Trans. Inform. Theory*, 19(1):101–110, 1973.
- [13] Ø. Ore. Über höhere kongruenzen. *Norsk Mat. Forenings Skrifter Ser. I*, 7(15):27, 1922.
- [14] D. Ruano. On the parameters of r-dimensional toric codes. *Finite Fields Appl.*, 13(4):962–976, 2007.
- [15] D. Ruano. On the structure of generalized toric codes. *J. Symbolic Comput.*, 44(5):499–506, 2009.
- [16] J. T. Schwartz. Fast probabilistic algorithms for verification of polynomial identities. *J ACM*, 27(4):701–7017, 1980.

- [17] A. B. Sørensen. Weighted Reed-Muller codes and algebraic-geometric codes. *IEEE Trans. Inform. Theory*, 38(6):1821–1826, 1992.
- [18] H. Stichtenoth. A note on Hermitian codes over $\text{GF}(q^2)$. *IEEE Trans. Inform. Theory*, 34(5):1345–1348, 1988.
- [19] H. Stichtenoth. *Algebraic function fields and codes*. Universitext. Springer-Verlag, Berlin, 1993.
- [20] K. Yang and P. V. Kumar. On the true minimum distance of Hermitian codes. In *Coding theory and algebraic geometry*, pages 99–107. Springer, 1992.
- [21] R. Zippel. Probabilistic algorithms for sparse polynomial. In *Proceedings of International Symposium on Symbolic and Algebraic Computations*, pages 216–226, 1979.