

# REALISING A FINITE GROUP AS A SUBGROUP OF A PRODUCT OF TWO GROUPS OF PERMUTATION MATRICES

MAHMOUD BENKHALIFA

ABSTRACT. In this paper we prove that any finite group of order  $n$  can be viewed as the group of the solutions of a certain matrix equation  $XB = BY$ , where the unknowns  $X, Y$  are two permutation matrices of order  $n$  and  $(1+k)n+2$  respectively and where  $k \in \mathbb{N}$  is given by Cayley's theorem. Moreover, we show that  $G$  is isomorphic to a certain subgroup formed by permutation matrices of order  $(1+k)n$  obtained by permuting all the rows of the identity matrix  $I_{(1+k)n}$ .

## 1. INTRODUCTION

Let  $\mathcal{P}(n)$  denote the group of permutation matrices of degree  $n$ . For a given matrix  $B$ , let us consider the group  $\Omega_B$  of the pairs  $(X, Y) \in \mathcal{P}(n) \times \mathcal{P}(m)$  which are solutions of the matrix equation  $XB = BY$ . Obviously,  $\Omega_B$  is finite group as  $\mathcal{P}(n)$  and  $\mathcal{P}(m)$  are finite and it is worth noting that if  $\lambda \in \mathbb{Q}$  and  $(X, Y) \in \Omega_B$ , then the pair  $(\lambda X, \lambda Y)$  needs not be in  $\Omega_B$  although that we have  $(\lambda X)B = B(\lambda Y)$  because  $\lambda X, \lambda Y$  are not permutation matrices for  $\lambda \neq 1$ .

A subgroup  $H$  of  $\mathcal{P}(n)$  is called *realisable* if each element  $M \in H$  is obtained by permuting the rows of the identity matrix  $I_n$  using a permutation  $\tau \in S_n$  satisfying  $\tau(i) \neq i$  for all  $1 \leq i \leq n$ . Here  $S_n$  denotes the symmetric group of order  $n$ .

Recall that, by Cayley's theorem, any finite group  $G$  of order  $n$  is isomorphic to realisable subgroup, denoted by  $\mathcal{C}_G$ , of  $\mathcal{P}(n)$  via the map

$$G = \{g_1, \dots, g_n\} \rightarrow S_n \cong \mathcal{P}(n) \quad , \quad g_j \mapsto \sigma_j = \begin{pmatrix} g_1 & g_2 & \dots & g_n \\ g_j & \sigma_j(g_2) & \dots & \sigma_j(g_n) \end{pmatrix} \leftrightarrow M_j$$

where  $M_j$  is the matrix obtained by permuting all the rows of the identity matrix  $I_n$  using  $\sigma_j$ .

Following the idea developed in [8] and inspired by the works done in [5, 7] regarding the so-called Kahn's realisability problem of groups ( see [9, 12] for more details), this paper is devoted to answer the question whether a given finite group  $G$  can occur as a group on the form  $\Omega_B$  and whether  $G$  can be embedded in  $\mathcal{P}(m)$ , where  $m > n$ , as a realisable subgroup. For this purpose we shall assign to  $G$  a matrix  $B_G$  and a realisable subgroup  $\mathcal{A}_G$  of  $\mathcal{P}((1+k)n+2)$ , where the number  $k$  is given by the decomposition of the permutation  $\sigma_2$  into product of disjoint cycles, i.e.,  $\sigma_2 = \tau_1 \tau_2 \dots \tau_k$  and we shall define  $\Omega_{B_G}$  as a certain subgroup of  $\mathcal{A}_G \times \mathcal{C}_G$ .

The group  $\mathcal{A}_G$  and the matrix  $B_G$  are defined using the framework of rational homotopy theory [11] and the ideas developed in [6, 8]. More precisely,  $\mathcal{A}_G$  is defined in

2000 *Mathematics Subject Classification.* 15A24, 55P10.

*Key words and phrases.* Finite groups, Permutation matrices, Graded commutative  $\mathbb{Q}$ -algebras, Group of homotopy self-equivalences, Matrix equation.

terms of the cohomology of a certain a free commutative cochain  $\mathbb{Q}$ -algebra associated with the group  $G$  and  $B_G$  is related to its differential.

Thus, in this paper we establish the following result

**Theorem 1.** *For any finite group  $G$  of order  $n$ , there exists a matrix  $B_G$  such that  $G$  is isomorphic to the group  $\Omega_{B_G}$  of the solutions of the matrix equation  $XB_G = B_GY$ , where the unknowns  $X, Y$  are two permutation matrices belonging to the groups  $\mathcal{A}_G$  and  $\mathcal{C}_G$  respectively.*

As a corollary we derive

**Corollary 2.** *Any finite group  $G$  of order  $n$  is isomorphic to a realisable subgroup of  $\mathcal{P}((1+k)n)$ .*

## 2. MAIN RESULTS

**2.1. Definition of the group  $\mathcal{A}_G$ .** Let us start by recalling the main construction in [8] on which this work is based. Indeed, let  $G = \{g_1, g_2, \dots, g_n\}$  be a finite group of order  $n$  and let  $S_n$  the symmetric group. By Cayley's theorem there is a monomorphism

$$\Psi : G \rightarrow S_n \quad , \quad g_j \mapsto \sigma_j : g_k \longrightarrow g_j g_k \quad , \quad 1 \leq k \leq n$$

For  $2 \leq j \leq n$ , write  $\sigma_j = \begin{pmatrix} 1 & 2 & \dots & n \\ j & \sigma_j(2) & \dots & \sigma_j(n) \end{pmatrix}$  and let:

$$\sigma_2 = \left( 1 2 \sigma_2(2) \dots, \sigma_2^{\kappa_2}(2) \right) \left( i_1 \sigma_2(i_1) \dots, \sigma_2^{\kappa_{i_1}}(i_1) \right) \dots \left( i_k \sigma_2(i_k) \dots, \sigma_2^{\kappa_{i_k}}(i_k) \right)$$

the decomposition of  $\sigma_2$  into a product of cycles.

Recall that in [8] we constructed a free commutative cochain  $\mathbb{Q}$ -algebra

$$\left( \Lambda(x_1, x_2, y_1, y_2, y_3, \{z_j, w_j\}_{g_i \in G}), \partial \right)$$

where the degrees of the elements in this graded algebra are

$$|x_1| = 8 \quad , \quad |x_2| = 10 \quad , \quad |w_j| = 40$$

and where the differential is given by:

$$\begin{aligned} \partial(x_1) &= \partial(x_2) = \partial(w_j) = 0, \quad \partial(y_1) = x_1^3 x_2, \quad \partial(y_2) = x_1^2 x_2^2, \quad \partial(y_3) = x_1 x_2^3 \\ \partial(z_j) &= w_j^3 + w_j w_{\sigma_{j+1}(1)} x_2^4 + \sum_{\tau=1}^k w_j w_{\sigma_{j+1}(i_\tau)} x_2^4 + u + x_1^{15} \quad , \quad 1 \leq j \leq n-1 \\ \partial(z_n) &= w_n^3 + w_n w_1 x_2^4 + \sum_{\tau=1}^k w_n w_{i_\tau} x_2^4 + u + x_1^{15} \end{aligned} \quad (1)$$

where  $u = y_1 y_2 x_1^4 x_2^2 - y_1 y_3 x_1^5 x_2 + y_2 y_3 x_1^6$ , and we proved that

$$\mathcal{E}(\Lambda(x_1, x_2, y_1, y_2, y_3, \{z_j, w_j\}_{g_i \in G})) \cong G$$

where  $\mathcal{E}(\Lambda(x_1, x_2, y_1, y_2, y_3, \{z_j, w_j\}_{1 \leq j \leq n}))$  denotes the group of self homotopy cochain equivalences of  $\Lambda(x_1, x_2, y_1, y_2, y_3, \{z_j, w_j\}_{g_i \in G})$  (see [6, 8] for more details).

Now let  $V^{119} = \mathbb{Q}\{z_1, \dots, z_n\}$  be the vector space spanned by the set  $\{z_1, \dots, z_n\}$ . Recall that  $|z_i| = 119$  for every  $1 \leq i \leq n$ . In ([8], Proposition 3.9), it is shown that

$$\mathcal{E}(\Lambda(x_1, x_2, y_1, y_2, y_3, \{z_j, w_j\}_{g_i \in G})) \cong \mathcal{D}_{40}^{119},$$

where  $\mathcal{D}_{40}^{119}$  is the subgroup of  $\text{aut}(V^{119}) \times \mathcal{E}(\Lambda(x_1, x_2, y_1, y_2, y_3, \{w_j\}_{g_i \in G}))$  consisting of the couples  $(\xi, [\alpha])$  making the following diagram commutes:

$$\begin{array}{ccc} V^{119} & \xrightarrow{\xi} & V^{119} \\ \downarrow b & & \downarrow b \\ \Gamma_G^{120} & \xrightarrow{H^{120}(\alpha)} & \Gamma_G^{120} \end{array} \quad (2)$$

where  $\Gamma_G^{120} = H^{120}(\Lambda(x_1, x_2, y_1, y_2, y_3, \{w_j\}_{g_i \in G}))$  and where  $b$  is defined by

$$b(z_i) = \widehat{\partial(z_i)} \quad , \quad 1 \leq j \leq n \quad (3)$$

Here  $\widehat{\partial(z_i)}$  is the cohomology class of  $\partial(z_i)$  in  $H^{120}(\Lambda(x_1, x_2, y_1, y_2, y_3, \{w_j\}_{g_i \in G}))$ . Moreover, it is shown that if  $(\xi, [\alpha]) \in \mathcal{D}_{40}^{119}$ , then there exists a unique permutation

$$\sigma_s = \begin{pmatrix} 1 & 2 & \dots & n \\ s & \sigma_s(2) & \dots & \sigma_s(n) \end{pmatrix}$$

such that

$$\xi(z_j) = z_{\sigma_s(j)} \quad , \quad \alpha(w_j) = w_{\sigma_s(j)} \quad , \quad \alpha = id \quad , \quad \text{on } x_1, x_2, y_1, y_2, y_3. \quad (4)$$

Thus, there is an isomorphism  $\Psi : \mathcal{D}_{40}^{119} \rightarrow G$  defined by  $\Psi((\xi, [\alpha])) = g_s$ , where the element  $g_s$  corresponds to the permutation  $\sigma_s$ , given in (2.1), via Cayley's theorem. Set  $u = y_1 y_2 x_1^4 x_2^2 - y_1 y_3 x_1^5 x_2 + y_2 y_3 x_1^6$ . As the following set of generators

$$\Sigma = \left\{ w_1^3 ; \dots ; w_n^3 ; w_j w_{\sigma_{j+1}(1)} x_2^4 ; w_j w_{\sigma_{j+1}(i_\tau)} x_2^4 ; u ; x_1^{15} \right\} \quad (5)$$

where  $1 \leq j \leq n$  and  $1 \leq \tau \leq k$ , is linearly independent in the vector space

$$\Gamma_G^{120} = H^{120}(\Lambda(x_1, x_2, y_1, y_2, y_3, \{w_j\}_{i \in G}))$$

it follows that  $\Sigma$  can be chosen, according the formulas (1) and (3), as a basis for the vector space  $b(V^{119}) \subseteq \Gamma_G^{120}$ . Notice that

$$\dim b(V^{119}) = \text{cardinal}(\Sigma) = (1+k)n + 2 \quad (6)$$

Thus, if  $B_G$  denotes the the matrix of order  $((1+k)n + 2) \times n$  which is associated to the linear map  $b$  defined in (3) with respects to the basis is  $\Sigma$ , then we can write

$$B_G = \begin{bmatrix} I_n \\ M \\ D \end{bmatrix} \quad \text{where} \quad D = \begin{bmatrix} 1 & 1 & \dots & 1 \\ 1 & 1 & \dots & 1 \end{bmatrix},$$

where the matrix  $M = [m_{ij}]$  is defined by

$$m_{ij} = \begin{cases} 1, & \text{if } i \in \{\sigma_{j+1}(1), \sigma_{j+1}(i_1), \dots, \sigma_{j+1}(i_k)\}, \\ 0, & \text{otherwise.} \end{cases}$$

Consequently, taking into construction (4), the matrices associated to the linear maps  $\xi$  and the restriction of the linear map  $H^{120}(\alpha)$  to  $b(V^{119})$ , given in the diagram (2)

and corresponding to the element  $(\xi, [\alpha]) \in \mathcal{D}_{40}^{119}$ , can be written, respectively, as

$$C_{g_s} = \sigma_s I_n \quad , \quad A_{g_s} = \begin{bmatrix} \sigma_s I_n & 0 & 0 \\ 0 & \tilde{A}_{g_s} & 0 \\ 0 & 0 & I_2 \end{bmatrix} , \quad (7)$$

where

$$\sigma_s I_n = [c_{i,j}]_{1 \leq i,j \leq (k+1)n} \quad , \quad c_{i,j} = \begin{cases} 1, & \text{if } i = \sigma_s(j) \\ 0, & \text{otherwise} \end{cases} ,$$

and where

$$\tilde{A}_{g_s} = [a_{n+i,n+j}]_{1 \leq i,j \leq (k+1)n} \quad , \quad a_{n+i,n+j} = \begin{cases} 1, & \text{if } i = \sigma_s(j) \\ 0, & \text{otherwise} \end{cases} .$$

Here  $\sigma_s$  is the permutation corresponds to  $g_s$  via Cayley's theorem.

*Remark 2.1.* From (7), it is clear to see that  $A_{g_s}$  is a permutation matrix. Recall that the commutativity of the diagram (2) implies that

$$A_{g_s} B_G = B_G C_{g_s} \quad , \quad \forall g_s \in G \quad (8)$$

**Definition 2.2.** Let  $G = \{g_1, \dots, g_n\}$  be a group, we define the following two sets

$$\mathcal{A}_G = \{A_{g_s} \ , \ g_s \in G\} \quad , \quad \Omega_G = \{(A_{g_s}, C_{g_s}) \in \mathcal{A}_G \times \mathcal{C}_G \ , \ g_s \in G\}$$

**Theorem 2.3.** *The sets  $\mathcal{A}_G$  and  $\Omega_G$  are groups isomorphic to  $G$ .*

*Proof.* First let us prove that  $\mathcal{A}_G$  is a group. Indeed, let  $A_{g_s}, A_{g_r} \in \mathcal{A}_G$ . By (8) there exist two matrices  $C_{g_s}, C_{g_r}$  such that

$$A_{g_s} B_G = B_G C_{g_s} \quad \text{and} \quad A_{g_r} B_G = B_G C_{g_r}$$

therefore  $A_{g_s} A_{g_r} B_G = A_{g_s} B_G C_{g_r} = B_G C_{g_s} C_{g_r}$ , it follows that  $A_{g_s} A_{g_r} \in \mathcal{A}_G$ . Here we use the that fact that

$$A_{g_s} A_{g_r} = A_{g_s g_r} \quad \text{and} \quad C_{g_s} C_{g_r} = C_{g_s g_r} \quad (9)$$

Next let  $A_{g_s} \in \mathcal{A}_G$ . As  $A_{g_s} B_G = B_G C_{g_s}$  and  $A_{g_s}, C_{g_s}$  are invertible, we deduce that  $B_G C_{g_s}^{-1} = (A_{g_s})^{-1} B_G$  implying that  $(A_{g_s})^{-1} \in \mathcal{A}_G$ . Notice that  $A_{g_s}^{-1} = A_{g_s^{-1}}$ .

Next, using the same arguments, it is easy to check that the set  $\Omega_G$  is a group. Finally, it is clear that the two maps  $\chi : G \rightarrow \mathcal{A}_G$  and  $\varphi : G \rightarrow \Omega_G$ , defined by  $\chi(g_s) = A_{g_s}$  and  $\varphi(g_s) = (A_{g_s}, C_{g_s})$  respectively, are isomorphisms of groups.  $\square$

## 2.2. Realisable subgroups.

**Definition 2.4.** A subgroup  $H$  of  $\mathcal{P}(n)$  is called *realisable* if each element  $M \in H$  is obtained by permuting the rows of the identity matrix  $I_n$  using a permutation  $\tau \in S_n$  satisfying  $\tau(i) \neq i$  for all  $1 \leq i \leq n$ .

Let  $G = \{g_1, \dots, g_n\}$  be a group. Based on the formula (7), let us define the following matrix

$$M_{g_s} = \begin{bmatrix} \sigma_s I_n & 0 \\ 0 & \tilde{A}_{g_s} \end{bmatrix} \quad , \quad g_s \in G \quad (10)$$

**Theorem 2.5.** *If  $H_G = \{M_{g_s} \ , \ g_s \in G\}$ , then  $H_G$  is a realisable subgroup of  $\mathcal{P}((1+k)n)$  isomorphic to  $G$*



and  $H_{\mathbb{Z}_4} = \{I_8, M_{(1234)}, M_{(13)(24)}, M_{(1432)}\}$ , where

$$M_{(1234)} = \begin{pmatrix} 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \end{pmatrix}, \quad M_{(13)(24)} = \begin{pmatrix} 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \end{pmatrix}, \quad M_{(1432)} = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \end{pmatrix}$$

Recall that  $A_{(1234)}$  is the matrix associate to the restriction of the linear map  $H^{120}(\alpha)$  to the vector space  $b(V^{119})$ , where the cochain map  $\alpha$  is given (4), with respects to the basis  $\Sigma$  in (11). Thus,  $A_{(1234)}$  is obtained by using the permutation  $\sigma_2 = (1234)$  as follows

$$w_1^3 \mapsto w_2^3, \quad w_2^3 \mapsto w_3^3, \quad w_3^3 \mapsto w_4^3, \quad w_4^3 \mapsto w_1^3, \quad w_1 w_2 x_2^4 \mapsto w_2 w_3 x_2^4$$

$$w_2 w_3 x_2^4 \mapsto w_3 w_4 x_2^4, \quad w_3 w_4 x_2^4 \mapsto w_4 w_1 x_2^4, \quad w_4 w_1 x_2^4 \mapsto w_1 w_4 x_2^4$$

$$y_1 y_2 x_1^4 x_2^2 - y_1 y_3 x_1^5 x_2 + y_2 y_3 x_1^6 \mapsto y_1 y_2 x_1^4 x_2^2 - y_1 y_3 x_1^5 x_2 + y_2 y_3 x_1^6, \quad x_1^{15} \mapsto x_1^{15}$$

and likewise we obtain the matrices  $A_{(13)(24)}$  and  $A_{(1432)}$ . Notice that matrices  $M_{(1234)}, M_{(13)(24)}, M_{(1432)}$  are constructed from the matrices  $A_{(1234)}, A_{(13)(24)}, A_{(1432)}$  using (9) and finally we have

$$\Omega_{\mathbb{Z}_4} = \left\{ (I_4, I_{12}), (A_{(1234)}, C_{(1234)}), (A_{(13)(24)}, C_{(13)(24)}), (A_{(1432)}, C_{(1432)}) \right\}$$

It is also worth noting to point out that the group  $M_{\mathbb{Z}_4}$ , which is isomorphic to  $\mathbb{Z}_4$  is a realisable subgroup of the group of permutation matrices  $\mathcal{P}(8)$ .

**Example 2.7.** In this example we use the same analysis and computation as in the example (2.6), but we omit all the details, to determine the two groups  $\mathcal{A}_{\mathbb{V}}, \mathcal{C}_{\mathbb{V}}, H(\mathbb{V})$  and  $\Omega_{\mathbb{V}}$  for the Klein group  $\mathbb{V}$ . Indeed, the monomorphism  $\mathbb{V} = \{g_1, g_2, g_3, g_4\} \hookrightarrow S_4$  is given by  $g_2 \leftrightarrow (12)(34)$ ,  $g_3 \leftrightarrow (13)(24)$ ,  $g_4 \leftrightarrow (14)(23)$ , so the model associated to  $\mathbb{V}$  is  $(\Lambda(x_1, x_2, y_1, y_2, y_3, w_1, w_2, w_3, w_4, z_1, z_2, z_3, z_4), \partial)$  where  $|x_1| = 8, |x_2| = 10, |w_j| = 40$ , and where the differential is given by

$$\partial(x_1) = \partial(x_2) = \partial(w_j) = 0, \quad \partial(y_1) = x_1^3 x_2, \quad \partial(y_2) = x_1^2 x_2^2, \quad \partial(y_3) = x_1 x_2^3$$

$$\partial(z_1) = w_1^3 + w_1 w_2 x_2^4 + w_1 w_4 x_2^4 + y_1 y_2 x_1^4 x_2^2 - y_1 y_3 x_1^5 x_2 + y_2 y_3 x_1^6 + x_1^{15}$$

$$\partial(z_2) = w_2^3 + w_2 w_3 x_2^4 + w_2 w_1 x_1^4 + y_1 y_2 x_1^4 x_2^2 - y_1 y_3 x_1^5 x_2 + y_2 y_3 x_1^6 + x_1^{15}$$

$$\partial(z_3) = w_3^3 + w_3 w_4 x_2^4 + w_3 w_2 x_2^4 + y_1 y_2 x_1^4 x_2^2 - y_1 y_3 x_1^5 x_2 + y_2 y_3 x_1^6 + x_1^{15}$$

$$\partial(z_4) = w_4^3 + w_4 w_1 x_2^4 + w_4 w_3 x_2^4 + y_1 y_2 x_1^4 x_2^2 - y_1 y_3 x_1^5 x_2 + y_2 y_3 x_1^6 + x_1^{15}$$

If  $u = y_1 y_2 x_1^4 x_2^2 - y_1 y_3 x_1^5 x_2 + y_2 y_3 x_1^6$ , then the basis  $\Sigma$  is given by

$$\Sigma = \left\{ w_1^3, w_2^3, w_3^3, w_4^3, w_1 w_2 x_2^4, w_2 w_3 x_2^4, w_3 w_4 x_2^4, w_4 w_1 x_2^4, w_1 w_4 x_2^4, w_2 w_1 x_2^4, w_3 w_2 x_2^4, w_4 w_3 x_2^4, u, x_1^{15} \right\} \quad (12)$$

implying that  $\dim b(V^{119}) = 14$  and the matrix  $B_{\mathbb{V}}$  is

$$B_{\mathbb{V}} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 0 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{pmatrix}$$



5. M. Benkhalifa, *The effect of cell-attachment on the group of self-equivalences of an elliptic space*, Michigan Mathematical Journal. DOI: 10.1307/mmj/20195840, (2021)
6. M. Benkhalifa, *On the group of self-homotopy equivalences of an elliptic space*, Proceedings of the American Mathematical Society. Vol.148 (6),2695-2706, 2020
7. M. Benkhalifa, , *Adams-Hilton model and the group of self-homotopy equivalences of a simply connected cw-complex*, Homology, Homotopy and Applications, vol. 21(2), 345–362, 2019
8. M. Benkhalifa, *Postnikov decomposition and the the group of self-equivalences of a rationalized space* . Homology, Homotopy and Applications, vol.19(1), pp.209-224, 2017
9. M. Benkhalifa and S. B. Smith, *The effect of cell attachment on the group of self-equivalences of an  $R$ -local space*. Journal of Homotopy and Related Structures. Vol. 4(1), (2015), 135-144.
10. M. Benkhalifa, *Whitehead exact sequence and differential graded free Lie algebra*, Internat. J. Math. 15 (2004), 987-1005
11. Y. Félix, S. Halperin, and J.-C. Thomas, *Rational homotopy theory*, Graduate Texts in Mathematics, Vol. 205, Springer-Verlag, New York, 2001.
12. D.W. Kahn, *Realization problem for the group of homotopy classes of self- homotopy equivalences*, Math. Annalen 220 (1976), 37-46.

DEPARTMENT OF MATHEMATICS. FACULTY OF SCIENCES, UNIVERSITY OF SHARJAH. SHARJAH, UNITED ARAB EMIRATES

*Email address:* mbenkhalifa@sharjah.ac.ae