

# A Control Theorem for Primitive ideals in Iwasawa algebras

Adam Jones

September 18, 2019

## Abstract

Let  $p$  be a prime,  $K \setminus \mathbb{Q}_p$  a finite extension,  $G$  a nilpotent, uniform pro- $p$  group. We prove that all faithful, primitive ideals in the Iwasawa algebra  $KG$  are controlled by  $C_G(Z_2(G))$ , the centraliser of the second term in the upper central series for  $G$ .

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
1.1	Background . . . . .	1
1.2	Main Results . . . . .	3
<b>2</b>	<b>Preliminaries</b>	<b>4</b>
2.1	The $C(G, \mathcal{O})$ -action and Mahler expansions . . . . .	4
2.2	Non-commutative valuations . . . . .	6
<b>3</b>	<b>Using Mahler expansions</b>	<b>8</b>
3.1	Mahler automorphisms . . . . .	8
3.2	Approximations . . . . .	10
3.3	Drawbacks in characteristic 0 . . . . .	12
3.4	Using Compactness . . . . .	12
3.5	The Controlling subgroup . . . . .	15
<b>4</b>	<b>Control Theorem</b>	<b>17</b>
4.1	Technical results in characteristic 0 . . . . .	17
4.2	$J$ -Ideals . . . . .	18
4.3	Controlling prime $J$ -ideals . . . . .	20

## 1 Introduction

### 1.1 Background

Fix  $p$  a prime,  $K \setminus \mathbb{Q}_p$  a finite extension,  $\mathcal{O}$  the ring of integers of  $K$ ,  $\pi$  a uniformiser for  $\mathcal{O}$ , and let  $G$  be a uniform pro- $p$  group in the sense of [7, Definition 4.1].

Define the *Iwasawa algebra* of  $G$  with coefficients in  $\mathcal{O}$  to be the  $\mathcal{O}$ -algebra defined by:

$$\mathcal{O}G := \varinjlim_{N \trianglelefteq_o G} \mathcal{O}[G/N]. \quad (1)$$

An ongoing project is to investigate the ideal structure of  $\mathcal{O}G$ , and provide an answer to the questions posed in [3, Section 7]. Much progress towards a complete classification of prime ideals was made in [1] and [9], but these results only apply in a characteristic  $p$  setting, i.e. they apply only to prime ideals of  $\mathcal{O}G$  which contain  $p$ . This paper will focus instead on the characteristic 0 case.

Define  $KG := \mathcal{O}G \otimes_{\mathcal{O}} K$ , then using [13, Lemma 21.1], this can be realised as the algebra of continuous,  $K$ -valued distributions on  $G$ , i.e. as a dense subalgebra of the distribution algebra  $D(G, K)$  of  $K$ -analytic distributions on  $G$  (see [14] for details). For this reason, we call  $KG$  the *Iwasawa algebra of continuous  $K$ -distributions* of  $G$ .

Note that  $\mathcal{O}G$  is an  $\mathcal{O}$ -lattice subalgebra of  $KG$  in the sense of [4, Definition 2.7], and that prime ideals of  $KG$  are in bijection with prime ideals of  $\mathcal{O}G$  that do not contain  $\pi$ . So our aim is to classify prime ideals in  $KG$ .

In the case where  $G = \mathbb{G}(\mathcal{O})$  for some simply connected, split-semisimple, affine algebraic group  $\mathbb{G}$ , it is strongly believed that the only non-zero prime ideals of  $KG$  have finite codimension (see [5] for more details). However, in this paper, we will be interested in the case when  $G$  is solvable, where there is a much richer prime ideal structure.

The general conjecture is that prime ideals in  $KG$  are all of a very predictable form, which is the essence of [3, Question G]. This is made precise by the following definition:

**Definition 1.1.** *Let  $P$  be a prime ideal of  $KG$ . We say that  $P$  is standard if there exists a closed, normal subgroup  $H$  of  $G$  such that:*

*i.  $H - 1 \subseteq P$ .*

*ii. Setting  $G' := \frac{G}{H}$ , the ideal  $\tilde{P} := P/(H - 1)KG$  of  $KG/(H - 1)KG \cong KG'$  is centrally generated.*

Standard prime ideals have very nice properties, in particular they are generally *completely prime*, i.e. if  $P$  is standard then  $KG/P$  is a domain.

**Conjecture 1.1.** *Let  $G$  be a solvable group and let  $P$  be a prime ideal of  $KG$ . Then  $P$  is standard.*

The following definition ([9, Definition 1.1]) will be essential throughout:

**Definition 1.2.** *Let  $I$  be a right ideal of  $KG$ :*

*1. We say that  $I$  is faithful if for all  $g \in G$ ,  $g - 1 \in I$  if and only if  $g = 1$ , i.e.  $G \rightarrow KG/I, g \mapsto g + I$  is injective.*

*2. We say that  $H \leq_c G$  controls  $I$  if  $I = (I \cap KH)KG$ .*

*Define the controller subgroup of  $I$  by  $I^\times := \bigcap \{U \leq_o G : U \text{ controls } I\}$ , and denote by  $\text{Spec}^f(KG)$  the set of all faithful prime ideals of  $KG$ .*

It follows from [2, Theorem A] that a closed subgroup  $H$  of  $G$  controls  $I$  if and only if  $I^\times \subseteq H$ , and we see immediately from Definition 1.1 that a faithful prime ideal  $P$  of  $KG$  is standard if and only if it is controlled by the centre.

Therefore, to prove Conjecture 1.1, it should remain only to prove that for any faithful prime ideal  $P$  of  $KG$ ,  $P$  is controlled by  $Z(G)$ .

## 1.2 Main Results

It was proved in [1] that Conjecture 1.1 holds in the analogous characteristic  $p$  setting if we assume further that  $G$  is *nilpotent*:

**Definition 1.3.** *Let  $H$  be a group,  $n \in \mathbb{N}$ . Define the  $n$ 'th centre  $Z_n(H)$  of  $H$  inductively by  $Z_0(H) := 1$ , and for  $n > 0$ ,  $Z_n(H) := \{h \in H : (h, H) \subseteq Z_{n-1}(H)\}$ .*

*Then each  $Z_n(H)$  is a normal subgroup of  $H$ ,  $Z_1(H) = Z(H)$ , and we define the upper central series of  $H$  to be the ascending chain of subgroups:*

$$1 = Z_0(H) \subseteq Z_1(H) \subseteq Z_2(H) \subseteq \dots$$

*We say that  $H$  is nilpotent if the upper central series terminates at  $H$ , i.e.  $Z_n(H) = H$  for some  $n \in \mathbb{N}$ . The nilpotency class of  $H$  is the smallest integer  $n$  such that  $Z_n(H) = H$ .*

It is not difficult to see that any nilpotent group is solvable, and note that for each  $i \geq 0$ ,  $Z_i(G)$  is a closed, isolated normal subgroup of  $G$ .

Recall that an ideal  $I$  of  $KG$  is *primitive* if  $I = \text{Ann}_{KG}(M)$  for some irreducible  $KG$ -module  $M$ . Here is our main result:

**Theorem 1.1.** *Let  $G$  be a nilpotent, uniform pro- $p$  group, and let  $P$  be a faithful, primitive ideal of  $KG$ . Then  $P$  is controlled by  $C_G(Z_2(G))$ , the centraliser of  $Z_2(G)$  in  $G$ .*

**Examples:** 1. If  $G$  is nilpotent and contains a closed, isolated, abelian normal subgroup  $H$  of codimension 1, it follows from Theorem 1.1 that all faithful primitive ideals of  $KG$  are controlled by  $H$ .

2. Let  $G$  be the group of unipotent, upper-triangular  $4 \times 4$  matrices over  $\mathbb{Z}_p$ , and let  $\mathfrak{g}$  be the  $\mathbb{Z}_p$ -Lie algebra of  $G$ .

Then  $\mathfrak{g}$  is the Lie algebra of strictly upper triangular matrices over  $p\mathbb{Z}_p$ . This has basis  $\{x_1, x_2, \dots, x_6\}$  such that  $x_3$  is central,  $[x_1, x_4] = px_2$ ,  $[x_1, x_5] = px_3$ ,  $[x_2, x_6] = px_3$ ,  $[x_4, x_6] = px_5$ , all other brackets are zero.

Then  $Z_2(\mathfrak{g}) = \text{Span}_{\mathbb{Z}_p}\{x_2, x_3, x_5\}$ , and  $C_{\mathfrak{g}}(Z_2(\mathfrak{g})) = \text{Span}_{\mathbb{Z}_p}\{x_2, x_3, x_4, x_5\}$ . This is abelian, so it follows from Theorem 1.1 that any primitive ideal of  $KG$  is controlled by an abelian subgroup of  $G$ .

3. More generally, suppose that  $G$  is the group of unipotent, upper triangular  $n \times n$  matrices, then the  $\mathbb{Z}_p$ -Lie algebra  $\mathfrak{g}$  of  $G$  is the Lie algebra of strictly upper triangular  $n \times n$  matrices over  $p\mathbb{Z}_p$ .

In this case,  $C_{\mathfrak{g}}(Z_2(\mathfrak{g}))$  is precisely the set of all such matrices where the first two columns and the bottom two rows are zero, and it follows that  $C_G(Z_2(G))$  is the group of unipotent, upper triangular matrices with zero in the (1,2)-position and in the (n-1,n)-position. For

$n > 4$ , this is non-abelian.

Now, Theorem 1.1 is not useful in the case when  $Z(G) = 1$ , because in this case  $C_G(Z_2(G)) = G$  so the statement is obvious. Therefore, we draw no new conclusion if  $G$  is semisimple, and in many cases when  $G$  is solvable.

However, if  $G$  is nilpotent, the result is highly useful, because for  $G$  non-abelian,  $C_G(Z_2(G))$  is always a proper subgroup of  $G$ . In particular, if  $G$  has nilpotency class 2, then  $Z_2(G) = G$  so  $C_G(Z_2(G)) = Z(G)$ , and thus we achieve the following result:

**Corollary 1.2.** *Let  $G$  be a uniform, nilpotent, pro- $p$  group of nilpotency class 2. Then all faithful, primitive ideals of  $KG$  are standard.*

To prove Theorem 1.1, we will adapt the approaches used in [1], [2] and [9] to study prime ideals in Iwasawa algebras in characteristic  $p$ :

Recall from [10, II 2.1.2] the definition of a  $p$ -valuation  $\omega$  on a group  $G$ . We say that  $G$  is  $p$ -valuable if it carries a complete  $p$ -valuation and has finite rank, note that any closed subgroup of a uniform group is  $p$ -valuable.

The following result is analogous to [1, Theorem B], and is the key step in the proof.

**Theorem 1.3.** *Let  $G$  be a  $p$ -valuable group, and let  $A$  be a closed, central subgroup of  $G$ . Let  $P$  be a faithful, prime ideal of  $\mathcal{O}G$  with  $\pi \notin P$ , and suppose there exists  $\varphi \in \text{Aut}^\omega(G)$  such that  $\varphi \neq 1$ ,  $\varphi(P) = P$  and  $\varphi(g)g^{-1} \in A$  for all  $g \in G$ .*

*Then if we assume that  $\mathcal{O}A/\mathcal{O}A \cap P$  is finitely generated over  $\mathcal{O}$ , we conclude that  $P$  is controlled by a proper, open subgroup of  $G$ .*

We will prove Theorem 1.3 in section 3, using the theory of Mahler expansions, and in section 4 we will show that this is enough to imply the appropriate control theorem for all prime ideals satisfying the appropriate finiteness condition, and ultimately prove Theorem 1.1.

## 2 Preliminaries

### 2.1 The $C(G, \mathcal{O})$ -action and Mahler expansions

Fix  $G$  a compact  $p$ -adic Lie group, and define:

$$C(G, \mathcal{O}) := \{f : G \rightarrow \mathcal{O} : f \text{ continuous}\}.$$

Then  $C(G, \mathcal{O})$  is an  $\mathcal{O}$ -algebra with pointwise addition and multiplication. Also, for each  $n \in \mathbb{N}$ , define:

$$C_n = C(G, \frac{\mathcal{O}}{\pi^n \mathcal{O}}) := \{f : G \rightarrow \frac{\mathcal{O}}{\pi^n \mathcal{O}} : f \text{ locally constant}\}.$$

Then each  $C_n$  is an  $\mathcal{O}$ -algebra, and there exists a surjective map:

$$c_{n+1,n} : C_{n+1} \rightarrow C_n, f \mapsto (h : G \rightarrow \frac{\mathcal{O}}{\pi^n \mathcal{O}}, g \mapsto f(g) + \pi^n \mathcal{O}).$$

Furthermore, there exists a surjective map  $c_n : C(G, \mathcal{O}) \rightarrow C_n, f \mapsto (h : G \rightarrow \frac{\mathcal{O}}{\pi^n \mathcal{O}}, g \mapsto f(g) + \pi^n \mathcal{O})$ , and clearly  $c_{n+1,n} \circ c_{n+1} = c_n$  for all  $n$ .

**Lemma 2.1.**  $C(G, \mathcal{O}) = \varprojlim_{n \in \mathbb{N}} C_n$  in the category of  $\mathcal{O}$ -algebras.

*Proof.* Firstly, note that  $C(G, \mathcal{O})$  is  $\pi$ -adically complete, and thus

$$C(G, \mathcal{O}) = \varprojlim_{n \in \mathbb{N}} C(G, \mathcal{O}) / \pi^n C(G, \mathcal{O}) \quad (2)$$

Consider the morphism of  $\mathcal{O}$ -algebras:

$$\Theta : C(G, \mathcal{O}) \rightarrow C_n, f \mapsto (f' : G \rightarrow \frac{\mathcal{O}}{\pi^n \mathcal{O}}, g \mapsto f(g) + \pi^n \mathcal{O})$$

It is clear that this map is surjective, and if  $\Theta(f) = 0$  then  $f(g) \in \pi^n \mathcal{O}$  for all  $g \in G$ , so  $f \in \pi^n C(G, \mathcal{O})$ . Thus  $\ker(\Theta) = \pi^n C(G, \mathcal{O})$ , so  $C(G, \mathcal{O}) / \pi^n C(G, \mathcal{O}) \cong C_n$ , and the result follows from (2).  $\square$

For convenience, set  $A_n := \frac{\mathcal{O}}{\pi^n \mathcal{O}}$  for each  $n \in \mathbb{N}$ , and note that  $\mathcal{O}G / \pi^n \mathcal{O}G \cong A_n G$ .

Recall from [2, Proposition 2.5] that there exists an action  $\rho_n : C_n \rightarrow \text{End}_{\mathcal{O}}(A_n G)$  for each  $n$  such that if  $U \leq_o G$  and  $f \in C_n$  is constant on the cosets of  $U$ , and  $r \in A_n G$  with  $r = \sum_{g \in G//U} s_g g$  for some  $s_g \in A_n U$ , then  $\rho_n(f)(r) = \sum_{g \in G//U} f(g) s_g g$ .

Consider the canonical homomorphisms:

$$\nu_n : \text{End}_{\mathcal{O}}(A_n G) \rightarrow \text{Hom}(\mathcal{O}G, A_n G), f \mapsto (g : \mathcal{O}G \rightarrow \mathcal{O}G / \pi^n \mathcal{O}G, r \mapsto f(r + \pi^n \mathcal{O}G)) \quad (3)$$

$$\mu_{n+1, n} : \text{Hom}(\mathcal{O}G, A_{n+1} G) \rightarrow \text{Hom}(\mathcal{O}G, A_n G), f \mapsto (g : \mathcal{O}G \rightarrow \mathcal{O}G / \pi^n \mathcal{O}G, r \mapsto f(r) + \pi^n \mathcal{O}G) \quad (4)$$

These give rise to the following commutative diagram for each  $n \in \mathbb{N}$ :

$$\begin{array}{ccccc} C_n & \xrightarrow{\rho_n} & \text{End}_{\mathcal{O}}(A_n G) & \xrightarrow{\nu_n} & \text{Hom}_{\mathcal{O}}(\mathcal{O}G, A_n G) \\ \uparrow c_{n+1, n} & & & & \uparrow \mu_{n+1, n} \\ C_{n+1} & \xrightarrow{\rho_{n+1}} & \text{End}_{\mathcal{O}}(A_{n+1} G) & \xrightarrow{\nu_{n+1}} & \text{Hom}_{\mathcal{O}}(A_{n+1} G) \end{array}$$

Now, using a similar argument to Lemma 2.1, we get  $\text{Hom}_{\mathcal{O}}(\mathcal{O}G, \mathcal{O}G) = \varprojlim_{n \in \mathbb{N}} \text{Hom}_{\mathcal{O}}(\mathcal{O}G, A_n G)$ , so it follows that there is a unique map from  $C(G, \mathcal{O}) = \varprojlim_{n \in \mathbb{N}} C_n$  to  $\text{Hom}_{\mathcal{O}}(\mathcal{O}G, \mathcal{O}G) = \text{End}_{\mathcal{O}} \mathcal{O}G$  making the corresponding diagrams commute.

**Definition 2.1.** Define  $\rho : C(G, \mathcal{O}) \rightarrow \text{End}_{\mathcal{O}} \mathcal{O}G$  to be the unique morphism defined above. We call this the canonical action of  $C(G, \mathcal{O})$  on  $\mathcal{O}G$ .

Note that for each  $n \in \mathbb{N}$ ,  $f \in C_n$ ,  $g \in G$ ,  $\rho_n(f)(g) = f(g)g$ , and it follows that for each  $f \in C(G, \mathcal{O})$ , we still have that  $\rho(f)(g) = f(g)g$ .

Also, note that if  $f \in C(G, \mathcal{O})$  is locally constant, then  $\rho(f)$  is the same as the endomorphism defined in [2, Proposition 2.5]. Therefore we have the following result [2, Proposition 2.8], which will be useful to us later:

**Proposition 2.2.** For each  $U \leq_o G$ , let  $C_U := \{f \in C(G, \mathcal{O}) : f \text{ is constant on the cosets of } U\}$ . Then  $C_U$  is an  $\mathcal{O}$ -subalgebra of  $C(G, \mathcal{O})$ , and for any right ideal  $I$  of  $\mathcal{O}G$ ,  $I$  is controlled by  $U$  if and only if  $\rho(C_U)(I) \subseteq I$ .

Now we will recap the notion of the Mahler expansion of an automorphism of  $G$ . This was defined in [1] working over a field of characteristic  $p$ , and we will now use the canonical action to extend the notion to fields of characteristic 0.

First recall the following result, first proved by Mahler in 1958, and given in full in [10, III 1.2.4]:

**Theorem 2.3.** *Let  $M$  be a complete topological  $\mathbb{Z}_p$ -module, and let  $f : \mathbb{Z}_p^d \rightarrow M$  be a continuous function. Then for each  $\alpha \in \mathbb{N}^d$ , define  $m_\alpha(f) := \sum_{\beta \leq \alpha} (-1)^{\alpha-\beta} \binom{\alpha}{\beta} f(\beta)$ , where  $\alpha \leq \beta$  if  $\alpha_i \leq \beta_i$  for all  $i$ , and  $\binom{\alpha}{\beta} = \prod_{i \leq d} \binom{\alpha_i}{\beta_i}$ .*

*Then  $m_\alpha(f) \rightarrow 0$  as  $|\alpha| \rightarrow \infty$ , and for all  $\gamma \in \mathbb{Z}_p^d$ ,  $f(\gamma) = \sum_{\alpha \in \mathbb{N}^d} m_\alpha(f) \binom{\gamma}{\alpha}$ .*

We call this  $m_\alpha(f)$  the  $\alpha$ -Mahler coefficient for  $f$ .

From now on, we will assume that  $G$  is  $p$ -valuable, with  $p$ -valuation  $\omega$ , and we fix an ordered basis  $\underline{g} = \{g_1, \dots, g_d\}$  for  $G$ .

Then given  $\varphi \in \text{Aut}^\omega(G) := \{\phi \in \text{Aut}(G) : \omega(\phi(g)g^{-1}) - \omega(g) > \frac{1}{p-1} \text{ for all } g \in G\}$ , we have a continuous map  $f : \mathbb{Z}_p^d \rightarrow \mathcal{O}G, \beta \mapsto \varphi(\underline{g}^\beta) \underline{g}^{-\beta}$ .

Using Theorem 2.3, define  $m_\alpha(\varphi) := m_\alpha(f)$  to be the  $\alpha$ -Mahler coefficient for  $\varphi$ , and we have that for all  $\beta \in \mathbb{Z}_p^d$ :

$$\varphi(\underline{g}^\beta) = \sum_{\alpha \in \mathbb{N}^d} m_\alpha(\varphi) \binom{\beta}{\alpha} \underline{g}^\alpha$$

Now, for each  $\alpha \in \mathbb{Z}_p^d$ , define  $i_{\underline{g}}^{(\alpha)} : G \rightarrow \mathcal{O}, \underline{g}^\beta \mapsto \binom{\beta}{\alpha}$ , then clearly  $i_{\underline{g}}^{(\alpha)} \in C(G, \mathcal{O})$ , so let  $\partial_{\underline{g}}^{(\alpha)} := \rho(i_{\underline{g}}^{(\alpha)}) \in \text{End}_{\mathcal{O}}(\mathcal{O}G)$ . We call this the  $\alpha$ -quantized divided power.

Then  $\partial_{\underline{g}}^{(\alpha)}(g) = i_{\underline{g}}^{(\alpha)}(g)g$  for all  $g \in G$ , i.e. for all  $\beta \in \mathbb{Z}_p^d$ ,  $\partial_{\underline{g}}^{(\alpha)}(\underline{g}^\beta) = \binom{\beta}{\alpha} \underline{g}^\beta$ .

Hence  $\varphi(g) = \sum_{\alpha \in \mathbb{N}^d} m_\alpha(\varphi) \partial_{\underline{g}}^{(\alpha)}(g)$  for all  $g \in G$ .

Therefore, after extending linearly to  $\mathcal{O}[G]$  and passing to the completion, we should get (after establishing a suitable convergence condition for the Mahler coefficients) that:

$$\varphi = \sum_{\alpha \in \mathbb{N}^d} m_\alpha(\varphi) \partial_{\underline{g}}^{(\alpha)} \tag{5}$$

as an endomorphism of  $\mathcal{O}G$ . This is the *Mahler expansion* of  $\varphi$ .

In section 3, we will see how to use Mahler expansions for suitable automorphisms to prove a control theorem.

## 2.2 Non-commutative valuations

Recall the definition of a *filtration*  $w : R \rightarrow \mathbb{Z} \cup \{\infty\}$  on a ring  $R$  from [9, Definition 2.2], and of the subgroups  $F_n R := \{r \in R : w(r) \geq n\}$ . Recall from [11] that  $w$  is a *Zariskian filtration* if  $F_1 R \subseteq J(F_0 R)$  and the *Rees ring*  $\bigoplus_{n \in \mathbb{N}} F_n R t^n$  is Noetherian.

Note that Zariskian filtrations are always separated.

An element  $x \in R \setminus 0$  is  $w$ -regular if  $w(xy) = w(x) + w(y)$  for all  $y \in R$ , and we say that  $w$  is a valuation if every non-zero element of  $R$  is  $w$ -regular. Finally, if  $S$  is a central subring of  $R$ , we say that  $w$  is  $S$ -linear if every non-zero element of  $S$  is  $w$ -regular.

**Example:** If  $\mathcal{O}$  is a complete, commutative DVR, and  $(G, \omega)$  is a  $p$ -valuable group with ordered basis  $\underline{g} = \{g_1, \dots, g_d\}$ , then  $\mathcal{O}G = \{ \sum_{\alpha \in \mathbb{N}^d} \lambda_\alpha b_1^{\alpha_1} \cdots b_d^{\alpha_d} : \lambda_\alpha \in \mathcal{O} \}$ , where  $b_i = g_i - 1$ .

By [10, III 2.3.3],  $\mathcal{O}G$  carries a complete, Zariskian valuation  $w$  given by

$$w\left(\sum_{\alpha \in \mathbb{N}^d} \lambda_\alpha b_1^{\alpha_1} \cdots b_d^{\alpha_d}\right) = \inf\{v_p(\lambda_\alpha) + \sum_{i \leq n} \alpha_i \omega(g_i) : \alpha \in \mathbb{N}^d\}$$

and we may assume that the associated graded ring is a commutative polynomial ring over  $k := \mathcal{O}/\pi\mathcal{O}$  in  $d + 1$  variables.

**Lemma 2.4.** *Suppose  $w$  is an  $\mathcal{O}$ -linear filtration on an  $\mathcal{O}$ -algebra  $A$  of characteristic 0, and suppose that  $x \in A$  with  $w(x - 1) > w(p)$ . Then  $w(x^{p^m} - 1) = mw(p) + w(x - 1)$  for all  $m \in \mathbb{N}$ .*

*Proof.* Using the binomial theorem, it is clear that

$$x^{p^m} - 1 = (x - 1 + 1)^{p^m} - 1 = \sum_{k \geq 1} \binom{p^m}{k} (x - 1)^k = p^m(x - 1) + \sum_{k \geq 2} \binom{p^m}{k} (x - 1)^k$$

Clearly  $w(p^m(x - 1)) = w(p^m) + w(x - 1) = mw(p) + w(x - 1)$  since  $w$  is  $\mathcal{O}$ -linear. So it remains to show that  $w(\binom{p^m}{k}(x - 1)^k) > mw(p) + w(x - 1)$  for all  $k \geq 2$ .

First, note that  $w(\binom{p^m}{p^m}(x - 1)^{p^m}) = w((x - 1)^{p^m}) \geq p^m w(x - 1) = (p^m - 1)w(x - 1) + w(x - 1) > (p^m - 1)w(p) + w(x - 1) \geq mw(p) + w(x - 1)$ .

So from now on, we may assume that  $k < p^m$ .

Now,  $k = a_0 + a_1 p + \cdots + a_t p^t$  for some integers  $0 \leq a_i < p$ , and since  $k \leq p^m - 1$  we may assume that  $t = m - 1$ . Define  $s(k) := a_1 + a_1 + \cdots + a_t$ .

Recall from [10, III 1.1.2.5] that  $v_p(k!) = \frac{k - s(k)}{p - 1}$ , and hence

$$v_p\left(\binom{p^m}{k}\right) = v_p(p^m!) - v_p(k!) - v_p((p^m - k)!) = \frac{p^m - s(p^m) - k + s(k) - (p^m - k) + s(p^m - k)}{p - 1} = \frac{s(k) + s(p^m - k) - s(p^m)}{p - 1}.$$

Now, let  $m \geq i \geq 1$  be maximal such that  $a_{m-i} \neq 0$ , then  $p^m = (p - 1)p^{m-i} + (p - 1)p^{m-i+1} + \cdots + (p - 1)p^{m-1} + p^{m-i}$ , so

$$\begin{aligned} p^m - k &= (p - 1 - a_{m-i})p^{m-i} + (p - 1 - a_{m-i+1})p^{m-i+1} + \cdots + (p - 1 - a_{m-1})p^{m-1} + p^{m-i} \\ &= (p - a_{m-i})p^{m-i} + (p - 1 - a_{m-i+1})p^{m-i+1} + \cdots + (p - 1 - a_{m-1})p^{m-1}. \end{aligned}$$

Hence  $s(p^m - k) = (p - a_{m-i}) + (p - 1 - a_{m-i+1}) + \cdots + (p - 1 - a_{m-1}) = i(p - 1) + 1 - s(k)$ .

Clearly  $s(p^m) = 1$ , so  $v_p\left(\binom{p^m}{k}\right) = \frac{s(k) + s(p^m - k) - s(p^m)}{p - 1} = \frac{s(k) + i(p - 1) + 1 - s(k) - 1}{p - 1} = \frac{i(p - 1)}{p - 1} = i$ .

Also,  $k = a_{m-i}p^{m-i} + \cdots + a_{m-1}p^{m-1} = p^{m-i}(a_{m-i} + \cdots + a_{m-1}p^{i-1})$ , so  $k \geq p^{m-i} \geq m - i + 1$  if  $i < m$ .

Now,  $w\left(\binom{p^m}{k}(x-1)^k\right) \geq w\left(\binom{p^m}{k}\right) + kw(x-1) \geq v_p\left(\binom{p^m}{k}\right)w(p) + kw(x-1)$

$$= iw(p) + (k-1)w(x-1) + w(x-1) > iw(p) + (k-1)w(p) + w(x-1), \text{ so:}$$

If  $i < m$  then this is at least  $iw(p) + (m-i)w(p) + w(x-1) = mw(p) + w(x-1)$ , and

if  $i = m$  then since  $k > 1$  we have  $iw(p) + (k-1)w(p) + w(x-1) = mw(p) + (k-1)w(p) + w(x-1) \geq mw(p) + w(x-1)$  as required.  $\square$

Recall the following definition ([9, Definition 3.1]):

**Definition 2.2.** *A non-commutative valuation on a simple artinian ring  $Q$  is a Zariskian filtration  $v : Q \rightarrow \mathbb{Z} \cup \{\infty\}$  such that if  $\widehat{Q}$  is the completion of  $R$  with respect to  $v$ , then  $\widehat{Q} \cong M_k(Q(D))$  for some  $k \in \mathbb{N}$  and some complete non-commutative DVR  $D$ , and  $v$  is induced by the  $J(D)$ -adic filtration on  $\widehat{Q}$ .*

It follows from this definition that if  $v$  is a non-commutative valuation on  $Q$ , then every element of  $Z(Q)$  is  $v$ -regular.

**Theorem 2.5.** *Let  $G$  be a  $p$ -valuable group,  $P$  a faithful prime ideal of  $\mathcal{O}G$  with  $\pi \notin P$ , then there exists a non-commutative valuation  $v$  on  $Q(\mathcal{O}G/P)$  such that the natural map  $\tau : (\mathcal{O}G, w) \rightarrow (Q(\mathcal{O}G/P), v)$  is continuous, and there exists  $m_1 \in \mathbb{N}$  such that for all  $g \in G$ ,  $v(\tau(g^{p^{m_1}} - 1)) > v(p)$ .*

*Proof.* Since  $\mathcal{O}G/P$  is a prime ring carrying a Zariskian filtration  $w$ , and the associated graded  $\text{gr } \mathcal{O}G/P \cong \text{gr } \mathcal{O}G/\text{gr } P$  is a commutative, infinite dimensional  $\mathbb{F}_p$ -algebra, it follows from [1, Theorem C] that there exists a non-commutative valuation  $v$  on  $Q(\mathcal{O}G/P)$  such that  $\tau$  is continuous.

Given  $g \in G$ , the sequence  $g^{p^m} - 1$  converges to zero in  $\mathcal{O}G$ , so since  $\tau$  is continuous, the sequence  $\tau(g^{p^m} - 1)$  converges to zero with respect to  $v$ , and hence there exists  $m_g \in \mathbb{N}$  such that  $v(\tau(g^{p^{m_g}} - 1)) > v(p)$ .

So, let  $\underline{g} = \{g_1, \dots, g_d\}$  be an ordered basis for  $G$ , and let  $m_1 := \max\{m_{g_i} : i = 1, \dots, d\}$ , then it follows that  $v(\tau(g^{p^{m_1}} - 1)) > v(p)$  for all  $g \in G$ .  $\square$

### 3 Using Mahler expansions

In section 2.1, we defined the Mahler expansion of an automorphism  $\varphi \in \text{Aut}^\omega(G)$  as a linear combination in the quantized divided powers  $\partial_{\underline{g}}^\alpha$ .

In this section, we will analyse convergence of Mahler expansions, and use this to prove our control Theorem 1.3.

#### 3.1 Mahler automorphisms

We now define a class of automorphisms which have a workable formula for their Mahler coefficients.



**Definition 3.1.** Suppose  $\underline{g} = \{g_1, \dots, g_d\}$  is an ordered basis for  $G$ , and  $\varphi \in \text{Aut}^\omega(G)$ . We say that  $\varphi$  is a Mahler automorphism with respect to  $\underline{g}$  if its Mahler coefficients satisfy:

$$m_\alpha(\varphi) = (\varphi(g_1)g_1^{-1} - 1)^{\alpha_1} \dots (\varphi(g_d)g_d^{-1} - 1)^{\alpha_d} \text{ for all } \alpha \in \mathbb{Z}_p^d.$$

It follows from the proof of [1, Corollary 6.6] and [1, Corollary 6.7] that if  $\varphi$  is a Mahler automorphism with respect to  $\underline{g}$ , then  $\varphi = \sum_{\alpha \in \mathbb{N}^d} m_\alpha(\varphi) \partial_{\underline{g}}^{(\alpha)}$  as an endomorphism of  $\mathcal{O}G$  as we require.

The following proposition simplifies the task of finding Mahler automorphisms. For convenience, we write  $\psi(g) := \varphi(g)g^{-1}$  for  $g \in G$ :

**Proposition 3.1.** *The following are equivalent:*

- i.  $\varphi$  is a Mahler automorphism with respect to  $\underline{g} = \{g_1, \dots, g_d\}$ .
- ii.  $\psi(\underline{g}^\beta) = \prod_{i=1}^d \psi(g_i)^{\beta_i}$  for all  $\beta \in \mathbb{N}^d$ .
- iii.  $\psi(g_i)$  commutes with  $g_j$  for all  $j \leq i$ .

*Proof.* (ii  $\implies$  i) Given by the proof of [1, Lemma 6.7]

(i  $\implies$  ii) Suppose that for all  $\alpha \in \mathbb{N}^d$ :

$$m_\alpha(\varphi) = \sum_{\beta \leq \alpha} (-1)^{\alpha-\beta} \binom{\alpha}{\beta} \psi(g_1^{\beta_1} g_2^{\beta_2} \dots g_d^{\beta_d}) = (\psi(g_1) - 1)^{\alpha_1} (\psi(g_2) - 1)^{\alpha_2} \dots (\psi(g_d) - 1)^{\alpha_d}.$$

Then given  $i = 1, \dots, d$ , suppose that  $\beta_j = 0$  for all  $j \neq i$ . Clearly if  $\beta_i = 0, 1$  then  $\psi(g_i^{\beta_i}) = \psi(g_i)^{\beta_i}$ , so suppose, for induction that  $\psi(g_i^k) = \psi(g_i)^k$  for all  $0 \leq k < \beta_i$ . Then:

$$\begin{aligned} \sum_{k \leq \beta_i} (-1)^{\beta_i-k} \binom{\beta_i}{k} \psi(g_i^k) &= \sum_{k < \beta_i} (-1)^{\beta_i-k} \binom{\beta_i}{k} \psi(g_i)^k + \psi(g_i^{\beta_i}) = (\psi(g_i) - 1)^{\beta_i} \\ &= \sum_{k \leq \beta_i} \binom{\beta_i}{k} (-1)^{\beta_i-k} \psi(g_i)^k \text{ using binomial expansion, hence } \psi(g_i^{\beta_i}) = \psi(g_i)^{\beta_i}. \end{aligned}$$

So, we have shown that  $\psi(g_i^{\beta_i}) = \psi(g_i)^{\beta_i}$  for all  $i = 1, \dots, d$ ,  $\beta_i \in \mathbb{Z}_p$ .

Now suppose that for  $i = 1, \dots, m$ ,  $\psi(g_1^{\beta_1} \dots g_m^{\beta_m}) = \psi(g_1)^{\beta_1} \dots \psi(g_m)^{\beta_m}$  for some  $1 \leq m < d$ . We have proved this for  $m = 1$ , so we will now show that it holds for  $m + 1$  and apply induction.

So now suppose that for all  $k < \beta_{m+1}$ ,  $\psi(g_1^{\beta_1} \dots g_m^{\beta_m} g_{m+1}^k) = \psi(g_1)^{\beta_1} \dots \psi(g_m)^{\beta_m} \psi(g_{m+1})^k$ . It is clear that this holds for  $\beta_{m+1} = 1$ , so we will apply a second induction.

$$\begin{aligned} \text{We know that } & \sum_{\gamma \leq \beta} (-1)^{\beta-\gamma} \binom{\beta}{\gamma} \psi(g_1^{\gamma_1} \dots g_{m+1}^{\gamma_{m+1}}) \\ &= \sum_{k=0}^{\beta_{m+1}} \sum_{\gamma \leq \beta} (-1)^{\beta-\gamma} \binom{\beta}{\gamma} (-1)^{\beta_{m+1}-k} \binom{\beta_{m+1}}{k} \psi(g_1^{\gamma_1} \dots g_m^{\gamma_m} g_{m+1}^k) \\ &= (\psi(g_1) - 1)^{\beta_1} (\psi(g_2) - 1)^{\beta_2} \dots (\psi(g_{m+1}) - 1)^{\beta_{m+1}}. \end{aligned}$$

So expanding out  $(\psi(g_{m+1}) - 1)^{\beta_{m+1}}$  and applying the first induction gives that:

$$\sum_{\gamma \leq \beta} (-1)^{\beta-\gamma} \binom{\beta}{\gamma} \psi(g_1)^{\gamma_1} \dots (g_m)^{\gamma_m} \sum_{k=0}^{\beta_{m+1}} (-1)^{\beta_{m+1}-k} \binom{\beta_{m+1}}{k} \psi(g_{m+1})^k$$

$$= \sum_{k=0}^{\beta_{m+1}} \sum_{\gamma \leq \beta} (-1)^{\beta-\gamma} (-1)^{\beta_{m+1}-k} \binom{\beta}{\gamma} \binom{\beta_{m+1}}{k} \psi(g_1^{\gamma_1} \cdots g_m^{\gamma_m} g_{m+1}^k).$$

So applying the second induction gives  $\psi(g_1^{\beta_1} \cdots g_{m+1}^{\beta_{m+1}}) = \psi(g_1)^{\beta_1} \cdots \psi(g_{m+1})^{\beta_{m+1}}$ , so we are done.

(ii  $\implies$  iii) Given  $i \leq j$ , we have that  $\psi(g_i g_j) = \psi(g_i) \psi(g_j)$ , so by the definition of  $\psi$ ,  $\varphi(g_i g_j) (g_i g_j)^{-1} = \varphi(g_i) g_i^{-1} \varphi(g_j) g_j^{-1}$ .

Therefore,  $\varphi(g_i) \varphi(g_j) g_j^{-1} g_i^{-1} = \varphi(g_i) g_i^{-1} \varphi(g_j) g_j^{-1}$ , so  $\varphi(g_j) g_j^{-1} g_i^{-1} = g_i^{-1} \varphi(g_j) g_j^{-1}$ , i.e.  $g_i^{-1} \psi(g_j) = \psi(g_j) g_i^{-1}$ .

So  $\psi(g_j)$  commutes with  $g_i^{-1}$ , and hence it commutes with  $g_i$ .

(iii  $\implies$  ii) This is clear from the definition of  $\psi$ . □

**Examples:** 1. Suppose that  $\varphi \in \text{Aut}^\omega Z(G)$  is trivial mod centre, i.e.  $\varphi(g) g^{-1} \in Z(G)$  for all  $g \in G$ . Then for any ordered basis  $\underline{g} = \{g_1, \dots, g_d\}$ ,  $\varphi(g_i) g_i^{-1}$  is central, and so commutes with  $g_1, \dots, g_i$ .

So by the proposition,  $\varphi$  is a Mahler automorphism with respect to any basis. In fact, the proposition shows that if  $\varphi$  is a Mahler automorphism with respect to any basis, then  $\varphi$  is trivial mod centre. But it is possible for  $\varphi$  to be a Mahler automorphism with respect to *some* basis, and yet not be trivial mod centre.

2. Recall from [9] that  $G$  is *abelian-by-procyclic* if  $G \cong \mathbb{Z}_p^d \rtimes \mathbb{Z}_p$ . Then if we take  $\varphi \in \text{Inn}(G)$  to be the automorphism defining the action of  $\mathbb{Z}_p$  on  $\mathbb{Z}_p^d$ , then  $\varphi$  is a Mahler automorphism with respect to any basis of the form  $\{h_1, \dots, h_d, X\}$ , where  $\{h_1, \dots, h_d\}$  is a basis for  $\mathbb{Z}_p^d$ .

Note that in this case, if  $G$  is not nilpotent, or has nilpotency class greater than 2, then  $\varphi$  will not be trivial mod centre.

Since we want  $\varphi$  to be a Mahler automorphism with respect to any basis, we will assume from now on that  $\varphi$  is trivial mod centre. However, it should be noted that to prove general control theorems for Iwasawa algebras in characteristic  $p$  for non-nilpotent groups, the theory of general Mahler automorphisms is highly useful, as demonstrated by the approach in [9], so this theory should be developed further.

## 3.2 Approximations

From now on, we will fix  $P$  a faithful, prime ideal of  $\mathcal{O}G$  such that  $\pi \notin P$ ,  $\varphi \in \text{Aut}^\omega(G)$  an automorphism, trivial mod centre,  $\varphi \neq \text{id}$ , and we assume that  $\varphi(P) = P$ .

Using Theorem 2.5, we fix a non-commutative valuation  $v$  on  $Q(\mathcal{O}G/P)$ , and let  $\tau : \mathcal{O}G \rightarrow Q(\mathcal{O}G/P)$  be the natural map, which is continuous.

Recall from [1, Proposition 4.9] that if we define  $z(\varphi) : G \rightarrow Z(G)$ ,  $g \mapsto \lim_{m \rightarrow \infty} (\varphi^{p^m}(g) g^{-1})^{p^{-m}}$ , then  $z(\varphi)$  is a group homomorphism,  $z(\varphi^{p^m})(g) = z(\varphi)(g)^{p^m}$  for all  $g \in G$ , and:

$$\varphi^{p^m}(g) g^{-1} \equiv z(\varphi)(g)^{p^m} \pmod{Z(G)^{p^{2m}}} \text{ for all } m \in \mathbb{N}.$$

Using Theorem 2.5, choose an integer  $m_1 > 0$  such that for all  $Z \in Z(G)$ ,  $v(\tau(Z^{p^{m_1}} - 1)) > v(p)$ , and from now on, fix  $z := z(\varphi^{p^{m_1}}) : G \rightarrow Z(G)$ .

Set  $\lambda := \inf\{v(\tau(z(g) - 1)) : g \in G\}$ , then since  $P$  is faithful and  $\varphi \neq id$ , the proof of [1, Lemma 7.5] shows that  $\lambda < \infty$ . Furthermore, since  $z(g) = z(\varphi^{p^{m_1}})(g) = z(\varphi)(g)^{p^{m_1}}$ , it follows from our choice of  $m_1$  that  $\lambda > v(p)$ .

Now, for any ordered basis  $\underline{g} = \{g_1, \dots, g_d\}$  for  $G$ , set  $q_{i,m} := \tau(z(g_i)^{p^m} - 1)$  for each  $i \leq d$ ,  $m \in \mathbb{N}$ . We write  $\underline{q}_m^\alpha := q_{1,m}^{\alpha_1} q_{2,m}^{\alpha_2} \cdots q_{d,m}^{\alpha_d}$  as usual, and it is clear that  $v(q_{i,0}) = \lambda$  for some  $i$ .

Moreover, if  $v(q_{i,0}) = \lambda$  then  $v(q_{i,m}) = v(\tau(z(g_i)^{p^m} - 1)) = \lambda + mv(p)$  by Lemma 2.4.

The following result is analogous to [1, Proposition 7.7]:

**Proposition 3.2.** *For any  $\alpha \in \mathbb{N}^d$  with  $\alpha \neq 0$ , we have that for sufficiently high  $m \in \mathbb{N}$ :*

$$v(\tau(m_\alpha(\varphi^{p^m})) - \underline{q}_m^\alpha) \geq (2m - m_1)v(p) + (|\alpha| - 1)\lambda$$

Where  $|\alpha| := \alpha_1 + \cdots + \alpha_d$ .

*Proof.* We have that  $m_\alpha(\varphi^{p^m}) = (\varphi^{p^m}(g_1)g_1^{-1} - 1)^{\alpha_1} \cdots (\varphi^{p^m}(g_d)g_d^{-1} - 1)^{\alpha_d}$  since  $\varphi^{p^m}$  is a Mahler automorphism, so let  $\delta_{i,m} := \tau(\varphi^{p^m}(g_i)g_i^{-1} - 1) - q_{i,m}$ , then:

$$\tau(m_\alpha(\varphi^{p^m})) = (\delta_{1,m} + q_{1,m})^{\alpha_1} \cdots (\delta_{d,m} + q_{d,m})^{\alpha_d}$$

Hence  $m_\alpha(\varphi^{p^m}) - \underline{q}_m^\alpha$  is a sum of words in  $q_{i,m}$  and  $\delta_{i,m}$  of length  $|\alpha|$ , each containing at least one  $\delta_{i,m}$ .

So, it remains to show that  $v(\delta_{i,m}) \geq (2m - m_1)v(p)$  for each  $i, m$ , and the result follows.

Since  $\varphi^{p^{m+m_1}}(g_i)g_i^{-1} \equiv z(g_i)^{p^m} \pmod{Z(G)^{p^{2(m+m_1)}}}$ , if we assume that  $m \geq m_1$  we have that  $\varphi^{p^m}(g_i)g_i^{-1} = z(g_i)^{p^{m-m_1}} \epsilon_{i,m}^{p^{2m}}$  for some  $\epsilon_{i,m} \in Z(G)$ . Hence:

$$\delta_{i,m} = \tau(z(g_i)^{p^{m-m_1}} \epsilon_{i,m}^{p^{2m}} - 1) - \tau(z(g_i)^{p^{m-m_1}} - 1) = \tau(z(g_i)^{p^{m-m_1}}) \tau(\epsilon_{i,m}^{p^{2m}} - 1)$$

Since  $\epsilon_{i,m} \in Z(G)$  and  $m > m_1$ , we have that  $v(\tau(\epsilon_{i,m}^{p^{m_1}} - 1)) \geq v(p)$ , so by Lemma 2.4:

$$v(\tau(\epsilon_{i,m}^{p^{2m}} - 1)) = (2m - m_1)v(p) + v(\tau(\epsilon_{i,m}^{p^{m_1}} - 1))$$

Hence  $v(\delta_{i,m}) \geq (2m - m_1)v(p)$  as required.  $\square$

Now, we know that  $\tau\varphi^{p^m} = \sum_{\alpha \in \mathbb{N}^d} \tau(m_\alpha(\varphi^{p^m})) \tau \partial_{\underline{g}}^{(\alpha)}$ , so using the proposition, we set  $\varepsilon_m := \sum_{\alpha \neq 0} (\tau(m_\alpha(\varphi^{p^m})) - \underline{q}_m^\alpha) \tau \partial_{\underline{g}}^{(\alpha)}$ , and we have that  $v(\varepsilon_m(y)) \geq (2m - m_1)v(p)$  for all  $y \in \mathcal{O}G$ , and:

$$\tau\varphi^{p^m} - \tau = \sum_{\alpha \neq 0} \underline{q}_m^\alpha \tau \partial_{\underline{g}}^{(\alpha)} + \varepsilon_m \tag{6}$$

Now, for each  $i = 1, \dots, d$ , set  $\partial_i := \partial_{\underline{g}}^{(e_i)}$ , where  $e_i$  is the standard  $i$ 'th basis vector. Then since  $\varphi(P) = P$ , we have that for any  $y \in P$ ,  $(\tau\varphi^{p^m} - \tau)(y) = 0$ , so:

$$0 = q_{1,m} \tau \partial_1(y) + \cdots + q_{d,m} \tau \partial_d(y) + \sum_{|\alpha| \geq 2} \underline{q}_m^\alpha \tau \partial_{\underline{g}}^{(\alpha)}(y) + \varepsilon_m(y) \tag{7}$$

We want to analyse convergence of this expression, from which we hope to deduce a control theorem.

### 3.3 Drawbacks in characteristic 0

So far, these techniques are identical to the methods outlined and used in [1] and [9], but in both these cases, the underlying ring had characteristic  $p$ . When working in characteristic zero, potentially fatal problems arise.

The key issue is the linear growth rate of the Mahler approximations  $q_{i,m} = \tau(z(g_i)^{p^m} - 1)$ , as described in Lemma 2.4, which arises due to our assumption that  $\pi \notin P$ . Suppose we assume instead that  $\pi \in P$ , and hence  $Q(\mathcal{O}G/P)$  has characteristic  $p$ . In this case:

$$q_{i,m} := \tau(z(g_i)^{p^m} - 1) = \tau(z(g_i) - 1)^{p^m} \text{ for each } i.$$

Therefore,  $v(q_{i,m}) = p^m \lambda$ , and hence  $v(q_m^\alpha) \geq p^m |\alpha| \lambda$ , so the Mahler approximations grow exponentially with  $m$ . In this case we may divide out our expression (7) by an expression involving only the lower order terms  $q_{i,m}$ , and this will not affect the convergence of the higher order terms.

Using this approach, we can deduce that  $\tau \partial_i(y) = 0$  for some  $i$ , and a control theorem will follow. This is outlined in full detail in [1, Section 7].

But in our case, we have that  $\pi \notin P$ , and hence  $Q(\mathcal{O}G/P)$  has characteristic 0, and we see that  $v(q_m^\alpha) \geq m |\alpha| v(p) + |\alpha| \lambda$  using Lemma 2.4, and equality holds for certain values of  $\alpha$ .

This means that the Mahler approximations grow linearly with  $m$ , and it is not possible to remove all the lower order terms in expression (7) without affecting convergence of the higher order terms.

We will now explore particular instances when we can deduce a convergence argument. These instances occur when we impose restrictions on the ideal  $P$  to ensure that the centre is as small as possible, and at present they seem like the strongest we can obtain from Mahler expansions in characteristic zero.

### 3.4 Using Compactness

From now on, we will assume, as in the statement of Theorem 1.3, that we can find a closed, isolated subgroup  $A$  of  $Z(G)$  such that  $\mathcal{O}A/\mathcal{O}A \cap P$  is finitely generated over  $\mathcal{O}$  for some  $n \in \mathbb{N}$ , and we will suppose that  $\varphi(g)g^{-1} \in A$  for all  $g \in G$ .

Then since  $\varphi^{p^{m+1}}(g)g^{-1} \in A$  for all  $g \in G$  and  $A$  is isolated, it follows that  $(\varphi^{p^{m+1}}(g)g^{-1})^{p^{-m}} \in A$  for all  $m$ . So since  $A$  is closed,  $z(g) = \lim_{m \rightarrow \infty} (\varphi^{p^{m+1}}(g)g^{-1})^{p^{-m}} \in A$ , and thus for all  $m \in \mathbb{N}$ :

$$q_{i,m} = \tau(z(g_i)^{p^m} - 1) \in \mathcal{O}A/\mathcal{O}A \cap P.$$

Now, fix  $R := \mathcal{O}A/\mathcal{O}A \cap P$ , then  $R$  is a subring of the  $K$ -algebra  $Q(\mathcal{O}G/P)$ . So let  $F$  be the  $K$ -span of  $R$ , then since  $R$  is a domain, finitely generated over  $\mathcal{O}$ ,  $F$  is a finite field extension of  $K$ , and clearly  $F \subseteq Z(Q(\mathcal{O}G/P))$ .

So, let  $\mathcal{V}$  be the valuation ring for  $F$ , and let  $\mu \in \mathcal{V}$  be a uniformiser. Then since  $v|_F$  is a valuation on  $F$ , it follows that  $v|_F$  is a scalar multiple of the standard  $\mu$ -adic valuation on  $F$ , and hence  $\mathcal{V} = \{\beta \in F : v(\beta) > 0\}$ .

Now, choose  $i$  such that  $q_{i,0}$  has value  $\lambda$ , we may assume without loss of generality that  $i = 1$ . Then since  $\lambda > v(p)$ ,  $v(q_{1,m}) = mv(p) + \lambda$  for all  $m$  by Lemma 2.4.

Now, the key property of  $\mathcal{V}$  which we can exploit is compactness, which implies that any sequence in  $\mathcal{V}$  has a convergent subsequence. Using this notion, we obtain the following result:

**Proposition 3.3.** *For any ordered basis  $\underline{g} = \{g_1, \dots, g_d\}$ , set  $q_{i,m} := \tau(z(g_i)^{p^m} - 1)$  with  $v(q_{1,0}) = \lambda$ . Then for each  $i = 1, \dots, d$ , the sequence  $q_{1,m}^{-1}q_{i,m}$  has a subsequence converging to some  $\beta_i \in \mathcal{V}$  as  $m \rightarrow \infty$ , and  $(\beta_1\tau\partial_1 + \dots + \beta_d\tau\partial_d)(P) = 0$ .*

*Proof.* For each  $m$ ,  $q_{1,m}$  is  $v$ -regular of value  $\lambda + mv(p)$ , so since  $v(q_{i,m}) \geq \lambda + mv(p)$ , it follows that  $v(q_{1,m}^{-1}q_{i,m}) \geq \lambda + mv(p) - \lambda - mv(p) = 0$ , so  $q_{1,m}^{-1}q_{i,m} \in \mathcal{V}$ .

Therefore, using compactness of  $\mathcal{V}$ , we can choose a subsequence  $\underline{a} := (m_1, m_2, \dots)$  with  $m_1 < m_2 < m_3 < \dots$  such that  $q_{1,m_j}^{-1}q_{i,m_j}$  converges in  $\mathcal{V}$  as  $j \rightarrow \infty$ .

Let  $\beta_i := \lim_{j \rightarrow \infty} q_{1,m_j}^{-1}q_{i,m_j} \in \mathcal{V}$ .

Now, given  $\alpha \in \mathbb{N}^d$  with  $|\alpha| \geq 2$ ;  $v(q_{1,m}^{-1}q_m^\alpha) = v(q_m^\alpha) - v(q_{1,m}) \geq m|\alpha|v(p) + |\alpha|\lambda - mv(p) - \lambda \geq mv(p) + \lambda \rightarrow \infty$  as  $m \rightarrow \infty$ .

Also,  $v(q_{1,m}^{-1}\varepsilon_m) = v(\varepsilon_m) - v(q_{1,m}) \geq (2m - m_1)v(p) - mv(p) - \lambda = (m - m_1)v(p) - \lambda \rightarrow \infty$  as  $m \rightarrow \infty$ .

Therefore,  $q_{1,m}^{-1}q_m^\alpha \rightarrow 0$  as  $m \rightarrow \infty$ ,  $q_{1,m}^{-1}\varepsilon_m \rightarrow 0$  as  $m \rightarrow \infty$ . So, dividing out our expression (7) by  $q_{1,m}$ , we obtain:

$$0 = q_{1,m}^{-1}q_{1,m}\tau\partial_1(y) + q_{1,m}^{-1}q_{2,m}\tau\partial_2(y) + \dots + q_{1,m}^{-1}q_{d,m}\tau\partial_d(y) + \sum_{|\alpha| \geq 2} q_{1,m}^{-1}q_m^\alpha \tau\partial_{\underline{g}}^{(\alpha)}(y) + q_{1,m}^{-1}\varepsilon_m(y) \quad (8)$$

So considering the subsequence associated with  $\underline{a} = (m_1, m_2, \dots)$ , and taking the limit, we get that  $\beta_1\tau\partial_1(y) + \beta_2\tau\partial_2(y) + \dots + \beta_d\tau\partial_d(y) = 0$ .

Since our choice of  $y \in P$  was arbitrary, it follows that  $(\beta_1\tau\partial_1 + \dots + \tau\partial_d)(P) = 0$  as required.  $\square$

**Note:** These elements  $\beta_i$  depend on the choice of ordered basis  $\underline{g}$ .

From now on, fix an ordered basis  $\underline{g} = \{g_1, \dots, g_d\}$ , and fix  $\beta_1, \dots, \beta_d$  as in the statement of Proposition 3.3, and define:

$$h_{\underline{g}} : \mathcal{O}G \rightarrow Q(\mathcal{O}G/P), x \mapsto (\beta_1\tau\partial_1 + \beta_2\tau\partial_2 + \dots + \beta_d\tau\partial_d)(x)$$

Then  $h_{\underline{g}} \in \text{Hom}_{\mathcal{O}}(\mathcal{O}G, Q(\mathcal{O}G/P))$ , and it follows from Proposition 3.3 that  $h_{\underline{g}}(P) = 0$ , and thus  $h_{\underline{g}} \in \text{Hom}_{\mathcal{O}}(\mathcal{O}G/P, Q(\mathcal{O}G/P))$ .

Also, since each  $\beta_i$  lies in  $F = (\mathcal{O}A/\mathcal{O}A \cap P) \otimes_{\mathcal{O}} K$ , it follows that the image of  $\mathcal{O}G/P$  under  $h_{\underline{g}}$  lies in  $(\mathcal{O}G/P) \otimes_{\mathcal{O}} K = KG/(P \otimes_{\mathcal{O}} K)$ .

So since it is clear that each  $\partial_i$  extends to a  $K$ -linear endomorphism of  $KG$ , we may assume that  $h_{\underline{g}}$  lies in  $\text{Hom}_{\mathcal{O}}(KG/(P \otimes_{\mathcal{O}} K), KG/(P \otimes_{\mathcal{O}} K))$ , and hence it makes sense to raise  $h_{\underline{g}}$  to integer powers.

We now need some technical results:

**Lemma 3.4.** Fix  $R = \frac{\mathcal{O}A}{\mathcal{O}A \cap P}$ , then there exists  $s \in \mathbb{N}$  such that  $\pi^s \mathcal{V} \subseteq R$ . Also, there exists  $t \in \mathbb{N}$  such that if  $x \in \mathcal{V}$  and  $v(x) > 0$  then  $x^t \in \pi R$ .

*Proof.* Since  $R$  is a lattice in  $F$ , which is a finite dimensional  $\mathbb{Q}_p$ -vector space, it follows that  $R$  is a free  $\mathbb{Z}_p$ -module of rank  $\dim_{\mathbb{Q}_p} F$ . This is also the rank of  $\mathcal{V}$ , and it follows that  $R$  has finite index in  $\mathcal{V}$ , and hence  $p^l \mathcal{V} \subseteq R$  for some  $l \in \mathbb{N}$ , and the first statement follows.

Now, if  $v(x) > 0$ , then  $x = \pi^{-k} r$  for some  $r \in R$ ,  $k \in \mathbb{N}$ , and  $v(r) - kv(\pi) \geq 1$ . We know that there exists  $s \in \mathbb{N}$  such that  $\pi^s \mathcal{V} \subseteq R$ , so choose  $t \in \mathbb{N}$  such that  $t \geq (s+1)v(\pi)$ .

Then  $x^t = \pi^{-kt} r^t = \pi^{s+1} (\pi^{-(kt+s+1)} r^t)$ , and note that:

$$v(\pi^{-(kt+s+1)} r^t) = t(v(r) - kv(\pi)) - (s+1)v(\pi) \geq t - (s+1)v(\pi) \geq 0$$

Thus  $\pi^{-(kt+s+1)} r^t \in \mathcal{V}$ , so since  $\pi^{s+1} \mathcal{V} \subseteq \pi R$  and  $x^t = \pi^{s+1} (\pi^{-(kt+s+1)} r^t)$ , it follows that  $x^t \in \pi R$  as required.  $\square$

Now, recall from the definition of  $\partial_1, \dots, \partial_d$ , that if  $g = \underline{g}^\alpha \in G$  for some  $\alpha \in \mathbb{Z}_p^d$ , then  $\partial_i(g) = \alpha_i g$ . So if we let  $k_{\underline{g}, \underline{g}} := \beta_1 \alpha_1 + \beta_2 \alpha_2 + \dots + \beta_d \alpha_d \in \mathcal{V}$ , then  $h_{\underline{g}}(g) = k_{\underline{g}, \underline{g}} g$  for all  $g \in G$ .

**Lemma 3.5.** For every ordered basis  $\underline{g}$  of  $G$ ,  $h_{\underline{g}}$  is  $F$ -linear.

*Proof.* Since  $h_{\underline{g}}(a) = k_{a, \underline{g}} a$  for every  $a \in A$ , and  $k_{a, \underline{g}} \in F$ , it follows that  $h_{\underline{g}}$  sends  $\mathcal{O}A$  to  $F = \mathcal{O}A / (\mathcal{O}A \cap P) \otimes_{\mathcal{O}} K$ , and hence it sends  $F$  to  $F$ , i.e.  $h_{\underline{g}}$  restricts to a  $K$ -linear endomorphism of  $F$ .

Also, for every  $i$ ,  $\partial_i$  is a  $K$ -linear derivation of  $\mathcal{O}G$ . So since each  $\beta_i$  is central, it follows that  $h_{\underline{g}} = \beta_1 \tau \partial_1 + \dots + \beta_d \tau \partial_d$  is also a  $K$ -linear derivation, so  $h_{\underline{g}}$  restricts to a  $K$ -linear derivation of  $F$ .

Using the derivation property, we see that to prove that  $h_{\underline{g}}$  is  $F$ -linear, it remains only to prove that  $h_{\underline{g}}(F) = 0$ . We will show, in fact, that all  $K$ -linear derivations of  $F$  are zero.

Suppose that  $\alpha \in F$ , then since  $F$  is a finite extension of  $K$ ,  $\alpha$  is the root of some polynomial  $f(x) = a_0 + a_1 x + \dots + a_n x^n$  with coefficients in  $K$ , and we will assume that  $n$  is minimal. Let  $\delta$  be a  $K$ -linear derivation of  $F$ , then:

$$\begin{aligned} 0 &= \delta(0) = \delta(a_0 + a_1 \alpha + \dots + a_n \alpha^n) = a_1 \delta(\alpha) + \dots + a_n \delta(\alpha^n) \\ &= a_1 \delta(\alpha) + \dots + n a_n \alpha^{n-1} \delta(\alpha) = \delta(\alpha) (a_1 + 2a_2 \alpha + \dots + n a_n \alpha^{n-1}). \end{aligned}$$

So if  $\delta(\alpha) \neq 0$  then  $a_1 + 2a_2 \alpha + \dots + n a_n \alpha^{n-1} = 0$ , contradicting minimality of  $n$ . Hence  $\delta(\alpha) = 0$ , meaning that  $\delta = 0$ .  $\square$

Now, let  $f$  be the degree of the residue field of  $F$ , then for all  $\beta \in \mathcal{V}$ :

$$\beta^{p^f - 1} \equiv \begin{cases} 1 & v(\beta) = 0 \\ 0 & v(\beta) > 0 \end{cases} \pmod{\mu \mathcal{V}} \quad (9)$$

It follows easily that for all  $n \in \mathbb{N}$ :

$$\beta^{p^n(p^f-1)} \equiv \begin{cases} 1 & v(\beta) = 0 \\ 0 & v(\beta) > 0 \end{cases} \pmod{\mu^{n+1}\mathcal{V}} \quad (10)$$

Using Lemma 3.4, choose an integer  $t_0 \geq 0$  such that  $\mu^{t_0+1}\mathcal{V} \subseteq \pi R$  where  $R = \mathcal{O}A/\mathcal{O}A \cap P$  and  $x^{p^{t_0}} \in \pi R$  for all  $x \in \mathcal{V}$  with  $v(x) > 0$ . Then examining (10) shows that  $\beta^{p^{t_0}(p^f-1)} \in R \subseteq \mathcal{O}G/P$  for all  $\beta \in \mathcal{V}$ .

Also, using Lemma 3.5, we see that  $h_{\underline{g}}$  is  $F$ -linear. So since  $k_{\underline{g},g} \in F$ , it follows that  $h_{\underline{g}}^n(g) = k_{\underline{g},g}^n g$  for all  $g \in G$ ,  $n \in \mathbb{N}$ .

Therefore, if we define  $w_{\underline{g}} := h_{\underline{g}}^{p^{t_0}(p^f-1)}$ , then for all  $g \in G$ ,  $n \in \mathbb{N}$ , since  $k_{\underline{g},g}^{p^{n+t_0}(p^f-1)} \in \mathcal{O}G/P$ , we have that  $w_{\underline{g}}^{p^n}(g) = k_{\underline{g},g}^{p^{n+t_0}(p^f-1)} g \in \mathcal{O}G/P$ , and thus  $w_{\underline{g}}$  sends  $\frac{\mathcal{O}G}{P}$  to  $\frac{\mathcal{O}G}{P}$ .

Fixing  $g \in G$ ,  $n \in \mathbb{N}$ , using (10) we have:

$$w_{\underline{g}}^{p^n}(g) \equiv \begin{cases} 1 & v(k_{\underline{g},g}) = 0 \\ 0 & v(k_{\underline{g},g}) > 0 \end{cases} \pmod{\pi^{n+1}\frac{\mathcal{O}G}{P}}$$

So, setting  $S := \frac{\mathcal{O}G}{P}$  for convenience, consider the composition:

$$\mathcal{O}G \xrightarrow{\tau} S \xrightarrow{w_{\underline{g}}^{p^n}} S$$

Let  $\iota_n$  be this composition, then  $\iota_n(\pi^m \mathcal{O}G) \subseteq \pi^m S$  for all  $m$ , and  $\iota_n(g) \equiv \begin{cases} g & v(k_{\underline{g},g}) = 0 \\ 0 & v(k_{\underline{g},g}) > 0 \end{cases} \pmod{\pi^{n+1}S}$ . Hence  $\iota_n(r) \equiv \iota_{n+1}(r) \pmod{\pi^{n+1}S}$  for all  $n \in \mathbb{N}$ ,  $r \in \mathcal{O}G$ .

Therefore, since  $S$  is  $\pi$ -adically complete, there exists a continuous,  $\mathcal{O}$ -linear morphism  $\iota : \mathcal{O}G \rightarrow S$  such that  $\iota(P) = 0$ ,  $\iota(x) \equiv \iota_n(x) \pmod{\pi^{n+1}S}$  for all  $n \in \mathbb{N}$ , hence for all  $g \in G$ :

$$\iota(g) = \begin{cases} g & v(k_{\underline{g},g}) = 0 \\ 0 & v(k_{\underline{g},g}) > 0 \end{cases} \quad (11)$$

### 3.5 The Controlling subgroup

We will now complete our proof of Theorem 1.3 by finding an appropriate subgroup of  $G$  controlling our ideal  $P$ .

Let  $U := \{g \in G : v(\tau(z(g) - 1)) > \lambda\}$ , then using the proof of [1, Lemma 7.6] we see that  $U$  is a proper, open subgroup of  $G$  containing  $G^p$ . So fix an ordered basis  $\underline{g} = \{g_1, \dots, g_d\}$  for  $G$  such that  $\{g_1^p, \dots, g_r^p, g_{r+1}, \dots, g_d\}$  is an ordered basis for  $U$ . We want to prove that  $P$  is controlled by  $U$ .

Recall from the previous subsection the definition of  $k_{\underline{g},g} \in \mathcal{V}$  for each  $g \in G$ :

**Lemma 3.6.**  $U = \{g \in G : v(k_{\underline{g},g}) > 0\}$

*Proof.* Using Proposition 3.3, we see that for each  $i = 1, \dots, d$ ,  $\beta_i = \lim_{j \rightarrow \infty} q_{1,m_j}^{-1} q_{i,m_j}$ , as  $m_j$  runs over some subsequence  $\underline{a} = (m_1, m_2, \dots)$ , where  $q_{i,m} = \tau(z(g_i)^{p^m} - 1)$ .

By Lemma 2.4,  $v(q_{i,m}) = \lambda + mv(p)$  for  $i \leq r$ , and  $v(q_{i,m}) > \lambda + mv(p)$  for all  $i > r$ . Hence  $v(\beta_i) = 0$  for all  $i \leq r$ ,  $v(\beta_i) > 0$  for all  $i > r$ .

Given  $g \in U$ ,  $g = g_1^{p\alpha_1} \cdots g_r^{p\alpha_r} g_{r+1}^{\alpha_{r+1}} \cdots g_d^{\alpha_d}$  for  $\alpha_i \in \mathbb{Z}_p$ , so:

$$v(k_{g,\underline{g}}) = v(p\alpha_1\beta_1 + p\alpha_2\beta_2 + \cdots + p\alpha_r\beta_r + \alpha_{r+1}\beta_{r+1} + \cdots + \alpha_d\beta_d) > 0$$

Conversely, if  $v(k_{g,\underline{g}}) > 0$ , suppose  $g = \underline{g}^\alpha$ , so  $k_{g,\underline{g}} = \beta_1\alpha_1 + \beta_2\alpha_2 + \cdots + \beta_d\alpha_d$ .

By the definition of  $\beta_i$ , this means that  $v(q_{1,m_j}^{-1}(\alpha_1 q_{1,m_j} + \cdots + \alpha_d q_{d,m_j})) > 0$  for sufficiently high  $j$ , and hence  $v(\alpha_1 q_{1,m_j} + \cdots + \alpha_r q_{d,m_j}) > \lambda + m_j v(p)$ .

But it is easily seen that  $\tau(z(g)^{p^{m_j}} - 1) \equiv \alpha_1 q_{1,m_j} + \cdots + \alpha_d q_{d,m_j} \pmod{\lambda + m_j v(p) + 1}$ , and hence  $v(\tau(z(g)^{p^{m_j}} - 1)) > \lambda + m_j v(p)$  for sufficiently high  $j$ .

But  $v(\tau(z(g)^{p^{m_j}} - 1)) = v(\tau(z(g) - 1)) + m_j v(p)$  by Lemma 2.4, and hence  $v(\tau(z(g) - 1)) > \lambda$  and  $g \in U$  as required.  $\square$

Now, recall the definition of the continuous  $\mathcal{O}$ -linear morphism  $\iota : \mathcal{O}G \rightarrow \frac{\mathcal{O}G}{P}$ . Then using

the lemma, we deduce that  $\iota(g) = \begin{cases} g & g \notin U \\ 0 & g \in U \end{cases}$ .

Define  $f : G \rightarrow \mathcal{O}, g \mapsto \begin{cases} 1 & g \notin U \\ 0 & g \in U \end{cases}$ , then clearly  $f \in C(G, \mathcal{O})$  is locally constant, so the

endomorphism  $\rho(f) \in \text{End}_{\mathcal{O}}(\mathcal{O}G)$  is well defined, and  $\rho(f)(g) = f(g)g = \begin{cases} g & g \notin U \\ 0 & g \in U \end{cases}$ .

Therefore  $\tau\rho(f) = \iota$  when restricted to  $\mathcal{O}[G]$ , so since  $\iota$  is continuous,  $\tau$  is continuous, and  $\rho(f)$  is continuous, it follows that  $\tau\rho(f) = \iota$ . Hence  $\tau\rho(f)(P) = 0$  and  $\rho(f)(P) \subseteq P$ .

**Proposition 3.7.** *P is controlled by U.*

*Proof.* Firstly, suppose that  $C = \{x_1, \dots, x_t\}$  is a complete set of coset representatives for  $U$  in  $G$ , then for all  $r \in \mathcal{O}G$ ,  $r = \sum_{i \leq t} r_i x_i$  for some  $r_i \in \mathcal{O}U$ .

Suppose we can choose  $C$  such that if  $r \in P$  then  $r_1 \in P \cap \mathcal{O}U$ . Then since  $rx_1^{-1}x_i \in P$  for all  $i = 1, \dots, t$  and  $rx_1^{-1}x_i$  has  $x_1$  component  $r_i$ , it follows that  $r_i \in P \cap \mathcal{O}U$  for each  $i$ , and hence  $P$  is controlled by  $U$ .

So it remains to prove that we can choose such a set  $C$  of coset representatives such that if  $\sum_{i \leq t} r_i x_i \in P$ , then at least one of the  $r_i$  lies in  $P \cap \mathcal{O}U$ .

Since  $U$  has ordered basis  $\{g_1^p, \dots, g_r^p, g_{r+1}, \dots, g_d\}$  it follows that  $C = \{g_1^{b_1} \cdots g_r^{b_r} : 0 \leq b_i < p\}$  is a complete set of coset representatives for  $U$  in  $G$ .

So for each  $\underline{b} \in [p-1]^r$ , let  $g_{\underline{b}} = g_1^{b_1} \cdots g_r^{b_r}$  (here  $[p-1] = \{0, 1, \dots, p-1\}$ ).

Then if  $r = \sum_{\underline{b} \in [p-1]^r} r_{\underline{b}} g_{\underline{b}} \in P$ , then  $\rho(f)(r) = \sum_{\underline{b} \in [p-1]^r} f(g_{\underline{b}}) r_{\underline{b}} g_{\underline{b}}$  by the proof of [1, Proposition 2.5], and this also lies in  $P$ .



But  $f(g_{\underline{b}}) = 1$  if  $\underline{b} \neq 0$ , and  $f(g_{\underline{0}}) = 0$  hence  $\rho(f)(r) = \sum_{\underline{b} \in [p-1]^r \setminus \{0\}} r_{\underline{b}} g_{\underline{b}} \in P$ .

Therefore,  $r_{\underline{0}} g_{\underline{0}} = r - \rho(f)(r) \in P$ , and thus  $r_{\underline{0}} \in P \cap \mathcal{O}U$  as required.  $\square$

The main theorem of this section follows immediately:

*Proof of Theorem 1.3.* Since we are assuming that  $\mathcal{O}A/\mathcal{O}A \cap P$  is finitely generated over  $\mathcal{O}$ , it follows from Proposition 3.7 that  $P$  is controlled by a proper open subgroup of  $G$ .  $\square$

## 4 Control Theorem

Now that we have established Theorem 1.3, we will conclude in this section with our main control theorem. Throughout, we will fix  $G$  a compact  $p$ -adic Lie group.

### 4.1 Technical results in characteristic 0

In [1, Section 5], a number of technical results were stated and proved for completed group algebras in characteristic  $p$ . These results are fundamental for the study of ideals in Iwasawa algebras, so we will first carry them over to a characteristic zero setting.

**Lemma 4.1.** *Let  $H$  be a closed subgroup of  $G$ , and let  $I_1, \dots, I_m, J$  be right ideals of  $KH$ . Then:*

- (i)  $I_1 KG \cap \dots \cap I_m KG = (I_1 \cap \dots \cap I_m) KG$ .
- (ii)  $JKG \cap KH = J$ .

*Proof.* We will prove that  $KG$  is faithfully flat over  $KH$ . Then part (i) follows from applying the functor  $- \otimes_{KH} KG$  to the short exact sequence  $0 \rightarrow I_1 \cap \dots \cap I_m \rightarrow KH \rightarrow \bigoplus_{j \leq m} \frac{KH}{I_j} \rightarrow 0$ , and part (ii) follows from [12, Lemma 7.2.5], taking  $R = KH$ ,  $S = KG$  and  $M = \frac{KH}{J}$ .

By [6, Lemma 4.5],  $\mathcal{O}G$  is projective over  $\mathcal{O}H$ , and hence it is flat. It follows immediately that  $KG$  is flat over  $KH$ , so it remains to prove that  $KG$  is faithful, i.e. if  $M \neq 0$  is a  $KH$ -module, then  $KG \otimes_{KH} M \neq 0$ .

Note that  $K \cong KH/(H-1)KH$  is a  $KH$ -module, so if  $KG \otimes_{KH} M = 0$  then we have the following isomorphisms of  $K$ -spaces:

$$0 = KG \otimes_{KH} M \cong KG \otimes_{KH} K \otimes_K M$$

But since  $K$  is a field, every module over  $K$  is faithfully flat, so if  $M \neq 0$ , it follows that  $KG \otimes_{KH} K = 0$ , which is impossible since the natural map  $KG \otimes_{KH} K \rightarrow K, r \otimes \alpha \rightarrow r\alpha + (G-1)KG$  is surjective. Hence  $KG$  is faithfully flat as required.  $\square$

Using this Lemma, we can now carry over the proofs of every result in [1, Chapter 5], with the exception of [1, Lemma 5.3], whose proof strongly depends on the assumption that the ground field has characteristic  $p$ .

**Lemma 4.2.** *Let  $G$  be a  $p$ -valuable group, let  $I$  be a two-sided ideal of  $KG$  such that  $I = J_1 \cap \dots \cap J_r$  for some right ideals  $J_i$  of  $KG$  forming a complete  $G$ -orbit via the conjugation action. Then if  $I$  is faithful, each  $J_i$  is faithful.*

*Proof.* For any ideal  $J$  of  $KG$ , let  $J^\dagger := \{g \in G : g - 1 \in J\}$ . Then clearly  $J^\dagger$  is a closed subgroup of  $G$ , and if  $J$  is a two-sided ideal then it is a normal subgroup. Clearly  $J$  is faithful if and only in  $J^\dagger = 1$ .

It is also clear that  $(J_1)^\dagger, \dots, (J_r)^\dagger$  form a single  $G$ -orbit, and  $I^\dagger = (J_1)^\dagger \cap \dots \cap (J_r)^\dagger$ .

But since  $G$  is  $p$ -valuable, it follows from [1, Proposition 5.9] that  $G$  is *orbitally sound*, i.e. for any closed subgroup  $H$  of  $G$  with finitely many  $G$ -conjugates, the intersection of these conjugates is open in  $H$ . Therefore  $I^\dagger$  has finite index in  $(J_i)^\dagger$  for each  $i$ .

So if  $I$  is faithful, then  $(J_i)^\dagger$  is a finite subgroup of  $G$ . So since  $G$  is torsionfree, this means that  $(J_i)^\dagger = 1$ , and  $J_i$  is faithful as required.  $\square$

Let us now reintroduce some definitions from [1, Section 5]:

**Definition 4.1.** (i). Given a prime ideal  $P$  of  $KG$ , we say that  $P$  is non-splitting if for any open normal subgroup  $U$  of  $G$  controlling  $P$ ,  $P \cap KU$  is prime in  $KU$ .

(ii). Let  $\mathcal{P}$  be a property satisfied by two-sided ideals in  $KH$ , for  $H$  any compact  $p$ -adic Lie group. Then given a right ideal  $I$  of  $KG$ , we say that  $I$  virtually satisfies  $\mathcal{P}$  if there exists an open subgroup  $U$  of  $G$  and a two sided ideal  $J$  of  $KU$  such that  $J$  satisfies  $\mathcal{P}$  and  $I = JKG$ .

In particular,  $I$  is virtually non-splitting if  $I = PKU$  for some non-splitting prime ideal  $P$  of  $KU$ .

**Proposition 4.3.** Let  $P$  be a non-splitting prime ideal of  $KG$ , then  $P \cap KP^\times$  is prime in  $KP^\times$

*Proof.* This is the proof of [1, Proposition 5.5], applied using Lemma 4.1.  $\square$

Now, recall from [1, Definition 5.6] that if  $R$  is a ring,  $J_1, \dots, J_r$  are right ideals of  $R$  with intersection  $I$ , then  $I = J_1 \cap \dots \cap J_r$  is an *essential decomposition* for  $I$  if the  $R$ -module embedding  $\frac{R}{I} \rightarrow \frac{R}{J_1} \times \dots \times \frac{R}{J_r}$  has essential image in the sense of [12, Definition 2.2.1].

**Proposition 4.4.** Suppose that  $G$  is pro- $p$ , let  $P$  be a prime ideal of  $KG$ , and let  $P = I_1 \cap \dots \cap I_r$  be an essential decomposition for  $P$  such that each  $I_j$  is virtually prime and  $I_1, \dots, I_r$  form a single  $G$ -orbit. If we assume that  $r$  is as large as possible then each  $I_j$  is virtually non-splitting.

*Proof.* This is the proof of [1, Theorem 5.7], applied using Lemma 4.1.  $\square$

## 4.2 $J$ -Ideals

Recall that we are interested in *primitive ideals* in  $KG$ , that is ideals of the form  $\text{Ann}_{KG} M$  for some irreducible  $KG$ -module  $M$ . It is well known that these ideals are prime.

**Proposition 4.5.** Let  $P$  be a primitive ideal of  $KG$ , then  $Z(KG/P)$  is finite dimensional over  $K$ .

*Proof.* By definition,  $P = \text{Ann}_{KG} M$  for some irreducible  $KG$ -module  $M$ . Then it follows from [8, Theorem 1.1(1)] that the algebra  $\text{End}_{K[G]} M$  of  $G$ -equivariant  $K$ -linear endomorphisms of  $M$  is a finite dimensional  $K$ -algebra.

Now, there exists a well defined, injective homomorphism of  $K$ -algebras:

$$Z(KG/P) \rightarrow \text{End}_{K[G]}M, z + P \mapsto (\phi_z : M \rightarrow M, m \mapsto zm)$$

so it follows immediately that  $Z(KG/P)$  is finite dimensional over  $K$ .  $\square$

So, we can safely assume that  $P$  is a prime ideal of  $KG$  such that  $Z(KG/P)$  is a finite field extension of  $K$ , however it will be necessary for us to work in slightly more generality:

**Definition 4.2.** *Let  $G$  be a compact  $p$ -adic Lie group,  $Z := Z(G)$ . Given a right ideal  $I$  of  $KG$ , we say that  $I$  is a  $J$ -ideal of  $KG$  if  $KZ(G)/I \cap KZ(G)$  is finite dimensional over  $K$ .*

Using Proposition 4.5, it is clear that primitive ideals are  $J$ -ideals. The reason why we prefer to work with ideals of this form is because even when  $Z(KG/P)$  is finite dimensional, after passing to a smaller subgroup  $H$ , there is no reason why  $Z(KH/P \cap KH)$  should be finite dimensional. So it makes sense to only consider elements of the full centre  $Z(G)$ .

**Lemma 4.6.** *Let  $U$  be an open normal subgroup of  $G$ ,  $P$  a prime ideal of  $KU$  such that  $PKU$  is a  $J$ -ideal of  $KG$ . Then  $P$  is a  $J$ -ideal of  $KU$ .*

*Proof.* Since  $PKG \cap KU = P$  by Lemma 4.1(ii), we have an injection of  $KU$ -modules  $KU/P \hookrightarrow KG/PKG$ , and this map sends  $KZ(U)/KZ(U) \cap P$  to  $\frac{KZ(U)+KZ(G) \cap PKG}{KZ(G) \cap PKG}$ .

But  $Z(U) \subseteq Z(G)$  by [1, Lemma 8.4(b)], so this image is contained in  $KZ(G)/KZ(G) \cap PKG$ , which is finite dimensional over  $K$  by the definition of a  $J$ -ideal.

Hence  $KZ(U)/KZ(U) \cap P$  is finite dimensional over  $K$  as required.  $\square$

**Lemma 4.7.** *Suppose  $I$  is a  $J$ -ideal in  $KG$  and  $I \subseteq J$  for some right ideal  $J$  of  $KG$ . Then  $J$  is a  $J$ -ideal of  $KG$ .*

*Proof.* Setting  $Z := Z(G)$ , since  $I$  is a  $J$ -ideal,  $KZ/I \cap KZ$  is finite dimensional over  $K$ .

But  $I \subseteq J$ , so there is a surjection  $KG/I \rightarrow KG/J$  of  $KG$ -modules, and  $KZ/KZ \cap J$  is the image of  $KZ/KZ \cap I$  under this map.

So since  $KZ/KZ \cap J$  is the image of a finite dimensional  $K$ -vector space under a  $K$ -linear map, it is also finite dimensional over  $K$ , making  $J$  a  $J$ -ideal.  $\square$

**Theorem 4.8.** *Let  $G$  be a  $p$ -valuable group, let  $A$  be a closed subgroup of  $G$ , and suppose that all faithful, virtually non-splitting right  $J$ -ideals of  $KG$  are controlled by  $A$ . Then all faithful, prime  $J$ -ideals of  $KG$  are controlled by  $A$ .*

*Proof.* This is similar to the proof of [1, Theorem 5.8].

Let  $P$  be a faithful, prime  $J$ -ideal of  $KG$ , and let  $P = I_1 \cap \dots \cap I_m$  be an essential decomposition for  $P$ , with each  $I_j$  virtually prime, and  $I_1, \dots, I_r$  forming a single  $G$ -orbit.

Setting  $r = 1$ ,  $I_1 = P$ , it is clear that such a decomposition exists, so we will assume that  $r$  is maximal such that a decomposition of this form exists. We know that  $r$  is finite because  $KG/P$  has finite uniform dimension in the sense of [12].

So, by Theorem 4.4, each  $I_j$  is a virtually non-splitting right ideal of  $KG$ , and since  $P \subseteq I_j$  it follows from Lemma 4.7 that  $I_j$  is a  $J$ -ideal. Furthermore, since  $P$  is faithful, it follows from Lemma 4.2 that each  $I_j$  is faithful.

Therefore, by assumption,  $I_j$  is controlled by  $A$ , so  $I_j = (I_j \cap KA)KG$  for each  $j$ . So since  $P = I_1 \cap \cdots \cap I_r$ , we have that

$$\begin{aligned} (P \cap KA)KG &= ((I_1 \cap KA) \cap \cdots \cap (I_r \cap KA))KG \\ &= (I_1 \cap KA)KG \cap \cdots \cap (I_r \cap KA)KG = I_1 \cap \cdots \cap I_r = P \text{ by Lemma 4.1(i)}. \end{aligned}$$

Thus  $P$  is controlled by  $A$  as required.  $\square$

### 4.3 Controlling prime $J$ -ideals

Recall from section 1 the definition of the second centre  $Z_2(G) = \{g \in G : (g, G) \subseteq Z(G)\}$ .

**Lemma 4.9.** *For  $G$  any  $p$ -valuable group,  $U \leq_o G$ ,  $C_U(Z_2(U)) = C_G(Z_2(G)) \cap U$ .*

*Proof.* Firstly, we will show that  $Z_2(U) \subseteq Z_2(G)$ :

If  $g \in Z_2(U)$  then  $(g, U) \subseteq Z(U)$  by definition, and  $Z(U) \subseteq Z(G)$  by [1, Lemma 8.4(b)], so  $(g, U) \subseteq Z(G)$ .

But we know that  $G^{p^n} \subseteq U$  for some  $n \in \mathbb{N}$ , so for all  $h \in G$ ,  $(g, h^{p^n}) \in Z(G)$ .

But  $Z(G)$  is an isolated normal subgroup of  $G$  by [1, Lemma 8.4(a)], so by [10, IV.3.4.2],  $\frac{G}{Z(G)}$  carries a  $p$ -valuation  $\bar{\omega}$ . So since  $(g, h^{p^n}) \in Z(G)$ , we have that  $\bar{\omega}((g, h^{p^n})) = \infty$ .

But  $\bar{\omega}((g, h^{p^n})) = \bar{\omega}(g, h) + n$  by [13, Proposition 25.1], and hence  $\bar{\omega}((g, h)) = \infty$ . Thus  $(g, h) \in Z(G)$  for all  $h \in G$ , and  $g \in Z_2(G)$  as required.

Now, given  $g \in C_U(Z_2(U))$ ,  $(g, Z_2(U)) = 1$ , so if  $h \in Z_2(G)$  then  $h^{p^n} \in Z_2(G) \cap U = Z_2(U)$ , so  $(g, h^{p^n}) = 1$ . So since  $G$  is  $p$ -valued, applying [13, Proposition 25.1] again gives that  $(g, h) = 1$ , thus  $g \in C_G(Z_2(G))$  and  $C_U(Z_2(U)) \subseteq C_G(Z_2(G))$ .

Conversely, if  $g \in C_G(Z_2(G)) \cap U$  then  $(g, Z_2(G)) = 1$ , so since  $Z_2(U) \subseteq Z_2(G)$ ,  $(g, Z_2(U)) = 1$  and  $g \in C_U(Z_2(U))$  as required.  $\square$

**Theorem 4.10.** *Let  $G$  be a nilpotent,  $p$ -valuable group, let  $P$  be a faithful prime  $J$ -ideal of  $KG$ . Then  $P$  is controlled by  $C_G(Z_2(G))$ .*

*Proof.* First, suppose that  $P$  is non-splitting, and let  $H := P^\times$  be the controller subgroup of  $P$ :

Then  $Q := P \cap KH$  is a faithful, prime ideal of  $KH$  by Proposition 4.3, and since  $H$  is the smallest subgroup of  $G$  controlling  $P$ ,  $Q$  is not controlled by any proper subgroup of  $H$ .

Also, note that  $H$  is a normal subgroup of  $G$  by the proof of [1, Lemma 5.2], so for any  $g \in G$ ,  $(g, H) \subseteq H$ , and clearly  $A := Z(G) \cap H \subseteq Z(H)$  is the subgroup of  $G$ -invariants in  $H$ .

Since  $P$  is a  $J$ -ideal of  $KG$ ,  $KZ(G)/KZ(G) \cap P$  is finite dimensional over  $K$  by definition. And:

$$\frac{KA}{KA \cap Q} = \frac{KA}{KA \cap P} \subseteq \frac{KZ(G)}{KZ(G) \cap P}.$$

Therefore,  $KA/KA \cap Q$  is finite dimensional over  $K$ , and hence  $\mathcal{O}A/\mathcal{O}A \cap Q$  is finitely generated over  $\mathcal{O}$ .

So, given  $g \in Z_2(G)$ ,  $(g, H) \subseteq Z(G) \cap H = A$ , so let  $\varphi$  be the automorphism of  $H$  induced by conjugation by  $g$ , then  $\varphi(Q) = Q$  and  $\varphi(h)h^{-1} \in A$  for all  $h \in H$ . So applying Theorem 1.3 gives that if  $\varphi \neq 1$ , then  $Q$  is controlled by a proper subgroup of  $H$  – contradiction.

Therefore  $\varphi = 1$ , i.e.  $g$  centralises  $H$ .

Since our choice of  $g$  was arbitrary, we have now proved that  $Z_2(G)$  centralises  $H$ , and hence  $H$  is contained in the centraliser  $C_G(Z_2(G))$  of  $Z_2(G)$  in  $G$  as required. Thus  $P$  is controlled by  $C_G(Z_2(G))$ .

Now suppose that  $I \trianglelefteq_r KG$  is a faithful and virtually non-splitting right  $J$ -ideal of  $KG$ . Then  $I = PKG$  for some open subgroup  $U$  of  $G$ , and some faithful, non-splitting prime  $P$  of  $KU$ , and  $P$  is a  $J$ -ideal of  $KU$  by Lemma 4.6.

We have proved that  $P$  is controlled by  $C_U(Z_2(U))$ , and  $C_U(Z_2(U)) = C_G(Z_2(G)) \cap U$  by Lemma 4.9, and hence  $I$  is controlled by  $C_G(Z_2(G))$ .

So, using Theorem 4.8, it follows that every faithful, prime  $J$ -ideal of  $KG$  is controlled by  $C_G(Z_2(G))$ .  $\square$

We can now prove our main control theorem:

*Proof of Theorem 1.1.* Let  $P$  be a faithful, primitive ideal of  $KG$ . Then using Proposition 4.5, we see that  $Z(KG/P)$  is finite dimensional over  $K$ .

Let  $Z = Z(G)$ , then since  $KZ/P \cap KZ \subseteq Z(KG/P)$ , it follows that  $KZ/P \cap KZ$  is also finite dimensional over  $K$ , thus  $P$  is a  $J$ -ideal of  $KG$ .

So since  $P$  is a faithful, prime  $J$ -ideal of  $KG$ , it follows from Theorem 4.10 that  $P$  is controlled by  $C_G(Z_2(G))$ .  $\square$

## References

- [1] K. Ardakov, Prime ideals in nilpotent Iwasawa algebras. *Invent. Math.* **190**(2), 439-503 (2012).
- [2] K. Ardakov, The Controller Subgroup of One-sided ideals in Completed Group Rings. *Contemporary Mathematics, vol 562* (2012).
- [3] K. Ardakov; K. A. Brown, Ring-theoretic properties of Iwasawa algebras: a survey. *Documenta Math.*, Extra Volume Coates, 7-33 (2006).
- [4] K. Ardakov; S. Wadsley, On irreducible representations of compact  $p$ -adic analytic groups. *Annals of Mathematics* **178**, 453-557 (2013).
- [5] K. Ardakov; S. Wadsley, Verma modules for Iwasawa algebras are faithful *Münster Journal of Mathematics* **7**, 5-26 (2014).
- [6] A. Brumer. Pseudocompact algebras, profinite groups and class formations, *J. Algebra* **4**, 442-470 (1966).
- [7] J.D. Dixon; M.P.F. du Sautoy; A. Mann; D. Segal, *Analytic Pro- $p$  Groups (second edition)*. *Cambridge Studies in Advanced Mathematics, vol 61*. Cambridge University Press, Cambridge, (1999).

- [8] G. Dopinescu; B. Schraen, Endomorphism algebras of  $p$ -adic representations of  $p$ -adic Lie groups. arXiv:1106.2442.
- [9] A. Jones, Completed Group Algebras of Abelian-by-procyclic Groups. *Advances in Mathematics*. **354** (to appear in 2019).
- [10] M. Lazard, Groupes analytiques  $p$ -adiques. *Publ. Math. IHÉS* **26**, 389-603 (1965).
- [11] H. Li; F. Van Oystaeyen, *Zariskian Filtrations*. Kluwer Academic, Dordrecht, (1996).
- [12] J.C. McConnell; J.C. Robson, *Noncommutative Noetherian Rings*. Graduate Studies in Mathematics, vol 30. American Mathematical Society, Providence, (2001).
- [13] P. Schneider,  *$p$ -adic Lie Groups*. Grundlehren der Mathematischen Wissenschaften, vol. 344. Springer, Heidelberg, (2011).
- [14] P. Schneider; J. Teitelbaum, Algebras of  $p$ -adic distributions and admissible representations, *Invent. Math.* **153**, 145-196 (2003).