

KATUGAMPOLA FRACTIONAL CALCULUS WITH GENERALIZED k -WRIGHT FUNCTION

AHMAD Y. A. SALAMOONI, D. D. PAWAR

ABSTRACT. In this article, we presented some properties of the Katugampola fractional integrals and derivatives. Also we studied the fractional calculus properties involving Katugampola Fractional integrals and derivatives of generalized k -Wright function ${}_n\Phi_m^k(z)$.

AMS classifications: 33B15; 33C20; 26A33.

1. INTRODUCTION AND PRELIMINARIES

In recent years, researchers and authors have introduced a new fractional integrators operators and fractional derivatives operators which are generalizations of the famous Riemann-Levielle and the Hadamard-type, for more details see [1-5] and references therein.

Definition 1.[5] Let $\Omega = [a, b]$, the Katugampola fractional integrals ${}_\rho I_{0+}^\gamma \varphi$ and ${}_\rho I_{-\infty}^\gamma \varphi$ of order $\gamma \in \mathbb{C}(\Re(\gamma) > 0)$ are defined for $\rho > 0$, $a = 0$ and $b = \infty$ as

$$({}_\rho I_{0+}^\gamma \varphi)(s) = \frac{\rho^{1-\gamma}}{\Gamma(\gamma)} \int_0^s \frac{\tau^{\rho-1} \varphi(\tau)}{(s^\rho - \tau^\rho)^{1-\gamma}} d\tau, \quad (s > 0), \quad (1.1)$$

and

$$({}_\rho I_{-\infty}^\gamma \varphi)(s) = \frac{\rho^{1-\gamma}}{\Gamma(\gamma)} \int_s^\infty \frac{\tau^{\rho-1} \varphi(\tau)}{(\tau^\rho - s^\rho)^{1-\gamma}} d\tau, \quad (s > 0), \quad (1.2)$$

and the corresponding Katugampola fractional derivatives ${}_\rho D_{0+}^\gamma \varphi$ and ${}_\rho D_{-\infty}^\gamma \varphi$ are defined with $(n = 1 + [\Re(\gamma)])$ as

$$\begin{aligned} ({}_\rho D_{0+}^\gamma \varphi)(s) &:= (s^{1-\rho} \frac{d}{ds})^{1+[\Re(\gamma)]} ({}_\rho I_{0+}^{1-\gamma+[\Re(\gamma)]} \varphi)(s) \\ &= \frac{\rho^{\gamma-[\Re(\gamma)]}}{\Gamma(1-\gamma+[\Re(\gamma)])} (s^{1-\rho} \frac{d}{ds})^{1+[\Re(\gamma)]} \int_0^s \frac{\tau^{\rho-1} \varphi(\tau)}{(s^\rho - \tau^\rho)^{\gamma-[\Re(\gamma)]}} d\tau, \quad (s > 0), \end{aligned} \quad (1.3)$$

and

$$\begin{aligned} ({}_\rho D_{-\infty}^\gamma \varphi)(s) &:= (-s^{1-\rho} \frac{d}{ds})^{1+[\Re(\gamma)]} ({}_\rho I_{-\infty}^{1-\gamma+[\Re(\gamma)]} \varphi)(s) \\ &= \frac{\rho^{\gamma-[\Re(\gamma)]}}{\Gamma(1-\gamma+[\Re(\gamma)])} (-s^{1-\rho} \frac{d}{ds})^{1+[\Re(\gamma)]} \int_s^\infty \frac{\tau^{\rho-1} \varphi(\tau)}{(\tau^\rho - s^\rho)^{\gamma-[\Re(\gamma)]}} d\tau, \quad (s > 0). \end{aligned} \quad (1.4)$$

Key words and phrases. Katugampola Fractional integral and derivative, k -Gamma function and k -Wright Function.

Definition 2.[6] The generalized K -Gamma function $\Gamma_k(y)$ defined by

$$\Gamma_k(y) = \lim_{n \rightarrow \infty} \frac{n!k^n(nk)^{\frac{y}{k}-1}}{(y)_{n,k}}, \quad (k > 0; y \in \mathbb{C} \setminus k\mathbb{Z}^-), \quad (1.5)$$

where $(y)_{n,k}$ is the k -Pochhammer symbol given as

$$(y)_{n,k} := \begin{cases} \frac{\Gamma_k(y+nk)}{\Gamma_k(y)} & (k \in \mathbb{R}; y \in \mathbb{C} \setminus \{0\}) \\ y(y+k)(y+2k)\dots(y+(n-1)k) & (n \in \mathbb{N}^+; y \in \mathbb{C}) \end{cases} \quad (1.6)$$

and for $\Re(y) > 0$, the K -Gamma function $\Gamma_k(y)$ defined by the integral

$$\Gamma_k(y) = \int_0^\infty x^{y-1} e^{-\frac{x}{k}} dx \quad (1.7)$$

this given relation with Euler's Gamma function as

$$\Gamma_k(y) = k^{\frac{y}{k}-1} \Gamma\left(\frac{y}{k}\right). \quad (1.8)$$

Also [7],

$$\Gamma(1-y)\Gamma(y) = \frac{\pi}{\sin(y\pi)}. \quad (1.9)$$

Definition 3. [8] The Beta function $B(v, \omega)$ is defined as

$$\begin{aligned} B(v, \omega) &= \int_0^1 z^{v-1} (1-z)^{\omega-1} dz, \quad \Re(v) > 0, \quad \Re(\omega) > 0, \\ &= \frac{\Gamma(v)\Gamma(\omega)}{\Gamma(v+\omega)} \end{aligned} \quad (1.10)$$

Furthermore,

$$\begin{aligned} \int_{\hat{x}}^\infty (z - \hat{x})^{v-1} (z - \hat{y})^{\omega-1} dz &= (\hat{x} - \hat{y})^{v+\omega-1} B(v, 1-v-\omega), \\ \hat{x} > \hat{y}, \quad 0 < \Re(v) < 1 - \Re(\omega). \end{aligned} \quad (1.11)$$

Recently the Generalized K -Wright function introduced by (Gehlot and Prajapati [9]) which is defined as following:

Definition 4. For $k \in \mathbb{R}^+$; $z \in \mathbb{C}$; $p_i, q_j \in \mathbb{C}$, $\alpha_i, \beta_j \in \mathbb{R}$ ($\alpha_i, \beta_j \neq 0$; $i = 1, 2, \dots, n$; $j = 1, 2, \dots, m$) and $(p_i + \alpha_i r)$, $(q_j + \beta_j r) \in \mathbb{C} \setminus k\mathbb{Z}^-$, the generalized k -Wright function ${}_n\Phi_m^k$ is defined by

$${}_n\Phi_m^k(z) = {}_n\Phi_m^k \left[\begin{matrix} (p_i, \alpha_i)_{1,n} \\ (q_j, \beta_j)_{1,m} \end{matrix} \middle| z \right] = \sum_{r=0}^{\infty} \frac{\prod_{i=1}^n \Gamma_k(p_i + \alpha_i r)}{\prod_{j=1}^m \Gamma_k(q_j + \beta_j r)} \frac{z^r}{r!}, \quad (1.12)$$

with the convergence conditions describing as

$$\Delta = \sum_{j=1}^m \left(\frac{\beta_j}{k} \right) - \sum_{i=1}^n \left(\frac{\alpha_i}{k} \right); \mu = \prod_{i=1}^n \left| \frac{\alpha_i}{k} \right|^{-\frac{\alpha_i}{k}} \prod_{j=1}^m \left| \frac{\beta_j}{k} \right|^{\frac{\beta_j}{k}}; \nu = \sum_{j=1}^m \left(\frac{q_j}{k} \right) - \sum_{i=1}^n \left(\frac{p_i}{k} \right) + \frac{n-m}{2}$$

Lemma 1. [9] For $k \in \mathbb{R}^+$; $z \in \mathbb{C}$; $p_i, q_j \in \mathbb{C}$, $\alpha_i, \beta_j \in \mathbb{R}$ ($\alpha_i, \beta_j \neq 0$; $i = 1, 2, \dots, n$; $j = 1, 2, \dots, m$) and $(p_i + \alpha_i r)$, $(q_j + \beta_j r) \in \mathbb{C} \setminus k\mathbb{Z}^-$

- (1) If $\Delta > -1$, then series (1.12) is absolutely convergent for all $z \in \mathbb{C}$ and generalized k -Wright function ${}_n\Phi_m^k(z)$ is an entire function of z .
(2) If $\Delta = -1$, then series (1.12) is absolutely convergent for all $|z| < \mu$ and of

$$|z| = \mu, \Re(\mu) > \frac{1}{2}.$$

2. PROPERTIES OF KATUGAMPOLA FRACTIONAL INTEGRAL AND DERIVATIVE

In this section, we investigated some properties of the Katugampola fractional integrals and derivatives (1.1), (1.2) and (1.3), (1.4) for the power function $\varphi(s) = s^{\alpha-1}$ and the exponential function $e^{-\lambda s^\rho}$.

Lemma 2. Let $\rho > 0, \Re(\gamma) \geq 0$ and $n = 1 + [\Re(\gamma)]$

- (1) If $\Re(\alpha) > 0$, then

$$({}_\rho I_{0+}^\gamma \tau^{\alpha-1})(s) = \frac{\rho^{-\gamma} \Gamma(1 + \frac{\alpha-1}{\rho})}{\Gamma(1 + \frac{\alpha-1}{\rho} + \gamma)} s^{\rho\gamma + (\alpha-1)} \quad (\Re(\gamma) \geq 0; \Re(\alpha) > 0) \quad (2.1)$$

$$({}_\rho D_{0+}^\gamma \tau^{\alpha-1})(s) = \frac{\rho^{\gamma-n} \Gamma(1 + \frac{\alpha-1}{\rho})}{\Gamma(1 + \frac{\alpha-1}{\rho} - \gamma)} s^{(\alpha-1)-\rho\gamma} \quad (\Re(\gamma) \geq 0; \Re(\alpha) > 0). \quad (2.2)$$

- (2) If $\alpha \in \mathbb{C}$, then

$$({}_\rho I_{-}^\gamma \tau^{\alpha-1})(s) = \frac{\rho^{-\gamma} \Gamma(\frac{1-\alpha}{\rho} - \gamma)}{\Gamma(\frac{1-\alpha}{\rho})} s^{\rho\gamma + (\alpha-1)} \quad (\Re(\gamma) \geq 0; \Re(\gamma + \alpha) < 1) \quad (2.3)$$

$$({}_\rho D_{-}^\gamma \tau^{\alpha-1})(s) = \frac{\rho^{\gamma-n} \Gamma(\frac{1-\alpha}{\rho} + \gamma)}{\Gamma(\frac{1-\alpha}{\rho})} s^{(\alpha-1)-\rho\gamma} \quad (\Re(\gamma) \geq 0; \Re(\gamma + \alpha - [\Re(\gamma)]) < 1). \quad (2.4)$$

- (3) If $\Re(\lambda) > 0$, then

$$({}_\rho I_{-}^\gamma e^{-\lambda \tau^\rho})(s) = (\lambda \rho)^{-\gamma} e^{-\lambda s^\rho} \quad (\Re(\gamma) \geq 0) \quad (2.5)$$

$$({}_\rho D_{-}^\gamma e^{-\lambda \tau^\rho})(s) = (\lambda \rho)^\gamma e^{-\lambda s^\rho} \quad (\Re(\gamma) \geq 0). \quad (2.6)$$

Proof. To prove this Lemma, let the substitution $x = \frac{\tau^\rho}{s^\rho}$, in parts (1) and (2).

- (1) Firstly, by the equation (1.1) and the given substitution we have

$$\begin{aligned} ({}_\rho I_{0+}^\gamma \tau^{\alpha-1})(s) &= \frac{\rho^{-\gamma} s^{\rho\gamma + \alpha-1}}{\Gamma(\gamma)} \int_0^1 \frac{x^{\frac{\alpha-1}{\rho}}}{(1-x)^{1-\gamma}} dx \\ &= \frac{\rho^{-\gamma} s^{\rho\gamma + \alpha-1}}{\Gamma(\gamma)} B\left(\gamma, 1 + \frac{\alpha-1}{\rho}\right), \end{aligned}$$

now, using equation (1.10), we obtain the result (2.1). \square

Secondly, by the equation (1.3), the given substitution and by using the result (2.1), we have

$$\begin{aligned} ({}_\rho D_{0+}^\gamma \tau^{\alpha-1})(s) &= \left(s^{1-\rho} \frac{d}{ds}\right)^n ({}_\rho I_{0+}^{n-\gamma} \tau^{\alpha-1})(s) \\ &= \frac{\rho^{\gamma-n} \Gamma(1 + \frac{\alpha-1}{\rho})}{\Gamma(1 + \frac{\alpha-1}{\rho} + n - \gamma)} \left(s^{1-\rho} \frac{d}{ds}\right)^n s^{\rho(n-\gamma)+\alpha-1} \end{aligned}$$

$$= \frac{\rho^{\gamma-n}\Gamma(1+\frac{\alpha-1}{\rho})}{\Gamma(1+\frac{\alpha-1}{\rho}-\gamma)} s^{(\alpha-1)-\rho\gamma}$$

□

(2) Firstly, by the equation (1.2) and the given substitution we have

$$({}_\rho I_-^\gamma \tau^{\alpha-1})(s) = \frac{\rho^{-\gamma} s^{\rho\gamma+\alpha-1}}{\Gamma(\gamma)} \int_1^\infty x^{\frac{\alpha-1}{\rho}} (x-1)^{\gamma-1} dx,$$

now, using the equation (1.11), with $\hat{x} = 1$ and $\hat{y} = 0$, we obtain

$$({}_\rho I_-^\gamma \tau^{\alpha-1})(s) = \frac{\rho^{-\gamma} s^{\rho\gamma+\alpha-1}}{\Gamma(\gamma)} B\left(\gamma, 1-\gamma - (1 + \frac{\alpha-1}{\rho})\right),$$

by using equation (1.10), we obtain the result (2.3). □

Secondly, by the equation (1.4), the given substitution and by using the result (2.3), we have

$$\begin{aligned} ({}_\rho D_-^\gamma \tau^{\alpha-1})(s) &= \left(-s^{1-\rho} \frac{d}{ds}\right)^n ({}_\rho I_-^{n-\gamma} \tau^{\alpha-1})(s) \\ &= \frac{(-1)^n \rho^{\gamma-n} \Gamma(\frac{1-\alpha}{\rho} + \gamma - n)}{\Gamma(\frac{1-\alpha}{\rho})} \left(s^{1-\rho} \frac{d}{ds}\right)^n s^{\rho(n-\gamma)+\alpha-1} \\ &= \frac{(-1)^n \rho^{\gamma-n} \Gamma(\frac{1-\alpha}{\rho} + \gamma - n) \Gamma(1 - [\frac{1-\alpha}{\rho} + \gamma - n])}{\Gamma(\frac{1-\alpha}{\rho}) \Gamma(1 - [\gamma - \frac{\alpha-1}{\rho}])} \end{aligned} \quad (2.7)$$

Also, by using (1.9), we have

$$\Gamma(\frac{1-\alpha}{\rho} + \gamma - n) \Gamma(1 - [\frac{1-\alpha}{\rho} + \gamma - n]) = \frac{\pi}{\sin([\frac{1-\alpha}{\rho} + \gamma - n]\pi)} = \frac{(-1)^n \pi}{\sin([\gamma - \frac{\alpha-1}{\rho}]\pi)} \quad (2.8)$$

and

$$\frac{1}{\Gamma(1 - [\gamma - \frac{\alpha-1}{\rho}])} = \frac{\Gamma(\gamma - \frac{\alpha-1}{\rho})}{\Gamma(\gamma - \frac{\alpha-1}{\rho}) \Gamma(1 - [\gamma - \frac{\alpha-1}{\rho}])} = \frac{\Gamma(\gamma - \frac{\alpha-1}{\rho})}{\pi} \sin([\gamma - \frac{\alpha-1}{\rho}]\pi) \quad (2.9)$$

Substituting relations (2.8) and (2.9) in (2.7), we obtain (2.4). □

(3) For this part let the substitution $x = \tau^\rho - s^\rho$.

Firstly, by the equation (1.2) and the given substitution in this part we have

$$({}_\rho I_-^\gamma e^{-\lambda\tau^\rho})(s) = \frac{\rho^{-\gamma}}{\Gamma(\gamma)} e^{-\lambda s^\rho} \int_0^\infty e^{-\lambda x} x^{\gamma-1} dx,$$

then use the substitution $\vartheta = \lambda x$, we obtain

$$({}_\rho I_-^\gamma e^{-\lambda\tau^\rho})(s) = \frac{\rho^{-\gamma}}{\Gamma(\gamma)} e^{-\lambda s^\rho} \lambda^{-\gamma} \int_0^\infty e^{-\vartheta} \vartheta^{\gamma-1} d\vartheta,$$

since $\int_0^\infty e^{-\vartheta} \vartheta^{\gamma-1} d\vartheta = \Gamma(\gamma)$ [7], then the result is satisfy. □

Secondly, by the equation (1.4) and by using the result (2.5), we have

$$({}_\rho D_-^\gamma e^{-\lambda\tau^\rho})(s) = \left(-s^{1-\rho} \frac{d}{ds}\right)^n ({}_\rho I_-^{n-\gamma} e^{-\lambda\tau^\rho})(s)$$

$$\begin{aligned}
&= (-1)^n \left(s^{1-\rho} \frac{d}{ds} \right)^n ((\lambda\rho)^{\gamma-n} e^{-\lambda s^\rho}) \\
&= (-1)^n s^{(1-\rho)n} (\lambda\rho)^{\gamma-n} \left(\frac{d^n}{ds^n} e^{-\lambda s^\rho} \right) \\
&= (\lambda\rho)^\gamma e^{-\lambda s^\rho}
\end{aligned}$$

□

Remark 1. (a) In Lemma 2, if the power function is $\varphi(s) = \left(\frac{s^\rho}{\rho}\right)^{\alpha-1}$, then

(1) If $\Re(\alpha) > 0$, then

$$\left({}_\rho I_{0+}^\gamma \left(\frac{\tau^\rho}{\rho} \right)^{\alpha-1} \right)(s) = \frac{\Gamma(\alpha)}{\Gamma(\alpha + \gamma)} \left(\frac{s^\rho}{\rho} \right)^{\alpha+\gamma-1} \quad (\Re(\gamma) \geq 0; \Re(\alpha) > 0)$$

$$\left({}_\rho D_{0+}^\gamma \left(\frac{\tau^\rho}{\rho} \right)^{\alpha-1} \right)(s) = \frac{\Gamma(\alpha)}{\Gamma(\alpha - \gamma)} \left(\frac{s^\rho}{\rho} \right)^{\alpha-\gamma-1} \quad (\Re(\gamma) \geq 0; \Re(\alpha) > 0).$$

(2) If $\alpha \in \mathbb{C}$, then

$$\left({}_\rho I_-^\gamma \left(\frac{\tau^\rho}{\rho} \right)^{\alpha-1} \right)(s) = \frac{\Gamma(1 - \gamma - \alpha)}{\Gamma(1 - \alpha)} \left(\frac{s^\rho}{\rho} \right)^{\alpha+\gamma-1} \quad (\Re(\gamma) \geq 0; \Re(\gamma + \alpha) < 1)$$

$$\left({}_\rho D_-^\gamma \left(\frac{\tau^\rho}{\rho} \right)^{\alpha-1} \right)(s) = \frac{\Gamma(1 + \gamma - \alpha)}{\Gamma(1 - \alpha)} \left(\frac{s^\rho}{\rho} \right)^{\alpha-\gamma-1} \quad (\Re(\gamma) \geq 0; \Re(\gamma + \alpha - [\Re(\gamma)]) < 1).$$

(b) If $\Re(\alpha) > \Re(\gamma) > 0$, then

$$({}_\rho I_-^\gamma \tau^{-\alpha})(s) = \frac{\rho^{-\gamma} \Gamma(\frac{\alpha}{\rho} - \gamma)}{\Gamma(\frac{\alpha}{\rho})} s^{\rho\gamma-\alpha} \quad (2.10)$$

3. KATUGAMPOLA FRACTIONAL INTEGRATION FOR GENERALIZED k -WRIGHT FUNCTION

In this section, we established the Katugampola fractional integration for generalized k -Wright function (1.12).

Theorem 1. Let $\gamma, \alpha \in \mathbb{C}$ such that $\Re(\gamma) > 0$, $\Re(\alpha) > 0$; $\lambda \in \mathbb{C}$, $\rho > 0$, $\nu > 0$, then for $\Delta > -1$, the Katugampola fractional integration ${}_\rho I_{0+}^\gamma$ for generalized k -Wright function ${}_n\Phi_m^k(z)$ is given as

$$\begin{aligned}
&\left({}_\rho I_{0+}^\gamma \left(\tau^{\frac{\alpha}{k}-1} {}_n\Phi_m^k \left[\begin{matrix} (p_i, \alpha_i)_{1,n} \\ (q_j, \beta_j)_{1,m} \end{matrix} \middle| \lambda \tau^{\frac{\nu}{k}} \right] \right) \right)(s) \\
&= \left(\frac{k}{\rho} \right)^\gamma s^{\frac{\alpha}{k} + \rho\gamma - 1} {}_{n+1}\Phi_{m+1}^k \left[\begin{matrix} (p_i, \alpha_i)_{1,n}, (\frac{1}{\rho}(\alpha + (\rho - 1)k), \frac{\nu}{\rho}) \\ (q_j, \beta_j)_{1,m}, (\frac{1}{\rho}(\alpha + (\rho(\gamma + 1) - 1)k), \frac{\nu}{\rho}) \end{matrix} \middle| \lambda s^{\frac{\nu}{k}} \right]
\end{aligned} \tag{3.1}$$

Proof. According to Lemma 1, a generalized k -Wright functions in both side of the equation (3.1), exist for $s > 0$. We consider that

$$M \equiv \left({}_\rho I_{0+}^\gamma \left(\tau^{\frac{\alpha}{k}-1} {}_n\Phi_m^k \left[\begin{matrix} (p_i, \alpha_i)_{1,n} \\ (q_j, \beta_j)_{1,m} \end{matrix} \middle| \lambda \tau^{\frac{\nu}{k}} \right] \right) \right)(s)$$

using (1.12), we can write the above equation as

$$M \equiv \left({}_{\rho}I_{0+}^{\gamma} \left(\tau^{\frac{\alpha}{k}-1} \sum_{r=0}^{\infty} \frac{\prod_{i=1}^n \Gamma_k(p_i + \alpha_i r)}{\prod_{j=1}^m \Gamma_k(q_j + \beta_j r)} \frac{(\lambda \tau^{\frac{\nu}{k}})^r}{r!} \right) \right) (s)$$

now, using the integration of the series term- by term we obtain

$$M \equiv \sum_{r=0}^{\infty} \frac{\prod_{i=1}^n \Gamma_k(p_i + \alpha_i r)}{\prod_{j=1}^m \Gamma_k(q_j + \beta_j r)} \frac{(\lambda)^r}{r!} \left({}_{\rho}I_{0+}^{\gamma} \left(\tau^{\frac{\alpha}{k} + \frac{\nu r}{k} - 1} \right) \right) (s)$$

applying (2.1), the above equation reduces to,

$$M \equiv \sum_{r=0}^{\infty} \frac{\prod_{i=1}^n \Gamma_k(p_i + \alpha_i r)}{\prod_{j=1}^m \Gamma_k(q_j + \beta_j r)} \frac{(\lambda)^r}{r!} \frac{\rho^{-\gamma} \Gamma(1 + \frac{\frac{\alpha}{k} + \frac{\nu r}{k} - 1}{\rho})}{\Gamma(1 + \frac{\frac{\alpha}{k} + \frac{\nu r}{k} - 1}{\rho} + \gamma)} s^{\frac{\alpha + \nu r}{k} + \rho \gamma - 1}$$

using (1.8), we obtain

$$M \equiv \left(\frac{k}{\rho} \right)^{\gamma} s^{\frac{\alpha}{k} + \rho \gamma - 1} {}_{n+1}\Phi_m^k \left[\begin{matrix} (p_i, \alpha_i)_{1,n}, (\frac{1}{\rho}(\alpha + (\rho - 1)k), \frac{\nu}{\rho}) \\ (q_j, \beta_j)_{1,m}, (\frac{1}{\rho}(\alpha + (\rho(\gamma + 1) - 1)k), \frac{\nu}{\rho}) \end{matrix} \middle| \lambda s^{\frac{\nu}{k}} \right] \quad \square$$

Theorem 2. Let $\gamma, \alpha \in \mathbb{C}$ such that $\Re(\gamma) > 0, \Re(\alpha) > 0$; $\lambda \in \mathbb{C}, \rho > 0, \nu > 0$, then for $\Delta > -1$, the Katugampola fractional integration ${}_{\rho}I_{-}^{\gamma}$ for generalized k -Wright function ${}_n\Phi_m^k(z)$ is given as

$$\begin{aligned} & \left({}_{\rho}I_{-}^{\gamma} \left(\tau^{-\frac{\alpha}{k}} {}_n\Phi_m^k \left[\begin{matrix} (p_i, \alpha_i)_{1,n} \\ (q_j, \beta_j)_{1,m} \end{matrix} \middle| \lambda \tau^{-\frac{\nu}{k}} \right] \right) \right) (s) \\ &= \left(\frac{k}{\rho} \right)^{\gamma} s^{\rho \gamma - \frac{\alpha}{k}} {}_{n+1}\Phi_m^k \left[\begin{matrix} (p_i, \alpha_i)_{1,n}, (\frac{\alpha}{\rho} - k\gamma, \frac{\nu}{\rho}) \\ (q_j, \beta_j)_{1,m}, (\frac{\alpha}{\rho}, \frac{\nu}{\rho}) \end{matrix} \middle| \lambda s^{-\frac{\nu}{k}} \right] \end{aligned} \quad (3.2)$$

Proof. According to Lemma 1, a generalized k -Wright functions in both side of the equation (3.2), exist for $s > 0$. We consider that

$$N \equiv \left({}_{\rho}I_{-}^{\gamma} \left(\tau^{-\frac{\alpha}{k}} {}_n\Phi_m^k \left[\begin{matrix} (p_i, \alpha_i)_{1,n} \\ (q_j, \beta_j)_{1,m} \end{matrix} \middle| \lambda \tau^{-\frac{\nu}{k}} \right] \right) \right) (s)$$

using (1.12), we can write the above equation as

$$N \equiv \sum_{r=0}^{\infty} \frac{\prod_{i=1}^n \Gamma_k(p_i + \alpha_i r)}{\prod_{j=1}^m \Gamma_k(q_j + \beta_j r)} \frac{(\lambda)^r}{r!} \left({}_{\rho}I_{-}^{\gamma} \left(\tau^{-\frac{\alpha + \nu r}{k}} \right) \right) (s)$$

applying (2.10), the above equation reduces to,

$$N \equiv \sum_{r=0}^{\infty} \frac{\prod_{i=1}^n \Gamma_k(p_i + \alpha_i r)}{\prod_{j=1}^m \Gamma_k(q_j + \beta_j r)} \frac{(\lambda)^r}{r!} \frac{\rho^{-\gamma} \Gamma(\frac{\alpha + \nu r}{\rho} - \gamma)}{\Gamma(\frac{\alpha + \nu r}{\rho})} s^{\rho \gamma - \frac{\alpha + \nu r}{k}}$$

using (1.8), we obtain

$$N \equiv \left(\frac{k}{\rho} \right)^{\gamma} s^{\rho \gamma - \frac{\alpha}{k}} {}_{n+1}\Phi_m^k \left[\begin{matrix} (p_i, \alpha_i)_{1,n}, (\frac{\alpha}{\rho} - k\gamma, \frac{\nu}{\rho}) \\ (q_j, \beta_j)_{1,m}, (\frac{\alpha}{\rho}, \frac{\nu}{\rho}) \end{matrix} \middle| \lambda s^{-\frac{\nu}{k}} \right] \quad \square$$

4. KATUGAMPOLA FRACTIONAL DIFFERENTIATION FOR GENERALIZED k -WRIGHT FUNCTION

This section deals with the Katugampola fractional differentiation for generalized k -Wright function (1.12).

Theorem 3. Let $\gamma, \alpha \in \mathbb{C}$ such that $\Re(\gamma) > 0, \Re(\alpha) > 0$; $\lambda \in \mathbb{C}, \rho > 0, \nu > 0$, then for $\Delta > -1$, the Katugampola fractional integration ${}_0D_{0+}^{\gamma}$ for generalized k -Wright function ${}_n\Phi_m^k(z)$ is given as

$$\begin{aligned} & \left({}_0D_{0+}^{\gamma} \left(\tau^{\frac{\alpha}{k}-1} {}_n\Phi_m^k \left[\begin{matrix} (p_i, \alpha_i)_{1,n} \\ (q_j, \beta_j)_{1,m} \end{matrix} \middle| \lambda \tau^{\frac{\nu}{k}} \right] \right) \right) (s) \\ &= \left(\frac{k}{\rho} \right)^{-\gamma} s^{\frac{\alpha}{k}-\rho\gamma-1} {}_{n+1}\Phi_{m+1}^k \left[\begin{matrix} (p_i, \alpha_i)_{1,n}, (\frac{1}{\rho}(\alpha + (\rho-1)k), \frac{\nu}{\rho}) \\ (q_j, \beta_j)_{1,m}, (\frac{1}{\rho}(\alpha + (\rho(1-\gamma)-1)k), \frac{\nu}{\rho}) \end{matrix} \middle| \lambda s^{\frac{\nu}{k}} \right] \end{aligned} \quad (4.1)$$

Proof. According to Lemma 1, a generalized k -Wright functions in both side of the equation (4.1), exist for $s > 0$. Let $n = 1 + [\Re(\gamma)]$, then we consider that

$$P \equiv \left({}_0D_{0+}^{\gamma} \left(\tau^{\frac{\alpha}{k}-1} {}_n\Phi_m^k \left[\begin{matrix} (p_i, \alpha_i)_{1,n} \\ (q_j, \beta_j)_{1,m} \end{matrix} \middle| \lambda \tau^{\frac{\nu}{k}} \right] \right) \right) (s)$$

using (1.3), we have

$$P \equiv (s^{1-\rho} \frac{d}{ds})^n \left({}_0I_{0+}^{n-\gamma} \left(\tau^{\frac{\alpha}{k}-1} {}_n\Phi_m^k \left[\begin{matrix} (p_i, \alpha_i)_{1,n} \\ (q_j, \beta_j)_{1,m} \end{matrix} \middle| \lambda \tau^{\frac{\nu}{k}} \right] \right) \right) (s)$$

using Theorem 1, we obtain

$$P \equiv (s^{1-\rho} \frac{d}{ds})^n \left(\left(\frac{k}{\rho} \right)^{n-\gamma} s^{\frac{\alpha}{k}+\rho(n-\gamma)-1} {}_{n+1}\Phi_{m+1}^k \left[\begin{matrix} (p_i, \alpha_i)_{1,n}, (\frac{1}{\rho}(\alpha + (\rho-1)k), \frac{\nu}{\rho}) \\ (q_j, \beta_j)_{1,m}, (\frac{1}{\rho}(\alpha + (\rho(n-\gamma+1)-1)k), \frac{\nu}{\rho}) \end{matrix} \middle| \lambda s^{\frac{\nu}{k}} \right] \right)$$

using (1.12), we can write the above equation as

$$P \equiv \left(\frac{k}{\rho} \right)^{n-\gamma} \sum_{r=0}^{\infty} \frac{\prod_{i=1}^n \Gamma_k(p_i + \alpha_i r) \Gamma_k(\frac{1}{\rho}(\alpha + (\rho-1)k) + \frac{\nu}{\rho} r)}{\prod_{j=1}^m \Gamma_k(q_j + \beta_j r) \Gamma_k(\frac{1}{\rho}(\alpha + (\rho(n-\gamma+1)-1)k) + \frac{\nu}{\rho} r)} \frac{(\lambda)^r}{r!} (s^{1-\rho} \frac{d}{ds})^n (s^{\frac{\alpha}{k}+\frac{\nu}{k}+\rho(n-\gamma)-1})$$

we can write the above equation as

$$\begin{aligned} P \equiv k^{n-\gamma} \rho^{\gamma} \sum_{r=0}^{\infty} & \frac{\prod_{i=1}^n \Gamma_k(p_i + \alpha_i r) \Gamma_k(\frac{1}{\rho}(\alpha + (\rho-1)k) + \frac{\nu}{\rho} r)}{\prod_{j=1}^m \Gamma_k(q_j + \beta_j r) \Gamma_k(\frac{1}{\rho}(\alpha + (\rho(n-\gamma+1)-1)k) + \frac{\nu}{\rho} r)} \frac{(\lambda)^r}{r!} \\ & \times \frac{\Gamma(\frac{1}{\rho}(\frac{\alpha}{k} + \frac{\nu r}{k} + (n-\gamma)\rho + \rho - 1))}{\Gamma(\frac{1}{\rho}(\frac{\alpha}{k} + \frac{\nu r}{k} - \gamma\rho + \rho - 1))} s^{\frac{\alpha}{k}+\frac{\nu}{k}-\rho\gamma-1} \end{aligned}$$

using (1.8), we obtain

$$P \equiv \left(\frac{k}{\rho} \right)^{-\gamma} s^{\frac{\alpha}{k}-\rho\gamma-1} {}_{n+1}\Phi_{m+1}^k \left[\begin{matrix} (p_i, \alpha_i)_{1,n}, (\frac{1}{\rho}(\alpha + (\rho-1)k), \frac{\nu}{\rho}) \\ (q_j, \beta_j)_{1,m}, (\frac{1}{\rho}(\alpha + (\rho(1-\gamma)-1)k), \frac{\nu}{\rho}) \end{matrix} \middle| \lambda s^{\frac{\nu}{k}} \right]$$

□

Theorem 4. Let $\gamma, \alpha \in \mathbb{C}$ such that $\Re(\gamma) > 0, \Re(\alpha) > 1 + [\Re(\gamma)] - \Re(\gamma)$; $\lambda \in$

\mathbb{C} , $\rho > 0$, $\nu > 0$, then for $\Delta > -1$, the Katugampola fractional integration ${}_0D_-^\gamma$ for generalized k -Wright function ${}_n\Phi_m^k(z)$ is given as

$$\begin{aligned} & \left({}_0D_-^\gamma \left(\tau^{-\frac{\alpha}{k}} {}_n\Phi_m^k \left[\begin{matrix} (p_i, \alpha_i)_{1,n} \\ (q_j, \beta_j)_{1,m} \end{matrix} \middle| \lambda \tau^{-\frac{\nu}{k}} \right] \right) \right) (s) \\ &= \left(\frac{k}{\rho} \right)^{-\gamma} s^{-\rho\gamma-\frac{\alpha}{k}} {}_{n+1}\Phi_{m+1}^k \left[\begin{matrix} (p_i, \alpha_i)_{1,n}, (\frac{\alpha}{\rho} + k\gamma, \frac{\nu}{\rho}) \\ (q_j, \beta_j)_{1,m}, (\frac{\alpha}{\rho}, \frac{\nu}{\rho}) \end{matrix} \middle| \lambda s^{-\frac{\nu}{k}} \right] \quad (4.2) \end{aligned}$$

Proof. According to Lemma 1, a generalized k -Wright functions in both side of the equation (4.2), exist for $s > 0$. Let $n = 1 + [\Re(\gamma)]$, then we consider that

$$Q \equiv \left({}_0D_-^\gamma \left(\tau^{-\frac{\alpha}{k}} {}_n\Phi_m^k \left[\begin{matrix} (p_i, \alpha_i)_{1,n} \\ (q_j, \beta_j)_{1,m} \end{matrix} \middle| \lambda \tau^{-\frac{\nu}{k}} \right] \right) \right) (s)$$

using (1.4), we have

$$Q \equiv \left(-s^{1-\rho} \frac{d}{ds} \right)^n \left({}_0I_-^{n-\gamma} \left(\tau^{-\frac{\alpha}{k}} {}_n\Phi_m^k \left[\begin{matrix} (p_i, \alpha_i)_{1,n} \\ (q_j, \beta_j)_{1,m} \end{matrix} \middle| \lambda \tau^{-\frac{\nu}{k}} \right] \right) \right) (s)$$

using Theorem 2, we obtain

$$Q \equiv \left(-s^{1-\rho} \frac{d}{ds} \right)^n \left(\frac{k}{\rho} \right)^{n-\gamma} s^{\rho(n-\gamma)-\frac{\alpha}{k}} {}_{n+1}\Phi_{m+1}^k \left[\begin{matrix} (p_i, \alpha_i)_{1,n}, (\frac{\alpha}{\rho} - k(n-\gamma), \frac{\nu}{\rho}) \\ (q_j, \beta_j)_{1,m}, (\frac{\alpha}{\rho}, \frac{\nu}{\rho}) \end{matrix} \middle| \lambda s^{-\frac{\nu}{k}} \right]$$

using (1.12), we can write the above equation as

$$Q \equiv (-1)^n \left(\frac{k}{\rho} \right)^{n-\gamma} \sum_{r=0}^{\infty} \frac{\prod_{i=1}^n \Gamma_k(p_i + \alpha_i r) \Gamma_k(\frac{\alpha}{\rho} - (n-\gamma)k + \frac{\nu}{\rho} r)}{\prod_{j=1}^m \Gamma_k(q_j + \beta_j r) \Gamma_k(\frac{\alpha}{\rho} + \frac{\nu}{\rho} r)} \frac{(\lambda)^r}{r!} \left(s^{1-\rho} \frac{d}{ds} \right)^n \left(s^{\rho(n-\gamma)-\frac{\alpha}{k}-\frac{\nu}{k}} \right)$$

on simplifying the above equation, we obtain

$$\begin{aligned} Q \equiv & (-1)^n k^{n-\gamma} \rho^\gamma \sum_{r=0}^{\infty} \frac{\prod_{i=1}^n \Gamma_k(p_i + \alpha_i r) \Gamma_k(\frac{\alpha}{\rho} - (n-\gamma)k + \frac{\nu}{\rho} r)}{\prod_{j=1}^m \Gamma_k(q_j + \beta_j r) \Gamma_k(\frac{\alpha}{\rho} + \frac{\nu}{\rho} r)} \frac{(\lambda)^r}{r!} \\ & \times \frac{\Gamma(1 + (n-\gamma) - \frac{\alpha}{\rho k} - \frac{\nu}{\rho k} r)}{\Gamma(1 - \gamma - \frac{\alpha}{\rho k} - \frac{\nu}{\rho k} r)} \left(s^{-\rho\gamma-\frac{\alpha}{k}-\frac{\nu}{k}} \right) \end{aligned}$$

using (1.8), we obtain

$$\begin{aligned} Q \equiv & (-1)^n \rho^\gamma \sum_{r=0}^{\infty} \frac{\prod_{i=1}^n \Gamma_k(p_i + \alpha_i r)}{\prod_{j=1}^m \Gamma_k(q_j + \beta_j r) \Gamma(\frac{\alpha}{\rho k} + \frac{\nu}{\rho k} r)} \frac{(\lambda)^r}{r!} \\ & \times \frac{\Gamma(\gamma - n + \frac{\alpha}{\rho k} + \frac{\nu}{\rho k} r) \Gamma(1 - (\gamma - n + \frac{\alpha}{\rho k} + \frac{\nu}{\rho k} r))}{\Gamma(1 - (\gamma + \frac{\alpha}{\rho k} + \frac{\nu}{\rho k} r))} \left(s^{-\rho\gamma-\frac{\alpha}{k}-\frac{\nu}{k}} \right) \quad (4.3) \end{aligned}$$

using (1.9), we have

$$\begin{aligned} & \Gamma(\gamma - n + \frac{\alpha}{\rho k} + \frac{\nu}{\rho k} r) \Gamma(1 - (\gamma - n + \frac{\alpha}{\rho k} + \frac{\nu}{\rho k} r)) \\ &= \frac{\pi}{\sin[(\gamma + \frac{\alpha}{\rho k} + \frac{\nu}{\rho k} r)\pi - n\pi]} \\ &= \frac{\pi}{\sin[(\gamma + \frac{\alpha}{\rho k} + \frac{\nu}{\rho k} r)\pi] \cos(n\pi)} \end{aligned}$$

$$= \frac{(-1)^n \pi}{\sin[(\gamma + \frac{\alpha}{\rho k} + \frac{\nu}{\rho k} r) \pi]} \quad (4.4)$$

and

$$\frac{1}{\Gamma(1 - (\gamma + \frac{\alpha}{\rho k} + \frac{\nu}{\rho k} r))} = \frac{\Gamma(\gamma + \frac{\alpha}{\rho k} + \frac{\nu}{\rho k} r) \sin[(\gamma + \frac{\alpha}{\rho k} + \frac{\nu}{\rho k} r) \pi]}{\pi} \quad (4.5)$$

Substituting (4.4) and (4.5) in (4.3), and finally using (1.8), we obtain

$$Q \equiv \left(\frac{k}{\rho} \right)^{-\gamma} s^{-\rho\gamma - \frac{\alpha}{k}} {}_{n+1}\Phi_{m+1}^k \left[\begin{matrix} (p_i, \alpha_i)_{1,n}, (\frac{\alpha}{\rho} + k\gamma, \frac{\nu}{\rho}) \\ (q_j, \beta_j)_{1,m}, (\frac{\alpha}{\rho}, \frac{\nu}{\rho}) \end{matrix} \middle| \lambda s^{-\frac{\nu}{k}} \right]$$

Remark 2. If $\rho = 1$, then

Theorems 1, 2, 3 and 4, are reduced to Theorems 2, 3, 4 and 5, respectively (see[10]).

REFERENCES

- [1] UDITA KATUGAMPOLA, *New approach to a generalized fractional integral.* arXiv:1010.0742v2[math.CA] 11 Oct 2010.
- [2] UDITA N. KATUGAMPOLA, *A new approach to generalized fractional derivatives. (Communicated by claudio r. henriquez).* Bulletin of Mathematical Analysis and Applications. Vol. 6 Issue 4 (2014), P. 1-15.
- [3] UDITA N. KATUGAMPOLA, *Existence and uniqueness results for a class of generalized fractional differential equations.* arXiv:1411.5229v2[math.CA] 9 Jun 2016.
- [4] Ricardo Almeida, Agnieszka B. Malinowska and Tatiana Odzijewicz, *Fractional differential equations with dependence on the Caputo-Katugampola derivative.* arXiv:1607.0691v1[math.CA] 23 Jun 2016.
- [5] D. S. OLIVEIRA1 and E. CAPELAS DE OLIVEIRA1, *Hilfer-Katugampola fractional derivative.* arXiv:1705.07733v2[math.CA] 10 Nov 2017.
- [6] R. Diaz and E. Pariguan, *On hypergeometric functions and Pochhammer k-symbol.* Divulgaciones Mathematicas, 15 (2) (2007) 179-192.
- [7] A. A. Kilbas, H. M. Srivastava and J. J. Trujillo, *Theory and Applications of Fractional Differential Equations.* Elsevier, Amsterdam (2006).
- [8] S. G. Samko, A. A. Kilbas and O. I. Marichev, *Fractional Integrals and Derivatives, Theory and Applications.* Gordon and Breach, New York (1993).
- [9] K. S. Gehlot and J. C. Prajapati, *On generalization of K-Wright function and its properties.* Pacific Journal of Applied Mathematics. Vol. 5, No. 2 (2013).
- [10] K. S. Gehlot and J. C. Prajapati, *Fractional calculus of generalized K-Wright function.* Journal of Fractional Calculus and Applications, Vol. 4(2) July 2013, pp. 283-289.

AHMAD Y. A. SALAMOONI

SCHOOL OF MATHEMATICAL SCIENCES, SWAMI RAMANAND TEERTH MARATHWADA UNIVERSITY,
NANDED-431606, INDIA

E-mail address: ayoussss83@gmail.com

D. D. PAWAR

SCHOOL OF MATHEMATICAL SCIENCES, SWAMI RAMANAND TEERTH MARATHWADA UNIVERSITY,
NANDED-431606, INDIA

E-mail address: dypawar@yahoo.com