

FS-INDICATORS OF pq -DIMENSIONAL POINTED HOPF ALGEBRAS

SI CHEN, TIAN TIAN LIU, LIN HONG WANG, AND XING TING WANG

ABSTRACT. We compute higher Frobenius-Schur indicators of pq -dimensional pointed Hopf algebras in characteristic p through their associated graded Hopf algebras. These indicators are gauge invariants for the monoidal categories of representations of these algebras.

1. INTRODUCTION

Higher Frobenius-Schur (FS) indicators of Hopf algebras were introduced in [4] and were further studied in [2, 3, 7] and other research works. These indicators provide gauge invariants of the monoidal representation categories of these Hopf algebras, see [2, Theorem 2.2]. The values of FS-indicators of certain Hopf algebras have been computed in, e.g., [1, 2, 9]. In this short note, we concern the indicators of pointed Hopf algebras of dimension pq over a field of positive characteristic. The classification of such Hopf algebras was recently obtained in [8]. Let r be a prime. We define a character function $\chi_r(n)$, associated to the prime r , in positive integers n such that

$$\chi_r(n) = \begin{cases} r, & \text{if } r \mid n \\ 1, & \text{if } r \nmid n. \end{cases}$$

Then our main result states as follows.

1.1. Theorem. *Let $q \neq p$ be primes, H be a pq -dimensional pointed Hopf algebra over characteristic p . If H is commutative and cocommutative, then the n -th indicator*

$$\nu_n(H) \equiv \chi_p(n)\chi_q(n) \pmod{p}.$$

Otherwise,

$$\nu_n(H) \equiv \chi_p(n) \pmod{p}.$$

In this section, we also set up notation and terminology. Readers unfamiliar with Hopf algebras are referred to standard textbooks such as [5, 6]. The proof of the main result is in section 2. Throughout, H is a finite-dimensional Hopf algebra over a base field \mathbf{k} . We use the standard notation $(H, m, u, \Delta, \varepsilon, S)$, where $m : H \otimes H \rightarrow H$ is the multiplication map, $u : \mathbf{k} \rightarrow H$ is the unit map, $\Delta : H \rightarrow H \otimes H$ is

2010 *Mathematics Subject Classification.* 16T05.

Key words and phrases. pointed Hopf algebras; FS-indicators; positive characteristic.

the comultiplication map, $\varepsilon : H \rightarrow \mathbf{k}$ is the counit map, and $S : H \rightarrow H$ is the antipode. The vector space dual of H is also a Hopf algebra and will be denoted by H^* . We use the Sweedler notation $\Delta(h) = \sum h_{(1)} \otimes h_{(2)}$. If $f, g \in H^*$, then $fg(h) = \sum f(h_{(1)})g(h_{(2)})$ for any $h \in H$ and $\varepsilon_{H^*}(f) = f(1)$.

Let n be a positive integer. Suppose $h_1, \dots, h_n \in H$. Then the n -th power of multiplication is defined as $m^{[n]}(h_1 \otimes \dots \otimes h_n) = h_1 \cdots h_n$. Let $h \in H$. The n -th power of comultiplication is defined to be

$$\Delta^{[n]}(h) = \begin{cases} h & n = 1 \\ (\Delta^{[n-1]} \otimes \text{id})(\Delta(h)) & n \geq 2 \end{cases}$$

The n -th Sweedler power of h is defined to be

$$P_n(h) = h^{[n]} = \begin{cases} \varepsilon(h)1_H & n = 0 \\ m^{[n]} \circ \Delta^{[n]}(h) & n \geq 1 \end{cases}$$

Then the n -th indicator [2, Definition 2.1] of H is given by

$$\nu_n(H) = \text{Tr}(S \circ P_{n-1}).$$

In particular, $\nu_1(H) = 1$ and $\nu_2(H) = \text{Tr}(S)$.

Acknowledgement. We began this work in an undergraduate research project at the University of Pittsburgh. We would like to express our gratitude to the math department for its supporting.

2. THE PROOF OF THE MAIN RESULT

In this section, $q \neq p$ are primes, the base field \mathbf{k} is algebraically closed of characteristic p , and ξ is a q th primitive root of unity.

2.1. Theorem. [8, Theorem 2.19] *Let $\delta = 0$ or 1 . Then a pq -dimensional pointed Hopf algebra H is isomorphic to one of the following*

- 1) $\mathbf{k}\langle g, x \rangle / (g^q - 1, x^p - \delta x)$,
 $\Delta(g) = g \otimes g, \Delta(x) = x \otimes 1 + 1 \otimes x$;
- 2) $\begin{cases} \mathbf{k}\langle g, x \rangle / (g^q - 1, x^p, xg - \xi gx), & q \nmid p - 1, \\ \mathbf{k}\langle g, x \rangle / (g^q - 1, x^p - \delta x, gx - \xi xg), & q \mid p - 1, \end{cases}$
 $\Delta(g) = g \otimes g, \Delta(x) = x \otimes 1 + 1 \otimes x$;
- 3) $\mathbf{k}\langle g, x \rangle / (g^q - 1, x^p - \delta(1 - g^p))$,
 $\Delta(g) = g \otimes g, \Delta(x) = x \otimes 1 + g \otimes x$;
- 4) $\begin{cases} \mathbf{k}\langle g, x \rangle / (g^q - 1, x^p - x, gx - xg - g + g^2), & q \nmid p - 1, \\ \mathbf{k}\langle g, x \rangle / (g^q - 1, x^p - x), & q \mid p - 1, \end{cases}$
 $\Delta(g) = g \otimes g, \Delta(x) = x \otimes 1 + g \otimes x$.

The approach of our proof is based on the above classification result and the following theorem and proposition. We also provide a lemma prior to the proof of Theorem 1.1. A *left integral* in a finite dimensional Hopf algebra H is an element $\Lambda \in H$ such that $h\Lambda = \varepsilon(h)\Lambda$, for all $h \in H$. The span of the left integrals in H is a one-dimensional space.

2.2. Theorem. [2, Corollary 2.6] *Suppose $\lambda \in H^*$ and $\Lambda \in H$ are both left integrals such that $\lambda(\Lambda) = 1$. Then $\nu_n(H) = \lambda(\Lambda^{[n]})$ for all positive integers n .*

2.3. Proposition. [7, Corollary 3.17, Corollary 4.6] *Let H and H' be finite-dimensional Hopf algebras. Then, for all positive integers n , we have the following:*

- 1) $\nu_n(H^*) = \nu_n(H)$ and $\nu_n(H \otimes H') = \nu_n(H) \cdot \nu_n(H')$.
- 2) If H is filtered, then $\nu_n(\text{gr}H) = \nu_n(H)$.
- 3) If $\text{char } \mathbf{k} > 0$, then $\{\nu_n(H)\}$ is periodic.

Although Proposition 2.3, 2) is true for any Hopf filtration, in this paper we use coradical filtration.

2.4. Lemma. *Let i, n be integers for $0 \leq i \leq q - 1$ and $n \geq 1$. Suppose nonnegative integers k_1, \dots, k_{n+1} form a partition of $p - 1$.*

- 1) If $xg = gx$, $\Delta(g) = g \otimes g$, and $\Delta(x) = x \otimes 1 + g \otimes x$, then

$$(g^i x^{p-1})^{[n+1]} = \sum_{0 \leq k_1, \dots, k_n \leq p-1} \binom{p-1}{k_1, \dots, k_{n+1}} g^{(n+1)i + nk_1 + (n-1)k_2 + \dots + k_n} x^{p-1}.$$

- 2) If $xg = \xi gx$, $\Delta(g) = g \otimes g$, and $\Delta(x) = x \otimes 1 + 1 \otimes x$, then

$$(g^i x^{p-1})^{[n+1]} = \left((\xi^i)^n + \dots + \xi^i + 1 \right)^{p-1} g^{(n+1)i} x^{p-1}.$$

Proof. 1) If $n = 1$, then $(g^i x^{p-1})^{[n+1]} = (g^i x^{p-1})^{[2]} = m \circ \Delta(g^i x^{p-1})$. Firstly,

$$\Delta(g^i x^{p-1}) = g^i \otimes g^i (x \otimes 1 + g \otimes x)^{p-1} = \sum_{k_1=0}^{p-1} \binom{p-1}{k_1} g^i x^{p-1-k_1} g^{k_1} \otimes g^i x^{k_1}.$$

Note that $xg = gx$. Then

$$(g^i x^{p-1})^{[2]} = \sum_{k_1=0}^{p-1} \binom{p-1}{k_1} g^{2i+k_1} x^{p-1}.$$

Now let $n = 2$, and k_1, k_2, k_3 form a partition of $p - 1$. Then

$$\Delta^{[3]}(g^i x^{p-1}) = \Delta \otimes \text{id} \circ \Delta(g^i x^{p-1}) = \sum_{k_1=0}^{p-1} \binom{p-1}{k_1} \Delta(g^i x^{p-1-k_1}) \Delta(g)^{k_1} \otimes \text{id}(g^i x^{k_1}).$$

Following the case when $n = 1$, we have

$$\Delta(g^i x^{p-1-k_1}) = \sum_{k_2=0}^{p-1-k_1} \binom{p-1-k_1}{k_2} g^i x^{k_3} g^{k_2} \otimes g^i x^{k_1}.$$

Hence,

$$\Delta^{[3]}(g^i x^{p-1}) = \sum_{k_1=0}^{p-1} \sum_{k_2=0}^{p-1-k_1} \binom{p-1}{k_1} \binom{p-1-k_1}{k_2} g^i x^{k_3} g^{k_1+k_2} \otimes g^i x^{k_2} g^{k_1} \otimes g^i x^{k_1}.$$

Thus, using multinomial coefficients, we have

$$\begin{aligned} (g^i x^{p-1})^{[3]} &= \sum_{0 \leq k_1, k_2 \leq p-1} \binom{p-1}{k_1, k_2, k_3} g^{3i+k_1+k_2+k_1} x^{p-1} \\ &= \sum_{0 \leq k_1, k_2 \leq p-1} \binom{p-1}{k_1, k_2, k_3} g^{3i+k_1+2k_2} x^{p-1}. \end{aligned}$$

Finally, one can show inductively that

$$(g^i x^{p-1})^{[n+1]} = \sum_{0 \leq k_1, \dots, k_n \leq p-1} \binom{p-1}{k_1, \dots, k_{n+1}} g^{(n+1)i+k_1+2k_2+\dots+nk_n} x^{p-1}.$$

2) It can be shown that

$$\begin{aligned} \Delta^{[n+1]}(x^{p-1}) &= (\Delta^{[n+1]}(x))^{p-1} = (x \otimes 1 \otimes \dots \otimes 1 + \dots + 1 \otimes \dots \otimes 1 \otimes x)^{p-1} \\ &= \sum_{0 \leq k_1, \dots, k_n \leq p-1} \binom{p-1}{k_1, \dots, k_{n+1}} x^{k_1} \otimes x^{k_2} \dots \otimes x^{k_{n+1}}. \end{aligned}$$

Note that we have $xg = \xi gx$. Hence,

$$\begin{aligned} (g^i x^{p-1})^{[n+1]} &= \sum_{0 \leq k_1, \dots, k_n \leq p-1} \binom{p-1}{k_1, \dots, k_{n+1}} g^i x^{k_1} g^i x^{k_2} \dots g^i x^{k_{n+1}} \\ &= \sum_{0 \leq k_1, \dots, k_n \leq p-1} \binom{p-1}{k_1, \dots, k_{n+1}} (\xi^i)^{nk_1} (\xi^i)^{(n-1)k_2} \dots (\xi^i)^{k_n} \cdot g^{(n+1)i} x^{p-1} \\ &= \left((\xi^i)^n + \dots + \xi^i + 1 \right)^{p-1} g^{(n+1)i} x^{p-1}. \end{aligned}$$

□

The proof of Theorem 1.1. Let H be a pq -dimensional pointed Hopf algebra H over \mathbf{k} given in Theorem 2.1. The linear basis of H can be chosen as $\{g^i x^j \mid 0 \leq i \leq p-1, 0 \leq j \leq q-1\}$. The associated graded Hopf algebras $\text{gr}H$ is one of the following, in the correspondence 1) $\rightsquigarrow A$, 3), 4) $\rightsquigarrow B$, and 2) $\rightsquigarrow C$.

$$A = \mathbf{k}[g, x]/(g^q - 1, x^p), \text{ with } \Delta(g) = g \otimes g, \Delta(x) = x \otimes 1 + 1 \otimes x.$$

$$B = \mathbf{k}[g, x]/(g^q - 1, x^p), \text{ with } \Delta(g) = g \otimes g, \Delta(x) = x \otimes 1 + g \otimes x.$$

$$C = \mathbf{k}\langle g, x \rangle/(g^q - 1, x^p, xg - \xi gx), \text{ with } \Delta(g) = g \otimes g, \Delta(x) = x \otimes 1 + 1 \otimes x.$$

Next we find case by case the indicators of these associated graded Hopf algebras.

Case A. Note that $A = \mathbf{k}[g]/(g^q - 1) \otimes \mathbf{k}[x]/(x^p)$ with $\Delta(g) = g \otimes g$, and $\Delta(x) = x \otimes 1 + 1 \otimes x$. Following from the counting argument

$$\nu_n(\mathbf{k}[g]/(g^q - 1)) = \#\{h \in \langle g \rangle \mid h^n = 1_{\langle g \rangle}\},$$

we have

$$\nu_n(\mathbf{k}[g]/(g^q - 1)) = \chi_q(n), \quad \text{for all } n \geq 1.$$

By [9, Corollary 3.9], it follows that

$$\nu_n(\mathbf{k}[x]/(x^p)) \equiv \chi_p(n) \pmod{p}, \quad \text{for all } n \geq 1.$$

Therefore

$$\nu_n(A) \equiv \chi_p(n)\chi_q(n) \pmod{p}, \quad \text{for all } n \geq 1.$$

Case B. The comultiplication on the basis elements is given by

$$\Delta(g^i x^j) = \sum_{k=0}^j \binom{j}{k} g^{k+i} x^{j-k} \otimes g^i x^k.$$

Let $\lambda_B = \delta_{g^{1-p}x^{p-1}} \in B^*$. Then

$$\delta_{g^{1-p}x^{p-1}}(g^i x^j) = \begin{cases} 1 & \text{if } j = p - 1 \text{ and } i \equiv 1 - p \pmod{q} \\ 0 & \text{otherwise.} \end{cases}$$

On the other hand,

$$\begin{aligned} & \sum_{k=0}^j \binom{j}{k} g^{k+i} x^{j-k} \cdot \delta_{g^{1-p}x^{p-1}}(g^i x^k) \\ &= \begin{cases} g^{p-1+1-p}x^0 = 1 & \text{if } j = k = p - 1 \text{ and } i \equiv 1 - p \pmod{q} \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

Therefore, by [1, Lemma 2.3], $\lambda_B = \delta_{g^{1-p}x^{p-1}}$ is a left integral of B^* . Next consider the element $\Lambda_B = (1 + g + \cdots + g^{q-1})x^{p-1} \in B$. It is clear that $x\Lambda_B = 0 = \varepsilon(x)\Lambda_B$ and $g\Lambda_B = \Lambda_B = \varepsilon(g)\Lambda_B$. Hence $\Lambda_B = (1 + g + \cdots + g^{q-1})x^{p-1}$ is a left integral of B . It is clear that $\lambda_B(\Lambda_B) = 1$. Following Theorem 2.2, for $n \geq 1$ we compute

$$\nu_{n+1}(B) = \lambda_B\left(\Lambda_B^{[n+1]}\right) = \delta_{g^{1-p}x^{p-1}}\left(\sum_{i=0}^{q-1} (g^i x^{p-1})^{[n+1]}\right).$$

Note that $gx = xg$ and $\delta(x) = x \otimes 1 + g \otimes x$. By Lemma 2.4,

$$\nu_{n+1}(B) = \sum_{\substack{0 \leq i \leq q-1 \\ 0 \leq k_1, \dots, k_n \leq p-1}} \binom{p-1}{k_1, \dots, k_{n+1}} \delta_{g^{1-p}}(g^{(n+1)i+k_1+2k_2+\dots+nk_n}).$$

For any fixed partition k_1, \dots, k_n, k_{n+1} of $p-1$ we have

$$\sum_{i=0}^{q-1} \delta_{g^{1-p}}(g^{(n+1)i+k_1+2k_2+\dots+nk_n}) = \begin{cases} 1 & \text{if } q \nmid n+1 \\ q \delta_{g^{1-p}}(g^{k_1+2k_2+\dots+nk_n}) & \text{if } q \mid n+1 \end{cases}.$$

Hence, for $q \nmid n+1$, we have

$$\nu_{n+1}(B) = \sum_{\substack{0 \leq i \leq q-1 \\ 0 \leq k_1, \dots, k_n \leq p-1}} \binom{p-1}{k_1, \dots, k_{n+1}} = (n+1)^{p-1}.$$

For $q \mid n+1$, we have

$$\nu_{n+1}(B) = \sum_{\substack{k_1+2k_2+\dots+nk_n \equiv 1-p \pmod{q} \\ 0 \leq k_1, \dots, k_n \leq p-1}} q \binom{p-1}{k_1, \dots, k_{n+1}}.$$

Case C. Since $\Delta(x) = x \otimes 1 + 1 \otimes x$, we have

$$\Delta(g^i x^j) = \sum_{k=0}^j \binom{j}{k} g^i x^{j-k} \otimes g^i x^k,$$

for the basis elements $g^i x^j$ for $0 \leq i \leq q-1$ and $0 \leq j \leq p-1$. By a similar argument as that for B , the element $\lambda_C = \delta_{x^{p-1}} \in C^*$ is a left integral of C^* . Next consider the element $\Lambda_C = (1 + g + \dots + g^{q-1})x^{p-1} \in C$. It is clear that $g\Lambda_C = \varepsilon(g)\Lambda_C$. Note that $xg = \xi gx$. Then

$$x\Lambda_C = x(1 + g + \dots + g^{q-1})x^{p-1} = (1 + \xi g + \dots + (\xi g)^{q-1})x^p = 0 = \varepsilon(x)\Lambda_C.$$

Hence $\Lambda_C = (1 + g + \dots + g^{q-1})x^{p-1}$ is a left integral of C . Again, by Theorem 2.2,

$$\begin{aligned} \nu_{n+1}(C) &= \delta_{x^{p-1}} \left(\sum_{i=0}^{q-1} (g^i x^{p-1})^{[n+1]} \right) = \delta_{x^{p-1}} \left(\sum_{i=0}^{q-1} ((\xi^i)^n + \dots + \xi^i + 1)^{p-1} g^{(n+1)i} \right) \\ &= \delta_{x^{p-1}} \left((n+1)^{p-1} x^{p-1} + \sum_{i=1}^{q-1} \frac{(\xi^i)^{n+1} - 1}{\xi^i - 1} g^{(n+1)i} x^{p-1} \right) = (n+1)^{p-1}. \end{aligned}$$

It follows from the Little Fermat's Theorem that $(n)^{p-1} \equiv \chi_p(n) \pmod{p}$. Hence we have $\nu_n(C) = \chi_p(n)$ for any $n \geq 1$.

Note that A is self-dual, B and C are dual Hopf algebras. Therefore, combining Proposition 2.3 and the three cases, we prove Theorem 1.1 and the following corollary.

2.5. **Corollary.** *The following combinatorics identity holds.*

$$\sum_{\substack{k_1+2k_2+\dots+(n-1)k_{n-1}\equiv 1-p \\ (\text{mod } q), \text{ for } 0\leq k_1,\dots,k_{n-1}\leq p-1}} q \binom{p-1}{k_1, \dots, k_n} = n^{p-1} \pmod{p}.$$

where p, q are primes, $n > 1$ is an integer, and $q \mid n$.

REFERENCES

- [1] H. Hu, X. Hu, L. Wang, and X. Wang, Computing indicators of Radford algebras, *Involve, a journal of mathematics*, 11 (2018), 325–334.
- [2] Y. Kashina, S. Montgomery, and S.-H. Ng, On the trace of the antipode and higher indicators, *Israel J. Math.*, 188(1):57–89, 2012.
- [3] Y. Kashina, Y. Sommerhäuser and Y. Zhu, On higher Frobenius-Schur indicators, *Memories of the American Mathematical Society*, 181 (2006), no. 855, viii+65 pp.
- [4] V. Linchenko and S. Montgomery, A Frobenius-Schur theorem for Hopf algebras, *Algebr. Represent. Theory*, 3(4):347–355, 2000.
- [5] S. Montgomery, *Hopf Algebras and Their Actions on Rings*, volume 82 of *CBMS Regional Conference Series in Mathematics*, published for the Conference Board of the Mathematical Sciences, Washington DC; by the American Mathematical Society, Providence, RI, 1993.
- [6] M. E. Sweedler, *Hopf algebras*, W. A. Benjamin, Inc., New York, 1969, Mathematics Lecture Note Series.
- [7] K. Shimizu, On indicators of Hopf algebras, *Israel J. Math.*, 207(1):155–201, 2015.
- [8] R. Xiong, Pointed p^2q -dimensional Hopf algebras in positive characteristic, preprint, arXiv1705.00339.
- [9] L. Wang and X. Wang, Indicators of Hopf algebras in positive characteristic, submitted preprint, arXiv1704.05382

E-mail address: sic26@pitt.edu, til43@pitt.edu, lhwang@pitt.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF PITTSBURGH, PITTSBURGH, PA 15260

E-mail address: xingting.wang@howard.edu

DEPARTMENT OF MATHEMATICS, HOWARD UNIVERSITY, WASHINGTON, DC 20059