

# Semigroups and Evolutionary Equations\*

Sascha Trostorff<sup>†</sup>

**Abstract.** We show how strongly continuous semigroups can be associated with evolutionary equations. For doing so, we need to define the space of admissible history functions and initial states. Moreover, the initial value problem has to be formulated within the framework of evolutionary equations, which is done by using the theory of extrapolation spaces. The results are applied to two examples. First, differential-algebraic equations in infinite dimensions are treated and it is shown, how a  $C_0$ -semigroup can be associated with such problems. In the second example we treat a concrete hyperbolic delay equation.

**Keywords:** Evolutionary equations,  $C_0$ -semigroups, admissible history and initial value

**2010 MSC:** 46N20, 47D06, 35F16

## 1 Introduction

In this article we bring together two theories for dealing with partial differential equations: the theory of  $C_0$ -semigroups on the one hand and the theory of evolutionary equations on the other hand. In particular, we show how  $C_0$ -semigroups can be associated with a given evolutionary equation.

The framework of evolutionary equations was introduced in the seminal paper [12]. Evolutionary equations are equations of the form

$$(\partial_t M(\partial_t) + A)U = F, \tag{1}$$

where  $\partial_t$  denotes the temporal derivative,  $M(\partial_t)$  is a bounded operator in space-time defined via a functional calculus for  $\partial_t$  and  $A$  is an, in general, unbounded spatial operator. The function  $F$  defined on  $\mathbb{R}$  and taking values in some Hilbert space is a given source term and one seeks for a solution  $U$  of the above equation. Here, the notion of solution is quite weak, since one just requires that the solution should belong to some exponentially weighted  $L_2$ -space. Thus, all operators have to be introduced in these spaces. Especially, the time derivative is introduced as an unbounded normal operator

---

\*Based on parts of the authors Habilitation thesis [23].

<sup>†</sup>Mathematisches Seminar, CAU Kiel, Germany, email: trostorff@math.uni-kiel.de

on such a space and so, in order to solve (1), one has to deal with the sum of two unbounded operators ( $\partial_t$  and  $A$ ). Problems of the form (1) cover a broad spectrum of different types of differential equations, such as hyperbolic, parabolic, elliptic and mixed-type problems, integro-differential equations [22], delay equations [8] and fractional differential equations [14]. Also, generalisations to some nonlinear [18, 19] and non-autonomous problems [15, 28, 29, 24] are possible. The solution theory is quite easy and just relies on pure Hilbert space theory.

On the other hand, there is the well-established theory of  $C_0$ -semigroups dealing with so-called Cauchy problems (see e.g. [7, 11, 5]). These are abstract equations of the form

$$\begin{aligned}(\partial_t + A)U &= F, \\ U(0) &= U_0,\end{aligned}\tag{2}$$

where  $A$  is a suitable operator acting on some Banach space. Although, (2) just seems to be a special case of (1) for  $M(\partial_t) = 1$ , the theories are quite different. While we focus on solutions lying in  $L_2$  in the theory of evolutionary equations, one seeks for continuous solutions in the framework of  $C_0$ -semigroups. Moreover, while (1) holds on  $\mathbb{R}$  as time horizon, (2) just holds on  $\mathbb{R}_{\geq 0}$  and is completed by an initial condition. Thus, in order to associate a  $C_0$ -semigroup with equations of the form (1) one has to find a way to formulate initial value problems and then derive assumptions, which would yield the additional regularity for the solutions (namely continuity with respect to time). This is the purpose of this work.

As we have indicated above, equations of the form (1) also cover delay equations, where it is more natural to prescribe histories instead of an initial state at time 0. Moreover, (1) also covers so-called differential algebraic equations (see [9] for the finite-dimensional case and [27, 26, 25] for infinite dimensions), where not every element of the underlying state space can be used as an initial state. Thus, one is confronted with the problem of defining the ‘right’ initial values and histories for (1) depending on the operators involved. Moreover, one has to incorporate these initial conditions within the framework of evolutionary equations, that is, initial conditions should enter the equation as a suitable source term on the right-hand side. This can be done by using extrapolation spaces and by extending the solution theory to those. Then it will turn out that initial conditions can be formulated by distributional right hand sides, which belong to a suitable extrapolation space associated with the time derivative operator  $\partial_t$ . Having the right formulation of initial value problems at hand, one can associate a  $C_0$ -semigroup on a product space consisting of the current state in the first and the past of the unknown in the second component. This idea was already used to deal with delay equations within the theory of  $C_0$ -semigroups, see [3]. As it turns out, this product space is not closed (as a subspace of a suitable Hilbert space) and in order to extend the associated  $C_0$ -semigroup to its closure one needs to impose similar conditions as in the Hille-Yosida Theorem. The key result, which will be used to extend the semigroup is the theorem of Widder-Arendt (see [1] or Theorem 6.4 below).

The paper is structured as follows: We begin by recalling the basic notions and well-posedness results for evolutionary problems (Section 2) and for extrapolation spaces

(Section 3). Then, in order to formulate initial value problems within the framework of evolutionary equations, we introduce a cut-off operator as an unbounded operator on the extrapolation space associated with the time derivative and discuss some of its properties (Section 4). Section 5 is then devoted to determine the ‘right’ space of admissible histories and initial values for a given evolutionary problem. We note here that we restrict ourselves to homogeneous problems in the sense that we do not involve an additional source term besides the given history. The main reason for that is that such source terms would restrict and change the set of admissible histories, a fact which is well-known in the theory of differential-algebraic equations. In Section 6 we associate a  $C_0$ -semigroup on the before introduced product space of admissible initial values and histories and prove the main result of this article (Theorem 6.7). In the last section we discuss two examples. First, we apply the results to abstract differential algebraic equations and thereby re-prove the Theorem of Hille-Yosida as a special case. In the second example, we discuss a concrete hyperbolic delay equation and prove that we can associate a  $C_0$ -semigroup with this problem.

Throughout, every Hilbert space is assumed to be complex and the inner product  $\langle \cdot, \cdot \rangle$  is conjugate-linear in the first and linear in the second argument.

## 2 Evolutionary Problems

We recall the basic notions and results for evolutionary problems, as they were introduced in [12] (see also [13, Chapter 6]). We begin by the definition of the time derivative operator on an exponentially weighted  $L_2$ -space (see also [16]).

**Definition.** Let  $\rho \in \mathbb{R}$  and  $H$  a Hilbert space. We set

$$L_{2,\rho}(\mathbb{R}; H) := \{f : \mathbb{R} \rightarrow H ; f \text{ measurable, } \int_{\mathbb{R}} \|f(t)\|^2 e^{-2\rho t} dt\}$$

with the common identification of functions coinciding almost everywhere. Then  $L_{2,\rho}(\mathbb{R}; H)$  is a Hilbert space with respect to the inner product

$$\langle f, g \rangle_\rho := \int_{\mathbb{R}} \langle f(t), g(t) \rangle e^{-2\rho t} dt \quad (f, g \in L_{2,\rho}(\mathbb{R}; H)).$$

Moreover, we define the operator

$$\partial_{t,\rho} : H_\rho^1(\mathbb{R}; H) \subseteq L_{2,\rho}(\mathbb{R}; H) \rightarrow L_{2,\rho}(\mathbb{R}; H), \quad f \mapsto f'$$

where

$$H_\rho^1(\mathbb{R}; H) := \{f \in L_{2,\rho}(\mathbb{R}; H) ; f' \in L_{2,\rho}(\mathbb{R}; H)\}$$

with  $f'$  denoting the usual distributional derivative.

We recall some facts on the operator  $\partial_{t,\rho}$  and refer to [8] for the respective proofs.

**Proposition 2.1.** *Let  $\rho \in \mathbb{R}$  and  $H$  a Hilbert space.*

(a) The operator  $\partial_{t,\rho}$  is densely defined, closed and linear and  $C_c^\infty(\mathbb{R}; H)$  is a core for  $\partial_{t,\rho}$ .

(b) The spectrum of  $\partial_{t,\rho}$  is given by

$$\sigma(\partial_{t,\rho}) = \{it + \rho; t \in \mathbb{R}\}.$$

(c) For  $\rho \neq 0$  the operator  $\partial_{t,\rho}$  is boundedly invertible with  $\|\partial_{t,\rho}^{-1}\| = \frac{1}{|\rho|}$  and the inverse is given by

$$(\partial_{t,\rho}^{-1} f)(t) = \begin{cases} \int_{-\infty}^t f(s) ds & \text{if } \rho > 0, \\ -\int_t^{\infty} f(s) ds & \text{if } \rho < 0 \end{cases}$$

for  $f \in L_{2,\rho}(\mathbb{R}; H)$  and  $t \in \mathbb{R}$ .

(d) The operator  $\partial_{t,\rho}$  is normal with  $\partial_{t,\rho}^* = -\partial_{t,\rho} + 2\rho$ .

(e) The following variant of Sobolev's embedding theorem holds:

$$H_\rho^1(\mathbb{R}; H) \hookrightarrow C_\rho(\mathbb{R}; H)$$

continuously, where

$$C_\rho(\mathbb{R}; H) := \{f : \mathbb{R} \rightarrow H; f \text{ continuous, } \sup_{t \in \mathbb{R}} \|f(t)\| e^{-\rho t} < \infty\}.$$

As a normal operator,  $\partial_{t,\rho}$  possesses a natural functional calculus, which can be described via the so-called Fourier-Laplace transform.

**Definition.** Let  $\rho \in \mathbb{R}$  and  $H$  a Hilbert space. We denote by  $\mathcal{L}_\rho$  the unitary extension of the mapping

$$C_c(\mathbb{R}; H) \subseteq L_{2,\rho}(\mathbb{R}; H) \rightarrow L_2(\mathbb{R}; H), f \mapsto \left( t \mapsto \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{-(it+\rho)s} f(s) ds \right).$$

*Remark 2.2.* Note that for  $\rho = 0$ , the operator  $\mathcal{L}_0$  is nothing but the classical Fourier transform, which is unitary due to Plancharel's Theorem (see e.g. [17, Theorem 9.13]). Since  $\mathcal{L}_\rho = \mathcal{L}_0 \exp(-\rho \cdot)$  with

$$\exp(-\rho \cdot) : L_{2,\rho}(\mathbb{R}; H) \rightarrow L_2(\mathbb{R}; H), f \mapsto (t \mapsto f(t)e^{-\rho t})$$

for  $t \in \mathbb{R}$ , it follows that  $\mathcal{L}_\rho$  is unitary as a composition of unitary operators.

**Proposition 2.3** ([8, Corollary 2.5]). *Let  $\rho \in \mathbb{R}$  and  $H$  a Hilbert space. We define the operator  $m$  by*

$$m : \text{dom}(m) \subseteq L_2(\mathbb{R}; H) \rightarrow L_2(\mathbb{R}; H), f \mapsto (t \mapsto tf(t))$$

with maximal domain

$$\text{dom}(m) := \{f \in L_2(\mathbb{R}; H); (t \mapsto tf(t)) \in L_2(\mathbb{R}; H)\}.$$

Then

$$\partial_{t,\rho} = \mathcal{L}_\rho^*(i m + \rho) \mathcal{L}_\rho.$$

Using the latter proposition, we can define an operator-valued functional calculus for  $\partial_{t,\rho}$  as follows.

**Definition.** Let  $\rho \in \mathbb{R}$  and  $H$  a Hilbert space. Let  $F : \{it + \rho; t \in \mathbb{R}\} \rightarrow L(H)$  be strongly measurable and bounded. Then we define

$$F(\partial_{t,\rho}) := \mathcal{L}_\rho^* F(\text{im} + \rho) \mathcal{L}_\rho \in L(L_{2,\rho}(\mathbb{R}; H)),$$

where

$$F(\text{im} + \rho)f := (t \mapsto F(it + \rho)f(t)) \quad (f \in L_2(\mathbb{R}; H)).$$

An important class of operator-valued function of  $\partial_{t,\rho}$  are those functions yielding causal operators.

**Proposition 2.4** ([13, Theorem 6.1.1, Theorem 6.1.4]). *Let  $\rho_0 \in \mathbb{R}$  and  $H$  a Hilbert space. If  $M : \mathbb{C}_{\text{Re} > \rho_0} \rightarrow L(H)$  is analytic and bounded, then  $M(\partial_{t,\rho})$  is causal for each  $\rho > \rho_0$ , i.e., for  $f \in L_{2,\rho}(\mathbb{R}; H)$  with  $\text{spt } f \subseteq \mathbb{R}_{\geq a}$  for some  $a \in \mathbb{R}$  it follows that*

$$\text{spt } M(\partial_{t,\rho})f \subseteq \mathbb{R}_{\geq a}.$$

Moreover,  $M(\partial_{t,\rho})$  is independent of the choice of  $\rho > \rho_0$  in the sense that

$$M(\partial_{t,\rho})f = M(\partial_{t,\mu})f \quad (f \in L_{2,\rho}(\mathbb{R}; H) \cap L_{2,\mu}(\mathbb{R}; H))$$

for each  $\rho, \mu > \rho_0$ .

*Remark 2.5.* (a) The proof of causality is based on a theorem by Paley and Wiener, which characterises the functions in  $L_2(\mathbb{R}_{\geq 0}; H)$  in terms of their Laplace transform (see [10] or [17, 19.2 Theorem]). The independence of  $\rho$  is a simple application of Cauchy's Theorem for analytic functions.

(b) It is noteworthy that causal, translation-invariant and bounded operators are always of the form  $M(\partial_{t,\rho})$  for some analytic and bounded mapping defined on a right half plane (see [6, 31]).

Finally, we are in the position to define well-posed evolutionary problems.

**Definition.** Let  $\rho_0 \in \mathbb{R}$  and  $H$  a Hilbert space. Moreover, let  $M : \mathbb{C}_{\text{Re} > \rho_0} \rightarrow L(H)$  be analytic and bounded and  $A : \text{dom}(A) \subseteq H \rightarrow H$  densely defined, closed and linear. Then we call an equation of the form

$$(\partial_{t,\rho} M(\partial_{t,\rho}) + A)u = f$$

the *evolutionary equation associated with  $(M, A)$* . The problem is called *well-posed* if there is  $\rho_1 > \rho_0$  such that  $zM(z) + A$  is boundedly invertible for each  $z \in \mathbb{C}_{\text{Re} \geq \rho_1}$  and

$$\mathbb{C}_{\text{Re} \geq \rho_1} \ni z \mapsto (zM(z) + A)^{-1}$$

is bounded. Moreover we set  $s_0(M, A)$  as the infimum over all such  $\rho_1 > \rho_0$ .

**Theorem 2.6.** *Let  $\rho_0 \in \mathbb{R}$  and  $H$  a Hilbert space. Moreover, let  $M : \mathbb{C}_{\operatorname{Re} > \rho_0} \rightarrow L(H)$  be analytic and bounded and  $A : \operatorname{dom}(A) \subseteq H \rightarrow H$  densely defined closed and linear. We assume that the evolutionary equation associated with  $(M, A)$  is well-posed. Then  $\overline{\partial_{t,\rho} M(\partial_{t,\rho})} + A$  is boundedly invertible as an operator on  $L_2(\mathbb{R}; H)$  for each  $\rho > s_0(M, A)$ . Moreover, the inverse*

$$S_\rho := \left( \overline{\partial_{t,\rho} M(\partial_{t,\rho})} + A \right)^{-1}$$

*is causal and independent of the choice of  $\rho > s_0(M, A)$  in the sense of Proposition 2.4.*

*Proof.* We note that the operator  $\overline{\partial_{t,\rho} M(\partial_{t,\rho})} + A$  for  $\rho > s_0(M, A)$  is unitarily equivalent to the multiplication operator on  $L_2(\mathbb{R}; H)$  associated with the operator-valued function

$$F(t) := (it + \rho) M(it + \rho) + A,$$

see [21, Lemma 2.2], which is boundedly invertible by assumption. The causality and independence of  $\rho$  are an immediate consequence of Proposition 2.4, since  $S_\rho = N(\partial_{t,\rho})$  for the analytic and bounded function  $N(z) := (zM(z) + A)^{-1}$  for  $z \in \mathbb{C}_{\operatorname{Re} > s_0(M, A)}$ .  $\square$

### 3 Extrapolation spaces

In this section we recall the notion of extrapolation spaces associated with a boundedly invertible operator on some Hilbert space  $H$ . We refer to [13, Section 2.1] for the proof of the results presented here.

**Definition.** Let  $C : \operatorname{dom}(C) \subseteq H \rightarrow H$  be a densely defined, closed, linear and boundedly invertible operator on some Hilbert space  $H$ . We define the Hilbert space

$$H^1(C) := \operatorname{dom}(C)$$

equipped with the inner product

$$\langle x, y \rangle_{H^1(C)} := \langle Cx, Cy \rangle \quad (x, y \in \operatorname{dom}(C)).$$

Moreover, we set

$$H^{-1}(C) := H^1(C^*)',$$

the dual space of  $H^1(C^*)$ .

*Remark 3.1.* Another way to introduce the space  $H^{-1}(C)$  is taking the completion of  $H$  with respect to the norm

$$x \mapsto \|C^{-1}x\|.$$

**Proposition 3.2** ([13, Theorem 2.1.6]). *Let  $C : \operatorname{dom}(C) \subseteq H \rightarrow H$  be a densely defined, closed, linear and boundedly invertible operator on some Hilbert space  $H$ . Then  $H^1(C) \hookrightarrow H \hookrightarrow H^{-1}(C)$  with dense and continuous embeddings. Here, the second embedding is given by*

$$H \rightarrow H^{-1}(C), \quad x \mapsto (\operatorname{dom}(C^*) \ni y \mapsto \langle x, y \rangle).$$

Moreover, the operator

$$C : H^1(C) \rightarrow H$$

is unitary and

$$C : \text{dom}(C) \subseteq H \rightarrow H^{-1}(C)$$

possesses a unitary extension, which will again be denoted by  $C$ .

**Example 3.3.** Let  $\rho \neq 0$  and  $H$  a Hilbert space. Then we set

$$\begin{aligned} H_\rho^1(\mathbb{R}; H) &:= H^1(\partial_{t,\rho}), \\ H_\rho^{-1}(\mathbb{R}; H) &:= H^{-1}(\partial_{t,\rho}). \end{aligned}$$

Moreover, the Dirac distribution  $\delta_t$  at a point  $t \in \mathbb{R}$  belongs to  $H_\rho^{-1}(\mathbb{R}; \mathbb{C})$  and

$$\partial_{t,\rho}^{-1} \delta_t = \begin{cases} e^{2\rho t} \chi_{\mathbb{R}_{\geq t}} & \text{if } \rho > 0, \\ -e^{2\rho t} \chi_{\mathbb{R}_{\leq t}} & \text{if } \rho < 0. \end{cases}$$

Indeed, for  $\rho > 0$  we have that

$$\begin{aligned} \langle \partial_{t,\rho} \chi_{\mathbb{R}_{\geq t}}, \varphi \rangle_{H_\rho^{-1}(\mathbb{R}; \mathbb{C}) \times H_\rho^1(\mathbb{R}; \mathbb{C})} &= \int_t^\infty (\partial_{t,\rho}^* \varphi)(s) e^{-2\rho s} ds \\ &= - \int_t^\infty (\varphi e^{-2\rho \cdot})'(s) ds \\ &= \varphi(t) e^{-2\rho t} \end{aligned}$$

for each  $\varphi \in C_c^\infty(\mathbb{R}; \mathbb{C})$ , which shows the asserted formula. The statement for  $\rho < 0$  follows by the same rationale.

**Proposition 3.4.** Let  $\rho_0 \geq 0$  and  $H$  a Hilbert space. Moreover, let  $M : \mathbb{C}_{\text{Re} > \rho_0} \rightarrow H$  be analytic and bounded and  $A : \text{dom}(A) \subseteq H \rightarrow H$  densely defined, linear and closed such that the evolutionary problem associated with  $(M, A)$  is well-posed. Then for each  $\rho > s_0(M, A)$  we obtain

$$S_\rho[H_\rho^1(\mathbb{R}; H)] \subseteq H_\rho^1(\mathbb{R}; H)$$

and

$$S_\rho : L_{2,\rho}(\mathbb{R}; H) \subseteq H_\rho^{-1}(\mathbb{R}; H) \rightarrow H_\rho^{-1}(\mathbb{R}; H)$$

is bounded and thus has a unique bounded extension to the whole  $H_\rho^{-1}(\mathbb{R}; H)$ .

*Proof.* The assertion follows immediately by realising that

$$(\partial_{t,\rho} M(\partial_{t,\rho}) + A) \partial_{t,\rho} \subseteq \partial_{t,\rho} (\partial_{t,\rho} M(\partial_{t,\rho}) + A). \quad \square$$

We recall that for a densely defined, closed, linear operator  $A : \text{dom}(A) \subseteq H_0 \rightarrow H_1$  between two Hilbert spaces  $H_0$  and  $H_1$ , the operators  $A^*A$  and  $AA^*$  are selfadjoint and positive. Then the moduli of  $A$  and  $A^*$  are defined by

$$|A| := \sqrt{A^*A}, \quad |A^*| := \sqrt{AA^*}$$

and are selfadjoint positive operators, too (see e.g. [30, Theorem 7.20]).

**Proposition 3.5** ([13, Lemma 2.1.16]). *Let  $H_0, H_1$  be Hilbert spaces and  $A : \text{dom}(A) \subseteq H_0 \rightarrow H_1$  densely defined, closed and linear. Then*

$$A : \text{dom}(A) \subseteq H_0 \rightarrow H^{-1}(|A^*| + 1)$$

*is bounded and hence, possesses a bounded extension to  $H_0$ .*

## 4 Cut-off operators

The main goal of the present section is to extend the cut-off operators  $\chi_{\mathbb{R}_{\geq t}}$  and  $\chi_{\mathbb{R}_{\leq t}}$  for some  $t \in \mathbb{R}$  defined on  $L_{2,\rho}(\mathbb{R}; H)$  to the extrapolation space  $H_\rho^{-1}(\mathbb{R}; H)$ . For doing so, we start with the following observation.

**Lemma 4.1.** *Let  $\rho > 0, t \in \mathbb{R}$  and  $H$  be a Hilbert space. We define the operators*

$$\begin{aligned} \chi_{\mathbb{R}_{\geq t}}(\mathbf{m}) : L_{2,\rho}(\mathbb{R}; H) &\rightarrow L_{2,\rho}(\mathbb{R}; H), & f &\mapsto (s \mapsto \chi_{\mathbb{R}_{\geq t}}(s)f(s)), \\ \chi_{\mathbb{R}_{\leq t}}(\mathbf{m}) : L_{2,\rho}(\mathbb{R}; H) &\rightarrow L_{2,\rho}(\mathbb{R}; H), & f &\mapsto (s \mapsto \chi_{\mathbb{R}_{\leq t}}(s)f(s)). \end{aligned}$$

*Then for  $f \in L_{2,\rho}(\mathbb{R}; H)$  we have<sup>1</sup>*

$$\begin{aligned} \chi_{\mathbb{R}_{\geq t}}(\mathbf{m})f &= \partial_{t,\rho}\chi_{\mathbb{R}_{\geq t}}(\mathbf{m})\partial_{t,\rho}^{-1}f - e^{-2\rho t}(\partial_{t,\rho}^{-1}f)(t+)\delta_t, \\ \chi_{\mathbb{R}_{\leq t}}(\mathbf{m})f &= \partial_{t,\rho}\chi_{\mathbb{R}_{\leq t}}(\mathbf{m})\partial_{t,\rho}^{-1}f + e^{-2\rho t}(\partial_{t,\rho}^{-1}f)(t-)\delta_t. \end{aligned}$$

*Proof.* We just prove the formula for  $\chi_{\mathbb{R}_{\geq t}}(\mathbf{m})$ . So, let  $f \in L_{2,\rho}(\mathbb{R}; H)$  and set  $F := \partial_{t,\rho}^{-1}f$ . We recall from Proposition 2.1 (c) that

$$F(t) = \int_{-\infty}^t f(s) \, ds \quad (t \in \mathbb{R}).$$

For  $g \in C_c^\infty(\mathbb{R}; H)$  we compute

$$\begin{aligned} &\langle \partial_{t,\rho}\chi_{\mathbb{R}_{\geq t}}(\mathbf{m})\partial_{t,\rho}^{-1}f, g \rangle_{H^{-1}(\partial_{t,\rho}) \times H^1(\partial_{t,\rho}^*)} \\ &= \langle \chi_{\mathbb{R}_{\geq t}}(\mathbf{m})\partial_{t,\rho}^{-1}f, \partial_{t,\rho}^*g \rangle_{L_{2,\rho}(\mathbb{R}; H)} \\ &= \int_t^\infty \langle F(s), -g'(s) + 2\rho g(s) \rangle e^{-2\rho s} \, ds \\ &= \int_t^\infty \langle f(s), g(s) \rangle e^{-2\rho s} \, ds + F(t+)g(t)e^{-2\rho t} \\ &= \langle \chi_{\mathbb{R}_{\geq t}}(\mathbf{m})f, g \rangle_{L_{2,\rho}(\mathbb{R}; H)} + \langle e^{-2\rho t}F(t+)\delta_t, g \rangle_{H^{-1}(\partial_{t,\rho}) \times H^1(\partial_{t,\rho}^*)}. \end{aligned}$$

Since  $C_c^\infty(\mathbb{R}; H)$  is dense in  $H^1(\partial_{t,\rho}^*)$  by Proposition 2.1 (a), we derive the asserted formula.  $\square$

The latter representation of the cut-off operators on  $L_{2,\rho}(\mathbb{R}; H)$  leads to the following definition on  $H_\rho^{-1}(\mathbb{R}; H)$ .

<sup>1</sup>Note that  $\partial_{t,\rho}^{-1}f$  has a continuous representer by Proposition 2.1 (e).



**Definition.** Let  $\rho > 0$  and  $H$  a Hilbert space. For  $t \in \mathbb{R}$  we define the operators

$$\begin{aligned} P_t &: \text{dom}(P_t) \subseteq H_\rho^{-1}(\mathbb{R}; H) \rightarrow H_\rho^{-1}(\mathbb{R}; H), \\ Q_t &: \text{dom}(Q_t) \subseteq H_\rho^{-1}(\mathbb{R}; H) \rightarrow H_\rho^{-1}(\mathbb{R}; H), \end{aligned}$$

with the domains

$$\begin{aligned} \text{dom}(P_t) &:= \{f \in H_\rho^{-1}(\mathbb{R}; H); (\partial_{t,\rho}^{-1} f)(t+) \text{ exists}\}, \\ \text{dom}(Q_t) &:= \{f \in H_\rho^{-1}(\mathbb{R}; H); (\partial_{t,\rho}^{-1} f)(t-) \text{ exists}\} \end{aligned}$$

by

$$P_t f := \partial_{t,\rho} \chi_{\mathbb{R}_{\geq t}}(\mathfrak{m}) \partial_{t,\rho}^{-1} f - e^{-2\rho t} (\partial_{t,\rho}^{-1} f)(t+) \delta_t \quad (f \in \text{dom}(P_t))$$

and

$$Q_t f := \partial_{t,\rho} \chi_{\mathbb{R}_{\leq t}}(\mathfrak{m}) \partial_{t,\rho}^{-1} f + e^{-2\rho t} (\partial_{t,\rho}^{-1} f)(t-) \delta_t \quad (f \in \text{dom}(Q_t)).$$

*Remark 4.2.* For a function  $f \in L_{1,\text{loc}}(\mathbb{R}; H)$  we say that  $a := f(t+)$  for some  $t \in \mathbb{R}$  if

$$\forall \varepsilon > 0 \exists \delta > 0 : \lambda(\{s \in [t, t + \delta]; |f(s) - a| > \varepsilon\}) = 0,$$

where  $\lambda$  denotes the Lebesgue measure on  $\mathbb{R}$ . The expression  $f(t-)$  is defined analogously.

We conclude this section by some properties of the so introduced cut-off operators.

**Proposition 4.3.** *Let  $H$  be a Hilbert space,  $\rho > 0$ ,  $y \in H$  and  $s, t \in \mathbb{R}$ . Then the following statements hold.*

(a)  $\delta_s y \in \text{dom}(P_t)$  and

$$P_t \delta_s y = \begin{cases} \delta_s y & \text{if } s > t, \\ 0 & \text{if } s \leq t. \end{cases}$$

(b) For  $f \in \text{dom}(P_t) \cap \text{dom}(Q_t)$  we obtain

$$f = P_t f + Q_t f + e^{-2\rho t} ((\partial_{t,\rho}^{-1} f)(t+) - (\partial_{t,\rho}^{-1} f)(t-)) \delta_t.$$

(c) For  $f \in H_\rho^{-1}(\mathbb{R}; H)$  we have  $\text{spt } f \subseteq \mathbb{R}_{\leq t}$  if and only if  $f \in \ker(P_t)$ . Here, the support  $\text{spt } f$  is meant in the sense of distributions.

*Proof.* (a) We note that  $\partial_{t,\rho}^{-1} \delta_s y = e^{2\rho s} \chi_{\mathbb{R}_{\geq s}} y$  and hence,  $\delta_s \in \text{dom}(P_t)$ . Moreover,

$$P_t \delta_s y = \partial_{t,\rho} \chi_{\mathbb{R}_{\geq t}}(\mathfrak{m}) \chi_{\mathbb{R}_{\geq s}} y e^{2\rho s} - e^{-2\rho t} (e^{2\rho s} \chi_{\mathbb{R}_{\geq s}} y)(t+) \delta_t = \begin{cases} \delta_s y & \text{if } s > t, \\ 0 & \text{if } s \leq t. \end{cases}$$

(b) If  $f \in \text{dom}(P_t) \cap \text{dom}(Q_t)$  we compute

$$\begin{aligned} P_t f + Q_t f &= \partial_{t,\rho} \chi_{\mathbb{R}_{\geq t}}(\mathfrak{m}) \partial_{t,\rho}^{-1} f - e^{-2\rho t} (\partial_{t,\rho}^{-1} f)(t+) \delta_t + \partial_{t,\rho} \chi_{\mathbb{R}_{\leq t}}(\mathfrak{m}) \partial_{t,\rho}^{-1} f + e^{-2\rho t} (\partial_{t,\rho}^{-1} f)(t-) \delta_t \\ &= \partial_{t,\rho} \partial_{t,\rho}^{-1} f - e^{-2\rho t} ((\partial_{t,\rho}^{-1} f)(t+) - (\partial_{t,\rho}^{-1} f)(t-)) \delta_t \\ &= f - e^{-2\rho t} ((\partial_{t,\rho}^{-1} f)(t+) - (\partial_{t,\rho}^{-1} f)(t-)) \delta_t. \end{aligned}$$

(c) Let  $f \in H_\rho^{-1}(\mathbb{R}; H)$  and assume first that  $\text{spt } f \subseteq \mathbb{R}_{\leq t}$ . We first prove that  $\partial_{t,\rho}^{-1} f$  is constant on  $\mathbb{R}_{\geq t}$ . For doing so, we define

$$V := \{\chi_{\mathbb{R}_{\geq t}} x; x \in H\} \subseteq L_{2,\rho}(\mathbb{R}; H).$$

Then  $V$  is a closed subspace and for  $g \in L_{2,\rho}(\mathbb{R}; H)$  we have that

$$g \in V^\perp \Leftrightarrow \int_t^\infty g(s) e^{-2\rho s} ds = 0.$$

For  $g \in L_{2,\rho}(\mathbb{R}; H)$  we obtain

$$\langle \chi_{\mathbb{R}_{\geq t}}(\mathbf{m}) \partial_{t,\rho}^{-1} f, g \rangle_{L_{2,\rho}(\mathbb{R}; H)} = \langle f, (\partial_{t,\rho}^*)^{-1} \chi_{\mathbb{R}_{\geq t}}(\mathbf{m}) g \rangle_{H_\rho^{-1}(\mathbb{R}; H) \times H_\rho^1(\mathbb{R}; H)}$$

and an elementary computation shows

$$\left( (\partial_{t,\rho}^*)^{-1} \chi_{\mathbb{R}_{\geq t}}(\mathbf{m}) g \right) (s) = \int_s^\infty \chi_{\mathbb{R}_{\geq t}}(r) g(r) e^{2\rho(s-r)} dr \quad (s \in \mathbb{R}).$$

Consequently, for  $g \in V^\perp$  we infer that  $(\partial_{t,\rho}^*)^{-1} \chi_{\mathbb{R}_{\geq t}}(\mathbf{m}) g = 0$  on  $\mathbb{R}_{\leq t}$ . Hence,  $\langle \chi_{\mathbb{R}_{\geq t}}(\mathbf{m}) \partial_{t,\rho}^{-1} f, g \rangle_{L_{2,\rho}(\mathbb{R}; H)} = 0$  for each  $g \in V^\perp$  and thus,  $\chi_{\mathbb{R}_{\geq t}}(\mathbf{m}) \partial_{t,\rho}^{-1} f \in V$ , which proves that  $\partial_{t,\rho}^{-1} f$  is constant on  $\mathbb{R}_{\geq t}$ . In particular, this shows  $f \in \text{dom}(P_t)$  and

$$\begin{aligned} P_t f &= \partial_{t,\rho} \chi_{\mathbb{R}_{\geq t}}(\mathbf{m}) \partial_{t,\rho}^{-1} f - e^{-2\rho t} (\partial_{t,\rho}^{-1} f)(t+) \delta_t \\ &= \partial_{t,\rho} \chi_{\mathbb{R}_{\geq t}}(\mathbf{m}) (\partial_{t,\rho}^{-1} f)(t+) - e^{-2\rho t} (\partial_{t,\rho}^{-1} f)(t+) \delta_t \\ &= 0. \end{aligned}$$

Assume on the other hand that  $f \in \ker(P_t)$  and let  $\varphi \in C_c^\infty(\mathbb{R}_{>t}; H)$ . We then compute, using that  $\text{spt } \partial_{t,\rho}^* \varphi \subseteq \mathbb{R}_{>t}$

$$\begin{aligned} \langle f, \varphi \rangle_{H_\rho^{-1}(\mathbb{R}; H) \times H_\rho^1(\mathbb{R}; H)} &= \langle \partial_{t,\rho}^{-1} f, \partial_{t,\rho}^* \varphi \rangle_{L_{2,\rho}(\mathbb{R}; H)} \\ &= \langle P_t f, \varphi \rangle_{H_\rho^{-1} \times H_\rho^1} + e^{-2\rho t} (\partial_{t,\rho}^{-1} f)(t+) \varphi(t) \\ &= 0, \end{aligned}$$

which gives  $\text{spt } f \subseteq \mathbb{R}_{\leq t}$ . □

## 5 Admissible histories for evolutionary equations

In this section we study evolutionary problems of the following form

$$\begin{aligned} (\partial_{t,\rho} M(\partial_{t,\rho}) + A) u &= 0 \quad \text{on } \mathbb{R}_{>0}, \\ u &= g \quad \text{on } \mathbb{R}_{<0}, \end{aligned} \tag{3}$$

where  $M$  and  $A$  are as in Theorem 2.6 and  $g$  is a given function on  $\mathbb{R}_{<0}$ . The first goal is to rewrite this ‘Initial value problem’ into a proper evolutionary equations as it

is introduced in Section 2. For doing so, we start with some heuristics to motivate the definition which will be made below. In particular, for the moment we will not care about domains of operators.

We will now write (3) as an evolutionary equation for the unknown  $v := u|_{\mathbb{R}_{\geq 0}}$ , which is the part of  $u$  to be determined. For doing so, we first assume that  $u \in H_\rho^1(\mathbb{R}; H)$  for some  $\rho > 0$ , which means that  $v + g \in H_\rho^1(\mathbb{R}; H)$ . We interpret the first line of (3) as

$$P_0 (\partial_{t,\rho} M(\partial_{t,\rho}) + A) u = 0,$$

where  $P_0$  is the cut-off operator introduced in Section 4. The latter gives

$$\begin{aligned} 0 &= P_0 (\partial_{t,\rho} M(\partial_{t,\rho}) + A) u \\ &= P_0 (\partial_{t,\rho} M(\partial_{t,\rho}) + A) v + P_0 (\partial_{t,\rho} M(\partial_{t,\rho}) + A) g \\ &= \partial_{t,\rho} P_0 M(\partial_{t,\rho}) v + A P_0 v - (M(\partial_{t,\rho}) v) (0+) \delta_0 + P_0 \partial_{t,\rho} M(\partial_{t,\rho}) g + A P_0 g \\ &= \partial_{t,\rho} P_0 M(\partial_{t,\rho}) v + A v + P_0 \partial_{t,\rho} M(\partial_{t,\rho}) g - (M(\partial_{t,\rho}) v) (0+) \delta_0. \end{aligned}$$

Since  $v$  is supported on  $\mathbb{R}_{\geq 0}$  by assumption and  $M(\partial_{t,\rho})$  is causal by Proposition 2.4, we infer that  $M(\partial_{t,\rho}) v$  is also supported on  $\mathbb{R}_{\geq 0}$  and so,  $P_0 M(\partial_{t,\rho}) v = M(\partial_{t,\rho}) v$ . Hence, we arrive at an evolutionary problem for  $v$  of the form

$$(\partial_{t,\rho} M(\partial_{t,\rho}) + A) v = (M(\partial_{t,\rho}) v) (0+) \delta_0 - P_0 \partial_{t,\rho} M(\partial_{t,\rho}) g.$$

Since  $u = v + g \in H_\rho^1(\mathbb{R}; H)$  by assumption, we infer that  $u$  is continuous by Proposition 2.1 (e) and hence, the limits  $v(0+)$  and  $g(0-)$  exist and coincide. Hence,  $v - \chi_{\mathbb{R}_{\geq 0}} g(0-) \in H_\rho^1(\mathbb{R}; H)$  and vanishes on  $\mathbb{R}_{< 0}$ . The latter gives

$$\begin{aligned} (M(\partial_{t,\rho}) v) (0+) &= (M(\partial_{t,\rho}) (v - \chi_{\mathbb{R}_{\geq 0}} g(0-))) (0+) + (M(\partial_{t,\rho}) \chi_{\mathbb{R}_{\geq 0}} g(0-)) (0+) \\ &= (M(\partial_{t,\rho}) \chi_{\mathbb{R}_{\geq 0}} g(0-)) (0+), \end{aligned}$$

where in the last equality we have used that  $M(\partial_{t,\rho}) (v - \chi_{\mathbb{R}_{\geq 0}} g(0-)) \in H_\rho^1(\mathbb{R}; H)$ , hence it is continuous, and vanishes on  $\mathbb{R}_{\leq 0}$  due to causality. Summarising, we end up with the following problem for  $v$

$$(\partial_{t,\rho} M(\partial_{t,\rho}) + A) v = (M(\partial_{t,\rho}) \chi_{\mathbb{R}_{\geq 0}} g(0-)) (0+) \delta_0 - P_0 \partial_{t,\rho} M(\partial_{t,\rho}) g. \quad (4)$$

Now, to make sense of (4) we need to ensure that the right hand side is well-defined. In particular, we need that  $(M(\partial_{t,\rho}) \chi_{\mathbb{R}_{\geq 0}} g(0-)) (0+)$  exists. In order to ensure that, we introduce the following notion.

**Definition.** Let  $H$  be a Hilbert space,  $\rho_0 \geq 0$  and  $M : \mathbb{C}_{\operatorname{Re} > \rho_0} \rightarrow L(H)$  be analytic and bounded. We call  $M$  *regularising*, if for all  $x \in H, \rho > \rho_0$  the limit

$$(M(\partial_{t,\rho}) \chi_{\mathbb{R}_{\geq 0}} x) (0+)$$

exists. Moreover, for  $\rho > 0$  we define the space

$$H_\rho^1(\mathbb{R}_{\leq 0}; H) := \{f|_{\mathbb{R}_{\leq 0}}; f \in H_\rho^1(\mathbb{R}; H)\}.$$

As it turns out, this assumption suffices to obtain a well-defined expression on the right hand side of (4).

**Proposition 5.1.** *Let  $H$  be a Hilbert space,  $\rho_0 \geq 0$  and  $M : \mathbb{C}_{\text{Re} > \rho_0} \rightarrow L(H)$  be analytic and bounded and assume that  $M$  is regularising. Then for each  $g \in H_\rho^1(\mathbb{R}_{\leq 0}; H)$  with  $\rho > \rho_0$  we have that*

$$\partial_{t,\rho} M(\partial_{t,\rho})g \in \text{dom}(P_0).$$

*Proof.* By assumption  $g = f|_{\mathbb{R}_{\leq 0}}$  for some  $f \in H_\rho^1(\mathbb{R}; H)$ . Hence,  $g(0-) = f(0)$  exists and hence, an easy computation shows that  $g - \chi_{\mathbb{R}_{\leq 0}}g(0-) \in H_\rho^1(\mathbb{R}; H)$ . Hence, also  $M(\partial_{t,\rho})(g - \chi_{\mathbb{R}_{\geq 0}}g(0-)) \in H_\rho^1(\mathbb{R}; H)$  and thus,

$$(M(\partial_{t,\rho})g)(0+) = (M(\partial_{t,\rho})(g - \chi_{\mathbb{R}_{\geq 0}}g(0-)))(0+) + (M(\partial_{t,\rho})\chi_{\mathbb{R}_{\geq 0}}g(0-))(0+)$$

exists and so,  $\partial_{t,\rho} M(\partial_{t,\rho})g \in \text{dom}(P_0)$ .  $\square$

We are now in the position to define the space of admissible history functions  $g$ .

**Definition.** Let  $H$  be a Hilbert space,  $\rho_0 \geq 0$  and  $M : \mathbb{C}_{\text{Re} > \rho_0} \rightarrow L(H)$  be analytic, bounded and regularising. Moreover, let  $A : \text{dom}(A) \subseteq H \rightarrow H$  be densely defined, closed and linear. For notational convenience, we set

$$\Gamma_\rho : H_\rho^1(\mathbb{R}_{\leq 0}; H) \rightarrow H, \quad g \mapsto (M(\partial_{t,\rho})\chi_{\mathbb{R}_{\geq 0}}g(0-))(0+)$$

and

$$K_\rho : H_\rho^1(\mathbb{R}_{\leq 0}; H) \rightarrow H_\rho^{-1}(\mathbb{R}; H), \quad g \mapsto P_0 \partial_{t,\rho} M(\partial_{t,\rho})g$$

for  $\rho > \rho_0$ . Furthermore, we assume that the evolutionary problem associated with  $(M, A)$  is well-posed and define

$$\text{His}_\rho := \{g \in H_\rho^1(\mathbb{R}_{\leq 0}; H); S_\rho(\Gamma_\rho g \delta_0 - K_\rho g) + g \in H_\rho^1(\mathbb{R}; H)\}$$

for each  $\rho > s_0(M, A)$ , the *space of admissible histories*. Here  $S_\rho$  denotes the extension of the solution operator  $(\partial_{t,\rho} M(\partial_{t,\rho}) + A)^{-1}$  to  $H_\rho^{-1}(\mathbb{R}; H)$  (cp. Proposition 3.4). Moreover, we set

$$\text{IV}_\rho := \{g(0-); g \in \text{His}_\rho\}$$

the space of *admissible initial values*.

*Remark 5.2.* We have

$$\Gamma_\rho g = (M(\partial_{t,\rho})g)(0-) - (M(\partial_{t,\rho})g)(0+)$$

for  $g \in H_\rho^1(\mathbb{R}_{\leq 0}; H)$ . Indeed, since  $M(\partial_{t,\rho})$  is causal we infer

$$\begin{aligned} (M(\partial_{t,\rho})g)(0-) &= (M(\partial_{t,\rho})(g + \chi_{\mathbb{R}_{\geq 0}}g(0-)))(0-) \\ &= (M(\partial_{t,\rho})(g + \chi_{\mathbb{R}_{\geq 0}}g(0-)))(0+), \end{aligned}$$

since  $g + \chi_{\mathbb{R}_{\geq 0}}g(0-) \in H_\rho^1(\mathbb{R}; H)$ . Thus,

$$(M(\partial_{t,\rho})g)(0-) - (M(\partial_{t,\rho})g)(0+) = (M(\partial_{t,\rho})\chi_{\mathbb{R}_{\geq 0}}g(0-))(0+) = \Gamma_\rho g.$$

We come back to the heuristic computation at the beginning of this section and show, that for  $g \in \text{His}_\rho$  the computation can be made rigorously.

**Proposition 5.3.** *Let  $H$  be a Hilbert space,  $\rho_0 \geq 0$  and  $M : \mathbb{C}_{\text{Re} > \rho_0} \rightarrow L(H)$  be analytic, bounded and regularising. Moreover, let  $A : \text{dom}(A) \subseteq H \rightarrow H$  be densely defined, closed and linear and assume that the evolutionary problem associated with  $(M, A)$  is well-posed. Let  $\rho > s_0(M, A)$  and  $g \in \text{His}_\rho$ . We set*

$$v := S_\rho(\Gamma_\rho g \delta_0 - K_\rho g)$$

and  $u := v + g$ . Then  $\text{spt } v \subseteq \mathbb{R}_{\geq 0}$ ,  $u \in H_\rho^1(\mathbb{R}; H)$  and satisfies (3).

*Proof.* Note that by assumption  $u = v + g \in H_\rho^1(\mathbb{R}; H)$  and thus,  $v = u - g \in L_{2,\rho}(\mathbb{R}; H)$ . We prove that  $\text{spt } v \subseteq \mathbb{R}_{\geq 0}$ . For doing so, we compute

$$\begin{aligned} \partial_{t,\rho}^{-1} v &= \partial_{t,\rho}^{-1} S_\rho(\Gamma_\rho g \delta_0 - K_\rho g) \\ &= S_\rho(\partial_{t,\rho}^{-1} \Gamma_\rho g \delta_0 - \partial_{t,\rho}^{-1} K_\rho g) \\ &= S_\rho(\Gamma_\rho g \chi_{\mathbb{R}_{\geq 0}} - \chi_{\mathbb{R}_{\geq 0}}(\mathfrak{m})M(\partial_{t,\rho})g + (M(\partial_{t,\rho})g)(0+)\chi_{\mathbb{R}_{\geq 0}}) \end{aligned}$$

and hence,  $\text{spt } \partial_{t,\rho}^{-1} v \subseteq \mathbb{R}_{\geq 0}$  by causality of  $S_\rho$ . The latter implies  $\text{spt } v \subseteq \mathbb{R}_{\geq 0}$ . Thus, we have  $u = g$  on  $\mathbb{R}_{< 0}$  and we are left to show

$$(\partial_{t,\rho} M(\partial_{t,\rho}) + A)u = 0 \quad \text{on } \mathbb{R}_{> 0}.$$

For doing so, let  $\varphi \in C_c^\infty(\mathbb{R}_{> 0}; \text{dom}(A^*))$ . We compute

$$\begin{aligned} &\langle (\partial_{t,\rho} M(\partial_{t,\rho}) + A)u, \varphi \rangle_{L_{2,\rho}(\mathbb{R}; H^{-1}(|A^*|+1)) \times L_{2,\rho}(\mathbb{R}; H^1(|A^*|+1))} \\ &= \langle u, (\partial_{t,\rho} M(\partial_{t,\rho}) + A)^* \varphi \rangle_{L_{2,\rho}(\mathbb{R}; H)} \\ &= \langle v, (\partial_{t,\rho} M(\partial_{t,\rho}) + A)^* \varphi \rangle_{L_{2,\rho}(\mathbb{R}; H)} + \langle g, (\partial_{t,\rho} M(\partial_{t,\rho}) + A)^* \varphi \rangle_{L_{2,\rho}(\mathbb{R}; H)} \\ &= \langle \Gamma_\rho g \delta_0 - K_\rho g, \varphi \rangle_{H_\rho^{-1}(\mathbb{R}; H) \times H_\rho^1(\mathbb{R}; H)} + \langle g, (\partial_{t,\rho} M(\partial_{t,\rho}))^* \varphi \rangle_{L_{2,\rho}(\mathbb{R}; H)}, \end{aligned}$$

where in the last line we have used  $\langle g, A^* \varphi \rangle = 0$ , since  $\text{spt } g \subseteq \mathbb{R}_{\leq 0}$ . Moreover, we compute

$$\begin{aligned} &\langle \Gamma_\rho g \delta_0 - K_\rho g, \varphi \rangle_{H_\rho^{-1}(\mathbb{R}; H) \times H_\rho^1(\mathbb{R}; H)} \\ &= -\langle K_\rho g, \varphi \rangle_{H_\rho^{-1}(\mathbb{R}; H) \times H_\rho^1(\mathbb{R}; H)} \\ &= -\langle \partial_{t,\rho} \chi_{\mathbb{R}_{\geq 0}}(\mathfrak{m})M(\partial_{t,\rho})g - (M(\partial_{t,\rho})g)(0+)\delta_0, \varphi \rangle_{H_\rho^{-1}(\mathbb{R}; H) \times H_\rho^1(\mathbb{R}; H)} \\ &= -\langle M(\partial_{t,\rho})g, \partial_{t,\rho}^* \varphi \rangle_{L_{2,\rho}(\mathbb{R}; H)}, \end{aligned}$$

where we have used two times that  $\varphi(0) = 0$ . Plugging this formula in the above computation, we infer that

$$\langle (\partial_{t,\rho} M(\partial_{t,\rho}) + A)u, \varphi \rangle_{L_{2,\rho}(\mathbb{R}; H^{-1}(|A^*|+1)) \times L_{2,\rho}(\mathbb{R}; H^1(|A^*|+1))} = 0,$$

which shows the claim.  $\square$

## 6 $C_0$ -semigroups associated with evolutionary problems

Throughout this section, let  $H$  be a Hilbert space,  $\rho_0 \geq 0$  and  $M : \mathbb{C}_{\text{Re} > \rho_0} \rightarrow L(H)$  analytic, bounded and regularising. Moreover, let  $A : \text{dom}(A) \subseteq H \rightarrow H$  be densely defined, closed and linear such that the evolutionary problem associated with  $(M, A)$  is well-posed.

In this section we aim for a  $C_0$ -semigroup associated with the evolutionary problem for  $(M, A)$  acting on a suitable subspace of  $\text{IV}_\rho \times \text{His}_\rho$  for  $\rho > s_0(M, A)$ . For doing so, we first need to prove that  $\text{His}_\rho$  is left invariant by the time evolution. The precise statement is as follows.

**Theorem 6.1.** *Let  $\rho > s_0(M, A)$  and  $g \in \text{His}_\rho$ . Moreover, let  $v := S_\rho(\Gamma_\rho g \delta_0 - K_\rho g)$  and  $u := v + g$ . For  $t > 0$  we set  $h := \chi_{\mathbb{R}_{\leq 0}}(\mathfrak{m})u(t + \cdot)$  and  $w := \chi_{\mathbb{R}_{\geq 0}}(\mathfrak{m})u(t + \cdot)$ . Then  $h \in \text{His}_\rho$  and*

$$w = S_\rho(\Gamma_\rho h \delta_0 - K_\rho h).$$

*In particular,  $w(0+) = h(0-) \in \text{IV}_\rho$ .*

*Proof.* We first note that

$$(\partial_{t,\rho}M(\partial_{t,\rho}) + A)\tau_t = \tau_t(\partial_{t,\rho}M(\partial_{t,\rho}) + A),$$

where  $\tau_t u := u(t + \cdot)$  for  $u \in L_{2,\rho}(\mathbb{R}; H)$ , and hence,

$$\text{spt}(\partial_{t,\rho}M(\partial_{t,\rho}) + A)\tau_t u \subseteq \mathbb{R}_{\leq 0}.$$

The latter gives, employing the causality of  $M(\partial_{t,\rho})$ ,

$$\begin{aligned} (\partial_{t,\rho}M(\partial_{t,\rho}) + A)\tau_t u &= \chi_{\mathbb{R}_{\leq 0}}(\mathfrak{m})(\partial_{t,\rho}M(\partial_{t,\rho}) + A)\tau_t u \\ &= \partial_{t,\rho}\chi_{\mathbb{R}_{\leq 0}}(\mathfrak{m})M(\partial_{t,\rho})\tau_t u + (M(\partial_{t,\rho})\tau_t u)(0-)\delta_0 + Ah \\ &= \partial_{t,\rho}\chi_{\mathbb{R}_{\leq 0}}(\mathfrak{m})M(\partial_{t,\rho})h + (M(\partial_{t,\rho})h)(0-)\delta_0 + Ah \\ &= Q_0\partial_{t,\rho}M(\partial_{t,\rho})h + Ah. \end{aligned}$$

The latter yields

$$\begin{aligned} (\partial_{t,\rho}M(\partial_{t,\rho}) + A)w &= (\partial_{t,\rho}M(\partial_{t,\rho}) + A)(\tau_t u - h) \\ &= Q_0\partial_{t,\rho}M(\partial_{t,\rho})h - \partial_{t,\rho}M(\partial_{t,\rho})h. \end{aligned}$$

Now, since  $\partial_{t,\rho}M(\partial_{t,\rho})h \in \text{dom}(P_0)$  by causality of  $M(\partial_{t,\rho})$ , we use Proposition 4.3 (b) and Remark 5.2 to derive

$$\begin{aligned} (\partial_{t,\rho}M(\partial_{t,\rho}) + A)w &= -P_0\partial_{t,\rho}M(\partial_{t,\rho})h - ((M(\partial_{t,\rho})h)(0+) - (M(\partial_{t,\rho})h)(0-))\delta_0 \\ &= \Gamma_\rho h \delta_0 - K_\rho h, \end{aligned}$$

which yields the desired formula for  $w$ . Now  $h \in \text{His}_\rho$  follows, since by definition

$$S_\rho(\Gamma_\rho h \delta_0 - K_\rho h) + h = w + h = \tau_t u \in H_\rho^1(\mathbb{R}; H). \quad \square$$

The latter theorem allows for defining a semigroup associated with  $(M, A)$ .

**Definition.** Let  $\rho > s_0(M, A)$  and set

$$D_\rho := \{(g(0-), g) ; g \in \text{His}_\rho\}.$$

For  $g \in \text{His}_\rho$  we set

$$v := S_\rho(\Gamma_\rho g \delta_0 - K_\rho g)$$

and  $u := v + g$ . For  $t \geq 0$  we define

$$\begin{aligned} T_1^\rho(t) : D_\rho \subseteq \text{IV}_\rho \times \text{His}_\rho &\rightarrow \text{IV}_\rho, & (g(0-), g) &\mapsto v(t+), \\ T_2^\rho(t) : D_\rho \subseteq \text{IV}_\rho \times \text{His}_\rho &\rightarrow \text{His}_\rho, & (g(0-), g) &\mapsto \chi_{\mathbb{R}_{\leq 0}}(\mathbf{m})\tau_t u \end{aligned}$$

and

$$T^\rho(t) := (T_1^\rho(t), T_2^\rho(t)) : D_\rho \subseteq \text{IV}_\rho \times \text{His}_\rho \rightarrow \text{IV}_\rho \times \text{His}_\rho.$$

We call  $(T^\rho(t))_{t \geq 0}$  the *semigroup associated with*  $(M, A)$ .

First we show that  $T^\rho$  defined above is indeed a strongly continuous semigroup.

**Proposition 6.2.** *Let  $\rho > s_0(M, A)$  and  $T^\rho$  be the semigroup associated with  $(M, A)$ . Then  $T^\rho$  is a strongly continuous semigroup. More precisely,*

$$T^\rho(t+s) = T^\rho(t)T^\rho(s) \quad (t, s \geq 0)$$

and

$$T^\rho(t)(g(0-), g) \rightarrow (g(0-), g) \quad (t \rightarrow 0+)$$

in  $H \times L_{2,\rho}(\mathbb{R}; H)$  for each  $g \in \text{His}_\rho$ .

*Proof.* Let  $g \in \text{His}_\rho$  and  $t, s \geq 0$ . We set  $v := S_\rho(\Gamma_\rho g \delta_0 - K_\rho g)$  and  $u := v + g$ . By Theorem 6.1 we have that

$$\chi_{\mathbb{R}_{\geq 0}}(\mathbf{m})\tau_s u = S_\rho(\Gamma_\rho(\chi_{\mathbb{R}_{\leq 0}}(\mathbf{m})\tau_s u)\delta_0 - K_\rho(\chi_{\mathbb{R}_{\leq 0}}(\mathbf{m})\tau_s u)).$$

and thus,

$$\begin{aligned} T^\rho(t)T^\rho(s)(g(0-), g) &= T^\rho(t)(u(s), \chi_{\mathbb{R}_{\leq 0}}(\mathbf{m})\tau_s u) \\ &= (u(t+s), \chi_{\mathbb{R}_{\leq 0}}(\mathbf{m})\tau_t \tau_s u) \\ &= T^\rho(t+s)(g(0-), g). \end{aligned}$$

Moreover,

$$\begin{aligned} &\|T^\rho(t)(g(0-), g) - (g(0-), g)\|_{H \times L_{2,\rho}(\mathbb{R}; H)}^2 \\ &= \|u(t) - g(0-)\|_H^2 + \|\chi_{\mathbb{R}_{\leq 0}}(\mathbf{m})\tau_t u - g\|_{L_{2,\rho}(\mathbb{R}; H)}^2 \\ &= \|u(t) - u(0)\|_H^2 + \|\chi_{\mathbb{R}_{\leq 0}}(\mathbf{m})(\tau_t u - u)\|_{L_{2,\rho}(\mathbb{R}; H)}^2 \\ &\leq \|u(t) - u(0)\|_H^2 + \|\tau_t u - u\|_{L_{2,\rho}(\mathbb{R}; H)}^2 \rightarrow 0 \quad (t \rightarrow 0+), \end{aligned}$$

by the continuity of  $u$  and the strong continuity of translation in  $L_{2,\rho}$ .  $\square$

In the rest of this section we show a characterisation result, when  $T^\rho$  can be extended to a  $C_0$ -semigroup on the space

$$X_\rho^\mu := \overline{D_\rho}^{H \times L_{2,\mu}(\mathbb{R}; H)} \subseteq H \times L_{2,\mu}(\mathbb{R}; H)$$

for some  $\mu \leq \rho$ . We first prove a result that suffices to consider the family  $T_1^\rho$ .

**Proposition 6.3.** *Let  $\rho > s_0(M, A)$  and  $\mu \leq \rho$ . Assume that*

$$T_1^\rho : D_\rho \subseteq X_\rho^\mu \rightarrow C_\omega(\mathbb{R}_{\geq 0}; H)$$

*is bounded for some  $\omega \in \mathbb{R}$ . Then*

$$T_2^\rho : D_\rho \subseteq X_\rho^\mu \rightarrow C_{\max\{\mu, \omega\} + \varepsilon}(\mathbb{R}_{\geq 0}; L_{2,\mu}(\mathbb{R}; H))$$

*is bounded for each  $\varepsilon > 0$ .*

*Proof.* Let  $\varepsilon > 0$  and  $g \in \text{His}_\rho$ . We note that

$$(T_2^\rho(t)(g(0-), g))(s) = \begin{cases} g(t+s) & \text{if } s < -t, \\ T_1^\rho(t+s)(g(0-), g) & \text{if } -t \leq s \leq 0 \end{cases} \quad (t \geq 0, s \leq 0).$$

Hence, we may estimate for  $\varepsilon > 0$

$$\begin{aligned} \|T_2^\rho(t)(g(0-), g)\|_{L_{2,\mu}(\mathbb{R}; H)}^2 &= \int_{-\infty}^{-t} \|g(t+s)\|^2 e^{-2\mu s} ds + \int_{-t}^0 \|T_1^\rho(t+s)(g(0-), g)\|^2 e^{-2\mu s} ds \\ &\leq \int_{-\infty}^0 \|g(s)\|^2 e^{-2\mu s} ds e^{2\mu t} + M \|(g(0-), g)\|_{X_\rho^\mu}^2 \int_{-t}^0 e^{2\omega(t+s)} e^{-2\mu s} ds \\ &= \|g\|_{L_{2,\mu}(\mathbb{R}; H)}^2 e^{2\mu t} + M \|(g(0-), g)\|_{X_\rho^\mu}^2 e^{2\omega t} \frac{1}{2(\omega - \mu)} (1 - e^{-2(\omega - \mu)t}) \\ &\leq \|g\|_{L_{2,\mu}(\mathbb{R}; H)}^2 e^{2\mu t} + M \|(g(0-), g)\|_{X_\rho^\mu}^2 t e^{2\max\{\mu, \omega\}t} \\ &\leq C e^{2(\max\{\mu, \omega\} + \varepsilon)t} \|(g(0-), g)\|_{X_\rho^\mu}^2 \end{aligned}$$

for each  $g \in \text{His}_\rho$ , where  $M$  denotes the norm of  $T_1^\rho$  and  $C := \max_{t \geq 0} (1 + Mt) e^{-2\varepsilon t}$ .  $\square$

In order to extend  $T_1^\rho$  to  $X_\rho^\mu$  we make use of the Widder-Arendt-Theorem.

**Theorem 6.4** (Widder-Arendt, [1],[2, Theorem 2.2.3]). *Let  $H$  be a Hilbert space and  $r \in C^\infty(\mathbb{R}_{> 0}; H)$  such that*

$$M := \sup_{\lambda > 0, k \in \mathbb{N}} \frac{\lambda^{k+1}}{k!} \|r^{(k)}(\lambda)\| < \infty.$$

*Then there is  $f \in L_\infty(\mathbb{R}_{\geq 0}; H)$  such that  $\|f\|_\infty = M$  and*

$$r(\lambda) = \int_0^\infty e^{-\lambda t} f(t) dt \quad (\lambda > 0).$$



*Remark 6.5.* The latter Theorem was first proved by Widder in the scalar-valued case [32] and then generalised by Arendt to the vector-valued case in [1]. It is noteworthy that the latter Theorem is also true in Banach spaces satisfying the Radon-Nikodym property (see [4, Chapter III]) and, in fact, this property of  $X$  is equivalent to the validity of Theorem 6.4, see [1, Theorem 1.4].

We now identify the function  $r$  mentioned in Theorem 6.4 within the presented framework.

**Proposition 6.6.** *Let  $\rho > s_0(M, A)$  and  $g \in \text{His}_\rho$ . We set  $v := S_\rho(\Gamma_\rho g \delta_0 - K_\rho g) \in L_{2,\rho}(\mathbb{R}; H)$  and*

$$r_g(\lambda) := \sqrt{2\pi}(\mathcal{L}_\lambda v)(0) \quad (\lambda > \rho).$$

*Then  $r_g \in C^\infty(\mathbb{R}_{>\rho}; H)$ . Moreover,*

$$r_g(\lambda) = (\lambda M(\lambda) + A)^{-1} \left( (M(\partial_{t,\rho})g)(0-) - \lambda\sqrt{2\pi}\mathcal{L}_\lambda(\chi_{\mathbb{R}_{\geq 0}}(\mathfrak{m})M(\partial_{t,\rho})g)(0) \right) \quad (\lambda > \rho).$$

*Proof.* We note that

$$(\mathcal{L}_\lambda v)(0) = \frac{1}{\sqrt{2\pi}} \int_0^\infty e^{-\lambda t} v(t) dt \quad (\lambda > \rho)$$

and hence, the regularity of  $r_g$  follows. Moreover,

$$\begin{aligned} \partial_{\lambda,t}^{-1} v &= \partial_{t,\rho}^{-1} v \\ &= S_\rho(\partial_{t,\rho}^{-1} \Gamma_\rho g \delta_0 - \partial_{t,\rho}^{-1} K_\rho g) \\ &= S_\rho(\Gamma_\rho g \chi_{\mathbb{R}_{\geq 0}} - \chi_{\mathbb{R}_{\geq 0}}(\mathfrak{m})M(\partial_{t,\rho})g + (M(\partial_{t,\rho})g)(0+) \chi_{\mathbb{R}_{\geq 0}}) \\ &= S_\lambda(\Gamma_\rho g \chi_{\mathbb{R}_{\geq 0}} - \chi_{\mathbb{R}_{\geq 0}}(\mathfrak{m})M(\partial_{t,\rho})g + (M(\partial_{t,\rho})g)(0+) \chi_{\mathbb{R}_{\geq 0}}), \end{aligned}$$

where we have used the independence of  $\rho$  stated in Theorem 2.6. Hence,

$$\begin{aligned} r_g(\lambda) &= \sqrt{2\pi}(\mathcal{L}_\lambda v)(0) \\ &= \lambda\sqrt{2\pi}(\mathcal{L}_\lambda \partial_{t,\lambda}^{-1} v)(0) \\ &= \lambda\sqrt{2\pi}(\lambda M(\lambda) + A)^{-1} \left( \frac{1}{\lambda\sqrt{2\pi}} \Gamma_\rho g - \mathcal{L}_\lambda(\chi_{\mathbb{R}_{\geq 0}}(\mathfrak{m})M(\partial_{t,\rho})g)(0) + \frac{1}{\lambda\sqrt{2\pi}} (M(\partial_{t,\rho})g)(0+) \right) \\ &= (\lambda M(\lambda) + A)^{-1} \left( \Gamma_\rho g + (M(\partial_{t,\rho})g)(0+) - \lambda\sqrt{2\pi}\mathcal{L}_\lambda(\chi_{\mathbb{R}_{\geq 0}}(\mathfrak{m})M(\partial_{t,\rho})g)(0) \right) \\ &= (\lambda M(\lambda) + A)^{-1} \left( (M(\partial_{t,\rho})g)(0-) - \lambda\sqrt{2\pi}\mathcal{L}_\lambda(\chi_{\mathbb{R}_{\geq 0}}(\mathfrak{m})M(\partial_{t,\rho})g)(0) \right) \end{aligned}$$

for each  $\lambda > \rho$ , where we have used the formula for  $\Gamma_\rho$  stated in Remark 5.2.  $\square$

With these preparations at hand, we can now state and prove the main result of this article.

**Theorem 6.7.** *Let  $\rho > s_0(M, A)$  and  $T^\rho$  be the semigroup on  $D_\rho$  associated with  $(M, A)$ . Moreover, for  $g \in \text{His}_\rho$  we set*

$$r_g(\lambda) := (\lambda M(\lambda) + A)^{-1} \left( (M(\partial_{t,\rho})g)(0-) - \lambda \sqrt{2\pi} \mathcal{L}_\lambda (\chi_{\mathbb{R}_{\geq 0}}(\mathfrak{m})M(\partial_{t,\rho})g)(0) \right) \quad (\lambda > \rho).$$

For  $\mu \leq \rho$  the following statements are equivalent:

(i)  $T^\rho$  can be extended to a  $C_0$ -semigroup on  $X_\rho^\mu = \overline{D_\rho}^{H \times L_{2,\mu}(\mathbb{R}; H)} \subseteq H \times L_{2,\mu}(\mathbb{R}; H)$ .

(ii) There exists  $M \geq 1$  and  $\omega \geq \rho$  such that

$$\frac{(\lambda - \omega)^{k+1}}{k!} \|r_g^{(k)}(\lambda)\| \leq M (\|g(0-)\|_H + \|g\|_{L_{2,\mu}(\mathbb{R}; H)})$$

for each  $\lambda > \omega, k \in \mathbb{N}$  and  $g \in \text{His}_\rho$ .

In this case

$$\begin{aligned} T_1^\rho : X_\rho^\mu &\rightarrow C_\omega(\mathbb{R}_{\geq 0}; H), \\ T_2^\rho : X_\rho^\mu &\rightarrow C_{\omega+\varepsilon}(\mathbb{R}_{\geq 0}; L_{2,\mu}(\mathbb{R}; H)) \end{aligned}$$

are bounded for each  $\varepsilon > 0$ .

*Proof.* (i)  $\Rightarrow$  (ii): Since  $T^\rho : X_\rho^\mu \rightarrow X_\rho^\mu$  is a  $C_0$ -semigroup, we find  $M \geq 1$  and  $\omega \geq \rho$  such that

$$\|T^\rho(t)\| \leq M e^{\omega t} \quad (t \geq 0).$$

In particular, we infer that

$$\|T_1^\rho(t)(g(0-), g)\| \leq M e^{\omega t} \|(g(0-), g)\|_{X_\rho^\mu} \quad (t \geq 0, g \in \text{His}_\rho).$$

Since  $r_g(\lambda) = \sqrt{2\pi} \mathcal{L}_\lambda (T_1^\rho(\cdot)(g(0-), g))(0)$  for  $\lambda > \omega$  by Proposition 6.6, we infer that

$$\begin{aligned} \|r_g^{(k)}(\lambda)\| &= \left\| \int_0^\infty e^{-\lambda t} (-t)^k T_1^\rho(t)(g(0-), g) dt \right\| \\ &\leq \int_0^\infty e^{-\lambda t} t^k M e^{\omega t} dt \|(g(0-), g)\|_{X_\rho^\mu} \\ &= M \frac{k!}{(\lambda - \omega)^{k+1}} \|(g(0-), g)\|_{X_\rho^\mu}, \end{aligned}$$

which shows (ii).

(ii)  $\Rightarrow$  (i): Let  $g \in \text{His}_\rho$  and define  $\tilde{r} : \mathbb{R}_{>0} \rightarrow H$  by  $\tilde{r}(\lambda) = r_g(\lambda + \omega)$  for  $\lambda > 0$ . Then  $\tilde{r}$  satisfies the assumptions of Theorem 6.4 and hence, there is  $f \in L_\infty(\mathbb{R}_{\geq 0}; H)$  with  $\|f\|_\infty \leq M (\|g(0-)\|_H + \|g\|_{L_{2,\mu}(\mathbb{R}; H)})$  such that

$$r_g(\lambda + \omega) = \int_0^\infty e^{-\lambda t} f(t) dt = \int_0^\infty e^{-(\lambda + \omega)t} e^{\omega t} f(t) dt$$

for each  $\lambda > 0$ . In particular, setting  $v := T_1^\rho(\cdot)(g(0-), g)$  we obtain

$$\int_0^\infty e^{-\lambda t} v(t) dt = r_g(\lambda) = \int_0^\infty e^{-\lambda t} e^{\omega t} f(t) dt \quad (\lambda > \omega)$$

and by analytic extension it follows that

$$\mathcal{L}_\lambda v = \mathcal{L}_\lambda(e^{\omega \cdot} f) \quad (\lambda > \omega).$$

Thus,  $v = e^{\omega \cdot} f$  and hence,

$$\|v(t)\| = e^{\omega t} \|f(t)\| \leq M e^{\omega t} (\|g(0-)\|_H + \|g\|_{L_{2,\mu}(\mathbb{R}; H)}).$$

Thus, since  $v$  is continuous on  $\mathbb{R}_{\geq 0}$ , we derive that

$$T_\rho^1 : D_\rho \subseteq X_\rho^\mu \rightarrow C_\omega(\mathbb{R}_{\geq 0}; H)$$

is bounded and can therefore be extended to a  $C_0$ -semigroup on  $X_\rho^\mu$ . Then, by Proposition 6.3 we obtain that

$$T_\rho^2 : D_\rho \subseteq X_\rho^\mu \rightarrow C_{\omega+\varepsilon}(\mathbb{R}_{\geq 0}; L_{2,\mu}(\mathbb{R}; H))$$

is also bounded for each  $\varepsilon > 0$  and hence, (i) follows.  $\square$

## 7 Applications

### 7.1 Differential-algebraic equations and classical Cauchy problems

In this section we consider initial value problems of the form

$$\begin{aligned} (\partial_{t,\rho} E + A) u &= 0 \quad \text{on } \mathbb{R}_{>0}, \\ u &= g \quad \text{on } \mathbb{R}_{\leq 0}, \end{aligned}$$

for a bounded operator  $E \in L(H)$ ,  $H$  a Hilbert space, and a densely defined linear and closed operator  $A : \text{dom}(A) \subseteq H \rightarrow H$ . We note that this corresponds to the abstract initial value problem (3) with

$$M(z) := E \quad (z \in \mathbb{C}).$$

We assume that the evolutionary problem is well-posed, that is we assume that there is  $\rho_1 \in \mathbb{R}_{\geq 0}$  such that  $zE + A$  is boundedly invertible for each  $z \in \mathbb{C}_{\text{Re} \geq \rho_1}$  and

$$\sup_{z \in \mathbb{C}_{\text{Re} \geq \rho_1}} \|(zE + A)^{-1}\| < \infty.$$

We again denote the infimum over all such  $\rho_1 \in \mathbb{R}_{\geq 0}$  by  $s_0(E, A)$ .

**Lemma 7.1.** *The function  $M$  given by  $M(z) := E$  for  $z \in \mathbb{C}$  is regularising. Moreover,  $\Gamma_\rho g = Eg(0-)$  and  $K_\rho g = 0$  for each  $g \in H_\rho^1(\mathbb{R}_{\leq 0}; H)$  and  $\rho \in \mathbb{R}_{> 0}$ . In particular, for  $\rho > s_0(E, A)$  we have that*

$$\text{IV}_\rho = \{x \in H; S_\rho(\delta Ex) - \chi_{\mathbb{R}_{\geq 0}}x \in H_\rho^1(\mathbb{R}; H)\}$$

and

$$\text{His}_\rho = \{g \in H_\rho^1(\mathbb{R}_{\leq 0}; H); g(0-) \in \text{IV}_\rho\}.$$

Moreover,  $X_\rho^\mu = \overline{\text{IV}_\rho} \times L_{2,\mu}(\mathbb{R}_{\leq 0}; H)$  for each  $\mu \leq \rho$ .

*Proof.* For  $x \in H$ ,  $\rho > 0$  we have

$$M(\partial_{t,\rho})\chi_{\mathbb{R}_{\geq 0}}x = \chi_{\mathbb{R}_{\geq 0}}Ex$$

and thus,  $M$  is regularising with  $\Gamma_\rho g = Eg(0-)$  for each  $g \in H_\rho^1(\mathbb{R}_{\leq 0}; H)$ . Moreover, we have

$$\begin{aligned} K_\rho g &= P_0 \partial_{t,\rho} Eg \\ &= \partial_{t,\rho} \chi_{\mathbb{R}_{\geq 0}}(m)Eg - \delta_0(Eg)(0+) \\ &= 0. \end{aligned}$$

Hence, for  $g \in H_\rho^1(\mathbb{R}_{\leq 0}; H)$ ,  $\rho > s_0(E, A)$ , we have

$$\begin{aligned} g \in \text{His}_\rho &\Leftrightarrow S_\rho(\delta_0 Eg(0-)) + g \in H_\rho^1(\mathbb{R}; H) \\ &\Leftrightarrow S_\rho(\delta_0 Eg(0-)) - \chi_{\mathbb{R}_{> 0}}g(0-) + g + \chi_{\mathbb{R}_{> 0}}g(0-) \in H_\rho^1(\mathbb{R}; H) \\ &\Leftrightarrow S_\rho(\delta_0 Eg(0-)) - \chi_{\mathbb{R}_{> 0}}g(0-) \in H_\rho^1(\mathbb{R}; H) \end{aligned}$$

which proves the asserted equalities for  $\text{His}_\rho$  and  $\text{IV}_\rho$ .

Finally, let  $x \in \overline{\text{IV}_\rho}$  and  $g \in L_{2,\mu}(\mathbb{R}_{\leq 0}; H)$  for some  $\mu \leq \rho$  with  $\rho > s_0(E, A)$ . Then we find a sequence  $(x_n)_{n \in \mathbb{N}}$  in  $\text{IV}_\rho$  and a sequence  $(\varphi_n)_{n \in \mathbb{N}}$  in  $C_c^\infty(\mathbb{R}_{< 0}; H)$  such that  $x_n \rightarrow x$  and  $\varphi_n \rightarrow g$  in  $H$  and  $L_{2,\mu}(\mathbb{R}_{\leq 0}; H)$ , respectively. Moreover, we set

$$\psi_n(t) := \begin{cases} (nt + 1)x_n & \text{if } -\frac{1}{n} \leq t \leq 0, \\ 0 & \text{else} \end{cases} \quad (t \in \mathbb{R}_{\leq 0}, n \in \mathbb{N})$$

and obtain a sequence  $(\psi_n)_{n \in \mathbb{N}}$  in  $H_\rho^1(\mathbb{R}_{\leq 0}; H)$  with  $\psi_n(0-) = x_n$  for  $n \in \mathbb{N}$  and  $\psi_n \rightarrow 0$  as  $n \rightarrow \infty$  in  $L_{2,\mu}(\mathbb{R}_{\leq 0}; H)$ . Consequently, setting  $g_n := \psi_n + \varphi_n \in H_\rho^1(\mathbb{R}_{\leq 0}; H)$  for  $n \in \mathbb{N}$  we obtain a sequence  $(x_n, g_n)_{n \in \mathbb{N}}$  in  $D_\rho$  with  $(x_n, g_n) \rightarrow (x, g)$  in  $H \times L_{2,\mu}(\mathbb{R}; H)$  and thus,  $(x, g) \in X_\rho^\mu$ . Since the other inclusion holds obviously, this proves the assertion.  $\square$

We now inspect the space  $\text{IV}_\rho$  a bit closer. In particular, we are able to determine its closure  $\overline{\text{IV}_\rho}$  and a suitable dense subset of  $\overline{\text{IV}_\rho}$ .

**Proposition 7.2.** *We set*

$$U := \{x \in \text{dom}(A); \exists y \in \text{dom}(A) : Ax = Ey\}.$$

*Then  $U \subseteq \text{IV}_\rho$  and  $\overline{U} = \overline{\text{IV}_\rho}$  for each  $\rho > s_0(E, A)$ . In particular,  $\overline{\text{IV}_\rho}$  does not depend on the particular choice of  $\rho > s_0(E, A)$ .*

*Proof.* Let  $\rho > s_0(E, A)$ ,  $x \in U$  and  $y \in \text{dom}(A)$  with  $Ax = Ey$ . Then we compute

$$\begin{aligned} S_\rho(\delta Ex) - \chi_{\mathbb{R}_{\geq 0}}x &= (\partial_{t,\rho}E + A)^{-1}(\delta Ex - \delta Ex - \chi_{\mathbb{R}_{\geq 0}}Ax) \\ &= -(\partial_{t,\rho}E + A)^{-1}(\chi_{\mathbb{R}_{\geq 0}}Ey) \\ &= -(\partial_{t,\rho}E + A)^{-1}(\partial_{t,\rho}E\partial_{t,\rho}^{-1}\chi_{\mathbb{R}_{\geq 0}}y) \\ &= -\partial_{t,\rho}^{-1}\chi_{\mathbb{R}_{\geq 0}}y + (\partial_{t,\rho}E + A)^{-1}(\partial_{t,\rho}^{-1}\chi_{\mathbb{R}_{\geq 0}}Ay) \in H_\rho^1(\mathbb{R}; H), \end{aligned}$$

which shows that  $x \in \text{IV}_\rho$  by Lemma 7.1. For showing the remaining assertion, we prove that  $\text{IV}_\rho \subseteq \overline{U}$ . For doing so, let  $x \in \text{IV}_\rho$  and set  $v := S_\rho(\delta Ex)$ . Then

$$\begin{aligned} \partial_{t,\rho}E(v - \chi_{\mathbb{R}_{\geq 0}}x) &= (\partial_{t,\rho}E + A)v - \delta Ex - Av \\ &= -Av, \end{aligned}$$

and since the left-hand side belongs to  $L_{2,\rho}(\mathbb{R}; H)$  we infer that  $v \in L_{2,\rho}(\mathbb{R}; \text{dom}(A))$ . Hence,  $\partial_{t,\rho}^{-1}v \in H_\rho^1(\mathbb{R}; \text{dom}(A)) \hookrightarrow C_\rho(\mathbb{R}; \text{dom}(A))$  and so  $\int_0^t v(s) ds = (\partial_{t,\rho}^{-1}v)(t) \in \text{dom}(A)$  for each  $t \geq 0$  and

$$A \int_0^t v(s) ds = Ev(t) - Ex \quad (t \geq 0).$$

Consequently,

$$\int_0^t v(s) ds \in A^{-1}[\text{ran}(E)] \quad (t \geq 0)$$

and since  $v$  is continuous on  $\mathbb{R}_{\geq 0}$  and hence,  $\frac{1}{t} \int_0^t v(s) ds \rightarrow v(0+) = x$  as  $t \rightarrow 0$ , it suffices to prove  $A^{-1}[\text{ran}(E)] \subseteq \overline{U}$ . For doing so, let  $y \in A^{-1}[\text{ran}(E)]$ , i.e.,  $y \in \text{dom}(A)$  and  $Ay = Ez$  for some  $z \in H$ . We choose a sequence  $(z_n)_{n \in \mathbb{N}}$  in  $\text{dom}(A)$  with  $z_n \rightarrow z$  as  $n \rightarrow \infty$  and define

$$y_n := (\lambda E + A)^{-1}(\lambda Ey + Ez_n) \quad (n \in \mathbb{N}),$$

where  $\lambda > s_0(E, A)$  is fixed. Then  $y_n \in U$ , since

$$Ay_n = A(\lambda E + A)^{-1}(\lambda Ey + Ez_n) = E(\lambda E + A)^{-1}(\lambda Ay + Az_n) \in E[\text{dom}(A)]$$

and since  $Ez_n \rightarrow Ez = Ay$ , we infer that  $y_n \rightarrow y$  and hence,  $y \in \overline{U}$ .  $\square$

**Theorem 7.3.** *Let  $M(z) := E$  for  $z \in \mathbb{C}$ ,  $\rho > s_0(E, A)$  and let  $T^\rho : D_\rho \subseteq \text{IV}_\rho \times \text{His}_\rho \rightarrow \text{IV}_\rho \times \text{His}_\rho$  denote the semigroup associated with  $(M, A)$ . Moreover, for  $x \in H$  we define*

$$f_x(t) := \begin{cases} (t+1)x & \text{if } t \in [-1, 0], \\ 0 & \text{else} \end{cases} \quad (t \in \mathbb{R}_{\leq 0}).$$

*Then the following statements are equivalent:*

(i)  $T^\rho$  extends to a  $C_0$ -semigroup in  $\overline{IV}_\rho \times L_{2,\mu}(\mathbb{R}_{\leq 0}; H)$  for some  $\mu \leq \rho$ .

(ii) There exists  $M \geq 1$  and  $\omega \geq \rho$  such that

$$\|((\lambda E + A)^{-1}E)^n\| \leq \frac{M}{(\lambda - \omega)^n} \quad (\lambda > \omega, n \in \mathbb{N}). \quad (5)$$

(iii)  $T^\rho$  extends to a  $C_0$ -semigroup in  $\overline{IV}_\rho \times L_{2,\mu}(\mathbb{R}_{\leq 0}; H)$  for each  $\mu \leq \rho$ .

(iv) The family of functions

$$S^\rho(t) : IV_\rho \subseteq \overline{IV}_\rho \rightarrow \overline{IV}_\rho, \quad x \mapsto T_1^\rho(t)(x, f_x)$$

for  $t \geq 0$  extends to a  $C_0$ -semigroup on  $\overline{IV}_\rho$ .

In the latter case,  $S^\rho(t)x = T^\rho(t)(x, 0)$  for each  $x \in \overline{IV}_\rho$  and  $t \geq 0$ .

*Proof.* We first compute the function  $r_g$  for  $g \in \text{His}_\rho$  as it was defined in Theorem 6.7. We have that

$$(M(\partial_{t,\rho})g)(0-) - \lambda\sqrt{2\pi}\mathcal{L}_\lambda(\chi_{\mathbb{R}_{\geq 0}}(\mathfrak{m})M(\partial_{t,\rho})g)(0) = Eg(0-) \quad (\lambda > \rho)$$

and hence,

$$r_g(\lambda) = (\lambda E + A)^{-1}Eg(0-) \quad (\lambda > \rho).$$

Consequently,

$$r_g^{(k)}(\lambda) = (-1)^k k! \left( (\lambda E + A)^{-1}E \right)^{k+1} g(0-) \quad (k \in \mathbb{N}_0, \lambda > \rho).$$

(i)  $\Rightarrow$  (ii): By Theorem 6.7 (note that  $X_\rho^\mu = \overline{IV}_\rho \times L_{2,\mu}(\mathbb{R}_{\leq 0}; H)$  by Lemma 7.1) we know that there exists  $M \geq 1$  and  $\omega \geq \rho$  such that

$$\frac{(\lambda - \omega)^{k+1}}{k!} \|r_g^{(k)}(\lambda)\| \leq M (\|g(0-)\|_H + \|g\|_{L_{2,\mu}(\mathbb{R}; H)})$$

for each  $\lambda > \omega, k \in \mathbb{N}$  and  $g \in \text{His}_\rho$ . Choosing now  $x \in \overline{IV}_\rho$  we infer that

$$\begin{aligned} \|((\lambda E + A)^{-1}E)^n x\| &= \frac{1}{(n-1)!} \|r_{f_x(k\cdot)}^{(n-1)}(\lambda)\| \\ &\leq \frac{M}{(\lambda - \omega)^n} (\|x\|_H + \|f_x(k\cdot)\|_{L_{2,\mu}(\mathbb{R}; H)}) \end{aligned}$$

for each  $\lambda > \omega, n, k \in \mathbb{N}$ . Since  $f_x(k\cdot) \rightarrow 0$  as  $k \rightarrow \infty$ , we infer that

$$\|((\lambda E + A)^{-1}E)^n\| \leq \frac{M}{(\lambda - \omega)^n} \quad (\lambda > \omega, n \in \mathbb{N}).$$

(ii)  $\Rightarrow$  (iii): Let  $\mu \leq \rho$ . By assumption, there exists  $M \geq 1, \omega \geq \rho$  such that

$$\begin{aligned} \frac{(\lambda - \omega)^{k+1}}{k!} \|r_g^{(k)}(\lambda)\| &= (\lambda - \omega)^{k+1} \|((\lambda E + A)^{-1} E)^{k+1} g(0-)\| \\ &\leq M \|g(0-)\|_H \\ &\leq M (\|g(0-)\| + \|g\|_{L_{2,\mu}(\mathbb{R};H)}) \end{aligned}$$

for each  $\lambda > \omega, k \in \mathbb{N}_0$  and  $g \in \text{His}_\rho$  and hence, the assertion follows from Theorem 6.7 and Lemma 7.1.

(iii)  $\Rightarrow$  (iv): Since  $T^\rho$  extends to a  $C_0$ -semigroup on  $\overline{\text{IV}_\rho} \times L_2(\mathbb{R}_{\leq 0}; H)$  and since

$$\|f_x\|_{L_2(\mathbb{R};H)} \leq \|x\|_H \quad (x \in H),$$

we infer that there is  $M \geq 1$  and  $\omega \in \mathbb{R}$  such that

$$\|S^\rho(t)x\| \leq 2Me^{\omega t} \|x\| \quad (x \in \text{IV}_\rho)$$

and thus,  $(S^\rho(t))_{t \geq 0}$  extends to a  $C_0$ -semigroup on  $\overline{\text{IV}_\rho}$ . Moreover, since

$$\begin{aligned} S^\rho(t)x &= T_1^\rho(t)(x, f_x) \\ &= ((\partial_{t,\rho} E + A)^{-1} (\delta_0 E x)) (t) \\ &= T_1^\rho(t)(x, 0) \end{aligned}$$

for each  $t \geq 0, x \in \overline{\text{IV}_\rho}$ , we obtain the at the end asserted formula .

(iv)  $\Rightarrow$  (i): By assumption, there is  $M \geq 1, \omega \in \mathbb{R}$  such that

$$\|T_1^\rho(t)(x, f_x)\| \leq Me^{\omega t} \|x\| \quad (x \in \text{IV}_\rho, t \geq 0).$$

Moreover, since

$$T_1^\rho(t)(x, g) = T_1^\rho(t)(x, f_x) \quad ((x, g) \in D_\rho),$$

we infer that

$$T_1^\rho : D_\rho \subseteq \overline{\text{IV}_\rho} \times L_{2,\mu}(\mathbb{R}_{\leq 0}; H) \rightarrow C_\omega(\mathbb{R}_{\geq 0}; H)$$

is continuous and hence, the assertion follows by Proposition 6.3.  $\square$

*Remark 7.4.* We remark that in the case of classical Cauchy problems, i.e.  $E = 1$ , condition (5) is nothing but the classical Hille-Yosida condition for generators of  $C_0$ -semigroups (see e.g. [5, Chapter II, Theorem 3.8]). Note that in this case,  $U = \text{dom}(A^2)$  in Proposition 7.2 and hence,  $\overline{\text{IV}_\rho} = \overline{U} = H$ .

## 7.2 A hyperbolic delay equation

As a slight generalisation of [3, Example 3.17] we consider a concrete delay equation of the form

$$\begin{aligned} \partial_{t,\rho}^2 u - \text{div } k \text{ grad } u - \sum_{i=1}^n c_i \tau_{-h_i} \partial_i u - c_0 \tau_{-h_0} \partial_{t,\rho} u &= 0 \quad \text{on } \mathbb{R}_{>0}, \\ u &= g \quad \text{on } \mathbb{R}_{<0}. \end{aligned} \tag{6}$$

Here,  $u$  attains values in  $L_2(\Omega)$  for some open set  $\Omega \subseteq \mathbb{R}^n$  as underlying domain,  $h_0, \dots, h_n > 0$  are given real numbers and  $k, c_0, \dots, c_n$  are bounded operators on  $L_2(\Omega)^n$  and  $L_2(\Omega)$ , respectively. The operators  $\text{grad}$  and  $\text{div}$  denote the usual gradient and divergence with respect to the spatial variables and will be introduced rigorously later. It is our first goal to rewrite this equation as a suitable evolutionary problem. For doing so, we need the following definition.

**Definition.** Let  $c_0, \dots, c_n \in L(L_2(\Omega))$  and  $k \in L(L_2(\Omega)^n)$  selfadjoint such that  $k \geq d$  for some  $d \in \mathbb{R}_{>0}$ . We define the function  $M_1 : \mathbb{C} \rightarrow L(L_2(\Omega) \times L_2(\Omega)^n; L_2(\Omega))$  by

$$M_1(z)q := c_0 e^{-h_0 z} q_0 - \sum_{i=1}^n c_i k^{-1} e^{-h_i z} q_i \quad (z \in \mathbb{C}, q \in L_2(\Omega) \times L_2(\Omega)^n).$$

Furthermore, we define  $M : \mathbb{C} \setminus \{0\} \rightarrow L(L_2(\Omega) \times L_2(\Omega)^n)$  by

$$M(z) \begin{pmatrix} v \\ q \end{pmatrix} := \begin{pmatrix} v + z^{-1} M_1(z)q \\ k^{-1}q \end{pmatrix}.$$

*Remark 7.5.* Since  $(\mathcal{L}_\rho \tau_h u)(t) = e^{(it+\rho)h} (\mathcal{L}_\rho u)(t)$  for each  $u \in L_{2,\rho}(\mathbb{R}; H)$  and  $t, h \in \mathbb{R}$ , we have that

$$M_1(\partial_{t,\rho})q = c_0 \tau_{-h_0} q_0 - \sum_{i=1}^n c_i k^{-1} \tau_{-h_i} q_i$$

for each  $q \in L_{2,\rho}(\mathbb{R}; L_2(\Omega)^n)$ .

Obviously, the so defined function  $M$  is analytic and if we restrict it to some open half plane  $\mathbb{C}_{\text{Re} > \rho_0}$  with  $\rho_0 > 0$ , it is bounded. Thus, we may consider the operator  $M(\partial_{t,\rho})$  for some  $\rho > 0$ .

**Lemma 7.6.** *The function  $M$  is regularising.*

*Proof.* We need to prove that  $(M(\partial_{t,\rho})\chi_{\mathbb{R}_{\geq 0}}x)(0+)$  exists for all  $x = (\tilde{x}, \hat{x}) \in L_2(\Omega) \times L_2(\Omega)^n$  and  $\rho > 0$ . We have that

$$M(\partial_{t,\rho})\chi_{\mathbb{R}_{\geq 0}}x = \begin{pmatrix} \chi_{\mathbb{R}_{\geq 0}}\tilde{x} + \partial_{t,\rho}^{-1}M_1(\partial_{t,\rho})\chi_{\mathbb{R}_{\geq 0}}x \\ k^{-1}\hat{x} \end{pmatrix}$$

and since  $M_1(\partial_{t,\rho})$  is causal, we infer that  $\partial_{t,\rho}^{-1}M_1(\partial_{t,\rho})\chi_{\mathbb{R}_{\geq 0}}x \in H_\rho^1(\mathbb{R}; L_2(\Omega))$  is supported on  $\mathbb{R}_{>0}$  and hence,  $(\partial_{t,\rho}^{-1}M_1(\partial_{t,\rho})\chi_{\mathbb{R}_{\geq 0}}x)(0+) = 0$ . Thus,  $M$  is regularising.  $\square$

We now rewrite (6) as an evolutionary equation. We introduce  $v := \partial_{t,\rho}u$  and  $q := k \text{grad } u$  as new unknowns, and rewrite (6) as

$$\left( \partial_{t,\rho} M(\partial_{t,\rho}) + \begin{pmatrix} 0 & \text{div} \\ \text{grad} & 0 \end{pmatrix} \right) \begin{pmatrix} v \\ q \end{pmatrix} = 0 \quad \text{on } \mathbb{R}_{>0}. \quad (7)$$

Of course (6) has to be completed by suitable boundary conditions. This will be done by introducing the differential operators  $\text{div}$  and  $\text{grad}$  in a suitable way.



**Definition.** We define  $\text{grad}_0 : \text{dom}(\text{grad}_0) \subseteq L_2(\Omega) \rightarrow L_2(\Omega)^n$  as the closure of the operator

$$C_c^\infty(\Omega) \subseteq L_2(\Omega) \rightarrow L_2(\Omega)^n, \varphi \mapsto (\partial_j \varphi)_{j \in \{1, \dots, n\}}$$

and similarly  $\text{div}_0 : \text{dom}(\text{div}_0) \subseteq L_2(\Omega)^n \rightarrow L_2(\Omega)$  as the closure of

$$C_c^\infty(\Omega)^n \subseteq L_2(\Omega)^n \rightarrow L_2(\Omega), (\varphi_j)_{j \in \{1, \dots, n\}} \mapsto \sum_{j=1}^n \partial_j \varphi_j.$$

Moreover, we set

$$\begin{aligned} \text{grad} &:= -(\text{div}_0)^* \\ \text{div} &:= -(\text{grad}_0)^*. \end{aligned}$$

*Remark 7.7.* We note that  $\text{dom}(\text{grad}_0)$  coincides with the classical Sobolev space  $H_0^1(\Omega)$  of weakly differentiable  $L_2$ -functions with vanishing Dirichlet trace. Moreover,  $\text{dom}(\text{grad})$  is nothing but the Sobolev space  $H^1(\Omega)$ . Thus, by Green's formula, elements in  $\text{dom}(\text{div}_0)$  may be interpreted as elements in  $L_2(\Omega)^n$  with distributional divergence also lying in  $L_2(\Omega)$  and whose normal trace vanishes, while elements in  $\text{dom}(\text{div})$  are just  $L_2(\Omega)$  vector fields with  $L_2(\Omega)$ -divergence. Note however, that these definitions are meaningful for arbitrary open sets  $\Omega$  and do not require any boundary regularity.

Thus, by replacing  $\text{div}$  by  $\text{div}_0$  or  $\text{grad}$  by  $\text{grad}_0$  in (7), we can model homogeneous Neumann- or Dirichlet conditions, respectively.

**Lemma 7.8.** *We set*

$$A_N := \begin{pmatrix} 0 & \text{div}_0 \\ \text{grad} & 0 \end{pmatrix} : \text{dom}(\text{grad}) \times \text{dom}(\text{div}_0) \subseteq L_2(\Omega) \times L_2(\Omega)^n \rightarrow L_2(\Omega) \times L_2(\Omega)^n$$

and

$$A_D := \begin{pmatrix} 0 & \text{div} \\ \text{grad}_0 & 0 \end{pmatrix} : \text{dom}(\text{grad}_0) \times \text{dom}(\text{div}) \subseteq L_2(\Omega) \times L_2(\Omega)^n \rightarrow L_2(\Omega) \times L_2(\Omega)^n.$$

Then both operators are skew-selfadjoint, i.e.  $A_N^* = -A_N$  and  $A_D^* = -A_D$ .

*Proof.* The claim follows immediately by the definitions of the differential operators.  $\square$

We now prove that the evolutionary problems associated with  $(M, A_{D/N})$  are well-posed.

**Proposition 7.9.** *Let  $c_0, \dots, c_n \in L(L_2(\Omega))$  and  $k \in L(L_2(\Omega)^n)$  selfadjoint such that  $k \geq d$  for some  $d \in \mathbb{R}_{>0}$ . Then the evolutionary problems associated with  $(M, A_{D/N})$  are well-posed.*

*Proof.* We first note that  $k^{-1}$  is selfadjoint and satisfies  $k^{-1} \geq \frac{1}{\|k\|}$ . Moreover, since  $A_{D/N}$  is skew-selfadjoint, we infer that

$$\text{Re}\langle A_{D/N}x, x \rangle = 0 \quad (x \in \text{dom}(A_{D/N})).$$

Hence, we may estimate for  $x = (x_1, x_2) \in \text{dom}(A_{D/N})$

$$\begin{aligned} \text{Re}\langle (zM(z) + A_{D/N})x, x \rangle &= \text{Re}\langle zM(z)x, x \rangle \\ &= \text{Re}\langle zx_1, x_1 \rangle + \text{Re}\langle zk^{-1}x_2, x_2 \rangle + \text{Re}\langle M_1(z)x, x_1 \rangle \\ &\geq \text{Re } z \min\left\{1, \frac{1}{\|k\|}\right\} \|x\|^2 - \|M_1(z)\| \|x\|^2. \end{aligned}$$

Moreover, we estimate

$$\|M_1(z)\| \leq \|c_0\| e^{-h_0 \text{Re } z} + \sum_{i=1}^n \|c_i\| \|k^{-1}\| e^{-h_i \text{Re } z}$$

and hence, we infer that  $\|M_1(z)\| \rightarrow 0$  as  $\text{Re } z \rightarrow \infty$ . Thus, we find  $c > 0$  and  $\rho_0 > 0$  such that

$$\text{Re}\langle (zM(z) + A_{D/N})x, x \rangle \geq c \|x\|^2 \quad (z \in \mathbb{C}_{\text{Re} \geq \rho_0}),$$

which yields the well-posedness for the evolutionary problem associated with  $(M, A_{D/N})$ .  $\square$

*Remark 7.10.* We note that the above proof also works for  $m$ -accretive operators  $A$  instead of  $A_{D/N}$ . This allows for the treatment of more general boundary conditions and we refer to [20] for a characterisation result about those boundary conditions (including also nonlinear ones).

Having these results at hand, we are now in the position to consider the history space for (7). From now on, to avoid cluttered notation, we will simply write  $A$  and note that  $A$  can be replaced by  $A_N$  and  $A_D$ , respectively.

**Proposition 7.11.** *Let  $\rho > s_0(M, A)$ . Then*

$$\Gamma_\rho g = \begin{pmatrix} 1 & 0 \\ 0 & k^{-1} \end{pmatrix} g(0-), \quad K_\rho g = \chi_{\mathbb{R}_{\geq 0}(\text{m})} \begin{pmatrix} M_1(\partial_{t,\rho}) \\ 0 \end{pmatrix} g$$

for each  $g \in H_\rho^1(\mathbb{R}_{\leq 0}; L_2(\Omega) \times L_2(\Omega)^n)$ . Moreover,

$$\left\{ g \in H_\rho^1(\mathbb{R}_{\leq 0}; \text{dom}(A)); \forall j \in \{0, \dots, n\} : g(-t_j) = 0, \begin{pmatrix} 1 & 0 \\ 0 & k \end{pmatrix} Ag(0-) \in \text{dom}(A) \right\} \subseteq \text{His}_\rho \quad (8)$$

and consequently,

$$X_\rho^\mu = (L_2(\Omega) \times L_2(\Omega)^n) \times L_{2,\mu}(\mathbb{R}_{\leq 0}; L_2(\Omega) \times L_2(\Omega)^n)$$

for each  $\mu \leq \rho$ .

*Proof.* Let  $g \in H_\rho^1(\mathbb{R}_{\leq 0}; L_2(\Omega) \times L_2(\Omega)^n)$ . Then

$$\begin{aligned}\Gamma_\rho g &= \left( \left( \begin{pmatrix} 1 & 0 \\ 0 & k^{-1} \end{pmatrix} + \partial_{t,\rho}^{-1} \begin{pmatrix} M_1(\partial_{t,\rho}) \\ 0 \end{pmatrix} \right) \chi_{\mathbb{R}_{\geq 0}} g(0-) \right) (0+) \\ &= \left( \begin{pmatrix} 1 & 0 \\ 0 & k^{-1} \end{pmatrix} \chi_{\mathbb{R}_{\geq 0}} g(0-) \right) (0+) \\ &= \begin{pmatrix} 1 & 0 \\ 0 & k^{-1} \end{pmatrix} g(0-),\end{aligned}$$

where we have used

$$\partial_{t,\rho}^{-1} \begin{pmatrix} M_1(\partial_{t,\rho}) \\ 0 \end{pmatrix} \chi_{\mathbb{R}_{\geq 0}} g(0-) \in H_\rho^1(\mathbb{R}; H)$$

and hence

$$\left( \partial_{t,\rho}^{-1} \begin{pmatrix} M_1(\partial_{t,\rho}) \\ 0 \end{pmatrix} \chi_{\mathbb{R}_{\geq 0}} g(0-) \right) (0+) = \left( \partial_{t,\rho}^{-1} \begin{pmatrix} M_1(\partial_{t,\rho}) \\ 0 \end{pmatrix} \chi_{\mathbb{R}_{\geq 0}} g(0-) \right) (0-) = 0$$

by causality. Moreover,

$$\begin{aligned}K_\rho g &= P_0 \partial_{t,\rho} M(\partial_{t,\rho}) g \\ &= P_0 \partial_{t,\rho} \begin{pmatrix} 1 & 0 \\ 0 & k^{-1} \end{pmatrix} g + \chi_{\mathbb{R}_{\geq 0}}(\mathbf{m}) \begin{pmatrix} M_1(\partial_{t,\rho}) \\ 0 \end{pmatrix} g \\ &= \chi_{\mathbb{R}_{\geq 0}}(\mathbf{m}) \begin{pmatrix} M_1(\partial_{t,\rho}) \\ 0 \end{pmatrix} g,\end{aligned}$$

since

$$\text{spt } \partial_{t,\rho} \begin{pmatrix} 1 & 0 \\ 0 & k^{-1} \end{pmatrix} g \subseteq \mathbb{R}_{\leq 0}$$

and thus,  $P_0 \partial_{t,\rho} \begin{pmatrix} 1 & 0 \\ 0 & k^{-1} \end{pmatrix} g = 0$  by Proposition 4.3 (c). Let now  $g$  be an element of the set on the left hand side of (8). Then, we compute

$$\begin{aligned}& S_\rho \left( \delta_0 \begin{pmatrix} 1 & 0 \\ 0 & k^{-1} \end{pmatrix} g(0-) - \chi_{\mathbb{R}_{\geq 0}}(\mathbf{m}) \begin{pmatrix} M_1(\partial_{t,\rho}) \\ 0 \end{pmatrix} g \right) - \chi_{\mathbb{R}_{\geq 0}} g(0-) \\ &= S_\rho \left( \delta_0 \begin{pmatrix} 1 & 0 \\ 0 & k^{-1} \end{pmatrix} g(0-) - \chi_{\mathbb{R}_{\geq 0}}(\mathbf{m}) \begin{pmatrix} M_1(\partial_{t,\rho}) \\ 0 \end{pmatrix} g - \right. \\ &\quad \left. - \partial_{t,\rho} \begin{pmatrix} 1 & 0 \\ 0 & k^{-1} \end{pmatrix} \chi_{\mathbb{R}_{\geq 0}} g(0-) - \begin{pmatrix} M_1(\partial_{t,\rho}) \\ 0 \end{pmatrix} \chi_{\mathbb{R}_{\geq 0}} g(0-) - \chi_{\mathbb{R}_{\geq 0}} A g(0-) \right) \\ &= -S_\rho \left( \chi_{\mathbb{R}_{\geq 0}}(\mathbf{m}) \begin{pmatrix} M_1(\partial_{t,\rho}) \\ 0 \end{pmatrix} (g + \chi_{\mathbb{R}_{\geq 0}} g(0-)) \right) - S_\rho (\chi_{\mathbb{R}_{\geq 0}} A g(0-)).\end{aligned}$$

We now treat both terms separately. We note that

$$(\chi_{\mathbb{R}_{\geq 0}}(\mathbf{m}) c_j \tau_{-h_j} f_j)(t) = \begin{cases} c_j f_j(t - h_j) & \text{if } t \geq 0, \\ 0 & \text{otherwise} \end{cases}$$

for  $f_j \in H_\rho^1(\mathbb{R}; L_2(\Omega))$  and thus,  $\chi_{\mathbb{R}_{\geq 0}}(\mathfrak{m})c_j\tau_{-h_j}f_j \in H_\rho^1(\mathbb{R}; L_2(\Omega))$  if  $f_j(-h_j) = 0$ . Thus, by the constraints on  $g$ , we infer that

$$\chi_{\mathbb{R}_{\geq 0}}(\mathfrak{m}) \begin{pmatrix} M_1(\partial_{t,\rho}) \\ 0 \end{pmatrix} (g + \chi_{\mathbb{R}_{\geq 0}}g(0-)) \in H_\rho^1(\mathbb{R}; L_2(\Omega) \times L_2(\Omega)^n).$$

Thus, we are left to consider the last term. By assumption, we find  $x \in \text{dom}(A)$  with  $Ag(0-) = \begin{pmatrix} 1 & 0 \\ 0 & k^{-1} \end{pmatrix} x$  and hence,

$$\begin{aligned} S_\rho(\chi_{\mathbb{R}_{\geq 0}}Ag(0-)) &= \partial_{t,\rho}^{-1}S_\rho \left( \partial_{t,\rho} \begin{pmatrix} 1 & 0 \\ 0 & k^{-1} \end{pmatrix} \chi_{\mathbb{R}_{\geq 0}}x \right) \\ &= \partial_{t,\rho}^{-1} \left( \chi_{\mathbb{R}_{\geq 0}}x - S_\rho \left( \begin{pmatrix} M_1(\partial_{t,\rho}) \\ 0 \end{pmatrix} + A \right) \chi_{\mathbb{R}_{\geq 0}}x \right) \\ &\in H_\rho^1(\mathbb{R}; L_2(\Omega) \times L_2(\Omega)^n), \end{aligned}$$

which proves the claim.  $\square$

We conclude this section by proving that the associated semigroup can be extended to  $X_\rho^\mu$  for each  $\mu \leq \rho$ .

**Theorem 7.12.** *Let  $\rho > s_0(M, A)$  and let  $T^\rho$  denote the associated semigroup with  $(M, A)$  on  $D_\rho$ . Then for large enough  $\rho$ ,  $T^\rho$  extends to a  $C_0$ -semigroup on  $(L_2(\Omega) \times L_2(\Omega)^n) \times L_{2,\mu}(\mathbb{R}_{\leq 0}; L_2(\Omega) \times L_2(\Omega)^n)$  for each  $\mu \leq \rho$ .*

*Proof.* The proof will be done by a perturbation argument. For doing so, we consider the evolutionary problem associated with  $(E, A)$ , where

$$E := \begin{pmatrix} 1 & 0 \\ 0 & k^{-1} \end{pmatrix}.$$

We note that this problem is well-posed with  $s_0(E, A) = 0$  (compare the proof of Proposition 7.9). We denote the associated semigroup by  $\tilde{T}^\rho$ . By Proposition 7.2 we know that the closure of the initial value space for  $\tilde{T}^\rho$  is given by

$$\overline{\{x \in \text{dom}(A); E^{-1}Ax \in \text{dom}(A)\}} = L_2(\Omega) \times L_2(\Omega)^n.$$

Moreover, by Theorem 7.3  $\tilde{T}^\rho$  extends to a  $C_0$ -semigroup on  $(L_2(\Omega) \times L_2(\Omega)^n) \times L_{2,\mu}(\mathbb{R}_{\leq 0}; L_2(\Omega) \times L_2(\Omega)^n)$  if and only if

$$\|((\lambda E + A)^{-1}E)^n\| \leq \frac{M}{(\lambda - \omega)^n} \quad (\lambda > \omega, n \in \mathbb{N})$$

for some  $M \geq 1$ ,  $\omega \geq \rho$ . We note that  $E$  is selfadjoint and strictly positive definite and thus,

$$(\lambda E + A)^{-1} = \sqrt{E^{-1}} \left( \lambda + \sqrt{E^{-1}}A\sqrt{E^{-1}} \right)^{-1} \sqrt{E^{-1}}.$$

The latter gives

$$((\lambda E + A)^{-1}E)^n = \sqrt{E^{-1}} \left( \lambda + \sqrt{E^{-1}}A\sqrt{E^{-1}} \right)^{-n} \sqrt{E}$$

for each  $n \in \mathbb{N}$ . Since  $A$  is skew-selfadjoint, so is  $\sqrt{E^{-1}}A\sqrt{E^{-1}}$  and thus,

$$\|((\lambda E + A)^{-1}E)^n\| \leq \frac{\|\sqrt{E}\| \|\sqrt{E^{-1}}\|}{\lambda^n} \quad (\lambda > 0, n \in \mathbb{N})$$

and hence,  $\tilde{T}^\rho$  extends to a bounded  $C_0$ -semigroup on  $(L_2(\Omega) \times L_2(\Omega)^n) \times L_{2,\mu}(\mathbb{R}_{\leq 0}; L_2(\Omega) \times L_2(\Omega)^n)$ . Now we come to the semigroup  $T^\rho$ . We will prove that  $T_1^\rho : D_\rho \subseteq (L_2(\Omega) \times L_2(\Omega)^n) \times L_{2,\mu}(\mathbb{R}_{\leq 0}; L_2(\Omega) \times L_2(\Omega)^n) \rightarrow C_\omega(\mathbb{R}_{\geq 0}; L_2(\Omega) \times L_2(\Omega)^n)$  is bounded for some  $\omega \in \mathbb{R}$ , which would imply the claim by Proposition 6.3. Let  $(g(0-), g) \in D_\rho$ . We then have, using the formulas in Proposition 7.11,

$$\begin{aligned} T_1^\rho(g(0-), g) &= (\partial_{t,\rho}M(\partial_{t,\rho}) + A)^{-1} \left( \delta_0 E g(0-) - \chi_{\mathbb{R}_{\geq 0}}(\mathfrak{m}) \begin{pmatrix} M_1(\partial_{t,\rho}) \\ 0 \end{pmatrix} g \right) \\ &= \left( \partial_{t,\rho}E + A + \begin{pmatrix} M_1(\partial_{t,\rho}) \\ 0 \end{pmatrix} \right)^{-1} (\delta_0 E g(0-)) - \\ &\quad - (\partial_{t,\rho}M(\partial_{t,\rho}) + A)^{-1} \left( \chi_{\mathbb{R}_{\geq 0}}(\mathfrak{m}) \begin{pmatrix} M_1(\partial_{t,\rho}) \\ 0 \end{pmatrix} g \right). \end{aligned}$$

The first term in the latter expression can be rewritten as

$$\left( \partial_{t,\rho}E + A + \begin{pmatrix} M_1(\partial_{t,\rho}) \\ 0 \end{pmatrix} \right)^{-1} (\delta_0 E g(0-)) = \left( 1 + (\partial_{t,\rho}E + A)^{-1} \begin{pmatrix} M_1(\partial_{t,\rho}) \\ 0 \end{pmatrix} \right)^{-1} \tilde{T}_1^\rho (\delta_0 E g(0-)).$$

Now, since  $\tilde{T}_1^\rho (\delta_0 E g(0-)) \in L_{2,\rho}(\mathbb{R}; L_2(\Omega) \times L_2(\Omega)^n)$  and  $\|M_1(\partial_{t,\rho})\| \rightarrow 0$  as  $\rho \rightarrow \infty$ , we infer that

$$\left( \partial_{t,\rho}E + A + \begin{pmatrix} M_1(\partial_{t,\rho}) \\ 0 \end{pmatrix} \right)^{-1} (\delta_0 E g(0-)) \in L_{2,\rho}(\mathbb{R}; L_2(\Omega) \times L_2(\Omega)^n)$$

for  $\rho$  large enough by the Neumann series. Since clearly

$$L_2(\Omega) \times L_2(\Omega)^n \ni x \mapsto \left( \partial_{t,\rho}E + A + \begin{pmatrix} M_1(\partial_{t,\rho}) \\ 0 \end{pmatrix} \right)^{-1} (\delta_0 E x) \in H_\rho^{-1}(\mathbb{R}; L_2(\Omega) \times L_2(\Omega)^n)$$

is bounded, we obtain

$$\left\| \left( \partial_{t,\rho}E + A + \begin{pmatrix} M_1(\partial_{t,\rho}) \\ 0 \end{pmatrix} \right)^{-1} (\delta_0 E g(0-)) \right\|_{L_{2,\rho}(\mathbb{R}; L_2(\Omega) \times L_2(\Omega)^n)} \leq C \|g(0-)\|_{L_2(\Omega) \times L_2(\Omega)^n}$$

for some  $C \geq 0$  by the closed graph theorem. Hence,

$$\begin{aligned} &\|T_1^\rho(g(0-), g)\|_{L_{2,\rho}(\mathbb{R}; L_2(\Omega) \times L_2(\Omega)^n)} \\ &\leq C \|g(0-)\|_{L_2(\Omega) \times L_2(\Omega)^n} + \left\| (\partial_{t,\rho}M(\partial_{t,\rho}) + A)^{-1} \left( \chi_{\mathbb{R}_{\geq 0}}(\mathfrak{m}) \begin{pmatrix} M_1(\partial_{t,\rho}) \\ 0 \end{pmatrix} g \right) \right\|_{L_{2,\rho}(\mathbb{R}; L_2(\Omega) \times L_2(\Omega)^n)} \\ &\leq C \|g(0-)\|_{L_2(\Omega) \times L_2(\Omega)^n} + C_1 \|g\|_{L_{2,\rho}(\mathbb{R}; L_2(\Omega) \times L_2(\Omega)^n)} \\ &\leq \tilde{C} \|(g(0-), g)\|_{X_\rho^\mu} \end{aligned}$$

for suitable  $C_1, \tilde{C} \geq 0$ . Thus,

$$T_1^\rho : D_\rho \subseteq X_\rho^\mu \rightarrow L_{2,\rho}(\mathbb{R}_{\geq 0}; L_2(\Omega) \times L_2(\Omega)^n)$$

is bounded and hence extends to a bounded operator on  $X_\rho^\mu$ . Moreover, for  $f \in C_c^\infty(\mathbb{R}_{\geq 0}; L_2(\Omega) \times L_2(\Omega)^n)$  we may estimate

$$\begin{aligned} \|((\partial_{t,\rho}E + A)^{-1}f)(t)\| &= \left\| \int_0^t \tilde{T}_\rho^{(1)}(t-s)E^{-1}f(s) \, ds \right\| \\ &\leq M\|E^{-1}\| \int_0^t \|f(s)\| \, ds \\ &\leq \frac{M\|E^{-1}\|}{\sqrt{2\rho}} \|f\|_{L_{2,\rho}(\mathbb{R}; L_2(\Omega) \times L_2(\Omega)^n)} e^{\rho t} \end{aligned}$$

for each  $t \geq 0$ , which proves that

$$(\partial_{t,\rho}E + A)^{-1} : L_{2,\rho}(\mathbb{R}_{\geq 0}; L_2(\Omega) \times L_2(\Omega)^n) \rightarrow C_\rho(\mathbb{R}_{\geq 0}; L_2(\Omega) \times L_2(\Omega)^n)$$

is bounded. Now, let  $(x, g) \in X_\rho^\mu$  and set  $u := T_1^\rho(x, g) \in L_{2,\rho}(\mathbb{R}_{\geq 0}; L_2(\Omega) \times L_2(\Omega)^n)$ . Then

$$(\partial_{t,\rho}E + A)u = \delta_0 E x - \chi_{\mathbb{R}_{\geq 0}}(\mathfrak{m}) \begin{pmatrix} M_1(\partial_{t,\rho}) \\ 0 \end{pmatrix} g - \begin{pmatrix} M_1(\partial_{t,\rho}) \\ 0 \end{pmatrix} u$$

and hence, we derive that

$$\begin{aligned} u &= (\partial_{t,\rho}E + A)^{-1} \left( \delta_0 E x - \chi_{\mathbb{R}_{\geq 0}}(\mathfrak{m}) \begin{pmatrix} M_1(\partial_{t,\rho}) \\ 0 \end{pmatrix} g - \begin{pmatrix} M_1(\partial_{t,\rho}) \\ 0 \end{pmatrix} u \right) \\ &= \tilde{T}_1^\rho(x, g) - (\partial_{t,\rho}E + A)^{-1} \left( \chi_{\mathbb{R}_{\geq 0}}(\mathfrak{m}) \begin{pmatrix} M_1(\partial_{t,\rho}) \\ 0 \end{pmatrix} g + \begin{pmatrix} M_1(\partial_{t,\rho}) \\ 0 \end{pmatrix} u \right) \\ &\in C_\rho(\mathbb{R}_{\geq 0}; L_2(\Omega) \times L_2(\Omega)^n) \end{aligned}$$

and hence, again by the closed graph theorem

$$T_1^\rho : X_\rho^\mu \rightarrow C_\rho(\mathbb{R}_{\geq 0}; L_2(\Omega) \times L_2(\Omega)^n)$$

is bounded. □

## References

- [1] W. Arendt. Vector-valued Laplace transforms and Cauchy problems. *Isr. J. Math.*, 59:327–352, 1987.
- [2] W. Arendt, C. J. Batty, M. Hieber, and F. Neubrander. *Vector-valued Laplace transforms and Cauchy problems. 2nd ed.* Basel: Birkhäuser, 2011.
- [3] A. Bátkai and S. Piazzera. *Semigroups for delay equations.* Research Notes in Mathematics 10. Wellesley, MA: A K Peters. xii, 2005.

- [4] J. Diestel and J. Uhl. Vector measures. Mathematical Surveys. No.15. Providence, R.I.: American Mathematical Society (AMS). XIII., 1977.
- [5] K. J. Engel and R. Nagel. *One-parameter semigroups for linear evolution equations*. Graduate texts in mathematics. Springer, 2000.
- [6] Y. Fourès and I. Segal. Causality and analyticity. *Trans. Am. Math. Soc.*, 78:385–405, 1955.
- [7] E. Hille and R. S. Phillips. *Functional analysis and semi-groups*. Colloquium Publications - American Mathematical Society. American Mathematical Society, 1957.
- [8] A. Kalauch, R. Picard, S. Siegmund, S. Trostorff, and M. Waurick. A Hilbert space perspective on ordinary differential equations with memory term. *J. Dyn. Differ. Equations*, 26(2):369–399, 2014.
- [9] P. Kunkel and V. Mehrmann. *Differential-algebraic equations. Analysis and numerical solution*. Zürich: European Mathematical Society Publishing House, 2006.
- [10] R. E. Paley and N. Wiener. Fourier transforms in the complex domain. (Am. Math. Soc. Colloq. Publ. 19) New York: Am. Math. Soc. VIII, 1934.
- [11] A. Pazy. Semigroups of linear operators and applications to partial differential equations. Applied Mathematical Sciences, 44. New York etc.: Springer-Verlag. VIII, 1983.
- [12] R. Picard. A structural observation for linear material laws in classical mathematical physics. *Math. Methods Appl. Sci.*, 32(14):1768–1803, 2009.
- [13] R. Picard and D. McGhee. *Partial differential equations. A unified Hilbert space approach*. de Gruyter Expositions in Mathematics 55. Berlin: de Gruyter. xviii, 2011.
- [14] R. Picard, S. Trostorff, and M. Waurick. On evolutionary equations with material laws containing fractional integrals. *Math. Methods Appl. Sci.*, 38(15):3141–3154, 2015.
- [15] R. Picard, S. Trostorff, M. Waurick, and M. Wehowski. On Non-autonomous Evolutionary Problems. *J. Evol. Equ.*, 13(4):751–776, 2013.
- [16] R. H. Picard. *Hilbert space approach to some classical transforms*. Pitman research notes in mathematics series. Longman Scientific & Technical, 1989.
- [17] W. Rudin. *Real and complex analysis*. Mathematics series. McGraw-Hill, 1987.
- [18] S. Trostorff. An alternative approach to well-posedness of a class of differential inclusions in Hilbert spaces. *Nonlinear Anal., Theory Methods Appl., Ser. A, Theory Methods*, 75(15):5851–5865, 2012.

- [19] S. Trostorff. Autonomous Evolutionary Inclusions with Applications to Problems with Nonlinear Boundary Conditions. *Int. J. Pure Appl. Math.*, 85(2):303–338, 2013.
- [20] S. Trostorff. A characterization of boundary conditions yielding maximal monotone operators. *J. Funct. Anal.*, 267(8):2787–2822, 2014.
- [21] S. Trostorff. Exponential stability for second order evolutionary problems. *J. Math. Anal. Appl.*, 429(2):1007–1032, 2015.
- [22] S. Trostorff. On Integro-Differential Inclusions with Operator-valued Kernels. *Math. Methods Appl. Sci.*, 38(5):834–850, 2015. doi:10.1002/mma.3111.
- [23] S. Trostorff. *Exponential Stability and Initial Value Problems for Evolutionary Equations*. Habilitation thesis, TU Dresden, 2018. URL: <https://nbn-resolving.org/urn:nbn:de:bsz:14-qucosa-236494>.
- [24] S. Trostorff. Well-posedness for a general class of differential inclusions. Technical report, TU Dresden, 2018. arXiv: 1808.00224, submitted.
- [25] S. Trostorff. Semigroups associated with differential-algebraic equations. Technical report, CAU Kiel, 2019. arXiv:1905.11197.
- [26] S. Trostorff and M. Waurick. On higher index differential-algebraic equations in infinite dimensions. In *The diversity and beauty of applied operator theory*, volume 268 of *Oper. Theory Adv. Appl.*, pages 477–486. Birkhäuser/Springer, Cham, 2018.
- [27] S. Trostorff and M. Waurick. On differential-algebraic equations in infinite dimensions. *J. Differ. Equations*, 266(1):526–561, 2019.
- [28] S. Trostorff and M. Wehowski. Well-posedness of non-autonomous evolutionary inclusions. *Nonlinear Anal., Theory Methods Appl., Ser. A, Theory Methods*, 101:47–65, 2014.
- [29] M. Waurick. On non-autonomous integro-differential-algebraic evolutionary problems. *Math. Methods Appl. Sci.*, 38(4):665–676, 2015.
- [30] J. Weidmann. *Linear operators in Hilbert spaces. Transl. by Joseph Szücs.*, volume 68. Springer, New York, NY, 1980.
- [31] G. Weiss. Representation of shift-invariant operators on  $L^2$  by  $H^\infty$  transfer functions: An elementary proof, a generalization to  $L^p$ , and a counterexample for  $L^\infty$ . *Math. Control Signals Syst.*, 4(2):193–203, 1991.
- [32] D. Widder. The inversion of the Laplace integral and the related moment problem. *Trans. Am. Math. Soc.*, 36:107–200, 1934.