

Stationary solutions of a free boundary problem modeling the growth of vascular tumors with a necrotic core

Huijuan Song^a, Bei Hu^b and Zejia Wang^a

^a *College of Mathematical and Informational Science, Jiangxi Normal University,
Nanchang 330022, China*

^b *Department of Applied Computational Mathematics and Statistics, University of Notre Dame,
Notre Dame, IN 46556, USA*

Abstract

In this paper, we present a rigorous mathematical analysis of a free boundary problem modeling the growth of a vascular solid tumor with a necrotic core. If the vascular system supplies the nutrient concentration σ to the tumor at a rate β , then $\frac{\partial \sigma}{\partial \mathbf{n}} + \beta(\sigma - \bar{\sigma}) = 0$ holds on the tumor boundary, where \mathbf{n} is the unit outward normal to the boundary and $\bar{\sigma}$ is the nutrient concentration outside the tumor. The living cells in the nonnecrotic region proliferate at a rate μ . We show that for any given $\rho > 0$, there exists a unique $R \in (\rho, \infty)$ such that the corresponding radially symmetric solution solves the steady-state necrotic tumor system with necrotic core boundary $r = \rho$ and outer boundary $r = R$; moreover, there exist a positive integer n^{**} and a sequence of μ_n , symmetry-breaking stationary solutions bifurcate from the radially symmetric stationary solution for each μ_n (even $n \geq n^{**}$).

Keywords: Stationary solution; Free boundary problem; Vascular tumor; Necrotic core; Bifurcation.

2000 MR Subject Classification: 35R35, 35K57, 35B35

1 Introduction

The process of tumor growth in vivo is a complicated phenomenon involving many inter-related processes, which can be divided into two phases: avascular and vascular growth. In both phases, when a tumor has grown to a detectable size, the inner far-from-surface region of the tumor may comprise only dead cells due to the nonuniform distribution of nutrient materials, which is called necrotic core; see [3, 11, 20] and the references cited therein. In this paper, we are interested in a model for the growth of a vascular solid tumor, which consists of a necrotic core of dead cells, and a surrounding shell which contains life-proliferating cells. Assume that there is no consumption of the nutrient as well as no vasculature in the necrotic core; on the other hand, as a result of angiogenesis, the proliferating rim possesses its own vasculature, and then the nutrient may be supplied to the nonnecrotic shell via the capillary network. Thus, as in [6, 9], the nutrient concentration σ satisfies the following reaction-diffusion equation

$$c \frac{\partial \sigma}{\partial t} = \Delta \sigma + [\Gamma(\sigma_B - \sigma) - \lambda_0 \sigma] I_{\Omega(t) \setminus D(t)} \quad \text{in } \Omega(t), \quad (1.1)$$

where $\Omega(t) \subset \mathbb{R}^3$ is the tumor domain at time t with a moving boundary $\partial\Omega(t)$, $D(t) \subset \Omega(t)$ is the necrotic core region, Γ is the transfer rate of nutrient-in-blood-tissue, σ_B is the concentration of nutrient in the vasculature, and so the term $\Gamma(\sigma_B - \sigma)$ accounts for the transfer of nutrient by means of the vasculature stemming from angiogenesis in the nonnecrotic region. The term $\lambda_0\sigma$ describes the nutrient consumption by proliferating cells at the rate of λ_0 , and $c = T_{\text{diffusion}}/T_{\text{growth}}$ represents the ratio of the nutrient diffusion time scale to the tumor growth (i.e., tumor doubling) time scale. Typically, $T_{\text{diffusion}} \approx 1$ minute while $T_{\text{growth}} \approx 1$ day [5, 9]. Thus, $c \ll 1$. Throughout this paper, the notation I_E will always be used to denote the indicator function of a subset E of \mathbb{R}^3 .

Using appropriate change of variables [9, 32], we can rewrite the equation (1.1) in the form

$$c \frac{\partial \sigma}{\partial t} = \Delta \sigma - \sigma I_{\Omega(t) \setminus D(t)} \quad \text{in } \Omega(t). \quad (1.2)$$

As mentioned above, it is natural to assume that

$$\sigma = \underline{\sigma} \quad \text{in } D(t), \quad (1.3)$$

where $\underline{\sigma} > 0$ plays the role of a threshold value in the sense that in the region where $\sigma > \underline{\sigma}$ nutrient is enough to sustain (at least a portion of) tumor cells alive and proliferating, whereas in the region where $\sigma \leq \underline{\sigma}$, nutrient is insufficient to sustain any tumor cell alive [51]. In addition, since the nutrient enters tumor by the vascular system, σ satisfies the boundary condition:

$$\frac{\partial \sigma}{\partial \mathbf{n}} + \beta(t)(\sigma - \bar{\sigma}) = 0 \quad \text{on } \partial\Omega(t), \quad (1.4)$$

where \mathbf{n} is the outward normal, $\bar{\sigma}$ is the nutrient concentration outside the tumor, $\beta(t)$ is the rate of nutrient supply to the tumor, which may vary in time. Angiogenesis results in an increase in $\beta(t)$; conversely, if the tumor is treated with anti-angiogenic drugs, $\beta(t)$ will decrease and the starved tumor will shrink.

The pressure p stems from the transport of cells which proliferate or die. Let \mathbf{V} be the velocity of tumor cells. If we assume that the density of tumor cells is constant, there is no proliferation in the necrotic core, and the proliferation rate within the nonnecrotic region is linearly dependent on the nutrient, then the conservation of mass gives that

$$\text{div} \mathbf{V} = \mu(\sigma - \tilde{\sigma}) I_{\Omega(t) \setminus D(t)} - \nu I_{D(t)},$$

where μ is a positive parameter expressing the ‘‘intensity’’ of the expansion by mitosis, the term $\mu(\sigma - \tilde{\sigma})$ means that the cell birthrate is $\mu\sigma$, while the cell death rate (apoptosis) is given by $\mu\tilde{\sigma}$, and ν is the dissolution rate of necrotic cells. Combining with Darcy’s law $\mathbf{V} = -\nabla p$, we obtain

$$-\Delta p = \mu(\sigma - \tilde{\sigma}) I_{\Omega(t) \setminus D(t)} - \nu I_{D(t)} \quad \text{in } \Omega(t). \quad (1.5)$$

Due to cell-to-cell adhesiveness, there holds

$$p = \kappa \quad \text{on } \partial\Omega(t), \quad (1.6)$$

where κ is the mean curvature, and the continuity of the velocity field up to the boundary of the tumor leads to

$$V_{\mathbf{n}} = \mathbf{V} \cdot \mathbf{n} = -\frac{\partial p}{\partial \mathbf{n}} \quad \text{on } \partial\Omega(t), \quad (1.7)$$

where $V_{\mathbf{n}}$ is the velocity of the free boundary in the direction \mathbf{n} .

The first reaction-diffusion mathematical model of tumor growth in the form of a free boundary problem of a system of partial differential equations was proposed in 1972 by Greenspan [33, 34], which was remarkably improved by Byrne and Chaplain [5, 6] during 1990's, where apoptosis is incorporated. It was Friedman and Reitich [22] who made the rigorous mathematical analysis of such free boundary problems in 1999. To date, mathematical modeling, numerical simulation and theoretical analysis on tumor models have been carried out in a large number of papers; see [3, 4, 8–20, 23, 26, 27, 29–32, 35–37, 41–56], the review articles [2, 25, 28, 39, 40] and the references therein.

If $D(t) = \emptyset$ and $\Omega(t) = B_{R(t)}(0)$, then the model (1.2)-(1.7) describes the growth of a nonnecrotic spherical tumor with angiogenesis, proposed by Friedman and Lam [32] as a revision to the Byrne-Chaplain inhibitor-free tumor model [5] in which instead of (1.4),

$$\sigma = \bar{\sigma} \quad (1.8)$$

is imposed on the tumor boundary. Biologically, the boundary condition (1.4) is more reasonable compared with (1.8). In fact, as explained in [55], tumor surface acts as a barrier to nutrient diffusion, and $1/\beta(t)$ reflects the reduction rate of nutrient by the tumor surface; particularly, $1/\beta(t) = 0$ means that tumor surface is obstacle-free to nutrient diffusion which is the case (1.8), whereas $1/\beta(t) = \infty$ means that tumor surface is a complete barrier to nutrient diffusion.

In the nonnecrotic case, assuming $\beta(t) \equiv \beta$ and $0 < \bar{\sigma} < \bar{\sigma}$, Friedman and Lam [32] showed that the system (1.2)-(1.7) allows a unique radially symmetric stationary solution $(\sigma_s(r), p_s(r), R_s)$; later, Huang, Zhang and Hu [36] proved that a branch of symmetry-breaking stationary solutions bifurcates from the radially symmetric stationary solution for each $\mu_n(R_s)$ (even $n \geq 2$) with free boundary

$$r = R_s + \varepsilon Y_{n,0}(\theta, \varphi) + o(\varepsilon),$$

where $Y_{n,0}$ is the spherical harmonic of order $(n, 0)$; very recently, they [37] made further efforts and found a threshold value $\mu^* = \mu^*(R_s)$ such that the radially symmetric stationary solution is linearly stable for $\mu < \mu^*$ and linearly unstable for $\mu > \mu^*$ with respect to nonradial perturbations. For the case where the nutrient consumption rate and the proliferation rate of tumor cells are both general functions, Zhuang and Cui established the existence and uniqueness of radially symmetric stationary solutions [55], and proved that there exists a branch of bifurcation solutions bifurcating from the radially symmetric stationary solution for γ_k (any $k \geq k_*$ in the 2 dimension case and even $k \geq k_*$ in the 3 dimension case) by taking γ (surface tension coefficient γ is defined when (1.6) is replaced by $p = \gamma\kappa$) as the bifurcation parameter [17]. The asymptotic stability of radially symmetric stationary solutions was also analyzed in [55, 56]. For cases in the presence of inhibitor, Wang et al. [44] obtained the existence of symmetric-breaking stationary solutions for μ_n (even $n > n^{**}$); also see a very recent paper [41] for the discussion on the existence of radially symmetric stationary solutions and the asymptotic behavior of radially symmetric transient solutions. Assuming $\beta(t) = \infty$, i.e., (1.8) holds, tumor models have been

intensively studied; we refer the reader to [12–14, 16, 19, 23, 26, 27, 29–31, 43, 49, 50, 52, 53] and the reference therein.

For the necrotic case, most of studies were on tumor models where the Dirichlet boundary condition (1.8) is imposed; see [3, 4, 6, 9–11, 15, 20, 35, 45, 47, 48]. Cui [11] proved the existence and uniqueness of radially symmetric stationary solutions under a crucial assumption $\underline{\sigma} < \tilde{\sigma} < \underline{\sigma} + \nu/\mu$, i.e., $\nu > \mu(\tilde{\sigma} - \underline{\sigma}) > 0$, improving earlier results in [9]. Under the assumption $\nu=0$, Hao et al. [35] derived the first bifurcation result for the tumor model with a necrotic core, in which they studied the two-dimensional case by taking μ as a bifurcation parameter with the aid of numerical calculations. Very recently, Wu [47] rigorously analyzed the necrotic multilayered tumor model, and obtained the existence of bifurcation branches of non-flat stationary solutions for γ_k ($k \geq K$). Asymptotic stability of stationary solutions to the above two types of necrotic tumor models was also studied, cf. [15, 48]. For necrotic tumor models with (1.4), Shen et al. [42] established the existence and uniqueness of radially symmetric stationary solutions to the tumor spheroid model with $\nu > 0$ under certain conditions on the parameters.

It is well known that the main feature of the free boundary problems modeling the growth of necrotic tumors is that they include two free boundaries, one for the outer tumor boundary, whose evolution is governed by an evolution equation (such as (1.7)), the other for the inner necrotic boundary, an obstacle-type free surface, whose evolution is implicit. Due to the presence of two free boundaries, the mathematical analysis turns out to be far more challenging.

Motivated by [35, 47], we shall perform rigorous mathematical analysis of the stationary state of the problem (1.2)-(1.7) in the three-dimensional case, under the assumption that

$$0 < \underline{\sigma} < \tilde{\sigma} < \bar{\sigma} = 1, \quad \beta(t) \equiv \beta, \quad \nu = 0,$$

where β is a positive constant. That is,

$$\Delta\sigma = \sigma I_{\Omega \setminus D} \quad \text{in } \Omega, \tag{1.9}$$

$$-\Delta p = \mu(\sigma - \tilde{\sigma}) I_{\Omega \setminus D} \quad \text{in } \Omega, \tag{1.10}$$

$$\sigma = \underline{\sigma} \quad \text{in } D, \tag{1.11}$$

$$[\partial_{\mathbf{n}}\sigma] = 0 \quad \text{on } \partial D, \tag{1.12}$$

$$[p] = 0, \quad [\partial_{\mathbf{n}}p] = 0 \quad \text{on } \partial D, \tag{1.13}$$

$$\partial_{\mathbf{n}}\sigma + \beta(\sigma - 1) = 0 \quad \text{on } \partial\Omega, \tag{1.14}$$

$$p = \kappa \quad \text{on } \partial\Omega, \tag{1.15}$$

$$\partial_{\mathbf{n}}p = 0 \quad \text{on } \partial\Omega. \tag{1.16}$$

Here and below, the notation $[p]|_{\partial D}$ denotes the jump of p as it crosses ∂D , i.e.,

$$[p] = p^+|_{\partial D} - p^-|_{\partial D} \quad \text{for } p^+ = p|_{\Omega \setminus D} \quad \text{and } p^- = p|_D.$$

Similarly, $[\partial_{\mathbf{n}}\sigma]|_{\partial D}$ and $[\partial_{\mathbf{n}}p]|_{\partial D}$ denote the jump of the normal derivatives of σ and p across ∂D respectively. It is not difficult to find out, the inner boundary conditions (1.12), (1.13) are implied by the equations (1.9), (1.10), and by the maximum principle, $\sigma > \underline{\sigma}$ in $\Omega \setminus \overline{D}$. For the problem (1.9)-(1.16), we shall first adopt a similar idea as that in [35] to study radially symmetric

solutions, then by choosing μ as a bifurcation parameter, show that there exist a positive integer n^{**} and a sequence of μ_n such that for each μ_n (even $n \geq n^{**}$), a branch of symmetry-breaking stationary solutions bifurcates from the radially symmetric solution (see Theorem 5.1 stated below).

We stress that in carrying out the bifurcation analysis based on the Crandall-Rabinowitz theorem (see Theorem 2.1), there are three major difficulties to be overcome. Firstly, since the problem (1.9)-(1.16) involves two free boundaries, there is a need for an appropriate Hanzawa-type transformation. Secondly, noting that the expansions (4.8), (4.9) are needed for the computation of Fréchet derivatives, in order to rigorously prove the expansions (4.8), (4.9) of σ , p with respect to ε , we have to analyze the dependence of the inner boundary on the outer boundary and the nutrient concentration (see (4.26), (4.36)), and establish Schauder estimates in each region (necrotic and nonnecrotic) for diffraction problems by using the Schauder estimates for an elliptic equation and that near the boundary for elliptic systems [1]. Thirdly, it is necessary to verify $B_0 < 0$ (see Lemma 5.1). However, as we shall see later, the expression of B_0 is complex. Inspired by [36], we will write B_0 in order of ascending powers of β . Then, thanks to the existence of explicit forms for the modified Bessel functions of order of half an odd integer, we can deduce that the coefficient (the function of ρ , R) of each term is negative by a lengthy calculation.

The rest of this paper is arranged as follows. In Section 2, we present some preliminary material which will be needed in the next sections. In Section 3, we study the radial symmetric solutions to the problem (1.9)-(1.16). In Section 4, we analyze the linearization of (1.9)-(1.16) about the radial solution by introducing an appropriate Hanzawa-type transformation. In the last section, we prove the existence of bifurcation solutions.

2 Preliminaries

In this section, we first collect some properties of the spherical harmonics and the modified spherical Bessel functions, then present an auxiliary lemma and end with the Crandall-Rabinowitz theorem.

In \mathbb{R}^3 , the family of the spherical harmonic functions $\{Y_{n,m}\}$ forms a complete orthonormal basis for $L^2(\Sigma)$, where Σ is the unit sphere, and

$$\Delta_\omega Y_{n,m} = -n(n+1)Y_{n,m}. \quad (2.1)$$

Here and below,

$$\Delta_\omega = \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial}{\partial \theta} \right) + \frac{1}{\sin^2 \theta} \frac{\partial^2}{\partial \varphi^2}$$

is the Laplace operator on Σ . The Laplace operator in \mathbb{R}^3 can be written as

$$\Delta = \frac{\partial^2}{\partial r^2} + \frac{2}{r} \frac{\partial}{\partial r} + \frac{1}{r^2} \Delta_\omega.$$

The modified spherical Bessel functions given by

$$i_n(s) = \sqrt{\frac{\pi}{2s}} I_{n+1/2}(s), \quad k_n(s) = \sqrt{\frac{\pi}{2s}} K_{n+1/2}(s), \quad s > 0, \quad n = 0, 1, 2, \dots,$$

form a fundamental solution set of the differential equation (cf. [21, (10.47.7)–(10.47.9)])

$$y''(s) + \frac{2}{s}y'(s) - \left[1 + \frac{n(n+1)}{s^2}\right]y(s) = 0, \quad (2.2)$$

where $I_{n+1/2}(s)$ and $K_{n+1/2}(s)$ are the modified Bessel functions, $I_\nu(s) > 0$, $I'_\nu(s) > 0$ for $s > 0$ [21, (10.25.2)], and $K_\nu(s) > 0$, $K'_\nu(s) < 0$ for $s > 0$ [21, (10.32.9)]. Let $g_n(s)$ denote $i_n(s)$ or $(-1)^n k_n(s)$. Then from [21, (10.51.4) and (10.51.5)],

$$g_{n-1}(s) - g_{n+1}(s) = \frac{2n+1}{s}g_n(s), \quad n \geq 1, \quad (2.3)$$

$$g'_n(s) = g_{n-1}(s) - \frac{n+1}{s}g_n(s), \quad n \geq 1, \quad (2.4)$$

$$g'_n(s) = g_{n+1}(s) + \frac{n}{s}g_n(s), \quad n \geq 0. \quad (2.5)$$

It follows from (2.4), (2.5) that

$$i'_n(s) > 0 \quad \text{and} \quad k'_n(s) < 0, \quad n \geq 0, \quad s > 0. \quad (2.6)$$

In addition, by [21, (10.28.2), (10.41.1) and (10.41.2)],

$$i_n(s)k_{n+1}(s) + i_{n+1}(s)k_n(s) = \frac{\pi}{2} \frac{1}{s^2}, \quad n \geq 0, \quad (2.7)$$

$$i_n(s) \sim \frac{1}{\sqrt{2(2n+1)s}} \left(\frac{es}{2n+1}\right)^{n+\frac{1}{2}} \quad \text{as } n \rightarrow \infty, \quad (2.8)$$

$$k_n(s) \sim \frac{\pi}{\sqrt{2(2n+1)s}} \left(\frac{es}{2n+1}\right)^{-n-\frac{1}{2}} \quad \text{as } n \rightarrow \infty. \quad (2.9)$$

Explicit expressions for i_n , k_n , $n = 0, 1$ are as follows [21, (10.49.9) and (10.49.13)]:

$$i_0(s) = \frac{\sinh s}{s}, \quad k_0(s) = \frac{\pi e^{-s}}{2s}, \quad (2.10)$$

$$i_1(s) = -\frac{\sinh s}{s^2} + \frac{\cosh s}{s}, \quad k_1(s) = \frac{\pi e^{-s}}{2} \left(\frac{1}{s} + \frac{1}{s^2}\right). \quad (2.11)$$

A direct calculation shows that

$$i_0(t)k_0(s) - i_0(s)k_0(t) = \frac{\pi}{2} \frac{1}{ts} \sinh(t-s), \quad (2.12)$$

$$i_1(t)k_1(s) - i_1(s)k_1(t) = \frac{\pi}{2} \frac{1}{t^2s^2} [(t-s) \cosh(t-s) + (ts-1) \sinh(t-s)], \quad (2.13)$$

$$i_0(t)k_1(s) + i_1(s)k_0(t) = \frac{\pi}{2} \frac{1}{ts^2} [\sinh(t-s) + s \cosh(t-s)]. \quad (2.14)$$

In order to establish the dependence of the inner boundary on the tumor outer boundary, we need the following lemma.

Lemma 2.1. ([24, Lemma 8.2]) Let s be a nonnegative integer, and let

$$f(\theta, \varphi) = \sum_{n \geq 0, m} f_{n,m} Y_{n,m}(\theta, \varphi).$$

Then there exist positive constants c_1, c_2 independent of f such that

$$c_1 \|f\|_{H^{s+1/2}(\Sigma)}^2 \leq \sum_{n \geq 0} (1 + n^{2s+1}) \sum_m |f_{n,m}|^2 \leq c_2 \|f\|_{H^{s+1/2}(\Sigma)}^2.$$

The bifurcation analysis in the present paper is based on an application of the following theorem of Crandall and Rabinowitz.

Theorem 2.1. ([7, Theorem 1.7]) Let X, Y be real Banach spaces and $F(x, \mu)$ a C^p map, $p \geq 3$, of a neighborhood $(0, \mu_0)$ in $X \times \mathbb{R}$ into Y . Suppose

- (i) $F(0, \mu) = 0$ for all μ in a neighborhood of μ_0 ;
- (ii) $\text{Ker}[F_x(0, \mu_0)]$ is a one dimensional space, spanned by x_0 ;
- (iii) $\text{Im}[F_x(0, \mu_0)] = Y_1$ has codimension 1;
- (iv) $[F_{\mu x}](0, \mu_0)x_0 \notin Y_1$.

Then $(0, \mu_0)$ is a bifurcation point of the equation $F(x, \mu) = 0$ in the following sense: In a neighborhood of $(0, \mu_0)$ the set of solutions of $F(x, \mu) = 0$ consists of two C^{p-2} smooth curves Γ_1 and Γ_2 which intersect only at the point $(0, \mu_0)$; Γ_1 is the curve $(0, \mu)$ and Γ_2 can be parameterized as follows:

$$\Gamma_2 : (x(\varepsilon), \mu(\varepsilon)), |\varepsilon| \text{ small}, (x(0), \mu(0)) = (0, \mu_0), x'(0) = x_0.$$

3 Radially symmetric stationary solutions

In this section, we study radially symmetric solutions to the system (1.9)-(1.16), denoted by $(\sigma_s(r), p_s(r), \rho, R)$, where $r = |x|$.

First, it follows from (1.9), (1.11), (1.12) and (1.14) that

$$\begin{cases} \sigma_s''(r) + \frac{2}{r}\sigma_s'(r) - \sigma_s(r) = 0, & \rho < r < R, \\ \sigma_s(\rho) = \underline{\sigma}, \quad \sigma_s'(\rho) = 0, \quad \sigma_s'(R) + \beta(\sigma_s(R) - 1) = 0. \end{cases} \quad (3.1)$$

Using (2.2), (2.5) and (2.7), we find

$$\sigma_s(r) = \begin{cases} \frac{2}{\pi}\underline{\sigma}\rho^2(k_1(\rho)i_0(r) + i_1(\rho)k_0(r)) & \text{for } \rho < r < R, \\ \underline{\sigma} & \text{for } 0 \leq r \leq \rho \end{cases} \quad (3.2)$$

with ρ, R satisfying

$$(k_1(\rho)i_1(R) - i_1(\rho)k_1(R)) + \beta(k_1(\rho)i_0(R) + i_1(\rho)k_0(R)) = \frac{\pi\beta}{2\underline{\sigma}\rho^2}. \quad (3.3)$$

Next we solve for p_s . In view of (1.10), (1.13), (1.15) and (1.16), we find that p_s satisfies

$$-p_s''(r) - \frac{2}{r}p_s'(r) = \mu(\sigma_s(r) - \tilde{\sigma}) \quad \text{for } \rho < r < R, \quad (3.4)$$

$$p'_s(\rho) = 0, \quad p_s(R) = \frac{1}{R}, \quad p'_s(R) = 0. \quad (3.5)$$

Let $q = p_s + \mu\sigma_s$. Then (3.4) reduces to

$$\Delta q = \mu\tilde{\sigma}.$$

Thus,

$$\begin{aligned} q(r) &= \frac{1}{6}\mu\tilde{\sigma}r^2 + C_1 + C_2\frac{1}{r}, \\ p_s(r) &= -\mu\sigma_s(r) + \frac{1}{6}\mu\tilde{\sigma}r^2 + C_1 + C_2\frac{1}{r}, \quad \rho < r < R, \end{aligned}$$

where C_1, C_2 are constants. Combining with the boundary condition (3.5), we solve for C_1, C_2 and derive

$$p_s(r) = \begin{cases} -\mu(\sigma_s(r) - \sigma_s(R)) + \frac{1}{6}\mu\tilde{\sigma}(r^2 - R^2) + \frac{1}{3}\mu\tilde{\sigma}\rho^3\left(\frac{1}{r} - \frac{1}{R}\right) + \frac{1}{R} & \text{for } \rho < r < R, \\ -\mu(\underline{\sigma} - \sigma_s(R)) + \frac{1}{6}\mu\tilde{\sigma}(\rho^2 - R^2) + \frac{1}{3}\mu\tilde{\sigma}\rho^3\left(\frac{1}{\rho} - \frac{1}{R}\right) + \frac{1}{R} & \text{for } 0 \leq r \leq \rho, \end{cases} \quad (3.6)$$

where ρ, R satisfy

$$\tilde{\sigma} = \frac{3R^2}{R^3 - \rho^3}\sigma'_s(R) = \frac{3R^2}{R^3 - \rho^3}\frac{2\underline{\sigma}\rho^2}{\pi}(k_1(\rho)i_1(R) - i_1(\rho)k_1(R)). \quad (3.7)$$

In the sequel, we should solve the system of equations (3.3) and (3.7) with respect to the variables ρ and R , for given $\beta, \underline{\sigma}$ and $\tilde{\sigma}$. However, for technical reasons, we will instead compute $R, \tilde{\sigma}$ for given $\rho, \beta, \underline{\sigma}$, as was done in [35]. More precisely, we will establish the following result.

Theorem 3.1. *Assume $\mu > 0, \beta > 0$ and $0 < \underline{\sigma} < 1$. If we fix $\rho > 0$, then there exist a unique R in (ρ, ∞) and a unique $\tilde{\sigma}$ in $(\underline{\sigma}, 1)$ such that $(\sigma_s(r), p_s(r), \rho, R)$ is a radially symmetric solution to the problem (1.9)-(1.16), where $\sigma_s(r), p_s(r)$ are given by (3.2), (3.6).*

Proof. In view of (3.3), we consider the function

$$f(s) = (k_1(\rho)i_1(s) - i_1(\rho)k_1(s)) + \beta(k_1(\rho)i_0(s) + i_1(\rho)k_0(s)) - \frac{\pi\beta}{2\underline{\sigma}\rho^2}, \quad s \geq \rho.$$

Then from (2.5), (2.7) it follows that

$$\begin{aligned} f(\rho) &= \frac{\pi\beta}{2\rho^2}\left(1 - \frac{1}{\underline{\sigma}}\right), \\ f'(s) &= (k_1(\rho)i'_1(s) - i_1(\rho)k'_1(s)) + \beta(i_1(s)k_1(\rho) - i_1(\rho)k_1(s)). \end{aligned}$$

Since $\beta > 0$ and $0 < \underline{\sigma} < 1$, $f(\rho) < 0$. By (2.6) we know that $f'(s) > 0$ for $s > \rho$. Thus, combining with $\lim_{s \rightarrow +\infty} f(s) = +\infty$, we deduce that there exists a unique $R > \rho$ such that (3.3) holds. In what follows, we prove that $\tilde{\sigma}$ given by (3.7) satisfies

$$\underline{\sigma} < \tilde{\sigma} < \sigma_s(R) < 1. \quad (3.8)$$

First, we show that $\underline{\sigma} < \tilde{\sigma}$, which reduces to

$$3(R - \rho) \cosh(R - \rho) + 3(R\rho - 1) \sinh(R - \rho) - R^3 + \rho^3 > 0 \quad (3.9)$$

by (2.13). As a matter of fact, define

$$g(s) = 3(s - \rho) \cosh(s - \rho) + 3(s\rho - 1) \sinh(s - \rho) - s^3 + \rho^3, \quad s \geq \rho.$$

Then, clearly, $g(\rho) = 0$ and

$$g'(s) = 3s\{\rho[\cosh(s - \rho) - 1] + [\sinh(s - \rho) - (s - \rho)]\}.$$

Since, as may easily be verified,

$$\cosh z > \frac{\sinh z}{z} > 1, \quad z > 0, \quad (3.10)$$

$g'(s) > 0$ for $s > \rho$. Thus, $g(s) > 0$ for every $s > \rho$, i.e., (3.9) is valid.

We next proceed to verify that $\tilde{\sigma} < \sigma_s(R)$, which is equivalent to

$$(R^3 - \rho^3)(k_1(\rho)i_0(R) + i_1(\rho)k_0(R)) > 3R^2(k_1(\rho)i_1(R) - i_1(\rho)k_1(R)) \quad (3.11)$$

by (3.2) and (3.7). Based on (2.13) and (2.14), (3.11) can be further simplified to the form

$$(\rho + 1)(R^3 - 3R^2 + 3R - \rho^3)e^{2(R-\rho)} + (\rho - 1)(R^3 + 3R^2 + 3R - \rho^3) > 0. \quad (3.12)$$

Consider the function

$$h(s) = (\rho + 1)(s^3 - 3s^2 + 3s - \rho^3)e^{2(s-\rho)} + (\rho - 1)(s^3 + 3s^2 + 3s - \rho^3) \quad \text{for } s \geq \rho.$$

A direct calculation shows that

$$\begin{aligned} h'(s) &= (\rho + 1)(2s^3 - 3s^2 - 2\rho^3 + 3)e^{2(s-\rho)} + 3(\rho - 1)(s + 1)^2, \\ h''(s) &= 2(\rho + 1)(2s^3 - 3s - 2\rho^3 + 3)e^{2(s-\rho)} + 6(\rho - 1)(s + 1), \\ h'''(s) &= 2(\rho + 1)(4s^3 + 6s^2 - 6s - 4\rho^3 + 3)e^{2(s-\rho)} + 6(\rho - 1), \\ h^{(4)}(s) &= 16(\rho + 1)(s^3 + 3s^2 - \rho^3)e^{2(s-\rho)} > 0 \quad \text{for } s > \rho, \end{aligned}$$

and in particular,

$$h(\rho) = h'(\rho) = h''(\rho) = 0, \quad h'''(\rho) = 12\rho^3.$$

Applying Taylor's formula with Lagrange's remainder-term

$$h(s) = h(\rho) + h'(\rho)(s - \rho) + \frac{h''(\rho)}{2!}(s - \rho)^2 + \frac{h'''(\rho)}{3!}(s - \rho)^3 + \frac{h^{(4)}(\xi)}{4!}(s - \rho)^4,$$

where $\xi \in (\rho, s)$, provided that $s > \rho$, we see that $h(s) > 0$ for $s > \rho$. Hence, (3.12) follows.

Finally, we prove that $\sigma_s(R) < 1$. Indeed, from (3.1) and the fact $\sigma'_s(R) > 0$, it is easy to see that

$$\sigma_s(R) = 1 - \frac{\sigma'_s(R)}{\beta} < 1.$$

The proof is complete. □

4 Linearized problem

In this section, we study the linearization of the problem (1.9)-(1.16) at the radially symmetric solution (σ_s, p_s, ρ, R) , where (ρ, R) satisfies (3.3), (3.7). In order to compute Fréchet derivatives in the next section, we shall derive rigorous mathematical estimates for this linearization.

Consider a family of domains with perturbed boundaries

$$\begin{aligned}\partial D_\varepsilon &: r = \rho + \varepsilon T(\theta, \varphi), \\ \partial \Omega_\varepsilon &: r = R + \varepsilon S(\theta, \varphi),\end{aligned}$$

where both S and T are unknown functions. Let (σ, p) be the solution to

$$\Delta \sigma = \sigma I_{\Omega_\varepsilon \setminus D_\varepsilon} \quad \text{in } \Omega_\varepsilon, \quad (4.1)$$

$$-\Delta p = \mu(\sigma - \tilde{\sigma}) I_{\Omega_\varepsilon \setminus D_\varepsilon} \quad \text{in } \Omega_\varepsilon, \quad (4.2)$$

$$\sigma = \underline{\sigma}, \quad [\partial_{\mathbf{n}} \sigma] = 0 \quad \text{on } \partial D_\varepsilon, \quad (4.3)$$

$$\partial_{\mathbf{n}} \sigma + \beta(\sigma - 1) = 0 \quad \text{on } \partial \Omega_\varepsilon, \quad (4.4)$$

$$[p] = 0, \quad [\partial_{\mathbf{n}} p] = 0 \quad \text{on } \partial D_\varepsilon, \quad (4.5)$$

$$p = \kappa \quad \text{on } \partial \Omega_\varepsilon. \quad (4.6)$$

Then the well-posedness of the problem (4.1)-(4.6) follows from the local flattening and diffraction problem for systems; see [38]. If we define

$$F(\tilde{R}, \mu) = \left. \frac{\partial p}{\partial \mathbf{n}} \right|_{\partial \Omega_\varepsilon}, \quad (4.7)$$

where $\tilde{R} = \varepsilon S$, then $(\sigma, p, \rho + \varepsilon T, R + \varepsilon S)$ is a solution to the system (1.9)-(1.16) if and only if $F(\tilde{R}, \mu) = 0$.

Let us formally write

$$\sigma(r, \theta, \varphi) = \begin{cases} \hat{\sigma}(r) + \varepsilon \sigma_1(r, \theta, \varphi) + O(\varepsilon^2) & \text{in } \Omega_\varepsilon \setminus D_\varepsilon, \\ \hat{\sigma}(\rho) + \varepsilon \sigma_1(r, \theta, \varphi) + O(\varepsilon^2) & \text{in } D_\varepsilon, \end{cases} \quad (4.8)$$

$$p(r, \theta, \varphi) = \begin{cases} \hat{p}(r) + \varepsilon p_1(r, \theta, \varphi) + O(\varepsilon^2) & \text{in } \Omega_\varepsilon \setminus D_\varepsilon, \\ \hat{p}(\rho) + \varepsilon p_1(r, \theta, \varphi) + O(\varepsilon^2) & \text{in } D_\varepsilon, \end{cases} \quad (4.9)$$

where

$$\begin{aligned}\hat{\sigma}(r) &= \frac{2}{\pi} \underline{\sigma} \rho^2 (k_1(\rho) i_0(r) + i_1(\rho) k_0(r)), \quad 0 < r < \infty, \\ \hat{p}(r) &= -\mu(\hat{\sigma}(r) - \hat{\sigma}(R)) + \frac{1}{6} \mu \tilde{\sigma} (r^2 - R^2) + \frac{1}{3} \mu \tilde{\sigma} \rho^3 \left(\frac{1}{r} - \frac{1}{R} \right) + \frac{1}{R}, \quad 0 < r < \infty.\end{aligned}$$

From (3.2) and (3.6) we can easily see that

$$\Delta \hat{\sigma} = \hat{\sigma} \quad \text{in } \mathbb{R}^3 \setminus \{0\}, \quad \hat{\sigma}(r) = \sigma_s(r) \quad \text{for } \rho < r < R, \quad (4.10)$$

$$-\Delta \hat{p} = \mu(\hat{\sigma} - \tilde{\sigma}) \quad \text{in } \mathbb{R}^3 \setminus \{0\}, \quad \hat{p}(r) = p_s(r) \quad \text{for } \rho < r < R. \quad (4.11)$$

In the following we shall first present a formal derivation of the linearized problem for (σ_1, p_1) . Recall that the gradient operator in \mathbb{R}^3 can be written as

$$\nabla = \vec{e}_r \partial_r + \vec{e}_\theta \frac{1}{r} \partial_\theta + \vec{e}_\varphi \frac{1}{r \sin \theta} \partial_\varphi = \vec{e}_r \partial_r + \frac{1}{r} \nabla_\omega,$$

and if a surface is given by $r = R + \varepsilon S(\theta, \varphi)$, then

$$\mathbf{n} = \vec{e}_r - \frac{\varepsilon}{R} \nabla_\omega S + \varepsilon^2 f_1,$$

where $\|f_1\|_{C^{l-1+\alpha}(\Sigma)} \leq C_l$ provided that $\|S\|_{C^{l+\alpha}(\Sigma)} \leq 1$. Thus, from (4.4), (4.8), (4.10) and (3.1), we derive

$$\begin{aligned} 0 &= \left(\frac{\partial \sigma}{\partial r} \Big|_{\partial \Omega_\varepsilon} \vec{e}_r + \frac{1}{R} \nabla_\omega \sigma \Big|_{\partial \Omega_\varepsilon} \right) \cdot \left(\vec{e}_r - \frac{\varepsilon}{R} \nabla_\omega S \right) + \beta \sigma \Big|_{\partial \Omega_\varepsilon} - \beta + O(\varepsilon^2) \\ &= \frac{\partial \sigma}{\partial r} \Big|_{\partial \Omega_\varepsilon} - \frac{\varepsilon}{R^2} \nabla_\omega \sigma \Big|_{\partial \Omega_\varepsilon} \cdot \nabla_\omega S + \beta \sigma \Big|_{\partial \Omega_\varepsilon} - \beta + O(\varepsilon^2) \\ &= \frac{\partial \hat{\sigma}}{\partial r} \Big|_{\partial \Omega_\varepsilon} + \varepsilon \frac{\partial \sigma_1}{\partial r} \Big|_{\partial \Omega_\varepsilon} + \beta \hat{\sigma} \Big|_{\partial \Omega_\varepsilon} + \beta \varepsilon \sigma_1 \Big|_{\partial \Omega_\varepsilon} - \beta + O(\varepsilon^2) \\ &= \frac{\partial \sigma_s}{\partial r} \Big|_{\partial B_R} + \frac{\partial^2 \sigma_s}{\partial r^2} \Big|_{\partial B_R} \varepsilon S + \varepsilon \frac{\partial \sigma_1}{\partial r} \Big|_{\partial B_R} + \beta \sigma_s \Big|_{\partial B_R} + \beta \frac{\partial \sigma_s}{\partial r} \Big|_{\partial B_R} \varepsilon S + \beta \varepsilon \sigma_1 \Big|_{\partial B_R} \\ &\quad - \beta + O(\varepsilon^2) \\ &= \varepsilon \left[\left(\frac{\partial \sigma_1}{\partial r} + \beta \sigma_1 \right) \Big|_{\partial B_R} + \lambda S \right] + O(\varepsilon^2), \end{aligned}$$

where

$$\lambda = \left(\frac{\partial^2 \sigma_s}{\partial r^2} + \beta \frac{\partial \sigma_s}{\partial r} \right) \Big|_{\partial B_R}.$$

Using (3.1), (3.2) and (2.5), we get

$$\frac{\partial \sigma_s}{\partial r} \Big|_{\partial B_R} = \frac{2}{\pi} \underline{\sigma} \rho^2 (k_1(\rho) i_1(R) - i_1(\rho) k_1(R)),$$

and

$$\frac{\partial^2 \sigma_s}{\partial r^2} \Big|_{\partial B_R} = \frac{2}{\pi} \underline{\sigma} \rho^2 \left[(i_0(R) k_1(\rho) + i_1(\rho) k_0(R)) - \frac{2}{R} (i_1(R) k_1(\rho) - i_1(\rho) k_1(R)) \right],$$

so that

$$\lambda = \frac{2}{\pi} \underline{\sigma} \rho^2 \left[i_0(R) k_1(\rho) + i_1(\rho) k_0(R) + \left(\beta - \frac{2}{R} \right) (i_1(R) k_1(\rho) - i_1(\rho) k_1(R)) \right]. \quad (4.12)$$

Substituting (4.8) into the boundary condition (4.3) and using (4.10), (3.1) again, we have

$$0 = \sigma^+ \Big|_{\partial D_\varepsilon} - \underline{\sigma} = \hat{\sigma} \Big|_{\partial D_\varepsilon} + \varepsilon \sigma_1^+ \Big|_{\partial D_\varepsilon} - \underline{\sigma} + O(\varepsilon^2)$$

$$\begin{aligned}
&= \sigma_s(\rho) + \sigma'_s(\rho)\varepsilon T + \varepsilon\sigma_1^+ \Big|_{\partial B_\rho} - \underline{\sigma} + O(\varepsilon^2) \\
&= \varepsilon\sigma_1^+ \Big|_{\partial B_\rho} + O(\varepsilon^2), \\
0 &= \sigma^- \Big|_{\partial D_\varepsilon} - \underline{\sigma} = \hat{\sigma}(\rho) + \varepsilon\sigma_1^- \Big|_{\partial D_\varepsilon} - \underline{\sigma} + O(\varepsilon^2) \\
&= \varepsilon\sigma_1^- \Big|_{\partial B_\rho} + O(\varepsilon^2).
\end{aligned}$$

Analogously,

$$\begin{aligned}
0 &= \frac{\partial\sigma^+}{\partial\mathbf{n}} \Big|_{\partial D_\varepsilon} = \frac{\partial\sigma^+}{\partial r} \Big|_{\partial D_\varepsilon} - \frac{\varepsilon}{\rho^2} \nabla_\omega \sigma^+ \Big|_{\partial D_\varepsilon} \cdot \nabla_\omega T + O(\varepsilon^2) \\
&= \frac{\partial\hat{\sigma}}{\partial r} \Big|_{\partial D_\varepsilon} + \varepsilon \frac{\partial\sigma_1^+}{\partial r} \Big|_{\partial D_\varepsilon} + O(\varepsilon^2) \\
&= \hat{\sigma}'(\rho) + \hat{\sigma}''(\rho)\varepsilon T + \varepsilon \frac{\partial\sigma_1^+}{\partial r} \Big|_{\partial B_\rho} + O(\varepsilon^2) \\
&= \varepsilon \left(\frac{\partial\sigma_1^+}{\partial r} \Big|_{\partial B_\rho} + \underline{\sigma} T \right) + O(\varepsilon^2).
\end{aligned}$$

Hence, σ_1 satisfies:

$$-\Delta\sigma_1 = 0 \quad \text{in } B_\rho, \quad (4.13)$$

$$-\Delta\sigma_1 + \sigma_1 = 0 \quad \text{in } B_R \setminus \overline{B}_\rho, \quad (4.14)$$

$$\sigma_1 = 0, \quad \frac{\partial\sigma_1^+}{\partial r} = -\underline{\sigma} T \quad \text{on } \partial B_\rho, \quad (4.15)$$

$$\frac{\partial\sigma_1}{\partial r} + \beta\sigma_1 = -\lambda S \quad \text{on } \partial B_R. \quad (4.16)$$

We now expand the boundary conditions for p . By the expression from [24]

$$\kappa \Big|_{\partial\Omega_\varepsilon} = \frac{1}{R} - \frac{\varepsilon}{R^2} \left(S + \frac{1}{2} \Delta_\omega S \right) + \varepsilon^2 f_2,$$

where $\|f_2\|_{C^{l-2+\alpha}(\Sigma)} \leq C_l$ if $\|S\|_{C^{l+\alpha}(\Sigma)} \leq 1$, we deduce from (4.6), (4.9), (4.11) and (3.5) that

$$\begin{aligned}
0 &= \hat{p} \Big|_{\partial\Omega_\varepsilon} + \varepsilon p_1 \Big|_{\partial\Omega_\varepsilon} - \frac{1}{R} + \frac{\varepsilon}{R^2} \left(S + \frac{1}{2} \Delta_\omega S \right) + O(\varepsilon^2) \\
&= p_s \Big|_{\partial B_R} + \frac{\partial p_s}{\partial r} \Big|_{\partial B_R} \varepsilon S + \varepsilon p_1 \Big|_{\partial B_R} - \frac{1}{R} + \frac{\varepsilon}{R^2} \left(S + \frac{1}{2} \Delta_\omega S \right) + O(\varepsilon^2) \\
&= \varepsilon \left[p_1 \Big|_{\partial B_R} + \frac{1}{R^2} \left(S + \frac{1}{2} \Delta_\omega S \right) \right] + O(\varepsilon^2).
\end{aligned}$$

The boundary condition (4.5) together with (4.9), (4.11) and (3.5) leads to

$$0 = p^+ \Big|_{\partial D_\varepsilon} - p^- \Big|_{\partial D_\varepsilon} = \hat{p} \Big|_{\partial D_\varepsilon} + \varepsilon p_1^+ \Big|_{\partial D_\varepsilon} - \hat{p}(\rho) - \varepsilon p_1^- \Big|_{\partial D_\varepsilon} + O(\varepsilon^2) = \varepsilon [p_1] \Big|_{\partial B_\rho} + O(\varepsilon^2),$$

and

$$\begin{aligned}
0 &= \frac{\partial p^+}{\partial \mathbf{n}} \Big|_{\partial D_\varepsilon} - \frac{\partial p^-}{\partial \mathbf{n}} \Big|_{\partial D_\varepsilon} \\
&= \frac{\partial p^+}{\partial r} \Big|_{\partial D_\varepsilon} - \frac{\partial p^-}{\partial r} \Big|_{\partial D_\varepsilon} - \frac{\varepsilon}{\rho^2} \left(\nabla_\omega p^+ \Big|_{\partial D_\varepsilon} \cdot \nabla_\omega T - \nabla_\omega p^- \Big|_{\partial D_\varepsilon} \cdot \nabla_\omega T \right) + O(\varepsilon^2) \\
&= \frac{\partial \hat{p}}{\partial r} \Big|_{\partial D_\varepsilon} + \varepsilon \frac{\partial p_1^+}{\partial r} \Big|_{\partial D_\varepsilon} - \varepsilon \frac{\partial p_1^-}{\partial r} \Big|_{\partial D_\varepsilon} + O(\varepsilon^2) \\
&= \frac{\partial^2 \hat{p}}{\partial r^2} \Big|_{\partial B_\rho} \varepsilon T + \varepsilon \frac{\partial p_1^+}{\partial r} \Big|_{\partial B_\rho} - \varepsilon \frac{\partial p_1^-}{\partial r} \Big|_{\partial B_\rho} + O(\varepsilon^2) \\
&= \varepsilon \left([\partial_{\mathbf{n}} p_1] \Big|_{\partial B_\rho} + \mu(\tilde{\sigma} - \underline{\sigma})T \right) + O(\varepsilon^2).
\end{aligned}$$

Therefore, p_1 satisfies

$$-\Delta p_1 = 0 \quad \text{in } B_\rho, \quad (4.17)$$

$$-\Delta p_1 = \mu\sigma_1 \quad \text{in } B_R \setminus \overline{B}_\rho, \quad (4.18)$$

$$[p_1] = 0, \quad [\partial_{\mathbf{n}} p_1] = -\mu(\tilde{\sigma} - \underline{\sigma})T \quad \text{on } \partial B_\rho, \quad (4.19)$$

$$p_1 = -\frac{1}{R^2} \left(S + \frac{1}{2} \Delta_\omega S \right) \quad \text{on } \partial B_R. \quad (4.20)$$

We next compute explicitly the functions σ_1 , p_1 and T . If we write

$$S(\theta, \varphi) = \sum_{n=0}^{\infty} \sum_{m=-n}^n a_{n,m} Y_{n,m}(\theta, \varphi), \quad (4.21)$$

then solving (4.14)-(4.16) by the separation of variables and using (2.1), we get

$$\sigma_1(r, \theta, \varphi) = -\lambda \sum_{n=0}^{\infty} \sum_{m=-n}^n a_{n,m} Q_n(r) Y_{n,m}(\theta, \varphi) \quad \text{in } B_R \setminus \overline{B}_\rho, \quad (4.22)$$

where $Q_n(r)$ satisfies

$$Q_n''(r) + \frac{2}{r} Q_n'(r) - \left(1 + \frac{n(n+1)}{r^2} \right) Q_n(r) = 0, \quad \rho < r < R, \quad (4.23)$$

$$Q_n(\rho) = 0, \quad Q_n'(R) + \beta Q_n(R) = 1, \quad (4.24)$$

i.e.,

$$Q_n(r) = \frac{i_n(r)k_n(\rho) - i_n(\rho)k_n(r)}{\left(\frac{n}{R} + \beta\right) (i_n(R)k_n(\rho) - i_n(\rho)k_n(R)) + i_n(\rho)k_{n+1}(R) + i_{n+1}(R)k_n(\rho)} \quad (4.25)$$

for $\rho < r < R$. Substituting (4.22) into the second equation in (4.15) yields

$$T(\theta, \varphi) = \frac{\lambda}{\underline{\sigma}} \sum_{n=0}^{\infty} \sum_{m=-n}^n a_{n,m} Q_n'(\rho) Y_{n,m}(\theta, \varphi). \quad (4.26)$$

Similarly, setting

$$p_1(r, \theta, \varphi) = \sum_{n=0}^{\infty} \sum_{m=-n}^n a_{n,m} P_n(r) Y_{n,m}(\theta, \varphi) \quad (4.27)$$

in (4.17)-(4.20), we are led to

$$\begin{aligned} P_n''(r) + \frac{2}{r} P_n'(r) - \frac{n(n+1)}{r^2} P_n(r) &= 0 \quad \text{for } 0 < r < \rho, \\ P_n''(r) + \frac{2}{r} P_n'(r) - \frac{n(n+1)}{r^2} P_n(r) &= \lambda \mu Q_n(r) \quad \text{for } \rho < r < R, \\ P_0'(0) &= 0, \quad P_n(0) = 0 \quad \text{for } n \geq 1, \\ P_n(\rho+) &= P_n(\rho-), \quad P_n'(\rho+) - P_n'(\rho-) = -\lambda \mu \frac{\tilde{\sigma} - \underline{\sigma}}{\underline{\sigma}} Q_n'(\rho), \\ P_n(R) &= -\frac{1}{R^2} \left(1 - \frac{n(n+1)}{2} \right). \end{aligned}$$

Solving the ODE, we obtain

$$P_n(r) = \frac{\lambda \mu \tilde{\sigma} \rho^{n+2} Q_n'(\rho)}{(2n+1)\underline{\sigma}} \left(\frac{1}{\rho^{2n+1}} - \frac{1}{R^{2n+1}} \right) r^n - \lambda \mu \frac{r^n}{R^n} Q_n(R) - \frac{r^n}{R^{n+2}} \left(1 - \frac{n(n+1)}{2} \right) \quad (4.28)$$

for $0 < r \leq \rho$, and

$$\begin{aligned} P_n(r) &= \frac{\lambda \mu \tilde{\sigma} \rho^{n+2} Q_n'(\rho)}{(2n+1)\underline{\sigma}} \left(\frac{1}{r^{2n+1}} - \frac{1}{R^{2n+1}} \right) r^n + \lambda \mu \left(Q_n(r) - \frac{r^n}{R^n} Q_n(R) \right) \\ &\quad - \frac{r^n}{R^{n+2}} \left(1 - \frac{n(n+1)}{2} \right) \quad \text{for } \rho < r < R. \end{aligned} \quad (4.29)$$

To proceed further, we need more information on Q_n .

Lemma 4.1. $\{Q_n'(\rho)\}$ is a positive, monotonically decreasing sequence in n , while $\{Q_n'(R)\}$ is a positive, monotonically increasing sequence in n .

Proof. From (4.25) it follows that

$$Q_n'(r) = \frac{i_n'(r)k_n(\rho) - i_n(\rho)k_n'(r)}{\left(\frac{n}{R} + \beta\right)(i_n(R)k_n(\rho) - i_n(\rho)k_n(R)) + i_n(\rho)k_{n+1}(R) + i_{n+1}(R)k_n(\rho)},$$

which together with (2.6) implies that $Q_n'(r) > 0$, $\rho \leq r \leq R$; particularly, $Q_n'(\rho) > 0$, $Q_n'(R) > 0$. Moreover, from (4.24),

$$0 = Q_n(\rho) < Q_n(r) \leq Q_n(R) < \frac{1}{\frac{n}{R} + \beta} \quad \text{for } \rho < r \leq R. \quad (4.30)$$

Let $F_n(r) = Q_{n+1}(r) - Q_n(r)$. Then by virtue of (4.23) and (4.24), F_n satisfies

$$\begin{cases} -\Delta F_n + \left(1 + \frac{n(n+1)}{r^2}\right) F_n = -\frac{2(n+1)}{r^2} Q_{n+1} & \text{in } B_R \setminus \overline{B}_\rho, \\ F_n = 0 & \text{on } \partial B_\rho, \\ \frac{\partial F_n}{\partial \mathbf{n}} + \beta F_n = 0 & \text{on } \partial B_R. \end{cases}$$

By (4.30), we apply the maximum principle to conclude that $F_n(r) < 0$ for $\rho < r \leq R$, $F'_n(\rho) < 0$ and $F'_n(R) > 0$. The proof is complete. \square

Remark 4.1. *In fact, from the proof of Lemma 4.1, one can also see that for every $\rho < r \leq R$, $\{Q_n(r)\}$ is a positive, monotonically decreasing sequence in n , and $\lim_{n \rightarrow \infty} Q_n(r) = 0$ by (4.30). In particular,*

$$\lim_{n \rightarrow \infty} Q_n(R) = 0.$$

As a result,

$$\lim_{n \rightarrow \infty} Q'_n(R) = \lim_{n \rightarrow \infty} (1 - \beta Q_n(R)) = 1. \quad (4.31)$$

Lemma 4.2. *Given $\rho < r \leq R$, $\{Q'_n(r)/Q_n(r) - n/r\}$ is a positive, monotonically decreasing sequence in n . Moreover,*

$$\lim_{n \rightarrow \infty} \left(\frac{Q'_n(R)}{Q_n(R)} - \frac{n}{R} \right) = 0. \quad (4.32)$$

Proof. For given $s \in (\rho, R]$, from (4.23), (4.24) we see that $G_n(r)$ defined by

$$G_n(r) = \frac{r}{s} Q_n(r) - \frac{Q_n(s)}{Q_{n+1}(s)} Q_{n+1}(r),$$

satisfies

$$\begin{cases} -\Delta G_n + \left[1 + \frac{(n+1)(n+2)}{r^2} \right] G_n = -\frac{2}{s} (Q'_n(r) - \frac{n}{r} Q_n(r)) & \text{in } B_s \setminus \overline{B}_\rho, \\ G_n = 0 & \text{on } \partial B_\rho, \quad G_n = 0 & \text{on } \partial B_s. \end{cases}$$

A direct calculation based on (4.25), (2.5) gives that

$$\begin{aligned} & Q'_n(r) \\ &= \frac{\frac{n}{r} [i_n(r)k_n(\rho) - i_n(\rho)k_n(r)] + [i_{n+1}(r)k_n(\rho) + i_n(\rho)k_{n+1}(r)]}{\left(\frac{n}{R} + \beta\right) [i_n(R)k_n(\rho) - i_n(\rho)k_n(R)] + i_n(\rho)k_{n+1}(R) + i_{n+1}(R)k_n(\rho)} \\ &= \frac{n}{r} Q_n(r) + \frac{i_{n+1}(r)k_n(\rho) + i_n(\rho)k_{n+1}(r)}{\left(\frac{n}{R} + \beta\right) [i_n(R)k_n(\rho) - i_n(\rho)k_n(R)] + i_n(\rho)k_{n+1}(R) + i_{n+1}(R)k_n(\rho)}, \end{aligned} \quad (4.33)$$

which implies that

$$Q'_n(r) - \frac{n}{r} Q_n(r) > 0, \quad \rho \leq r \leq R.$$

Thus, $G_n(r) < 0$ for $\rho < r < s$ and $G'_n(s) > 0$ by the maximum principle. Combining with the definition of $G_n(r)$, we further have

$$0 < \frac{Q'_{n+1}(s)}{Q_{n+1}(s)} - \frac{n+1}{s} < \frac{Q'_n(s)}{Q_n(s)} - \frac{n}{s},$$

which proves the monotonicity result.

Now we turn to the proof of (4.32). Using (4.25), (4.33), we compute

$$\frac{Q'_n(R)}{Q_n(R)} - \frac{n}{R} = \frac{i_{n+1}(R)k_n(\rho) + i_n(\rho)k_{n+1}(R)}{i_n(R)k_n(\rho) - i_n(\rho)k_n(R)} = \frac{\frac{i_{n+1}(R)}{i_n(R)} + \frac{i_n(\rho)}{i_n(R)} \frac{k_{n+1}(R)}{k_n(\rho)}}{1 - \frac{i_n(\rho)}{i_n(R)} \frac{k_n(R)}{k_n(\rho)}}.$$

By (2.8), (2.9) and the fact $0 < \rho < R$, letting $n \rightarrow \infty$ in the above equality yields (4.32). The proof is complete. \square

We conclude this section by rigorously establishing the expansions (4.8), (4.9) by estimating the $O(\varepsilon^2)$ terms in the $C^{2+\alpha}$ -norm. To do that, observing that (σ, p) is defined in Ω_ε while (σ_1, p_1) is defined only in B_R , we will first transform the domain B_R into Ω_ε by introducing the following Hanzawa-type transformation

$$(r, \theta, \varphi) = H_\varepsilon(r', \theta', \varphi') = (r' + \chi(\rho - r')\varepsilon T(\theta', \varphi') + \chi(R - r')\varepsilon S(\theta', \varphi'), \theta', \varphi'), \quad (4.34)$$

where

$$\chi \in C^\infty, \quad \chi(z) = \begin{cases} 0, & \text{if } |z| \geq 3\delta_0/4 \\ 1, & \text{if } |z| < \delta_0/4 \end{cases}, \quad \left| \frac{d^k \chi}{dz^k} \right| \leq \frac{C}{\delta_0^k}$$

with δ_0 being a small positive constant satisfying $\delta_0 < \min\{\frac{4}{3}\rho, \frac{2}{3}(R - \rho)\}$. It is easy to see that H_ε maps B_R (B_ρ) into Ω_ε (D_ε) while keeping the ball $\{r < \rho - \frac{3}{4}\delta_0\}$ fixed, and the inverse transformation H_ε^{-1} maps Ω_ε (D_ε) into B_R (B_ρ). Let

$$\begin{aligned} \tilde{\sigma}_1(r, \theta, \varphi) &= \sigma_1(H_\varepsilon^{-1}(r, \theta, \varphi)) \quad \text{in } \Omega_\varepsilon, \\ \tilde{p}_1(r, \theta, \varphi) &= p_1(H_\varepsilon^{-1}(r, \theta, \varphi)) \quad \text{in } \Omega_\varepsilon. \end{aligned} \quad (4.35)$$

Then $(\tilde{\sigma}_1, \tilde{p}_1)$ is well defined in Ω_ε .

Next, if S , given by (4.21), belongs to $C^{4+\alpha}(\Sigma)$, and is π -periodic in θ and 2π -periodic in φ , then T admits the representation (4.26) and satisfies

$$\|T\|_{C(\Sigma)} \leq C\|T\|_{H^{3/2}(\Sigma)} \leq C\|S\|_{H^{3/2}(\Sigma)}$$

by Lemmas 2.1 and 4.1. Furthermore,

$$\|T\|_{C^{2+\alpha}(\Sigma)} \leq C\|T\|_{C^3(\Sigma)} \leq C\|S\|_{H^{9/2}(\Sigma)} \leq C\|S\|_{C^{4+\alpha}(\Sigma)}, \quad (4.36)$$

provided that $1/2 < \alpha < 1$.

We now proceed to establish the following.

Lemma 4.3. *Assume that $S \in C^{4+\alpha}(\Sigma)$ with $1/2 < \alpha < 1$ and $\|S\|_{C^{4+\alpha}(\Sigma)} \leq 1$, and is given by (4.21), π -periodic in θ and 2π -periodic in φ . Then the estimates*

$$\|\sigma - \hat{\sigma}(\rho) - \varepsilon \tilde{\sigma}_1\|_{C^{2+\alpha}(\overline{D_\varepsilon})} + \|\sigma - \hat{\sigma} - \varepsilon \tilde{\sigma}_1\|_{C^{2+\alpha}(\overline{\Omega_\varepsilon \setminus D_\varepsilon})} \leq C|\varepsilon|^2 \|S\|_{C^{4+\alpha}(\Sigma)}, \quad (4.37)$$

$$\|p - \hat{p}(\rho) - \varepsilon \tilde{p}_1\|_{C^{2+\alpha}(\overline{D_\varepsilon})} + \|p - \hat{p} - \varepsilon \tilde{p}_1\|_{C^{2+\alpha}(\overline{\Omega_\varepsilon \setminus D_\varepsilon})} \leq C|\varepsilon|^2 \|S\|_{C^{4+\alpha}(\Sigma)} \quad (4.38)$$

are valid uniformly for small $|\varepsilon|$ with C being independent of ε and S .

Proof. By (4.13)-(4.16), we see that $\sigma_1 \equiv 0$ in B_ρ . Applying the Schauder estimates, we obtain

$$\|\sigma_1\|_{C^{2+\alpha}(\overline{B_R \setminus B_\rho})} \leq C\|S\|_{C^{1+\alpha}(\Sigma)}.$$

Moreover, we derive for $\tilde{\sigma}_1$ the problem

$$\tilde{\sigma}_1 \equiv 0 \quad \text{in } D_\varepsilon,$$

$$\begin{aligned}
-\Delta \tilde{\sigma}_1 + \tilde{\sigma}_1 &= \tilde{f}_1 \quad \text{in } \Omega_\varepsilon \setminus D_\varepsilon, \\
\tilde{\sigma}_1 &= 0 \quad \text{on } \partial D_\varepsilon, \\
\frac{\partial \tilde{\sigma}_1}{\partial \mathbf{n}} + \beta \tilde{\sigma}_1 &= -\lambda S + \tilde{g}_1 \quad \text{on } \partial \Omega_\varepsilon,
\end{aligned}$$

where \tilde{f}_1/ε involves at most second order derivatives of T , S and σ_1 , \tilde{g}_1/ε involves at most first order derivatives of S and σ_1 , and

$$\|\tilde{f}_1\|_{C^\alpha(\overline{\Omega_\varepsilon \setminus D_\varepsilon})} \leq C|\varepsilon|(\|T\|_{C^{2+\alpha}(\Sigma)} + \|S\|_{C^{2+\alpha}(\Sigma)}), \quad (4.39)$$

$$\|\tilde{g}_1\|_{C^{1+\alpha}(\Sigma)} \leq C|\varepsilon|\|S\|_{C^{2+\alpha}(\Sigma)}. \quad (4.40)$$

Set

$$\psi = \begin{cases} \sigma - \hat{\sigma} - \varepsilon \tilde{\sigma}_1 & \text{in } \Omega_\varepsilon \setminus D_\varepsilon, \\ \sigma - \hat{\sigma}(\rho) - \varepsilon \tilde{\sigma}_1 & \text{in } D_\varepsilon. \end{cases}$$

Then, we find

$$\begin{aligned}
\psi &\equiv 0 \quad \text{in } D_\varepsilon, \\
-\Delta \psi + \psi &= -\varepsilon \tilde{f}_1 \quad \text{in } \Omega_\varepsilon \setminus D_\varepsilon, \\
\psi^+ &= \hat{\sigma}(\rho) - \hat{\sigma}(\rho + \varepsilon T) \quad \text{on } \partial D_\varepsilon, \\
\frac{\partial \psi}{\partial \mathbf{n}} + \beta \psi &= -\varepsilon \tilde{g}_1 + \tilde{g}_2 \quad \text{on } \partial \Omega_\varepsilon,
\end{aligned}$$

where

$$\|\hat{\sigma}(\rho) - \hat{\sigma}(\rho + \varepsilon T)\|_{C^{2+\alpha}(\Sigma)} \leq C|\varepsilon|^2\|T\|_{C^{2+\alpha}(\Sigma)}, \quad (4.41)$$

and similarly to \tilde{k}_1 in [36, Lemma 3.1],

$$\|\tilde{g}_2\|_{C^{1+\alpha}(\Sigma)} \leq C|\varepsilon|^2\|S\|_{C^{2+\alpha}(\Sigma)}. \quad (4.42)$$

Combining (4.39)-(4.42), we get

$$\|\psi\|_{C^{2+\alpha}(\overline{\Omega_\varepsilon \setminus D_\varepsilon})} \leq C|\varepsilon|^2(\|T\|_{C^{2+\alpha}(\Sigma)} + \|S\|_{C^{2+\alpha}(\Sigma)}), \quad (4.43)$$

which together with (4.36) proves (4.37).

We now proceed with the proof of (4.38). It should be pointed out that, being different from that for σ (which equals to a constant in the dead-core region), here we need establish Schauder estimates for diffraction problems by using the estimates near the boundary for elliptic systems [1]. Precisely, for the problem (4.17)-(4.20) there holds

$$\|p_1\|_{C^{2+\alpha}(\overline{B_\rho})} + \|p_1\|_{C^{2+\alpha}(\overline{B_R \setminus B_\rho})} \leq C(\|T\|_{C^{2+\alpha}(\Sigma)} + \|S\|_{C^{4+\alpha}(\Sigma)}). \quad (4.44)$$

Denote

$$\phi = \begin{cases} p - \hat{p} - \varepsilon \tilde{p}_1 & \text{in } \Omega_\varepsilon \setminus D_\varepsilon, \\ p - \hat{p}(\rho) - \varepsilon \tilde{p}_1 & \text{in } D_\varepsilon. \end{cases}$$

By a similar procedure as before, we obtain

$$\begin{aligned}
-\Delta\phi &= -\varepsilon\tilde{f}_2 \quad \text{in } D_\varepsilon, \\
-\Delta\phi &= \mu(\sigma - \hat{\sigma} - \varepsilon\tilde{\sigma}_1) - \varepsilon\tilde{f}_3 \quad \text{in } \Omega_\varepsilon \setminus D_\varepsilon, \\
[\phi] &= \hat{p}(\rho) - \hat{p}(\rho + \varepsilon T), \quad [\partial_{\mathbf{n}}\phi] = \tilde{g}_3 \quad \text{on } \partial D_\varepsilon, \\
\phi &= \tilde{g}_4 \quad \text{on } \partial\Omega_\varepsilon,
\end{aligned}$$

where

$$\begin{aligned}
\|\tilde{f}_2\|_{C^\alpha(\overline{D_\varepsilon})} + \|\tilde{f}_3\|_{C^\alpha(\overline{\Omega_\varepsilon \setminus D_\varepsilon})} &\leq C|\varepsilon|(\|T\|_{C^{2+\alpha}(\Sigma)} + \|S\|_{C^{4+\alpha}(\Sigma)}), \\
\|\hat{p}(\rho) - \hat{p}(\rho + \varepsilon T)\|_{C^{2+\alpha}(\Sigma)} &\leq C|\varepsilon|^2\|T\|_{C^{2+\alpha}(\Sigma)}, \\
\|\tilde{g}_3\|_{C^{1+\alpha}(\Sigma)} &\leq C|\varepsilon|^2\|T\|_{C^{2+\alpha}(\Sigma)}, \\
\|\tilde{g}_4\|_{C^{2+\alpha}(\Sigma)} &\leq C|\varepsilon|^2\|S\|_{C^{4+\alpha}(\Sigma)}.
\end{aligned}$$

Using (4.43), the Schauder estimates for diffraction problems yield

$$\|\phi\|_{C^{2+\alpha}(\overline{D_\varepsilon})} + \|\phi\|_{C^{2+\alpha}(\overline{\Omega_\varepsilon \setminus D_\varepsilon})} \leq C|\varepsilon|^2(\|T\|_{C^{2+\alpha}(\Sigma)} + \|S\|_{C^{4+\alpha}(\Sigma)}),$$

and then (4.38) follows immediately from (4.36). The proof is complete. \square

5 Symmetry-breaking solutions

In this section, regarding (1.9)-(1.16) as a bifurcation problem with a bifurcation parameter μ (see (4.7)), we prove the existence of symmetry-breaking solutions by using the Crandall-Rabinowitz theorem.

By (4.36) and (4.44),

$$\left. \frac{\partial(\hat{p} + \varepsilon\tilde{p}_1)}{\partial \mathbf{n}} \right|_{\partial\Omega_\varepsilon} = \varepsilon \left[\left. \frac{\partial^2 \hat{p}}{\partial r^2} \right|_{\partial B_R} S + \left. \frac{\partial p_1}{\partial r} \right|_{\partial B_R} \right] + O(|\varepsilon|^2\|S\|_{C^{4+\alpha}(\Sigma)}),$$

which together with (4.11) and (4.38) leads to

$$F(\tilde{R}, \mu) = \varepsilon \left[\left. \frac{\partial^2 p_s}{\partial r^2} \right|_{\partial B_R} S + \left. \frac{\partial p_1}{\partial r} \right|_{\partial B_R} \right] + O(|\varepsilon|^2\|S\|_{C^{4+\alpha}(\Sigma)}). \quad (5.1)$$

As in [19, 36], we introduce the Banach spaces

$$X^{l+\alpha} = \{\tilde{R} \in C^{l+\alpha}(\Sigma), \tilde{R} \text{ is } \pi\text{-periodic in } \theta, 2\pi\text{-periodic in } \varphi\},$$

$$X_2^{l+\alpha} = \text{closure of the linear space spanned by } \{Y_{n,0}(\theta), n = 0, 2, 4, \dots\} \text{ in } X^{l+\alpha}.$$

Take $X = X_2^{4+\alpha}$ and $Y = X_2^{1+\alpha}$ with $1/2 < \alpha < 1$. Noticing that $Y_{n,0}(\pi - \theta) = Y_{n,0}(\theta)$ if and only if n is even, $X_2^{l+\alpha}$ coincides with the subspace of the $C^{l+\alpha}(\Sigma)$ -closure of the smooth functions consisting of those functions u that are independent of φ and satisfy $u(\theta) = u(\pi - \theta)$.

Thus, F maps X into Y . The relation (5.1) shows that the mapping $(\tilde{R}, \mu) \rightarrow F(\tilde{R}, \mu)$ from $X_2^{l+3+\alpha}$ to $X_2^{l+\alpha}$ is bounded if $l = 1$, and the same argument shows that the same is true for any $l \geq 1$. A similar argument shows that this mapping is Fréchet differentiable in (\tilde{R}, μ) ; furthermore $\partial F(\tilde{R}, \mu)/\partial \tilde{R}$ (or $\partial F(\tilde{R}, \mu)/\partial \mu$) is obtained by solving a linearized problem about (\tilde{R}, μ) with respect to \tilde{R} (or μ). By using the Schauder estimates we can then further obtain differentiability of $F(\tilde{R}, \mu)$ to any order.

In view of (5.1), the Fréchet derivative of $F(\tilde{R}, \mu)$ in \tilde{R} at $(0, \mu)$ is given as follows:

$$[F_{\tilde{R}}(0, \mu)]S = \left. \frac{\partial^2 p_s}{\partial r^2} \right|_{\partial B_R} S + \left. \frac{\partial p_1}{\partial r} \right|_{\partial B_R}. \quad (5.2)$$

Using (3.4), (3.5), we have

$$\left. \frac{\partial^2 p_s}{\partial r^2} \right|_{\partial B_R} = -\mu(\sigma_s(R) - \tilde{\sigma}), \quad (5.3)$$

and by (4.27), (4.29),

$$\begin{aligned} \left. \frac{\partial p_1}{\partial r} \right|_{\partial B_R} &= \sum_{n=0}^{\infty} \sum_{m=-n}^n a_{n,m} \left[\frac{n}{R^3} \left(\frac{n(n+1)}{2} - 1 \right) + \lambda \mu \left(Q'_n(R) - \frac{n}{R} Q_n(R) \right) \right. \\ &\quad \left. - \lambda \mu \frac{\tilde{\sigma}}{\underline{\sigma}} \frac{\rho^{n+2}}{R^{n+2}} Q'_n(\rho) \right] Y_{n,m}(\theta, \varphi). \end{aligned} \quad (5.4)$$

Substituting (5.3) and (5.4) into (5.2), we arrive at

$$\begin{aligned} [F_{\tilde{R}}(0, \mu)]S &= \sum_{n=0}^{\infty} \sum_{m=-n}^n a_{n,m} \left[\frac{n}{R^3} \left(\frac{n(n+1)}{2} - 1 \right) + \lambda \mu \left(Q'_n(R) - \frac{n}{R} Q_n(R) \right) \right. \\ &\quad \left. - \lambda \mu \frac{\tilde{\sigma}}{\underline{\sigma}} \frac{\rho^{n+2}}{R^{n+2}} Q'_n(\rho) - \mu(\sigma_s(R) - \tilde{\sigma}) \right] Y_{n,m}(\theta, \varphi). \end{aligned}$$

In particular,

$$[F_{\tilde{R}}(0, \mu)]Y_{n,m} = (A_n - \mu B_n)Y_{n,m},$$

where

$$A_n = \frac{n}{R^3} \left(\frac{n(n+1)}{2} - 1 \right), \quad (5.5)$$

$$B_n = \sigma_s(R) - \tilde{\sigma} - \lambda \left[\left(Q'_n(R) - \frac{n}{R} Q_n(R) \right) - \frac{\tilde{\sigma}}{\underline{\sigma}} \frac{\rho^{n+2}}{R^{n+2}} Q'_n(\rho) \right]. \quad (5.6)$$

Obviously,

$$A_0 = A_1 = 0, \quad A_n > 0 \quad \text{for } n \geq 2.$$

As will be shown in the next lemma, $B_0 < 0$ and $B_1 = 0$.

Lemma 5.1. $B_0 < 0$ and $B_1 = 0$.

Proof. Combining (3.2), (3.7), (4.12) and (5.6), we derive

$$B_n = \frac{2}{\pi} \underline{\sigma} \rho^2 H_n \quad (5.7)$$

with

$$\begin{aligned} H_n = & i_0(R)k_1(\rho) + i_1(\rho)k_0(R) - \frac{3R^2}{R^3 - \rho^3} (i_1(R)k_1(\rho) - i_1(\rho)k_1(R)) \\ & - \left[i_0(R)k_1(\rho) + i_1(\rho)k_0(R) + \left(\beta - \frac{2}{R} \right) (i_1(R)k_1(\rho) - i_1(\rho)k_1(R)) \right] \\ & \left[\left(Q'_n(R) - \frac{n}{R} Q_n(R) \right) - \frac{\tilde{\sigma}}{\underline{\sigma}} \frac{\rho^{n+2}}{R^{n+2}} \left(Q'_n(\rho) - \frac{n}{\rho} Q_n(\rho) \right) \right], \end{aligned} \quad (5.8)$$

where we have employed the fact that $Q_n(\rho) = 0$. Besides, (3.7) also implies

$$\frac{\tilde{\sigma}}{\underline{\sigma}} = \frac{3R^2}{R^3 - \rho^3} \frac{2}{\pi} \rho^2 (i_1(R)k_1(\rho) - i_1(\rho)k_1(R)), \quad (5.9)$$

and (4.33) together with (2.7) gives

$$\begin{aligned} & Q'_n(R) - \frac{n}{R} Q_n(R) \\ = & \frac{i_{n+1}(R)k_n(\rho) + i_n(\rho)k_{n+1}(R)}{\left(\frac{n}{R} + \beta \right) [i_n(R)k_n(\rho) - i_n(\rho)k_n(R)] + i_n(\rho)k_{n+1}(R) + i_{n+1}(R)k_n(\rho)}, \end{aligned} \quad (5.10)$$

$$\begin{aligned} & Q'_n(\rho) - \frac{n}{\rho} Q_n(\rho) \\ = & \frac{i_{n+1}(\rho)k_n(\rho) + i_n(\rho)k_{n+1}(\rho)}{\left(\frac{n}{R} + \beta \right) [i_n(R)k_n(\rho) - i_n(\rho)k_n(R)] + i_n(\rho)k_{n+1}(R) + i_{n+1}(R)k_n(\rho)} \\ = & \frac{\pi}{2\rho^2 \left(\frac{n}{R} + \beta \right) [i_n(R)k_n(\rho) - i_n(\rho)k_n(R)] + i_n(\rho)k_{n+1}(R) + i_{n+1}(R)k_n(\rho)}. \end{aligned} \quad (5.11)$$

Thus, substituting (5.9), (5.10) and (5.11) into (5.8), we get

$$H_n = \frac{1}{\left(\frac{n}{R} + \beta \right) [i_n(R)k_n(\rho) - i_n(\rho)k_n(R)] + i_n(\rho)k_{n+1}(R) + i_{n+1}(R)k_n(\rho)} W_n, \quad (5.12)$$

where

$$\begin{aligned} W_n = & \left[\left(\frac{n}{R} + \beta \right) (i_n(R)k_n(\rho) - i_n(\rho)k_n(R)) + (i_n(\rho)k_{n+1}(R) + i_{n+1}(R)k_n(\rho)) \right] \\ & \left[(i_0(R)k_1(\rho) + i_1(\rho)k_0(R)) - \frac{3R^2}{R^3 - \rho^3} (i_1(R)k_1(\rho) - i_1(\rho)k_1(R)) \right] \\ & - \left[(i_0(R)k_1(\rho) + i_1(\rho)k_0(R)) + \left(\beta - \frac{2}{R} \right) (i_1(R)k_1(\rho) - i_1(\rho)k_1(R)) \right] \\ & \left[(i_n(\rho)k_{n+1}(R) + i_{n+1}(R)k_n(\rho)) - \frac{3R^2}{R^3 - \rho^3} \frac{\rho^{n+2}}{R^{n+2}} (i_1(R)k_1(\rho) - i_1(\rho)k_1(R)) \right] \\ = & M_{1,n} + M_{2,n}\beta \end{aligned} \quad (5.13)$$

with

$$\begin{aligned}
M_{1,n} = & \frac{n}{R}(i_n(R)k_n(\rho) - i_n(\rho)k_n(R)) \\
& \left[(i_0(R)k_1(\rho) + i_1(\rho)k_0(R)) - \frac{3R^2}{R^3 - \rho^3}(i_1(R)k_1(\rho) - i_1(\rho)k_1(R)) \right] \\
& + \frac{i_1(R)k_1(\rho) - i_1(\rho)k_1(R)}{R(R^3 - \rho^3)} \left[3R^3 \frac{\rho^{n+2}}{R^{n+2}}(i_0(R)k_1(\rho) + i_1(\rho)k_0(R)) \right. \\
& \quad - 6R^2 \frac{\rho^{n+2}}{R^{n+2}}(i_1(R)k_1(\rho) - i_1(\rho)k_1(R)) \\
& \quad \left. - (R^3 + 2\rho^3)(i_n(\rho)k_{n+1}(R) + i_{n+1}(R)k_n(\rho)) \right], \tag{5.14}
\end{aligned}$$

$$\begin{aligned}
M_{2,n} = & (i_n(R)k_n(\rho) - i_n(\rho)k_n(R)) \\
& \left[(i_0(R)k_1(\rho) + i_1(\rho)k_0(R)) - \frac{3R^2}{R^3 - \rho^3}(i_1(R)k_1(\rho) - i_1(\rho)k_1(R)) \right] \\
& - (i_1(R)k_1(\rho) - i_1(\rho)k_1(R)) \\
& \left[(i_n(\rho)k_{n+1}(R) + i_{n+1}(R)k_n(\rho)) - \frac{3R^2}{R^3 - \rho^3} \frac{\rho^{n+2}}{R^{n+2}}(i_1(R)k_1(\rho) - i_1(\rho)k_1(R)) \right]. \tag{5.15}
\end{aligned}$$

In the sequel, we shall first prove that $B_1 = 0$. By (5.7), (5.12) and (5.13), it suffices to verify that $M_{1,1} = 0$ and $M_{2,1} = 0$. As a matter of fact, by virtue of (5.14), (5.15), we compute

$$\begin{aligned}
M_{1,1} = & \frac{R^3 + 2\rho^3}{R(R^3 - \rho^3)}(i_1(R)k_1(\rho) - i_1(\rho)k_1(R)) \\
& \left[k_1(\rho) \left(i_0(R) - \frac{3}{R}i_1(R) - i_2(R) \right) + i_1(\rho) \left(k_0(R) + \frac{3}{R}k_1(R) - k_2(R) \right) \right], \\
M_{2,1} = & (i_1(R)k_1(\rho) - i_1(\rho)k_1(R)) \\
& \left[k_1(\rho) \left(i_0(R) - \frac{3}{R}i_1(R) - i_2(R) \right) + i_1(\rho) \left(k_0(R) + \frac{3}{R}k_1(R) - k_2(R) \right) \right].
\end{aligned}$$

From (2.3),

$$i_2(s) = i_0(s) - \frac{3}{s}i_1(s), \quad k_2(s) = k_0(s) + \frac{3}{s}k_1(s),$$

from which, $M_{1,1} = M_{2,1} = 0$ follows.

Next, we show $B_0 < 0$ by showing that $M_{1,0} < 0$ and $M_{2,0} < 0$. In view of (5.14),

$$\begin{aligned}
M_{1,0} = & \frac{i_1(R)k_1(\rho) - i_1(\rho)k_1(R)}{R(R^3 - \rho^3)} \left[3R\rho^2(i_0(R)k_1(\rho) + i_1(\rho)k_0(R)) \right. \\
& \left. - 6\rho^2(i_1(R)k_1(\rho) - i_1(\rho)k_1(R)) - (R^3 + 2\rho^3)(i_0(\rho)k_1(R) + i_1(R)k_0(\rho)) \right].
\end{aligned}$$

Denote

$$\begin{aligned}
\xi(s) = & 3Rs^2(i_0(R)k_1(s) + i_1(s)k_0(R)) - 6s^2(i_1(R)k_1(s) - i_1(s)k_1(R)) \\
& - (R^3 + 2s^3)(i_0(s)k_1(R) + i_1(R)k_0(s)) \quad \text{for } 0 < s \leq R.
\end{aligned}$$

Then, by (2.4), (2.5),

$$\xi'(s) = (R^3 + 2s^3)(i_1(R)k_1(s) - i_1(s)k_1(R)) - 3Rs^2(i_0(R)k_0(s) - i_0(s)k_0(R))$$

for $0 < s < R$. Furthermore, observing that

$$R^3 + 2s^3 = (R - s)^2(R + 2s) + 3Rs^2,$$

and applying (2.12) and (2.13), we get

$$\begin{aligned} \xi'(s) &> 3Rs^2[(i_1(R)k_1(s) - i_1(s)k_1(R)) - (i_0(R)k_0(s) - i_0(s)k_0(R))] \\ &= \frac{3\pi}{2R}[(R - s) \cosh(R - s) - \sinh(R - s)] \end{aligned}$$

for $0 < s < R$. Thus, by (3.10), we deduce that $\xi'(s) > 0$ for $0 < s < R$, which together with $\xi(R) = 0$ implies that $\xi(s) < 0$, $0 < s < R$; in particular, $\xi(\rho) < 0$. Hence, $M_{1,0} < 0$.

It remains to show that $M_{2,0} < 0$. By (5.15),

$$\begin{aligned} M_{2,0} &= (i_0(R)k_0(\rho) - i_0(\rho)k_0(R)) \\ &\quad \left[(i_0(R)k_1(\rho) + i_1(\rho)k_0(R)) - \frac{3R^2}{R^3 - \rho^3}(i_1(R)k_1(\rho) - i_1(\rho)k_1(R)) \right] \\ &\quad - (i_1(R)k_1(\rho) - i_1(\rho)k_1(R)) \\ &\quad \left[(i_0(\rho)k_1(R) + i_1(R)k_0(\rho)) - \frac{3\rho^2}{R^3 - \rho^3}(i_1(R)k_1(\rho) - i_1(\rho)k_1(R)) \right]. \end{aligned}$$

Using (2.12)-(2.14), we further obtain

$$\begin{aligned} M_{2,0} &= \frac{\pi^2}{8R^4\rho^3(R^3 - \rho^3)} \left[-2R^5 + 3R^4\rho - 2R^3 - R^2\rho^3 + 3R^2\rho + 2\rho^3 - 3\rho \right. \\ &\quad \left. - (R^4 - 2R^3\rho - 6R^2\rho^2 + 8R\rho^3 + 6R\rho - \rho^4 - 6\rho^2) \sinh(2(R - \rho)) \right. \\ &\quad \left. - (R^4\rho - 2R^3 - 3R^2\rho^3 - 3R^2\rho + 2R\rho^4 + 12R\rho^2 - 4\rho^3 - 3\rho) \cosh(2(R - \rho)) \right]. \end{aligned}$$

Set

$$\begin{aligned} \eta(s) &= [s^4 - 8Rs^3 + 6(R^2 + 1)s^2 + 2(R^3 - 3R)s - R^4] \sinh(2(R - s)) \\ &\quad + [-2Rs^4 + (3R^2 + 4)s^3 - 12Rs^2 + (-R^4 + 3R^2 + 3)s + 2R^3] \cosh(2(R - s)) \\ &\quad - (R^2 - 2)s^3 + (3R^4 + 3R^2 - 3)s - 2R^5 - 2R^3 \quad \text{for } 0 < s \leq R. \end{aligned}$$

Then

$$M_{2,0} = \frac{\pi^2}{8R^4\rho^3(R^3 - \rho^3)} \eta(\rho). \quad (5.16)$$

Notice that

$$\eta''(s) = 2s\zeta(s) \quad (5.17)$$

with

$$\begin{aligned} \zeta(s) &= [2s^3 + (-6R^2 - 6)s + 4R^3 + 12R] \sinh(2(R - s)) \\ &\quad + [-4Rs^3 + 6R^2s^2 + 12Rs - 2R^4 - 9R^2 - 6] \cosh(2(R - s)) - 3(R^2 - 2), \end{aligned}$$

and

$$\zeta'(s) = e^{-2(R-s)}w(s), \quad (5.18)$$

where

$$\begin{aligned} w(s) = & [(4R-2)s^3 + (-6R^2 - 6R + 3)s^2 + (12R^2 - 12R + 6)s + 2R^4 - 4R^3 + 6R^2 \\ & - 6R + 3]e^{4(R-s)} - (4R+2)s^3 + (6R^2 - 6R - 3)s^2 + (12R^2 + 12R + 6)s \\ & - 2R^4 - 4R^3 - 6R^2 - 6R - 3. \end{aligned}$$

It is easy to see that

$$\eta(R) = \eta'(R) = \zeta(R) = 0. \quad (5.19)$$

Thus, we only need to verify that $w(s) > 0$ for $0 < s < R$, which together with (5.17)-(5.19) implies

$$\eta(s) < 0 \quad \text{for } 0 < s < R.$$

In particular, $\eta(\rho) < 0$ and by (5.16), there holds $M_{2,0} < 0$.

We consider, instead of $w(s)$,

$$\begin{aligned} u(R) = & [2R^4 - 4R^3 + (-6s^2 + 12s + 6)R^2 + (4s^3 - 6s^2 - 12s - 6)R - 2s^3 + 3s^2 \\ & + 6s + 3]e^{4(R-s)} - 2R^4 - 4R^3 + (6s^2 + 12s - 6)R^2 \\ & - (4s^3 + 6s^2 - 12s + 6)R - 2s^3 - 3s^2 + 6s - 3 \end{aligned}$$

for $R \geq s > 0$. A direct calculation gives that

$$u(s) = u'(s) = u''(s) = 0, \quad u^{(3)}(s) = 240s^2 > 0, \quad u^{(4)}(s) = 16(120s^2 + 48s) > 0,$$

and

$$u^{(5)}(R) = 64e^{4(R-s)}v(R)$$

with

$$\begin{aligned} v(R) = & 32R^4 + 96R^3 + (-96s^2 + 192s + 96)R^2 + (64s^3 - 336s^2 + 288s + 24)R \\ & + 48s^3 - 192s^2 + 96s + 3. \end{aligned}$$

Furthermore, there hold

$$v(s) = 192s^2 + 120s + 3 > 0, \quad v'(s) = 8(42s^2 + 60s + 3) > 0$$

and

$$v''(R) = 192(2R^2 + 3R - s^2 + 2s + 1) > 0 \quad \text{for } R > s > 0.$$

On the basis of the above analysis, we conclude that $u(R) > 0$ for $R > s > 0$, i.e., $w(s) > 0$ for $0 < s < R$. The proof is complete. \square

Lemma 5.2. (i) $\lim_{n \rightarrow \infty} B_n = \sigma_s(R) - \tilde{\sigma}$;

(ii) There exists $n^* \in \mathbb{N}$ such that $\mu_n = A_n/B_n$ is positive and monotonically increasing for $n \geq n^*$; moreover, $\lim_{n \rightarrow \infty} \mu_n = +\infty$.

Proof. (i) Applying Remark 4.1 and Lemma 4.2, we get

$$\lim_{n \rightarrow \infty} \left(Q'_n(R) - \frac{n}{R} Q_n(R) \right) = \lim_{n \rightarrow \infty} Q_n(R) \left(\frac{Q'_n(R)}{Q_n(R)} - \frac{n}{R} \right) = 0. \quad (5.20)$$

Besides, from Lemma 4.1 and $0 < \rho < R$, we see that

$$\lim_{n \rightarrow \infty} Q'_n(\rho) \frac{\rho^{n+2}}{R^{n+2}} = 0.$$

The assertion (i) then follows from the definition of B_n .

(ii) Since $\sigma_s(R) - \tilde{\sigma} > 0$ by (3.8), the assertion (i) implies that there exists a positive integer $N_1 \geq 2$ such that $B_n > 0$ for every $n \geq N_1$. Consequently,

$$\mu_n = \frac{A_n}{B_n} \quad (5.21)$$

is well defined and positive for $n \geq N_1$, and

$$\lim_{n \rightarrow \infty} \mu_n = +\infty, \quad \lim_{n \rightarrow \infty} \frac{\mu_n}{n^3} = \frac{1}{2R^3(\sigma_s(R) - \tilde{\sigma})}.$$

We claim that

$$\lim_{n \rightarrow \infty} \frac{\frac{\mu_n}{n^3} - \frac{1}{2R^3(\sigma_s(R) - \tilde{\sigma})}}{\frac{1}{n}} = \frac{1}{2R^3(\sigma_s(R) - \tilde{\sigma})}. \quad (5.22)$$

In fact, using (5.5), (5.6) and (5.21), we compute for $n \geq N_1$,

$$\begin{aligned} \frac{\frac{\mu_n}{n^3} - \frac{1}{2R^3(\sigma_s(R) - \tilde{\sigma})}}{\frac{1}{n}} &= \frac{1}{2R^3} \frac{\frac{1}{B_n} \left(1 + \frac{1}{n} - \frac{2}{n^2} \right) - \frac{1}{\sigma_s(R) - \tilde{\sigma}}}{\frac{1}{n}} \\ &= \frac{1}{2R^3} \left[\frac{1}{B_n} + \frac{\frac{1}{B_n} - \frac{1}{\sigma_s(R) - \tilde{\sigma}}}{\frac{1}{n}} - \frac{2}{n} \frac{1}{B_n} \right] \\ &= \frac{1}{2R^3} \left[\frac{1}{B_n} + \lambda \frac{nQ_n(R) \left(\frac{Q'_n(R)}{Q_n(R)} - \frac{n}{R} \right) - \frac{\tilde{\sigma}}{\sigma} n \frac{\rho^{n+2}}{R^{n+2}} Q'_n(\rho)}{B_n(\sigma_s(R) - \tilde{\sigma})} - \frac{2}{n} \frac{1}{B_n} \right]. \end{aligned} \quad (5.23)$$

Notice that

$$nQ_n(R) = R \left[Q'_n(R) - \left(Q'_n(R) - \frac{n}{R} Q_n(R) \right) \right].$$

Thus, from (4.31) and (5.20), we derive

$$\lim_{n \rightarrow \infty} nQ_n(R) = R. \quad (5.24)$$

Hence, by Lemmas 4.1, 4.2, the assertion (i) and (5.24), sending $n \rightarrow \infty$ in (5.23) yields (5.22). Therefore, there exists a positive integer $n^* > N_1$ such that for each $n \geq n^*$,

$$\frac{1}{4R^3(\sigma_s(R) - \tilde{\sigma})} \leq \frac{\frac{\mu_n}{n^3} - \frac{1}{2R^3(\sigma_s(R) - \tilde{\sigma})}}{\frac{1}{n}} \leq \frac{3}{4R^3(\sigma_s(R) - \tilde{\sigma})},$$

i.e.,

$$\frac{2n^3 + n^2}{4R^3(\sigma_s(R) - \tilde{\sigma})} \leq \mu_n \leq \frac{2n^3 + 3n^2}{4R^3(\sigma_s(R) - \tilde{\sigma})}.$$

Subsequently, there holds

$$\mu_{n+1} - \mu_n \geq \frac{4n^2 + 8n + 3}{4R^3(\sigma_s(R) - \tilde{\sigma})} > 0$$

for $n \geq n^*$, which completes the proof of the lemma. \square

Denote

$$n^{**} = \min \left\{ n : n \geq n^*, \mu_n > \max \left\{ \frac{A_n}{B_n} : B_n \neq 0, n = 0, 1, \dots, n^* - 1 \right\} \right\}.$$

Recall that we have computed

$$[F_{\tilde{R}}(0, \mu)]Y_{n,0} = (A_n - \mu B_n)Y_{n,0}, \quad n = 0, 1, 2, \dots.$$

Thus, from the definition of n^{**} , we derive that for even $n \geq n^{**}$,

$$\text{Ker}[F_{\tilde{R}}(0, \mu_n)] = \text{span}\{Y_{n,0}\},$$

i.e.,

$$\dim(\text{Ker}[F_{\tilde{R}}(0, \mu_n)]) = 1.$$

Next, since

$$[F_{\tilde{R}}(0, \mu_n)]Y_{k,0} = (A_k - \mu_n B_k)Y_{k,0}, \quad k = 0, 2, 4, \dots,$$

and $A_k - \mu_n B_k \neq 0$ for $k = 0, 2, \dots, n-2, n+2, \dots$,

$$Y_1 = \text{Im}[F_{\tilde{R}}(0, \mu_n)] = \text{span}\{Y_{0,0}, Y_{2,0}, \dots, Y_{n-2,0}, Y_{n+2,0}, \dots\},$$

namely,

$$\text{codim}Y_1 = 1.$$

Finally, it is easy to see that

$$[F_{\mu\tilde{R}}(0, \mu_n)]Y_{n,0} = -B_n Y_{n,0} \notin Y_1.$$

To sum up, we have the following result by the Crandall-Rabinowitz theorem.

Theorem 5.1. *There exists a positive integer n^{**} such that for every even integer $n \geq n^{**}$, μ_n defined by (5.21) is a bifurcation point of the symmetry-breaking solutions to the system (1.9)-(1.16) with free boundary*

$$\partial\Omega_\varepsilon : r = R + \varepsilon Y_{n,0}(\theta) + O(\varepsilon^2), \quad \partial D_\varepsilon : r = \rho + \varepsilon \frac{\lambda}{\underline{\sigma}} Q'_n(\rho) Y_{n,0}(\theta) + O(\varepsilon^2),$$

where λ, Q_n are respectively given by (4.12), (4.25).

Acknowledgments

This work was done in part while the first author was visiting Department of Applied and Computational Mathematics and Statistics, University of Notre Dame, under the support of China Scholarship Council (201708360059). The kind hospitality and the financial support are gratefully acknowledged. This work is also supported by the National Natural Science Foundation of China (No. 11601200, No. 11861038 and No. 11771156), and the Science and Technology Planning Project from Educational Commission of Jiangxi Province, China (No. GJJ160299).

References

- [1] S. Agmon, A. Douglis and L. Nirenberg, Estimates near the boundary for solutions of elliptic partial differential equations satisfying general boundary conditions II, *Comm. Pure Appl. Math.*, **17**(1964), 35–92.
- [2] R. P. Araujo, D. L. S. McElwain, A history of the study of solid tumour growth: the contribution of mathematical modeling, *Bull. Math. Biol.*, **66**(2004), 1039–1091.
- [3] M. Bodnar, U. Forys, Time delay in necrotic core formation, *Math. Biosci. Eng.*, **2**(2005), 461–472.
- [4] H. Bueno, G. Ercole, A. Zumpano, Stationary solutions of a model for the growth of tumors and a connection between the nonnecrotic and necrotic phases, *SIAM J. Appl. Math.*, **68**(2008), 1004–1025.
- [5] H. M. Byrne, M. A. J. Chaplain, Growth of nonnecrotic tumors in the presence and absence of inhibitors, *Math. Biosci.*, **130**(1995), 151–181.
- [6] H. M. Byrne, M. A. J. Chaplain, Growth of necrotic tumors in the presence and absence of inhibitors, *Math. Biosci.*, **135**(1996), 187–216.
- [7] M. G. Crandall, P. H. Rabinowitz, Bifurcation from simple eigenvalues, *J. Functional Analysis*, **8**(1971), 321–340.
- [8] V. Cristini, J. Lowengrub, Q. Nie, Nonlinear simulation of tumor growth, *J. Math. Biol.*, **46**(2003), 191–224.
- [9] S. Cui and A. Friedman, Analysis of a mathematical model of the growth of necrotic tumors, *J. Math. Anal. Appl.*, **255**(2001), 636–677.
- [10] S. Cui, Global existence of solutions for a free boundary problem modeling the growth of necrotic tumors, *Interfaces Free Bound.*, **7**(2005), 147–159.
- [11] S. Cui, Formation of necrotic cores in the growth of tumors: analytic results, *Acta Math. Sci.*, **26B**(2006), 781–796.

- [12] S. Cui, J. Escher, Bifurcation analysis of an elliptic free boundary problem modelling the growth of avascular tumors, *SIAM J. Math. Anal.*, **39** (2007), 210–235.
- [13] S. Cui, J. Escher, Asymptotic behaviour of solutions of a multidimensional moving boundary problem modeling tumor growth, *Comm. Partial Differential Equations*, **33**(2008), 636–655.
- [14] S. Cui, Analysis of a mathematical model for the growth of tumors under the action of external inhibitors, *J. Math. Biol.*, **44**(2002), 395–426.
- [15] S. Cui, Analysis of a free boundary problem modeling the growth of necrotic tumors, arXiv:1902.04066.
- [16] S. Cui, A. Friedman, Analysis of a mathematical model of the effect of inhibitors on the growth of tumors, *Math. Biosci.*, **164**(2000), 103–137.
- [17] S. Cui, Y. Zhuang, Bifurcation solutions of a free boundary problem modeling tumor growth with angiogenesis, *J. Math. Anal. Appl.*, **468**(2018), 391–405.
- [18] J. Escher, A. V. Matioc, Bifurcation analysis for a free boundary problem modeling tumor growth, *Arch. Math.*, **97**(2011), 79–90.
- [19] M. A. Fontelos, A. Friedman, Symmetry-breaking bifurcations of free boundary problems in three dimensions, *Asymptot. Anal.*, **35**(2003), 187–206.
- [20] U. Foryś, A. Mokwa-Borkowska, Solid tumour growth analysis of necrotic core formation, *Math. Comput. Modelling*, **42**(2005), 593–600.
- [21] Frank W. J. Olver, Daniel W. Lozier, Ronald F. Boisvert and Charles W. Clark, *NIST Handbook of Mathematical Functions*, Cambridge University Press, Cambridge, 2010.
- [22] A. Friedman, F. Reitich, Analysis of a mathematical model for the growth of tumors, *J. Math. Biol.*, **38**(1999), 262–284.
- [23] A. Friedman, F. Reitich, Symmetry-breaking bifurcation of analytic solutions to free boundary problems: an application to a model of tumor growth, *Trans. Amer. Math. Soc.*, **353**(2001), 1587–1634.
- [24] A. Friedman, F. Reitich, Nonlinear stability of a quasi-static Stefan problem with surface tension: a continuation approach, *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, **30**(2001), 341–403.
- [25] A. Friedman, A hierarchy of cancer models and their mathematical challenges, *Discrete Contin. Dyn. Syst. Ser. B*, **4**(2004), 147–159.
- [26] A. Friedman, B. Hu, Bifurcation from stability to instability for a free boundary problem arising in a tumor model, *Arch. Ration. Mech. Anal.*, **180**(2006), 293–330.
- [27] A. Friedman, B. Hu, Asymptotic stability for a free boundary problem arising in a tumor model, *J. Differential Equations*, **227**(2006), 598–639.

- [28] A. Friedman, Mathematical analysis and challenges arising from models of tumor growth, *Math. Models Methods Appl. Sci.*, **17**(2007), 1751–1772.
- [29] A. Friedman, B. Hu, Bifurcation for a free boundary problem modeling tumor growth by Stokes equation, *SIAM J. Math. Anal.*, **39**(2007), 174–194.
- [30] A. Friedman, B. Hu, Bifurcation from stability to instability for a free boundary problem modeling tumor growth by Stokes equation, *J. Math. Anal. Appl.*, **327**(2007), 643–664.
- [31] A. Friedman, B. Hu, Stability and instability of Liapunov-Schmidt and Hopf bifurcation for a free boundary problem arising in a tumor model, *Trans. Amer. Math. Soc.*, **360**(2008), 5291–5342.
- [32] A. Friedman, K-Y. Lam, Analysis of a free-boundary tumor model with angiogenesis, *J. Differential Equations*, **259**(2015), 7636–7661.
- [33] H. P. Greenspan, Models for the growth of a solid tumor by diffusion, *Studies Appl. Math.*, **51**(1972), 317–340.
- [34] H. P. Greenspan, On the growth and stability of cell cultures and solid tumors, *J. Theor. Biol.*, **56**(1976), 229–242.
- [35] W. Hao, J. D. Hauenstein, B. Hu, Y. Liu, A. J. Somnese, Y. Zhang, Bifurcation for a free boundary problem modeling the growth of a tumor with a necrotic core, *Nonlinear Anal. Real World Appl.*, **13**(2012), 694–709.
- [36] Y. Huang, Z. Zhang, B. Hu, Bifurcation for a free-boundary tumor model with angiogenesis, *Nonlinear Anal. Real World Appl.*, **35**(2017), 483–502.
- [37] Y. Huang, Z. Zhang, B. Hu, Bifurcation from stability to instability for a free boundary tumor model with angiogenesis, *Discrete Contin. Dyn. Syst.*, **39**(2019), 2473–2510.
- [38] O. A. Ladyzhenskaya and N. N. Ural'tseva, *Linear and Quasilinear Elliptic Equations*, Academic Press, New York, 1968.
- [39] J. S. Lowengrub, H. B. Frieboes, F. Jin, Y-L. Chuang, X. Li, P. Macklin, S. M. Wise, V. Cristini, Nonlinear modelling of cancer: bridging the gap between cells and tumours, *Nonlinearity*, **23**(2010), 1–91.
- [40] J. Nagy, The ecology and evolutionary biology of cancer: a review of mathematical models of necrosis and tumor cell diversity, *Math. Biosci. Eng.*, **2**(2005), 381–418.
- [41] H. Shen, X. Wei, A qualitative analysis of a free boundary problem modeling tumor growth with angiogenesis, *Nonlinear Anal. Real World Appl.*, **47**(2019), 106–126.
- [42] H. Shen, X. Wei, et al., Existence and uniqueness of the stationary solution of mathematical model of necrotic tumor with the third boundary, *Acta Sci. Natur. Univ. Sunyatseni*, **57**(2018), 140–144.

- [43] Z. Wang, Bifurcation for a free boundary problem modeling tumor growth with inhibitors, *Nonlinear Anal. Real World Appl.*, **19**(2014), 45–53.
- [44] Z. Wang, S. Xu, H. Song, Stationary solutions of a free boundary problem modeling growth of angiogenesis tumor with inhibitor, *Discrete Contin. Dyn. Syst. Ser. B*, **23**(2018), 2593–2605.
- [45] X. Wei, Global existence for a free boundary problem modelling the growth of necrotic tumors in the presence of inhibitors, *Int. J. Pure Appl. Math.*, **28**(2006), 321–338.
- [46] J. Wu, Stationary solutions of a free boundary problem modeling the growth of tumors with Gibbs-Thomson relation, *J. Differential Equations*, **260**(2016), 5875–5893.
- [47] J. Wu, Bifurcation for a free boundary problem modeling the growth of necrotic multilayered tumors, *Discrete Contin. Dyn. Syst.*, **39**(2019), 3399–3411.
- [48] J. Wu, Asymptotic behavior of a free boundary problem for the growth of multi-layer tumors in necrotic phase, arXiv: 1802.03112.
- [49] J. Wu, S. Cui, Asymptotic behaviour of solutions of a free boundary problem modelling the growth of tumours in the presence of inhibitors, *Nonlinearity*, **20**(2007), 2389–2408.
- [50] J. Wu, S. Cui, Bifurcation analysis of a mathematical model for the growth of solid tumors in the presence of external inhibitors, *Math. Methods Appl. Sci.*, **38**(2015), 1813–1823.
- [51] S. Xu, M. Bai, Stability of solutions to a mathematical model for necrotic tumor growth with time delays in proliferation, *J. Math. Anal. Appl.*, **421**(2015), 955–962.
- [52] F. Zhou, S. Cui, Bifurcation for a free boundary problem modeling the growth of multi-layer tumors, *Nonlinear Anal.*, **68**(2008), 2128–2145.
- [53] F. Zhou, J. Escher, S. Cui, Bifurcation for a free boundary problem with surface tension modeling the growth of multi-layer tumors, *J. Math. Anal. Appl.*, **337**(2008), 443–457.
- [54] F. Zhou, J. Wu, Stability and bifurcation analysis of a free boundary problem modelling multi-layer tumours with Gibbs-Thomson relation, *European J. Appl. Math.*, **26**(2015), 401–425.
- [55] Y. Zhuang, S. Cui, Analysis of a free boundary problem modeling the growth of multicell spheroids with angiogenesis, *J. Differential Equations*, **265**(2018), 620–644.
- [56] Y. Zhuang, S. Cui, Analysis of a free boundary problem modeling the growth of spherically symmetric tumors with angiogenesis, *Acta Appl. Math.*, (2018), 1–17.