

Blocks with abelian defect groups of rank 2 and one simple module

Xueqin Hu

School of Mathematics and Statistics, Central China Normal University, Wuhan 430079, China

Abstract

In this paper, we investigate the block that has an abelian defect group of rank 2 and its Brauer correspondent has only one simple module. We will get an isotypy between the block and its Brauer correspondent. It will generalize the result of Kessar and Linckelmann ([5]).

1 Introduction

Let p be a prime and \mathcal{O} a complete discrete valuation ring having an algebraically closed residue field k of characteristic p and a quotient field \mathcal{K} of characteristic 0. We will always assume that \mathcal{K} is big enough for the finite groups below.

Let G be a finite group and b a block of $\mathcal{O}G$ with a defect group P . Denote by $\text{Irr}_{\mathcal{K}}(G, b)$ and $\text{IBr}(G, b)$ the set of irreducible ordinary characters in b and the set of irreducible Brauer characters in b respectively. Set $l_G(b) = |\text{IBr}(G, b)|$. Let c be the Brauer correspondent of b in $N_G(P)$. In [5], Kessar and Linckelmann investigated the block b under the assumptions that $l_{N_G(P)}(c) = 1$ and P is elementary abelian of rank 2. They showed that the inertial quotient of b is abelian and there is an isotypy between b and c all of whose signs are positive.

In this note, we will generalize these results to the blocks with defect groups of rank 2.

Theorem 1.1. *Keep the notation as above. Assume that P is abelian of rank 2 and $l_{N_G(P)}(c) = 1$. Then the inertial quotient of b is abelian and there is an isotypy between b and c .*

These results are well-known when either p is 2 or the inertial quotient of b is trivial. Therefore, we may assume that p is odd and the inertial quotient of b is non-trivial throughout this paper.

2 The structure of the block c

Keep the notation as above. In this section, we will investigate the structure of the inertial quotient of b and irreducible ordinary characters of the block c .

Given a positive integer a , denote by C_a the cyclic group of order a . We will use $[-, -]$ to represent the commutator. Assume that $P = C_{p^n} \times C_{p^m}$ for some positive integers n, m . We will fix a maximal b -Brauer pair (P, b_P) . For any $Q \leq P$, denote by (Q, b_Q) the unique b -Brauer pair contained in (P, b_P) . Let E be the inertial quotient of b associated with (P, b_P) , namely, $E = N_G(P, b_P)/C_G(P)$.

Lemma 2.1. *The inertial quotient E is abelian if $l_{N_G(P)}(c) = 1$.*

Proof. Let $\Phi(P)$ be the Frattini subgroup of P . So $P/\Phi(P)$ is $C_p \times C_p$. Set H to be $N_G(P, b_P)$. Then $\Phi(P) \trianglelefteq H$ and denote $H/\Phi(P)$ by \bar{H} . For any subset X of $\mathcal{O}H$, \bar{X} denotes the image of X under the canonical map $\mathcal{O}H \rightarrow \mathcal{O}\bar{H}$.

Since $l_H(b_P) = 1$, $l_{\bar{H}}(\bar{b}_P) = 1$ and \bar{b}_P is a block of \bar{H} with defect group $\bar{P} = C_p \times C_p$. Let \hat{C} be the subgroup of H such that $\hat{C}/\Phi(P) = C_{\bar{H}}(\bar{P})$. Hence, $\hat{C} = \{x \in H \mid [P, x] \subseteq \Phi(P)\}$. It is clear that $P = [P, \hat{C}] \times C_P(\hat{C})$. So $P = C_P(\hat{C})$ since $[P, \hat{C}] \leq \Phi(P)$. This means $C_{\bar{H}}(\bar{P}) = \bar{C}_G(P)$. Hence, (\bar{P}, \bar{b}_P) is a maximal \bar{b}_P -Brauer pair of $\mathcal{O}\bar{H}\bar{b}_P$. By [5, Proposition 5.2], $N_{\bar{H}}(\bar{P}, \bar{b}_P)/C_{\bar{H}}(\bar{P})$ is abelian. It is evident that E is isomorphic to $N_{\bar{H}}(\bar{P}, \bar{b}_P)/C_{\bar{H}}(\bar{P})$. We are done. \square

By [4, Lemma 2] and the structure of blocks with normal defect groups, E is a direct product of two isomorphic groups. Next, we will show that E acts diagonally on P . This can be deduced from the following general fact.

Lemma 2.2. *Let D be an abelian p -group of rank 2 and $F \leq \text{Aut}(P)$ an abelian p' -group which is a direct product of two isomorphic subgroups. Then we have the decompositions $F = F_1 \times F_2$ and $D = D_1 \times D_2$ such that F_1 acts faithfully on D_1 and centralises D_2 and F_2 acts faithfully on D_2 and centralises D_1 and $F_1 \cong F_2$. In particular, F_1 and F_2 are cyclic groups of order dividing $(p-1)$.*

Proof. We will exhibit it by induction on $|D|$. When D is elementary abelian, it is actually done in [5, Proposition 5.3]. We may assume that $n \geq 2$ or $m \geq 2$. Let $\Phi(D)$ be the Frattini subgroup of D . So $D/\Phi(D)$ is $C_p \times C_p$. Let π be the canonical map from F to $\text{Aut}(D/\Phi(D))$. For any subset X of F , \bar{X} denotes the image of X under π . It is clear that π is injective. So there exist two subgroups F_1 and F_2 of F and two subgroups D_1 and D_2 of D containing $\Phi(D)$ satisfying the properties $\bar{F} = \bar{F}_1 \times \bar{F}_2$ and $D/\Phi(D) = D_1/\Phi(D) \times D_2/\Phi(D)$ and \bar{F}_1 acts faithfully on $D_1/\Phi(D)$ and centralises $D_2/\Phi(D)$ and \bar{F}_2 acts faithfully on $D_2/\Phi(D)$ and centralises $D_1/\Phi(D)$ and $\bar{F}_1 \cong \bar{F}_2$. Hence, D_1 and D_2 are F -stable and they fulfill

- (i) $D_1 = [D_1, F_1] \cdot \Phi(D)$ and $[D_1, F_2] \subseteq \Phi(D)$ and F_1 acts faithfully on D_1 ;
- (ii) $D_2 = [D_2, F_2] \cdot \Phi(D)$ and $[D_2, F_1] \subseteq \Phi(D)$ and F_2 acts faithfully on D_2 ;
- (iii) $D_1 \cap D_2 = \Phi(D)$ and $D_1/\Phi(D) \cong C_p \cong D_2/\Phi(D)$ and $D = D_1 \cdot D_2$.

Suppose that $\Phi(D)$ is cyclic. Then $D = C_p \times C_{p^m}$ with $m \geq 2$ and $\Phi(D) = C_{p^{m-1}}$. Since $D_2 = [D_2, F_1] \times C_{D_2}(F_1)$ and $[D_2, F_1] \subseteq \Phi(D)$, $\Phi(D) = [D_2, F_1] \times C_{\Phi(D)}(F_1)$. Then either $[D_2, F_1] = 1$ or $C_{\Phi(D)}(F_1) = 1$ by the assumption that $\Phi(D)$ is cyclic. If $[D_2, F_1] = 1$, then $\Phi(D) \leq D_2 \leq C_D(F_1)$. Clearly, $D = [D, F_1] \times C_D(F_1)$ and D_2 is a maximal subgroup of D . Thus, $D_2 = C_D(F_1)$ and $[D, F_1] = [D_1, F_1]$. Since F_1 and F_2 commute with each other and $[D_1, F_2] \subseteq \Phi(D) \subseteq C_D(F_1)$, $[[D_1, F_1], F_2] = 1$. So $[D_1, F_1] \leq C_P(F_2)$. Since F_1 acts faithfully on D_1 and $D_1 = [D_1, F_1] \times C_{D_1}(F_1)$, F_1 acts faithfully on $[D_1, F_1]$. Thus, the decompositions $F = F_1 \times F_2$ and $D = [D_1, F_1] \times D_2$ are what we want. We may assume that $C_{\Phi(D)}(F_1) = 1$. Then $\Phi(D) = [D_2, F_1]$ and $D_2 = \Phi(D) \times C_{D_2}(F_1)$. If $C_{\Phi(D)}(F_2) = 1$, we can get $\Phi(D) = [D_1, F_2]$ and $D_1 = \Phi(D) \times C_{D_1}(F_2)$ similarly. Then $D = C_{D_1}(F_2) \times C_{D_2}(F_1)$ which is impossible. So $C_{\Phi(D)}(F_2) \neq 1$. Then replacing D_2 by D_1 in the previous argument, we can obtain the decompositions that we need.

Suppose $\Phi(D)$ is of rank 2. Then both D_1 and D_2 are of rank 2. Let K be subgroup of F consisting of automorphisms acting trivially on D_1 . Then $D = [D, K] \times C_D(K)$ and $D_1 \leq C_D(K)$. Hence, K has to be trivial since D_1 has rank 2. This means F acts faithfully on D_1 . By induction, we have $D_1 = D_{11} \times D_{12}$ and $F = F_{11} \times F_{12}$ such that F_{11} acts faithfully on D_{11} and centralises D_{12} and F_{12} acts faithfully on D_{12} and centralises D_{11} and $F_{11} \cong F_{12}$. Then $D = [D, F_{11}] \times C_D(F_{11})$ and $D_{11} = [D_{11}, F_{11}] \leq [D, F_{11}]$ and $D_{11} \leq C_D(F_{12})$. In particular, $C_{[D, F_{11}]}(F_{12}) \neq 1$. But $[D, F_{11}]$ is cyclic. Then $[D, F_{11}] \leq C_D(F_{12})$ and moreover

$C_D(F_{12}) = [D, F_{11}] \times (C_D(F_{11}) \cap C_D(F_{12}))$. But $C_D(F_{12})$ is also cyclic. We have $[D, F_{11}] = C_D(F_{12})$. Similarly, we can prove that $[D, F_{12}] = C_D(F_{11})$. Then the decompositions $D = [D, F_{11}] \times [D, F_{12}]$ and $F = F_{11} \times F_{12}$ are what we want. We are done. \square

Hence, by Lemma 2.2, we have $E = E_1 \times E_2$ and $P = P_1 \times P_2$ such that

- (i) E_1 acts faithfully on P_1 and centralises P_2 ;
- (ii) E_2 acts faithfully on P_1 and centralises P_1 ;
- (iii) $E_1 \cong E_2$ are cyclic groups of order l , which l is a positive integer dividing $(p-1)$.

We can easily describe the source algebra of the block c by the structure theory of blocks with normal defect groups and the structure of inertial quotient E . It is well-known that there exists a central extension

$$1 \longrightarrow Z \longrightarrow \tilde{E} \longrightarrow E \longrightarrow 1$$

with Z cyclic p' -group such that there is an irreducible ordinary character θ of Z which is covered by a unique irreducible character of \tilde{E} . Let $e_\theta \in \mathcal{O}Z$ be the central idempotent corresponding to θ . Set $N = P \rtimes \tilde{E}$. Then $\mathcal{O}Ne_\theta$ is the source algebra of the block c . Note that e_θ is still a block of $C_N(R)$ for any $R \leq P$. The following lemma gives some information about the degrees and number of irreducible ordinary characters of $\mathcal{O}Ne_\theta$, which is similar with [5, Proposition 5.3]. We will skip the proof.

Lemma 2.3. *Set A to be $\mathcal{O}Ne_\theta$. Then the degree of an element of $\text{Irr}_\mathcal{K}(A)$ is either l or l^2 and $\text{Irr}_\mathcal{K}(A)$ has $p^n + p^m - 1$ elements of degree l and $\frac{p^n-1}{l} \cdot \frac{p^m-1}{l}$ elements of degree l^2 .*

3 The extension of local system

Keep the notation as above. In this section, we will use the so-called (G, b) -local system introduced by Puig and Usami in [9] to prove the main theorem.

First, let us recall some notation and state the definition of (G, b) -local system under our setting (see [9]).

Let $\mathcal{CF}_\mathcal{K}(G)$ be the vector space of \mathcal{K} -valued class functions of G and $\mathcal{BCF}_\mathcal{K}(G)$ be the vector space of \mathcal{K} -valued class functions on the set $G_{p'}$ of p' -elements of G . It is clear that the set of irreducible ordinary characters of G is a \mathcal{K} -basis of $\mathcal{CF}_\mathcal{K}(G)$ and the set of irreducible Brauer characters of G is a \mathcal{K} -basis of $\mathcal{BCF}_\mathcal{K}(G)$. For $\chi, \chi' \in \mathcal{CF}_\mathcal{K}(G)$, we denote by $\langle \chi, \chi' \rangle$ the inner product of χ and χ' .

Let u be a p -element of G . we have the well-known surjective \mathcal{K} -linear map $d_G^u : \mathcal{CF}_\mathcal{K}(G) \longrightarrow \mathcal{BCF}_\mathcal{K}(C_G(u))$ defined by $d_G^u(\chi)(s) = \chi(us)$ for any $\chi \in \mathcal{CF}_\mathcal{K}(G)$ and $s \in C_G(u)_{p'}$. It has a section $e_G^u : \mathcal{BCF}_\mathcal{K}(C_G(u)) \longrightarrow \mathcal{CF}_\mathcal{K}(G)$ such that for $\varphi \in \mathcal{BCF}_\mathcal{K}(C_G(u))$, $e_G^u(\varphi)(g) = 0$ if the p -part of g is not conjugate to u in G .

For the block b , let $\mathcal{CF}_\mathcal{K}(G, b)$ be the subspace of $\mathcal{CF}_\mathcal{K}(G)$ generated by the elements in $\text{Irr}_\mathcal{K}(G, b)$ and $\mathcal{L}_\mathcal{K}(G, b)$ the group of generalized characters in b . Also, let $\mathcal{CF}_\mathcal{K}^\circ(G, b) = \mathcal{CF}_\mathcal{K}(G, b) \cap \text{Ker}(d_G^1)$ and $\mathcal{L}_\mathcal{K}^\circ(G, b) = \mathcal{L}_\mathcal{K}(G, b) \cap \text{Ker}(d_G^1)$.

Definition 3.1. (Puig-Usami [9, 3.2]) *With the above notation and assumption. Let X be an E -stable non-empty set of subgroups of P and assume that X contains any subgroup of P containing an element of X . Let Γ be a map over X sending $Q \in X$ to a bijective isometry*

$$\Gamma_Q : \mathcal{BCF}_\mathcal{K}(C_N(Q), e_\theta) \longrightarrow \mathcal{BCF}_\mathcal{K}(C_G(Q), b_Q).$$

If Γ satisfies the following conditions, then Γ is called a (G, b) -local system over X .

(i) For any $Q \in X$, any $\eta \in \mathcal{BCF}_K(C_N(Q), e_\theta)$ and any $s \in E$, we have $\Gamma_Q(\eta)^s = \Gamma_{Q^s}(\eta^s)$.

(ii) For any $Q \in X$ and any $\eta \in \mathcal{L}_K(C_N(Q), e_\theta)$, the sum

$$\sum_u e_{C_G(Q)}^u(\Gamma_{Q \cdot \langle u \rangle}(d_{C_N(Q)}^u(\eta)))$$

where u runs over a set of representatives U_Q for the orbits of $C_E(Q)$ in P , is a generalized character of $C_G(Q)$.

Let Γ be a (G, b) -local system over X . Such Γ always exists by [9, 3.4.2]. For any $Q \in X$, we have a map $\Delta_Q : \mathcal{CF}_K(C_N(Q), e_\theta) \rightarrow \mathcal{CF}_K(C_G(Q), b_Q)$ defined by

$$\Delta_Q(\eta) = \sum_{u \in U_Q} e_{C_G(Q)}^u(\Gamma_{Q \cdot \langle u \rangle}(d_{C_N(Q)}^u(\eta))).$$

Then by [9, 3.3 and 3.4] Δ_Q gives a perfect isometry between the block e_θ of $C_N(Q)$ and the block b_Q of $C_G(Q)$ and $\Delta_Q(\lambda * \eta) = \lambda * \Delta_Q(\eta)$ for any $\lambda \in \mathcal{CF}_K(P)^{C_E(Q)}$ and $\eta \in \mathcal{CF}_K(C_N(Q))$. Here, $\mathcal{CF}_K(P)^{C_E(Q)}$ denotes the set of $C_E(Q)$ -stable elements of $\mathcal{CF}_K(P)$ and $*$ denotes the $*$ -construction of characters due to Broué and Puig (see [2]). Hence, if X contains the trivial subgroup 1 of P , then Δ_1 induces a perfect isometry between the block e_θ of N and the block b of G . Moreover, this is an isotypy in the sense of [1] by [13, Proposition 2.7].

In [9], Puig and Usami developed a criterion for the extendibility of the (G, b) -local system. With the notation above. Suppose that $1 \notin X$ and let Q be a maximal subgroup of P such that $Q \notin X$. Denote by X' the union of X and the E -orbit of Q . For any subset Y of $\mathcal{OC}_N(Q)$, denote by \bar{Y} the image of Y under the canonical map from $\mathcal{OC}_N(Q)$ to $\mathcal{OC}_N(Q)/Q$. We have the similar notation for $\mathcal{OC}_G(Q)$. So \bar{e}_θ and \bar{b}_Q are the blocks of $\bar{C}_N(Q)$ and $\bar{C}_G(Q)$ respectively. Set $\Delta_Q^\circ = \sum_{u \in U_{Q-Q}} e_{C_G(Q)}^u \circ \Gamma_{Q \cdot \langle u \rangle} \circ d_{C_N(Q)}^u$ (see [9, 3.6.2]). By [9, Proposition 3.7 and Remark 3.8], Δ_Q° induces a bijective isometry

$$\bar{\Delta}_Q^\circ : \mathcal{CF}_K(\bar{C}_N(Q), \bar{e}_\theta) \cong \mathcal{CF}_K(\bar{C}_G(Q), \bar{b}_Q)$$

such that $\bar{\Delta}_Q^\circ(\mathcal{L}_K(\bar{C}_N(Q), \bar{e}_\theta)) = \mathcal{L}_K(\bar{C}_G(Q), \bar{b}_Q)$. Clearly, $\bar{\Delta}_Q^\circ(\lambda * \eta) = \lambda * \bar{\Delta}_Q^\circ(\eta)$ for $\lambda \in \text{Irr}_K(\bar{P})^{C_E(Q)}$ and $\eta \in \mathcal{L}_K(\bar{C}_N(Q), \bar{e}_\theta)$ (see [11, Case 2.2]) and $\bar{\Delta}_Q^\circ$ is $N_E(Q)$ -stable. The following is the key criterion of extendibility.

Proposition 3.2. ([9, Proposition 3.11]) *With the notation above, the (G, b) -local system Γ over X can be extended to a (G, b) -local system Γ' over X' if and only if $\bar{\Delta}_Q^\circ$ can be extended to an $N_E(Q)$ -stable bijective isometry*

$$\bar{\Delta}_Q : \mathcal{CF}_K(\bar{C}_N(Q), \bar{e}_\theta) \cong \mathcal{CF}_K(\bar{C}_G(Q), \bar{b}_Q)$$

such that $\bar{\Delta}_Q(\mathcal{L}_K(\bar{C}_N(Q), \bar{e}_\theta)) = \mathcal{L}_K(\bar{C}_G(Q), \bar{b}_Q)$.

In order to prove Theorem 1.1, it suffices to show that there is a (G, b) -local system over the set of all the subgroups P . Hence, by Proposition 3.2, we can assume that there is a (G, b) -local system Γ over X such that $1 \notin X$ and Q is a maximal subgroup of P such that $Q \notin X$.

Theorem 3.3. *With the notation above and assumptions of Section 2. Then $\bar{\Delta}_Q^\circ$ can be extended to an $N_E(Q)$ -stable bijective isometry*

$$\bar{\Delta}_Q : \mathcal{CF}_K(\bar{C}_N(Q), \bar{e}_\theta) \cong \mathcal{CF}_K(\bar{C}_G(Q), \bar{b}_Q)$$

such that $\bar{\Delta}_Q(\mathcal{L}_K(\bar{C}_N(Q), \bar{e}_\theta)) = \mathcal{L}_K(\bar{C}_G(Q), \bar{b}_Q)$.

Proof. By the structure of E and P , $C_E(Q)$ has only three possibilities: 1, E and E_1 or E_2 . So we will divided the proof into 3 cases.

Case 1 Assume that $C_E(Q) = 1$.

Then the blocks e_θ of $C_N(Q)$ and b_Q of $C_G(Q)$ are nilpotent. By the same argument as in [9, 4.4], $\bar{\Delta}_Q^\circ$ can be extended to an $N_E(Q)$ -stable bijective isometry $\bar{\Delta}_Q$.

Case 2 Assume that $C_E(Q) = E$.

Then Q has to be trivial subgroup of P and $N_E(Q) = E$. So $\bar{C}_N(Q) = N$ and $\bar{C}_G(Q) = G$ and we have a bijective isometry

$$\bar{\Delta}^\circ : \mathcal{CF}_K^\circ(N, e_\theta) \longrightarrow \mathcal{CF}_K^\circ(G, b)$$

such that $\bar{\Delta}^\circ(\mathcal{L}_K^\circ(N, e_\theta)) = \mathcal{L}_K^\circ(G, b)$.

The following technique we adopt to extend $\bar{\Delta}^\circ$ is essentially due to Kessar and Linckelmann (see [5, Theorem 4.1]).

By Lemma 2.3, we have the following disjoint union

$$\text{Irr}_K(N, e_\theta) = \Lambda_1 \cup \Lambda_2,$$

where Λ_1 consists of irreducible ordinary characters of dimension l and Λ_2 consists of irreducible ordinary characters of dimension l^2 . Hence, $|\Lambda_1| = p^n + p^m - 1$ and $|\Lambda_2| = \frac{p^n-1}{l} \cdot \frac{p^m-1}{l}$. We can assume that $n \geq 2$. Then $|\Lambda_1| > 2$ and $|\Lambda_2| > 2$. Choose an element $\psi_i \in \Lambda_i$ and set $\Lambda'_i = \Lambda_i - \{\psi_i\}$ for $i = 1, 2$. Since $l_N(e_\theta) = 1$, it is easy to see

$$\mathcal{B} = \{\psi_1 - \psi'_1 \mid \psi'_1 \in \Lambda'_1\} \cup \{\psi_2 - \psi'_2 \mid \psi'_2 \in \Lambda'_2\} \cup \{\psi_2 - l\psi_1\}$$

is a \mathbb{Z} -basis of $\mathcal{L}_K^\circ(N, e_\theta)$. Since p is odd, $|\Lambda'_i| \geq 3$ for $i = 1, 2$. So by the same argument in [9, 4.4], for any $i = 1, 2$, there exists a subset $\Omega_i = \{\chi_{\psi_i}, \chi_{\psi'_i} \mid \psi'_i \in \Lambda'_i\}$ of $\text{Irr}_K(G, b)$ and $\delta_i \in \{\pm 1\}$ such that $\bar{\Delta}^\circ(\psi_i - \psi'_i) = \delta_i(\chi_{\psi_i} - \chi_{\psi'_i})$. Since $\langle \psi_1 - \psi'_1, \psi_2 - \psi'_2 \rangle = 0$ for any $\psi'_1 \in \Lambda'_1$ and $\psi'_2 \in \Lambda'_2$, $\{\chi_{\psi_1}, \chi_{\psi'_1} \mid \psi'_1 \in \Lambda'_1\}$ and $\{\chi_{\psi_2}, \chi_{\psi'_2} \mid \psi'_2 \in \Lambda'_2\}$ have trivial intersection. Denote $\psi_2 - l\psi_1$ by μ . Then $\langle \mu, \psi_1 - \psi'_1 \rangle = -l$ for all $\psi'_1 \in \Lambda'_1$. Thus

$$\bar{\Delta}^\circ(\mu) = \delta_1(a - l)\chi_{\psi_1} + \delta_1 a \sum_{\psi'_1 \in \Lambda'_1} \chi_{\psi'_1} + \Xi \quad (3.1)$$

for some integer a and some element $\Xi \in \mathcal{L}_K(N, e_\theta)$ not involving any of elements in Ω_1 . Since $\langle \mu, \psi_2 - \psi'_2 \rangle = 1$ and $\bar{\Delta}^\circ(\psi_2 - \psi'_2) = \delta_2(\chi_{\psi_2} - \chi_{\psi'_2})$, Ξ must involve one of the two characters occuring in $\bar{\Delta}^\circ(\psi_2 - \psi'_2)$ for any $\psi'_2 \in \Lambda'_2$. Taking norms on both sides in equation (3.1), we have

$$\begin{aligned} 1 + l^2 &\geq (a - l)^2 + (p^n + p^m - 2)a^2 = (p^n + p^m - 1)a^2 - 2la + l^2 \\ \iff 1 &\geq (p^n + p^m - 1)a^2 - 2la \end{aligned} \quad (3.2)$$

Suppose that $a \leq 0$. Since a is integer and $p^n + p^m - 1, l$ are positive integers, a has to be 0.

Suppose that $a > 0$. Since $p^n + p^m - 1 > 2l$, $(p^n + p^m - 1)a^2 - 2la > (p^n + p^m - 1)(a^2 - a)$. This forces $a = 1$. Hence, $a = 0$ or 1. Notice that $\Xi \neq 0$. This implies (3.2) is a proper inequality. So a must be 0. Then equation (3.1) becomes

$$\bar{\Delta}^\circ(\mu) = -\delta_1 l \chi_{\psi_1} + \Xi.$$

Comparing norms, we have $\langle \Xi, \Xi \rangle = 1$.

For any $\psi'_2, \psi''_2 \in \Lambda'_2$,

$$\langle \bar{\Delta}^\circ(\mu), \delta_2(\chi_{\psi_2} - \chi_{\psi'_2}) \rangle = \langle \mu, \psi_2 - \psi'_2 \rangle = 1$$

and

$$\langle \bar{\Delta}^\circ(\mu), \delta_2(\chi_{\psi'_2} - \chi_{\psi''_2}) \rangle = \langle \mu, \psi'_2 - \psi''_2 \rangle = 0.$$

Then $\Xi = \delta_2 \chi_{\psi_2}$. But $\bar{\Delta}^\circ(\mu)(1) = 0$. This forces $\delta_1 = \delta_2$. Since \mathcal{B} is a \mathbb{Z} -basis of $\mathcal{L}_K(N, e_\theta)$, $\text{Irr}_K(G, b) = \Omega_1 \cup \Omega_2$. Hence, we get a bijective isometry $\bar{\Delta}$ from $\mathcal{L}_K(N, e_\theta)$ to $\mathcal{L}_K(G, b)$ mapping ψ_i and ψ'_i to χ_{ψ_i} and $\chi_{\psi'_i}$ respectively, where $i = 1, 2$. In particular, $l_G(b) = l_N(e_\theta) = 1$. Clearly, it is an extension of $\bar{\Delta}^\circ$. Since $\bar{\Delta}^\circ$ is E -stable and $l_G(b) = l_N(e_\theta) = 1$, $\bar{\Delta}$ is also E -stable.

Case 3 Assume that $C_E(Q) = E_i$ for some $i = 1, 2$.

We can assume that $C_E(Q) = E_1$ and then $1 \neq Q \leq P_2$ and $N_E(Q) = E$. It suffices to prove that $\bar{\Delta}_Q^\circ$ can extend to an E_2 -stable bijective isometry $\bar{\Delta}_Q : \mathcal{L}_K(\bar{C}_N(Q), \bar{e}_\theta) \rightarrow \mathcal{L}_K(\bar{C}_G(Q), \bar{b}_Q)$.

By [12, Theorem 1], $|\text{Irr}_K(\bar{C}_N(Q), \bar{e}_\theta)| = |\text{Irr}_K(\bar{C}_G(Q), \bar{b}_Q)|$ and $l_{\bar{C}_N(Q)}(\bar{e}_\theta) = l_{\bar{C}_G(Q)}(\bar{b}_Q)$ since the block \bar{b}_Q of $\bar{C}_G(Q)$ has a cyclic hyperfocal subgroup. It is clear $C_N(Q) = (P_1 \rtimes \tilde{E}_1) \times P_2$ and $C_N(Q) \trianglelefteq N$, where \tilde{E}_1 is the preimage of E_1 in \tilde{E} . Hence, E_1 is the inertial quotient of the block e_θ of $C_N(Q)$ and P_1 is a hyperfocal subgroup with respect to E_1 . By [12, Theorem 1], $l_{C_N(Q)}(e_\theta) = l$. We will claim that N acts transitively on $\text{IBr}(C_N(Q), e_\theta)$. Indeed, this holds because $l_N(e_\theta) = 1$ by the assumption and $N/C_N(Q) \cong E_2$ is a cyclic group of order l .

Denote by $\text{Irr}_K(\tilde{E}_1)_\theta$ the subset of $\text{Irr}_K(\tilde{E}_1)$ consisting of characters covering θ . Then $|\text{Irr}_K(\tilde{E}_1)_\theta| = l$ and we set $\text{Irr}_K(\tilde{E}_1)_\theta = \{\tau_i \mid i = 1, 2, \dots, l\}$, which is transitively acted by N . Hence, we can write $\text{Irr}_K(\tilde{E}_1)_\theta$ as $\{\tau^a \mid a \in E_2\}$ for any $\tau \in \text{Irr}_K(\tilde{E}_1)_\theta$. By Clifford theorem, we have $\text{Res}_Z^{\tilde{E}_1}(\tau_i) = \theta$ for any i and $\text{Ind}_Z^{\tilde{E}_1}(\theta) = \sum_{i=1}^l \tau_i$. Let M be a representative of \tilde{E}_1 -orbit of $\text{Irr}_K(P_1) - \{1_{P_1}\}$, where 1_{P_1} is the trivial character of P_1 . Then

$$\text{Irr}_K(\bar{C}_N(Q), \bar{e}_\theta) = \{\tau_i \bar{\zeta}_j \mid \bar{\zeta}_j \in \text{Irr}_K(\bar{P}_2), i = 1, 2, \dots, l\} \cup \{\text{Ind}_{P_1 \times Z}^{P_1 \rtimes \tilde{E}_1}(\xi \theta) \bar{\zeta}_j \mid \xi \in M, \bar{\zeta}_j \in \text{Irr}_K(\bar{P}_2)\}.$$

We will write $\text{Ind}_{P_1 \times Z}^{P_1 \rtimes \tilde{E}_1}(\xi \theta)$ and $\bar{\chi} \cdot 1_{\bar{P}_2}$ as $\text{Ind}(\xi)$ and $\bar{\chi}$ respectively for simplicity. Here, $\bar{\chi}$ is an element of $\mathcal{CF}_K(P_1 \rtimes \tilde{E}_1)$. Clearly, $\text{Ind}(\xi)$ is N and E_2 -stable for any $\xi \in M$. Similar to the argument of [11, Case 2],

$$\left\{ \left(\sum_{i=1}^l \tau_i - \text{Ind}(\xi) \right) \bar{\zeta} \mid \xi \in M, \bar{\zeta} \in \text{Irr}_K(\bar{P}_2) \right\} \cup \{ \tau_i - \tau_i \bar{\zeta} \mid i = 1, 2, \dots, l, 1_{\bar{P}_2} \neq \bar{\zeta} \in \text{Irr}_K(\bar{P}_2) \}$$

is a \mathbb{Z} -basis of $\mathcal{L}_K(\bar{C}_N(Q), \bar{e}_\theta)$.

Case 3.1 Assume that $\bar{P}_2 = 1$, i.e., $Q = P_2$.

Set $H = N_G(Q, b_Q)$. Then $H = C_G(Q)N_G(P, b_P)$ and b_Q is still a block of H . Let d be the Brauer correspondent of the block b_Q of H in $N_H(P)$. Then $l_{N_H(P)}(d) = 1$ by the assumption. We claim that $l_H(b_Q) = 1$.

Indeed, considering the canonical map from $\mathcal{O}H$ to $\mathcal{O}(H/Q)$, denote by \bar{X} the image of X under this canonical map for any subset X of $\mathcal{O}H$. Then \bar{b}_Q is still a block of $\bar{C}_G(Q)$ and $\bar{H}/\bar{C}_G(Q)$ is a cyclic group of order l . By [6, Lemma 3.5], $\text{Br}_{\bar{P}}(\bar{b}_Q) = \overline{\text{Br}_P(b_Q)}$. Since $l_{N_H(P)}(d) = 1$, \bar{d} is still a block of $\bar{N}_H(P)$. Therefore, $\text{Br}_{\bar{P}}(\bar{b}_Q) = \bar{d}$ is a block of $\bar{N}_H(P)$. Suppose that the blocks of \bar{H} covering the block \bar{b}_Q of $\bar{C}_G(Q)$ have the same defect group \bar{P} . Then \bar{b}_Q is a block of \bar{H} since $\text{Br}_{\bar{P}}(\bar{b}_Q)$ is a block of $\bar{N}_H(P)$ and $N_{\bar{H}}(\bar{P}) = \bar{N}_H(P)$. Hence, it has a defect group \bar{P} which is cyclic by our assumption. In particular, we have $l_H(b_Q) = l_{\bar{H}}(\bar{b}_Q) = l_{N_H(P)}(d) = 1$ since $N_{\bar{H}}(\bar{P}) = \bar{N}_H(P)$. Consequently, the argument follows from the lemma below.

Lemma 3.4 *Let L be a normal subgroup of K such that K/L is a cyclic p' -group. Let i be a K -stable block of L with defect group D . For any block e of K covering i , e has defect group D .*

Proof. We will prove it by induction on K/L . Let $M \leq K$ such that M contains L and $|M/L|$ is a prime. Then $M \trianglelefteq K$ and K/M is still a cyclic p' -group. Denote by $M[i]$ the subgroup of M consisting of elements acting on $\mathcal{O}Li$ as inner automorphisms. Therefore, $M[i] = M$ or L . Let f be a block of M covered by e . So f covers the block i of L . If $M[i] = M$, then $\mathcal{O}Mf$ and $\mathcal{O}Li$ are source algebra equivalent by [7, Theorem 7]. In particular, the block f has defect group D . If $M[i] = L$, then $f = i$ by [3, Theorem 3.5] and certainly they have the same defect group. In conclusion, D is a defect group of the block f . Let K_f be the stabilizer of f in K . Then blocks of K_f covering f have defect group D by induction. So is e .

Moreover, we claim that there is a regular E_2 -orbit of $\text{Irr}_{\mathcal{K}}(\bar{C}_G(Q), \bar{b}_Q)$, namely, H acts transitively on it.

Indeed, since the block \bar{b}_Q of \bar{H} has a cyclic defect group, it must be nilpotent. By [8, Theorem 3.13], the block \bar{b}_Q of $\bar{C}_G(Q)$ is basic Morita equivalent to its Brauer correspondent. Note that the block \bar{b}_Q of $\bar{C}_G(Q)$ is not nilpotent since $l > 1$. This implies that every irreducible Brauer character of the block \bar{b}_Q of $\bar{C}_G(Q)$ can be uniquely lifted to an irreducible ordinary character by the theory of cyclic blocks.

On the other hand, since $l_{C_G(Q)}(b_Q) = l$ and $l_H(b_Q) = 1$ and $H/C_G(Q) \cong E_2$ has order l , H acts transitively on $\text{IBr}(C_G(Q), b_Q)$. Combining this with the argument above, there exists a regular H -orbit of $\text{Irr}_{\mathcal{K}}(\bar{C}_G(Q), \bar{b}_Q)$. We are done.

Case 3.1.1 Assume that $|M| = 1$.

Then $\text{rank}_{\mathcal{O}}(\mathcal{L}_{\mathcal{K}}^{\circ}(\bar{C}_N(Q), \bar{e}_{\theta})) = 1$ and $\mathcal{L}_{\mathcal{K}}^{\circ}(\bar{C}_N(Q), \bar{e}_{\theta}) = \mathbb{Z}(\text{Ind}(\xi) - \sum_{i=1}^l \tau_i)$. Since there is a regular E_2 -orbit of $\text{Irr}_{\mathcal{K}}(\bar{C}_G(Q), \bar{b}_Q)$, $\text{Irr}_{\mathcal{K}}(\bar{C}_G(Q), \bar{b}_Q) = \{\chi_0\} \cap \{\chi_1, \chi_2, \dots, \chi_l\}$ such that χ_0 is E_2 -stable and E_2 acts regularly on $\{\chi_1, \chi_2, \dots, \chi_l\}$. Then we have

$$\bar{\Delta}_Q^{\circ}(\text{Ind}(\xi) - \sum_{i=1}^l \tau_i) = \delta_0 \chi_0 - \sum_{i=1}^l \delta_i \chi_i$$

for some $\delta_0, \delta_i \in \{\pm 1\}$, $i = 1, 2, \dots, l$. Since $\bar{\Delta}_Q^{\circ}$ is E_2 -stable, we have $\delta_1 = \delta_2 = \dots = \delta_l = \delta_0$. If we write $\{\tau_1, \tau_2, \dots, \tau_l\}$ and $\{\chi_1, \chi_2, \dots, \chi_l\}$ as $\{\tau^a \mid a \in E_2\}$ and $\{\chi^a \mid a \in E_2\}$ respectively, then we can define a bijective isometry as below

$$\bar{\Delta}_Q^{\circ} : \mathcal{L}_{\mathcal{K}}(\bar{C}_N(Q), \bar{e}_{\theta}) \longrightarrow \mathcal{L}_{\mathcal{K}}(\bar{C}_G(Q), \bar{b}_Q)$$

$$\text{Ind}(\xi) \mapsto \delta_0 \chi_0$$

$$\tau^a \mapsto \delta_0 \chi^a.$$

It is evident that it is an extension of $\bar{\Delta}_Q^{\circ}$ and E_2 -stable. We are done for this case.

Case 3.1.2 Assume that $|M| \geq 2$.

Then there are at least two different $\xi_1, \xi_2 \in M$. So $\text{Ind}(\xi_1) - \text{Ind}(\xi_2) \in \mathcal{L}_{\mathcal{K}}^{\circ}(\bar{C}_N(Q), \bar{e}_{\theta})$ and $\langle \text{Ind}(\xi_1) - \text{Ind}(\xi_2), \text{Ind}(\xi_1) - \text{Ind}(\xi_2) \rangle = 2$. Then there exist $\chi_1 \neq \chi_2 \in \text{Irr}_{\mathcal{K}}(\bar{C}_G(Q), \bar{b}_Q)$ such that

$$\bar{\Delta}_Q^{\circ}(\text{Ind}(\xi_1) - \text{Ind}(\xi_2)) = \delta(\chi_1 - \chi_2)$$

for some $\delta \in \{\pm 1\}$. Since $\bar{\Delta}_Q^{\circ}$ is E_2 -stable, we have ${}^a(\delta\chi_1 - \delta\chi_2) = \delta(\chi_1 - \chi_2)$ for any $a \in E_2$. This means that χ_1 and χ_2 are both E_2 -stable.

If there is a $\xi_3 \in M$ different from ξ_1 and ξ_2 , then there is a $\chi_3 \in \text{Irr}_{\mathcal{K}}(\bar{C}_G(Q), \bar{b}_Q)$ different from χ_1 and χ_2 such that

$$\bar{\Delta}_Q^\circ(\text{Ind}(\xi_1) - \text{Ind}(\xi_3)) = \delta\chi_1 - \delta\chi_3 \text{ or } -\delta\chi_2 + \delta\chi_3$$

and χ_3 is E_2 -stable; then we may choose the notation in such a way that

$$\bar{\Delta}_Q^\circ(\text{Ind}(\xi_1) - \text{Ind}(\xi_2)) = \delta(\chi_1 - \chi_2) \text{ and } \bar{\Delta}_Q^\circ(\text{Ind}(\xi_1) - \text{Ind}(\xi_3)) = \delta(\chi_1 - \chi_3)$$

for some E_2 -stable elements χ_1, χ_2, χ_3 of $\text{Irr}_{\mathcal{K}}(\bar{C}_G(Q), \bar{b}_Q)$.

If $|M| \geq 4$, then for any $\xi \in M - \{\xi_1, \xi_2, \xi_3\}$, there is a unique $\chi \in \text{Irr}_{\mathcal{K}}(\bar{C}_G(Q), \bar{b}_Q) - \{\chi_1, \chi_2, \chi_3\}$ such that

$$\bar{\Delta}_Q^\circ(\text{Ind}(\xi_1) - \text{Ind}(\xi)) = \delta(\chi_1 - \chi)$$

and χ is E_2 -stable.

In conclusion, we have an injective isometry

$$\Phi : \mathbb{Z}\{\text{Ind}(\xi) \mid \xi \in M\} \longrightarrow \mathcal{L}_{\mathcal{K}}(\bar{C}_G(Q), \bar{b}_Q)$$

mapping $\text{Ind}(\xi)$ to $\delta\chi_\xi$ such that

$$\Phi(\text{Ind}(\xi) - \text{Ind}(\xi')) = \bar{\Delta}_Q^\circ(\text{Ind}(\xi) - \text{Ind}(\xi'))$$

and χ_ξ is E_2 -stable for any $\xi, \xi' \in M$.

Denote $\text{Irr}_{\mathcal{K}}(\bar{C}_G(Q), \bar{b}_Q) - \{\chi_\xi \mid \xi \in M\}$ by Ω . Then $|\Omega| = l$ and E_2 acts on Ω . Since there is a regular E_2 -orbit of $\text{Irr}_{\mathcal{K}}(\bar{C}_G(Q), \bar{b}_Q)$, E_2 acts regularly on Ω . This means that Ω can be represented as $\{\chi^a \mid a \in E_2\}$ for some $\chi \in \Omega$.

Now we fix an element ξ of M . Suppose that χ does not get involved in $\bar{\Delta}_Q^\circ(\text{Ind}(\xi) - \sum_{i=1}^l \tau_i)$. Then there is $\xi' \in M$ such that $\langle \chi, \bar{\Delta}_Q^\circ(\text{Ind}(\xi') - \sum_{i=1}^l \tau_i) \rangle \neq 0$ since $\{\text{Ind}(\xi) - \sum_{i=1}^l \tau_i \mid \xi \in M\}$ is a \mathbb{Z} -basis of $\mathcal{L}_{\mathcal{K}}^\circ(\bar{C}_N(Q), \bar{e}_\theta)$. Hence, χ has to get involved in $\bar{\Delta}_Q^\circ(\text{Ind}(\xi) - \sum_{i=1}^l \tau_i) - \bar{\Delta}_Q^\circ(\text{Ind}(\xi') - \sum_{i=1}^l \tau_i)$ which is $\delta(\chi_\xi - \chi_{\xi'})$. This is impossible. So χ must get involved in $\bar{\Delta}_Q^\circ(\text{Ind}(\xi) - \sum_{i=1}^l \tau_i)$ for any $\xi \in M$. Since $\bar{\Delta}_Q^\circ$ and $\text{Ind}(\xi) - \sum_{i=1}^l \tau_i$ are E_2 -stable, χ^a has to get involved in $\bar{\Delta}_Q^\circ(\text{Ind}(\xi) - \sum_{i=1}^l \tau_i)$ for any $a \in E_2$ and $\xi \in M$. Since $\langle \text{Ind}(\xi) - \sum_{i=1}^l \tau_i, \text{Ind}(\xi) - \sum_{i=1}^l \tau_i \rangle = 1 + l$ and $\langle \text{Ind}(\xi) - \sum_{i=1}^l \tau_i, \text{Ind}(\xi) - \text{Ind}(\xi') \rangle = 1$, $\bar{\Delta}_Q^\circ(\text{Ind}(\xi) - \sum_{i=1}^l \tau_i) = \delta\chi_\xi - \sum_{a \in E_2} \delta_a \chi^a$ or $-\delta\chi_{\xi'} - \sum_{a \in E_2} \delta_a \chi^a$, where $\delta_a \in \{\pm 1\}$ for any $a \in E_2$. Note that the last situation can happen if and only if $|M| = 2$. By switching χ_ξ and $\chi_{\xi'}$ if necessary, we can assume that $\bar{\Delta}_Q^\circ(\text{Ind}(\xi) - \sum_{i=1}^l \tau_i) = \delta\chi_\xi - \sum_{a \in E_2} \delta_a \chi^a$. Since $\bar{\Delta}_Q^\circ$ is E_2 -stable and E_2 acts regularly on Ω , δ_a is equal to δ for any $a \in E_2$. Then we can define an E_2 -stable bijective isometry as follows

$$\bar{\Delta}_Q : \mathcal{L}_{\mathcal{K}}(\bar{C}_N(Q), \bar{e}_\theta) \longrightarrow \mathcal{L}_{\mathcal{K}}(\bar{C}_G(Q), \bar{b}_Q)$$

$$\text{Ind}(\xi) \mapsto \delta\chi_\xi$$

$$\tau^a \mapsto \delta\chi^a.$$

It is clear that $\bar{\Delta}_Q$ is an extension of $\bar{\Delta}_Q^\circ$.

Case 3.2 $\bar{P}_2 > 1$, namely, Q is a non-trivial proper subgroup of P_2 .

Then $\text{Ind}(\xi) - \text{Ind}(\xi)\bar{\zeta} \in \mathcal{L}_{\mathcal{K}}^{\circ}(\bar{C}_N(Q), \bar{e}_{\theta})$ for any $\xi \in M$ and $1_{\bar{P}_2} \neq \bar{\zeta} \in \text{Irr}_{\mathcal{K}}(\bar{P}_2)$.

Now we fix an element $\xi \in M$. Since p is odd, $|\bar{P}_2| \geq 3$. Then there are at least two elements $\bar{\zeta}$ and $\bar{\zeta}'$ of $\text{Irr}_{\mathcal{K}}(\bar{P}_2)$ different from $1_{\bar{P}_2}$. With the same argument in the first three paragraphs in Case 3.1.2, we can get a subset $\{\chi_{\xi}, \chi_{\bar{\zeta}} \mid 1_{\bar{P}_2} \neq \bar{\zeta} \in \text{Irr}_{\mathcal{K}}(\bar{P}_2)\}$ of $\text{Irr}_{\mathcal{K}}(\bar{C}_G(Q), \bar{b}_Q)$ such that

$$\bar{\Delta}_Q^{\circ}(\text{Ind}(\xi) - \text{Ind}(\xi)\bar{\zeta}) = \delta(\chi_{\xi} - \chi_{\bar{\zeta}})$$

for any $1_{\bar{P}_2} \neq \bar{\zeta} \in \text{Irr}_{\mathcal{K}}(\bar{P}_2)$, where $\delta \in \{\pm 1\}$.

Given any $1 \neq a \in E_2$ and $1_{\bar{P}_2} \neq \bar{\zeta} \in \text{Irr}_{\mathcal{K}}(\bar{P}_2)$, ${}^a(\text{Ind}(\xi)\bar{\zeta}) = \text{Ind}(\xi)({}^a\bar{\zeta})$. Since $\bar{\Delta}_Q^{\circ}$ is E_2 -stable, this means ${}^a\chi_{\xi} - {}^a\chi_{\bar{\zeta}} = \chi_{\xi} - \chi_{a\bar{\zeta}}$. Hence, we have χ_{ξ} is E_2 -stable and ${}^a\chi_{\bar{\zeta}} = \chi_{a\bar{\zeta}}$. On the other hand,

$$(\text{Ind}(\xi) - \text{Ind}(\xi)\bar{\zeta})\bar{\zeta} = (\text{Ind}(\xi) - \text{Ind}(\xi)\bar{\zeta}^2) - (\text{Ind}(\xi) - \text{Ind}(\xi)\bar{\zeta}).$$

Since $\bar{\Delta}_Q^{\circ}$ is compatible with $*$ -structure, using $\bar{\Delta}_Q^{\circ}$ on both sides in the above equality, we can get

$$\delta(\chi_{\xi} - \chi_{\bar{\zeta}}) * \bar{\zeta} = \delta(\chi_{\bar{\zeta}} - \chi_{\bar{\zeta}^2}).$$

Therefore, $\chi_{\bar{\zeta}} = \chi_{\xi} * \bar{\zeta}$ for any $1_{\bar{P}_2} \neq \bar{\zeta} \in \text{Irr}_{\mathcal{K}}(\bar{P}_2)$.

Suppose that there is another element ξ' of M different from ξ . Similarly, we can get a subset $\{\chi_{\xi'} * \bar{\zeta} \mid \bar{\zeta} \in \text{Irr}_{\mathcal{K}}(\bar{P}_2)\}$ of $\text{Irr}_{\mathcal{K}}(\bar{C}_G(Q), \bar{b}_Q)$ such that $\bar{\Delta}_Q^{\circ}(\text{Ind}(\xi') - \text{Ind}(\xi')\bar{\zeta}) = \delta'(\chi_{\xi'} - \chi_{\xi'} * \bar{\zeta})$ for any $1_{\bar{P}_2} \neq \bar{\zeta} \in \text{Irr}_{\mathcal{K}}(\bar{P}_2)$ and $\chi_{\xi'}$ is E_2 -stable, where $\delta' \in \{\pm 1\}$. Assume that $\{\chi_{\xi} * \bar{\zeta} \mid \bar{\zeta} \in \text{Irr}_{\mathcal{K}}(\bar{P}_2)\} \cap \{\chi_{\xi'} * \bar{\zeta} \mid \bar{\zeta} \in \text{Irr}_{\mathcal{K}}(\bar{P}_2)\} \neq \emptyset$. Then there is $\bar{\zeta}_0 \in \text{Irr}_{\mathcal{K}}(\bar{P}_2)$ such that $\chi_{\xi} = \chi_{\xi'} * \bar{\zeta}_0$. If $\bar{\zeta}_0 = 1_{\bar{P}_2}$, then $\chi_{\xi} = \chi_{\xi'}$. This implies that $\text{Ind}(\xi) - \text{Ind}(\xi)\bar{\zeta} = \pm(\text{Ind}(\xi') - \text{Ind}(\xi')\bar{\zeta})$ for any $1_{\bar{P}_2} \neq \bar{\zeta} \in \text{Irr}_{\mathcal{K}}(\bar{P}_2)$. This is impossible. Then $\bar{\zeta}_0$ is non-trivial. But it implies that $\chi_{\xi'} * \bar{\zeta}_0^2 = \chi_{\xi'}$ since $\langle \text{Ind}(\xi) - \text{Ind}(\xi)\bar{\zeta}_0, \text{Ind}(\xi') - \text{Ind}(\xi')\bar{\zeta}_0 \rangle = 0$. It is well-known that $\text{Irr}_{\mathcal{K}}(\bar{P}_2) \setminus \{1_{\bar{P}_2}\}$ acts freely on irreducible ordinary characters of height zero in the block \bar{b}_Q of $\bar{C}_G(Q)$ (see [10, §1]). Hence, $\bar{\zeta}_0^2 = 1_{\bar{P}_2}$ since the defect group of the block \bar{b}_Q of $\bar{C}_G(Q)$ is cyclic. But it is impossible because p is odd. Then

$$\{\chi_{\xi} * \bar{\zeta} \mid \bar{\zeta} \in \text{Irr}_{\mathcal{K}}(\bar{P}_2)\} \cap \{\chi_{\xi'} * \bar{\zeta} \mid \bar{\zeta} \in \text{Irr}_{\mathcal{K}}(\bar{P}_2)\} = \emptyset$$

for any different $\xi, \xi' \in M$. It is clear that $\chi_{\xi} * \bar{\zeta}$ is an irreducible ordinary character in the block \bar{b}_Q of $\bar{C}_G(Q)$ by [2, Corollary]. Then we get an injective isometry

$$\Psi : \mathbb{Z}\{\text{Ind}(\xi)\bar{\zeta} \mid \xi \in M, \bar{\zeta} \in \text{Irr}_{\mathcal{K}}(\bar{P}_2)\} \longrightarrow \mathcal{L}_{\mathcal{K}}(\bar{C}_G(Q), \bar{b}_Q)$$

mapping $\text{Ind}(\xi)\bar{\zeta}$ to $\delta_{\xi}(\chi_{\xi} * \bar{\zeta})$ such that $\Psi(\text{Ind}(\xi) - \text{Ind}(\xi)\bar{\zeta}) = \bar{\Delta}_Q^{\circ}(\text{Ind}(\xi) - \text{Ind}(\xi)\bar{\zeta})$ and χ_{ξ} is E_2 -stable for any $\xi \in M$ and $1_{\bar{P}_2} \neq \bar{\zeta} \in \text{Irr}_{\mathcal{K}}(\bar{P}_2)$, where $\delta_{\xi} \in \{\pm 1\}$.

At the same time, $\tau - \tau\bar{\zeta} \in \mathcal{L}_{\mathcal{K}}^{\circ}(\bar{C}_N(Q), \bar{e}_{\theta})$ for any $\tau \in \text{Irr}_{\mathcal{K}}(\tilde{E}_1)_{\theta}$ and $1_{\bar{P}_2} \neq \bar{\zeta} \in \text{Irr}_{\mathcal{K}}(\bar{P}_2)$. Take an element τ of $\text{Irr}_{\mathcal{K}}(\tilde{E}_1)_{\theta}$. With the same arguments as above, we can get an element χ_{τ} of $\text{Irr}_{\mathcal{K}}(\bar{C}_G(Q), \bar{b}_Q)$ and $\delta_{\tau} \in \{\pm 1\}$ such that $\bar{\Delta}_Q^{\circ}(\tau - \tau\bar{\zeta}) = \delta_{\tau}(\chi_{\tau} - \chi_{\tau} * \bar{\zeta})$ for any $1_{\bar{P}_2} \neq \bar{\zeta} \in \text{Irr}_{\mathcal{K}}(\bar{P}_2)$. Choosing any $1 \neq a \in E_2$ and $1_{\bar{P}_2} \neq \bar{\zeta} \in \text{Irr}_{\mathcal{K}}(\bar{P}_2)$, since $\bar{\Delta}_Q^{\circ}$ is E_2 -stable, we have

$$\delta_{a\tau}(\chi_{a\tau} - \chi_{a\tau} * {}^a\bar{\zeta}) = \bar{\Delta}_Q^{\circ}({}^a\tau - {}^a\tau({}^a\bar{\zeta})) = {}^a(\bar{\Delta}_Q^{\circ}(\tau - \tau\bar{\zeta})) = \delta_{\tau}({}^a\chi_{\tau} - {}^a\chi_{\tau} * {}^a\bar{\zeta}).$$

Then $\chi_{a\tau} = {}^a\chi_{\tau}$ or $\chi_{a\tau} = {}^a\chi_{\tau} * {}^a\bar{\zeta}$. If $\chi_{a\tau} = {}^a\chi_{\tau} * {}^a\bar{\zeta}$, then ${}^a\chi_{\tau} = \chi_{a\tau} * {}^a\bar{\zeta}$. Therefore, $\chi_{a\tau} = \chi_{a\tau} * {}^a(\bar{\zeta}^2)$, which is impossible. Hence, $\chi_{a\tau} = {}^a\chi_{\tau}$ and $\delta_{a\tau} = \delta_{\tau}$ for any $a \in E_2$ since E_2 acts transitively on $\text{Irr}_{\mathcal{K}}(\tilde{E}_1)_{\theta}$. And we denote δ_{τ} by δ . By the facts that $\langle \tau - \tau\bar{\zeta}, \tau' - \tau'\bar{\zeta}' \rangle = 0$ and $\langle \text{Ind}(\xi) - \text{Ind}(\xi)\bar{\zeta}, \tau - \tau\bar{\zeta}' \rangle = 0$ for any $\tau \neq \tau' \in \text{Irr}_{\mathcal{K}}(\tilde{E}_1)_{\theta}$ and $\xi \in M$ and $\bar{\zeta}, \bar{\zeta}' \in \text{Irr}_{\mathcal{K}}(\bar{P}_2) - \{1_{\bar{P}_2}\}$, we can get

$$\{\chi_{\tau} * \bar{\zeta} \mid \bar{\zeta} \in \text{Irr}_{\mathcal{K}}(\bar{P}_2)\} \cap \{\chi_{\tau'} * \bar{\zeta} \mid \bar{\zeta} \in \text{Irr}_{\mathcal{K}}(\bar{P}_2)\} = \emptyset$$

and

$$\{\chi_\xi * \bar{\zeta} \mid \xi \in M, \bar{\zeta} \in \text{Irr}_{\mathcal{K}}(\bar{P}_2)\} \cap \{\chi_\tau * \bar{\zeta} \mid \tau \in \text{Irr}_{\mathcal{K}}(\tilde{E}_1)_\theta, \bar{\zeta} \in \text{Irr}_{\mathcal{K}}(\bar{P}_2)\} = \emptyset.$$

Hence, we have a well-defined E_2 -stable bijective isometry as below

$$\begin{aligned} \bar{\Delta}_Q : \mathcal{L}_{\mathcal{K}}(\bar{C}_N(Q), \bar{e}_\theta) &\longrightarrow \mathcal{L}_{\mathcal{K}}(\bar{C}_G(Q), \bar{b}_Q) \\ \text{Ind}(\xi)\bar{\zeta} &\mapsto \delta_\xi \chi_\xi * \bar{\zeta} \\ {}^a \tau \bar{\zeta} &\mapsto \delta^a \chi_\tau * \bar{\zeta}. \end{aligned}$$

It suffices to show that $\bar{\Delta}_Q$ is an extension of $\bar{\Delta}_Q^\circ$, namely,

$$\bar{\Delta}_Q^\circ(\text{Ind}(\xi) - \sum_{i=1}^l \tau_i) = \delta_\xi \chi_\xi - \delta \sum_{i=1}^l \chi_{\tau_i}$$

for any $\xi \in M$.

Choose an element ξ of M . Since $\langle \text{Ind}(\xi) - \sum_{i=1}^l \tau_i, \tau - \tau\bar{\zeta} \rangle = -1$, then at least χ_τ and $\chi_\tau * \bar{\zeta}$ must get involved in $\bar{\Delta}_Q^\circ(\text{Ind}(\xi) - \sum_{i=1}^l \tau_i)$ for any $\tau \in \text{Irr}_{\mathcal{K}}(\tilde{E}_1)_\theta$ and $\bar{\zeta} \in \text{Irr}_{\mathcal{K}}(\bar{P}_2) - \{1_{\bar{P}_2}\}$.

Keep the notation as above. Suppose that there are τ and $\bar{\zeta}$ such that $\chi_\tau * \bar{\zeta}$ gets involved in $\bar{\Delta}_Q^\circ(\text{Ind}(\xi) - \sum_{i=1}^l \tau_i)$. Since $\langle \text{Ind}(\xi) - \sum_{i=1}^l \tau_i, \tau\bar{\zeta} - \tau\bar{\zeta}' \rangle = 0$ for any $\bar{\zeta}' \in \text{Irr}_{\mathcal{K}}(\bar{P}_2)$ different from $\bar{\zeta}$ and $1_{\bar{P}_2}$, $\chi_\tau * \bar{\zeta}$ must get involved in $\bar{\Delta}_Q^\circ(\text{Ind}(\xi) - \sum_{i=1}^l \tau_i)$ for any $\bar{\zeta}$. At the same time, since $\bar{\Delta}_Q^\circ$ is E_2 -stable and $\text{Ind}(\xi) - \sum_{i=1}^l \tau_i$ is E_2 -stable, we have ${}^a(\chi_\tau * \bar{\zeta})$ must get involved in $\bar{\Delta}_Q^\circ(\text{Ind}(\xi) - \sum_{i=1}^l \tau_i)$ for any $a \in E_2$ and $\bar{\zeta}$. Then there are at least $l \cdot (|\bar{P}_2| - 1)$ different irreducible characters involved in $\bar{\Delta}_Q^\circ(\text{Ind}(\xi) - \sum_{i=1}^l \tau_i)$. This is impossible since $\langle \text{Ind}(\xi) - \sum_{i=1}^l \tau_i, \text{Ind}(\xi) - \sum_{i=1}^l \tau_i \rangle = 1 + l$ and $|\bar{P}_2| - 1 \geq 2$ and $l > 1$.

So for any $\xi \in M$, $\bar{\Delta}_Q^\circ(\text{Ind}(\xi) - \sum_{i=1}^l \tau_i) = a_\chi \chi - \delta \sum_{\tau} \chi_\tau$. Here, $a_\chi \in \{\pm 1\}$ and χ is an element of $\text{Irr}_{\mathcal{K}}(\bar{C}_G(Q), \bar{b}_Q) - \{\chi_\tau \mid \tau \in \text{Irr}_{\mathcal{K}}(\tilde{E}_1)_\theta\}$. Since $\langle \text{Ind}(\xi) - \sum_{i=1}^l \tau_i, \text{Ind}(\xi) - \text{Ind}(\xi)\bar{\zeta} \rangle = 1$ for any $\bar{\zeta} \neq 1_{\bar{P}_2}$, we have $a_\chi = \delta_\xi$ and $\chi = \chi_\xi$. We are done. \square

Then the proof of Theorem 1.1 will follow by Theorem 3.3 and [9, 3.4.2].

References

- [1] M. Broué, Isométries parfaites, types de blocs, catégories dérivées, *Astérisque*, **181-182** (1990), 61-92.
- [2] M. Broué, L. Puig, Characters and local structure in G -algebras, *J. Algebra* **63** (1980), 306-317.
- [3] E. C. Dade, Block extensions, *Illinois, J. Math.* **17** (1973), 198-272.
- [4] F. DeMeyer, G. Janusz, Finite groups with an irreducible representation of large degree, *Math. Z.* **108** (1969), 145-153.
- [5] R. Kessar, M. Linckelmann, On stable equivalences and blocks with one simple module, *J. Algebra* **323** (2010), 1607-1621.

- [6] R. Knörr, G. Robinson, Some remarks on a conjecture of Alperin, *J. London Math. Soc. (2)* **39** (1989), 48-60.
- [7] B. Külshammer, Morita equivalent blocks in Clifford theory of finite groups, *Astérisque*, **181-182** (1990), 209-215.
- [8] L. Puig, Nilpotent extensions of blocks, *Math. Z.* **269** (2011), 115-136.
- [9] L. Puig, Y. Usami, Perfect isometries for blocks with abelian defect groups and Klein four inertial quotients, *J. Algebra* **160** (1993), 192-225.
- [10] G. Robinson, On the local defect group of a block, characters of height zero, and lower defect group multiplicities, *J. Algebra* **320** (2008), 2624-2628.
- [11] A. Watanabe, On perfect isometries for blocks with abelian defect groups and cyclic hyperfocal subgroups, *Kumamoto J. Math.* **18** (2005), 85-92.
- [12] A. Watanabe, The number of irreducible Brauer characters in a p -block of a finite group with cyclic hyperfocal subgroup, *J. Algebra* **416** (2014), 167-183.
- [13] C. Wu, K. Zhang, Y. Zhou, Blocks with defect group $\mathbb{Z}_{2^n} \times \mathbb{Z}_{2^n} \times \mathbb{Z}_{2^m}$, *J. Algebra* **510** (2018), 469-498.