

ON THE MONOID OF COFINITE PARTIAL ISOMETRIES OF \mathbb{N}^n WITH THE USUAL METRIC

OLEG GUTIK AND ANATOLII SAVCHUK

ABSTRACT. In this paper we study the structure of the monoid \mathbf{IN}_∞^n of cofinite partial isometries of the n -th power of the set of positive integers \mathbb{N} with the usual metric for a positive integer $n \geq 2$. We describe the elements of the monoid \mathbf{IN}_∞^n as partial transformation of \mathbb{N}^n , the group of units and the subset of idempotents of the semigroup \mathbf{IN}_∞^n , the natural partial order and Green's relations on \mathbf{IN}_∞^n . In particular we show that the quotient semigroup $\mathbf{IN}_\infty^n/\mathfrak{C}_{\text{mg}}$, where \mathfrak{C}_{mg} is the minimum group congruence on \mathbf{IN}_∞^n , is isomorphic to the symmetric group \mathcal{S}_n and $\mathcal{D} = \mathcal{J}$ in \mathbf{IN}_∞^n . Also, we prove that for any integer $n \geq 2$ the semigroup \mathbf{IN}_∞^n is isomorphic to the semidirect product $\mathcal{S}_n \times_{\text{h}} (\mathcal{P}_\infty(\mathbb{N}^n), \cup)$ of the free semilattice with the unit $(\mathcal{P}_\infty(\mathbb{N}^n), \cup)$ by the symmetric group \mathcal{S}_n .

1. INTRODUCTION AND PRELIMINARIES

In this paper we shall follow the terminology of [3, 13]. We shall denote the the cardinality of the set A by $|A|$. For any positive integer n by \mathcal{S}_n we denote the group of permutations of the set $\{1, \dots, n\}$.

A semigroup S is called *inverse* if for any element $x \in S$ there exists a unique $x^{-1} \in S$ such that $xx^{-1}x = x$ and $x^{-1}xx^{-1} = x^{-1}$. The element x^{-1} is called the *inverse of $x \in S$* . If S is an inverse semigroup, then the function $\text{inv}: S \rightarrow S$ which assigns to every element x of S its inverse element x^{-1} is called an *inversion*.

If S is a semigroup, then we shall denote the subset of all idempotents in S by $E(S)$. If S is an inverse semigroup, then $E(S)$ is closed under multiplication and we shall refer to $E(S)$ as a *band* (or the *band of S*). Then the semigroup operation on S determines the following partial order \preceq on $E(S)$: $e \preceq f$ if and only if $ef = fe = e$. This order is called the *natural partial order* on $E(S)$. A *semilattice* is a commutative semigroup of idempotents.

A linearly ordered subset of a poset is called a *chain*. An ω -*chain* is a chain which is order isomorphic to the set of all negative integers with the usual order \leq .

If S is an inverse semigroup then the semigroup operation on S determines the following partial order \preceq on S : $s \preceq t$ if and only if there exists $e \in E(S)$ such that $s = te$. This order is called the *natural partial order* on S [17].

By $(\mathcal{P}_\infty(X), \cup)$ we shall denote the *free semilattice with identity* over a set X of cardinality $\geq \omega$, i.e., $(\mathcal{P}_\infty(X), \cup)$ is the set of all finite subsets (with the empty set) of X with the semilattice operation “union”.

A congruence \mathfrak{C} on a semigroup S is called *non-trivial* if \mathfrak{C} is distinct from universal and identity congruences on S , and a *group congruence* if the quotient semigroup S/\mathfrak{C} is a group. If \mathfrak{C} is a congruence on a semigroup S then by \mathfrak{C}^\sharp we denote the natural homomorphism from S onto the quotient semigroup S/\mathfrak{C} . Every inverse semigroup S admits a least (minimum) group congruence \mathfrak{C}_{mg} :

$$a\mathfrak{C}_{\text{mg}}b \text{ if and only if there exists } e \in E(S) \text{ such that } ae = be$$

(see [15, Lemma III.5.2]).

Date: September 20, 2019.

2010 Mathematics Subject Classification. 20M18, 20M20, 20M30.

Key words and phrases. Partial isometry, inverse semigroup, partial bijection, natural partial order, Green's relations, least group congruence, F -inverse semigroup, semidirect product, free semilattice, symmetric group.

If S is a semigroup, then we shall denote the Green relations on S by \mathcal{R} , \mathcal{L} , \mathcal{J} , \mathcal{D} and \mathcal{H} (see [4] or [3, Section 2.1]):

$$\begin{aligned} a\mathcal{R}b &\text{ if and only if } aS^1 = bS^1; \\ a\mathcal{L}b &\text{ if and only if } S^1a = S^1b; \\ a\mathcal{J}b &\text{ if and only if } S^1aS^1 = S^1bS^1; \\ \mathcal{D} &= \mathcal{L} \circ \mathcal{R} = \mathcal{R} \circ \mathcal{L}; \\ \mathcal{H} &= \mathcal{L} \cap \mathcal{R}. \end{aligned}$$

If $\alpha: X \rightarrow Y$ is a partial map, then we shall denote the domain and the range of α by $\text{dom } \alpha$ and $\text{ran } \alpha$, respectively. A partial map $\alpha: X \rightarrow Y$ is called *cofinite* if both sets $X \setminus \text{dom } \alpha$ and $Y \setminus \text{ran } \alpha$ are finite.

Let \mathcal{I}_λ denote the set of all partial one-to-one transformations of a non-zero cardinal λ together with the following semigroup operation:

$$x(\alpha\beta) = (x\alpha)\beta \quad \text{if } x \in \text{dom}(\alpha\beta) = \{y \in \text{dom } \alpha : y\alpha \in \text{dom } \beta\}, \quad \text{for } \alpha, \beta \in \mathcal{I}_\lambda.$$

The semigroup \mathcal{I}_λ is called the *symmetric inverse (monoid) semigroup* over cardinal λ (see [3]). The symmetric inverse semigroup was introduced by Wagner [17] and it plays a major role in the theory of semigroups. By $\mathcal{I}_\lambda^{\text{cf}}$ is denoted a subsemigroup of injective partial selfmaps of λ with cofinite domains and ranges in \mathcal{I}_λ . Obviously, $\mathcal{I}_\lambda^{\text{cf}}$ is an inverse submonoid of the semigroup \mathcal{I}_λ . The semigroup $\mathcal{I}_\lambda^{\text{cf}}$ is called the *monoid of injective partial cofinite selfmaps* of λ [9].

A partial transformation $\alpha: (X, d) \rightarrow (X, d)$ of a metric space (X, d) is called *isometric* or a *partial isometry*, if $d(x\alpha, y\alpha) = d(x, y)$ for all $x, y \in \text{dom } \alpha$. It is obvious that the composition of two partial isometries of a metric space (X, d) is a partial isometry, and the converse partial map to a partial isometry is a partial isometry, too. Hence the set of partial isometries of a metric space (X, d) with the operation the composition of partial isometries is an inverse submonoid of the symmetric inverse monoid over the cardinal $|X|$. Also, it is obvious that the set of partial cofinite isometries of a metric space (X, d) with the operation the composition of partial isometries is an inverse submonoid of the monoid of injective partial cofinite selfmaps of the cardinal $|X|$.

The semigroup \mathbf{ID}_∞ of all partial cofinite isometries of the set of integers \mathbb{Z} with the usual metric $d(n, m) = |n - m|$, $n, m \in \mathbb{Z}$ established in the Bezushchak papers [1, 2]. In [1] the generators of the semigroup \mathbf{ID}_∞ are described and there proved that \mathbf{ID}_∞ has the exponential growth. We remark that the semigroup \mathbf{ID}_∞ is inverse submonoid of the monoid of all partial cofinite bijections of \mathbb{Z} , and elements of \mathbf{ID}_∞ are restrictions of isometries of \mathbb{Z} onto its cofinite subsets in the Lawson interpretation (see [13, p. 9]). Green's relations and principal ideals of \mathbf{ID}_∞ are described in [2]. In [10] is shown that the quotient semigroup $\mathbf{ID}_\infty/\mathbf{C}_{\text{mg}}$ is isomorphic to the group $\mathbf{Iso}(\mathbb{Z})$ of all isometries of \mathbb{Z} , the semigroup \mathbf{ID}_∞ is F -inverse, and \mathbf{ID}_∞ is isomorphic to the semidirect product $\mathbf{Iso}(\mathbb{Z}) \rtimes_{\mathfrak{h}} \mathcal{P}_\infty(\mathbb{Z})$ of the free semilattice $(\mathcal{P}_\infty(\mathbb{Z}), \cup)$ by the group $\mathbf{Iso}(\mathbb{Z})$. Also in [10] established semigroup and shift-continuous topologies on \mathbf{ID}_∞ and embedding of the discrete semigroup \mathbf{ID}_∞ into compact-like topological semigroups.

Let \mathbf{IN}_∞ be the set of all partial cofinite isometries of the set of positive integers \mathbb{N} with the usual metric $d(n, m) = |n - m|$, $n, m \in \mathbb{N}$. Then \mathbf{IN}_∞ with the operation of composition of partial isometries is an inverse submonoid of \mathcal{I}_ω . The semigroup \mathbf{IN}_∞ of all partial co-finite isometries of positive integers is studied in [11]. There we describe the Green relations on the semigroup \mathbf{IN}_∞ , its band and proved that \mathbf{IN}_∞ is a simple E -unitary F -inverse semigroup. Also in [11], the least group congruence \mathbf{C}_{mg} on \mathbf{IN}_∞ is described and proved that the quotient-semigroup $\mathbf{IN}_\infty/\mathbf{C}_{\text{mg}}$ is isomorphic to the additive group of integers $\mathbb{Z}(+)$. An example of a non-group congruence on the semigroup \mathbf{IN}_∞ is presented. Also we proved that a congruence on the semigroup \mathbf{IN}_∞ is group if and only if its restriction onto an isomorphic copy of the bicyclic semigroup in \mathbf{IN}_∞ is a group congruence.

For an arbitrary positive integer $n \geq 2$ by \mathbb{N}^n we denote the n -th power of the set of positive integers \mathbb{N} with the usual metric:

$$d((x_1, \dots, x_n), (y_1, \dots, y_n)) = \sqrt{(x_1 - y_1)^2 + \dots + (x_n - y_n)^2}.$$

Let \mathbf{IN}_∞^n be the set of all partial cofinite isometries of \mathbb{N}^n . It is obvious that \mathbf{IN}_∞^n with the operation of composition of partial isometries is an inverse submonoid of \mathcal{S}_ω and later by \mathbf{IN}_∞^n we shall denote the monoid of all partial cofinite isometries of \mathbb{N}^n .

By \mathbb{I} we denote the identity map of \mathbb{N}^n which obviously is the unit of the semigroup \mathbf{IN}_∞^n . Later by $H(\mathbb{I})$ we shall denote the group of units of \mathbf{IN}_∞^n .

In the paper we study the structure of the monoid \mathbf{IN}_∞^n . We describe the elements of the monoid \mathbf{IN}_∞^n as partial transformations of \mathbb{N}^n , the group of units and the subset of idempotents of \mathbf{IN}_∞^n , the natural partial order and Green's relations on \mathbf{IN}_∞^n . In particular we show that the quotient semigroup $\mathbf{IN}_\infty^n / \mathcal{C}_{\text{mg}}$ is isomorphic to the symmetric group \mathcal{S}_n and $\mathcal{D} = \mathcal{J}$ in \mathbf{IN}_∞^n . Also, we prove that for any integer $n \geq 2$ the semigroup \mathbf{IN}_∞^n is isomorphic to the semidirect product $\mathcal{S}_n \ltimes_{\mathfrak{h}} (\mathcal{P}_\infty(\mathbb{N}^n), \cup)$ of free semilattice with the unit $(\mathcal{P}_\infty(\mathbb{N}^n), \cup)$ by the symmetric group \mathcal{S}_n .

2. PROPERTIES OF PARTIAL COFINITE ISOMETRIES OF \mathbb{N}^n

The definition of an isometry implies the following proposition.

Proposition 2.1. *Let n be a positive integer ≥ 2 . Then every permutation σ of the set $\{1, \dots, n\}$ induces an isometry $\alpha_\sigma: \mathbb{N}^n \rightarrow \mathbb{N}^n$, $\alpha_\sigma: x \mapsto x \circ \alpha$, of the set \mathbb{N}^n .*

We denote $\mathbf{1} = (1, \dots, 1)$ and $\vec{\mathbf{1}}_i = \{(1, \dots, \underbrace{x_i}_{i\text{-th}}, \dots, 1) \in \mathbb{N}^n: x_i \in \mathbb{N}\}$ for any $i \in \{1, \dots, n\}$. Also for any $\mathbf{x} = (x_1, \dots, x_n) \in \mathbb{N}^n$ and any positive real number r we put

$$\mathcal{M}_r(\mathbf{x}) = |\{\mathbf{y} \in \mathbb{N}^n: d(\mathbf{x}, \mathbf{y}) = r\}|.$$

Lemma 2.2. *Let n be a positive integer ≥ 2 . Then the following statements holds.*

- (i) $\mathcal{M}_1(\mathbf{1}) = n$.
- (ii) *If for $\mathbf{x} \in \mathbb{N}^n$ exactly k coordinates of \mathbf{x} distinct from 1 for some $k = 1, \dots, n$, then $\mathcal{M}_1(\mathbf{x}) = n + k$.*

Proof. (i) It is obvious that the set $\{\mathbf{y} \in \mathbb{N}^n: d(\mathbf{1}, \mathbf{y}) = 1\}$ consists of

$$\left\{ \left(\underbrace{2}_{1\text{-st}}, 1, \dots, 1 \right), \left(1, \underbrace{2}_{2\text{-nd}}, \dots, 1 \right), \dots, \left(1, \dots, \underbrace{2}_{n\text{-th}} \right) \right\},$$

which implies the equality $\mathcal{M}_1(\mathbf{1}) = n$.

(ii) Fix an arbitrary $\mathbf{x} \in \mathbb{N}^n$ such that exactly k coordinates of \mathbf{x} distinct from 1 for some $k \in \{1, \dots, n\}$. By Proposition 2.1 without loss of generality we may assume that only the first k coordinates of \mathbf{x} distinct from 1, i.e., $\mathbf{x} = (x_1, \dots, x_k, 1, \dots, 1)$ for some $x_1 > 1, \dots, x_k > 1$. This implies that the set $\{\mathbf{y} \in \mathbb{N}^n: d(\mathbf{x}, \mathbf{y}) = 1\}$ coincides with the following set

$$\begin{aligned} & \{(x_1 - 1, \dots, x_k, 1, \dots, 1), (x_1 + 1, \dots, x_k, 1, \dots, 1), \dots, \\ & (x_1, \dots, x_k - 1, 1, \dots, 1), (x_1, \dots, x_k + 1, 1, \dots, 1), \\ & (x_1, \dots, x_k, \underbrace{2}_{(k+1)\text{-th}}, \dots, 1), \dots, (x_1, \dots, x_k, 1, \dots, \underbrace{2}_{n\text{-th}})\}, \end{aligned}$$

and hence $\mathcal{M}_1(\mathbf{x}) = n + k$. □

Lemma 2.2(i) implies the following corollary.

Corollary 2.3. *Let n be a positive integer ≥ 2 . Then $(\mathbf{1})\alpha = \mathbf{1}$ for every isometry α of \mathbb{N}^n .*

For any positive integers $k > 1$ and m we denote

$$\mathbf{k}_1 = (\underbrace{k}_{1\text{-st}}, 1, \dots, 1), \dots, \mathbf{k}_j = (1, \dots, 1, \underbrace{k}_{j\text{-th}}, 1, \dots, 1), \dots, \mathbf{k}_n = (1, \dots, 1, \underbrace{k}_{n\text{-th}})$$

and

$$\mathbf{C}_m = \{(x_1, \dots, x_n) \in \mathbb{N}^n : x_1 \leq m, \dots, x_n \leq m\}.$$

Later we need the following technical lemma.

Lemma 2.4. *Let x and y be any positive integers. If the sequence $\{a_i\}_{i \in \mathbb{N}}$, where $a_i = \sqrt{(x+i)^2 + y}$, contains an integer then $\{a_i\}_{i \in \mathbb{N}}$ has infinitely many non-integer members.*

Proof. Suppose that $n = \sqrt{x^2 + y}$ is an integer. Since $x < \sqrt{x^2 + y}$ we get that

$$\begin{aligned} 2x < 2\sqrt{x^2 + y} &\iff \\ \iff x^2 + 2x + 1 + y < x^2 + y + 2\sqrt{x^2 + y} + 1 &\iff \\ \iff (x+1)^2 + y < (\sqrt{x^2 + y} + 1)^2 &\iff \\ \iff \sqrt{(x+1)^2 + y} < n + 1, & \end{aligned}$$

which implies that $n < \sqrt{(x+1)^2 + y} < n + 1$. Thus, $\sqrt{(x+1)^2 + y}$ is irrational, and hence the statement of the lemma holds. \square

Lemma 2.5. *Let n be a positive integer ≥ 2 . Let $\alpha: \mathbb{N}^n \rightarrow \mathbb{N}^n$ be a partial cofinite isometry of \mathbb{N}^n . Then for every $i \in \{1, \dots, n\}$ there exists a unique $j(i) \in \{1, \dots, n\}$ and an integer $q(i)$ such that $(\mathbf{m}_i)\alpha = (\mathbf{m} + \mathbf{q}(i))_{j(i)}$ for any $\mathbf{m}_i \in \text{dom } \alpha$. Moreover α determines the permutation $i \mapsto j(i)$ of the set $\{1, \dots, n\}$.*

Proof. Since α is a partial cofinite isometry of \mathbb{N}^n there exists a positive integer $m \geq 3$ such that $\mathbb{N}^n \setminus \text{dom } \alpha \subseteq \mathbf{C}_{m-1}$ and $\mathbb{N}^n \setminus \text{ran } \alpha \subseteq \mathbf{C}_{m-1}$. Since the set \mathbf{C}_{m+1} is finite there exists a positive integer k such that $\mathbf{p}_i \notin \mathbf{C}_{m+1}$ and $(\mathbf{p}_i)\alpha \notin \mathbf{C}_{m+1}$ for any integer $p \geq k$. By Proposition 2.1 without loss of generality we may assume that $i = j(i) = 1$, i.e., $(\mathbf{k}_1)\alpha = \mathbf{q}_1$ for some positive integer q .

We claim that $((\mathbf{k} + \mathbf{s})_1)\alpha = (\mathbf{q} + \mathbf{s})_1$ for any positive integer s . Since $\mathbf{k}_1 \notin \mathbf{C}_{m+1}$, we have that

$$|\{\mathbf{x} \in \text{dom } \alpha : d(\mathbf{x}, (\mathbf{k} - \mathbf{2} + \mathbf{t})_1) = 1\}| = n + 1,$$

for every positive integer t . Since α is a partial cofinite isometry of \mathbb{N}^n and $(\mathbf{k} - \mathbf{1})_1 \in \text{dom } \alpha$, the above implies that either $((\mathbf{k} + \mathbf{1})_1)\alpha = (\mathbf{q} - \mathbf{1})_1$ or $((\mathbf{k} + \mathbf{1})_1)\alpha = (\mathbf{q} + \mathbf{1})_1$. We claim that $((\mathbf{k} + \mathbf{1})_1)\alpha = (\mathbf{q} + \mathbf{1})_1$. Suppose to the contrary that $((\mathbf{k} + \mathbf{1})_1)\alpha = (\mathbf{q} - \mathbf{1})_1$. Next we consider the following sequence $\{(\mathbf{k} + \mathbf{p})_1\}_{p \in \mathbb{N}}$ in \mathbb{N}^n . By the above assumption there exists a positive integer s such that $((\mathbf{k} + \mathbf{p})_1)\alpha \notin \mathbf{C}_{m+1} \cup \vec{\mathbf{I}}_1 \cup \dots \cup \vec{\mathbf{I}}_n$ for all $p \geq s$. By Lemma 2.4 the sequence $\{d(((\mathbf{k} + \mathbf{p})_1)\alpha, (\mathbf{k}_1))\}_{p \in \mathbb{N}}$ contains non-integer numbers. But the sequence $\{d((\mathbf{k} + \mathbf{p})_1, \mathbf{k}_1)\}_{p \in \mathbb{N}}$ has only positive integers, which contradicts that α is a partial isometry of \mathbb{N}^n . The obtained contradiction implies that $((\mathbf{k} + \mathbf{1})_1)\alpha = (\mathbf{q} + \mathbf{1})_1$. Next, similar arguments and induction imply that $((\mathbf{k} + \mathbf{p})_1)\alpha = (\mathbf{q} + \mathbf{p})_1$ for any $p \in \mathbb{N}$.

Fix an arbitrary integer $l < k$ such that $\mathbf{l}_1 \in \text{dom } \alpha$. We claim that $(\mathbf{l}_1)\alpha = (\mathbf{q} - \mathbf{k} + \mathbf{l})_1$. Indeed, in the other case we have that $d(\mathbf{l}_1, (\mathbf{k} + \mathbf{p})_1) = k + p - l$ for any $p \in \mathbb{N}$, and hence the sequence $\{d(\mathbf{l}_1, (\mathbf{k} + \mathbf{p})_1)\}_{p \in \mathbb{N}}$ contains only integer elements. But since $(\mathbf{l}_1)\alpha = (x_1, x_2, \dots, x_n)$ with some $x_i \neq 1$, $i = 2, \dots, n$, we have that

$$d((\mathbf{l}_1)\alpha, ((\mathbf{k} + \mathbf{p})_1)\alpha) = \sqrt{(x_1 - k + p)^2 + (x_2 - 1)^2 + \dots + (x_n - 1)^2},$$

and hence by Lemma 2.4 the sequence $\{d((\mathbf{l}_1)\alpha, ((\mathbf{k} + \mathbf{p})_1)\alpha)\}_{p \in \mathbb{N}}$ contains non-integer elements. This contradicts that α is a partial isometry of \mathbb{N}^n , which completes the proof of the lemma. \square

Lemma 2.6. *Let n be a positive integer ≥ 2 . Let $\alpha: \mathbb{N}^n \rightarrow \mathbb{N}^n$ be a partial cofinite isometry of \mathbb{N}^n such that $(\mathbf{m}_i)\alpha \in \vec{\mathbf{1}}_i$ for some $\mathbf{m}_i \in \text{dom } \alpha \setminus \{\mathbf{1}\}$, $m \in \mathbb{N}$, and any $i \in \{1, \dots, n\}$. Then $(\mathbf{x}_i)\alpha = \mathbf{x}_i$ for any $\mathbf{x}_i \in \text{dom } \alpha \cap \vec{\mathbf{1}}_i$.*

Proof. Suppose to the contrary that there exists $i \in \{1, \dots, n\}$ such that $(\mathbf{x}_i)\alpha \neq \mathbf{x}_i$ for some $\mathbf{x}_i \in \vec{\mathbf{1}}_i$. Without loss of generality we may assume that $i = 1$. Then there exists a non-zero integer p such that $(\mathbf{x}_1)\alpha = (\mathbf{x} + \mathbf{p})_1$. We assume that $p > 0$.

Fix an arbitrary $\mathbf{y}_2 \in \text{dom } \alpha \cap \vec{\mathbf{1}}_2 \setminus \{\mathbf{1}\}$. Lemma 2.5 implies that without loss of generality we may assume that $y < x$. Since α is a partial isometry of \mathbb{N}^n ,

$$(1) \quad d((\mathbf{x}_1)\alpha, (\mathbf{y}_2)\alpha) = d(\mathbf{x}_1, \mathbf{y}_2) = \sqrt{(x-1)^2 + (y-1)^2}.$$

This and the equality $(\mathbf{x}_1)\alpha = (\mathbf{x} + \mathbf{p})_1$ imply that there exists a positive integer q such that $(\mathbf{y}_2)\alpha = (\mathbf{x} - \mathbf{q})_2$. Then we have that

$$(2) \quad d((\mathbf{x}_1)\alpha, (\mathbf{y}_2)\alpha) = d((\mathbf{x} + \mathbf{p})_1, (\mathbf{y} - \mathbf{q})_2) = \sqrt{(x+p-1)^2 + (y-q-1)^2}.$$

By equalities (1) and (2) we get that

$$(x-1)^2 + (y-1)^2 = (x-1)^2 + 2(x-1)p + p^2 + (y-1)^2 - 2(y-1)q + q^2,$$

and hence

$$(3) \quad p^2 + q^2 + 2(xp - yq) - 2(p - q) = 0.$$

Again, since α is a partial isometry of \mathbb{N}^n , Lemma 2.5 implies that

$$(4) \quad d(((\mathbf{x} + \mathbf{1})_1)\alpha, ((\mathbf{y} + \mathbf{1})_2)\alpha) = d((\mathbf{x} + \mathbf{1})_1, (\mathbf{y} + \mathbf{1})_2) = \sqrt{(x+1-1)^2 + (y+1-1)^2} = \sqrt{x^2 + y^2},$$

$((\mathbf{x} + \mathbf{1})_1)\alpha = (\mathbf{x} + \mathbf{1} + \mathbf{p})_1$ and $((\mathbf{y} + \mathbf{1})_2)\alpha = (\mathbf{y} + \mathbf{1} - \mathbf{q})_2$. Then

$$(5) \quad \begin{aligned} d(((\mathbf{x} + \mathbf{1})_1)\alpha, ((\mathbf{y} + \mathbf{1})_2)\alpha) &= d((\mathbf{x} + \mathbf{1} + \mathbf{p})_1, (\mathbf{y} + \mathbf{1} - \mathbf{q})_2) = \\ &= \sqrt{(x+1+p-1)^2 + (y+1-q-1)^2} = \\ &= \sqrt{(x+p)^2 + (y-q)^2}. \end{aligned}$$

By equalities (4) and (5) we have that

$$x^2 + y^2 = x^2 + 2xp + p^2 + y^2 - 2yq + q^2,$$

and after some arithmetic simplifications we get that

$$(6) \quad p^2 + q^2 + 2(xp - yq) = 0.$$

Then equalities (3) and (6) imply that $p = q$. Therefore, $(\mathbf{y}_2)\alpha = (\mathbf{x} - \mathbf{p})_2$. Then equalities (1) and (2) imply that

$$(x-1)^2 + (y-1)^2 = (x-1)^2 + 2(x-1)p + p^2 + (y-1)^2 - 2(y-1)p + p^2,$$

and hence

$$2p(x - y + p) = 0.$$

This contradicts the assumptions $p > 0$ and $x > y$.

In the case when $p < 0$ we choose $x < y$, and by similar way as in the above we get a contradiction. The obtained contradictions imply that $(\mathbf{x}_i)\alpha = \mathbf{x}_i$ for any $\mathbf{x}_i \in \vec{\mathbf{1}}_i$ with $x \geq k$, $i = 1, \dots, n$. This completes the proof of the lemma. \square

Theorem 2.7. *Let n be a positive integer ≥ 2 and $\alpha: \mathbb{N}^n \rightarrow \mathbb{N}^n$ be a partial cofinite isometry of \mathbb{N}^n such that $(\mathbf{m}_i)\alpha \in \vec{\mathbf{1}}_i$ for some $\mathbf{m}_i \in \text{dom } \alpha \setminus \{\mathbf{1}\}$, $m \in \mathbb{N}$, and any $i \in \{1, \dots, n\}$. Then $(\mathbf{x})\alpha = \mathbf{x}$ for any $\mathbf{x} \in \text{dom } \alpha$.*

Proof. By Lemma 2.6 it is sufficiently to show that $(\mathbf{x})\alpha = \mathbf{x}$ for any $\mathbf{x} \in \text{dom } \alpha \setminus (\vec{\mathbf{1}}_1 \cup \dots \cup \vec{\mathbf{1}}_n)$.

Suppose to the contrary that there exists $\mathbf{x} = (x_1, \dots, x_n) \in \text{dom } \alpha \setminus (\vec{\mathbf{1}}_1 \cup \dots \cup \vec{\mathbf{1}}_n)$ with $x_1, \dots, x_n \neq 1$ such that $(\mathbf{x})\alpha = (y_1, \dots, y_n) \neq \mathbf{x}$. Lemma 2.5 implies that $(\mathbf{p}_1)\alpha = \mathbf{p}_1$ for all positive integers p such that $\mathbf{p}_1 \in \text{dom } \alpha$. Then we have that

$$\begin{aligned} d(\mathbf{x}, \mathbf{p}_1) &= d((\mathbf{x})\alpha, (\mathbf{p}_1)\alpha) \iff \\ (x_1 - p)^2 + (x_2 - 1)^2 + \dots + (x_n - 1)^2 &= (y_1 - p)^2 + (y_2 - 1)^2 + \dots + (y_n - 1)^2 \iff \\ x_1^2 - 2x_1p + p^2 + (x_2 - 1)^2 + \dots + (x_n - 1)^2 - y_1^2 + 2y_1p - p^2 - (y_2 - 1)^2 - \dots - (y_n - 1)^2 &= 0 \iff \\ (7) \quad 2(y_1 - x_1)p + (x_2 - 1)^2 + \dots + (x_n - 1)^2 + x_1^2 - y_1^2 - (y_2 - 1)^2 - \dots - (y_n - 1)^2 &= 0. \end{aligned}$$

Since the partial map α is cofinite equation (7) has infinitely many solutions in \mathbb{N} with the respect to the variable p . This implies that $y_1 = x_1$ and

$$(x_2 - 1)^2 + \dots + (x_n - 1)^2 + x_1^2 - y_1^2 - (y_2 - 1)^2 - \dots - (y_n - 1)^2 = 0.$$

Similar arguments and Lemma 2.5 imply that $y_i = x_i$ for any other $i \in \{2, \dots, n\}$. \square

3. ALGEBRAIC PROPERTY OF THE SEMIGROUP \mathbf{IN}_∞^n

Theorem 2.7 implies the following corollary.

Corollary 3.1. *Let n be any positive integer ≥ 2 and $\alpha: \mathbb{N}^n \rightarrow \mathbb{N}^n$ be an isometry such that $(\mathbf{2}_i)\alpha = \mathbf{2}_i$ for all $i = 1, \dots, n$. Then α is the identity map of \mathbb{N}^n .*

The following theorem describes the structure of the group of units $H(\mathbb{I})$ of the semigroup \mathbf{IN}_∞^n .

Theorem 3.2. *For any positive integer $n \geq 2$ the group of units $H(\mathbb{I})$ of the semigroup \mathbf{IN}_∞^n is isomorphic to the group \mathcal{S}_n . Moreover, every element of $H(\mathbb{I})$ is induced by a permutation of the set $\{1, \dots, n\}$.*

Proof. It is obvious that every element of the group of units $H(\mathbb{I})$ of the semigroup \mathbf{IN}_∞^n is an isometry of \mathbb{N}^n .

Fix an arbitrary isometry α of \mathbb{N}^n . Since the set $\{\mathbf{y} \in \mathbb{N}^n: d(\mathbf{1}, \mathbf{y}) = 1\}$ coincides with

$$\mathbf{D}_1(\mathbf{1}) = \left\{ \left(\underbrace{2}_{1\text{-st}}, 1, \dots, 1 \right), \left(1, \underbrace{2}_{2\text{-nd}}, \dots, 1 \right), \dots, \left(1, \dots, \underbrace{2}_{n\text{-th}} \right) \right\},$$

we have that $(\mathbf{D}_1(\mathbf{1}))\alpha = \mathbf{D}_1(\mathbf{1})$. Then there exists a permutation $\sigma: \{1, \dots, n\} \rightarrow \{1, \dots, n\}$ such that $(\mathbf{2}_i)\alpha = \mathbf{2}_{(i)\sigma}$ for any $i = 1, \dots, n$. By Proposition 2.1 the permutation σ induces the isometry $\alpha_\sigma: \mathbb{N}^n \rightarrow \mathbb{N}^n$, $x \mapsto x \circ \alpha$ of the set \mathbb{N}^n . It is obvious that $(\mathbf{2}_i)\alpha\alpha_\sigma^{-1} = \mathbf{2}_i$ and $(\mathbf{2}_j)\alpha_\sigma^{-1}\alpha = \mathbf{2}_j$ for all $i, j = 1, \dots, n$. Since $\alpha\alpha_\sigma^{-1}$ and $\alpha_\sigma^{-1}\alpha$ are isometries of \mathbb{N}^n , Corollary 3.1 implies that $\alpha\alpha_\sigma^{-1} = \alpha_\sigma^{-1}\alpha = \mathbb{I}$. This implies that the isometry α is induced by some permutation of the set $\{1, \dots, n\}$, which completes the proof of the theorem. \square

Proposition 3.3. *Let n be any positive integer ≥ 2 . Then the following statements hold.*

- (i) *An element α of the semigroup \mathbf{IN}_∞^n is an idempotent if and only if $(x)\alpha = x$ for every $x \in \text{dom } \alpha$.*
- (ii) *If $\varepsilon, \iota \in E(\mathbf{IN}_\infty^n)$, then $\varepsilon \leq \iota$ if and only if $\text{dom } \varepsilon \subseteq \text{dom } \iota$.*
- (iii) *The semilattice $E(\mathbf{IN}_\infty^n)$ is isomorphic to the free semilattice $(\mathcal{P}_\infty(\mathbb{N}^n), \cup)$ under the mapping $(\varepsilon)h = \mathbb{N} \setminus \text{dom } \varepsilon$.*
- (iv) *Every maximal chain in $E(\mathbf{IN}_\infty^n)$ is an ω -chain.*
- (v) *\mathbf{IN}_∞^n is an inverse semigroup.*
- (vi) *$\alpha \mathcal{R} \beta$ in \mathbf{IN}_∞^n if and only if $\text{dom } \alpha = \text{dom } \beta$.*
- (vii) *$\alpha \mathcal{L} \beta$ in \mathbf{IN}_∞^n if and only if $\text{ran } \alpha = \text{ran } \beta$.*
- (viii) *$\alpha \mathcal{H} \beta$ in \mathbf{IN}_∞^n if and only if $\text{dom } \alpha = \text{dom } \beta$ and $\text{ran } \alpha = \text{ran } \beta$.*

Proof. Statements (i) – (iv) are trivial and their proofs follow from the definition of the semigroup \mathbf{IN}_∞^n .

(v) Fix an arbitrary $\alpha \in \mathbf{IN}_\infty^n$. Since α is a partial cofinite isometry of \mathbb{N}^n , we conclude that so is its inverse α^{-1} . This implies that \mathbf{IN}_∞^n is regular, and hence by statement (iii) and the Wagner-Preston Theorem (see [3, Theorem 1.17]), \mathbf{IN}_∞^n is an inverse semigroup.

Statements (vi) – (viii) follow from the description of Green's relations \mathcal{R} , \mathcal{L} and \mathcal{H} on the symmetric inverse monoid \mathcal{I}_ω and Proposition 3.2.11 of [13]. \square

Lemma 3.4. *Let n be any positive integer ≥ 2 and α be an arbitrary element of the semigroup \mathbf{IN}_∞^n . Then there exist unique elements σ_l and σ_r of the group of units $H(\mathbb{I})$ of the semigroup \mathbf{IN}_∞^n such that $\sigma_l\alpha$ and $\alpha\sigma_r$ are idempotents in \mathbf{IN}_∞^n .*

Proof. Suppose that $\sigma: \{1, \dots, n\} \rightarrow \{1, \dots, n\}$, $i \mapsto j(i)$ is a permutation defined in Lemma 2.5 for the partial isometry α . We put σ_α is an element of $H(\mathbb{I})$ which is induced by the permutation σ of the set $\{1, \dots, n\}$ (see Theorem 3.2) and σ_α^{-1} is its inverse. Then the assumption of Theorem 2.7 holds for $\alpha\sigma_\alpha^{-1}$, and hence by Proposition 3.3(i), $\alpha\sigma_\alpha^{-1}$ is an idempotent of \mathbf{IN}_∞^n . Next we put $\sigma_r = \sigma_\alpha^{-1}$.

By a similar way we suppose that $\bar{\sigma}: \{1, \dots, n\} \rightarrow \{1, \dots, n\}$, $i \mapsto j(i)$ is a permutation defined in Lemma 2.5 for α^{-1} , i.e., $\bar{\sigma}$ is inverse of σ . We put $\bar{\sigma}_\alpha$ is an element of $H(\mathbb{I})$ which is induced by the permutation $\bar{\sigma}$ of the set $\{1, \dots, n\}$ and $\bar{\sigma}_\alpha^{-1}$ is its inverse. Then by Proposition 3.3(i) the element $\alpha^{-1}\bar{\sigma}_\alpha^{-1}$ is an idempotent of \mathbf{IN}_∞^n , and hence so is its inverse $(\alpha^{-1}\bar{\sigma}_\alpha^{-1})^{-1} = \bar{\sigma}_\alpha\alpha$, because by Proposition 3.3(v) the semigroup \mathbf{IN}_∞^n is inverse. Hence, we put $\sigma_l = \sigma_\alpha$.

We observe that the uniqueness of the elements σ_l and σ_r follows from Lemma 2.5. \square

Lemma 3.5. *Let n be any positive integer ≥ 2 and α be an arbitrary element of the semigroup \mathbf{IN}_∞^n . Then there exists the unique element σ_α of the group of units $H(\mathbb{I})$ and the unique idempotents $\varepsilon_{l(\alpha)}$ and $\varepsilon_{r(\alpha)}$ of the semigroup \mathbf{IN}_∞^n such that*

$$\alpha = \sigma_\alpha \varepsilon_{l(\alpha)} = \varepsilon_{r(\alpha)} \sigma_\alpha.$$

Proof. By Lemma 3.4 there exists unique elements σ_{l_α} and σ_{r_α} of the group of units $H(\mathbb{I})$ such that $\sigma_{l_\alpha}\alpha$ and $\alpha\sigma_{r_\alpha}$ are idempotents of the semigroup \mathbf{IN}_∞^n . This implies that

$$(8) \quad \sigma_{l_\alpha}\alpha = \alpha^{-1}\alpha \quad \text{and} \quad \alpha\sigma_{r_\alpha} = \alpha\alpha^{-1},$$

because \mathbf{IN}_∞^n is an inverse semigroup, $\text{ran}(\sigma_{l_\alpha}\alpha) = \text{ran}(\alpha^{-1}\alpha)$ and $\text{dom}(\alpha\sigma_{r_\alpha}) = \text{dom}(\alpha\alpha^{-1})$. Hence we get that

$$\alpha = \sigma_{l_\alpha}\sigma_{l_\alpha}\alpha = \sigma_{l_\alpha}\alpha^{-1}\alpha \quad \text{and} \quad \alpha = \alpha\sigma_{r_\alpha}\sigma_{r_\alpha}^{-1} = \alpha\alpha^{-1}\sigma_{r_\alpha}^{-1}.$$

We put

$$\sigma_{l(\alpha)} = \sigma_{l_\alpha}, \quad \sigma_{r(\alpha)} = \sigma_{r_\alpha}^{-1}, \quad \varepsilon_{l(\alpha)} = \alpha^{-1}\alpha, \quad \text{and} \quad \varepsilon_{r(\alpha)} = \alpha\alpha^{-1}.$$

We claim that $\sigma_{l(\alpha)} = \sigma_{r(\alpha)}$. Indeed, the inequality $\sigma_{l(\alpha)} \neq \sigma_{r(\alpha)}$ and Theorem 3.2 imply that the restriction $\sigma_{l(\alpha)}^{-1}\sigma_{r(\alpha)}|_{\mathbb{N}^n \setminus \mathbf{C}_m} : \mathbb{N}^n \setminus \mathbf{C}_m \rightarrow \mathbb{N}^n \setminus \mathbf{C}_m$ is not the identity map for any positive integer m . This implies that $\varepsilon_{l(\alpha)} \neq \sigma_{l(\alpha)}^{-1}\varepsilon_{r(\alpha)}\sigma_{r(\alpha)}$, which contradicts the equalities $\alpha = \sigma_{l(\alpha)}\varepsilon_{l(\alpha)} = \varepsilon_{r(\alpha)}\sigma_{r(\alpha)}$. Hence $\sigma_{l(\alpha)} = \sigma_{r(\alpha)}$ and we put $\sigma_\alpha = \sigma_{r(\alpha)}$.

The uniqueness of the elements σ_α , $\varepsilon_{l(\alpha)}$ and $\varepsilon_{r(\alpha)}$ follows from Lemma 2.5 and equalities (8), because \mathbf{IN}_∞^n is an inverse semigroup. \square

Lemma 3.5 implies the following corollary.

Corollary 3.6. *For any positive integer $n \geq 2$ the subset $H(\mathbb{I}) \cup E(\mathbf{IN}_\infty^n) \subset \mathbf{IN}_\infty^n$ is a set of generators of \mathbf{IN}_∞^n . Moreover, so is any subset $A = A_1 \sqcup A_2 \subset \mathbf{IN}_\infty^n$ such that A_1 generates the group $H(\mathbb{I})$ and A_2 generates the set $E(\mathbf{IN}_\infty^n)$.*

The following theorem describes the natural partial order on the semigroups \mathbf{IN}_∞^n in the terms of Lemma 3.5.

Theorem 3.7. *Let n be any positive integer ≥ 2 . Let α and β be elements of the semigroup \mathbf{IN}_∞^n . Let*

$$\alpha = \sigma_\alpha \varepsilon_{l(\alpha)} = \varepsilon_{r(\alpha)} \sigma_\alpha \quad \text{and} \quad \beta = \sigma_\beta \varepsilon_{l(\beta)} = \varepsilon_{r(\beta)} \sigma_\beta$$

for some elements σ_α and σ_β of the group of units $H(\mathbb{I})$ and idempotents $\varepsilon_{l(\alpha)}$, $\varepsilon_{r(\alpha)}$, $\varepsilon_{l(\beta)}$ and $\varepsilon_{r(\beta)}$ of the semigroup \mathbf{IN}_∞^n . Then $\alpha \preceq \beta$ in \mathbf{IN}_∞^n if and only if $\sigma_\alpha = \sigma_\beta$, $\varepsilon_{l(\alpha)} \preceq \varepsilon_{l(\beta)}$ and $\varepsilon_{r(\alpha)} \preceq \varepsilon_{r(\beta)}$ in $E(\mathbf{IN}_\infty^n)$.

Proof. (\Rightarrow) Suppose that $\alpha \preceq \beta$ in \mathbf{IN}_∞^n . We consider the case $\alpha = \sigma_\alpha \varepsilon_{l(\alpha)}$ and $\beta = \sigma_\beta \varepsilon_{l(\beta)}$. Then there exists an idempotent $\varepsilon \in \mathbf{IN}_\infty^n$ such that $\alpha = \beta \varepsilon$. Then $\sigma_\alpha \varepsilon_{l(\alpha)} = \sigma_\beta \varepsilon_{l(\beta)} \varepsilon$ and hence

$$\varepsilon_{l(\alpha)} = \sigma_\alpha^{-1} \sigma_\beta \varepsilon_{l(\beta)} \varepsilon = \sigma_\alpha^{-1} \sigma_\beta \varepsilon_{l(\beta)} \varepsilon,$$

because $\sigma_\alpha \in H(\mathbb{I})$. Since $\varepsilon_{l(\alpha)}$, $\varepsilon_{l(\beta)}$ and ε are idempotents of \mathbf{IN}_∞^n there exists a positive integer m such that

$$\mathbb{N}^n \setminus \text{dom } \varepsilon_{l(\alpha)} \cup \mathbb{N}^n \setminus \text{dom } \varepsilon_{l(\beta)} \cup \mathbb{N}^n \setminus \text{dom } \varepsilon \subseteq \mathbf{C}_m,$$

and hence by Proposition 3.3, $\mathbb{N}^n \setminus \text{dom } (\varepsilon_{l(\beta)} \varepsilon) \subseteq \mathbf{C}_m$. Then the equality $\varepsilon_{l(\alpha)} = \sigma_\alpha^{-1} \sigma_\beta \varepsilon_{l(\beta)} \varepsilon$ implies that the restriction of $\sigma_\alpha^{-1} \sigma_\beta |_{\mathbb{N}^n \setminus \mathbf{C}_m} : \mathbb{N}^n \setminus \mathbf{C}_m \rightarrow \mathbb{N}^n$ is the identity partial map of $\mathbb{N}^n \setminus \mathbf{C}_m$, and hence by Theorem 3.2 we get that $\sigma_\alpha = \sigma_\beta$. Then

$$\varepsilon_{l(\alpha)} = \sigma_\alpha^{-1} \sigma_\beta \varepsilon_{l(\beta)} \varepsilon = \sigma_\alpha^{-1} \sigma_\alpha \varepsilon_{l(\beta)} \varepsilon = \mathbb{I} \varepsilon_{l(\beta)} \varepsilon = \varepsilon_{l(\beta)} \varepsilon,$$

and hence $\varepsilon_{l(\alpha)} \preceq \varepsilon_{l(\beta)}$ in $E(\mathbf{IN}_\infty^n)$.

The proof in the case $\alpha = \varepsilon_{r(\alpha)} \sigma_\alpha$ and $\beta = \varepsilon_{r(\beta)} \sigma_\beta$ is similar using Lemma 1.4.6 of [13].

The implication (\Leftarrow) is trivial and it follows from Lemma 1.4.6 and Proposition 1.4.7 of [13]. \square

Since $\alpha \mathbf{C}_{\text{mg}} \beta$ in \mathbf{IN}_∞^n if and only if there exists $\gamma \in \mathbf{IN}_\infty^n$ such that $\gamma \preceq \alpha$ and $\gamma \preceq \beta$ (see [13, Section 2.4, p. 62]), Theorem 3.7 implies that $\alpha \mathbf{C}_{\text{mg}} \beta$ in \mathbf{IN}_∞^n if and only if $\sigma_\alpha = \sigma_\beta$. Hence the following theorem holds.

Theorem 3.8. *Let n be any positive integer ≥ 2 . Then the quotient semigroup $\mathbf{IN}_\infty^n / \mathbf{C}_{\text{mg}}$ is isomorphic to the group \mathcal{S}_n and the natural homomorphism $\mathbf{C}_{\text{mg}}^\sharp : \mathbf{IN}_\infty^n \rightarrow \mathbf{IN}_\infty^n / \mathbf{C}_{\text{mg}}$ is defined in the following way: $\alpha \mapsto \sigma_\alpha$.*

The following theorem gives more detail description of Green's relations \mathcal{R} , \mathcal{L} , \mathcal{H} , \mathcal{D} and \mathcal{J} on the semigroup \mathbf{IN}_∞^n .

Theorem 3.9. *Let n be any positive integer ≥ 2 and $\alpha, \beta \in \mathbf{IN}_\infty^n$. Then the following statements hold:*

- (i) $\alpha \mathcal{L} \beta$ if and only if there exists $\sigma \in H(\mathbb{I})$ such that $\alpha = \sigma \beta$;
- (ii) $\alpha \mathcal{R} \beta$ if and only if there exists $\sigma \in H(\mathbb{I})$ such that $\alpha = \beta \sigma$;
- (iii) $\alpha \mathcal{H} \beta$ if and only if there exist $\sigma_1, \sigma_2 \in H(\mathbb{I})$ such that $\alpha = \sigma_1 \beta$ and $\alpha = \beta \sigma_2$;
- (iv) $\alpha \mathcal{D} \beta$ if and only if there exist $\sigma_1, \sigma_2 \in H(\mathbb{I})$ such that $\alpha = \sigma_1 \beta \sigma_2$;
- (v) $\mathcal{D} = \mathcal{J}$ on \mathbf{IN}_∞^n ;
- (vi) every \mathcal{J} -class in \mathbf{IN}_∞^n is finite and consists of incomparable elements with the respect to the natural partial order \preceq on \mathbf{IN}_∞^n .

Proof. (i) (\Rightarrow) Suppose that $\alpha \mathcal{L} \beta$ in \mathbf{IN}_∞^n . Then $\alpha^{-1} \alpha = \beta^{-1} \beta$ and by Lemma 3.4 there exists unique elements σ_{l_α} and σ_{l_β} of the group of units $H(\mathbb{I})$ such that $\sigma_{l_\alpha} \alpha$ and $\sigma_{l_\beta} \alpha$ are idempotents of the semigroup \mathbf{IN}_∞^n . This implies that

$$\sigma_{l_\alpha} \alpha = \alpha^{-1} \alpha = \beta^{-1} \beta = \sigma_{l_\beta} \beta,$$

because \mathbf{IN}_∞^n is an inverse semigroup and

$$\text{ran}(\sigma_{l_\alpha} \alpha) = \text{ran}(\alpha^{-1} \alpha) = \text{ran}(\beta^{-1} \beta) = \text{ran}(\sigma_{l_\beta} \beta).$$

Hence we get that $\alpha = \mathbb{I} \alpha = \sigma_{l_\alpha}^{-1} \sigma_{l_\alpha} \alpha = \sigma_{l_\alpha}^{-1} \sigma_{l_\beta} \beta$, and the element $\sigma = \sigma_{l_\alpha}^{-1} \sigma_{l_\beta} \in H(\mathbb{I})$ is requested.

(\Leftarrow) Suppose that there exists $\sigma \in H(\mathbb{I})$ such that $\alpha = \sigma \beta$. Then

$$\alpha^{-1} \alpha = (\sigma \beta)^{-1} \sigma \beta = \beta^{-1} \sigma^{-1} \sigma \beta = \beta^{-1} \mathbb{I} \beta = \beta^{-1} \beta,$$

and hence $\alpha \mathcal{L} \beta$ in \mathbf{IN}_∞^n .

The proof of statement (ii) is similar to (i).

Statement (iii) follows from (i) and (ii).

(iv) (\Rightarrow) Suppose that $\alpha \mathcal{D} \beta$ in \mathbf{IN}_∞^n . Then there exists $\gamma \in \mathbf{IN}_\infty^n$ such that $\alpha \mathcal{L} \gamma$ and $\gamma \mathcal{R} \beta$ and by statements (i) and (ii) we get that there exist $\sigma_1, \sigma_2 \in H(\mathbb{I})$ such that $\alpha = \sigma_1 \gamma$ and $\gamma = \beta \sigma_2$, and hence $\alpha = \sigma_1 \beta \sigma_2$.

(\Leftarrow) Suppose that there exist $\sigma_1, \sigma_2 \in H(\mathbb{I})$ such that $\alpha = \sigma_1 \beta \sigma_2$. Then $\alpha = \sigma_1 \gamma$ for $\gamma = \beta \sigma_2$, which implies by statements (i) and (ii) that $\alpha \mathcal{L} \gamma$ and $\gamma \mathcal{R} \beta$, and hence $\alpha \mathcal{D} \beta$ in \mathbf{IN}_∞^n .

(v) By Proposition 3.2.17 of [13] it is sufficient to show that every \mathcal{D} -class D in \mathbf{IN}_∞^n contains a minimal element with the respect to the partial natural order \preceq on \mathbf{IN}_∞^n . Fix an arbitrary \mathcal{D} -class D in \mathbf{IN}_∞^n and any $\alpha \in D$. Then by Lemma 3.5 there exists the unique element σ_α of the group of units $H(\mathbb{I})$ and the unique idempotent $\varepsilon_{l(\alpha)}$ of \mathbf{IN}_∞^n such that $\alpha = \sigma_\alpha \varepsilon_{l(\alpha)}$. Then $\sigma_\alpha^{-1} \alpha = \sigma_\alpha^{-1} \sigma_\alpha \varepsilon_{l(\alpha)} = \mathbb{I} \varepsilon_{l(\alpha)} = \varepsilon_{l(\alpha)}$ and $\varepsilon_{l(\alpha)} \in D$ by statement (iv). Also statement (iv) and Theorem 3.2 imply that if an idempotent ε belongs to D then $|\mathbb{N}^n \setminus \text{dom } \varepsilon| = |\mathbb{N}^n \setminus \text{dom } \varepsilon_{l(\alpha)}|$, and hence all idempotents in \mathcal{D} -class D are incomparable with respect to the natural partial order \preceq on \mathbf{IN}_∞^n . Thus, every \mathcal{D} -class D in \mathbf{IN}_∞^n contains a minimal element.

(vi) The first statement follows from (v), (iv) and Theorem 3.2.

Let α and β be elements of a some \mathcal{J} -class j in \mathbf{IN}_∞^n such that $\alpha \preceq \beta$. Then by Lemma 3.5 there exist the unique elements $\sigma_\alpha, \sigma_\beta \in H(\mathbb{I})$ and $\varepsilon_{l(\alpha)}, \varepsilon_{l(\beta)} \in E(\mathbf{IN}_\infty^n)$ such that $\alpha = \sigma_\alpha \varepsilon_{l(\alpha)}$ and $\beta = \sigma_\beta \varepsilon_{l(\beta)}$. By Theorem 3.7, $\sigma_\alpha = \sigma_\beta$ and $\varepsilon_{l(\alpha)} \preceq \varepsilon_{l(\beta)}$ in $E(\mathbf{IN}_\infty^n)$. Then $\sigma_\alpha^{-1} \alpha = \sigma_\alpha^{-1} \sigma_\alpha \varepsilon_{l(\alpha)} = \varepsilon_{l(\alpha)}$ and $\sigma_\alpha^{-1} \beta = \sigma_\alpha^{-1} \sigma_\beta \varepsilon_{l(\beta)} = \varepsilon_{l(\beta)}$ are \mathcal{J} -equivalent idempotents in \mathbf{IN}_∞^n . By the proof of statement (v) we get that $\varepsilon_{l(\alpha)} = \varepsilon_{l(\beta)}$, and hence by Lemma 3.5, $\alpha = \beta$. \square

4. THE STRUCTURE THEOREM FOR THE SEMIGROUP \mathbf{IN}_∞^n

Recall [14], an inverse semigroup S is said to be *F-inverse* if every element of S has a unique maximal element above it in the natural partial order \preceq , i.e. every \mathfrak{C}_{mg} -class has a maximum element. It is obvious that every *F-inverse* semigroup contains a unit. An inverse semigroup S is *E-unitary* if, whenever e is an idempotent in S , $s \in S$ and $e \preceq s$, then s is an idempotent in S [16].

Lemma 3.5 implies the following two corollaries.

Corollary 4.1. \mathbf{IN}_∞^n is an *F-inverse* semigroup for any positive integer $n \geq 2$.

Corollary 4.2. \mathbf{IN}_∞^n is an *E-unitary* inverse semigroup for any positive integer $n \geq 2$.

For any element s of an inverse semigroup S we denote $\downarrow s = \{x \in S : x \preceq s\}$, where \preceq is the natural partial order on S .

Let S be any *F-inverse* semigroup. Then for any $s \in S$ we denote by e_s the idempotent $ss^{-1} \in S$ and by t_s the maximum element in the \mathfrak{C}_{mg} -class of $s \in S$, and by T_S the set $\{t_s : s \in S\}$. We note that T is the set of maximal elements of S , and that T_S need not be closed under multiplication [14]. For each $x \in S$, as in [14] let e_x denote the idempotent xx^{-1} .

The structure of *F-inverse* semigroups is described in [14], and later we need the following two statements for the description of the structure of the semigroup \mathbf{IN}_∞^n .

Lemma 4.3 ([14, Lemma 3]). *Let S be an *F-inverse* semigroup and 1_S denote the identity element of S . Then*

- (i) $E(S)$ is a semilattice with 1_S for an identity element.
- (ii) On T_S define the following multiplication:

$$u * v = t_{uv}, \quad \text{for } u, v \in T_S.$$

Then $(T_S, *)$ is a group with 1_S for an identity element, and each $t \in T_S$ has t^{-1} as its group inverse in $(T_S, *)$.

(iii) For each $t \in T_S$ the map $\mathfrak{F}_t: E(S) \rightarrow \downarrow e_t$, defined by

$$(f)\mathfrak{F}_t = tft^{-1}, \quad f \in E(S),$$

is a homomorphism onto $\downarrow e_t$, and moreover \mathfrak{F}_{1_S} is the identity map on $E(S)$.

(iv) For each $t \in T_S$ $(1_S)\mathfrak{F}_t = e_t$ and $(e_t)\mathfrak{F}_{t^{-1}} = e_{t^{-1}}$.

(v) For each $u, v \in T_S$

$$((1_S)\mathfrak{F}_u)\mathfrak{F}_v \cdot (f)\mathfrak{F}_{u*v} = ((f)\mathfrak{F}_u)\mathfrak{F}_v, \quad \text{for all } f \in E(S);$$

(vi) If $u, v \in T_S$ then

$$f \cdot (g)\mathfrak{F}_u \leq e_{u*v},$$

for all idempotents $f \leq e_u$ and $g \leq e_v$ of S .

Theorem 4.4 ([14, Theorem 3]). Let S be an F -inverse semigroup and let $\mathcal{S} = \bigcup_{t \in T_S} (\downarrow e_t \times \{t\})$. Define the multiplication \circ on \mathcal{S} as follows. If $u, v \in T_S$ then for idempotents $f \leq e_u$ and $g \leq e_v$ put

$$(9) \quad (f, u) \circ (g, v) = (f \cdot (g)\mathfrak{F}_u, u * v).$$

Then \circ is a well-defined multiplication on \mathcal{S} and (\mathcal{S}, \circ) is isomorphic to S under the map $\mathfrak{H}: S \rightarrow \mathcal{S}$, $s \mapsto (ss^{-1}, t_s)$.

Let A and B be semigroups, $\text{End}(B)$ be the semigroup of endomorphisms of B and the following homomorphism $\mathfrak{h}: A \rightarrow \text{End}(B): b \mapsto \mathfrak{h}_b$ is defined. Then the set $A \times B$ with the semigroup operation

$$(a_1, b_1) \cdot (a_2, b_2) = (a_1 a_2, (b_1)\mathfrak{h}_{a_2} b_2), \quad a_1, a_2 \in A, b_1, b_2 \in B,$$

is called the *semidirect product* of the semigroup A by B with the respect to the homomorphism \mathfrak{h} and it is denoted by $A \rtimes_{\mathfrak{h}} B$ [13]. In this case we say that the right action of the semigroup A is defined on the semigroup of endomorphisms of B . We remark that a semidirect product of two inverse semigroups is not need an inverse semigroup (see [13, Section 5.3]).

Lemma 4.5. The map $\mathfrak{h}: H(\mathbb{I}) \rightarrow \text{End}(E(\mathbb{I}\mathbb{N}_{\infty}^n))$, $\sigma \mapsto \mathfrak{h}_{\sigma}$, where $(\alpha)\mathfrak{h}_{\sigma} = \sigma^{-1}\alpha\sigma$ is the automorphism of the semilattice $E(\mathbb{I}\mathbb{N}_{\infty}^n)$, is a homomorphism, and moreover $\mathfrak{h}_{\mathbb{I}}$ is the identity automorphism of $E(\mathbb{I}\mathbb{N}_{\infty}^n)$.

Proof. For any $\sigma \in H(\mathbb{I})$, $\varepsilon, \iota \in E(\mathbb{I}\mathbb{N}_{\infty}^n)$ we have that

$$(\varepsilon\iota)\mathfrak{h}_{\sigma} = \sigma^{-1}\varepsilon\iota\sigma = \sigma^{-1}\varepsilon\sigma\sigma^{-1}\iota\sigma = (\varepsilon)\mathfrak{h}_{\sigma}(\iota)\mathfrak{h}_{\sigma},$$

and hence \mathfrak{h}_{σ} is an endomorphism of the semilattice $E(\mathbb{I}\mathbb{N}_{\infty}^n)$. Also, since for any $\sigma \in H(\mathbb{I})$ and $\varepsilon \in E(\mathbb{I}\mathbb{N}_{\infty}^n)$ the element $\sigma\varepsilon\sigma^{-1}$ is an idempotent of the semigroup $\mathbb{I}\mathbb{N}_{\infty}^n$ and $(\sigma\varepsilon\sigma^{-1})\mathfrak{h}_{\sigma} = \sigma^{-1}(\sigma\varepsilon\sigma^{-1})\sigma = \varepsilon$, the homomorphism \mathfrak{h}_{σ} is a surjective map. It is obvious that $\mathfrak{h}_{\mathbb{I}}$ is the identity automorphism of the semilattice $E(\mathbb{I}\mathbb{N}_{\infty}^n)$.

Suppose that $(\varepsilon)\mathfrak{h}_{\sigma} = (\iota)\mathfrak{h}_{\sigma}$ for some $\gamma \in H(\mathbb{I})$ and $\varepsilon, \iota \in E(\mathbb{I}\mathbb{N}_{\infty}^n)$. Since $H(\mathbb{I})$ is the group of units of the semigroup $\mathbb{I}\mathbb{N}_{\infty}^n$, the following equalities

$$\sigma^{-1}\varepsilon\sigma = (\varepsilon)\mathfrak{h}_{\sigma} = (\iota)\mathfrak{h}_{\sigma} = \sigma^{-1}\iota\sigma$$

imply that

$$\varepsilon = 1\varepsilon 1 = \sigma\sigma^{-1}\varepsilon\sigma\sigma^{-1} = \sigma\sigma^{-1}\iota\sigma\sigma^{-1} = 1\iota 1 = \iota,$$

and hence \mathfrak{h}_{σ} is an automorphism of the semilattice $E(\mathbb{I}\mathbb{N}_{\infty}^n)$.

Fix arbitrary $\sigma_1, \sigma_2 \in H(\mathbb{I})$. Then for any idempotent $\varepsilon \in \mathbb{I}\mathbb{N}_{\infty}^n$ we have that

$$(\varepsilon)\mathfrak{h}_{\sigma_1\sigma_2} = (\sigma_1\sigma_2)^{-1}\varepsilon\sigma_1\sigma_2 = \sigma_2^{-1}\sigma_1^{-1}\varepsilon\sigma_1\sigma_2 = \sigma_2^{-1}(\varepsilon)\mathfrak{h}_{\sigma_1}\sigma_2 = ((\varepsilon)\mathfrak{h}_{\sigma_1})\mathfrak{h}_{\sigma_2} = (\varepsilon)(\mathfrak{h}_{\sigma_1} \cdot \mathfrak{h}_{\sigma_2}),$$

and hence so defined map $\mathfrak{h}: H(\mathbb{I}) \rightarrow \text{End}(E(\mathbb{I}\mathbb{N}_{\infty}^n))$ is a homomorphism. \square

The following theorem describes the structure of the semigroup $\mathbb{I}\mathbb{N}_{\infty}^n$.

Theorem 4.6. *Let n be any positive integer ≥ 2 . Then the semigroup \mathbf{IN}_∞^n is isomorphic to the semidirect product $\mathcal{S}_n \rtimes_{\mathfrak{h}} (\mathcal{P}_\infty(\mathbb{N}^n), \cup)$ of free semilattice with the unit $(\mathcal{P}_\infty(\mathbb{N}^n), \cup)$ by the symmetric group \mathcal{S}_n .*

Proof. Since by Theorem 3.2 the group of units $H(\mathbb{I})$ of the semigroup \mathbf{IN}_∞^n is isomorphic to the symmetric group \mathcal{S}_n , it is sufficient to show that the semigroup \mathbf{IN}_∞^n is isomorphic to the semidirect product $H(\mathbb{I}) \rtimes_{\mathfrak{h}} E(\mathbf{IN}_\infty^n)$ the semilattice $E(\mathbf{IN}_\infty^n)$ by the group of units $H(\mathbb{I})$ of \mathbf{IN}_∞^n with the respect to the homomorphism $\mathfrak{h}: H(\mathbb{I}) \rightarrow \text{End}(E(\mathbf{IN}_\infty^n))$, $\sigma \mapsto \mathfrak{h}_\sigma$, where $(\varepsilon)\mathfrak{h}_\sigma = \sigma^{-1}\varepsilon\sigma$, $\varepsilon \in E(\mathbf{IN}_\infty^n)$.

We define a map $\mathfrak{T}: \mathbf{IN}_\infty^n \rightarrow H(\mathbb{I}) \rtimes_{\mathfrak{h}} E(\mathbf{IN}_\infty^n)$ in the following way

$$(\alpha)\mathfrak{T} = (\sigma_\alpha, \alpha^{-1}\alpha),$$

where the element σ_α of the group of units $H(\mathbb{I})$ of \mathbf{IN}_∞^n is defined by formula $\alpha \mapsto \sigma_\alpha$ in Theorem 3.8. By Lemma 3.5 and Theorem 3.8 the map $\mathfrak{T}: \mathbf{IN}_\infty^n \rightarrow H(\mathbb{I}) \rtimes_{\mathfrak{h}} E(\mathbf{IN}_\infty^n)$ is well defined and by Theorem 3.9(iv) it is surjective. Suppose there exist $\alpha, \beta \in \mathbf{IN}_\infty^n$ such that $(\alpha)\mathfrak{T} = (\beta)\mathfrak{T}$. Then $(\sigma_\alpha, \alpha^{-1}\alpha) = (\sigma_\beta, \beta^{-1}\beta)$ and by Theorem 3.7 we get that

$$\alpha = \sigma_\alpha \alpha^{-1} \alpha = \sigma_\beta \beta^{-1} \beta = \beta,$$

and hence the map $\mathfrak{T}: \mathbf{IN}_\infty^n \rightarrow H(\mathbb{I}) \rtimes_{\mathfrak{h}} E(\mathbf{IN}_\infty^n)$ is injective.

Fix arbitrary $\alpha, \beta \in \mathbf{IN}_\infty^n$. Then Lemma 3.5 and Theorem 3.8 imply that $\alpha\beta\mathfrak{C}_{\text{mg}}\gamma_\alpha\gamma_\beta$, and since $\sigma_\alpha, \sigma_\beta \in H(\mathbb{I})$, we get that $\sigma_\alpha\sigma_\beta = \sigma_{\alpha\beta}$. This implies that

$$(\alpha)\mathfrak{T}(\beta)\mathfrak{T} = (\sigma_\alpha, \alpha^{-1}\alpha) (\sigma_\beta, \beta^{-1}\beta) = (\sigma_\alpha\sigma_\beta, \sigma_\beta^{-1}\alpha^{-1}\alpha\sigma_\beta\beta^{-1}\beta) = (\sigma_{\alpha\beta}, \sigma_\beta^{-1}\alpha^{-1}\alpha\sigma_\beta\beta^{-1}\beta).$$

By Lemma 4.5 the map $\mathfrak{h}_\sigma: E(\mathbf{IN}_\infty^n) \rightarrow E(\mathbf{IN}_\infty^n): \alpha \mapsto \sigma^{-1}\alpha\sigma$ is an automorphism of the semilattice $E(\mathbf{IN}_\infty^n)$, and hence we get that $\sigma_\beta^{-1}\alpha^{-1}\alpha\sigma_\beta$ is an idempotent of the semigroup \mathbf{IN}_∞^n . By Lemma 3.5 and Theorem 3.7 for any $\alpha \in \mathbf{IN}_\infty^n$ there exists the unique element σ_α of the group of units $H(\mathbb{I})$ such that $\alpha \preceq \sigma_\alpha$. Lemma 1.4.6 from [13] implies that $\beta = \sigma_\beta\beta^{-1}\beta$, and hence

$$\begin{aligned} \sigma_\beta^{-1}\alpha^{-1}\alpha\sigma_\beta\beta^{-1}\beta &= (\sigma_\beta^{-1}\alpha^{-1}\alpha\sigma_\beta) (\beta^{-1}\beta) (\beta^{-1}\beta) = \\ &= (\beta^{-1}\beta\gamma_\beta^{-1}) (\alpha^{-1}\alpha) (\sigma_\beta\beta^{-1}\beta) = \\ &= (\sigma_\beta\beta^{-1}\beta)^{-1} (\alpha^{-1}\alpha) (\sigma_\beta\beta^{-1}\beta) = \\ &= \beta^{-1} (\alpha^{-1}\alpha) \beta = \\ &= (\beta^{-1}\alpha^{-1}) (\alpha\beta) = \\ &= (\alpha\beta)^{-1} (\alpha\beta). \end{aligned}$$

Therefore, we have that

$$(\alpha\beta)\mathfrak{T} = (\sigma_{\alpha\beta}, (\alpha\beta)^{-1}\alpha\beta) = (\alpha)\mathfrak{T}(\beta)\mathfrak{T},$$

and hence the map $\mathfrak{T}: \mathbf{IN}_\infty^n \rightarrow H(\mathbb{I}) \rtimes_{\mathfrak{h}} E(\mathbf{IN}_\infty^n)$ is a homomorphism, which completes the proof of the theorem. \square

Remark 4.7. The monoid \mathbf{IN}_∞ of all partial cofinite isometries of the set of positive integers \mathbb{N} is a submonoid of the monoid $\mathcal{S}_\infty^{\nearrow}(\mathbb{N})$ of cofinite, monotone, non-decreasing, injective partial transformations of \mathbb{N} [12]. The structure of the semigroup is described in [8]. Also, Lemma 3.5 implies that for any positive integer $n \geq 2$ the semigroup \mathbf{IN}_∞^n is an inverse submonoid of the monoid $\mathcal{PO}_\infty(\mathbb{N}_{\leq}^n)$ of monotone injective partial selfmaps of \mathbb{N}_{\leq}^n with the order product having cofinite domain and image [5, 6, 7].

ACKNOWLEDGEMENTS

The authors acknowledge Alex Ravsky for his comments and suggestions.

REFERENCES

- [1] O. Bezushchak, *On growth of the inverse semigroup of partially defined co-finite automorphisms of integers*, Algebra Discrete Math. (2004), no. 2, 45–55.
- [2] O. Bezushchak, *Green's relations of the inverse semigroup of partially defined co-finite isometries of discrete line*, Visn., Ser. Fiz.-Mat. Nauky, Kyiv. Univ. Im. Tarasa Shevchenka (2008), no. 1, 12–16.
- [3] A. H. Clifford and G. B. Preston, *The Algebraic Theory of Semigroups*, Vol. I., Amer. Math. Soc. Surveys 7, Providence, R.I., 1961; Vol. II., Amer. Math. Soc. Surveys 7, Providence, R.I., 1967.
- [4] J. A. Green, *On the structure of semigroups*, Ann. Math. (2) **54** (1951), no. 1, 163–172.
- [5] O. Gutik and O. Krokmalna, *The monoid of monotone injective partial selfmaps of the poset (\mathbb{N}^n, \leq) with cofinite domains and images*, Preprint.
- [6] O. Gutik and I. Pozdniakova, *On the monoid of monotone injective partial selfmaps of N_{\leq}^2 with co-finite domains and images*, Visn. L'viv. Univ., Ser. Mekh.-Mat. **81** (2016), 100–116.
- [7] O. Gutik and I. Pozdniakova, *On the monoid of monotone injective partial selfmaps of N_{\leq}^2 with co-finite domains and images, II*, Visn. L'viv. Univ., Ser. Mekh.-Mat. **82** (2016), 109–127.
- [8] O. Gutik and D. Repovš, *Topological monoids of monotone, injective partial selfmaps of \mathbb{N} having cofinite domain and image*, Stud. Sci. Math. Hungar. **48** (2011), no. 3, 342–353.
- [9] Gutik O., Repovš D. On monoids of injective partial cofinite selfmaps, Math. Slovaca **65** (2015), no. 5, 981–992.
- [10] O. Gutik and A. Savchuk, *On the semigroup \mathbf{ID}_{∞}* , Visn. Lviv. Univ., Ser. Mekh.-Mat. **83** (2017), 5–19 (in Ukrainian).
- [11] O. Gutik and A. Savchuk, *The semigroup of partial co-finite isometries of positive integers*, Bukovyn. Mat. Zh. **6** (2018), no. 1–2, 42–51 (in Ukrainian).
- [12] O. Gutik and A. Savchuk, *On inverse submonoids of the monoid of almost monotone injective co-finite partial selfmaps of positive integers*, Preprint (arXiv:1904.11802).
- [13] M. Lawson, *Inverse Semigroups. The Theory of Partial Symmetries*, Singapore: World Scientific, 1998.
- [14] R. McFadden and L. O'Carroll, *F-inverse semigroups*, Proc. Lond. Math. Soc., III. Ser. **22** (1971), no. 4, 652–666.
- [15] M. Petrich, *Inverse Semigroups*, John Wiley & Sons, New York, 1984.
- [16] T. Saitô, *Proper ordered inverse semigroups*, Pacif. J. Math. **15** (1965), no. 2, 649–666.
- [17] V. V. Wagner, *Generalized groups*, Dokl. Akad. Nauk SSSR **84** (1952), 1119–1122 (in Russian).

FACULTY OF MATHEMATICS, IVAN FRANKO NATIONAL UNIVERSITY OF LVIV, UNIVERSYTETSKA 1, LVIV, 79000, UKRAINE

E-mail address: oleg.gutik@lnu.edu.ua, ovgutik@yahoo.com, asavchuk3333@gmail.com