A RIGID THEOREM FOR DEFORMED HERMITIAN-YANG-MILLS EQUATION

XIAOLI HAN AND XISHEN JIN

ABSTRACT. In this paper, we study the deformed Hermitian-Yang-Mills equation on compact Kähler manifold with non-negative orthogonal bisectional curvature. We prove that the curvatures of deformed Hermitian-Yang-Mills metrics are parallel with respect to the background metric if there exists a positive constant C such that $-\frac{1}{C}\omega < \sqrt{-1}F < C\omega$. We also study the self-shrinker over \mathbb{C}^n to the corresponding parabolic flow. We prove that the self-shrinker over \mathbb{C}^n is a quadratic polynomial function. We also show the similar rigid theorem for the J-equations and the self-shrinkers over \mathbb{C}^n to J-flow.

1. INTRODUCTION

Let (X, ω) be a connected compact Kähler manifold of complex dimension n and L be a holomorphic line bundle over X. Given a metric h on L, we define the complex function

$$\zeta := \frac{(\omega - F)^n}{\omega^n},$$

where F is the curvature of the Chern connection with respect to the metric h. It is easy to see that the average of this function is a fixed complex number

$$Z_{L,[\omega]} := \int_X \zeta \frac{\omega^n}{n!}$$

dependent only the cohomology classes on $c_1(L)$ and $[\omega] \in H^{1,1}(X, \mathbb{R})$. Let θ denote the argument of ζ and $\hat{\theta}$ the argument of $Z_{L,[\omega]}$. Here the branch cut is specified.

Definition 1.1. A Hermitian metric h on L is said to be deformed Hermitian-Yang-Mills metric if it satisfies

(1.1)
$$\theta = \hat{\theta}.$$

or equivalently

(1.2)
$$\operatorname{Im}(\omega - F)^n = \tan(\hat{\theta}) \operatorname{Re}(\omega - F)^n$$

The deformed Hermitian-Yang-Mills equation was discovered by Marino et all [21] as the requirement for a D-brane on the B-model of mirror symmetry to be supersymmetric. Recently, it has been studied by Collins-Jacob-Yau [4], Collins-Xie-Yau [6], Jacob-Yau [18] and some other people. According to superstring theory, the spacetime of the universe is constrained to be a product of a compact Calabi-Yau threefold and a four-dimensional Lorentzian manifold. A 'duality' relates the geometry of one Calabi-Yau manifold to another 'mirror' Calabi-Yau manifold. From a differential geometry viewdpoint this might be thought of a relationship between the existence of 'nice' metrics on the line bundle over one Calabi-Yau manifold

and the special Lagrangian submanifolds in another Calabi-Yau manifold. In [19], Leung-Yau-Zaslow showed that the deformed Hermitian-Yang-Mills equation on a line bundle corresponds to the special Lagrangian equation in the mirror.

In [6], Collins-Xie-Yau addressed a Chern number inequality on 3-dimensional Kähler manifold (X, Ω, L) admitting a deformed Hermitian-Yang-Mills metric by considering the position and winding angle of

$$\gamma(t) = -\int_X e^{-t\sqrt{-1}\omega} ch(L).$$

As consequences of this Chern number inequality, they obtained some algebraic stability conditions to the existence of deformed Hermitian-Yang-Mills metrics.

In [4], Collins-Jacob-Yau showed the existence of deformed Hermitian-Yang-Mills metrics under the assumption of the existence of a C-subsolution with supercritical phase using the method of continuity. They also conjectured some stability-type cohomological obstructions to the existence of deformed Hermitian-Yang-Mills metrics. In particular, they got a Liouville theorem for the deformed Hermitian-Yang-Mills metrics with bounded complex Hessian on \mathbb{C}^n under the assumption of supercritical phase, i.e. $\hat{\theta} \in (\frac{n-2}{2}\pi, \frac{n}{2}\pi)$, by applying the Evans-Krylov theory.

In [25], Pingali viewed the deformed Hermitian-Yang-Mills equation as a generalized type of Monge-Ampère equation PDE with "non-constant coefficients" and got some existence results to the deformed Hermitian-Yang-Mills equation for some ranges of the phase angle assuming the existence of a subsolution. In particular, Pingali generalised the work of [5] on toric varieties to the deformed Hermitian-Yang-Mills equation under some algebraic geometric stability condition and addressed a conjecture of Collins-Jacob-Yau [4].

In [18], Jacob-Yau provided a necessary and sufficient criterion for the existence of dHYM metrics in the case that X is a Kähler surface. They also introduced a line bundle version of the Lagrangian mean curvature flow and proved the convergence of the flow when L is sufficiently ample and X has non-negative orthogonal bisectional curvature. In particular, they required that the initial data satisfies the hypercritical phase condition so that the Evans-Krylov theory works. In [17], Han-Yamamoto established a ε -regularity theorem for this flow in the semi-flat case. We also remark that the criterion of Jacbo-Yau on Kähler surfaces was generalized to Kähler manifold of complex dimension 3 by viewing the deformed Hermitian-Yang-Mills equation as a generalized type of Monge-Amère equation in [24].

In the first part of this paper, we mainly consider the rigidity of deformed Hermitian-Yang-Mills metrics on (X, ω) with nonnegative orthogonal bisectional curvature.

Classically, rigid theorems in Kähler geometry aim to show that the Kähler manifolds satisfying some natural curvature conditions are the known examples. A famed example is the complex projective space $\mathbb{C}P^N$ equipped with the standard Fubini-Study metric g_{FS} whose bisectional curvature is positive. The Frankel conjecture states that every compact Kähler manifold with positive bisectional curvature is biholomorphic to $\mathbb{C}P^N$. This was established independently by Mori [23] and Siu-Yau [26]. An interesting new proof of the Frankel conjecture was given by Chen-Tian [7] with the help of Kähler-Ricci flow. In their sense, positivity of bisectional curvature is preserved under the Kähler-Ricci flow and the flow converges to a Kähler-Einstein metric exponentially. Then a classical result of Berger [1] and Golberg-Kobayashi [11] on rigidity of closed constant scalar curvature Kähler manifold with positive bisectional curvature implies the Frankel conjecture. The results of Berger [1] and Golberg-Kobayashi [11] was generalized to compact Kähler manifolds with nonnegative bisectional curvature in [14] by Howard-Smyth-Wu. In details, Howard-Smyth-Wu proved that a compact csck Kähler manifold with nonnegative bisectional curvature is automatically a Kähler-Einstein manifold in [14]. We also remark that the csck condition above was generalized to some fully nonlinear conditions on the Ricci tensor in [12] and the key technology is that the Ricci tensor on Kähler manifold satisfies the Codazzi condition.

Definition 1.2. A Kähler metric ω is said to have nonnegative orthogonal bisectional curvature, if for any orthonormal tangent frame $\{e_1, \dots, e_n\}$ at any $x \in X$, the curvature tensor of ω satisfies that

$$R_{i\bar{i}j\bar{j}} = R(e_i, \bar{e}_i, e_j, \bar{e}_j) \ge 0$$
, for all $1 \le i, j \le n$ and $i \ne j$.

We remark that nonnegativity of the orthogonal bisectional curvature is weaker than nonnegativity of the bisectional curvature. In fact, the former condition is satisfied by not only complex projective spaces and the Hermitian symmetric spaces, but also some compact Kähler manifolds of dimension ≥ 2 whose holomorphic sectional curvature is strictly negative somewhere. We refer the readers to the work of Gu-Zhang [13], Mok [22] and Siu-Yau [26].

The main result of this paper is the following theorem.

Theorem 1.1. Let (X, ω) be a Kähler manifold with nonnegative orthogonal bisectional curvature and (L, h) be a holomorphic line bundle on X. Suppose h solves (1.1). If there exists a positive constant C such that $-\frac{1}{C}\omega < \sqrt{-1}F < C\omega$, i.e. the eigenvalues λ_i is in $(-\frac{1}{C}, C)$, then $\lambda_i (i = 1, \dots, n)$ are all constants. In particular, if the orthogonal bisectional curvature is strictly positive at some point $p \in X$, then $\sqrt{-1}F = c\omega$ for some constant c.

The deformed Hermitian-Yang-Mills equation can be seen as a complex analogy to the special Lagrangian equation. In [31, 32], Yuan proved that viewed as a graph over \mathbb{R}^n , the entire special Lagrangian submanifold in \mathbb{C}^n must be a plane under some assumptions of the Hessian of the potential function. Here our assumptions can be interpreted as the complex Hessian of the Hermitian metric h on line bundle L. The method that we adopt to establish the main theorem is the maximal principle by choosing a suitable auxiliary function.

Another interesting equation in Kähler geometry and mirror symmetry is the J-equation. Given Kähler metrics ω and χ on X, the J-equation is defined as

$$\operatorname{Tr}_{\omega_{\varphi}}\chi = c$$

where $\omega_{\varphi} = \omega + \sqrt{-1}\partial\bar{\partial}\varphi$ is the solution to be found. The J-equation can also be rewritten in the following version

$$\sum_{i=1}^{n} \frac{1}{\lambda_i} = c$$

where $\{\lambda_1, \dots, \lambda_n\}$ are the eigenvalues of $\chi^{-1}\omega_{\varphi}$ as an endomorphism on $T^{1,0}X$. The J-equation was introduced by Donaldson [8] from the viewpoint of moment maps. In [27], Song-Weinkove gave a necessary and sufficient condition for the existence of the solution of J-equation. In [5], Collins-Székelyhidi considered the solvability of J-equation on toric varieties by the J-flow. In [3], Chen gave a new numerical condition for the solvability of J-equation.

Observed by Collins-Jacob-Yau [4], for any fixed *n*-tuples $(\lambda_1, \dots, \lambda_n)$ with all $\lambda_i > 0$,

$$\lim_{k \to \infty} \sum_{i=1}^{n} k(\frac{\pi}{2} - \arctan(k\lambda_i)) = \sum_{i=1}^{n} \frac{1}{\lambda_i}$$

i.e. the J-equation can be exactly seen as the limit of the deformed Hermitian-Yang-Mills equation. So we hope that the rigid theorem above should also hold for J-equation. In fact, we obtain the following theorem.

Theorem 1.2. If (X, χ) is a compact Kähler manifold with non-negative orthogonal bisectional curvature and ω is a Kähler metric satisfying the J-equation, then the eigenvalues of $K = \chi^{-1}\omega$ are constants. In particular, if we also assume the orthogonal bisectional curvature is strictly positive at some point, then $\omega = c\chi$ for some constant c.

Coming back to the corresponding parabolic flows to the deformed Hermitian-Yang-Mills equation and J-equation, we want to consider the self- shrinkers of these flows. In fact, we prove the following rigid theorems for the self-shrinkers on \mathbb{C}^n .

Theorem 1.3. If u is a smooth function from \mathbb{C}^n to \mathbb{R} and satisfies the following equation

(1.3)
$$\sum_{i=1}^{n} \arctan \lambda_i(z) - \theta_0 = \frac{1}{2} (\langle z, \partial u \rangle + \langle \bar{z}, \bar{\partial} u \rangle) - u(z),$$

where λ_i is the eigenvalue of $\{u_{i\bar{j}}\}$, then u is quadratic.

Any function u satisfying (1.3) leads to an entire self-similar solution

$$v(z,t) = -tu(\frac{z}{\sqrt{-t}})$$

to the line-bundle mean curvature flow defined by Jacob-Yau in [18]

$$\frac{\partial v}{\partial t} = \sum_{i=1}^{n} \arctan \lambda_i - \theta_0.$$

This type of rigid results for self-shrinkers to parabolic flow have been studied in [2, 9, 10, 16, 29, 30]. In particular, Chau-Chen-Yuan proved the rigid theorem for entire smooth Lagrangian self-shrinkers over \mathbb{R}^n . Theorem 1.3 can be seen as a complex analogy to their results.

We also consider the self-shrinkers to the J-flow

$$\frac{\partial u}{\partial t} = c - \mathrm{Tr}_{\omega_{\varphi}} \chi$$

We can prove the following rigid result for entire solutions to self-shrinkers with respect to the J-flow.

Theorem 1.4. $u : \mathbb{C}^n \to \mathbb{R}$ is a smooth strictly pluri-subharmonic function satisfying the following equation

(1.4)
$$c - \sum_{i=1}^{n} \frac{1}{\lambda_i(z)} = \frac{1}{2} (\langle z, \partial u \rangle + \langle \bar{z}, \bar{\partial}u \rangle) - u(z)$$

where $\lambda_i(z)$ is the eigenvalue of $\{u_{i\bar{j}}\}$. If the complex hessian of u satisfies

$$\{u_{i\bar{j}}(z)\} \geq \frac{\sqrt{2n-1+\delta}}{|z|}I$$

for any $\delta > 0$ as $|z| \to \infty$, then u is quadratic.

Similarly, any function u in the theorem above leads to an entire self-similar solution

$$v(z,t) = -tu(\frac{z}{\sqrt{-t}})$$

to the J-flow

$$\frac{\partial v}{\partial t} = c - \sum_{i=1}^{n} \frac{1}{\lambda_i}.$$

Remark 1.1. In [15], Huang-Ou-Wang considered the similar rigid theorem for real version J-type equation(Theorem 1.1 in [15]). And they did not need the extra growing condition of the Hessian. The key observation in [15] is that the self-shrinker equation has a good expression after the Legendre transformation. However, there is no such complex analog Legendre transformation. This difficulty was also discussed in [28].

We will organize this paper as following. In section 2, we give some background knowledge to the deformed Hermitian-Yang-Mills equations and prove Theorem 1.1 and Theorem 1.2. In section 3, we prove the rest results for self-shrinkers.

Acknowledgements: Both authors are grateful to Prof. Jiayu Li and Prof. Hikaru Yamamoto for helpful discussions.

2. Rigid Theorem for Deformed Hermitian-Yang-Mills Equation and J-equation

2.1. **Preliminaries.** Let (X, ω) be a compact Kähler manifold of complex dimension n and (L, h) be a holomorphic line bundle over X. F is the Chern curvature with respect to the Hermitian metric h on L. In local coordinates, we write

$$\omega = \frac{\sqrt{-1}}{2} g_{i\bar{j}} dz^i \wedge d\bar{z}^j$$

and

$$F = \frac{1}{2} F_{i\bar{j}} dz^i \wedge d\bar{z}^j = -\frac{1}{2} \partial_i \partial_{\bar{j}} \log(h) dz^i \wedge d\bar{z}^j.$$

Since ω is Kähler, we have the so-called second Bianchi equality

$$\begin{aligned} F_{i\bar{j},k} &= -\partial_k \partial_{\bar{j}} \partial_i \log(h) + \Gamma^s_{ik} F_{s\bar{j}} \\ &= -\partial_i \partial_{\bar{j}} \partial_k \log(h) + \Gamma^s_{ki} F_{s\bar{j}} = F_{k\bar{j},i}, \end{aligned}$$

i.e. $F_{i\bar{j},k} = F_{k\bar{j},i}$.

Using F, we introduce a Hermitian(usually not Kähler) metric

$$\eta = \frac{\sqrt{-1}}{2} \eta_{\bar{k}j} dz^j \wedge d\bar{z}^k$$

and an endomorphism K of $T^{(1,0)}(X)$ that are defined by

$$\eta_{\bar{k}j} = g_{\bar{k}j} + F_{\bar{k}l} g^{l\bar{m}} F_{\bar{m}j},$$

and

$$K := \omega^{-1}F = g^{i\bar{j}}F_{k\bar{j}}dz^i \otimes \frac{\partial}{\partial z^k}$$

Then the complex-valued (n, n)-form $(\omega - F)^n$ can be locally written as

$$(\omega - F)^n = n! \det(g_{\bar{k}j} + \sqrt{-1}F_{\bar{k}j})(\frac{i}{2})^n dz^1 \wedge d\bar{z}^1 \wedge \dots \wedge dz^n \wedge d\bar{z}^n$$

=
$$\det(g^{j\bar{k}}) \det(g_{\bar{k}j} + \sqrt{-1}F_{\bar{k}j})\omega^n$$

=
$$\det(I + \sqrt{-1}K)\omega^n.$$

Thus the complex function ζ can be expressed as following

(2.1)
$$\zeta = \det(I + \sqrt{-1}K)$$

and the argument of ζ is

(2.2)
$$\theta = -i \log \frac{\det \left(I + \sqrt{-1}K\right)}{\sqrt{\det(I + K^2)}},$$

where the endomorphism $I + K^2$ can be expressed locally as

$$I + K^2 = g^{p\bar{q}} \eta_{\bar{q}l} \frac{\partial}{\partial z^p} \otimes dz^l.$$

We choose the normal coordinates around some point $p \in X$ so that the endomorphism K is diagonal at p with eigenvalues $\{\lambda_j\}$ for $1 \leq j \leq n$, then the function θ can be expressed as

$$\theta := \arctan \lambda_1 + \dots + \arctan \lambda_n.$$

Here θ takes value in $\left(-\frac{n}{2}\pi, \frac{n}{2}\pi\right)$. Then the equation (1.1) is equivalent to

(2.3)
$$\theta = \hat{\theta}(mod \ 2\pi).$$

Hence the deformed Hermitian-Yang-Mills equation (1.1) can be written as

(2.4)
$$\sum_{i=1}^{n} \arctan \lambda_i = \theta.$$

As described in [6], we also remark the constant θ in (2.4) can be obtained by considering the "winding angle" of

$$\gamma(t) = \int_X e^{-t\sqrt{-1}\omega} Ch(L)$$

as t runs from $+\infty$ to 1.

Taking the derivative of (2.2) , we get (cf. Lemma 3.3 in [18]) the first variation of θ as following

(2.5)
$$\delta\theta = Tr((I+K^2)^{-1}\delta K).$$

Thus if we take the derivatives ∂_j on both sides of the deformed Hermitian-Yang-Mills equation, we have the equality

(2.6)
$$0 = \partial_j \theta = Tr(I + K^2)^{-1} \nabla_j K) = \eta^{p\bar{q}} g_{\bar{q}l} \nabla_j (g^{l\bar{m}} F_{\bar{m}p}) = \eta^{p\bar{q}} \nabla_j F_{\bar{q}p}.$$

where $\{\eta^{p\bar{q}}\}\$ is the inverse matrix of $\{\eta_{p\bar{q}}\}$.

2.2. **Proof of Theorem 1.1.** With notations above, we give a proof of Theorem 1.1 in this subsection.

Proof. It is easy to see that $\log \frac{\eta^n}{\omega^n}$ is a well-defined function on X. Indeed, it is nothing but $\log \det(I + K^2)$ if we view the endomorphism K as a matrix function on X. We use the Hermitian metric η to define the following Laplacian on $C^{\infty}(X)$

$$\Delta_{\eta} = \eta^{i\bar{j}} \nabla_i \nabla_{\bar{j}} : C^{\infty}(X) \to \mathbb{R},$$

where ∇ is the covariant derivative with respect to ω . This operator is nothing but the linearization operator of the fully nonlinear second order operator

$$\Theta(u) = \sum_{i=1}^{n} \arctan \lambda_i(u_{\alpha\bar{\beta}}).$$

We remark that the operator Δ_{η} is elliptic as long as F is bounded. Then we compute $\Delta_{\eta} \log \frac{\eta^n}{\omega^n}$ step by step.

(2.7)
$$\Delta_{\eta} \log \frac{\eta^{n}}{\omega^{n}} = \eta^{p\bar{q}} (\eta^{i\bar{j}} \eta_{i\bar{j},p})_{\bar{q}} \\ = -\eta^{p\bar{q}} \eta^{i\bar{t}} \eta^{s\bar{j}} \eta_{s\bar{t},\bar{q}} \eta_{i\bar{j},p} + \eta^{p\bar{q}} \eta^{i\bar{j}} \eta_{i\bar{j},p\bar{q}}$$

By the expression of η , we have

$$\eta_{i\bar{j},p} = F_{i\bar{b},p}g^{a\bar{b}}F_{a\bar{j}} + F_{i\bar{b}}g^{a\bar{b}}F_{a\bar{j},p}$$

and

$$\begin{aligned} \eta_{i\bar{j},p\bar{q}} &= (g_{i\bar{j}} + F_{i\bar{t}}g^{st}F_{s\bar{j}})_{,p\bar{q}} \\ &= F_{i\bar{t},p}g^{s\bar{t}}F_{s\bar{j},\bar{q}} + F_{i\bar{t},\bar{q}}g^{s\bar{t}}F_{s\bar{j},p} + F_{i\bar{t},p\bar{q}}g^{s\bar{t}}F_{s\bar{j}} + F_{i\bar{t}}g^{s\bar{t}}F_{s\bar{j},p\bar{q}} \end{aligned}$$

Taking trace with respect to η , we obtain

(2.8)
$$\Delta_{\eta} \log \frac{\eta^{n}}{\omega^{n}} = -\eta^{p\bar{q}} \eta^{i\bar{t}} \eta^{s\bar{j}} \eta_{s\bar{t},\bar{q}} \eta_{i\bar{j},p} + \eta^{p\bar{q}} \eta^{i\bar{j}} (F_{i\bar{t},p} g^{s\bar{t}} F_{s\bar{j},\bar{q}} + F_{i\bar{t},\bar{q}} g^{s\bar{t}} F_{s\bar{j},p}) + \eta^{p\bar{q}} \eta^{i\bar{j}} (F_{i\bar{t},p\bar{q}} g^{s\bar{t}} F_{s\bar{j}} + F_{i\bar{t}} g^{s\bar{t}} F_{s\bar{j},p\bar{q}}).$$

By the second Bianchi equality and communicating law of the covariant derivatives, we have

$$\begin{split} F_{i\bar{t},p\bar{q}} &= F_{p\bar{t},i\bar{q}} \\ &= F_{p\bar{t},\bar{q}i} + F_{a\bar{t}}g^{a\bar{b}}R_{p\bar{b}i\bar{q}} - F_{p\bar{b}}g^{a\bar{b}}R_{a\bar{t}i\bar{q}} \\ &= F_{p\bar{q},\bar{t}i} + F_{a\bar{t}}g^{a\bar{b}}R_{p\bar{b}i\bar{q}} - F_{p\bar{b}}g^{a\bar{b}}R_{a\bar{t}i\bar{q}}. \end{split}$$

Thus, we have

$$(2.9) \qquad \Delta_{\eta} \log \frac{\eta^{n}}{\omega^{n}} = -\eta^{p\bar{q}} \eta^{i\bar{t}} \eta^{s\bar{j}} \eta_{s\bar{t},\bar{q}} \eta_{i\bar{j},p} + \eta^{p\bar{q}} \eta^{i\bar{j}} (F_{i\bar{t},p} g^{s\bar{t}} F_{s\bar{j},\bar{q}} + F_{i\bar{t},\bar{q}} g^{s\bar{t}} F_{s\bar{j},p}) + \eta^{p\bar{q}} \eta^{i\bar{j}} g^{s\bar{t}} F_{s\bar{j}} (F_{p\bar{q},\bar{t}i} + F_{a\bar{t}} g^{a\bar{b}} R_{p\bar{b}i\bar{q}} - F_{p\bar{b}} g^{a\bar{b}} R_{a\bar{t}i\bar{q}}) + \eta^{p\bar{q}} \eta^{i\bar{j}} g^{s\bar{t}} F_{i\bar{t}} (F_{p\bar{q},\bar{j}s} + F_{a\bar{j}} g^{a\bar{b}} R_{p\bar{b}s\bar{q}} - F_{p\bar{b}} g^{a\bar{b}} R_{a\bar{j}s\bar{q}}).$$

On the other hand, taking derivative ∇_p on the both sides of (2.6), we get

(2.10)
$$\eta^{i\bar{j}}F_{i\bar{j},\bar{q}p} = -\eta^{ij}_{,p}F_{i\bar{j},\bar{q}}$$
$$= \eta^{i\bar{t}}\eta^{s\bar{j}}\eta_{s\bar{t},p}F_{i\bar{j},\bar{q}}$$
$$= \eta^{i\bar{t}}\eta^{s\bar{j}}F_{i\bar{j},\bar{q}}g^{a\bar{b}}(F_{s\bar{b},p}F_{a\bar{t}} + F_{s\bar{b}}F_{a\bar{t},p}).$$

Applying (2.6) and (2.10) to the equation (2.9), we have

$$\begin{split} \Delta_{\eta} \log \frac{\eta^{n}}{\omega^{n}} &= -\eta^{p\bar{q}} \eta^{i\bar{t}} \eta^{s\bar{j}} g^{a\bar{b}} g^{c\bar{d}} (F_{s\bar{b},\bar{q}} F_{a\bar{t}} + F_{s\bar{b}} F_{a\bar{t},\bar{q}}) (F_{i\bar{d},p} F_{c\bar{j}} + F_{i\bar{d}} F_{c\bar{j},p}) \\ &+ \eta^{p\bar{q}} \eta^{i\bar{j}} (F_{i\bar{t},p} g^{s\bar{t}} F_{s\bar{j},\bar{q}} + F_{i\bar{t},\bar{q}} g^{s\bar{t}} F_{s\bar{j},p}) \\ &+ \eta^{i\bar{j}} g^{s\bar{t}} F_{s\bar{j}} \eta^{p\bar{d}} \eta^{c\bar{q}} F_{p\bar{q},\bar{t}} g^{a\bar{b}} (F_{c\bar{b},i} F_{a\bar{d}} + F_{c\bar{b}} F_{a\bar{d},i}) \\ &+ \eta^{p\bar{q}} \eta^{i\bar{j}} g^{s\bar{t}} F_{s\bar{j}} (F_{a\bar{t}} g^{a\bar{b}} R_{p\bar{b}i\bar{q}} - F_{p\bar{b}} g^{a\bar{b}} R_{a\bar{t}i\bar{q}}) \\ &+ \eta^{i\bar{j}} g^{s\bar{t}} F_{i\bar{t}} \eta^{p\bar{d}} \eta^{c\bar{q}} F_{p\bar{q},\bar{j}} g^{a\bar{b}} (F_{c\bar{b},s} F_{a\bar{d}} + F_{c\bar{b}} F_{a\bar{d},s}) \\ &+ \eta^{p\bar{q}} \eta^{i\bar{j}} g^{s\bar{t}} F_{i\bar{t}} (F_{a\bar{j}} g^{a\bar{b}} R_{p\bar{b}s\bar{q}} - F_{p\bar{b}} g^{a\bar{b}} R_{a\bar{j}s\bar{q}}). \end{split}$$

For convenience, we take the normal coordinates around some point $p \in X$ such that $g_{i\bar{j}}(p) = \delta_{ij}$ and $F_{i\bar{j}}(p) = \lambda_i \delta_{ij}$. In particular, under this coordinates system, $\eta_{i\bar{j}}(p) = \theta_i \delta_{ij}$ where $\theta_i = 1 + \lambda_i^2$. We also denote $\theta^i = 1/\theta_i$.

Then at $p \in X$, we have

(2.11)

$$\Delta_{\eta} \log \frac{\eta^{\cdot \cdot}}{\omega^{n}}(p) = -\theta^{p} \theta^{i} \theta^{j} (\lambda_{i} + \lambda_{j})^{2} F_{i\bar{j},\bar{p}} F_{j\bar{i},p}$$

$$+ \theta^{p} \theta^{i} F_{i\bar{t},p} F_{t\bar{i},\bar{p}} + \theta^{p} \theta^{i} F_{i\bar{t},\bar{p}} F_{t\bar{i},p}$$

$$+ 2\theta^{i} \theta^{p} \theta^{q} \lambda_{i} (\lambda_{p} + \lambda_{q}) F_{p\bar{q},\bar{i}} F_{q\bar{p},i}$$

$$+ 2\theta^{i} \theta^{p} \lambda_{i}^{2} R_{i\bar{i}p\bar{p}} - 2\theta^{i} \theta^{p} \lambda_{i} \lambda_{p} R_{i\bar{i}p\bar{p}}.$$

Inserting the formula $\theta^i(1+\lambda_i^2)=1$ to the second and third items, we get

$$\begin{split} \Delta_{\eta} \log \frac{\eta^{n}}{\omega^{n}}(p) &= -\theta^{p} \theta^{i} \theta^{j} (\lambda_{i}^{2} + 2\lambda_{i}\lambda_{j} + \lambda_{j}^{2}) F_{i\bar{j},\bar{p}} F_{j\bar{i},p} \\ &+ \theta^{p} \theta^{i} \theta^{j} (1 + \lambda_{j}^{2}) F_{i\bar{j},\bar{p}} F_{j\bar{i},p} + \theta^{p} \theta^{i} \theta^{j} (1 + \lambda_{j}^{2}) F_{i\bar{j},\bar{p}} F_{j\bar{i},p} \\ &+ 2\theta^{i} \theta^{j} \theta^{p} \lambda_{i} \lambda_{j} F_{j\bar{i},\bar{p}} F_{i\bar{j},p} + 2\theta^{i} \theta^{j} \theta^{p} \lambda_{i} \lambda_{j} F_{i\bar{j},\bar{p}} F_{j\bar{i},p} \\ &+ 2\sum_{i < p} (\theta^{i} \theta^{p} \lambda_{i}^{2} R_{i\bar{i}p\bar{p}} - 2\theta^{i} \theta^{p} \lambda_{i} \lambda_{p} R_{i\bar{i}p\bar{p}} + \theta^{i} \theta^{p} \lambda_{p}^{2} R_{i\bar{i}p\bar{p}}) \\ &= 2\theta^{i} \theta^{j} \theta^{p} (1 + \lambda_{i} \lambda_{j}) F_{p\bar{j},\bar{i}} F_{j\bar{p},i} + 2\theta^{i} \theta^{p} R_{i\bar{i}p\bar{p}} (\lambda_{i} - \lambda_{p})^{2}. \end{split}$$

Since $-\frac{1}{C}\omega < \sqrt{-1}F < C\omega$, we know that $\lambda_i\lambda_j > -1$. From $R_{i\bar{i}p\bar{p}} \ge 0$, we know that $\log \frac{\eta^n}{\omega^n}$ is a subharmonic function on X. By maximal principle, we know that $\log \frac{\eta^n}{\omega^n}$ is constant. Hence, $F_{i\bar{j},p} = 0$, i.e. all λ_i are constants(maybe different for each i).

In particular, if $R_{i\bar{i}p\bar{p}}$ is strictly positive at some point $p_0 \in M$, then $\lambda_i = \lambda_p$ for any $i \neq p$, i.e. $\sqrt{-1}F = C\omega$.

Remark 2.1. Jacob-Yau [18] can prove the existence of dHYM metrics on ample line bundle over a compact Kähler manifold with non-negative orthogonal bisectional curvature under the assumption of hypercritical phase $(\theta \in (\frac{n-1}{2}\pi, \frac{n}{2}\pi))$ using the line bundle mean curvature flow. In particular, the hypercritical phase condition implies $\sqrt{-1}F$ is positive.

More pricisely, we have the following property for Kähler manifold with nonnegative orthogonal bisectional holomorphic curvature which is strictly positive at some point. **Proposition 2.1.** If a compact Kähler manifold (M, ω) has nonnegative orthogonal bisectional curvature and its orthogonal bisectional curvature is strictly positive at some point $p \in M$, then the second betti number of M is 1, i.e.

$$\dim H^{1,1}(M,\mathbb{R}) = 1$$

Proof. For any $[\alpha] \in H^{1,1}(M, \mathbb{R})$, we know that there exists a smooth function on M such that

$$\operatorname{Tr}_{\omega}(\alpha + \sqrt{-1}\partial\bar{\partial}u) = c$$

where $c = \frac{n \int_M \alpha \wedge \omega^{n-1}}{\int_M \omega^n}$ is a constant dependent only on $[\alpha]$ and $[\omega]$. By direct computation and communicating law of covariant derivatives, we have

$$\begin{split} \Delta_{\omega} \left| \alpha_{u} \right|_{\omega}^{2} &= 2 \langle \Delta_{\omega} \alpha_{u}, \alpha_{u} \rangle_{\omega} + 2 \left| \nabla_{\omega} \alpha_{u} \right|_{\omega}^{2} \\ &= 2 g^{i\bar{j}} g^{k\bar{n}} g^{m\bar{l}} \alpha_{m\bar{n}} (\alpha_{i\bar{j},\bar{l}k} + \alpha_{p\bar{l}} R_{i\bar{p}k\bar{j}} - \alpha_{i\bar{p}} R_{p\bar{l}k\bar{j}}) + 2 \left| \nabla_{\omega} \alpha_{u} \right|_{\omega}^{2}. \end{split}$$

where $g_{i\bar{j}}$ and $\alpha_{i\bar{j}}$ are the coefficients with respect to ω and $\alpha_u = \alpha + \sqrt{-1}\partial\bar{\partial}u$. Since $g^{i\bar{j}}\alpha_{i\bar{j}} = c$, we know

$$g^{i\bar{j}}\alpha_{i\bar{i},k\bar{l}} = 0.$$

Then we have

$$\Delta_{\omega} |\alpha_{u}|_{\omega}^{2} = 2g^{i\bar{j}}g^{k\bar{n}}g^{m\bar{l}}\alpha_{m\bar{n}}(\alpha_{p\bar{l}}R_{i\bar{p}k\bar{j}} - \alpha_{i\bar{p}}R_{p\bar{l}k\bar{j}}) + 2|\nabla_{\omega}\alpha_{u}|_{\omega}^{2}$$
$$= R_{i\bar{i}p\bar{p}}(\lambda_{i} - \lambda_{p})^{2} + 2|\nabla_{\omega}\alpha_{u}|_{\omega}^{2}$$

where λ_i are the eigenvalues of $\omega^{-1}\alpha_u$ as an endomorphism on $T^{1,0}M$. By our assumption $R_{i\bar{i}j\bar{j}} \geq 0$, we know $\nabla_{\omega}\alpha_u = 0$, i.e. all λ_i are constants. Since $R_{i\bar{i}j\bar{j}}$ is strictly positive at some point, we know $\lambda_i = C$ for some constant C, i.e. $\alpha_u = C\omega$. In conclusion, $[\alpha] = C[\omega]$, i.e. dim $H^{1,1}(M, \mathbb{R}) = 1$.

Remark 2.2. In [12], Guan-Li-Zhang consider a uniqueness result of

$$F(Ric) = f(\lambda_1(Ric), \cdots, \lambda_n(Ric)) = 0$$

for some fully nonlinear function F under the assumption that non-negative orthogonal bisectional curvature is strictly positive at some point. They require fsatisfies one of the following conditions

- (1) f is concave,
- (2) as a function acted on the eigenvalues of A, f satisfies that if $\forall 0 \leq l \leq n$, for all $0 \leq \lambda_1 \leq \cdots \leq \lambda_n$ and $\lambda_i > 0$, $\forall i \geq n l + 1$, there holds

$$\sum_{j,k=n-l+1}^{n} \ddot{f}^{jk}(A) X_{j\bar{j}} X_{k\bar{k}} + 2 \sum_{n-l+1 \le j < k} \frac{\dot{f}^j - \dot{f}^k}{\lambda_j - \lambda_k} |X_{j\bar{k}}|^2 + \sum_{i,k=n-l+1}^{n} \frac{\dot{f}^i(A)}{\lambda_k} |X_{i\bar{k}}|^2 \ge 0$$

for every Hermitian matrix $X = (X_{j\bar{k}})$ with $X_{j\bar{k}} = 0$ if $j \le n - l$.

Here \cdot means the derivatives of f about the eigenvalues of A.

We can check that f in our theorem does not satisfy these two conditions above if we assume $-\frac{1}{C}\omega < \sqrt{-1}F < C\omega$ for C > 1. In fact,

(1) if at least one of $\lambda_i < 0$, then f is not concave, since

$$\ddot{f}(\cdots,\lambda_i+t,\cdots)|_{t=0} = \frac{-2\lambda_i}{(1+\lambda_i^2)^2} > 0.$$

(2) if we assume $\lambda_n > 1$ and $\lambda_i = 0$ for $i \le n-1$, $X_{j\bar{k}} = 0$ for $j \le n-1$, $k \le n-1$ and $X_{n\bar{n}} = 1$, then the left-hand side of the inequality in condition (2) is $\frac{1-\lambda_n^2}{\lambda_n(1+\lambda_n^2)^2}$ which is negative.

2.3. **Proof of Theorem 1.2.** We can prove Theorem 1.2 by computing the same auxiliary function as in the proof of Theorem 1.1. In this subsection, we will take

$$\eta = \frac{\sqrt{-1}}{2} \eta_{i\bar{j}} dz^i \wedge d\bar{z}^j = \frac{\sqrt{-1}}{2} \omega_{i\bar{l}} \chi^{k\bar{l}} \omega_{k\bar{j}} dz^i \wedge d\bar{z}^j$$

where $\omega = \frac{\sqrt{-1}}{2} \omega_{i\bar{j}} dz^i \wedge d\bar{z}^j$ and $\chi = \frac{\sqrt{-1}}{2} \chi_{i\bar{j}} dz^i \wedge d\bar{z}^j$. We choose the normal coordinates near a point $p \in M$ such that

$$\chi(p) = \frac{\sqrt{-1}}{2} \delta_{ij} dz^i \wedge d\bar{z}^j \text{ and } \omega(p) = \frac{\sqrt{-1}}{2} \delta_{ij} \lambda_i dz^i \wedge d\bar{z}^j$$

In particular, we have $\eta_{i\bar{j}}(p) = \delta_{ij}\lambda_i^2$. We also denote $\Delta_{\eta} = \eta^{i\bar{j}}\nabla_{\bar{j}}\nabla_i$ where ∇ is the covariant derivative with respect to χ . Indeed, Δ_{η} is the linearization operator of the *J*-equation. By direct computation, we have

$$\Delta_{\eta} \log \frac{\eta^n}{\chi^n} = \frac{2}{\lambda_i \lambda_j \lambda_p^2} \omega_{p\bar{j},i} \omega_{j\bar{p},\bar{i}} + \frac{2}{\lambda_i^2 \lambda_p^2} R_{i\bar{i}p\bar{p}} (\lambda_i - \lambda_p)^2 \ge 0.$$

Same arguments as in the proof of Theorem 1.1 implies the result needed.

Remark 2.3. Theorem 1.2 can be proved by the same argument in [12] since the operator

$$f(A) = \sum_{i=1}^{n} \frac{1}{\lambda_i}$$

is concave at any positive Hermitian matrix A. So our proof can be seen as a new proof of the result in [12] when the operator is the special one above.

3. RIGID RESULTS FOR SELF-SHRINKERS

In this section we choose the suitable barrier function to prove Theorem 1.3.

Proof. We denote
$$\Theta(z) = \sum_{i=1}^{n} \arctan \lambda_i(z)$$
. Then we have

(3.1)
$$\Theta_{\bar{i}} = \eta^{k\bar{l}} u_{k\bar{l}\bar{i}}, \Theta_i = \eta^{k\bar{l}} u_{k\bar{l}i}.$$

On the other hand, by taking derivatives on the right side of (1.3), we get that

(3.2)
$$(\frac{1}{2}(\langle z, \partial u \rangle + \langle \bar{z}, \bar{\partial} u \rangle) - u)_{i\bar{j}} = \frac{1}{2}(z_k u_{ki\bar{j}} + \bar{z}_k u_{\bar{k}i\bar{j}}).$$

1

Combining the equations above, we have

$$\eta^{i\bar{j}}\Theta_{i\bar{j}} - \frac{1}{2}(\langle z, \partial \Theta \rangle + \langle \bar{z}, \bar{\partial} \Theta \rangle) = 0.$$

We construct a barrier function as follow

$$v(r) = \varepsilon r^2 + \max_{\partial B_{r_0}} \{\Theta\}$$

where $r_0 = \sqrt{n}$ and $\varepsilon \in \mathbb{R}^+$. By direct computation, we have

(3.3)
$$\frac{1}{4}(w_{rr} + \frac{2n-1}{r}w_r) - \frac{r}{2}w_r = (\frac{n}{2r} - \frac{r}{2})w_r \le 0$$

for any $r \geq r_0$. Furthermore,

$$\sqrt{-1}\partial\bar{\partial}w = \varepsilon \sum \sqrt{-1}dz^i \wedge d\bar{z}^i > 0.$$

Hence, for z outside of the ball B_{r_0} , we have

$$\eta^{i\bar{j}}w_{i\bar{j}} - \frac{1}{2}(\langle z, \partial w \rangle + \langle \bar{z}, \bar{\partial}w \rangle) \le \frac{1}{4}\Delta w - \frac{1}{2}(\langle z, \partial w \rangle + \langle \bar{z}, \bar{\partial}w \rangle) \le 0$$

where we use the inequality (3.3) and the fact that $\eta^{i\bar{j}} \leq I$.

So far we have

$$\eta^{i\bar{j}}w_{i\bar{j}} - \frac{1}{2}(\langle z, \partial w \rangle + \langle \bar{z}, \bar{\partial}w \rangle) \le \eta^{i\bar{j}}\Theta_{i\bar{j}} - \frac{1}{2}(\langle z, \partial \Theta \rangle + \langle \bar{z}, \bar{\partial}\Theta \rangle)$$

if $|z| \ge r_0$. On the other hand, we know

$$w(r_0) = \varepsilon r_0^2 + \max_{\partial B_{r_0}} \{\Theta\} \ge \Theta \text{ on } \partial B_{r_0}$$

and

$$w(|x|) > \Theta(x)$$
 when $|x| \to \infty$

since $w(|x|) \to \infty$ as $|x| \to \infty$ while Θ is bounded. Hence $w \ge \Theta$ outside B_{r_0} by maximum principle. As ε tends to 0, we know that Θ attains its global maximum on \mathbb{C}^n in the closure of B_{r_0} . By the strong maximum principle, we know Θ is a constant.

According to the equation (3.2), we know that $z \cdot \partial u_{i\bar{j}} + \bar{z} \cdot \bar{\partial} u_{i\bar{j}} = 0$. That implies that $r(u_{i\bar{j}})_r = 0$. Thus $u_{i\bar{j}} = c$ except for 0. However, $u_{i\bar{j}}$ is smooth on the whole \mathbb{C}^n , therefore $\{u_{i\bar{j}}\} = \{C_{i\bar{j}}\}$ on \mathbb{C}^n where $\{C_{i\bar{j}}\}$ is a constant Hermitian matrix. Then we know that the function

$$v = u - \sum_{i,j=1}^{n} C_{i\bar{j}} z^i \bar{z}^j$$

is a pluriharmonic function, i.e. $\partial \overline{\partial} v = 0$. In particular, there exists a holomorphic f on \mathbb{C}^n such that

$$u = \sum_{i,j=1}^{n} C_{i\bar{j}} z^{i} \bar{z}^{j} + f - \bar{f}.$$

Considering the Laurent series of f and the self-shrinker equation (1.3), we can prove that $f - \overline{f}$ is quadratic. And so is u.

Remark 3.1. In [17], a rigid result for self-shrinker to the flow is proved. However, the function (or as an Hermitian metric of the trivial line bundle) should satisfy the so-called graphical condition in [17]. Here, we do not need any condition on u.

We can prove Theorem 1.4 by choosing analogous auxiliary function as in [2].

Proof. We denote
$$J(z) = -\sum_{i=1}^{n} \frac{1}{\lambda_i(z)}$$
. Then
(3.4) $J_{\overline{i}} = \eta^{k\overline{l}} u_{k\overline{l}\overline{i}}, J_i = \eta^k$

3.4)
$$J_{\overline{i}} = \eta^{k\overline{l}} u_{k\overline{l}\overline{i}}, J_{i} = \eta^{k\overline{l}} u_{k\overline{l}i}$$

where $\eta^{k\bar{l}} = \frac{\delta_{kl}}{\lambda_l^2}$. Combining the equations (3.2) and (1.4), we get that

$$\eta^{i\bar{j}}J_{i\bar{j}} - \frac{1}{2}(\langle z, \partial J \rangle + \langle \bar{z}, \bar{\partial}J \rangle) = 0.$$

We assume that $(u_{i\bar{j}}) \geq \frac{\sqrt{2(2n-1+\delta)}}{r}I$ if $r \geq 1$. We choose a barrier function as follow

$$w(r) = \varepsilon r^{1+\delta} + \max_{\partial B_1} \{-J\}.$$

By direct computation, we have

(3.5)
$$\frac{r^2}{2(2n-1+\delta)}\Delta\omega - \frac{r}{2}w_r = 0.$$

Furtheremore,

$$(\omega_{i\bar{j}}) = \frac{\omega_r}{2r} (\delta_{ij} + \frac{(\delta - 1)\bar{z}^i z^j}{2r^2}),$$

since for any $X = \sum_{i=1}^{n} X^{i} \frac{\partial}{\partial z^{i}}$,

$$\sqrt{-1}\partial\bar{\partial}w(X,\bar{X}) = X^{i}(\delta_{ij} + \frac{(\delta-1)\bar{z}^{i}z^{j}}{2r^{2}})\bar{X}^{j} \ge |X|^{2} + \frac{\delta-1}{2}|X|^{2} > 0.$$

Thus for any z outside of B_1 , we have

$$\eta^{i\bar{j}}w_{i\bar{j}} - \frac{1}{2}(\langle z, \partial w \rangle + \langle \bar{z}, \bar{\partial}w \rangle) \le \frac{r^2}{2(2n-1+\delta)}\Delta w - \frac{1}{2}(\langle z, \partial w \rangle + \langle \bar{z}, \bar{\partial}w \rangle) = 0$$

where we use the fact that $\eta^{i\bar{j}} \leq \frac{r^2}{2n-1+\delta}I$. So far we have

$$\eta^{i\bar{j}}w_{i\bar{j}} - \frac{1}{2}(\langle z, \partial w \rangle + \langle \bar{z}, \bar{\partial}w \rangle) \leq \eta^{i\bar{j}}(-J)_{i\bar{j}} - \frac{1}{2}(\langle z, \partial(-J) \rangle + \langle \bar{z}, \bar{\partial}(-J) \rangle)$$

if $|z| \ge 1$. On the other hand, we know on ∂B_1

$$w = \varepsilon + \max_{\partial B_1} \{-J\} \ge -J$$

and

$$w(|z|) > -J(z)$$
 when $|z| \to \infty$

by our assumption on $\sqrt{-1}\partial \bar{\partial} u$. Hence $w \geq -J$ outside B_1 by maximum principle. As $\varepsilon \to 0$, we know that -J attains its global maximum in the closure of B_1 . By the strong maximum principle, we know -J is a constant.

By similar argument as in the proof of Theorem 1.3, we know that u is a quadratic polynomial function.

A direct consequence of the rigid theorem above is the following rigid result for self-shrinkers of J-flow.

Corollary 3.1. If $u : \mathbb{C}^n \to \mathbb{R}$ is a smooth solution to (1.4) satisfying

$$\{u_{i\bar{j}}\} \ge CI$$

for some constant C, then u is quadratic.

Remark 3.2. We should also remark that the rigid theorem for a class of fully nonlinear second elliptic operator is proved in Theorem 1.2 [30]. However, the operator in [30] acts on the real symmetric matrices. Using the canonical relation between Hermitian matrices and real symmetric matrices, we can regard the operator facting on $n \times n$ Hermitian matrices as an operator \tilde{f} acting on $2n \times 2n$ real symmetric matrices. In fact, if we write Hermitian matrix A as $A = E + \sqrt{-1}F$ where E and F are real matrices such that $E^T = E$ and $F^T = -F$, then the real symmetric matrix B related to A can be taken as

$$B = \left(\begin{array}{cc} E/2 & F^T/2 \\ F/2 & E/2 \end{array}\right)$$

Then the operator $f: A \to \mathbb{R}$

$$f(A) = \sum_{i=1}^{n} \arctan \lambda_i(A)$$

can be regarded as a operator \tilde{f} on B

$$\tilde{f}(B) = \frac{1}{2} \sum_{i=1}^{2n} \arctan \lambda_i (B + J \cdot B \cdot J^T)$$

where

$$J = \left(\begin{array}{cc} 0 & I \\ -I & 0 \end{array}\right)$$

is the standard complex structure on \mathbb{C}^n . In particular, the new operator \tilde{f} does not satisfies condition (iii) of Theorem 1.2 in [30], i.e. for any positive B

$$\left\| D\tilde{f}(B) \cdot B \right\| \le k_2,$$

where $D\tilde{f}(B) = (\frac{\partial \tilde{f}}{\partial B_{ij}}), \|\cdot\|$ is a fixed norm on the vector space of the matrices and k_2 is a certain constant. In particular, we consider the case $n = 1, A = a + \sqrt{-1c}$ and we denote

$$B = \left(\begin{array}{cc} a & c \\ c & a \end{array}\right)$$

for $a, c \in \mathbb{R}$. Then \tilde{f} can be expressed by

$$\tilde{f}(B) = \arctan(a).$$

In particular, $D\tilde{f}(B) \cdot B = \frac{B}{1+a^2}$ and its norm can not be controlled by a fixed constant.

References

- M. Berger, Sur les variétés d'Einstein compactes, Comptes Rendus de la IIIe Réunion du Groupement des Mathématiciens d'Expression Latine, pp. 35-55, (1965).
- [2] A. Chau, J.Y. Chen and Y. Yuan Rigidity of entire self-shrinking solutions to curvature flows, J. Reine Angew. Math, vol. 664, pp.229C239, (2012).
- [3] G. Chen, On J-equation, arxiv:1905.10222, (2019).
- [4] T. Collins, A. Jacob and S.-T. Yau, (1,1)-forms with specified Lagrangian phase: a priori estimates and algebraic obstructions, arxiv: 1508.01934, (2015).
- [5] T. Collins, G. Székelyhidi, Convergence of the J-flow on toric manifolds, Journal of Differential Geometry, vol. 107(1), pp. 47-81, (2017).
- [6] T. Collins, D. Xie and S.-T. Yau, The deformed Hermitian-Yang-Mills in geometry and Physics, arxiv:1712.00893, (2017)

- [7] X.X. Chen and G. Tian, Ricci flow on Kähler-Einstein manifolds, Duke Math. Journal, vol. 131(1), pp. 17-73, (2006).
- [8] S.K. Donaldson, Moment maps and diffeomorphisms, Asian J. Math, vol. 3(1), pp. 1-16, (1999).
- [9] G. Drugan, P. Lu and Y. Yuan, Rigidity of complete entire self-shrinking solutions to Khler-Ricci flow, Int. Math. Res. Not, vol 12, pp. 3908-3916, (2015).
- [10] Q. Ding and Y.L. Xin, The rigidity theorems for Lagrangian self shrinkers, J. Reine Angew. Math, vol. 692, pp. 109-123, (2014).
- [11] S. Golberg and S. Kobayashi, Holomorphic Bisectional Curvature, J. Diiferential Geom. vol. 1, pp. 225-233, (1967).
- [12] P.F. Guan, Q. Li and X. Zhang, A uniqueness theorem in Kähler geometry, Math. Ann. vol. 345(2), pp. 377-393, (2009).
- [13] H.L. Gu and Z.H. Zhang, An extension of Moks Theorem on the generalized Frankel conjecture, Sci. China, Math. vol. 53, pp. 1253C1264,(2010).
- [14] A. Howard, B. Smyth and H. Wu, On compact Kähler manifolds of nonnegative bisectional, I, Acta Math. vol. 147, pp. 51-56, (1981).
- [15] R.L. Huang, Q.Z. Ou and W.L. Wang, On the entire self-shrinking solutions to Lagrangian mean curvature flow II, arXiv:1904.07713.
- [16] R.L. Huang and Z.Z. Wang, On the entire self-shrinking solution to Lagrangian mean curvature, alc. Var. Partial Differential Equations, vol. 41, pp. 321-339, (2011).
- [17] X.L. Han and H. Yamamoto, A $\varepsilon\text{-regularity theorem for line bundle mean curvature flow arxiv1904.02391.$
- [18] A. Jacob and S.-T. Yau, A special Lagrangian type equation for holomorphic line bundle, Math. Ann. vol. 369(1-2), pp. 869-898, (2017).
- [19] C. Leung, S.-T. Yau and E. Zaslow, From special Lagrangian to Hermitian-Yang-Mills via Fourier-Mukai transform, Winter School on Mirror Symmetry, Vector Bundles and Lagrangian Submanifolds(Cambridge,MA,1990).AMS/IP Stud. Adv.Math., vol. 23, pp. 209-225. American Mathematical Society, Providence, RI(2001)
- [20] Q. Li, D. Wu and F. Zheng, An example of compact Kähler manifold with nonnegative quadratic bisectional curvature, Proceedings of the American Mathematical Society. vol. 141(6), pp. 2117-2126, (2013).
- M. Marino, R. Minasian, G. Moore, A. Stromiger, Nonlinear instantons from supersymmetric p-Branes, arXiv:hep-th/9911206
- [22] N. Mok, The Uniformization Theorem for Compact Kähler Manifolds of Nonnegative Holomorphic Bisectional Curvature, J. Diff. Geom. vol. 27, pp. 179C214,(1988).
- [23] S. Mori Projective manifolds with ample tangent bundles, Ann. of Math. vol. 110, pp. 593-606, (1979).
- [24] V. P. Pingali A priori estimates for a generalised Monge-Ampère PDE on some compact K?hler manifolds, Complex Variables and Elliptic Equations, vol. 24, pp. 1-15, (2016).
- [25] V. P. Pingali A note on the deformed Hermitian Yang-Mills PDE, Complex Variables and Elliptic Equations, vol. 64(3), pp. 503-518, (2019).
- [26] Y.T. Siu and Y.-T. Yau, Compact Kähler manifolds of positive bisectional curvature, Invent. Math. vol. 59(2), pp. 189-204, (1980)
- [27] J. Song and B. Weinkove, On the convergence and singularities of the J-flow with applications to the Mabuchi energy, Comm. Pure. Appl. Math, vol. 61(2), pp. 210-229, (2008).
- [28] J. Streets and M. Warren, EvansCKrylov Estimates for a nonconvex MongeCAmpère equation, Math. Ann. vol. 365, pp. 805-834, (2016)
- [29] W.L. Wang, Rigidity of entire self-shrinking solutions to Kähler-Ricci flow on complex plane, Proc. Amer. Math. Soc, vol. 145(7), pp. 3105-3108, (2017)
- [30] W.L. Wang, Rigidity of entire convex self-shrinking solutions to Hessian quotient flows, Int. Math. Res. Not, vol. 24, pp. 7755-7775, (2018)
- [31] Y. Yuan, A Bernstein problem for special Lagrangian equations, Invent. Math, vol. 150(1), pp. 117-125, (2002)
- [32] Y. Yuan, Global solutions to special Lagrangian equations, Proc. Amer. Math. Soc, vol. 134(5), pp. 1355-1358, (2006)

XIAOLI HAN, MATH DEPARTMENT OF TSINGHUA UNIVERSITY, BEIJING, 100084, CHINA, E-mail address: hanxiaoli@mail.tsinghua.edu.cn

Xishen Jin, Department of Mathematics, Renmin University of China, Beijing, 100872, China,

 $E\text{-}mail \ address: \texttt{jinxishen} \texttt{Cruc.edu.cn}$