

# BERGMAN AND SZEGÖ PROJECTIONS, EXTREMAL PROBLEMS, AND SQUARE FUNCTIONS

TIMOTHY FERGUSON

ABSTRACT. We study estimates for Hardy space norms of analytic projections. We first find a sufficient condition for the Bergman projection of a function in the unit disc to belong to the Hardy space  $H^p$  for  $1 < p < \infty$ . We apply the result to prove a converse to an extension of Ryabykh's theorem about Hardy space regularity for Bergman space extremal functions. We also prove that the  $H^q$  norm of the Szegő projection of  $F^{p/2}\bar{F}^{(p/2)-1}$  cannot be too small if  $F$  is analytic, for certain values of  $p$  and  $q$ . We apply this to show that the best analytic approximation in  $L^p$  of a function in both  $L^p$  and  $L^q$  will also lie in  $L^q$ , for certain values of  $p$  and  $q$ .

## 1. INTRODUCTION

This paper deals with estimates for Hardy space norms of analytic projections. One of the main tools is a result (due to Calderon) about square functions of functions of the form  $|F|^\delta$ , where  $F$  is analytic in the unit disc. We find a sufficient condition for the Bergman projection of a function to lie in a Hardy space. We also apply our results to extremal problems in spaces of analytic functions.

We begin with some background. Let  $\mathcal{P}(f)$  denote the Bergman projection of a function  $f \in L^1(\mathbb{D})$ . This can be defined by the formula

$$\mathcal{P}(f)(z) = \int_{\mathbb{D}} f(w) \frac{1}{(1 - z\bar{w})^2} \frac{dA(w)}{\pi}.$$

The Bergman projection is the projection of  $L^2(\mathbb{D}, dA/\pi)$  onto the Bergman space  $A^2(\mathbb{D})$  consisting of all analytic functions in  $L^2(\mathbb{D})$ . The Bergman spaces  $A^p$  for  $1 < p < \infty$  are defined similarly.

There are many results about when a Bergman projection lies in certain weighted Lebesgue or Sobolev spaces, but we want to find a condition for a Bergman projection to lie in a Hardy space. Recall that  $f$  is in the Hardy space  $H^p$  if it is analytic in the unit disc and

$$\|f\|_{H^p} = \sup_{0 \leq r < 1} M_p(r, f) < \infty$$

---

*Date:* September 24, 2019.

where

$$M_p(r, f) = \left\{ \frac{1}{2\pi} \int_0^{2\pi} |f(re^{i\theta})|^p d\theta \right\}^{1/p}.$$

For a function  $f \in W_{\text{loc}}^{1,2}(\mathbb{D})$  we may define

$$S_z(f)(e^{i\theta})^2 = \iint_{\Gamma(\theta)} |\partial_z f(re^{i\phi})|^2 dr d\theta,$$

where  $\Gamma_\theta$  is a cone centered at  $e^{i\theta}$ , to be specified later. For an analytic function, this is equivalent to the Zygmund  $S$  function. Our first result says that, under suitable conditions, if  $S_z(f) \in L^p(\partial\mathbb{D})$  then  $\mathcal{P}(f) \in H^p$  for  $1 < p < \infty$ .

An application is as follows. Let  $1 < p < \infty$ . Then the dual space  $(A^p)^*$  is isomorphic to  $A^{p'}$ . Given a nonzero  $k \in A^{p'}$ , we say the extremal function  $F$  associated with the integral kernel  $k$  is the function of unit  $A^p$  norm such that

$$\operatorname{Re} \frac{1}{\pi} \int_{\mathbb{D}} F \bar{k} dA = \sup_{f \in A^p, \|f\|=1} \operatorname{Re} \frac{1}{\pi} \int_{\mathbb{D}} f \bar{k} dA.$$

It is known that  $F$  exists and is unique. Also,

$$k = \|k\|_{(A^p)^*} \mathcal{P}(F^{p/2} \bar{F}^{(p/2)-1}).$$

One important result about these problems is that if  $k \in H^q$  for some  $q$  such that  $p' \leq q < \infty$  then  $F \in H^{(p-1)q}$ . This was first proved by Ryabykh [10] in the case  $q = p'$  for  $1 < p < \infty$ . The full result was proved by Ferguson [3, 4].

It was previously known that the converse of this statement holds for  $p$  an even integer [3]. In fact, if  $0 < q < \infty$  and  $p$  is an even integer and  $F \in H^{(p-1)q}$  then  $k \in H^q$ . We prove that this converse result in fact holds for  $1 < p < \infty$ .

For our next result, we prove that for certain  $p$  and  $q$ , there is a constant  $0 \leq C < 1$  such that  $\|P_S f\|_q \geq (1 - C)\|f\|_q$ , where  $f = F^{p/2} \bar{F}^{(p/2)-1}$  for analytic  $F$  and  $P_S$  is the Szegő projection. The Szegő projection is the orthogonal projection of  $L^2(\partial\mathbb{D})$  onto  $H^2$ . We then apply this result to extremal problems for Hardy spaces. It is known that  $(H^p)^* = H^{p'}$  for  $1 < p < \infty$ . As before, we can define a Hardy space extremal function  $F$  for  $k \in H^{p'}$  by

$$\frac{1}{2\pi} \operatorname{Re} \int_{\mathbb{D}} F \bar{k} d\theta = \sup_{f \in A^p, \|f\|=1} \frac{1}{2\pi} \operatorname{Re} \int_{\mathbb{D}} f \bar{k} d\theta.$$

As before, the extremal function will exist and be unique for  $1 < p < \infty$  and

$$k = \|k\|_{(H^p)^*} P_S(F^{p/2} \bar{F}^{(p/2)-1}).$$

Using this with our result shows that if  $k \in H^q \cap H^{p'}$  and  $F$  is the  $H^p$  extremal function for  $k$ , then for certain values of  $p$  and  $q$  we have that  $F \in H^{q(p-1)}$ .

We then apply the theory of dual extremal problems developed by S. Ya. Khavinson [6–8], and Rogosinski and Shapiro [9] (see [1, Chapter 8] for a good introduction). One of the results from this theory is that for  $k \in L^{p'}$  and  $1 < p < \infty$  we have

$$\max_{F \in H^p, \|F\|_p=1} \operatorname{Re} \frac{1}{2\pi} \int_0^{2\pi} F(e^{i\theta}) k(e^{i\theta}) d\theta = \min_{g \in H_0^{p'}} \|k - g\|_{p'}$$

where both the minimum and the maximum are attained, and  $H_0^{p'}$  consists of those functions in  $H^{p'}$  that vanish at the origin. Moreover,

$$k - g = \|k\|_{(H^p)^*} F^{p/2} \overline{F}^{(p/2)-1}$$

by the conditions for equality in Hölder's inequality. The function  $k - g$  is called the extremal kernel.

Using this theory of dual extremal problems, we derive a theorem on best approximation of functions in  $L^p$  spaces. Namely, for certain values of  $p$  and  $q$ , if  $k \in H^q \cap H^p$  and  $f$  is the analytic function in  $H^p$  that is closest to  $k$ , then  $f \in H^q$  as well.

We now state the result of Calderon that we will use. Let

$$S(G)(e^{i\theta})^2 = \iint_{\Gamma_\theta} |\nabla G(re^{i\phi})|^2 dr d\theta$$

where  $\Gamma_\theta$  is the cone  $\{(r, \phi) : |\theta - \phi| < (1 - r)/2, 0 < r < 1\}$ .

**Theorem A.** *Let  $F$  be analytic and let  $G = |F|^\delta$  for  $\delta > 0$ . Let  $0 < p < \infty$ . Then there are constants  $C_{p,\delta}$  and  $\widehat{C}_{p,\delta}$  such that*

$$\|S(G)\|_p \leq C_{p,\delta} \|G\|_p$$

and

$$\|G\|_p \leq \widehat{C}_{p,\delta} \|S(G)\|_p \text{ if } F(0) = 0.$$

We remark that the constants in this theorem may be explicitly estimated by following Calderon's proof.

## 2. BERGMAN PROJECTIONS IN HARDY SPACES

Our first result is as follows.

**Theorem 2.1.** *If  $1 < p < \infty$  and  $f \in W^{1,1}(\mathbb{D}) \cap L^{2p/(p+1)}(\mathbb{D})$  and  $M_1(r, f) = o((1 - r)^{-1})$  and  $S_z(f) \in L^p$ , then  $\mathcal{P}f \in H^p$  and*

$$\|\mathcal{P}(f)\|_{H^p} \leq C_p (\|S_z(f)\|_p + \|f\|_{L^{2p/(p+1)}(\mathbb{D})})$$

*Proof.* Let  $h$  be an analytic polynomial of degree at most  $n$ . Let  $\mathcal{P}_n$  be the orthogonal projection from  $L^2$  onto the polynomials in  $z$  of degree at most  $n$ . The Cauchy-Green theorem shows that

$$\frac{i}{2\pi} \int_0^{2\pi} \mathcal{P}_n(f) \bar{h} d\theta = \frac{1}{2\pi} \int_0^{2\pi} \mathcal{P}_n(f) z \bar{h} dz = \frac{1}{\pi} \int_{\mathbb{D}} \mathcal{P}_n(f) \overline{(zh)'} dA(z).$$

But since  $(zh)'$  has degree at most  $n$ , the right side of the above equals

$$\frac{1}{\pi} \int_{\mathbb{D}} f \overline{(zh)'} dA(z).$$

Now we may use the product rule and integrate one of the resulting integrals by parts to see that the above expression equals

$$\frac{1}{\pi} \int_{\mathbb{D}} \partial_z(f) \overline{h'(z)} (1 - |z|^2) dA(z) + \frac{1}{\pi} \int_{\mathbb{D}} f \bar{h} dA(z).$$

Technically, to prove the above equality, we apply integration by parts over a disc of radius  $r$  and let  $r \rightarrow 1$ , using the fact that  $M_1(r, f) = o((1-r)^{-1})$ .

By writing the integral on the left above in polar coordinates we see that it equals

$$\begin{aligned} & \int_0^1 \frac{1}{2\pi} \int_0^{2\pi} \partial_z(f(re^{i\theta})) \overline{h'(re^{i\theta})} (1+r)r \int_{\theta-(1-r)/2}^{\theta+(1-r)/2} d\phi dr d\theta \\ &= \int_0^1 \frac{1}{2\pi} \int_0^{2\pi} \int_{\phi-(1-r)/2}^{\phi+(1-r)/2} \partial_z(f(re^{i\theta})) \overline{h'(re^{i\theta})} (1+r)r d\theta dr d\phi \\ &= \frac{1}{2\pi} \int_0^{2\pi} \iint_{\Gamma_\phi} \partial_z(f(re^{i\theta})) \overline{h'(re^{i\theta})} (1+r)r dr d\theta d\phi \end{aligned}$$

where the region

$$\Gamma_\phi = \{(r, \theta) : \phi - (1-r)/2 < \theta < \phi + (1-r)/2, 0 < r < 1\}.$$

Now we may apply the Cauchy-Schwarz inequality and the fact that  $r(1+r) \leq 2$  to deduce the following inequality:

$$(2.1) \quad \left| \frac{1}{2\pi} \int_0^{2\pi} \mathcal{P}_n(f) \bar{h} d\theta \right| \leq \frac{2}{2\pi} \int_0^{2\pi} S_z(f) S_z(h) d\theta + \|f\|_{L^{2p/(p+1)}(\mathbb{D})} \|h\|_{A^{2p'}}$$

where

$$S_z(f)(e^{i\theta})^2 = \iint_{\Gamma_\theta} |\partial_z f(re^{i\phi})|^2 dr d\phi.$$

We now see that

$$\begin{aligned} \left| \frac{1}{2\pi} \int_0^{2\pi} \mathcal{P}_n(f) \bar{h} d\theta \right| &\leq 2 \|S_z(f)\|_p \|S_z(h)\|_{p'} + \|f\|_{L^{2p/(p+1)}(\mathbb{D})} \|h\|_{A^{2p'}} \\ &\leq C(\|S_z(f)\|_p + \|f\|_{L^{2p/(p+1)}(\mathbb{D})}) \|h\|_{H^{p'}} \end{aligned}$$

by Theorem A and the fact that  $\|h\|_{A^{2p'}} \leq \|h\|_{H^{p'}}$  [11]. We may replace  $h$  by any polynomial  $\hat{h}$  whose first  $n$  terms agree with  $h$  without changing the value of the left hand side of the above inequality. We also have that  $\|\hat{h}\|_{H^{p'}} \leq C\|h\|_{H^{p'}}$  where  $C$  depends only on  $q$ , due to the fact that Fourier partial summation operators are uniformly bounded on  $L^q$  for  $1 < q < \infty$ . Taking the supremum over all  $h$  with  $\|h\|_{H^{p'}} \leq 1$  thus shows that

$$\sup_{\|h\|_{H^{p'}} \leq 1} \left| \frac{1}{2\pi} \int_0^{2\pi} \mathcal{P}_n(f) \bar{h} d\theta \right| \leq C(\|S_z(f)\|_p + \|f\|_{L^{2p/(p+1)}(\mathbb{D})}).$$

But the boundedness of the Szegő projection (or equivalently the fact that  $(H^p)^*$  is isomorphic to  $H^{p'}$  for  $1 < p < \infty$ ) now implies that

$$\|\mathcal{P}_n(f)\|_p \leq C(\|S_z(f)\|_p + \|f\|_{L^{2p/(p+1)}(\mathbb{D})}).$$

Since integral means are increasing, this means that for any fixed  $r$ ,

$$M_p(r, \mathcal{P}_n(f)) \leq C(\|S_z(f)\|_p + \|f\|_{L^{2p/(p+1)}(\mathbb{D})}).$$

Since  $\mathcal{P}_n(f)$  converges to  $\mathcal{P}(f)$  uniformly on compact subsets, this implies that

$$M_p(r, \mathcal{P}(f)) \leq C(\|S_z(f)\|_p + \|f\|_{L^{2p/(p+1)}(\mathbb{D})}).$$

□

We note that a similar argument to the one above shows that

$$\frac{1}{2\pi} \int_0^{2\pi} P_n(\bar{z}f) \bar{h} d\theta \leq C \|S_z(f)\|_p \|S_z(zh)\|_{p'}$$

for a polynomial  $h$  of degree at most  $n$ , so that

$$\|\mathcal{P}(\bar{z}f)\|_{H^p} \leq C_p \|S_z(f)\|_p.$$

We will not need this in the future, but we note that its proof does not use  $f \in L^{2p/(p+1)}$ , unlike our proof of the theorem.

We next apply Theorem 2.1 to functions  $f$  of the form  $f = F^{p/2} \bar{F}^{(p/2)-1}$  using Theorem A. Note that in this case  $\partial_z(f) = \frac{p}{2}|F|^{p-2}F'$ . Now note that  $\| |F|^{p-2}F' \| = |\nabla(|F|^{p-1})|/(p-1)$  and thus

$$S_z(f) \leq \frac{p/2}{p-1} S(|F|^{p/2}).$$

**Theorem 2.2.** *Let  $k \in A^q$  be not identically 0 and let  $F$  be the Bergman space extremal function for  $k$ . If  $1 \leq p < \infty$  and  $0 < q < \infty$  and  $F \in H^q$  then  $k \in H^{q/(p-1)}$ .*

Combining this with a result in [4] yields the following.

**Corollary 2.3.** *If  $1 \leq p < \infty$  and  $p \leq q < \infty$  then  $F \in H^q$  if and only if  $k \in H^{q/(p-1)}$ .*

### 3. SZEGÖ PROJECTIONS AND EXTREMAL PROBLEMS

We will state our main result for this section. An examination of the proof and the fact that we could compute an estimate for the  $C_{q,p}$  shows that we could compute an estimate for the constants  $\tilde{C}_{q,p}$  in the theorem if we wanted.

**Theorem 3.1.** *Let  $1 < p < \infty$  and  $1 < q < \infty$ . Let  $\tilde{C}_{q,p}$  be as defined in equation (3.1). Suppose  $\tilde{C}_{q,p} < 1$ . Let  $k \in H^{p'} \cap H^q$  and let  $F$  be the  $H^p$  extremal function for  $k$ . Then  $F \in H^{(p-1)q}$  and*

$$\|F\|_{(p-1)q}^{p-1} \leq \frac{1}{1 - \tilde{C}_{p,q}} \frac{\|k\|_q}{\|k\|_{(H^p)^*}}.$$

Before proving the theorem, we mention some consequences. Let  $K$  be the extremal kernel corresponding to  $k$  from the theory of dual extremal problems. Then

$$K = \|k\|_{(H^p)^*} F^{p/2} \overline{F}^{(p/2)-1}.$$

But  $K$  is by definition the solution to the problem of finding the function of smallest  $L^{p'}$  norm of the form  $\bar{k} - g$ , where  $g$  is analytic and vanishes at the origin. Changing  $p'$  to  $p$  gives the following result.

**Lemma 3.2.** *Let  $1 < p < \infty$  and  $1 < q < \infty$ . Suppose that  $\tilde{C}_{q,p'} < 1$  and that  $k \in H^p \cap H^q$  and let  $f \in H^p$  satisfy  $\|f - \bar{k}\|_{H^p} \leq \|g - \bar{k}\|_{H^p}$  for all  $g \in H^p$  that vanish at the origin. Then  $f \in H^q$  and*

$$\|f - \bar{k}\|_{H^q} \leq \frac{1}{1 - \tilde{C}_{q,p'}} \|k\|_{H^q}$$

and thus

$$\|f\|_{H^q} \leq \left( 1 + \frac{1}{1 - \tilde{C}_{q,p'}} \right) \|k\|_{H^q}.$$

Suppose now that  $k \in L^p$  and we want to find the function  $f$  that minimizes  $\|f - k\|_p$ , where  $f$  is in  $H^p$ . This is equivalent to finding the function  $h \in H^p$  which minimizes the norm of  $h - P_S^\perp k$ , where  $P_S^\perp = I - P_S$  is the projection of  $L^2$  onto  $(H^2)^\perp$ . Indeed, we have the

relation  $f = h + P_S(k)$ . But this is equivalent to minimizing the norm of  $g - zP_S^\perp k$ , where  $g$  is analytic and vanishes at the origin. Note that  $zP_S^\perp k$  is an anti-analytic function and that  $g = zh$ . Since then

$$\|g - zP_S^\perp k\|_{H^q} \leq \left(1 + \frac{1}{1 - \tilde{C}_{q,p'}}\right) \|P_S^\perp k\|_{H^q}$$

and  $\|P_S^\perp k\|_q \leq \mathfrak{s}_q \|k\|_q$  and finally  $f = g/z + P_S(k)$  we have the following result.

**Theorem 3.3.** *Let  $1 < p < \infty$  and  $1 < q < \infty$ . Suppose that  $\tilde{C}_{q,p'} < 1$  and that  $k \in L^p \cap L^q$  and let  $f \in H^p$  satisfy  $\|f - k\|_{H^p} \leq \|g - k\|_{H^p}$  for all  $g \in H^p$ . Then  $f \in H^q$  and*

$$\|f\|_{H^q} \leq \left(2 + \frac{1}{1 - \tilde{C}_{q,p'}}\right) \mathfrak{s}_q \|k\|_{L^q}.$$

*Proof of Theorem 3.1.* Let  $F \in H^p$  and let  $f = F^{p/2} \overline{F}^{(p/2)-1}$ . Suppose that  $h$  is analytic and 0 at the origin. For the moment assume that both  $F$  and  $h$  are analytic in the closed unit disc. Then an application of Green's formula [5, VI.3] shows that

$$\frac{1}{2\pi} \int_0^{2\pi} fh \, d\theta = \frac{1}{2\pi} \int_{\mathbb{D}} \Delta(fh) \log(1/|z|) \, dA.$$

Let

$$A = \frac{1}{2\pi} \int_{\frac{1}{4}\mathbb{D}} \Delta(fh) \log(1/|z|) \, dA$$

and

$$B = \frac{1}{2\pi} \int_{\mathbb{D} \setminus \frac{1}{4}\mathbb{D}} \Delta(fh) \log(1/|z|) \, dA.$$

Now note that  $\Delta(fh) = 4(\partial_{\bar{z}}f)h' + \Delta(f)h$ . First note that ,

$$\partial_{\bar{z}}f = ((p/2) - 1)F^{p/2} \overline{F}^{(p/2)-2} \overline{F}'.$$

Now this means that

$$|\partial_{\bar{z}}f| = |(p/2) - 1| |F|^{p-2} |F'| = \frac{|p-2|}{2\sqrt{2}(p-1)} |\nabla|F|^{p-1}|.$$

Also,

$$|\Delta(f)| = 4 \left| \frac{p}{2} - 1 \right| (p/2) |F|^{p-3} |F'|^2.$$

But this equals

$$\frac{p|p-2|}{2(p-1)} |\nabla|F|^{(p-1)/2}|^2$$

So we have that

$$|\Delta(fh)| \leq \frac{|p-2|}{4(p-1)} |\nabla|F|^{p-1}| |\nabla|h|| + \frac{p|p-2|}{2(p-1)} |\nabla|F|^{(p-1)/2}|^2 |h|.$$

Because of the above formulas, the integrability of  $\log(1/|z|)$  and the fact that point evaluation (and point evaluation of derivatives) is a bounded linear functional in all Hardy spaces, there is a constant  $\mathfrak{k}_q$  depending only on  $q$  such that

$$A \leq \frac{p|p-2|}{p-1} \mathfrak{k}_q \|F\|_{(p-1)q}^{p-1} \|k\|_{q'},$$

where the norms are with respect to Hardy spaces. We can also note that

$$|B| \leq \frac{2}{2\pi} \int_{\mathbb{D}} |\Delta(fh)|(1-|z|) dA$$

A calculation similar to that in the last section shows that the right hand side of the above expression equals

$$\frac{2}{2\pi} \int_0^{2\pi} \iint_{\Gamma_\theta} |\Delta(fh)| r d\phi dr d\theta$$

Putting this all together gives

$$\begin{aligned} \frac{1}{2\pi} \int_0^{2\pi} fh d\theta &\leq \frac{p|p-2|}{p-1} \mathfrak{k}_q \|F\|_{(p-1)q}^{p-1} \|h\|_{q'} + \\ &\quad \frac{|p-2|}{4(p-1)} \|S(|F|^{p-1})\|_q \|S(h)\|_{q'} + \\ &\quad \frac{p|p-2|}{2(p-1)} \|S(|F|^{(p-1)/2})^2\|_q \|h^*\|_{q'} \end{aligned}$$

where  $h^*$  is the nontangential maximal function. Thus

$$\begin{aligned} \frac{1}{2\pi} \int_0^{2\pi} fh d\theta &\leq \left( \frac{p|p-2|}{p-1} \mathfrak{k}_q + \frac{p-2}{4(p-1)} C_{q,p-1} C_{q',1} + \right. \\ &\quad \left. \mathfrak{m}_{q'} \frac{p|p-2|}{2(p-1)} C_{2q,(p-1)/2}^2 \right) \|F\|_{(p-1)q}^{p-1} \|h\|_{q'} \end{aligned}$$

where  $\mathfrak{m}_{q'}$  is defined by  $\|h^*\|_{q'} \leq \mathfrak{m}_{q'} \|h\|$ . Applying the above result to dilations allows us to remove the regularity assumptions on  $f$  and  $h$ .

We now show how to apply the above to extremal problems. It is here we will need the assumption  $q > 1$ . Let  $q \geq p$  and suppose that  $f \in H^q$  where  $q \geq p/(p-1)$ . Note that this is equivalent to  $F \in H^{q(p-1)}$ . Note that if  $h$  is analytic and 0 at the origin that

$$\frac{1}{2\pi} \int_0^{2\pi} fh d\theta = \frac{1}{2\pi} \int_0^{2\pi} (f - P_S(f))h d\theta.$$



Let  $\mathfrak{s}_p = \csc(\pi/p)$  be the norm of the Szegő projection from  $L^p$  to  $H^p$ . Now,

$$\begin{aligned} \|f - P_S(f)\|_q &= \sup_{\substack{h \text{ harmonic} \\ \|h\|_{q'} \leq 1}} \frac{1}{2\pi} \int_0^{2\pi} (f - P_S(f))z h \, d\theta \\ &= \sup_{\substack{h \text{ harmonic} \\ \|h\|_{q'} \leq 1}} \frac{1}{2\pi} \int_0^{2\pi} (f - P_S(f))z P_S(h) \, d\theta \\ &\leq \sup_{\substack{h \text{ analytic} \\ \|h\|_{q'} \leq \mathfrak{s}_{q'}}} \frac{1}{2\pi} \int_0^{2\pi} (f - P_S(f))z h \, d\theta \end{aligned}$$

Let

$$(3.1) \quad \tilde{C}_{q,p} = \mathfrak{s}_{q'} \left( \frac{p|p-2|}{p-1} \mathfrak{k}_q + \frac{p-2}{4(p-1)} C_{q,p-1} C_{q',1} + \mathfrak{m}_{q'} \frac{p|p-2|}{2(p-1)} C_{2q,(p-1)/2}^2 \right)$$

where  $\mathfrak{m}$  is constant for the nontangential maximal operator. But our work above now shows that  $\|f - P_S(f)\|_q$  is at most  $\tilde{C}_{q,p} \|f\|_q$ . If this constant  $\tilde{C}_{q,p}$  is less than 1, then we have  $\|P_S f\|_q \geq (1 - \tilde{C}_{q,p}) \|f\|_q$ , as long as  $\|f\|_q$  is finite.

We can now phrase the above result in terms of extremal problems. If  $F$  is the  $H^p$  extremal function for  $k$  and  $\|F\|_{q(p-1)}$  is finite then

$$\|F\|_{q(p-1)}^{p-1} \leq \frac{1}{1 - \tilde{C}_{q,p}} \frac{\|k\|_q}{\|k\|_{(H^p)^*}}.$$

All that is left to do is show that  $F \in H^{(p-1)q}$  if  $k \in H^q$ .

To this end, let  $k_n$  be a sequence of polynomials approaching  $k$  in the  $H^q$  and  $H^{p'}$  norms. We may assume that  $\|k_n\|_q \leq 2\|k\|_q$  for all  $n$  and  $\|k_n\|_{(H^p)^*} \geq \|k\|_{(H^p)^*}/2$  for all  $n$ . It is known that the extremal function  $F_n$  corresponding to  $k_n$  is continuous in the closed unit disc. (In fact, an explicit form is known for such  $F_n$  from the theory of dual extremal problems). Thus for any fixed  $0 \leq r < 1$  we have

$$M_{q(p-1)}^{p-1}(r, F_n) \leq \|F_n\|_{q(p-1)}^{p-1} \leq \frac{4}{1 - \tilde{C}_{q,p}} \frac{\|k\|_q}{\|k\|_{(H^p)^*}}.$$

It is known that  $F_n \rightarrow F$  in  $H^p$  [2], and thus  $F_n \rightarrow F$  uniformly on compact subsets of the unit disc. This means that  $M_{q(p-1)}(r, F_n) \rightarrow$

$M_{q(p-1)}(r, F)$ , and so

$$M_{q(p-1)}^{p-1}(r, F) \leq \frac{4}{1 - \tilde{C}_{q,p}^r} \frac{\|k\|_q}{\|k\|_{(H^p)^*}}.$$

Since  $r$  is arbitrary, this proves that  $F \in H^{q(p-1)}$ .  $\square$

#### REFERENCES

- [1] Peter Duren, *Theory of  $H^p$  spaces*, Pure and Applied Mathematics, Vol. 38, Academic Press, New York, 1970. MR 0268655 (42 #3552)
- [2] Timothy Ferguson, *Continuity of extremal elements in uniformly convex spaces*, Proc. Amer. Math. Soc. **137** (2009), no. 8, 2645–2653. MR 2497477
- [3] Timothy Ferguson, *Extremal problems in Bergman spaces and an extension of Ryabykh’s theorem*, Illinois J. Math. **55** (2011), no. 2, 555–573 (2012). MR 3020696
- [4] ———, *Extremal problems in Bergman spaces and an extension of Ryabykh’s Hardy space regularity theorem for  $1 < p < \infty$* , Indiana Univ. Math. J. **66** (2017), no. 1, 259–274. MR 3623410
- [5] John B. Garnett, *Bounded analytic functions*, Pure and Applied Mathematics, vol. 96, Academic Press Inc. [Harcourt Brace Jovanovich Publishers], New York, 1981. MR 628971 (83g:30037)
- [6] S. Ya. Havinson, *On some extremal problems of the theory of analytic functions*, Amer. Math. Soc. Transl. (2) **32** (1963), 139–154.
- [7] S. Ya. Khavinson, *On an extremal problem of the theory of analytic functions*, Uspehi Matem. Nauk (N.S.) **4** (1949), no. 4(32), 158–159. MR 0033887 (11,508e)
- [8] ———, *On some extremal problems of the theory of analytic functions*, Moskov. Gos. Univ. Učenyje Zapiski Matematika **148(4)** (1951), 133–143. MR 0049322 (14,155f)
- [9] W. W. Rogosinski and H. S. Shapiro, *On certain extremum problems for analytic functions*, Acta Math. **90** (1953), 287–318. MR 0059354
- [10] V. G. Ryabykh, *Extremal problems for summable analytic functions*, Sibirsk. Mat. Zh. **27** (1986), no. 3, 212–217, 226 ((in Russian)). MR 853902 (87j:30058)
- [11] Dragan Vukotić, *The isoperimetric inequality and a theorem of Hardy and Littlewood*, Amer. Math. Monthly **110** (2003), no. 6, 532–536. MR 1984405

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ALABAMA, TUSCALOOSA, AL

*E-mail address:* tjferguson1@ua.edu