

Camassa-Holm and M-CIV equations with self-consistent sources: geometry and peakon solutions

Gulmira Yergaliyeva*, Tolkynay Myrzakul†, Gulgassyl Nugmanova‡
Kuralay Yesmakhanova§ and Ratbay Myrzakulov¶
Eurasian International Center for Theoretical Physics,
Eurasian National University, Nur-Sultan, 010008, Kazakhstan

Abstract

In this paper, we study one of generalized Heisenberg ferromagnet equations with self-consistent sources, namely, the so-called M-CIV equation with self-consistent sources (M-CIVESCS). The Lax representation of the M-CIVESCS is presented. We have shown that the M-CIVESCS and the CH equation with self-consistent sources (CHESCS) is geometrically equivalent each to other. The gauge equivalence between these equations is proved. Soliton (peakon) and pseudo-spherical surfaces induced by these equations are considered. The one peakon solution of the M-CIVESCS is presented.

KEYWORDS: Camassa-Holm equation with self-consistent sources; Heisenberg ferromagnet equation; Heisenberg ferromagnet equation with self-consistent sources; Heisenberg ferromagnet equation with self-consistent potentials; Lax representation; conservation laws; peakon; soliton.

1 Introduction

Camassa-Holm equation (CHE) has the form

$$u_t + 2\omega u_x - u_{xxt} + 3uu_x = 2u_x u_{xx} + uu_{xxx}, \quad (1.1)$$

where $u = u(x, t)$ is the fluid velocity in the x direction and $\omega = const$ is related to the critical shallow water wave speed. This equation has several equivalent forms, for example, the following ones

$$q_t + 2u_x q + u q_x = 0, \quad (1.2)$$

$$q - u + u_{xx} - \omega = 0, \quad (1.3)$$

*Email: gyergaliyeva@gmail.com

†Email: trmyrzakul@gmail.com

‡Email: nugmanovagn@gmail.com

§Email: kryesmakhanova@gmail.com

¶Email: rmyrzakulov@gmail.com

or

$$\kappa_t + (u\kappa)_x = 0, \quad (1.4)$$

$$u - u_{xx} + \omega - \nu\kappa^2 = 0. \quad (1.5)$$

The CHE was implicitly contained in the class of multi-Hamiltonian system introduced by Fuchssteiner and Fokas [1]. It explicitly derived as a shallow water wave equation by Camassa and Holm [2] and after their works, the CHE and its different modifications have been studied from many kinds of views [3]-[18]. The CHE shares most of the properties of the integrable system of KdV type and possesses Lax representation (LR), the bi-Hamiltonian structure, smooth solitary wave solutions (as $\omega > 0$). For example, the solitary wave solutions become piecewise smooth and have cusps at their peaks when $\omega \rightarrow 0$. When $\omega = 0$, these kind of solutions are weak solutions and are called "peakons".

Integrable Heisenberg ferromagnet type equations play important role in modern physics and mathematics (see, e.g., Refs. [20]-[61]). Recently, integrable generalized Heisenberg ferromagnet equations which are equivalent to Camassa-Holm type equations were presented (see, e.g., refs. [20]-[25]). In particular, it is shown that the CHE is (geometrically and gauge) equivalent to the following Myrzakulov-CIV (M-CIV) equation [20]-[27]

$$A_{xt} + (uA_x)_x + (uA_x^2 + 0.5\{A_x, A_t\})A - \frac{1}{4\beta^2}[A, A_x] = 0, \quad (1.6)$$

$$\text{tr}(A_x^2) + 8\beta^2(u - u_{xx}) = 0 \quad (1.7)$$

or

$$(A_t + uA_x)_x + (u_{xx} - u_x - 2u)A - \frac{1}{4\beta^2}[A, A_x] = 0, \quad (1.8)$$

$$\text{tr}(A_x^2) + 8\beta^2(u - u_{xx}) = 0. \quad (1.9)$$

Here

$$A = \begin{pmatrix} A_3 & A^- \\ A^+ & -A_3 \end{pmatrix}, \quad A^\pm = A_1 \pm iA_2, \quad A^2 = I, \quad \mathbf{A} = (A_1, A_2, A_3), \quad \mathbf{A}^2 = 1, \quad (1.10)$$

$$A_x^2 = -4\beta^2 qI, \quad \{A_t, A_x\} = [(8\beta^2 u - 4)q - 2(u_x + u_{xx})]I. \quad (1.11)$$

There are many integrable generalizations of the CHE. One of such generalizations is the CHE with self-consistent sources (CHESCS) [3]. Such type integrable equations with self-consistent sources have attracted much attention in recent years [15]. They have important applications in many branches of physics. For example, the nonlinear Schrödinger equation with self-consistent sources describes the nonlinear interaction of an electrostatic high-frequency wave with the ion acoustic wave in a two component homogeneous plasma. Another example is the KdV equation with self-consistent sources. It describes the interaction of long and short capillary-gravity waves. The famous KP equation with self-consistent sources represents the nonlinear interaction of a long wave with a short wave packet propagating on the x - y plane at some angle to each other. In this paper, we would like to study the M-CIV equation with self-consistent sources (M-CIVESCS) and its relation with the CHESCS.

This paper is organized as follows. In Section 2, we present the M-CIVESCS and its Lax representation. In Section 3, the integrable motion of curves induced by the M-CIVESCS are

constructed. The gauge equivalence between the M-CIVESCS and CHESCS is established in Section 4. The peakon (soliton) and pseudo-spherical surfaces induced by the M-CIVESCS and the CHESCS are presented in Section 5 and in Section 6, respectively. In Section 7, the formulas of the one peakon solution of the M-CIVESCS is presented. The M-CIVESCS and the CHESCS with N self-consistent sources is given in Section 8. In Section 9, the conclusion is presented.

2 M-CIV equation with self-consistent sources

One of examples of the peakon spin systems is the following Myrzakulov-CIV equation with self-consistent sources (M-CIVESCS)

$$[A, A_{xt}] + (u[A, A_x])_x - \frac{1}{\beta^2}A_x - \alpha(\omega[A, A_x])_x + \frac{2\alpha^2\chi}{\beta^2 + \alpha}A_x + \frac{2\alpha\beta^2\chi_x}{\beta^2 + \alpha}A = 0, \quad (2.12)$$

$$\psi_{1x} - \left(\frac{\alpha}{4\beta} - \frac{1}{4}\right)[(A_x^+ A^- - A^+ A_x^-)\psi_1 + 2(A_x^- A_3 - A^- A_{3x})\psi_2] = 0, \quad (2.13)$$

$$\psi_{2x} - \left(\frac{\alpha}{4\beta} - \frac{1}{4}\right)[2(A^+ A_{3x} - A_x^+ A_3)\psi_1 + (A^+ A_x^- - A_x^+ A^-)\psi_2] = 0. \quad (2.14)$$

This equation can be written in the following equivalent form

$$A_{xt} + uA_{xx} + (uA_x^2 + 0.5\{A_x, A_t\})A + u_1A_x + u_3I + u_2[A, A_x] = 0, \quad (2.15)$$

$$\psi_{1x} - \left(\frac{\alpha}{4\beta} - \frac{1}{4}\right)[(A_x^+ A^- - A^+ A_x^-)\psi_1 + 2(A_x^- A_3 - A^- A_{3x})\psi_2] = 0, \quad (2.16)$$

$$\psi_{2x} - \left(\frac{\alpha}{4\beta} - \frac{1}{4}\right)[2(A^+ A_{3x} - A_x^+ A_3)\psi_1 + (A^+ A_x^- - A_x^+ A^-)\psi_2] = 0, \quad (2.17)$$

where $\omega = \phi_1^2$, $\chi = \omega_x + \omega$, $\alpha, \beta = \text{consts}$ and

$$u_1 = u - \alpha\omega, \quad u_2 = \frac{\alpha^2\chi}{\beta^2 + \alpha} - \frac{1}{4\beta^2}, \quad u_3 = \frac{\alpha\beta^2\chi_x}{\beta^2 + \alpha}, \quad (2.18)$$

$$A = \begin{pmatrix} A_3 & A^- \\ A^+ & -A_3 \end{pmatrix}, \quad A^\pm = A_1 \pm iA_2, \quad A^2 = I, \quad \mathbf{A} = (A_1, A_2, A_3), \quad \mathbf{A}^2 = 1. \quad (2.19)$$

The LR of the M-CIVESCS reads as

$$\Psi_x = U_1\Psi, \quad (2.20)$$

$$\Psi_t = V_1\Psi. \quad (2.21)$$

Here

$$U_1 = \left(\frac{\lambda}{4\beta} - \frac{1}{4}\right)[A, A_x], \quad (2.22)$$

$$V_1 = \left(\frac{1}{4\beta^2} - \frac{1}{4\lambda^2}\right)A + \left[\frac{1}{8\beta\lambda} - \frac{1}{8\beta^2} - \left(\frac{\lambda}{4\beta} - \frac{1}{4}\right)u\right][A, A_x] + \left(\frac{1}{2\lambda} - \frac{1}{2\beta}\right)Z, \quad (2.23)$$

where

$$Z = \frac{1}{2} \left[\frac{u_x + u_{xx}}{\beta} - \frac{\alpha\beta\chi_x}{\beta^2 + \alpha} \right]^{-1} \left[A, A_t - \left(\frac{1}{2\beta} - u + \frac{\alpha\beta\omega}{\beta^2 + \alpha} \right) A_x \right]. \quad (2.24)$$

3 Integrable motion of space curves induced by the M-CIVESCS

In this section, we consider the integrable motion of space curves induced by the M-CIVESCS. As usual, let us consider a smooth space curve $\gamma(x, t) : [0, X] \times [0, T] \rightarrow R^3$ in R^3 . Let x is the arc length of the curve at each time t . The corresponding Frenet-Serret equation and its temporal counterpart look like

$$\begin{pmatrix} \mathbf{e}_1 \\ \mathbf{e}_2 \\ \mathbf{e}_3 \end{pmatrix}_x = C \begin{pmatrix} \mathbf{e}_1 \\ \mathbf{e}_2 \\ \mathbf{e}_3 \end{pmatrix}, \quad \begin{pmatrix} \mathbf{e}_1 \\ \mathbf{e}_2 \\ \mathbf{e}_3 \end{pmatrix}_t = G \begin{pmatrix} \mathbf{e}_1 \\ \mathbf{e}_2 \\ \mathbf{e}_3 \end{pmatrix}, \quad (3.25)$$

where \mathbf{e}_j are the unit tangent vector ($j = 1$), principal normal vector ($j = 2$) and binormal vector ($j = 3$) which given by $\mathbf{e}_1 = \gamma_x$, $\mathbf{e}_2 = \frac{\gamma_{xx}}{|\gamma_{xx}|}$, $\mathbf{e}_3 = \mathbf{e}_1 \wedge \mathbf{e}_2$, respectively. Here

$$C = \begin{pmatrix} 0 & \kappa_1 & \kappa_2 \\ -\kappa_1 & 0 & \tau \\ -\kappa_2 & -\tau & 0 \end{pmatrix}, \quad G = \begin{pmatrix} 0 & \omega_3 & \omega_2 \\ -\omega_3 & 0 & \omega_1 \\ -\omega_2 & -\omega_1 & 0 \end{pmatrix}, \quad (3.26)$$

where τ , κ_1, κ_2 are the "torsion", "geodesic curvature" and "normal curvature" of the curve, respectively; ω_j are some functions. The compatibility condition of the equations (3.25) reads as

$$C_t - G_x + [C, G] = 0 \quad (3.27)$$

or in elements

$$\kappa_{1t} - \omega_{3x} - \kappa_2\omega_1 + \tau\omega_2 = 0, \quad (3.28)$$

$$\kappa_{2t} - \omega_{2x} + \kappa_1\omega_1 - \tau\omega_3 = 0, \quad (3.29)$$

$$\tau_t - \omega_{1x} - \kappa_1\omega_2 + \kappa_2\omega_3 = 0. \quad (3.30)$$

We now assume $\mathbf{A} \equiv \mathbf{e}_1$. Let take place the following expressions

$$\kappa_1 = i, \quad \kappa_2 = \lambda(q - 1), \quad \tau = -i\lambda(q + 1), \quad (3.31)$$

where $q = 0.5\lambda(\kappa_2 + i\tau)$. Then we have

$$\omega_1 = i[(0.5\lambda^{-1} - \lambda u)(q + 1) + 0.5\lambda^{-2}(u_x + u_{xx})], \quad (3.32)$$

$$\omega_2 = [(0.5\lambda^{-1} - \lambda u)(q + 1) + 0.5\lambda^{-1}(u_x + u_{xx})], \quad (3.33)$$

$$\omega_3 = i[0.5\lambda^{-2} - u - u_x]. \quad (3.34)$$

Now Eqs.(3.28)-(3.30) give us the following equations for q, u, ϕ_1 :

$$q_t + 2qu_x + uq_x - \omega_x + \omega_{xxx} = 0, \quad (3.35)$$

$$\phi_{1xx} - (\alpha^2 q + \frac{1}{4})\phi_1 = 0 \quad (3.36)$$

or

$$q_t + 2qu_x + uq_x - [(\phi_1^2)_x - (\phi_1^2)_{xxx}] = 0, \quad (3.37)$$

$$\phi_{1xx} - (\alpha^2 q + \frac{1}{4})\phi_1 = 0. \quad (3.38)$$

It is nothing but the CHESCS [3]-[4]. So, we have proved that the M-CIVESCS is the Lakshmanan (geometrical) equivalent to the CHESCS.

4 Gauge equivalence between the M-CIVESCS and the CHESCS

Above, we have shown that the M-CIVESCS and the CHESCS are the geometrical (Lakshmanan) equivalent each to other. Now we consider the possible gauge equivalence between these equations [26]. First, we note that from the results of the previous section and from the isomorphism $so(3) \approx su(2)$ follow the LR for the CHESCS of the form [3]

$$\Phi_x = U_2 \Phi, \quad (4.39)$$

$$\Phi_t = V_2 \Phi, \quad (4.40)$$

where

$$\Phi = \begin{pmatrix} \phi_1 \\ \phi_2 \end{pmatrix}, \quad U_2 = \begin{pmatrix} -0.5 & \lambda \\ \lambda q & 0.5 \end{pmatrix}, \quad V_2 = V_{20} + V_2'. \quad (4.41)$$

Here

$$V_{20} = \begin{pmatrix} \frac{u_x + u_{xx}}{2} - \frac{1}{4\lambda^2} & \frac{1}{2\lambda} - u\lambda \\ \frac{u_x + u_{xx} + q}{2\lambda} - uq\lambda & \frac{1}{4\lambda^2} - \frac{u_x + u_{xx}}{2} \end{pmatrix}, \quad (4.42)$$

$$V_2' = -\frac{\alpha\lambda^2\chi}{2(\lambda^2 + \alpha)}\sigma_3 + \frac{\alpha\lambda^3\omega}{\lambda^2 + \alpha} \begin{pmatrix} 0 & 1 \\ q & 0 \end{pmatrix} - \frac{\alpha\lambda\chi_x}{2(\lambda^2 + \alpha)}\Sigma, \quad \Sigma = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}. \quad (4.43)$$

The compatibility condition $\Phi_{xt} = \Phi_{tx}$ given by

$$U_{2t} - V_{2x} + [U_2, V_2] = 0 \quad (4.44)$$

is equivalent to the CHESCS (3.35)-(3.36). Consider the transformation $\Psi = g^{-1}\Phi$, where Ψ is the solution of the equations (2.20)-(2.21) and $g = \Phi|_{\lambda=\beta}$. Then the Lax pairs of the M-CIVESCS and CHESCS is related by the following equations

$$U_1 = g^{-1}U_2g - g^{-1}g_x, \quad V_1 = g^{-1}V_2g - g^{-1}g_t. \quad (4.45)$$

Note that these LR can be rewritten in the equivalent scalar forms. For example, the equivalent scalar form of the LR for the CHESCS is given by [3]

$$\phi_{1xx} = \left(\lambda^2 q + \frac{1}{4}\right)\phi_1, \quad (4.46)$$

$$\phi_{1t} = \left(\frac{1}{2\lambda^2} - u + \frac{\alpha\lambda^2\omega}{\lambda^2 + \alpha}\right)\phi_{1x} + \left(\frac{1}{2}u_x - \frac{\alpha\lambda^2\omega_x}{2(\lambda^2 + \alpha)}\right)\phi_1. \quad (4.47)$$

The compatibility condition $\phi_{1xxt} = \phi_{1txx}$ is equivalent to the CHESCS (3.35)-(3.36). Finally we present the following important relation between the solutions of the M-CIVESCS and the CHESCS:

$$tr(A_x^2) = -8\beta^2 q = -8\beta^2(u - u_{xx}) \quad (4.48)$$

or

$$\mathbf{A}_x^2 = -4\beta^2 q = -4\beta^2(u - u_{xx}). \quad (4.49)$$

5 Peakon (soliton) surfaces

5.1 Peakon surfaces corresponding to the CHESCS

As well-known, the Sym-Tafel formula gives an interesting connection between the classical geometry of manifolds immersed in R^n and the integrable systems. Using the Sym-Tafel formula, here we want to construct the soliton (peakon) surface induced by the CHESCS and the M-CIVESCS. To this end, let us consider a λ -family of parametric surfaces given by the matrix $r = r(x, t, \lambda)$. According to the Sym-Tafel formula, this matrix defines as

$$r = \Phi^{-1}\Phi_\lambda, \quad (5.50)$$

where Φ is the solution of the equations (4.39)-(4.40). We have

$$r_x = \Phi^{-1}U_{2\lambda}\Phi, \quad r_t = \Phi^{-1}V_{2\lambda}\Phi. \quad (5.51)$$

Then the components of the metric tensor define as

$$g_{ij} = \mathbf{r}_{,i} \cdot \mathbf{r}_{,j}, \quad (5.52)$$

where $\mathbf{r} = (r_1, r_2, r_3)$ is the position vector, $\mathbf{r}_{,1} \equiv \mathbf{r}_x$, $\mathbf{r}_{,2} \equiv \mathbf{r}_t$. Then for example, the first fundamental form of the soliton surface is given by

$$I = g_{ij}dx^i dx^j = \mathbf{r}_x^2 dx^2 + 2\mathbf{r}_x \mathbf{r}_t dx dt + \mathbf{r}_t^2 dt^2, \quad (5.53)$$

where

$$\mathbf{r}_x^2 = \frac{1}{2}tr(r_x^2), \quad \mathbf{r}_x \mathbf{r}_t = \frac{1}{2}tr(r_x r_t), \quad \mathbf{r}_t^2 = \frac{1}{2}tr(r_t^2). \quad (5.54)$$

For example,

$$g_{11} = \mathbf{r}_x^2 = q. \quad (5.55)$$

Similarly, we can construct the second fundamental form in the standard way (see, e.g., [19]).

5.2 Peakon surfaces corresponding to the M-CIVESCS

Let us now we present the main elements of the soliton (peakon) surfaces induced by the M-CIVESCS. In this case, the matrix r is given by

$$r = \Psi^{-1}\Psi_\lambda. \quad (5.56)$$

Here Ψ is the solution of the equations (2.20)-(2.21). These equations give

$$r_x = \Psi^{-1}U_{1\lambda}\Psi, \quad r_t = \Psi^{-1}V_{1\lambda}\Psi. \quad (5.57)$$

Then the components of the metric tensor is given by the formula (5.52). For example,

$$g_{11} = -\frac{1}{8\beta^2}tr(A_x^2). \quad (5.58)$$

Now it is not difficult to construct the first and second fundamental forms of the peakon (soliton) surfaces that we can do in the standard way (see, e.g., [19]).

6 Pseudo-spherical surfaces

In this section, we briefly present the main facts on the pseudo-spherical surfaces induced by the M-CIVESCO and the CHESCO.

6.1 Pseudo-spherical surfaces related with the CHESCO

Consider the following linear problem

$$d\Phi = Y_2\Phi, \quad (6.59)$$

where

$$Y_2 = \frac{1}{2} \begin{pmatrix} \omega^2 & \omega^1 - \omega^3 \\ \omega^1 + \omega^3 & -\omega^2 \end{pmatrix} = U_2 dx + V_2 dt. \quad (6.60)$$

The integrable condition of the 1-form Y_2 is given by

$$d^2 Y_2 = (U_{2t} - V_{2x} + [U_2, V_2]) dt \wedge dx = 0. \quad (6.61)$$

This means that the one-forms ω_j satisfy the following structure equations

$$d\omega^1 = \omega^3 \wedge \omega^2, \quad (6.62)$$

$$d\omega^2 = \omega^1 \wedge \omega^3, \quad (6.63)$$

$$d\omega^3 = \omega^1 \wedge \omega^2. \quad (6.64)$$

These equations are equivalent to the CHESCO.

6.2 Pseudo-spherical surfaces related with the M-CIVESCO

We now return to the M-CIVESCO. To construct the pseudo-spherical surfaces induced by this equation, let us consider the linear problem

$$d\Psi = Y_1\Psi, \quad (6.65)$$

where the 1-form Y_1 reads as

$$Y_1 = \frac{1}{2} \begin{pmatrix} \sigma^2 & \sigma^1 - \sigma^3 \\ \sigma^1 + \sigma^3 & -\sigma^2 \end{pmatrix} = U_1 dx + V_1 dt. \quad (6.66)$$

As in the previous subsection, we consider the integrable condition of the 1-form Y_1 :

$$d^2 Y_1 = (U_{1t} - V_{1x} + [U_1, V_1]) dt \wedge dx = 0 \quad (6.67)$$

which in components takes the form

$$d\sigma^1 = \sigma^3 \wedge \sigma^2, \quad (6.68)$$

$$d\sigma^2 = \sigma^1 \wedge \sigma^3, \quad (6.69)$$

$$d\sigma^3 = \sigma^1 \wedge \sigma^2. \quad (6.70)$$

These equations is equivalent to the M-CIVESCO. Thus the M-CIVESCO and the CHESCO describe some kind pseudo-spherical surfaces as their analogies without sources (see, e.g., refs. [16]-[17]).

7 One peakon solution of the M-CIV ESCS

As the integrable equation, the M-CIVESCS has all ingredients of integrable systems like LR, conservation laws, bi-Hamiltonian structure, soliton solutions and so on. In particular, it admits the peakon solutions. Here let us present a one peakon solution of the M-CIVESCS. To construct this 1-peakon solution, we use the corresponding 1-peakon solution of the CHESCS [3]. The 1-peakon solution of the M-CIVESCS has the form

$$A^+ = \frac{2g_1g_2}{|g_1|^2 + |g_2|^2}, \quad A_3 = \frac{|g_1|^2 - |g_2|^2}{|g_1|^2 + |g_2|^2}, \quad (7.71)$$

$$\psi_1 = \frac{\bar{g}_1\phi_1 + \bar{g}_2\phi_2}{|g_1|^2 + |g_2|^2}, \quad \psi_2 = \frac{-g_2\phi_1 + g_1\phi_2}{|g_1|^2 + |g_2|^2}. \quad (7.72)$$

Here

$$\phi_2 = \lambda^{-1}(\phi_{1x} + 0.5\phi_1), \quad g_j = \phi_j|_{\lambda=\beta}, \quad (7.73)$$

where

$$\phi_1 = \sqrt{\sigma'(t)}ce^{-\frac{1}{2}|x-ct+\sigma(t)|}, \quad u = ce^{-|x-ct+\sigma(t)|} \quad (7.74)$$

is the 1-peakon solution of the CHESCS [3].

8 Integrable self-consistent N sources case

In the previous sections, we have considered the M-CIVESCS and the CHESCS with the one self-consistent source. In this section, we present, in short form, integrable generalizations of these equations with N self-consistent sources.

8.1 M-CIV equation with self-consistent N -sources

The M-CIVE with N self-consistent sources has the form

$$[A, A_{xt} + (uA_x)_x] - \frac{1}{\beta^2}A_x - \sum_{j=1}^N \left(\lambda_j(\omega_j[A, A_x])_x + \frac{2\lambda_j^2\chi_j}{\beta^2 + \lambda_j}A_x + \frac{2\lambda_j\beta^2\chi_{jx}}{\beta^2 + \lambda_j}A \right) = 0, \quad (8.75)$$

$$\psi_{1jx} - \left(\frac{\lambda_j}{4\beta} - \frac{1}{4} \right) [(A_x^+ A^- - A^+ A_x^-) \psi_{1j} + 2(A_x^- A_3 - A^- A_{3x}) \psi_{2j}] = 0, \quad (8.76)$$

$$\psi_{2jx} - \left(\frac{\lambda_j}{4\beta} - \frac{1}{4} \right) [2(A^+ A_{3x} - A_x^+ A_3) \psi_{1j} + (A^+ A_x^- - A_x^+ A^-) \psi_{2j}] = 0, \quad (8.77)$$

where $\omega_j = \phi_{1j}^2$, $\chi_j = \omega_{jx} + \omega_j$, $\alpha, \beta = \text{consts}$ and

$$A = \begin{pmatrix} A_3 & A^- \\ A^+ & -A_3 \end{pmatrix}, \quad A^\pm = A_1 \pm iA_2, \quad A^2 = I, \quad \mathbf{A} = (A_1, A_2, A_3), \quad \mathbf{A}^2 = 1. \quad (8.78)$$

The LR of the M-CIVESCS reads as

$$\bar{\Psi}_x = U_1 \bar{\Psi}, \quad (8.79)$$

$$\bar{\Psi}_t = V_1 \bar{\Psi}. \quad (8.80)$$

Here

$$U_1 = \left(\frac{\lambda}{4\beta} - \frac{1}{4} \right) [A, A_x], \quad (8.81)$$

$$V_1 = \left(\frac{1}{4\beta^2} - \frac{1}{4\lambda^2} \right) A + \left[\frac{1}{8\beta\lambda} - \frac{1}{8\beta^2} - \left(\frac{\lambda}{4\beta} - \frac{1}{4} \right) u \right] [A, A_x] + \left(\frac{1}{2\lambda} - \frac{1}{2\beta} \right) \sum_{j=1}^N Z_j, \quad (8.82)$$

where

$$Z_j = \frac{1}{2} \left[\frac{u_x + u_{xx}}{\beta} - \frac{\lambda_j \beta \chi_{jx}}{\beta^2 + \lambda_j} \right]^{-1} \left[A, A_t - \left(\frac{1}{2\beta} - u + \frac{\lambda_j \beta \omega_j}{\beta^2 + \lambda_j} \right) A_x \right]. \quad (8.83)$$

8.2 CHE with self-consistent N -sources

The CHE with N self-consistent sources has the form [3]

$$q_t + 2qu_x + uq_x - \sum_{j=1}^N [(\varphi_j^2)_x - (\varphi_j^2)_{xxx}] = 0, \quad (8.84)$$

$$\varphi_{j,xx} - (\lambda_j q + \frac{1}{4}) \varphi_j = 0. \quad (8.85)$$

Its LR reads as [3]

$$\phi_{xx} = \left(\frac{1}{4} + \lambda q \right) \phi, \quad (8.86)$$

$$\phi_t = \frac{u_x}{2} \phi + \left(\frac{1}{2\lambda} - u \right) \phi_x + 2 \sum_{j=1}^N \frac{\lambda \lambda_j \phi_j}{\lambda - \lambda_j} (\phi_{jx} \phi - \phi_j \phi_x), \quad (8.87)$$

which means that the CHESCS is Lax integrable [3].

9 Conclusion

The integrable generalized Heisenberg ferromagnet equation with self-consistent sources, namely, the M-CIVESCS is investigated. The integrable motion of space curves induced by the M-CIVESCS is constructed. Using this result, the geometrical equivalence between the M-CIVESCS and the CHESCS is established. It is shown that the M-CIVESCS and the CHESCS is gauge equivalent each to other. The simplest conservation law and the one peakon solution are constructed. The peakon (soliton) surfaces induced by the M-CIVESCS and the CHESCS are presented.

Acknowledgements

This work was supported by the Ministry of Education and Science of Kazakhstan under grants 011800935 and 011800693.

References

- [1] Fuchssteiner B., Fokas A.S. *Physica D*, **4**, 47 (1981)
- [2] Camassa R., Holm D. *Phys. Rev. Lett.*, **71**, 1661 (1993)
- [3] Huang Y., Yao Y., Zeng Y. *On Camassa-Holm equation with self-consistent sources and its solutions*, [arXiv:0811.2552]
- [4] Baltueva I.I., Urazboev G.U. *About the Camassa-Holm equation with a self-consistent source*, Ufa mathematical journal, **3**, N2, 10-18 (2011)
- [5] Yehui Huang. *On Multi-Soliton Solution of Degasperis-Procesi Equation with Self-Consistent Sources*, International Journal of Nonlinear Science, **18**, No.3, 217-223 (2014)
- [6] Yehui Huang, Yunbo Zeng, Orlando Ragnisco. *The Degasperis-Procesi equation with self-consistent sources*, [arXiv:0807.0085]
- [7] Qi Li, Wen Zhang, Qiu-Yuan Duan, Deng-Yuan Chen. *The transformation between the AKNS hierarchy and the KN hierarchy with self-consistent sources*, Journal of Nonlinear Mathematical Physics, **18**, No.4, 483490 (2011)
- [8] Runliang Lin, Haishen Yao, Yunbo Zeng. *Restricted Flows and the Soliton Equation with Self-Consistent Sources*, SIGMA, **2**, 096 (2006)
- [9] Yuqin Yao, Yehui Huang, Yunbo Zeng. *The two-component Camassa-Holm equation with self-consistent sources and its multisoliton solutions*, Theoretical and Mathematical Physics, **162**, No1, 63-73 (2010)
- [10] Wen-Xiu Ma, Ruguang Zhou. *A coupled AKNS-Kaup-Newell soliton hierarchy*, Journal of Mathematical Physics, (1999).[arXiv:solv-int/9908005]
- [11] Yuqin Yao, Yunbo Zeng. *Integrable Rosochatius deformations of higher-order constrained flows and the soliton hierarchy with self-consistent sources*, [arXiv:0806.2251]
- [12] Zhihua Yang, Ting Xiao, Yunbo Zeng. *Integrable dispersionless KdV hierarchy with sources*, [arXiv:0606047]
- [13] Yuqin Yao, Yehui Huang, Guixiang Dong, Yunbo Zeng. *The new integrable deformations of short pulse equation and sine-Gordon equation, and their solutions* , [arXiv:1012.0462]
- [14] Yuqin Yao, Yunbo Zeng. *The generalized Kupershmidt deformation for constructing new integrable systems from integrable bi-Hamiltonian systems*, [arXiv:1005.0281]
- [15] Myrzakulov R. *Soliton Equations with Self-Consistent Sources*, [arXiv:1409.0486]

- [16] Reyes E.G. *Geometric integrability of the Camassa-Holm Equation*, Letters in Mathematical Physics, **59**, No2, 117131 (2002)
- [17] Reyes E.G. *Pseudo-potentials, nonlocal symmetries and integrability of some shallow water equations*, [arXiv:nlin/0212045]
- [18] Ivanov R., Lyons T., Orr N. *A dressing method for soliton solutions of the Camassa-Holm equation*, AIP Conference Proceedings 1895, 040003 (2017): <https://doi.org/10.1063/1.5007370>
- [19] Cieslinski J. *The Darboux-Bianchi-Backlund transformation and soliton surfaces*, [arXiv:1303.5472]
- [20] Mussatayeva A., Myrzakul T., Nugmanova G., Yesmakhanova K. *Gauge equivalence between the Myrzakulov-CIV equation and the Camassa-Holm Equation*
- [21] Mussatayeva A., Myrzakul T., Nugmanova G., Yesmakhanova K., Myrzakulov R. *Integrable Motion of Curves, Spin Equation and Camassa-Holm Equation*, [arXiv:1907.10910]
- [22] Kutum B. *Gauge equivalence between the M-CV equation and the two-component Camassa-Holm equation*
- [23] Taishiyeva A., Myrzakul T., Nugmanova G., Yesmakhanova K. *On the Gauge Equivalence between the 2-component Camassa-Holm and Myrzakulov-CVI equations*
- [24] Taishiyeva A., Myrzakul T., Nugmanova G., Yesmakhanova K., Myrzakulov R. *Geometric Flows of Curves, Two-Component Camassa-Holm Equation and Generalized Heisenberg Ferromagnet Equation*
- [25] Kutum B., Nugmanova G., Myrzakul T., Yesmakhanova K., Myrzakulov R. *Integrable Deformation of Space Curves, Generalized Heisenberg Ferromagnet Equation and Two-Component Modified Camassa-Holm Equation*, [arXiv:1908.01371]
- [26] Yergaliyeva G., Yesmakhanova K., Nugmanova G., Myrzakul T. *Gauge equivalence between the Myrzakulov-CIV equation with self-consistent sources and Camassa-Holm equation with self-consistent sources*
- [27] Mussatayeva A., Myrzakul T., Nugmanova G., Yesmakhanova K., Myrzakulov R. *Integrable Motion of Curves, Spin Equation and Camassa-Holm Equation*, [arXiv:1907.10910]
- [28] G. Nugmanova, Z. Zhunussova, K. Yesmakhanova, G. Mamyrbekova, R. Myrzakulov. International Journal of Mathematical, Computational, Statistical, Natural and Physical Engineering, **9**, N8, 328-331 (2015).
- [29] U. Saleem, M. Hasan. J. Phys. A: Math. Theor., **43**, 045204 (2010).
- [30] R. Myrzakulov, S. Vijayalakshmi, G. Nugmanova, M. Lakshmanan Physics Letters A, **233**, 14-6, 391-396 (1997).
- [31] R. Myrzakulov, S. Vijayalakshmi, R. Syzdykova, M. Lakshmanan, J. Math. Phys., **39**, 2122-2139 (1998).

- [32] R. Myrzakulov, M. Lakshmanan, S. Vijayalakshmi, A. Danlybaeva, J. Math. Phys., **39**, 3765-3771 (1998).
- [33] Myrzakulov R., Danlybaeva A.K., Nugmanova G.N. Theoretical and Mathematical Physics, V.118, 13, P. 441-451 (1999).
- [34] Myrzakulov R., Nugmanova G., Syzdykova R. Journal of Physics A: Mathematical & Theoretical, V.31, 147, P.9535-9545 (1998).
- [35] Myrzakulov R., Daniel M., Amuda R. Physica A., V.234, 13-4, P.715-724 (1997).
- [36] Myrzakulov R., Makhankov V.G., Pashaev O.?. Letters in Mathematical Physics, V.16, N1, P.83-92 (1989)
- [37] Myrzakulov R., Makhankov V.G., Makhankov A. Physica Scripta, V.35, N3, P. 233-237 (1987)
- [38] Myrzakulov R., Pashaev O.K., Kholmurodov Kh. Physica Scripta, V.33, N4, P. 378-384 (1986)
- [39] Anco S.C., Myrzakulov R. Journal of Geometry and Physics, v.60, 1576-1603 (2010)
- [40] Myrzakulov R., Rahimov F.K., Myrzakul K., Serikbaev N.S. *On the geometry of stationary Heisenberg ferromagnets*. In: "Non-linear waves: Classical and Quantum Aspects", Kluwer Academic Publishers, Dordrecht, Netherlands, P. 543-549 (2004)
- [41] Myrzakulov R., Serikbaev N.S., Myrzakul Kur., Rahimov F.K. *On continuous limits of some generalized compressible Heisenberg spin chains*. Journal of NATO Science Series II. Mathematics, Physics and Chemistry, V 153, P. 535-542 (2004)
- [42] R.Myrzakulov, G. K. Mamyrbekova, G. N. Nugmanova, M. Lakshmanan. Symmetry, **7**(3), 1352-1375 (2015). [arXiv:1305.0098]
- [43] R.Myrzakulov, G. K. Mamyrbekova, G. N. Nugmanova, K. Yesmakhanova, M. Lakshmanan. Physics Letters A, **378**, N30-31, 2118-2123 (2014). [arXiv:1404.2088]
- [44] Myrzakulov R., Martina L., Kozhamkulov T.A., Myrzakul Kur. *Integrable Heisenberg ferromagnets and soliton geometry of curves and surfaces*. In book: "Nonlinear Physics: Theory and Experiment. II". World Scientific, London, P. 248-253 (2003)
- [45] Myrzakulov R. *Integrability of the Gauss-Codazzi-Mainardi equation in 2+1 dimensions*. In "Mathematical Problems of Nonlinear Dynamics", Proc. of the Int. Conf. "Progress in Non-linear sciences", Nizhny Novgorod, Russia, July 2-6, 2001, V.1, P.314-319 (2001)
- [46] Chen Chi, Zhou Zi-Xiang. *Darboux Transformation and Exact Solutions of the Myrzakulov-I Equations*. Chin. Phys. Lett., **26**, N8, 080504 (2009)
- [47] Chen Hai, Zhou Zi-Xiang. *Darboux Transformation with a Double Spectral Parameter for the Myrzakulov-I Equation*. Chin. Phys. Lett., **31**, N12, 120504 (2014)
- [48] Zhao-Wen Yan, Min-Ru Chen, Ke Wu, Wei-Zhong Zhao. J. Phys. Soc. Jpn., **81**, 094006 (2012)
- [49] Yan Zhao-Wen, Chen Min-Ru, Wu Ke, Zhao Wei-Zhong. Commun. Theor. Phys., **58**, 463-468 (2012)

- [50] K.R. Ysmakhanova, G.N. Nugmanova, Wei-Zhong Zhao, Ke Wu. *Integrable inhomogeneous Lakshmanan-Myrzakulov equation*, [nlin/0604034]
- [51] Zhen-Huan Zhang, Ming Deng, Wei-Zhong Zhao, Ke Wu. *On the integrable inhomogeneous Myrzakulov-I equation*, [arXiv: nlin/0603069]
- [52] Martina L, Myrzakul Kur., Myrzakulov R, Soliani G. *Journal of Mathematical Physics*, V.42, 13, P.1397-1417 (2001).
- [53] Xiao-Yu Wu, Bo Tian, Hui-Ling Zhen, Wen-Rong Sun and Ya Sun. *Journal of Modern Optics*, 2015.
- [54] Z.S. Yersultanova, M. Zhassymbayeva, K. Yesmakhanova, G. Nugmanova, R. Myrzakulov. *International Journal of Geometric Methods in Modern Physics*, **13**, N1, 1550134 (2016).. [arXiv:1404.2270]
- [55] Myrzakul Akbota and Myrzakulov Ratbay. *Integrable Motion of Two Interacting Curves and Heisenberg Ferromagnetic Equations*, Abstracts of XVIII-th Intern. Conference "Geometry, Integrability and Quantization", June 3-8, 2016, Bulgaria.
- [56] Myrzakul Akbota and Myrzakulov Ratbay. *Integrable motion of two interacting curves, spin systems and the Manakov system*. *International Journal of Geometric Methods in Modern Physics*, **13**, N1, 1550134 (2016). [arXiv:1606.06598]
- [57] Myrzakul Akbota and Myrzakulov Ratbay. *Darboux transformations and exact soliton solutions of integrable coupled spin systems related with the Manakov system*, [arXiv:1607.08151]
- [58] Myrzakul Akbota and Myrzakulov Ratbay. *Integrable geometric flows of interacting curves/surfaces, multilayer spin systems and the vector nonlinear Schrodinger equation*. *International Journal of Geometric Methods in Modern Physics*, **13**, N1, 1550134 (2016). [arXiv:1608.08553]
- [59] Myrzakulova Zh., Myrzakul A., Nugmanova G., Myrzakulov R. *Notes on Integrable Motion of Two Interacting Curves and Two-layer Generalized Heisenberg Ferromagnet Equations*, [arXiv:1811:12216]
- [60] Hussien R.A., Mohamed S.G. *Generated Surfaces via Inextensible Flows of Curves in R^3* . *Journal of Applied Mathematics*, v.2016, Article ID 6178961 (2016).
- [61] Bekova G., Yesmakhanova K., Shaikhova G., Nugmanova G., Myrzakulov R. *Geometric formulation and soliton solutions of the Myrzakulov-LXXIII equation and the complex short pulse equation*,
- [62] C. Qu, J. Song and R. Yao. *Multi-Component Integrable Systems and Invariant Curve Flows in Certain Geometries*, *Symmetry, Integrability and Geometry: Methods and Applications SIGMA* 9 (2013), 001, 19 pages [arXiv:1301.0180]