

# A Stochastic production planning problem

Elena Cristina Canepa

University Politehnica of Bucharest, Bucharest, Romania.

E-mail address: cristinacanepa@yahoo.com

Dragos-Patru Covei

The Bucharest University of Economic Studies, Bucharest, Romania.

E-mail address: dragos.covei@csie.ase.ro

Traian A. Pirvu

McMaster University, Hamilton, Canada.

E-mail address: tpirvu@math.mcmaster.ca

December 2, 2019

## Abstract

Stochastic production planning problems were studied in several works; the model with one production good was discussed in [3]. The extension to several economic goods is not a trivial issue as one can see from the recent works [4], [5] and [6]. The following qualitative aspects of the problem are analyzed in [5]; the existence of a solution and its characterization through dynamic programming/HJB equation, as well as the verification (i.e., the solution of the HJB equation yields the optimal production of the goods). In this paper, we stylize the model of [4] and [5] in order to provide some quantitative answers to the problem. This is possible especially because we manage to solve the HJB equation in closed form. Among other results, we find that the optimal production rates are the same across all the goods and they also turn to be independent of some model parameters. Moreover we show that production rates are increasing in the aggregate number of goods produced, and they are also uniformly bounded. Numerical experiments show some patterns of the output.

## 1 Introduction

Production planning problems were studied for quite some time. [15] considered a stochastic production-inventory model to determine optimal production rates, i.e., the ones which minimize a discounted quadratic loss function. Their solution has three terms: the initial inventory, a steady state of the solution and a correction term which kicks in when time approaches maturity. This work was extended from a deterministic to a stochastic framework by [3] and [14] who added randomness to the

dynamics of the inventory process. The work of [7] looks at the infinite horizon stochastic production planning problem in which a continuous-time Markov chain models the demand.

The aforementioned papers consider in general the production planning problem with one economic good only. The extension to several economic goods makes the problem more mathematically involved as one can see from the recent works of [4] and [5]. Moreover, [5] characterized the solution through dynamic programming/HJB equation; using regularity and estimate results from the area of partial differential equations a classical solution of the HJB was established, and the verification result was proved. Since these works deal with the infinite horizon, a transversality condition was imposed on the value function, and it was shown that the value function verifies it. The paper [6] is within the paradigm of multiple goods' production. Because of the complexity of HJB equations, the goal is not to solve the HJB equations, but to offer an approximate solution.

In this paper we specialized the model of [4] and [5] to make it more tractable and to obtain quantitative results. Our main contribution is that we solved in closed form the HJB equation and the optimal production rate. The solution displays a mean field structure; the optimal production rate of some good is a function of the number of that specific produced good and an average of all the goods produced (this average is expressed by a norm of the vector of goods produced). By exploiting the structure of our closed form solution we can see that the optimal production rates are the same across all goods and they do not depend on some model parameters. Moreover, the optimal production rates are zeros when there are no goods produced, and they are of order  $O(\frac{1}{N})$  ( $N$  here stands for the number of goods). We show that production rates are increasing in the aggregate number of goods produced, and they are also uniformly bounded. Numerical experiments reveal that the production rate is a decreasing function of the number of goods' type  $N$  and, the variance of the number of goods produced.

Finally, the HJB equation characterizing the optimal production rates appears in other practical applications as we mention in the last section of the paper.

Now we are ready to present the organization of this paper. Section 2 describes the model. Section 3 provides the methodology. Section 4 presents other practical applications of the mathematics developed. The paper ends with an appendix containing a technical proof.

## 2 The model

Consider a factory producing  $N$  types of economic goods which stores them in an inventory designated place. Next, we describe the model mathematically. There exists a complete probability space  $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{0 \leq t < \infty}, P)$ , on which lives a  $N$ -dimensional Brownian motion denoted by  $w = (w_1, \dots, w_N)$ .

The filtration  $\{\mathcal{F}_t\}_{0 \leq t < \infty}$ , is the natural filtration of the Brownian motion. Let  $p(t) = (p_1(t), \dots, p_N(t))$ , represent the production rate at time  $t$  (control variable). Next, let us introduce the control variables. Let the

threshold  $p^0 = (p_1^0, \dots, p_N^0)$  be a vector standing for the factory optimal production level. This level can be optimal from a technological standpoint, but its implementation may not be optimal because of inventory costs.

Next, let  $l = (l_1, \dots, l_N)$  be the factory-optimal inventory level which can be attained but not maintained since there is noise in the system. In order to simplify the notations we assume that  $p^0 = l = (0, \dots, 0)$ . This simplification is obtained by considering deviations from the factory-optimal inventory level and the factory-optimal production level. The deviations may be negative.

Next, let us describe the inventories. There exists a constant demand rate for every economic good, demand rate represented by the vector  $\xi = (\xi_1, \dots, \xi_N)$ . Again, to simplify the notations we take  $\xi = (0, \dots, 0)$  meaning that we consider deviations from the constant demand rate.

Let  $y_i^0$  denote the initial inventory level of good  $i$ , and  $y_i(t)$  the inventory level of good  $i$ , at time  $t$ . These inventory levels are modelled by the following system of stochastic differential equations

$$dy_i(t) = (p_i - \xi_i)dt + \sigma dw_i, \quad y_i(0) = y_i^0, \quad i = 1, \dots, N, \quad (1)$$

where  $\sigma$  is a constant (non-zero) diffusion coefficient. Let us recall that the stochasticity here is due to inventory spoilages which are random in nature.

Let  $\tau$  be the stopping time representing the moment when the inventory level reaches some threshold  $R$ , i.e.,

$$\tau = \inf_{t > 0} \{|y(t)| \geq R\}.$$

Here,  $|\cdot|$  stands for the Euclidian norm, and this way of limiting the inventory level is imposed for tractability. The factory may consider stopping the production when the inventory level  $R$  is attained and/or exceeded.

## 2.1 The Objective

The performance over time of a production  $p(t) = (p_1(t), \dots, p_N(t))$  is measured by means of its cost. At this point we introduce the cost functional which yields the cost:

$$J(p_1, \dots, p_N) := E \int_0^\tau (|p(t)|^2 + |y(t)|^2) dt, \quad (2)$$

which measures the quadratic loss. Again let us recall that we measure deviations from an optimal state, whence the loss. At this point we are ready to frame our objective, which is to minimize the cost functional. i.e.,

$$\inf \{J(p_1, \dots, p_N) \mid p_i, \forall i = 1, 2, \dots, N\}, \quad (3)$$

subject to the Itô equation (1).

### 3 The Methodology

Having presented the problem we want to solve, now we provide our means to tackle it. Our approach is based on the value function and dynamic programming which leads to the HJB equation. Let  $z$  denote the value function, i.e.,

$$z(y_1^0, y_2^0, \dots, y_N^0) = \inf\{J(p_1, \dots, p_N) \mid p_i, \forall i = 1, 2, \dots, N\},$$

subject to the Itô equation (1). We apply probabilistic techniques to characterize the value function; that is we search for a function  $U(x)$  such that the stochastic process  $M^p(t)$  defined below

$$M^p(t) = U(y(t)) - \int_0^t [f_1(p(s)) + f_2(y(s))] ds,$$

is supermartingale for all  $p(t) = (p_1(t), \dots, p_N(t))$  and martingale for the optimal control  $p^*(t) = (p_1^*(t), \dots, p_N^*(t))$ . Once such a function is found it turns out that  $-U = z$ . We search for  $U$  a  $C^2[0, R]$  function and the supermartingale/martingale requirement yields by means of Ito's Lemma the Hamilton-Jacobi-Bellman (HJB) equation which characterizes the value function

$$-\frac{\sigma^2}{2}\Delta z - |x|^2 = \inf\{p\nabla z + |p|^2 \mid \forall i = 1, 2, \dots, N\}. \quad (4)$$

This HJB can be turned into a partial differential equation (PDE) since a simple calculation yields

$$\inf\{p\nabla z + |p|^2 \mid p_i \forall i = 1, 2, \dots, N\} = -\frac{1}{4}|\nabla z|^2. \quad (5)$$

Thus, the HJB equation becomes the PDE

$$-\frac{\sigma^2}{2}\Delta z - |x|^2 = -\frac{1}{4}|\nabla z|^2 \text{ for } x \in \mathbb{R}^N, |x| \leq R,$$

or, equivalently

$$2\sigma^2\Delta z + 4|x|^2 = |\nabla z|^2 \text{ for } x \in \mathbb{R}^N, |x| \leq R. \quad (6)$$

The change of variable  $z = -v$ , yields the PDE

$$\Delta v = \frac{4|x|^2 - |\nabla v|^2}{2\sigma^2} \text{ for } x \in \mathbb{R}^N, |x| \leq R. \quad (7)$$

The gradient term in the above PDE can be removed by the change of variable  $u(x) = e^{\frac{v(x)}{2\sigma^2}}$ , to get a simpler PDE

$$\begin{cases} \Delta u(x) = \frac{1}{\sigma^4}|x|^2 u(x) \text{ for } x \in \mathbb{R}^N, |x| \leq R, \\ u(x) > 0 \text{ for } x \in \mathbb{R}^N, |x| \leq R. \end{cases} \quad (8)$$

The value function will give us in turn the candidate optimal control. The first order optimality conditions on the lefthand side of (5) are sufficient for optimality since we deal with a quadratic (convex) function and they produce the candidate optimal control as follows:

$$p_i^* = \bar{p}_i(y_1(t), \dots, y_N(t)), \quad i = 1, \dots, N,$$

and

$$\bar{p}_i = \frac{1}{2} \frac{\partial v}{\partial x_i}(x_1, \dots, x_N), \text{ for } i = 1, \dots, n. \quad (9)$$

### 3.1 The Equation of Value Function

Let  $B_R(0)$  be the ball in  $\mathbb{R}^N$  centered at the origin and radius  $R > 0$ . The equation of the value function according to (8) is

$$\Delta u(x) = \frac{1}{\sigma^4} |x|^2 u(x) \text{ in } B_R(0). \quad (10)$$

The boundary condition is taken to be

$$u(0) = \alpha, \quad (11)$$

where  $\alpha$  is a positive constant. The following result concerns the equation of value function.

**Theorem 3.1.** *Given the positive constant  $\alpha$ , there exists a unique positive radially symmetric solution  $u_\alpha \in C^2[0, R]$ , to the problem (10) subject to the Dirichlet boundary condition (11). Moreover, the solution is convex and increasing, and the following holds true*

$$u'_\alpha(0) = 0, \quad (12)$$

$$u_\alpha(r) = \alpha \left( 1 + \sum_{j=1}^{\infty} \frac{1}{j!(N+2)(N+6)\dots(N+4j-2)} \left( \frac{r^2}{2\sigma^2} \right)^{2j} \right) \quad (13)$$

$$u'_\alpha(r) = \alpha \sum_{j=1}^{\infty} \frac{4jr}{2\sigma^2 j!(N+2)(N+6)\dots(N+4j-2)} \left( \frac{r^2}{2\sigma^2} \right)^{2j-1}, \quad (14)$$

for all  $r := |x| \in [0, R]$ . In addition,

$$u_\alpha(r) \leq \alpha e^{\frac{r^4}{4\sigma^4(N+2)}}, \quad r \in [0, R], \quad (15)$$

$$(u_\alpha)'(r) \leq \frac{\alpha r^3}{\sigma^4(N+2)} e^{\frac{r^4}{4\sigma^4(N+2)}}, \quad r \in [0, R], \quad (16)$$

hold.

**Proof. It is done in the appendix**

### 3.2 Verification

In this subsection we show that the control of (18) is indeed optimal. In a first step let us show that  $M^P(t)$

$$M^P(t) = U(y(t)) - \int_0^t (|p(s)|^2 + |y(s)|^2) ds,$$

is supermartingale for all

$$p(t) = (p_1(t), \dots, p_N(t))$$

and martingale for the optimal control

$$p^*(t) = (p_1^*(t), \dots, p_N^*(t)).$$

Indeed, Ito' Lemma yields for the optimal control candidate

$$dM^p(t) = \left( \frac{\sigma^2}{2} \Delta U(y(s)) - |y(s)|^2 + p(s) \nabla U(s) - |p(s)|^2 \right) ds + \sigma p(s) \nabla z(y(s)) dw(s).$$

Then, the claim yields in light of HJB equation (4).

In a second step let us establish the optimality of  $(p_1^*, \dots, p_N^*)$ . The martingale/supermartingale principle yields

$$EU(y^*(\tau^*)) - E \int_0^{\tau^*} (|p^*(u)|^2 + |y^*(u)|^2) du = U(y^*(0)) = U(y(0)),$$

and

$$EU(y(\tau)) - E \int_0^\tau (|p(u)|^2 + |y(u)|^2) du \leq U(y(0)).$$

Here, let us recall that  $\tau^* = \inf_{t>0} \{|y^*(t)| \geq R\}$  and  $\tau = \inf_{t>0} \{|y(t)| \geq R\}$ . Moreover,

$$EU(y^*(\tau^*)) = EU(y(\tau)) = 2\sigma^2 \ln u(R),$$

and this finishes the proof.

### 3.3 Optimal Control

Let us notice that equations (9) become

$$\bar{p}_i(y_1, \dots, y_N) = \sigma^2 \frac{u'_\alpha(r)}{ru_\alpha(r)} y_i, \quad r \neq 0, \quad i = 1, 2, \dots, N, \quad (17)$$

and  $r = |y|$ . The optimal control is given by

$$p_i^* = \bar{p}_i(y_1(t), \dots, y_N(t)), \quad i = 1, \dots, N,$$

and

$$dy_i(t) = p_i^* dt + \sigma dw_i, \quad y_i(0) = y_i^0, \quad i = 1, \dots, N. \quad (18)$$

This SDE system has a unique solution since the map  $y \rightarrow \bar{p}_i(y)$ ,  $i = 1, \dots, N$ , is Lipschitz on  $[0, R]$ . Let us notice that the production rate

$$\frac{\bar{p}_i}{y_i} = \sigma^2 \frac{u'_\alpha(r)}{ru_\alpha(r)}, \quad r \neq 0, \quad (19)$$

is the same across **all** goods. Let us notice the connection with **mean field models**, with the key quantity being  $r = |y|$ .

*Remark 3.2.* The choice of  $\alpha > 0$  is irrelevant because the value function equation admits the following symmetry; if  $u$  is the solution with  $\alpha = 1$ , then  $\alpha u$  is the solution for arbitrary  $\alpha > 0$ . However, both  $u$  and  $\alpha u$  yield the same optimal control (see (19)). Let us notice that if we impose the boundary condition  $\bar{u}(R) = \alpha > 0$  instead of (11) then we get a solution  $\bar{u}$  which is a scalar multiple of  $u$ , i.e.,  $\bar{u} = Ku$ , for some constant  $K > 0$ . Thus,  $\bar{u}$  yields the same optimal control (see (19)). Therefore, the optimal control does not depend on the choices of  $\alpha$  and  $R$ .

In light of this remark we set  $\alpha = 1$ , so that

$$u(r) := u_1(r) = 1 + \sum_{j=1}^{\infty} \frac{1}{j!(N+2)(N+6)\dots(N+4j-2)} \left(\frac{r^2}{2\sigma^2}\right)^{2j}, \quad (20)$$

for all  $r \geq 0$ , whence we can get the production rate  $\sigma^2 \frac{u'(r)}{ru(r)}$ ,  $r \neq 0$ , in closed form. Moreover, from (13) we get that  $\lim_{r \rightarrow 0} \frac{u'(r)}{ru(r)} = 0$ , thus the optimal production rates are zeros when there are no goods produced.

Using (20) and operations with power series (see [12] Chapter 1), we get the optimal production rate in closed form.

**Theorem 3.3.** *The optimal production rate is given by*

$$\frac{\bar{p}_i}{y_i} = \sigma^2 \frac{u'(r)}{ru(r)} = \frac{4\sigma^2}{r^2} \sum_{j=0}^{\infty} c_j \left[\frac{r^4}{4\sigma^4}\right]^j, \quad r \neq 0,$$

where

$$\begin{aligned} a_0 &= 1, \quad c_0 = 0, \quad c_j = \frac{1}{a_0} \left[ b_j - \sum_{i=1}^j c_{j-i} a_i \right], \quad j = 1, 2, 3, \dots \\ a_j &= \frac{1}{j!(N+2)(N+6)\dots(N+4j-2)}, \quad j = 1, 2, \dots \\ b_j &= \frac{j}{j!(N+2)(N+6)\dots(N+4j-2)}, \quad j = 1, 2, \dots \end{aligned}$$

The production rate is increasing and bounded. This fact will be made precise in the following Lemma.

**Lemma 3.4.** *The function*

$$r \rightarrow \frac{u'(r)}{ru(r)}$$

is increasing and

$$\frac{u'(r)}{ru(r)} \leq \frac{1}{\sigma^2}. \quad (21)$$

*Proof.* The first part of the claim yields if the derivative of this function is positive which boils down to

$$u''(r) \geq \frac{(u'(r))^2}{u(r)} + \frac{u'(r)}{r}.$$

Next we use the fact that  $u$  solves the following ODE

$$u''(r) + \frac{N-1}{r} u'(r) = \frac{1}{\sigma^4} r^2 u(r),$$

whence, the claim becomes

$$\frac{1}{\sigma^4} r^2 u(r) \geq N \frac{u'(r)}{r} + \frac{(u'(r))^2}{u(r)}.$$

This is equivalent to

$$\frac{u'(r)}{u(r)} \leq \frac{\sqrt{\frac{N^2}{r^2} + \frac{4r^2}{\sigma^4}} - \frac{N}{r}}{2},$$

or

$$\frac{u'(r)}{ru(r)} \leq \frac{\sqrt{\frac{N^2}{r^2} + \frac{4r^2}{\sigma^4}} - \frac{N}{r}}{2r}. \quad (22)$$

This argument shows that

$$r \rightarrow \frac{u'(r)}{ru(r)},$$

is increasing if and only if (22) holds true. However, the function

$$r \rightarrow \frac{\sqrt{\frac{N^2}{r^2} + \frac{4r^2}{\sigma^4}} - \frac{N}{r}}{2r},$$

is increasing, both functions are 0 when  $r = 0$  (since  $u'(0) = 0$ ) and  $r \rightarrow \frac{u'(r)}{ru(r)}$  is increasing on some small interval  $[0, \epsilon]$  in light of  $u$  being convex (for this see Theorem 3.1). This shows that  $r \rightarrow \frac{u'(r)}{ru(r)}$  is increasing and (22) holds true. Moreover, since

$$r \rightarrow \frac{\sqrt{\frac{N^2}{r^2} + \frac{4r^2}{\sigma^4}} - \frac{N}{r}}{2r}$$

is increasing and has as asymptote at infinity  $\frac{1}{\sigma^2}$  we also get the second part of the claim.  $\square$

### 3.4 Asymptotic Analysis

Let us recall the estimate for large  $N$  from [4]

$$\frac{u'(r)}{r} \leq \frac{K}{N-1}, \quad r \neq 0.$$

Thus, for big  $N$  an approximate solution is

$$\frac{u'(r)}{ru(r)} \leq \frac{K}{u_0(N-1)} \approx 0, \quad r \neq 0,$$

which says that the optimal control  $p^* \approx 0$ , since

$$\bar{p}_i = \sigma^2 \frac{u'(r)}{ru(r)} x_i, \quad r \neq 0, \quad i = 1, \dots, N.$$

This means that if the number of goods is big then  $p^* = 0$  is an approximate solution.

Next, we prove an asymptotical result.

**Lemma 3.5.** *The following result hold true*

$$\lim_{r \rightarrow \infty} \left[ \frac{u'(r)}{ru(r)} \right] = \frac{1}{\sigma^2}. \quad (23)$$



*Proof.* Because the function  $u$  is convex and increasing (for this see Theorem 3.1) it follows that

$$\lim_{r \rightarrow \infty} u(r) = \lim_{r \rightarrow \infty} u'(r) = \infty.$$

In light of Lemma 3.4 the limit exists and is finite. Let us denote it by  $l$ . L'Hospital rule yields

$$l = \lim_{r \rightarrow \infty} \left[ \frac{u'(r)}{ru(r)} \right] = \lim_{r \rightarrow \infty} \left[ \frac{u''(r)}{u(r) + ru'(r)} \right]. \quad (24)$$

Next we use the fact that  $u$  solves the following ODE

$$u''(r) + \frac{N-1}{r}u'(r) = \frac{1}{\sigma^4}r^2u(r),$$

whence

$$u''(r) = \frac{1}{\sigma^4}r^2u(r) - \frac{N-1}{r}u'(r).$$

Inserting this into (24) we get

$$l = \frac{1}{l\sigma^4}.$$

Therefore

$$l = \lim_{r \rightarrow \infty} \left[ \frac{u'(r)}{ru(r)} \right] = \frac{1}{\sigma^2}.$$

□

### 3.5 Simulation of the optimal inventory

Let us recall the SDE system

$$dy_i(t) = p_i^* dt + \sigma dw_i, \quad y_i(0) = y_i^0, \quad i = 1, \dots, N, \quad (25)$$

governing the optimal inventory.

This SDE system can be simulated numerically. It can be done using a Euler scheme as follows: start with  $y_i^0, i = 1, \dots, N$ , and

$$r = \sum_{i=1}^N [y_i^0]^2, \quad r \neq 0.$$

On  $[0, \Delta t]$  we approximate

$$y_i(\Delta t) \simeq \sigma^2 \frac{u'(r)}{ru(r)} y_i^0 + \sigma \sqrt{\Delta t} Z_i^0, \quad r \neq 0,$$

where  $Z_i^0$  is standard normal.

Next repeat this on  $[\Delta t, 2\Delta t]$  as follows:

$$r(\Delta t) = \sum_{i=1}^N [y_i^{\Delta t}]^2,$$

and

$$y_i(2\Delta t) \simeq \sigma^2 \frac{u'(r(\Delta t))}{r(\Delta t)u(r(\Delta t))} y_i^{\Delta t} + \sigma \sqrt{\Delta t} Z_i^1,$$

where  $Z_i^1$  is standard normal. The process is then repeated on  $[2\Delta t, 3\Delta t]$ , and so on. In the following we present two plots resulting from this simulation procedure. We considered  $N = 2$  (two economic goods) and  $\sigma = 2$  in the first plot  $\sigma = 5$  in the second plot.

### 3.6 Numerical Experiments

In the first set of experiments we set  $\sigma = 0.5$ , and vary  $N$  the number of goods' type.

We observe from these set of plots the following patterns:

- 1) the production rate is an increasing function of the total number of goods produced, fact explained by Lemma 3.4;
- 2) when the total number of goods produced exceed a certain threshold the production rate converges to 1, fact explained by Lemma 3.5;
- 3) the production rate is a decreasing function of the total number of goods produced.

In the next set of plots we set  $N = 100$ , and vary  $\sigma$ .

We observe from these set of plots the following patterns:

- 1) the production rate is an increasing function of the total number of goods produced, fact explained by Lemma 3.4;
- 2) when the total number of goods produced exceed a certain threshold the production rate converges, fact explained by Lemma 3.5;
- 3) the production rate is a decreasing function of  $\sigma$ .

## 4 Other Applications

The value function equation characterizing the optimal control, i.e., (10), appears naturally in other practical applications. There is by now a vast literature concerning on the existence of positive solutions and their behaviour for the partial differential equation

$$\Delta u(x) = f(x, u(x)) \text{ for } x \in \Omega, \quad (26)$$

where  $\Omega$  is a bounded or unbounded domain of  $\mathbb{R}^N$  ( $N \geq 1$ ) or the all space  $\mathbb{R}^N$  and  $f$  is a function suitable chosen.

The interest in studying the above equation comes, for instance, from various physical situations, such as quantum mechanics, quantum optics, nuclear physics and reaction-diffusion processes (cf. [1, 9, 10, 11]). For instance, a basic preoccupation for the study of problem (26) is the time-independent Schrödinger equation (single non-relativistic particle)

$$\Delta u = \frac{2m}{\hbar^2}(V(x) - E)u, \quad \hbar = h/2\pi, \quad (27)$$

where  $h$  is Planck's constant,  $\hbar$  is the reduced Planck constant (or the Dirac constant),  $E$  and  $V(x)$  are the total (non relativistic) and potential energies of a particle of mass  $m$ , respectively.

Besides the importance in applications, the equation (26) also raises many difficult mathematical problems that need to be solved. In general, the existence of the solutions and numerical approximation of the elliptic problem (26) is widely open. See the paper of Santos, Zhou and Santos [11], which includes a nice survey and recent progresses for Eq. (26).

Let us mention this result which is interesting in itself.

**Theorem 4.1.** (see [13, Theorem 2.1, p. 199]) *The problem (10) subject to the Dirichlet boundary condition*

$$u(x) \rightarrow \infty \text{ as } |x| \rightarrow R, \quad (28)$$

*has no positive solutions.*

Even if the next result has no importance in economic theories, it helps us to understand the beauty of this problem and to discover other questions that will need to be solved by the researchers.

**Theorem 4.2.** (see [11]) *The problem (10) with  $B_R(0)$  replaced with  $\mathbb{R}^N$ , admits a sequence of symmetric radial solutions  $u_k(|x|) \in C^2(\mathbb{R}^N)$  with*

$$u_k(0) = \infty \text{ as } k \rightarrow \infty.$$

*Besides this,  $u'_k \geq 0$  in  $[0, \infty)$ .*

In the next, we provide two exact solutions for the problem (10) with  $B_R(0)$  replaced with  $\mathbb{R}^4 \setminus \{0_{\mathbb{R}^4}\}$ . They are:

$$u_1(x) = \alpha e^{\frac{1}{2\sigma^2}|x|^2} |x|^{-2}, \quad \alpha \in \mathbb{R} \text{ and } x \in \mathbb{R}^4 \setminus \{0_{\mathbb{R}^4}\}, \quad (29)$$

$$u_2(x) = \alpha e^{-\frac{1}{2\sigma^2}|x|^2} |x|^{-2}, \quad \alpha \in \mathbb{R} \text{ and } x \in \mathbb{R}^4 \setminus \{0_{\mathbb{R}^4}\}. \quad (30)$$

The solutions (29) and (30) were determined by analyzing the series in (13) and can be used by physicists in the study of the time-independent Schrodinger equation (27). Moreover, reasoning in the same manner we think that similar solutions can be constructed for the total (non relativistic) and potential energies of a particle of mass  $m$  in (27).

Next, we posit the following open problems inspired by the two solutions and [11].

**Problem 4.1.** *Assume that  $g \in C^1([0, \infty), [0, \infty))$  is a non-decreasing function satisfying*

$$\int_{\gamma}^{\infty} \frac{1}{\sqrt{\int_0^t g(s) ds}} dt = \infty, \text{ for } t \geq \gamma > 0,$$

*and  $p$  is a non-negative continuous symmetric radially function such that*

$$\int_0^{\infty} t^{1-N} \int_0^t s^{N-1} p(s) ds dt = \infty.$$

*Then, there exists at least one positive radially symmetric solution  $u \in C^2(\mathbb{R}^N \setminus \{0_{\mathbb{R}^N}\})$  for the problem*

$$\Delta u(x) = p(r) g(u(x)) \text{ in } \mathbb{R}^N, \quad r = |x|, \quad (31)$$

*subject to the Dirichlet boundary condition*

$$u(x) \rightarrow \infty \text{ as } |x| \rightarrow \infty, \quad (32)$$

*such that*

$$u(x) \rightarrow \infty \text{ as } |x| \rightarrow 0.$$

*Moreover,  $\partial u / \partial r \geq 0$  on  $[t_0, \infty)$  and  $\partial u / \partial r < 0$  on  $[0, t_0)$ , for some  $t_0 \geq 0$ .*

**Problem 4.2.** Under the same assumptions on  $p$  and  $g$  as in Problem 4.1, there exists at least one positive radially symmetric solution  $u \in C^2(\mathbb{R}^N \setminus \{0_{\mathbb{R}^N}\})$  of (31) subject to the Dirichlet boundary condition

$$u(x) \rightarrow 0 \text{ as } |x| \rightarrow \infty, \quad (33)$$

such that

$$u(x) \rightarrow \infty \text{ as } |x| \rightarrow 0.$$

Moreover,  $\partial u / \partial r \leq 0$  on  $[0, \infty)$ .

Example of solutions for problems 4.1, and 4.2 are the ones given in (29), and (30). To the best of our knowledge the only result for the problems 4.1, and 4.2 is Theorem 4.2.

## 5 Appendix

### 5.1 Proof of Theorem 3.1

We consider the radial form of the problem (10) subject to the Dirichlet boundary condition (11), i.e.,

$$\begin{cases} u''_{\alpha}(r) + \frac{N-1}{r}u'_{\alpha}(r) = \frac{1}{\sigma^4}r^2u_{\alpha}(r) & \text{in } [0, R], \\ u_{\alpha}(0) = \alpha. \end{cases} \quad (34)$$

We show that the solution  $u_{\alpha}(r)$  of (34) can be obtained successively in the following way

$$\begin{cases} u_{\alpha}^0(r) = u_{\alpha}(0) = \alpha, \\ u_{\alpha}^k(r) = \alpha + \int_0^r t^{1-N} \int_0^t s^{N-1} \frac{1}{\sigma^4} s^2 u_{\alpha}^{k-1}(s) ds dt & \text{for } 0 < r \leq R \text{ and } k \in \mathbb{N}^*. \end{cases} \quad (35)$$

It is easy to see that  $\{u_{\alpha}^k(r)\}_{k \geq 0}$  is a nondecreasing sequence of functions satisfying

$$\begin{aligned} u_{\alpha}^{k+1}(r) - u_{\alpha}^k(r) &\leq \frac{\alpha}{(k+1)!} \left( \frac{r^4}{4\sigma^4(N+2)} \right)^{k+1} & (36) \\ &\leq \frac{\alpha}{(k+1)!} \left( \frac{R^4}{4\sigma^4(N+2)} \right)^{k+1} \xrightarrow{k \rightarrow \infty} 0, & (37) \end{aligned}$$

for all  $r \in [0, R]$ . Then  $\{u_{\alpha}^k(r)\}_{k \geq 0}$  is a Cauchy sequence of functions on  $[0, R]$ . It is a straightforward argument to prove that

$$u_{\alpha}^k \in C^2[0, R], \quad k \in \mathbb{N}.$$

Since a Cauchy sequence of functions is convergent, it has a limit function  $u_{\alpha}(r)$  and the convergence is uniform. Moreover, since a uniformly Cauchy sequence of continuous functions has a continuous limit, then  $u_{\alpha}(r)$  is a continuous function on  $[0, R]$ .

By passing to the limit in (35) we obtain that  $u_{\alpha}(r)$  verifies the integral form of the problem (10) subject to the Dirichlet boundary condition (11)

$$u_{\alpha}(r) = \alpha + \int_0^r t^{1-N} \int_0^t s^{N-1} \frac{1}{\sigma^4} s^2 u_{\alpha}(s) ds dt, \text{ in } [0, R]. \quad (38)$$

Hence, the limit function  $u_\alpha(r)$  is the solution of (10) subject to the Dirichlet boundary condition (11).

Next, we examine the sequence  $\{(u_\alpha^k(r))'\}_{k \geq 0}$ . We note first that

$$0 \leq (u_\alpha^k)'(r) = r^{1-N} \int_0^r t^{N-1} t^2 u_\alpha^{k-1}(t) dt. \quad (39)$$

Thus, the function  $r \rightarrow u_\alpha^k(r)$  is nondecreasing for all  $k \in \mathbb{N}$ . Using (36) and (37) we get

$$\begin{aligned} \left| (u_\alpha^{k+1})'(r) - (u_\alpha^k)'(r) \right| &\leq \frac{r^3 \alpha}{\sigma^4(N+2)k!} \left( \frac{r^4}{4\sigma^4(N+2)} \right)^k \\ &\leq \frac{R^3 \alpha}{\sigma^4(N+2)k!} \left( \frac{R^4}{4\sigma^4(N+2)} \right)^k \xrightarrow{k \rightarrow \infty} 0. \end{aligned} \quad (40)$$

Consequently,

$$(u_\alpha^k(r))' \xrightarrow{k \rightarrow \infty} (u_\alpha(r))' \text{ uniformly in } [0, R],$$

which implies that  $(u_\alpha(r))'$  is a continuous function on  $[0, R]$ . A direct computation shows that

$$u_\alpha \in C^2[0, R].$$

Next, let us prove (15). To do this we use (36) successively

$$\begin{aligned} u_\alpha^{k+1}(r) &\leq \frac{\alpha}{(k+1)!} \left( \frac{r^4}{4\sigma^4(N+2)} \right)^{k+1} + u_\alpha^k(r) \\ &\leq \frac{\alpha}{(k+1)!} \left( \frac{r^4}{4\sigma^4(N+2)} \right)^{k+1} + \frac{\alpha}{k!} \left( \frac{r^4}{4\sigma^4(N+2)} \right)^k + u_\alpha^{k-1}(r) \\ &\quad \dots \\ &\leq \sum_{j=0}^{k+1} \frac{\alpha}{j!} \left( \frac{r^4}{4\sigma^4(N+2)} \right)^j. \end{aligned} \quad (41)$$

On the other hand, we note that

$$u_\alpha(r) = \lim_{k \rightarrow \infty} u_\alpha^{k+1}(r) \leq \sum_{j=0}^{\infty} \frac{\alpha}{j!} \left( \frac{r^4}{4\sigma^4(N+2)} \right)^j = \alpha e^{\frac{r^4}{4\sigma^4(N+2)}}, \quad (42)$$

for all  $r \in [0, R]$ .

Next, let us prove (16). We observe that  $\{(u_\alpha^k)'(r)\}_{k \geq 0}$  is a nondecreasing sequence of continuous functions. Following the proof in (41), and using (40) successively it can be shown the inequality

$$\begin{aligned} (u_\alpha^{k+1})'(r) &\leq \frac{\alpha r^3}{\sigma^4(N+2)k!} \left( \frac{r^4}{4\sigma^4(N+2)} \right)^k + (u_\alpha^k)'(r) \\ &\quad \dots \\ &\leq \frac{\alpha r^3}{\sigma^4(N+2)} \sum_{j=0}^k \frac{1}{j!} \left( \frac{r^4}{4\sigma^4(N+2)} \right)^j. \end{aligned}$$

Repeating the arguments of (42) we notice that

$$\begin{aligned}
(u_\alpha)'(r) &= \lim_{k \rightarrow \infty} \left( u_\alpha^{k+1} \right)'(r) \\
&\leq \frac{\alpha r^3}{\sigma^4(N+2)} \sum_{j=0}^{\infty} \frac{1}{j!} \left( \frac{r^4}{4\sigma^4(N+2)} \right)^j \\
&= \frac{\alpha r^3}{\sigma^4(N+2)} e^{\frac{r^4}{4\sigma^4(N+2)}},
\end{aligned}$$

for all  $r \in [0, R]$ .

Next, let us prove (13). To do this, we observe that

$$\begin{aligned}
u_\alpha^1(r) &= \alpha + \int_0^r t^{1-N} \int_0^t s^{N+1} \frac{1}{\sigma^4} u_\alpha^0(r) ds dt \\
&= \alpha + \int_0^r t^{1-N} \int_0^t s^{N+1} \frac{1}{\sigma^4} \alpha ds dt \\
&= \alpha \left( 1 + \frac{1}{\sigma^4} \int_0^r \frac{t^3}{N+2} dt \right) \\
&= \alpha \left( 1 + \frac{1}{4\sigma^4} \frac{r^4}{N+2} \right).
\end{aligned}$$

Substituting  $u_\alpha^1(r)$  into

$$u_\alpha^2(r) = \alpha + \int_0^r t^{1-N} \int_0^t s^{N+1} \frac{1}{\sigma^4} u_\alpha^1(r) ds dt,$$

we obtain

$$u_\alpha^2(r) = \alpha \left( 1 + \frac{1}{4\sigma^4} \frac{r^4}{N+2} + \frac{r^8}{\sigma^8 \cdot 4 \cdot 8 \cdot (N+2)(N+6)} \right).$$

Continuing this process we get

$$\begin{aligned}
u_\alpha^k(r) &= \alpha + \frac{1}{\sigma^4} \int_0^r t^{1-N} \int_0^t s^{N+1} u_\alpha^{k-1}(s) ds dt \\
&= \alpha \left( 1 + \sum_{j=1}^k \frac{1}{j! (N+2)(N+6)\dots(N+4j-2)} \left( \frac{r^2}{2\sigma^2} \right)^{2j} \right).
\end{aligned}$$

Since the sequence of functions  $\{u_\alpha^k\}_{k \geq 0}$  is uniform convergent to the limit function  $u_\alpha(r)$  then (13) is proved.

The power series representation of function  $u_\alpha(r)$  can be differentiated to obtain a power series representation of its derivative  $u'_\alpha(r)$ . Thus, we obtain that  $u_\alpha(r)$  is differentiable on  $[0, R]$  and (14) holds true. In addition, the term-by-term derivative of a power series has the same interval of convergence as the original power series.

Next, (14) leads to  $u'_\alpha(0) = 0$ , whence (12) is proved. A direct computation shows that

$$u_\alpha \in C^2([0, R]).$$

The convexity of the solution is proved in [4] and the uniqueness of solution follows from Remark 3.2. The monotonicity of the solution is now obvious. This completes the proof.

**Acknowledgements.** This work was supported by a mobility grant of the Romanian Ministry of Research and Innovation, CNCS-UEFISCDI, project number PN-III-P1-1.1-MCD-2019-0151, within PNCDI III, and NSERC grant 5-36700.

## References

- [1] O. Alvarez, *A quasilinear elliptic equation in  $R^N$* , Proc. Roy. Soc. Edinburgh Sect. A, 126 (1996) 911-921.
- [2] L. Arnold, *Stochastic Differential Equations*, Wiley, New York, 1974.
- [3] A. Bensoussan, S.P. Sethi, R. Vickson and N. Derzko, *Stochastic production planning with production constraints*, SIAM J. Control Optim., 22 (1984) 920-935.
- [4] D.-P. Covei, *Symmetric solutions for an elliptic partial differential equation that arises in stochastic production planning with production constraints*, Appl. Math. Comput., 350 (2019) 190-197.
- [5] D.-P. Covei, and T. A. Pirvu, *An elliptic partial differential equation and its application*, Appl. Math. Lett., 101 (2020) 1-7.
- [6] A. Gharbi and J. P. Kenne, *Optimal production control problem in stochastic multiple-product multiple-machine manufacturing systems*, IIE Transactions, 35 (2003) 941-952.
- [7] W. H. Fleming, S. P. Sethi and H. M. Soner, *An Optimal Stochastic Production Planning Problem with Randomly Fluctuating Demand*, SIAM J. Control Optim., 25 (1987) 1494-1502.
- [8] J.M. Lasry, P. L. Lions, *Nonlinear Elliptic Equations with Singular Boundary conditions and Stochastic Control with State Constraints*, Math. Ann., 283 (1989) 583-630.
- [9] T. Leonori, *Large solutions for a class of nonlinear elliptic equations with gradient terms*, Adv. Nonlinear Stud., 7 (2007) 237-269.
- [10] A. Porretta, *Some uniqueness results for elliptic equations without condition at infinity*, Commun. Contemp. Math., 5 (2003) 705-717.
- [11] C.A. Santos, J. Zhou and J. A. Santos, *Necessary and sufficient conditions for existence of blow-up solutions for elliptic problems in Orlicz-Sobolev spaces*, Complex Var. Elliptic Equ., 62 (2017) 887-899.
- [12] D. G. Simpson, *Power series*, NASA Goddard Space Flight Center Greenbelt, Maryland 20771. <https://caps.gsfc.nasa.gov/simpson/pubs/series.pdf>
- [13] H. Yang, *On the existence and asymptotic behavior of large solutions for a semilinear elliptic problem in  $\mathbb{R}^N$* , Commun. Pure Appl. Anal., 4 (2005) 187-198.
- [14] S. P. Sethi and G. L. Thompson, *Applied Optimal Control: Applications to Management Science*, Nijhoff, Boston, 1981.
- [15] G. L. Thompson and S. P. Sethi, *Turnpike horizons for production planning*, Management Sci., 26 (1980) 229-241.