Wide Class of Logarithmic Potentials with Power-Tower Kink Tails

Avinash Khare

Physics Department, Savitribai Phule Pune University Pune 411007, India

Avadh Saxena

Theoretical Division and Center for Nonlinear Studies, Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA

Abstract:

We present a wide class of potentials which admit kinks and corresponding mirror kinks with either a power law or an exponential tail at the two extreme ends and a power-tower form of tails at the two neighbouring ends, i.e. of the forms *ette* or *pttp* where e, p and t denote exponential, power law and power-tower tail, respectively. We analyze kink stability equation in all these cases and show that there is no gap between the zero mode and the beginning of the continuum. Finally, we provide a recipe for obtaining logarithmic potentials with power-tower kink tails and estimate kink-kink interaction strength.

1 Introduction

A vast majority of kink solutions obtained during the last four decades for a variety of field theory potentials, e.g. sine-Gordon, double sine-Gordon, ϕ^4 , ϕ^6 , etc. harbor kinks with an exponential tail [1]. Recently we and others have presented a wide class of kink-bearing potentials for which one has a power law kink tail [2, 3, 4, 5, 6]. Very recently we have also presented a model with a super-exponential profile with one of the tails also being super-exponential [7, 8, 9, 10]. Thus by now we have models where one has a variety of kink tails such as of power law, exponential or super-exponential form. The obvious question is if there are models with still different types of kink tails. The purpose of this paper is to present an entirely different and novel class of potentials with *power-tower* kink tails, thus further expanding the type of kink asymptotes one could realize. One of the logarithmic potentials with super-exponential kink tails arises in the context of infinite order phase

transitions [7] and, therefore, conceivably the family of potentials considered here may have similar physical relevance.

The plan of the paper is as follows. In Section 2 we discuss a oneparameter family of potentials with kink tails of the form *ette*. We discuss the stability analysis of these kink solutions, which are expressed in terms of the exponential integral function Ei(x) [11, 12] and show that there is no gap between the zero mode and the beginning of the continuum. In Section 3 we consider a two-parameter family of potentials which lead to kink tails of the form *pttp*. We also discuss the stability analysis of these solutions and show that even in this case there is no gap between the zero mode and the beginning of the continuum. Our main conclusions are summarized in Section 4 where we also discuss how to obtain potentials with power-tower kink tails and estimate interaction between such kinks.

2 Models With Tails of the Form *ette*

In this section we consider a one parameter family of potentials of the form

$$V(\phi) = (1/2)\phi^{2m+2}[(1/2)\ln(\phi^2)]^2, \quad m \ge 1.$$
(1)

These potentials have degenerate minima at $\phi = 0, \pm 1$ with $V_{min} = 0$ while they have degenerate maxima at

$$\phi_{max} = \pm e^{-1/(m+1)}, \quad V_{max} = \frac{1}{2e^2(m+1)^2}.$$
 (2)

Thus notice that while $\phi_{max}(m=1) = \pm e^{-1/2}$, as *m* becomes larger, then ϕ_{max} moves towards ± 1 . On the other hand while, $V_{max}(m=1) = \frac{1}{8e^2}$, as *m* becomes larger, V_{max} decreases progressively towards zero. All these models for any integer *m* admit a kink from 0 to 1 and a mirror kink from -1 to 0 (and corresponding antikinks) with tails of the form *ette*. As an illustration we first discuss the case of m = 1, 2 and then generalize to arbitrary *m*. The potential given by Eq. (1) is shown in Fig. 1 for different values of *m*.

Case I: m = 1

Consider the potential

$$V(\phi) = (1/2)\phi^4 [(1/2)\ln(\phi^2)]^2.$$
(3)

Thus the self-dual equation we need to solve is

$$\frac{d\phi}{dx} = \pm \phi^2 [(1/2)\ln(\phi^2)].$$
(4)



Figure 1: Potential $V(\phi)$ for m = 1, m = 2 and m = 4 (see Eq. (1)).

For the kink between 0 and 1 we need to solve the self-dual Eq. (4) with negative sign. In fact this is true no matter what $m (\geq 1)$ is. This is easily integrated by making the substitution $t = (1/2) \ln(\phi^2)$ and we obtain the implicit kink solution

$$-x = \int \frac{e^{-t}}{t} dt = Ei(-t), \qquad (5)$$

where Ei(x) denotes the exponential integral function [11, 12]. Unfortunately, we do not know how to invert this function analytically [13] and obtain t and hence ϕ in terms of x. However, using the Taylor series expansion of Ei(x) as given in [11]

$$Ei(x) = \gamma + \ln|x| + x + \frac{x^2}{22!} + \dots,$$
(6)

as well as the asymptotic formula [11]

$$Ei(x) \equiv e^x \left[\frac{1}{x} + \frac{1}{x^2} + \frac{2!}{x^3} + \frac{3!}{x^4} + \dots \right],$$
(7)

we can estimate the tail behaviour around $\phi = 0$ as $x \to -\infty$ and around $\phi = 1$ as $x \to +\infty$. Here $\gamma = 0.577$ is the Euler's constant.



Figure 2: Kink solution $\phi(x)$ for m = 1, m = 2 and m = 4 (see Eq. (17)).

We find that

$$\lim_{x \to -\infty} \phi(x) \ln[\phi(x)] = \frac{1}{x}, \quad \lim_{x \to +\infty} \phi(x) = 1 - e^{-(x+\gamma)}.$$
 (8)

It is worth pointing out that the asymptotic behaviour around $\phi = 0$ (as $x \to -\infty$) can also be written as

$$\lim_{x \to -\infty} \phi(x)^{\phi(x)} = e^{1/x}, \qquad (9)$$

which is known in the literature as the power-tower function [14] of order two. It is also related to the iterated or repeated exponentiation, i.e. tetration [15]. We can invert Eq. (5) numerically and obtain the kink solution as given in Fig. 2. For $x \to \infty$ the kink tail approaches $\phi = 1$ asymptotically as an *exponential* tail whereas for $x \to -\infty$ it approaches $\phi = 0$ asymptotically as a *power-tower* tail.

Case II: m = 2

Consider the potential

$$V(\phi) = (1/2)\phi^6[(1/2)\ln(\phi^2)]^2.$$
(10)

Thus the self-dual equation we need to solve is

$$\frac{d\phi}{dx} = \pm \phi^3[(1/2)\ln(\phi^2)].$$
 (11)

This is easily integrated by making the substitution $t = (1/2) \ln(\phi^2)$ and we obtain the implicit kink solution

$$-x = \int \frac{e^{-2t}}{t} dt = Ei(-2t).$$
 (12)

From here using Eqs. (6) and (7) it is easily checked that

$$\lim_{x \to -\infty} \phi^2(x) \ln[\phi(x)] = \frac{1}{2x}, \quad \lim_{x \to +\infty} \phi(x) = 1 - \frac{1}{2} e^{-(x+\gamma)}.$$
(13)

It is worth pointing out that the asymptotic behaviour around $\phi = 0$ (as $x \to -\infty$) in Eq. (13) can also be written as

$$\lim_{x \to -\infty} \phi(x)^{\phi(x)^2} = e^{1/2x}, \qquad (14)$$

which is known in the literature as the power-tower function of order two [14] or tetration [15]. The kink solution is obtained by numerically inverting Eq. (12) as shown in Fig. 2.

Case III: Arbitrary m

The generalization to arbitrary m is now straightforward. Consider the potential

$$V(\phi) = (1/2)\phi^{2m+2}[(1/2)\ln(\phi^2)]^2.$$
(15)

Thus the self-dual equation we need to solve is

$$\frac{d\phi}{dx} = \pm \phi^{m+1}[(1/2)\ln(\phi^2)].$$
(16)

This is easily integrated by making the substitution $t = (1/2) \ln(\phi^2)$ and we obtain the implicit kink solution

$$-x = \int \frac{e^{-mt}}{t} dt = Ei(-mt).$$
(17)

From here using Eqs. (6) and (7) it is easy to see that

$$\lim_{x \to -\infty} \phi^m(x) \ln[\phi(x)] = \frac{1}{mx}, \quad \lim_{x \to +\infty} \phi(x) = 1 - \frac{1}{m} e^{-(x+\gamma)}.$$
 (18)

It is worth pointing out that the asymptotic behaviour around $\phi = 0$ (as $x \to -\infty$) in Eq. (18) can also be written as

$$\lim_{x \to -\infty} \phi(x)^{\phi(x)^m} = e^{1/mx} \,, \tag{19}$$

which is known in the literature as the power-tower function of order two [14] or tetration [15]. Kink profiles for three different values of m are depicted in Fig. 2. Note that with increasing m, the approach to $\phi = 0$ for large negative x becomes progressively slower in accordance with the power-tower function. In other words, for large m the kink profile tends to become symmetric.

Kink Mass:

One can easily calculate the kink mass for the entire family of potentials. In particular, for the kink potential as given by Eq. (15), the kink mass is given by

$$M_K = \int_0^1 \sqrt{2V(\phi)} \, d\phi = \int_0^1 d\phi \, \phi^{m+1} \ln(\phi) d\phi = \frac{1}{(m+2)^2} \,. \tag{20}$$

Note that the kink mass decreases as m increases.

2.1 Stability Analysis

We can perform the stability analysis of all the above kink solutions and show that, akin to the kinks with the power law tail [4], for all the above kink solutions, there is no gap between the zero mode and the beginning of the continuum.

As an illustration, we discuss the m = 1 case in detail, the generalization to the arbitrary m case is then straightforward. In this case the self-dual equation is as given by Eq. (4) with minus sign. Thus the kink zero mode is given by

$$\psi_0(x) = \frac{d\phi_k}{dx} \propto \left[(\phi_k(x))^2 \ln[\phi_k(x)] \right],\tag{21}$$

where ϕ_k is the kink solution. The above zero mode ψ_0 is clearly nodeless and vanishes as $x \to \pm \infty$ since as x goes from $-\infty$ to ∞ , ϕ varies from 0 to 1. We can also calculate the corresponding kink potential $V_K(x)$ which appears in the stability equation

$$-\frac{d^2\psi}{dx^2} + V_K(x)\psi = \omega^2\psi, \qquad (22)$$

where $V_K(x) = \frac{d^2 V(\phi)}{d\phi^2}$, evaluated at $\phi = \phi_k(x)$. On using the potential for m = 1 as given by Eq. (3) we find that

$$V_K(x) = \frac{d^2 V(\phi_k)}{d\phi^2} = 6(\phi_k)^2 [\ln(\phi_k)]^2 + 7(\phi_k)^2 \ln(\phi_k) + (\phi_k)^2, \qquad (23)$$

and hence it is clear that while $V_K(x = \infty) = 1$, $V_K(-\infty) = 0$. Thus the continuum begins at $\omega^2 = 0$, i.e. there is no gap between the zero mode and the beginning of the continuum.

The generalization to the arbitrary m case is now straightforward. In particular, we find that the kink zero mode is given by

$$\psi_0(x) = \frac{d\phi_k}{dx} = [\phi_k(x)]^{m+1} \ln[\phi_k(x)], \qquad (24)$$

which is clearly nodeless. The corresponding kink stability potential is given by

$$V_K(x) = \frac{d^2 V(\phi_k)}{d\phi^2} = (m+1)(2m+1)(\phi_k)^{2m}[\ln(\phi_k)]^2 + (4m+3)(\phi_k)^{2m}\ln(\phi_k) + (\phi_k)^{2m},$$
(25)

from where again it is clear that there is no gap between the zero mode and the beginning of the continuum.

2.2 Nature of Kink-Kink and Kink-Antikink Interactions

In this section we have obtained kink and mirror kink solutions (and corresponding antikinks) for a one-parameter family of potentials as given by Eq. (1). The kinks are from 0 to 1 (and mirror kinks are from -1 to 0) as x goes from $-\infty$ to ∞ . We have seen that while the kink tail around $\phi = 1$ or $\phi = -1$ is exponential, the kink tail around $\phi = 0$ has a power-tower form. Using this information, let us try to qualitatively understand the nature of kink-kink (KK) and kink-antikink (K-AK) interactions.

Let us first consider the kink-kink interaction between the (-1, 0) mirror kink and the (0, 1) kink. From Eq. (1) it is clear that around $\phi = 0$ the kink

potential is as given by Eq. (1). Notice that if there were no $\ln(\phi^2)$ term in Eq. (1) then using the recent approach of Manton for potentials with a power law tail [16, 17] one would have immediately predicted that the KK force would be

$$F_{KK} = \left[\frac{\Gamma[m/2(m+1)]\Gamma[1/2(m+1)]}{2m\sqrt{\pi}}\right]^{2(m+1)/m} \frac{1}{2R^{2(m+1)/m}}, \qquad (26)$$

where R is the distance between the two kinks. So the question is: what is the effect of the $[(1/2)\ln(\phi^2)]$ term multiplying the potential in Eq. (1)? In this connection we notice that in a recent publication by the present authors [7] we have shown that in the case of the potential $V(\phi) = (1/2)\phi^2[(1/2)\ln(\phi^2)]^2$, while the KK force would have been exponentially small if there were no $\ln(\phi^2)$ term present, because of the $\ln(\phi^2)$ term, the force actually gets even weaker and is in fact super-exponential. Taking this as a guide, we would therefore expect that the KK force in the case of potential (1) will still have a power law fall off but perhaps with a slower fall off and the strength of the force too would be different. Only either a new formalism or numerical estimation can decide the issue.

Similar conclusion is also true concerning the force between (1,0) AK and (0,1) K. On the other hand the force between (0,1) K and (1,0) AK will be exponentially small and using the original Manton formalism [18] is of the form e^{-R} .

3 Models With Tails of the Form *pttp*

In this section we present a two-parameter family of potentials of the form

$$V(\phi) = (1/2)\phi^{2m+2}[(1/2)\ln(\phi^2)]^{2n+2}, \quad m, n \ge 1.$$
(27)

These potentials have degenerate minima at $\phi = 0, \pm 1$ with $V_{min} = 0$ while they have degenerate maxima at

$$\phi_{max} = \pm e^{-(n+1)/(m+1)}, \quad V_{max} = \frac{1}{2e^{2(n+2)}} \left[\frac{(n+1)}{(m+1)}\right]^{2(n+1)}.$$
 (28)

Notice that both ϕ_{max} and V_{max} depend on two parameters m and n. Notice also that for a fixed m, as $n \to \infty$, $\phi_{max} \to 0$ and $V_{max} \to 0$. On the other hand, for a fixed n, as $m \to \infty$, $\phi_{max} \to 1$ and $V_{max} \to 0$. Finally, for m = n, $\phi_{max} = \pm e^{-1}$ and the corresponding $V_{max} = \frac{1}{2e^{2(n+1)}}$. It is interesting to note that for a given m, all the potentials as given by Eq. (27) with arbitrary integral n have the same value $V(\phi) = \frac{1}{2e^{2(m+1)}}$ in case $\phi = \pm 1/e$ or $V(\phi) = \frac{e^{2(m+1)}}{2}$ in case $\phi = \pm e$. All these models for any integers m and n admit a kink from 0 to 1 and

All these models for any integers m and n admit a kink from 0 to 1 and a mirror kink from -1 to 0 (and corresponding antikinks) with tails of the form *pttp*. As an illustration we first discuss the case of arbitrary m and n = 1, 2 and then generalize to arbitrary n.

Case I: n = 1, m Arbitrary

Consider the potential

$$V(\phi) = (1/2)\phi^{2m+2}[(1/2)\ln(\phi^2)]^4.$$
(29)

The potential in Eq. (29) is shown in Fig. 3 for different values of m. Thus the self-dual equation we need to solve is

$$\frac{d\phi}{dx} = \pm \phi^{m+1} [(1/2)\ln(\phi^2)]^2 \,. \tag{30}$$



Figure 3: Potential $V(\phi)$ for n = 1 and m = 1, m = 2 and m = 4 (see Eq. (29)).

For the kink between 0 and 1 we need to solve the self-dual Eq. (30) with positive sign. This is easily integrated by making the substitution $t = (1/2) \ln(\phi^2)$ and we obtain the implicit kink solution

$$x = \int \frac{e^{-mt}}{t^2} dt = -\frac{e^{-mt}}{t} - m Ei(-mt).$$
 (31)

Using Eqs. (6) and (7) we then find that

$$\lim_{x \to -\infty} \phi(x)^m (\ln[\phi(x)])^2 = -\frac{1}{mx}, \quad \lim_{x \to +\infty} \phi(x) = 1 - \frac{1}{(x + m\gamma)}.$$
 (32)

It is worth pointing out that the asymptotic behaviour around $\phi = 0$ (as $x \to -\infty$) can also be written as

$$\lim_{x \to -\infty} \phi(x)^{\phi(x)^{m/2}} = e^{1/\sqrt{-mx}}.$$
(33)

Equation (31) can be inverted numerically and the kink profile is depicted in Fig. 4. For $x \to \infty$ the kink tails approach $\phi = 1$ as a power-law whereas for $x \to -\infty$ the kink tails approach $\phi = 0$ as a power-tower function. With increasing *m* for large negative *x* the tails approach $\phi = 0$ progressively slowly. In other words, for large *m* the kink profile tends to become symmetric.

Case II: n = 2

Consider the potential

$$V(\phi) = (1/2)\phi^{2m+2}[(1/2)\ln(\phi^2)]^6.$$
(34)

Thus the self-dual equation we need to solve is

$$\frac{d\phi}{dx} = \pm \phi^{m+1} ([(1/2)\ln(\phi^2)])^3.$$
(35)

For the kink between 0 and 1, unlike the n = 1 case, we need to solve the self-dual Eq. (35) with negative sign. This is easily integrated by making the substitution $t = (1/2) \ln(\phi^2)$ and we obtain the implicit kink solution

$$-x = \int \frac{e^{-mt}}{t^3} dt = -\frac{e^{-mt}}{2t^2} + \frac{me^{-mt}}{2t} + \frac{m^2}{2}Ei(-mt).$$
(36)

Using Eqs. (6) and (7) we then find that

$$\lim_{x \to -\infty} \phi^m(x) (\ln[\phi(x)])^3 = \frac{1}{mx}, \quad \lim_{x \to +\infty} \phi(x) = 1 - \frac{1}{[2x + m^2 \gamma]^{1/2}}.$$
 (37)



Figure 4: Kink solution $\phi(x)$ for n = 1 and m = 1, m = 2 and m = 4 (see Eq. (31)). Inset shows in detail how the three profiles cross each other for x > 0.

It is worth pointing out that the asymptotic behaviour around $\phi = 0$ (as $x \to -\infty$) in Eq. (37) can also be written as

$$\lim_{x \to -\infty} \phi(x)^{\phi(x)^{m/3}} = e^{1/(-mx)^{1/3}}, \qquad (38)$$

which is known in the literature as the power-tower function of order two [14] or tetration [15].

Case III: Arbitrary n

The generalization to arbitrary n is now straightforward. Consider the potential

$$V(\phi) = (1/2)\phi^{2m+2}[(1/2)\ln(\phi^2)]^{2n+2}, \qquad (39)$$

which is shown in Fig. 5 for m = 1 and general n. Thus the self-dual equation we need to solve is

$$\frac{d\phi}{dx} = \pm \phi^{m+1} ([(1/2)\ln(\phi^2)])^{n+1}.$$
(40)



Figure 5: Potential $V(\phi)$ for m = 1, and n = 1, n = 2, n = 3 and n = 4 (see Eq. (39)).

This is easily done by making the substitution $t = (1/2) \ln(\phi^2)$ and we obtain

$$\pm x = \int \frac{e^{-mt}}{t^{n+1}} \, dt \,. \tag{41}$$

This is easily integrated using [12]

$$\int \frac{e^{ax}}{x^n} dx = -e^{ax} \sum_{k=1}^{k=n-1} \frac{a^{k-1}}{(n-1)(n-2)\dots(n+1-k)x^{n-k}} + \frac{a^{n-1}}{(n-1)!} Ei(ax) \,.$$
(42)

We obtain

$$\pm x = -e^{-mt} \sum_{k=1}^{n} \frac{(-m)^{k-1}}{n(n-1)\dots(n+1-k)t^{n+1-k}} + \frac{(-m)^n}{n!} Ei(-mt), \quad (43)$$

which can be numerically inverted to obtain the kink solutions, as depicted in Fig. 6 for m = 1 and different values of n.

Using Eqs. (6) and (7) we can now find the kink tail around both $\phi = 0$ and $\phi = 1$. We note that in order to find the self-dual kink solution between



Figure 6: Kink solution $\phi(x)$ for m = 1 and n = 1, n = 2, n = 3 and n = 4 (see Eq. (43)).

0 and 1, we need to take +x (-x) in Eq. (43) depending on whether n is an odd (or even) integer. Note that this is consistent with what we have used for n = 1, 2. Using Eq. (7) we then find that for any integer n

$$\lim_{x \to -\infty} \phi^m(x) (\ln[\phi(x)])^{n+1} = (-1)^n \frac{1}{mx} \,. \tag{44}$$

On the other hand, for both odd and even integer n using Eq. (6) we find that

$$\lim_{x \to +\infty} \phi(x) = 1 - \frac{1}{\left[nx + \frac{m^n \gamma}{(n-1)!}\right]^{1/n}}.$$
(45)

It is worth pointing out that the asymptotic behaviour around $\phi = 0$ (as $x \to -\infty$) in Eq. (44) (i.e. for even integer n) can also be written as

$$\lim_{x \to -\infty} \phi(x)^{\phi(x)^{m/(n+1)}} = e^{1/(mx)^{1/(n+1)}}.$$
(46)

On the other hand the asymptotic behaviour around $\phi = 0$ (as $x \to -\infty$) in Eq. (45) (i.e. for odd integer n) can also be written as

$$\lim_{x \to -\infty} \phi(x)^{\phi(x)^{m/(n+1)}} = e^{1/(-mx)^{1/(n+1)}}, \qquad (47)$$

which is known in the literature as the power-tower function of order two [14] or tetration [15].

Kink Mass

One can easily calculate the kink mass for the entire family of potentials. In particular, for the kink potential given by Eq. (39), the kink mass is given by

$$M_K = \int_0^1 \sqrt{2V(\phi)} \, d\phi = \int_0^1 d\phi \, \phi^{m+1} [\ln(\phi)]^{n+1} d\phi = \frac{(n+1)!}{(m+2)^2} \,. \tag{48}$$

Note that the kink mass decreases as m increases keeping n fixed. On the other hand the kink mass increases as n increases keeping m fixed.

3.1 Stability Analysis

We can perform the stability analysis of the kink solutions discussed in this section and show that like the previous section (as well as the kinks with the power law tail), for all the kink solutions of this section, there is no gap between the zero mode and the onset of the continuum.

As an illustration, we discuss the m = n = 1 case in detail, the generalization to the arbitrary m, n case is then straightforward. In the case of m = n = 1, the self-dual equation is as given by (30). Thus the kink zero mode is given by

$$\psi_0(x) = \frac{d\phi_k}{dx} \propto [\phi_k(x)]^2 (\ln[\phi_k(x)])^2, \qquad (49)$$

(where ϕ_k is the kink solution) which clearly is nodeless and vanishes as $x \to \pm \infty$ since as x goes from $-\infty$ to ∞ , ϕ varies from 0 to 1.

We can also calculate the corresponding potential $V_K(x)$ which occurs in the stability equation (25). On using the potential for m = n = 1 as given by Eq. (29) we find that

$$V_K(x) = \frac{d^2 V(\phi_k)}{d\phi^2} = 6(\phi_k)^2 [\ln(\phi_k)]^4 + 14(\phi_k)^2 [\ln(\phi_k)]^3 + 6(\phi_k)^2 [\ln(\phi_k)]^2,$$
(50)

and hence it is clear that $V_K(x = \infty) = V_K(-\infty) = 0$. Thus the continuum begins at $\omega^2 = 0$, i.e. there is no gap between the zero mode and the onset of the continuum.

The generalization to the arbitrary m, n case is now straightforward. In particular, we find that the kink zero mode in that case is given by

$$\psi_0(x) = \frac{d\phi_k}{dx} = [\phi_k(x)]^{m+1}(x)(\ln[\phi_k(x)])^{n+1}, \qquad (51)$$

which is clearly nodeless. The corresponding kink stability potential is given by

$$V_K(x) = \frac{d^2 V(\phi_k)}{d\phi^2} = (m+1)(2m+1)[\phi_k(x)]^{2m}[\ln(\phi_k)]^{2n+2}$$

$$+ (4m+3)(n+1)[\phi_k(x)]^{2m}[\ln(\phi_k)]^{2n+1} + (n+1)(2n+1)(\phi_k)^{2m}[\ln(\phi_k)]^{2n},$$
(52)

from where again it is clear that there is no gap between the zero mode and the onset of the continuum.

3.2 Nature of Kink-Kink and Kink-Antikink Interactions

In this section we have obtained the kink and the mirror kink solutions (and the corresponding antikinks) for the two-parameter family of potentials as given by Eq. (27). The kinks are from 0 to 1 (and the mirror kinks are from -1 to 0) as x goes from $-\infty$ to ∞ . We have seen that while around $\phi = 1$ or $\phi = -1$ one has a power law tail, the kink tail around $\phi = 0$ has a power-tower form. Using this information we attempt to qualitatively understand the nature of the KK and the K-AK interactions.

Let us first consider the kink-kink interaction between the (-1, 0) mirror kink and the (0, 1) kink. From Eq. (1) it is clear that around $\phi = 0$ the kink potential is as given by Eq. (27). Notice that if there were no $[\ln(\phi^2)]^{2n+2}$ term in Eq. (27) then using the recent approach of Manton for potentials with a power law tail [16, 17] one would have immediately predicted that the KK force would be as given by Eq. (26). So the question is: what is the effect of the $[(1/2)\ln(\phi^2)]^{2(n+1)}$ term multiplying the potential in Eq. (27)? In this connection as noted earlier, in a recent publication [7] we have shown that in the case of the potential $V(\phi) = (1/2)\phi^2[(1/2)\ln(\phi^2)]^2$, while the kink-kink force would have been exponentially small if there were no $\ln(\phi^2)$ term present, because of the $\ln(\phi^2)$ term, the force actually gets even weaker and is in fact super-exponential. Taking this as a guide, we would therefore expect that the KK force in the case of the potential (27) will still have a power law fall off but perhaps with a slower fall off and the strength of the force too would be different. Only either a new formalism or numerical estimation can decide the issue.

Similar conclusion is also true concerning the force between (1,0) AK and (0,1) K. On the other hand the force between (0,1) K and (1,0) AK can be immediately estimated using the recent Manton formalism [16, 17] and is given by

$$F_{KK} = \left[-\frac{\sqrt{\pi}\Gamma[n/2(n+1)]}{\Gamma[-1/(2(n+1))]} \right]^{2(n+1)/n} \frac{1}{2R^{2(n+1)/n}}$$
(53)

where R denotes the distance between a kink and an antikink.

4 Summary

In this paper we have considered a continuous one-parameter family of potentials as given by Eq. (1), all of which have kink tails of the form *ette* where e and t correspond to exponential and power-tower type of tail, respectively. Similarly, in Sec. III we have constructed a two-parameter family of potentials given by Eq. (27), all of which admit kink tails of the form *pttp* where p corresponds to power law type of tail. For all these cases we have calculated the corresponding kink masses. Further, we have shown that the kink stability equation in all these cases is such that there is no gap between the zero mode and the beginning of the continuum in the Schrödinger-like equation. Finally, we have also qualitatively discussed the nature of the kink-kink and the kink-antikink interactions in all these cases.

By now we have a large number of kink bearing models which admit kinks with a variety of tails such as exponential [1], power-law [2, 3, 5, 6], super-exponential [7, 8, 9, 10] and power-tower. It is then natural to enquire if there is a recipe for constructing models which admit such a diverse variety of kink tails. In this context we might add that the recipe for constructing kink solutions with an exponential or a power law tail is well known [4]. For completeness we mention it first and then give the recipe for constructing the kink solutions with either super-exponential or power-tower type of tail.

Since a kink has finite energy it implies that the solution must approach one of the minima (vacua), say ϕ_0 , of the theory as $x \to \pm \infty$. If the lowest non-vanishing derivative of the potential at the minimum has order m, then by Taylor series expansion of the potential at the minimum and writing the field close to it as $\phi = \phi_0 + \eta$, one finds that the self-dual first order equation in η implies that (assuming that the potential vanishes at the minimum)

$$\frac{d\eta}{dx} \propto \eta^{m/2} \,. \tag{54}$$

Thus if m = 2 then $\eta \propto e^{-\alpha x}$ (i.e. exponential tail) while if m > 2 then $\eta \propto 1/x^{2/(m-2)}$ (i.e. power law tail).

In our recent paper about the super-exponential tail [7], we have shown that if instead

$$\frac{d\eta}{dx} \propto \eta^2 \ln(\eta^2) \,, \tag{55}$$

then $\eta \propto e^{-e^{-\alpha x}}$, so that there is a super-exponential tail.

On the other hand using the results of this paper it is clear that if

$$\frac{d\eta}{dx} \propto \eta^{m+1} [\ln(\eta^2)]^{n+1}, \quad m, n \ge 1,$$
 (56)

then η is a solution of the equation

$$\eta^m [\ln \eta]^{n+1} = (-1)^n \frac{1}{mx} \,, \tag{57}$$

which leads to power-tower kink tails.

Before ending this discussion, it is worth pointing out some of the open problems in the context of power-tower type kink tails.

- In this paper we have constructed one- and two-parameter family of potentials which lead to kink tails of the form *ette* and *pttp*, respectively. The obvious question is, can one similarly construct at least a oneparameter family of potentials which gives tails of the form *teet*, *tppt*, *tttt* as well as the mixed tails of the form *ettp*, *eppt* and *peet*? Finally, can one construct models with an admixture of super-exponential tails and exponential and/or power law and/or power-tower type of tails?
- 2. In this paper we have not been able to explicitly calculate the force between the (-1,0) kink and the (0,1) kink since the two ends facing each other have power-tower type of tails and in this case it is not straightforward to invert and obtain the behaviour of the tail as a function of x when $x \to -\infty$. As an illustration, consider the asymptotic behaviour around $\phi = 0$ in case $x \to -\infty$ as given by Eq. (8), that is

$$\lim_{x \to -\infty} \phi(x) \ln(\phi(x)) = \frac{1}{x}.$$
(58)



Figure 7: $\phi(x)$ vs x obtained by inverting Eq. (58) and compared with $\phi = 1/x$ and $\phi = 1/x^{1.25}$.

If $\ln(\phi(x))$ were not there then we know that for large negative x, $\phi(x) \propto -1/x$. In this connection we notice that in a recent publication [7] we have shown that in the case of the potential $V(\phi) = (1/2)\phi^2[(1/2)\ln(\phi^2)]^2$, while the kink tail around $\phi = 0$ would have been an exponential tail in case there were no $\ln(\phi^2)$ term present, because of the $\ln(\phi^2)$ term, the kink tail actually gets even weaker and is in fact super-exponential. Taking this as a guide, we speculate that corresponding to the power-tower form as given by Eq. (58), the behaviour of $\phi(x)$ for large negative x should be of the form

$$\lim_{x \to -\infty} \phi(x) = \frac{1}{(-x)^{1+\epsilon_{1,0}}}, \quad \epsilon_{1,0} > 0.$$
(59)

Here by $\epsilon_{1,0}$ one means $\epsilon_{m=1,n=0}$ corresponding to ϕ^m and $[\ln(\phi)]^{n+1}$ in Eq. (58) with m = 1, n = 0. We have inverted Eq. (58) numerically and from Fig. 7 we see that it can be fitted in the form of Eq. (59) with $\epsilon_{1,0}$ approximately equal to 0.25. This would imply that the potential around $\phi = 0$ is of the form ϕ^{2k} with k = 9/5. We might add here that



Figure 8: $\phi(x)$ vs 4x obtained by inverting Eq. (60) for the case m = 4.

the new Manton formalism [16, 17], even though developed for integral k is also valid for any real number k. Using this information, one can estimate the force between the (-1,0) K and the (0,1) K using the new Manton formalism and show that the KK force would vary like $R^{-9/2}$, where R is the distance between the two kinks.

In the same way, one can numerically invert for any m the equation around $\phi = 0$ for large negative x as given by Eq. (18), i.e.

$$\lim_{x \to -\infty} \phi(x)^m \ln(\phi(x)) = \frac{1}{mx}, \qquad (60)$$

and try to numerically estimate the corresponding exponent. As an illustration, in Fig. 8 we have inverted Eq. (60) for the case of m = 4.

3. Generalization of the above discussion in the case of the power-tower Eq. (44) with arbitrary m and n is now straightforward. In particular, if the power-tower equation is given by

$$\lim_{x \to -\infty} \phi^m(x) [\ln(\phi(x))]^{n+1} = (-1)^n \frac{1}{mx}, \qquad (61)$$

then we speculate that the behaviour of $\phi(x)$ for large negative x should be of the form

$$\lim_{x \to -\infty} \phi(x) = \frac{1}{(-mx)^{1/m + \epsilon_{m,n}}}, \quad \epsilon_{m,n} > 0.$$
 (62)

As an illustration, in Fig. 9 we have numerically inverted Eq. (61) in case m = 1, n = 1 and we find that $\epsilon_{1,1}$ is approximately equal to 0.5 which is larger than $\epsilon_{1,0}$, which is approximately 0.25.

In case the exponent is 0.5, this would imply that the potential around $\phi = 0$ is of the form ϕ^{2k} with k = 5/3. Using this information, one can estimate the force between (-1,0) K and (0,1) K using the new Manton formalism [16, 17] and show that the KK force would go like R^{-5} , where R is the distance between the two kinks.



Figure 9: $\phi(x)$ vs x obtained by inverting Eq. (61) for the case m = 1 and n = 2, and compared with $\phi = 1/x$, $\phi = 1/x^{1.2}$ and $\phi = 1/x^{1.5}$.

4. Looking at the two examples of m = 1, n = 0 and m = 1, n = 1 it is immediately clear that $\epsilon_{1,0} < \epsilon_{1,1}$. We speculate that in general



Figure 10: $\phi(x)$ vs x obtained by inverting Eq. (61) with m = 1 and n = 9.

for arbitrary m, n we will have the inequality $\epsilon_{m,n_1} < \epsilon_{m,n_2}$ in case $n_1 < n_2$.

5. In the same way, one can invert for any n and m the equation around $\phi = 0$ for large negative x as given by Eq. (61). As an illustration, in Fig. 10 we have numerically inverted Eq. (61) in case m = 1, n = 9. It is clear from the figure that the exponent $\epsilon_{1,9}$ is significantly bigger than 0.5. It thus appears that as n becomes progressively larger, effectively the kink tail around $\phi = 0$ for large negative x will approach an exponential tail.

We first elaborate our argument in the case of m = 1 and arbitrary n. Generalization to arbitrary m and n is then straightforward. For m = 1 and arbitrary n case, the corresponding exponent is $\epsilon_{1,n}$. This would imply that the potential around $\phi = 0$ is of the form ϕ^{2k} with

$$k = \frac{2 + \epsilon_{1,n}}{1 + \epsilon_{1,n}}.\tag{63}$$

Now we have seen from the examples of m = 1, n = 0; m = 1, n = 1and m = 1, n = 10 that as n increases $\epsilon_{1,n}$ becomes progressively larger. In other words, for very large n we expect that $\epsilon_{1,n} \gg 2$ and hence for very large n, k as defined by Eq. (63) tends to 1 which corresponds to an exponential tail. However, we would like to emphasize that no matter how large n is, so long as it is finite, k is strictly greater than one such that for all finite n, the kink tail has power law fall off thereby justifying the name power-tower. Therefore, kinks with power-tower type of tails provide a bridge between kinks with power law type of tails and kinks with exponential tails.

Generalization to arbitrary m is now straightforward. In this case the corresponding exponent is $\epsilon_{m,n}$. This would imply that the potential around $\phi = 0$ is of the form ϕ^{2k} with

$$k = \frac{m+1+m\epsilon_{m,n}}{1+m\epsilon_{m,n}}.$$
(64)

We surmise that for very large n, no matter what m is, $m\epsilon_{m,n} \gg m+1$, so that even in this case k would tend to one which corresponds to an exponential tail, although for any large but finite m, n, it will strictly be greater than one.

Also as shown in Sec. III, for arbitrary n and m, the kink stability equation is such that there is no gap between the zero mode and the beginning of the continuum, which is the hallmark of kink solutions with a power law tail.

It is worth pointing out that since for arbitrary m and n, the potential around $\phi = 0$ is of the form ϕ^{2k} with k as given by Eq. (64) hence using the new Manton formalism [16, 17] one can show that in that case the KK force would go like R^{-d} , where $d = 2[1 + \epsilon_{m,n} + 1/m]$.

6. In this paper by numerically inverting power-tower equations in a few cases we have shown that the exponent $\epsilon_{1,n}$ satisfies the inequality $\epsilon_{1,0} < \epsilon_{1,1}$. From here it is natural to speculate that even for arbitrary m, the exponents will satisfy the inequality $\epsilon_{m,n_1} < \epsilon_{m,n_2}$ in case $n_1 < n_2$. It would be desirable if one can prove this rigorously.

One of the obvious open problems is about the exponent for fixed n but varying m. In particular, what can one say about $\epsilon_{m_1,n} - \epsilon_{m_2,n}$ in case $m_1 < m_2$? Similarly proceeding further, can one say anything about $\epsilon_{m_1,n_2} - \epsilon_{m_2,n_1}$ where $n_1 < n_2$ and $m_1 < m_2$?

We hope to address some of these issues in the near future.

Acknowledgement: We acknowledge useful discussions with Ayhan Duzgun. A.K. is grateful to Indian National Science Academy (INSA) for the award of INSA Scientist position at Savitribai Phule Pune University. This work was supported in part by the U.S. Department of Energy.

References

- [1] Rajaraman R (1987) Solitons and Instantons 15 North Holland
- [2] Khare A, Christov I C, and Saxena A 2014 Phys. Rev. E 90 023208
- [3] Saxena A, Christov I C, Khare A, "Higher-order field theories: φ⁶, φ⁸ and beyond", in: A dynamical perspective on the φ⁴ model: Past, present and future, eds. P. G. Kevrekidis and J. Cuevas-Maraver, Chap. 11 (Springer, 2018) [arXiv:1806.06693].
- [4] Khare A and Saxena A 2019 J. Phys. A 52 365401
- [5] Gomes A R, Menezes R and Oliveira J C R E 2012 Phys. Rev. D 86 025008
- [6] Bazeia D, Menezes R and Moreira D C 2018 J. Phys. Commun 2 055019
- [7] Kumar P, Khare A and Saxena A arXiv:1908.04978
- [8] Bazeia D, Marques M A and Menezes R 2015 Phys. Rev. D 92 084058
- [9] Marques M A 2019 *EPL* **125** 11001
- [10] Bazeia D and Marques M A 2019 EPL **127** 21001
- [11] Harris F E 1957 Mathematical Tables and Other Aids to Computation 11 (no. 57), pp. 9-16, AMS, January
- [12] Gradshteyn I S and Ryzhik I M 1979 Table of Integrals, Series and Products (Academic Press).
- [13] Pecina P 1986 Bull. Astron. Inst. Czechosl. 37 8
- [14] Wolfram MathWorld: http://mathworld.wolfram.com/PowerTower.html
- [15] https://en.wikipedia.org/wiki/Tetration

- [16] Manton N S 2019 J. Phys. A 52 065401
- [17] Christov I C. Decker R J, Demirkaya A, Gani V A, Kevrekidis P G, Khare A and Saxena A 2019 Phys. Rev. Lett. 122 171601
- [18] Manton N S 1979 Nuc. Phys. B 150 397