

On the degree growth of iterated birational maps

Claude M. Viallet*

Sorbonne Université, Centre National de la Recherche Scientifique

October 1, 2019

Abstract

We construct a family of birational maps acting on two dimensional projective varieties, for which the growth of the degrees of the iterates is cubic. It is known that this growth can be bounded, linear, quadratic or exponential for such maps acting on two dimensional compact Kähler varieties. The example we construct goes beyond this limitation, thanks to the presence of a singularity on the variety where the maps act. We provide all details of the calculations.

1 Introduction

A simple characterisation of the complexity of birational self-maps of projective spaces is given by the nature of the growth of the sequence of degrees $\{d_n\}$ of their iterates: generic maps have exponential growth, and some exceptional maps have polynomial growth [1–3], [4–6], the latter behaviour being linked with the notion of discrete integrability [7].

A straightforward test of integrability is then the vanishing of the algebraic entropy [8]

$$\epsilon = \lim_{n \rightarrow \infty} \frac{1}{n} \log(d_n).$$

This global index of complexity (and equivalently its exponential called the dynamical degree) is always defined, and is conjectured to be the logarithm of an algebraic integer for any birational map [8] (see also [9]). It is clearly invariant by birational changes of coordinates. Moreover, if the degree growth happens to be polynomial (vanishing entropy), i.e. $d_n \simeq \alpha n^\nu$ when $n \rightarrow \infty$, then the power ν is itself invariant by birational changes of coordinates.

The singularity structure is known to play an essential role in the behaviour of the iterates. Singularities appear in two instances:

- birational self-maps of projective spaces of degree higher than 2 have a non empty singular locus. A large number of results, most notably the Lie algebraic characterisation of discrete Painlevé equations [10] inspired by [11], have been obtained from the description of this singular locus and the construction by blow ups of rational varieties where the maps are diffeomorphisms [12].

*claude.viallet@upmc.fr

– one may also consider the action of maps on algebraic subvarieties of projective spaces, and these varieties may or may not be smooth.

Both these types of singularities (of maps and of varieties) are present in our analysis.

We construct a family of birational maps acting on two dimensional projective varieties, having vanishing entropy, for which the growth of the degrees of the iterates is *cubic*. This goes beyond the result of [3], which would prevent anything between quadratic and exponential growth. The existence of a singular point on the varieties where the maps act is playing an important role in the result. Such a behaviour has been observed previously [13–15]. Our point here is to provide a complete proof that the growth is cubic, using a method which can be applied in all dimensions and for non-vanishing entropy.

Our construction proceeds as follows: we start from a pencil of elliptic curves in \mathbb{CP}_2 . All curves of this pencil are invariant by a birational map φ , which has a specific form, related to a second order recurrence relation. Such maps and pencils are typical ingredients of the so called QRT maps [12, 16, 17]. We then construct a third order recurrence through the inflation process described in [14]. This new recurrence defines birational maps in \mathbb{CP}_3 , which automatically have a rational invariant. The action on the two dimensional invariant projective varieties obtained in this way is the one we analyse, proving that the growth of the degree of the iterates is cubic.

2 A pencil of elliptic curves

Consider the recurrence relation

$$x_{n+1} + x_{n-1} = \frac{a + b x_n + d x_n^2}{c + d x_n}. \quad (1)$$

where a, b, c, d are independent parameters. It defines a birational map on the projectivised space of initial conditions \mathbb{CP}_2 :

$$\varphi : [\xi, \zeta, \tau] \mapsto [-a\tau^2 - (b\xi + c\zeta)\tau - d\xi(\xi + \zeta), \xi(c\tau + d\xi), \tau(c\tau + d\xi)] \quad (2)$$

written in homogeneous coordinates $[\xi, \zeta, \tau]$ of \mathbb{CP}_2 . The map φ has a rational invariant Π

$$\tau^3 \Pi = a(\xi + \zeta)\tau^2 + b\tau\xi\zeta + c(\xi^2 + \zeta^2)\tau + d\xi\zeta(\xi + \zeta) \quad (3)$$

defining a pencil of invariant elliptic curves in \mathbb{CP}_2 .

The map φ is integrable, but we will not go into this aspect. We will concentrate on the nature of the growth of the degrees of the iterates of the maps we consider. For (2), this sequence is

$$1, 2, 4, 8, 13, 20, 28, 38, \dots \quad (4)$$

which has quadratic growth. The proof results from the construction of an elliptic surface R obtained by a finite number of blow ups of \mathbb{CP}_2 , and the properties of the linear map induced by φ on the Picard group of R [12, 18, 19].

3 Inflation

We apply to (1) the inflation process described in [14] with the simple transformation

$$x_n = y_n + y_{n-1} \quad (5)$$

We get a recurrence relation of order three on y_n , and the corresponding map in \mathbb{CP}_3 is $\tilde{\varphi} : [x, y, z, t] \mapsto [x', y', z', t']$ with:

$$\begin{cases} x' = -a t^2 - b(x+y)t - c(x+y+z)t - d(x+y)(2x+2y+z), \\ y' = x(ct+d(x+y)), z' = y(ct+d(x+y)), t' = t(ct+d(x+y)). \end{cases} \quad (6)$$

which automatically has an invariant, constructed from the one of φ by setting $\xi = x + y$, $\zeta = y + z$, and $\tau = t$.

In other words the new map $\tilde{\varphi}$ leaves the pencil of two dimensional surfaces $\tilde{\Pi}(k) = 0$ invariant

$$\begin{aligned} \tilde{\Pi}(k) = & k_0(a t^2(2y+x+z) + b t(z+y)(x+y) + c t((x+y)^2 + (y+z)^2) \\ & + d(z+y)(x+y)(2y+x+z)) + k_1 t^3. \end{aligned} \quad (7)$$

with $k = (k_0, k_1)$

Lemma 1 The varieties $Q(k)$ of equation $\tilde{\Pi}(k) = 0$ are ruled surfaces. They are cones with one singular point, the tip $\Sigma = [1, -1, 1, 0]$.

Proof: Choosing a point $M \in \mathbb{CP}_3$ (different from Σ) determines a value $k(M)$ such that $M \in Q(k(M))$. It is immediate to see that the line $\Lambda(M) = \tau\Sigma + (1 - \tau)M$ is entirely contained in $Q(k(M))$. It is also straightforward to check that the surfaces $Q(k) = 0$ have Σ as a unique singular point.

Lemma 2 The map $\tilde{\varphi}$ sends the line $\Lambda(M)$ onto the line $\Lambda(\tilde{\varphi}(M))$

Proof: Straightforward.

Lemma 1 and 2 give the geometrical explanation of the fact that, loosely speaking, Σ is a fixed singular point of $\tilde{\varphi}$. We will give a precise meaning to this statement in section 6.

Remark The singular locus of $\tilde{\varphi}$ is the line Δ , which is intersection of the two hyperplanes $\{x + y = 0\}$ and $\{t = 0\}$. Notice that $\Delta \subset Q(k)$ for all k .

4 The degree sequence, a fast analysis

The simplest way to get the exact value of the degree of the iterates of $\tilde{\varphi}$ is to compute explicitly a sufficient number of terms of the sequence of degrees, and try to fit the generating function of their sequence with a rational function. We get

$$\{d_n\} = 1, 2, 4, 8, 14, 23, 35, 51, 71, 96, 126, 162, 204, \dots \quad (8)$$

which is fitted by the rational generating function

$$g(s) = \sum_{k=0}^{\infty} d_k s^k = \frac{1 - s + 2s^3 - s^4}{(1 + s)(1 - s)^4} \quad (9)$$

showing vanishing entropy [8] with *cubic* growth, since all poles of $g(s)$ are on the unit circle and the order of the pole at $s = 1$ is 4.

It is straightforward to conjecture that the sequence of degrees is given by

$$d_n = 17/16 + 5/12 n + 3/8 n^2 + 1/12 n^3 - 1/16 (-1)^n \quad (10)$$

We will now prove this result.

5 Preliminaries

5.1 The \mathbf{K} varieties

The rational map $\tilde{\varphi}$ has a rational inverse $\tilde{\psi}$, which can be written $\tilde{\psi} = \lambda \cdot \tilde{\varphi} \cdot \lambda$ with the linear involution $\lambda : [x, y, z, t] \mapsto [z, y, x, t]$. The composition of $\tilde{\varphi} \cdot \tilde{\psi}$ is the identity, but when written with homogeneous coordinates, it appears as a multiplication of all the coordinates by a common factor:

$$\tilde{\psi} \cdot \tilde{\varphi} = \mathbf{K}_{\tilde{\varphi}} id, \quad \tilde{\varphi} \cdot \tilde{\psi} = \mathbf{K}_{\tilde{\psi}} id. \quad (11)$$

The factors $\mathbf{K}_{\tilde{\varphi}}$ and $\mathbf{K}_{\tilde{\psi}}$ define varieties of codimension 1 in \mathbb{CP}_3 :

$$\mathbf{K}_{\tilde{\varphi}} = (ct + d(x + y))^3 = B_1^3, \quad \mathbf{K}_{\tilde{\psi}} = (ct + d(z + y))^3 = C_1^3 \quad (12)$$

We know that the map $\tilde{\varphi}$ blows the hyperplane $\{B_1 = 0\}$ (resp. $\tilde{\psi}$, $\{C_1 = 0\}$) down to something of smaller dimension and the fate of the images under further iterates of $\tilde{\varphi}$ (resp. $\tilde{\psi}$) eventually shapes the sequence of degrees [8].

The factors $\mathbf{K}_{\tilde{\varphi}}$ also appear in the calculations of the iterates (see section 5.2).

5.2 Two ways of calculating iterates

We will denote by $\tilde{\varphi}^k$ the *regularised* k -th power of $\tilde{\varphi}$. By this we mean that when computing the k th power with its homogeneous polynomial form, we remove all common factors.

We will make use of the seemingly trivial relation $\tilde{\varphi}^{k+1} = \tilde{\varphi} \cdot \tilde{\varphi}^k = \tilde{\varphi}^k \cdot \tilde{\varphi}$ to calculate iterates in two different ways:

1- Direct image (i.e. $\tilde{\varphi}^{k+1} = \tilde{\varphi} \cdot \tilde{\varphi}^k$)

- a) Apply the polynomial formulae giving the image by $\tilde{\varphi}$. At each step this will provide $N + 1$ polynomials in the homogeneous coordinates of the initial point.
- b) Remove the common factors.
- c) Proceed.

2 - Pull-back (i.e. $\tilde{\varphi}^{k+1} = \tilde{\varphi}^k \cdot \tilde{\varphi}$)

- a) Get the coordinates of $\tilde{\varphi}^{k+1}$ as the pull-backs by $\tilde{\varphi}$ of the coordinates of $\tilde{\varphi}^k$.
- b) Remove the common factors.
- c) Proceed.

Steps a) of the two methods give the same point in projective space, but they possibly produce different determinations, as the common factors differ in general. Step b) is crucial. Once the common factors are removed of course the determinations coincide.

The interest of the pull-back method is the following:

Suppose S is an indecomposable variety of codimension 1 of equation $E_S = 0$. The pullback by $\tilde{\varphi}$ of the equation of S gives the equation $E_{S'}$ of the image S' of S by $\tilde{\psi}$.

Claim: This pull-back may contain additional factors, which are part of the total transform, but not part of the proper transform, *and the only possible factors are powers of B_1 .*

$$\tilde{\varphi}^*(E_S) = B_1^{\nu_S} E_{S'} \quad (13)$$

with ν_S some integer depending on S .

6 The fate of the K varieties under the action of $\tilde{\varphi}$

The map $\tilde{\varphi}$ sends the whole hyperplane $\{B_1 = 0\}$ to the point $S_1 = [1, 0, 0, 0]$. Further images are evaluated by first a calculation of the image of a generic point and then the specialisation of the coordinates to the ones of the running point $[cx, cy, cz, -d(x+y)]$ of B_1 . We get successively $S_2 = [2, -1, 0, 0]$, $S_3 = [2, -2, 1, 0]$, $S_4 = \Sigma = [1, -1, 1, 0]$, and then $S_5 = \Sigma$ again. The points S_3 and Σ are singular points of $\tilde{\varphi}$. As mentioned above, Σ is a “singular fixed point” of $\tilde{\varphi}$. By this we mean the following:

Theorem 1 Σ is a singular point of $\tilde{\varphi}$ and $\tilde{\varphi}^{k+4}(B_1) = \Sigma$ for $k \geq 0$.

Proof of Theorem 1: The proof uses blow-ups intended to desingularise the map as in [20]. Since the hyperplane $\{B_1 = 0\}$ is blown down to the point S_1 by $\tilde{\varphi}$, the first step is to blow S_1 up and see what is the (new) image of B_1 . To be complete, the coordinatisation of the blow up should be done in more than one coordinate patch but the main point is to ensure that the induced map in the blown-up space does not send the two dimensional hyperplane $\{B_1 = 0\}$ into a smaller dimensional space.

In the transformed space obtained by blowing up S_1 , the image of $\{B_1 = 0\}$ is a curve. Therefore we have to go on and blow-up again (along that curve). A possible choice of coordinates for the explicit realisation of the blow ups is given in the Appendix. The result of the calculation is that we need three successive blow ups $\beta_{i,j}, j = 1 \dots 3$ at each point S_i . We denote by $\delta_{i,j}$ the inverse of $\beta_{i,j}$ and by $F_{i,j}$ the images of $\{B_1 = 0\}$ in the j -th blow up at S_i , with evident notations (for example $F_{4,2}$ is the image of $\{B_1 = 0\}$ by $\tilde{\varphi}^4$ in $\beta_{4,2} \cdot \beta_{4,1}(\mathbb{CP}_3)$).

We find that the $F_{i,1}$ and $F_{i,2}$ are curves, while the $F_{i,3}$ are two dimensional.

To obtain the action of $\tilde{\varphi}^{k+4}$ on $\{B_1 = 0\}$ for $k \geq 0$, we consider $\tilde{\varphi} \cdot \delta_{4,1} \cdot \delta_{4,2} \cdot \delta_{4,3}(F_{4,3})$. It is just $S_4 = \Sigma$, which proves the theorem.

Remark The plane $\{w_{4,1} = 0\}$ is sent to Σ by $\tilde{\varphi} \cdot \delta_{4,1}$. The map on this plane induced by $\tilde{\varphi}$ is isomorphic to the one we started with (Eq. 2). One may indeed go further and blow up $S_5 = S_4$ with $\beta_{5,1} = \beta_{4,1}$. The image of $\{B_1 = 0\}$ by $\tilde{\varphi}^5$ is a curve $F_{5,1}$ (see Appendix) *This curve is different from $F_{4,1}$, of which it is the image by $\tilde{\varphi}$.* As a consequence, if we wanted to desingularise the iterates of $\tilde{\varphi}$ by blow ups, one would need an infinite number of them, along the various curves $F_{n,1}$. Note that $F_{5,2}$ is a curve, and $F_{5,3}$ is two dimensional, as expected.

Corollary 1.1 *The action of the inverse $\tilde{\psi}^k$ on the hyperplane C_1 being obtained via the linear symmetry λ , Σ is also a “singular fixed point” of $\tilde{\psi}$ after the 4th iterate.*

7 Stabilisation of the iterates

Start from a generic point $p_0 = [x, y, z, t]$ and examine the structure of the iterates of the map $\tilde{\varphi}$, calculated explicitly, keeping track of the product structure of the components, along the lines of [21], see also [22]:

$$\begin{aligned} p_1 &= [A_1, x B_1, y B_1, t B_1] \\ p_2 &= [A_2, A_1 B_2, x B_1 B_2, t B_1 B_2] \\ p_3 &= [A_3, A_2 B_3, A_1 B_2 B_3, t B_1 B_2 B_3] \\ p_4 &= [A_4, A_3 B_4, A_2 B_3 B_4, t B_1 B_2 B_3 B_4] \\ p_5 &= [A_5, A_4 B_5, A_3 B_4 B_5, t B_1 B_2 B_3 B_4 B_5] \end{aligned}$$

Theorem 2 *The general form of the iterates p_k for $k \geq 3$ is*

$$p_k = [A_k, A_{k-1} B_k, A_{k-2} B_{k-1} B_k, t B_1 B_2 \dots B_k], \quad (14)$$

or, defining $\Gamma_k = t B_1 B_2 \dots B_k$, $p_k = [A_k, A_{k-1} B_k, A_{k-2} B_{k-1} B_k, \Gamma_k]$.

We will prove Theorem 2 and give the recurrence relation between the successive A_k and B_k using both the direct way and the pull back way to express p_{k+1} in terms of p_k . Denoting $[P_k^i]$, $i = 1..4$ the coordinates of p_k , we have

$$[P_{k+1}^i] \simeq \tilde{\varphi}([P_k^i]) \quad (\text{direct image}) \quad \text{and} \quad [P_{k+1}^i] \simeq [\tilde{\varphi}^*(P_k^i)] \quad (\text{pull-back}), \quad (15)$$

where the equivalence \simeq means equality up to common factors.

Proof of Theorem 2:

Defining the sequences $\{\alpha\}$ and $\{\beta\}$ by $\tilde{\varphi}^*(A_k) = B_1^{\alpha(k)} A_{k+1}$ and $\tilde{\varphi}^*(B_k) = B_1^{\beta(k)} B_{k+1}$, we may evaluate $\tilde{\varphi}^*(p_k)$ (up to common factors):

$$[\mathbf{B}_1^{\alpha(k)} A_{k+1}, \mathbf{B}_1^{\alpha(k-1)+\beta(k)} A_k B_{k+1}, \mathbf{B}_1^{\alpha(k-2)+\beta(k-1)+\beta(k)} A_{k-1} B_k B_{k+1}, \mathbf{B}_1^{(\sum_{j=1}^k \beta(j))} \Gamma_{k+1}] \quad (16)$$

We know from theorem 1 that no factor B_1 is left on the three first components for $k \geq 3$. This gives:

$$\alpha(k) = \alpha(k-1) + \beta(k) \quad (17)$$

which in turn proves the stability of the form of p_k since

$$p_{k+1} = [A_{k+1}, A_k B_{k+1}, A_{k-1} B_k B_{k+1}, \Gamma_{k+1}]. \quad (18)$$

8 Back to the sequence of degrees

Theorem 3 *The sequence of degrees $\{d_n\}$ is the one given in section 4, i.e.*

$$d_n = 17/16 + 5/12 n + 3/8 n^2 + 1/12 n^3 - 1/16 (-1)^n \quad (19)$$

Proof of Theorem 3: We may apply directly $\tilde{\varphi}$ to p_k , equate it (up to factors) to p_{k+1} , and get the new recurrence relations determining A_{k+1} and B_{k+1} in terms of the previous A 's and B 's.

$$\begin{cases} d D_k + c \Gamma_k - B_1 B_2 \dots B_{k-3} B_{k-2}^2 \mathbf{B}_{k+1} = 0 \\ \mathbf{A}_{k+1} \Gamma_{k-2} B_{k-2} + a \Gamma_k^2 + B_k D_{k-1} (d D_k + c \Gamma_k) + D_k \left(b \Gamma_k + d (D_k + A_k) \right) = 0 \end{cases} \quad (20)$$

with $D_k = A_k + A_{k-1} B_k$.

These relations give us the additional information on the degrees d_A and d_B of the A 's and the B 's:

$$d_B(n+1) - d_B(n) - d_B(n-1) + d_B(n-2) - 1 = 0, \quad d_B(1) = 1, \quad d_B(2) = 2, \quad d_B(3) = 4,$$

giving

$$\begin{aligned} d_B(n) &= 1/8 + n/2 + 1/4 n^2 - 1/8 (-1)^n, \\ d_A(n) &= 1 + \sum_{k=1}^n d_B(k), \\ d_A(n) &= 17/16 + 5/12 n + 3/8 n^2 + 1/12 n^3 - 1/16 (-1)^n. \end{aligned}$$

Όπερ ἔδει δεῖξαι.

9 Conclusion and perspectives

- The heuristic method used in section 4 proves to be extremely efficient: one gets the results in a fraction of a second with a small laptop. The reason is the rational nature of the generating function of the sequence of degrees (8), a property which is not exceptional. Although it is not true for all birational maps (see [23]), it is verified for a large number of them. This moreover supports our conjecture that the entropy is the logarithm of an algebraic integer, *keeping in mind that birationality is essential*, as is clear from sections 5 and 7. We will address this question in a forthcoming publication.
- An important result of [3] is that the sequence of degrees of the iterates of birational maps on compact Kähler varieties of dimension 2 have bounded, linear, quadratic or exponential growth. What we have produced here is an example of birational maps acting on 2-dimensional algebraic hypersurfaces in $\mathbb{C}P_3$ and having cubic growth. The surfaces where the maps act have one singular point, and that point is a sink for the maps, allowing this different behaviour. One could also view this as an example of birational maps acting on a Kähler variety of dimension 3 with cubic growth.
- The fact that the growth is cubic implies in particular that the maps we constructed *cannot be obtained by a reduction of the known integrable quad equations [24] since they all have at most quadratic growth [25]*.
- Another interesting feature of the maps (6) is obtained by a simple graphical analysis of the orbits, showing the existence of an additional invariant, *which cannot be*

algebraic [26,27]. It is worth noticing that the existence of a “fixed singular point” as shown in section 6 can be observed for the linearisable two dimensional example given in [28]. This similarity is not accidental. The integrability aspects will be treated elsewhere.

- The model can be made non-autonomous by setting $a = \alpha + \beta n + \gamma(-1)^n$. The sequence of degrees of the iterates *remains unchanged*, meaning that this non-autonomous generalisation is integrable as well. The proof goes exactly along the same line as above. The fact that the sequence of degree is unchanged between the autonomous case and the non-autonomous case comes from the fact that the singularity structures are equivalent.

Acknowledgements

Parts of these results have been presented at the SIDE 13 meeting in Fukuoka (Japan) November 2018 (<http://www.side-conferences.net/>). The author would like to thank the Sydney Mathematical Research Institute (University of Sydney), for hospitality and support during the final elaboration of this work, and N. Joshi for a number of challenging discussions and fruitful suggestions during the stay at the SMRI.

Appendix: Local coordinates for the blow-ups

We first have to deal with blow-ups at the points $S_i, i = 1..4$. Denoting $[x, y, z, t]$ the original homogeneous components in \mathbb{CP}_3 , we see that they all have a non vanishing first coordinate (x), and we will use local (affine) coordinates obtained by normalising this first coordinate to 1. There is a great arbitrariness in the choice of coordinates used to write the blow ups as explicit birational transformations. A possible choice for the first blow-up is

$$\beta_{1,1} : [x, y, z, t] \mapsto [u_{1,1}, v_{1,1}, w_{1,1}] = [z/y, t/y, t/x] \quad (21)$$

with $[u_{1,1}, v_{1,1}, w_{1,1}]$ as affine (local) coordinates in the blown-up space. The image of B_1 is the curve $F_{1,1} = \{d(u_{1,1} + 1) + cv_{1,1} = 0, w_{1,1} = 0\}$, and we need to perform a second blow-up along this curve. This is done by

$$\beta_{1,2} : [u_{1,1}, v_{1,1}, w_{1,1}] \mapsto [u_{1,2}, v_{1,2}, w_{1,2}] = [u_{1,1}, v_{1,1}, \frac{w_{1,1}}{d u_{1,1} + c v_{1,1} + d}] \quad (22)$$

The image of B_1 is still a curve $F_{1,2} = \{d(u_{1,2} + 1) + cv_{1,2} = 0, (ad - bc + c^2)v_{1,2}w_{1,2} + d = 0\}$, and we need one more blow up to have B_1 sent to a two dimensional image by $\tilde{\varphi}$:

$$\beta_{1,3} : [u_{1,2}, v_{1,1}, w_{1,2}] \mapsto [u_{1,3}, v_{1,2}, \frac{(ad - bc + c^2)v_{1,2}w_{1,2} + d}{d u_{1,2} + c v_{1,2} + d}] \quad (23)$$

After this third blow up, B_1 is sent to a two dimensional image $F_{1,3}$ by $\tilde{\varphi}$.

$$F_{1,3} = \{d(u_{1,3} + 1) + cv_{1,3} = 0\} \quad (24)$$

We may then proceed and blow up the point S_2 . We again have to do three successive blow ups

$$\begin{cases} \beta_{2,1} : [x, y, z, t] \mapsto [u_{2,1}, v_{2,1}, w_{2,1}] = [2z/(x+2y), 2t/(x+2y), t/x] \\ \beta_{2,2} : [u_{2,1}, v_{2,1}, w_{2,1}] \mapsto [u_{2,2}, v_{2,2}, w_{2,2}] = [u_{2,1}, w_{2,1}, \frac{d(u_{2,1}+2)+(b-2c)v_{2,1}}{w_{2,1}}] \\ \beta_{2,3} : [u_{2,2}, v_{2,2}, w_{2,2}] \mapsto [u_{2,3}, v_{2,3}, w_{2,3}] = [u_{2,2}, v_{2,2}, v_{2,2}/w_{2,2}] \end{cases} \quad (25)$$

Next are the blow ups over S_3 and S_4

$$\begin{cases} \beta_{3,1} : [x, y, z, t] \mapsto [u_{3,1}, v_{3,1}, w_{3,1}] = [(2z - x)/2(x + y), t/(x + y), t/x] \\ \beta_{3,2} : [u_{3,1}, v_{3,1}, w_{3,1}] \mapsto [u_{3,2}, v_{3,2}, w_{3,2}] = [u_{3,1}, v_{3,1}, w_{3,1}/(v_{3,1} + d/c)] \\ \beta_{3,3} : [u_{3,2}, v_{3,2}, w_{3,2}] \mapsto [u_{3,3}, v_{3,3}, w_{3,3}] = [u_{3,2}, v_{3,2}, \frac{cv_{3,2}+d}{2(ad-bc+c^2)w_{3,2}+c^2}] \end{cases} \quad (26)$$

$$\begin{cases} \beta_{4,1} : [x, y, z, t] \mapsto [u_{4,1}, v_{4,1}, w_{4,1}] = [(z - x)/(x + y), t/(x + y), t/x] \\ \beta_{4,2} : [u_{4,1}, v_{4,1}, w_{4,1}] \mapsto [u_{4,2}, v_{4,2}, w_{4,2}] = [u_{4,1}, v_{4,1}, \frac{w_{4,1}}{d(u_{4,1}+1)+cv_{4,1}}] \\ \beta_{4,3} : [u_{4,2}, v_{4,2}, w_{4,2}] \mapsto [u_{4,3}, v_{4,3}, w_{4,3}] = [u_{4,2}, v_{4,2}, \frac{d(u_{4,2}+1)+cv_{4,2}}{2(ad-bc+c^2)v_{4,2}w_{4,2}+d}] \end{cases} \quad (27)$$

$$\begin{cases} F_{4,1} = \{du_{4,1} + cv_{4,1} + d = 0, \quad w_{4,1} = 0\}. \\ F_{4,2} = \{du_{4,2} + cv_{4,2} + d = 0, \quad v_{4,2}w_{4,2} + 2d/(ad - bc + c^2) = 0\}. \\ F_{4,3} = \{du_{4,3} + cv_{4,3} + d = 0\}. \end{cases} \quad (28)$$

and

$$F_{5,1} = \{u_{5,1}^2 d^2 + d(b - c)u_{5,1}v_{5,1} + 3d^2u_{5,1} + (ad - c^2)v_{5,1}^2 + bdv_{5,1} + 2d^2, \quad w_{5,1} = 0\}, \quad (29)$$

recalling that $[u_{4+k,1}, v_{4+k,1}, w_{4+k,1}] = [u_{4,1}, v_{4,1}, w_{4,1}]$, but $[u_{4+k,j}, v_{4+k,j}, w_{4+k,j}] \neq [u_{4,j}, v_{4,j}, w_{4,j}]$ for $j = 2, 3, k \geq 0$. The curve $F_{5,1}$ is rational. It is just the transform of $F_{4,1}$ by the map induced by $\tilde{\varphi}$.

References

- [1] A.P. Veselov, *Growth and Integrability in the Dynamics of Mappings*. Comm. Math. Phys. **145** (1992), pp. 181–193.
- [2] G. Falqui and C.-M. Viallet, *Singularity, complexity, and quasi-integrability of rational mappings*. Comm. Math. Phys. **154** (1993), pp. 111–125. hep-th/9212105.
- [3] J. Diller and C. Favre, *Dynamics of bimeromorphic maps of surfaces*. Amer. J. Math. **123**(6) (2001), pp. 1135–1169.
- [4] Y. Yomdin, *Volume growth and entropy*. Israel J. Math. **57** (1987), pp. 285–299.
- [5] V.I. Arnold, *Dynamics of complexity of intersections*. Bol. Soc. Bras. Mat. **21** (1990), pp. 1–10.
- [6] A. Russakovskii and B. Shiffman, *Value distribution of sequences of rational mappings and complex dynamics*. Indiana U. Math. J. **46** (1997), pp. 897–932.
- [7] J. Hietarinta, N. Joshi, and F.W. Nijhoff. *Discrete Systems and Integrability*. Cambridge texts in applied mathematics. Cambridge University Press, (2016).
- [8] M.P. Bellon and C.-M. Viallet, *Algebraic Entropy*. Comm. Math. Phys. **204** (1999), pp. 425–437. chao-dyn/9805006.

- [9] R. Benedetto, P. Ingram, R. Jones, M. Manes, J.H.Silverman, and T.J. Tucker, *Current trends and open problem in arithmetic dynamics*. Bull. Amer. Math. Soc. (2019). DOI: <https://doi.org/10.1090/bull/1665>.
- [10] H. Sakai, *Rational Surfaces Associated with Affine Root Systems and Geometry of the Painlevé Equations*. Comm. Math. Phys. **220**(1) (2001), pp. 165–229.
- [11] K. Okamoto, *Sur les feuilletages associés aux équations du second ordre à points critiques fixes de P. Painlevé*. Jap. J. Math (1979), pp. 1–79.
- [12] J.J. Duistermaat. *Discrete Integrable Systems: QRT Maps and Elliptic Surfaces*. Springer Monographs in Mathematics. Springer New York, (2010).
- [13] J-C. Anglès d’Auriac, J-M. Maillard, and C-M. Viallet, *A classification of four-state spin edge Potts models*. J. Phys. A: Math. Gen. **35** (2002), pp. 9251–9272.
- [14] N. Joshi and C-M. Viallet, *Rational maps with invariant surfaces*. Journal of Integrable Systems **3**(1) (2018), p. xxy017. arXiv:1706.00173.
- [15] G. Gubbiotti, N. Joshi, D.T. Tran, and C-M. Viallet. *Complexity and integrability in 4D bi-rational maps with two invariants*. arXiv:1808.04942.
- [16] G.R.W. Quispel, J.A.G. Roberts, and C.J. Thompson, *Integrable Mappings and Soliton Equations*. Phys. Lett. **A 126** (1988), p. 419.
- [17] G.R.W. Quispel, J.A.G. Roberts, and C.J. Thompson, *Integrable Mappings and Soliton Equations II*. Physica **D34** (1989), pp. 183–192.
- [18] T. Takenawa, *Discrete dynamical systems associated with root systems of indefinite type*. Comm. Math. Phys. **224**(3) (2001), pp. 657–681.
- [19] C.T. McMullen, *Dynamics on blowups of the projective plane*. Publ. Math. Inst. Hautes Etudes Sci. **105** (2007), pp. 49–89.
- [20] A.S. Carstea and T. Takenawa. *Space of initial conditions and geometry of two 4-dimensional discrete Painlevé equations*. arXiv:1810.01664.
- [21] C. M. Viallet, *On the algebraic structure of rational discrete dynamical systems*. J. Phys. A: Math. Theor. **48** (2015), p. 16FT01.
- [22] M. Kanki, T. Mase, and T. Tokihiro, *On the coprimeness property of discrete systems without the irreducibility condition*. SIGMA **14** (2018), p. 065.
- [23] B. Hasselblatt and J. Propp, *Degree-growth of monomial maps*. Ergodic Theory and Dynamical Systems **27**(05) (2007), pp. 1375–1397. arXiv:math.DS/0604521.
- [24] V.E. Adler, A.I. Bobenko, and Yu.B. Suris, *Classification of integrable equations on quad-graphs. The consistency approach*. Comm. Math. Phys. **233**(3) (2003), pp. 513–543. arXiv:nlin.SI/0202024.
- [25] C-M. Viallet, *Integrable lattice maps: Q_V , a rational version of Q_4* . Glasgow Math. J. **51 A** (2009), pp. 157–163. arXiv:0802.0294.

- [26] M.K. Gizatullin, *Rational G -surface*. Izv. Akad. Nauk SSSR. Ser. Mat. **44**(1) (1980), pp. 110–144. English translation is in "Math. USSR, Izvestiya", vol. 16 (1981), no.1, p.103-134: MR.
- [27] M.P. Bellon, *Algebraic entropy of birational maps with invariant curves*. Lett. Math. Phys. **50** (1999), pp. 79–90.
- [28] M.J. Ablowitz, R. Halburd, and B. Herbst, *On the extension of the Painlevé property to difference equations*. Nonlinearity **13** (2000), pp. 889–905.