

ON THE FRACTIONAL SUSCEPTIBILITY FUNCTION OF PIECEWISE EXPANDING MAPS

MAGNUS ASPENBERG[†], VIVIANE BALADI^{††}, JUHO LEPPÄNEN^{††},
AND TOMAS PERSSON[†]

ABSTRACT. We associate to a perturbation (f_t) of a (stably mixing) piecewise expanding unimodal map f_0 a two-variable fractional susceptibility function $\Psi_\phi(\eta, z)$, depending also on a bounded observable ϕ . For fixed $\eta \in (0, 1)$, we show that the function $\Psi_\phi(\eta, z)$ is holomorphic in a disc $D_\eta \subset \mathbb{C}$ centered at zero of radius > 1 , and that $\Psi_\phi(\eta, 1)$ is the Marchaud fractional derivative of order η of the function $t \mapsto \mathcal{R}_\phi(t) := \int \phi(x) d\mu_t$, at $t = 0$, where μ_t is the unique absolutely continuous invariant probability measure of f_t . In addition, we show that $\Psi_\phi(\eta, z)$ admits a holomorphic extension to the domain $\{(\eta, z) \in \mathbb{C}^2 \mid 0 < \Re \eta < 1, z \in D_\eta\}$. Finally, if the perturbation (f_t) is horizontal, we prove that $\lim_{\eta \in (0, 1), \eta \rightarrow 1} \Psi_\phi(\eta, 1) = \partial_t \mathcal{R}_\phi(t)|_{t=0}$.

1. INTRODUCTION

1.1. Linear response and violation thereof. Response theory describes how the “physical measure” μ_t of a dynamical system f_0 responds to perturbations $t \mapsto f_t$, given a class of observables (test functions) ϕ . Classically, one studies *linear response*, where the goal is to express the derivative of $\mathcal{R}_\phi(t) := \int \phi d\mu_t$, for a fixed ϕ in a suitable class, in terms of $(\phi$ together with) f_0 , the vector field $v_0 := \partial_t f_t|_{t=0}$, and the measure μ_0 .

Linear response was first investigated [14, 23, 30, 19] for smooth hyperbolic dynamics (Anosov or Axiom A). In the smooth mixing hyperbolic case, the physical measure μ_t is the SRB measure [42] and corresponds to the fixed point of a transfer operator \mathcal{L}_t (whose dual preserves Lebesgue measure) on a suitable Banach space \mathcal{B} (see e.g. [5]). This fixed point is a simple isolated eigenvalue in the spectrum of \mathcal{L}_t , and linear response can be proved via perturbation theory for simple eigenvalues (see e.g. [5, §2.5 and §5.3]).

To avoid technicalities, we write the key formulas in the easy case of smooth expanding circle maps (see e.g. [4] for more details): Then, the physical measure is the unique absolutely continuous invariant probability measure, that is, $\mu_t = \rho_t dx$, with ρ_t smooth. The operator \mathcal{L}_t acts on smooth functions, and we have the trivial but key identity

$$(1) \quad \rho_t - \rho_0 = (I - \mathcal{L}_t)^{-1}(\mathcal{L}_t - \mathcal{L}_0)\rho_0.$$

Date: August 18, 2022.

We are grateful to Daniel Smania, some of whose ideas in the collaboration [11] were very useful here. Part of this work was carried out at the Centre for Mathematical Sciences, Lund University, during VB’s Knut and Alice Wallenberg Guest Professorship. VB’s and JL’s research is supported by the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (grant agreement No 787304).

Next, assuming that $v_0 = X_0 \circ f_0$, it is not hard to show that

$$\lim_{t \rightarrow 0} \frac{(\mathcal{L}_t - \mathcal{L}_0)\rho_0}{t} = -(X_0\rho_0)'.$$

Then, since $\int \phi \mathcal{L}_0^k(\psi) dx = \int (\phi \circ f_0^k)\psi dx$, we have,¹ for continuous ϕ (say),

$$\int \phi \cdot (1 - \mathcal{L}_0)^{-1}(X_0\rho_0)' dx = \sum_{k=0}^{\infty} \int \phi \cdot \mathcal{L}_0^k(X_0\rho_0)' dx = \sum_{k=0}^{\infty} \int (\phi \circ f_0^k)(X_0\rho_0)' dx.$$

Finally, using (1), we get the *fluctuation–dissipation formula* (expressing the derivative as a Green–Kubo [21] sum of decorrelations)

$$(2) \quad \partial_t \mathcal{R}_\phi(t)|_{t=0} = \partial_t \left(\int \phi d\mu_t \right) |_{t=0} = - \sum_{k=0}^{\infty} \int (\phi \circ f_0^k)(X_0\rho_0)' dx,$$

and, if ϕ is differentiable, integrating by parts, we get the *linear response formula*

$$(3) \quad \partial_t \mathcal{R}_\phi(t)|_{t=0} = \partial_t \left(\int \phi d\mu_t \right) |_{t=0} = \sum_{k=0}^{\infty} \int (\phi \circ f_0^k)' \cdot (X_0\rho_0) dx.$$

Exponential decay of the series in the right-hand side of (3) is not as apparent as in (2), since the derivative $(\phi \circ f_0^k)'(x) = \phi'(f_0^k(x)) \cdot (f_0^k)'(x)$ grows exponentially. However, the presence of this derivative should be expected² when describing response, and Ruelle [31] pointed out that it was meaningful to view the linear response formula (3) as the value at $z = 1$ of a natural power series, the *susceptibility function* of f_0 and $v_0 = X_0 \circ f_0$. In the present smooth one-dimensional expanding case, setting $z = e^{i\omega}$, the susceptibility function $\Psi_\phi(z)$ is the Fourier transform of the *response function* $k \mapsto \int X_0(\phi \circ f_0^k)' \rho_0 dx$, that is:

$$(4) \quad \Psi_\phi(z) := \sum_{k=0}^{\infty} z^k \int (\phi \circ f_0^k)' (X_0\rho_0) dx.$$

We have, integrating by parts,

$$\Psi_\phi(z) = - \sum_{k=0}^{\infty} z^k \int (\phi \circ f_0^k) (X_0\rho_0)' dx,$$

so that exponential mixing implies that the susceptibility function $\Psi_\phi(z)$ is holomorphic in a disc of radius larger than one, using that $X_0\rho_0$ is smooth.

In situations when the map $t \mapsto \mathcal{R}_\phi(t)$ is not differentiable (either due to bifurcations in the dynamics [27, 3, 9, 6], or to singularities [7, 29] of the test function ϕ), it is natural to consider *fractional response*, i.e., to investigate weaker moduli of continuity of this map. The simplest situation where linear response breaks down is that of piecewise expanding unimodal maps. In this case, the transfer operator has a spectral gap when acting on BV . However, the derivative ρ_0' of the invariant density ρ_0 involves a sum of Dirac masses, and thus does not belong to BV , or to a space on which decorrelations are summable. Keller [24] showed in 1982 that $|\rho_t - \rho_0|_{L^1} = O(|t| \log |t|)$. Examples

¹Note that $\int_{S^1} (X_0\rho_0)' dx = 0$ because the circle has no boundary.

²See e.g. the heuristic argument leading to [33, (2)].

of families (f_t) and smooth functions ϕ such that $|\mathcal{R}_\phi(t) - \mathcal{R}_\phi(0)| \geq |t| |\log |t||$ were described in [27] and [3, Theorem 6.1]. (See also the previous work of Ershov [17].) Baladi and Smania [9] showed that if the family f_t is *tangential* to³ the topological class of f_0 , then for any continuous function ϕ , the map $\mathcal{R}_\phi(t)$ is differentiable at $t = 0$. They also showed that when horizontality does not hold, then there exist smooth observables ϕ such that $\mathcal{R}_\phi(t)$ is *not* Lipschitz at $t = 0$. More recently, de Lima and Smania [16] showed central limit theorems which imply that for a generic map f_0 , if $v_0 = \partial_t f_t|_{t=0}$ is not horizontal, then for a generic ϕ , the map $t \mapsto \int \phi d\mu_t$ cannot be Lipschitz on a set of parameters t of positive Lebesgue measure. (See also [15] for a related result.)

In the piecewise expanding unimodal case, the susceptibility function $\Psi_\phi(z)$ can also be defined by the formal power series (4), and it has been studied in [3, 9, 8]. In particular [8], if the postcritical orbit is dense (a generic condition) then $\Psi_\phi(z)$ has a strong natural boundary on the unit circle, while if the perturbation $v_0 = X_0 \circ f$ is horizontal and, in addition (a generic condition) the postcritical orbit is Birkhoff typical, then the nontangential limit of $\Psi_\phi(z)$ as z tends to 1 coincides with $\partial_t \mathcal{R}_\phi(t)|_{t=0}$.

In the present paper, we introduce and study a two-variable fractional susceptibility function $\Psi_\phi(\eta, z)$, for $\Re \eta \in (0, 1)$, in the setting of piecewise expanding unimodal maps. The initial motivation for this work comes from the following paradox:

In 2005, Ruelle and Jiang considered [32, 22] finite Misiurewicz–Thurston (MT) parameters t_0 in the quadratic family $f_t(x) = t - x^2$. By definition of MT, there exists a repelling periodic point x_0 and $M \geq 2$ such that $f_{t_0}^M(0) = x_0$. Such parameters t_0 are Collet–Eckmann, so f_{t_0} admits a unique absolutely continuous invariant probability measure μ_{t_0} . The map f_{t_0} also has a finite Markov partition, which simplifies the analysis. Ruelle and Jiang proved that, for any C^1 observable ϕ , the susceptibility function $\Psi_\phi(z)$ defined by (4) is meromorphic in the whole complex plane, and that $z = 1$ is not a pole. Since $\Psi_\phi(1)$ is the natural candidate for the derivative of $\mathcal{R}_\phi(t)$, this raised the hope that there could exist a “large” subset Ω of Collet–Eckmann parameters, containing t_0 , and such that $t \mapsto \mathcal{R}_\phi(t)$ would be differentiable in the sense of Whitney on Ω at t_0 . However, Baladi, Benedicks, and Schnellmann [6] later showed that for any mixing (non horizontal⁴) MT parameter t_0 , there exist a C^∞ observable ϕ , a sequence $t_n \rightarrow t_0$ of Collet–Eckmann parameters (with bounded constants), and $C > 1$ such that

$$(5) \quad \frac{\sqrt{|t_n - t_0|}}{C} \leq |\mathcal{R}_\phi(t_n) - \mathcal{R}_\phi(t_0)| \leq C \sqrt{|t_n - t_0|}, \quad \forall n.$$

It is not known whether t_0 is a Lebesgue density point in the set of t_n such that (5) holds. Also, the analogue of the de Lima–Smania [16] central limit theorems is not known in this setting. However, we expect that a similar CLT holds, and in particular that $\Psi_\phi(1)$ *cannot* be interpreted as the derivative of $\mathcal{R}_\phi(t)$ in the sense of Whitney on a set of parameters containing t_0 as a Lebesgue density point. The fact that $\Psi_\phi(z)$ is holomorphic at $z = 1$ can thus be viewed as a paradox.

Baladi and Smania [11] very recently introduced two-variable fractional susceptibility functions $\Psi_\phi(\eta, z)$ for the quadratic family, with the goal of resolving this paradox. The

³We refer to the beginning of §4.4 for a definition of tangentiality, which is equivalent to the *horizontality* condition (16) on v_0 .

⁴All MT parameters are non horizontal by [1] or [26].

piecewise expanding case is a toy model for the quadratic setting, and indeed, several ideas previously developed for piecewise expanding families [9] were crucial to obtain the breakthrough result (5) of [6] for the quadratic family. Although we believe it is interesting in its own right, the analysis carried out in the present paper for this toy model can also be viewed as a “proof of concept,” establishing the feasibility of the fractional susceptibility function approach.

1.2. Informal statement of the results. We next describe briefly our main results. It is well-known that there is no canonical notion of a fractional derivative, see [35] and [28] for a presentation of the theory. We use here the Marchaud fractional derivative, defined for suitable functions g by

$$(M_t^\eta g(t))|_{t=0} = \frac{\eta}{2\Gamma(1-\eta)} \int_{-\infty}^{\infty} \frac{g(t) - g(0)}{|t|^{1+\eta}} \text{sign}(t) dt,$$

where Γ denotes Euler’s Gamma function. Our motivation for using this particular fractional derivative is threefold: First, $M^\eta g$ can be well-defined even if g does not decay at infinity. Second, the Marchaud fractional derivative of a constant function vanishes, while this is not the case for other fractional derivatives, in particular for the Bessel potential derivative defined by $\mathcal{F}^{-1}(1 + |\xi|^2)^{\eta/2} \mathcal{F}g$, where \mathcal{F} is the Fourier transform. Finally, the expression of Marchaud derivatives in terms of differences $g(t) - g(-t) = g(t) - g(0) + g(0) - g(-t)$ is convenient in view of (1). Nevertheless, we expect that fractional susceptibility functions defined via (e.g.) Bessel, or Riemann–Liouville fractional derivatives would enjoy similar properties as those we establish here using the Marchaud derivative.

We define for $\eta \in (0, 1)$ the *fractional susceptibility function* of the perturbation (f_t) and the observable $\phi \in L^\infty$, to be the formal power series

$$(6) \quad \Psi_\phi(\eta, z) := \sum_{k=0}^{\infty} z^k \int_I \frac{\eta}{2\Gamma(1-\eta)} \int_{-\infty}^{\infty} (\phi \circ f_t^k)(x) \cdot \frac{((\mathcal{L}_t - \mathcal{L}_0)\rho_0)(x)}{|t|^{1+\eta}} \text{sign}(t) dt dx,$$

and the *frozen fractional susceptibility function* of (f_t) and ϕ to be the formal power series

$$(7) \quad \Psi_\phi^{\text{fr}}(\eta, z) := \sum_{k=0}^{\infty} z^k \int_I \frac{\eta}{2\Gamma(1-\eta)} (\phi \circ f_0^k)(x) \int_{-\infty}^{\infty} \frac{((\mathcal{L}_t - \mathcal{L}_0)\rho_0)(x)}{|t|^{1+\eta}} \text{sign}(t) dt dx.$$

In both susceptibility functions, \mathcal{L}_t is the transfer operator $\mathcal{L}_t\varphi(x) = \sum_{f_t(y)=x} \frac{\varphi(y)}{|f_t'(y)|}$.

Our first main result (Theorem 2.3) says that if ϕ is bounded, then for any $\eta \in (0, 1)$ the improper integrals in the two above formal power series are convergent, and that $\Psi_\phi(\eta, z)$ and $\Psi_\phi^{\text{fr}}(\eta, z)$ are holomorphic functions of z in a disc D_η of radius (which may depend on f_0) larger than one. In addition,

$$\Psi_\phi^{\text{fr}}(\eta, z) = \int_I \phi \cdot (I - z\mathcal{L}_0)^{-1} M_t^\eta(\mathcal{L}_t\rho_0(x))|_{t=0} dx = \sum_k z^k \int_I (\phi \circ f_0^k) M_t^\eta(\mathcal{L}_t\rho_0(x))|_{t=0} dx,$$

and

$$\Psi_\phi(\eta, 1) = M_t^\eta \mathcal{R}_\phi(t)|_{t=0}.$$

Remark 1.1. *Our lower bound for the radius of D_η tends to 1 as $\eta \rightarrow 1$. If the critical point of f_0 is preperiodic, then we expect $\Psi_\phi(\eta, z)$ to be holomorphic in a disc of radius strictly larger than 1 and meromorphic in the entire complex plane for all $0 \leq \eta \leq 1$. However, we believe that, generically, the radius of convergence of $\Psi_\phi(\eta, z)$ should tend to 1 as $\eta \rightarrow 1$.*

As a corollary of our main theorem and the results of [9], we obtain (Corollary 2.4) that, if v_0 is horizontal and ϕ is continuous, then

$$\lim_{\eta \rightarrow 1} \Psi_\phi(\eta, 1) = \lim_{\eta \rightarrow 1} \Psi_\phi^{\text{fr}}(\eta, 1) = \Psi_\phi(1) = \partial_t \mathcal{R}_\phi(t)|_{t=0}.$$

Our second result, Theorem 2.6, is about more general fractional moduli of continuity: We consider there weighted Marchaud derivatives, replacing $|t|^{-1-\eta}$ by $(\log |t|)^{-\beta} |t|^{-1-\eta}$. Applying Theorem 2.6 to $\beta > 1$ and $\eta = 1$ gives a modulus of continuity $|t| \log |t|^\beta$, almost reproducing the $|t| \log |t|$ estimates from [24, 27, 3]. Applying Theorem 2.6 to $\beta < 0$, we show (Corollary 2.7) that $\Psi_\phi(\eta, z)$ and $\Psi_\phi^{\text{fr}}(\eta, z)$ are holomorphic in the domain $\{(\eta, z) \in \mathbb{C}^2 \mid 0 < \Re \eta < 1, z \in D_\eta\}$.

The fractional susceptibility functions in (6) and (7) are of “fluctuation–dissipation” type. It is tempting to consider the *response fractional susceptibility function*⁵

$$\begin{aligned} \Psi_\phi^{\text{rsp}}(\eta, z) &:= - \sum_{k=0}^{\infty} z^k \int_I (\phi \circ f_0^k) \cdot M_x^\eta(X_0 \rho_0) \, dx \\ (8) \qquad \qquad &= \sum_{k=0}^{\infty} z^k \int_I (M_x^\eta(\phi \circ f_0^k)) \cdot X_0 \rho_0 \, dx, \end{aligned}$$

where the Marchaud derivative is now taken with respect to x . The arguments in the present paper easily show that for all $\eta \in (0, 1)$ the function $\Psi_\phi^{\text{rsp}}(\eta, z)$ is holomorphic in a disc of radius larger than 1 for all bounded ϕ , and that in the horizontal case we have $\lim_{\eta \rightarrow 1} \Psi_\phi^{\text{rsp}}(\eta, 1) = \partial_t \mathcal{R}_\phi|_{t=0}$. The function $\Psi_\phi^{\text{rsp}}(\eta, z)$ is at first sight the most seductive fractional susceptibility function. However, $\Psi_\phi^{\text{rsp}}(\eta, 1)$ has no reason to coincide in general with any fractional derivative of order η of $\mathcal{R}_\phi(t)$ (except in the limit $\eta \rightarrow 1$, even in the linear examples studied in §2.3). In this respect, $\Psi_\phi^{\text{rsp}}(\eta, z)$ is not better than the frozen susceptibility function $\Psi_\phi^{\text{fr}}(\eta, z)$. Keeping also in mind that the goal of this fractional approach is to resolve the paradox described above for the quadratic family, we focus on the definitions (6) and (7) in the present paper, referring to [11] for more on the fractional response susceptibility function.

Observe also that fractional derivatives do not enjoy a Leibniz formula with finitely many terms for the derivative of a product, so we cannot expect fractional susceptibility functions to be as well-behaved as ordinary ones.

We next make a few comments about our method of proof. We shall consider transfer operators acting on Sobolev spaces $\mathcal{H}^{\tau, p}$ with $p > 1$, close to 1, and $0 < \tau < 1/p$. These spaces give us more flexibility than the BV spaces classically used for piecewise expanding interval maps. We exploit the bounds of Thomine [38] for the essential spectral radius of \mathcal{L}_t on such spaces $\mathcal{H}^{\tau, p}$. In particular, we have $\rho_0 \in \mathcal{H}^{\tau, p}$ (see the beginning of

⁵The equality (8) follows by integration by parts for the Marchaud derivative [35, (6.27)] for $\phi \in C^1$.

§3.1), so that, for each $\eta < 1$, the function $M_x^\eta \rho_0$ belongs to a Sobolev space on which the transfer operator \mathcal{L}_t associated to f_t has a spectral gap (see (22)). In order to apply the stable exponential decorrelation result of Keller–Liverani [25], we need Lasota–Yorke bounds which are uniform in t . Such bounds were known for the BV norm, but we have to carry out the corresponding estimates for the Sobolev spaces.

The paper is organised as follows: Section 2 starts with definitions and formal statements. In §2.1, we state precisely Theorem 2.3 on fractional susceptibility functions, followed by its consequence (Corollary 2.4) in the horizontal case. In §2.2 we state Theorem 2.6 about other moduli of continuity and Corollary 2.7 about holomorphic extensions of the fractional susceptibility functions. In §2.3, we discuss two linear examples. Section 3 introduces the key tools: In §3.1, we recall properties of the transfer operator acting on Sobolev spaces, from the work of Thomine [38]. In §3.2, we discuss stability of mixing and mixing rates for good families, recalling in particular the results of Keller and Liverani [25] that we shall use, and stating the relevant technical lemmas (Lemma 3.2 and Lemma 3.1). Subsection 3.3 contains the proof of the perturbation Lemma 3.2 for Sobolev spaces which is needed to apply results of Keller and Liverani. In Section 4, we prove Theorems 2.3 and 2.6, as well as Corollaries 2.4 and 2.7, using the techniques presented in Section 3. Appendix A contains the simple proof that the limit of the Marchaud derivatives as $\eta \rightarrow 1$ is the ordinary derivative. In Appendix B, we show the uniform Lasota–Yorke estimates (Lemma 3.1) needed to apply the results of Keller and Liverani to Sobolev spaces.

Throughout, we shall use the notation C for a finite positive constant which can vary from place to place.

2. DEFINITIONS AND FORMAL STATEMENT OF RESULTS

2.1. Fractional susceptibility functions for piecewise expanding interval maps.

Let I be the compact interval $[-1, 1]$ and let $r \in \{2, 3\}$. We say that a continuous map $f: I \rightarrow I$ is a piecewise C^r unimodal map if there exists $-1 < c < 1$ such that f is increasing on $I_+ = [-1, c]$, decreasing on $I_- = [c, 1]$, and $f|_{I_\sigma}$ extends as a C^r map denoted f_σ to a neighbourhood \tilde{I}_σ of I_σ . If, in addition, $\lambda(f) := \inf_{\sigma=\pm} |f'_\sigma(x)| > 1$, we say that $f: I \rightarrow I$ is a piecewise C^r expanding unimodal map, and we define

$$(9) \quad \lambda_n(f) := \inf_{\vec{\sigma} \in \{\pm\}^n} \inf_x |(f_{\vec{\sigma}}^n)'(x)|, \quad \Lambda(f) = \left(\lim_{n \rightarrow \infty} \lambda_n(f)^{-1/n} \right)^{-1},$$

for all $n \geq 1$, where $f_{\vec{\sigma}}^n$ is the composition of the maps f_{σ_i} , $i = 1, \dots, n$, and the points x in (9) are restricted to those $x \in \tilde{I}_{\sigma_1}$ for which the composition exists. Following [9], we say that a piecewise C^r expanding unimodal map f is *good*⁷ if, either c is not periodic for f , or $|(f^{P_f})'(c)| > 2$ where $P_f \geq 2$ is the minimal period of c . Finally, we say that a piecewise C^r expanding unimodal map f is *mixing* if f is topologically mixing on $[f^2(c), f(c)]$.

⁶The results of Thomine [38] hold for piecewise $C^{1+\eta_0}$ maps, up to replacing the condition $\tau < 1/p$ there by $\tau < \max(\eta_0, 1/p)$. It would be interesting to adapt our results to this setting, for $0 < \eta < \eta_0$.

⁷Goodness will ensure the uniform Lasota–Yorke Lemma 3.1 and thus stable mixing. It is not the weakest possible condition, see e.g. [18, Example 5.6] but some assumption is needed [24, Remark 15]. See also [10, Proposition 2.1].

Definition 2.1 (Perturbation (f_t) of a piecewise C^r expanding unimodal map). *Let $r \in \{2, 3\}$ and let f be a piecewise C^r expanding unimodal map. Given $\varepsilon > 0$ and a family $(f_t)_{|t| \leq \varepsilon}$ of piecewise C^r unimodal maps (for fixed \tilde{I}_\pm) is called a C^r perturbation of $f_0 = f$ if the C^r norm of the extension $f_{t,\sigma}$ of f_t to \tilde{I}_\pm is uniformly bounded, with $\|(f - f_t)|_{\tilde{I}_\sigma}\|_{C^{r-1}} = O(|t|)$ as $t \rightarrow 0$ for $\sigma = \pm$. If all f_t are topologically conjugated, or if c is not periodic for f , set $\Lambda := \inf_{|t| < \varepsilon} \Lambda(f_t)$, otherwise, if f is good, with c periodic of minimal period $P_f \geq 2$, set*

$$(10) \quad \Lambda^{-1} := \sup_{|t| < \varepsilon} \lim_{k \rightarrow \infty} \left(\frac{\lambda_{kP_f}(f_t)}{2^k} \right)^{-\frac{1}{kP_f}};$$

in all these cases, up to taking a smaller ε , we may and shall assume that $\Lambda > 1$.

Any piecewise C^2 expanding unimodal map f_t admits a unique absolutely continuous invariant probability measure $\mu_t = \rho_t dx$. The measure μ_t is ergodic, and it is mixing if f_t is mixing. The density ρ_t is the unique fixed point (see e.g. [2]) of the transfer operator \mathcal{L}_t defined on BV by

$$\mathcal{L}_t \varphi(x) = \sum_{f_t(y)=x} \frac{\varphi(y)}{|f_t'(y)|}.$$

We now recall the definition of the Marchaud fractional derivatives in order to introduce fractional susceptibility functions. Let $0 < \eta < 1$, and set $\Gamma_\eta = \frac{\eta}{\Gamma(1-\eta)}$ where Γ is Euler's function. Let $g: \mathbb{R} \rightarrow \mathbb{C}$ be a bounded globally $\bar{\eta}$ -Hölder function with $\bar{\eta} \in (0, 1)$. Recall [35, p. 110, Theorem 5.9] that for any $\eta \in (0, \bar{\eta})$, the left-sided and right-sided Marchaud derivatives of g at $t_0 \in \mathbb{R}$ are

$$M_+^\eta g(t_0) = \Gamma_\eta \int_0^\infty \frac{g(t_0) - g(t_0 - t)}{t^{1+\eta}} dt, \quad M_-^\eta g(t_0) = \Gamma_\eta \int_0^\infty \frac{g(t_0) - g(t_0 + t)}{t^{1+\eta}} dt.$$

The two-sided Marchaud derivative is then defined by

$$\begin{aligned} M^\eta g(t_0) &:= \frac{1}{2}(M_+^\eta g(t_0) - M_-^\eta g(t_0)) \\ &= \frac{\Gamma_\eta}{2} \left(\int_0^\infty \frac{g(t_0) - g(t_0 - t)}{t^{1+\eta}} dt - \int_0^\infty \frac{g(t_0) - g(t_0 + t)}{t^{1+\eta}} dt \right) \\ &= \frac{\Gamma_\eta}{2} \int_{-\infty}^\infty \frac{g(t_0 + t) - g(t_0)}{|t|^{1+\eta}} \text{sign}(t) dt. \end{aligned}$$

The choice of the normalisation Γ_η ensures the following key property:

Lemma 2.2. *Assume that g is bounded on \mathbb{R} and differentiable at t_0 . Then*

$$\lim_{\eta \uparrow 1} M_+^\eta g(t_0) = g'(t_0) \quad \text{and} \quad \lim_{\eta \uparrow 1} M_-^\eta g(t_0) = -g'(t_0), \quad \text{so that} \quad \lim_{\eta \uparrow 1} M^\eta g(t_0) = g'(t_0).$$

(The above lemma is certainly well-known, we provide a proof in Appendix A.)

Given a C^2 perturbation $(f_t)_{|t| \leq \varepsilon}$ of a piecewise C^2 expanding unimodal map f , and given $\varepsilon_1 \leq \varepsilon$, we put⁸

$$(11) \quad f_t^{(\varepsilon_1)} = f_t, \forall |t| \leq \varepsilon_1, \quad f_t^{(\varepsilon_1)} = f_{\varepsilon_1}, \forall t \geq \varepsilon_1, \quad f_t^{(\varepsilon_1)} = f_{-\varepsilon_1}, \forall t \leq -\varepsilon_1.$$

⁸See also (26) for an alternative approach.

Then, for any function $\phi \in L^1(I)$, we define the η -fractional susceptibility function of $(f_t^{(\varepsilon_1)})$ for the observable ϕ to be the formal power series

$$(12) \quad \Psi_\phi(\eta, z) = \sum_{k=0}^{\infty} \frac{\Gamma_\eta}{2} z^k \int_I \phi \int_{-\infty}^{\infty} \mathcal{L}_t^k \left(\frac{(\mathcal{L}_t - \mathcal{L}_0)\rho_0}{|t|^{1+\eta}} \right) \text{sign}(t) dt dx,$$

and we define the frozen η -fractional susceptibility function of $(f_t^{(\varepsilon_1)})$ and ϕ to be the formal series

$$(13) \quad \Psi_\phi^{\text{fr}}(\eta, z) = \sum_{k=0}^{\infty} \frac{\Gamma_\eta}{2} z^k \int_I \phi \int_{-\infty}^{\infty} \mathcal{L}_0^k \left(\frac{(\mathcal{L}_t - \mathcal{L}_0)\rho_0}{|t|^{1+\eta}} \right) \text{sign}(t) dt dx.$$

The coefficient of z^k in each of the two formal power series above is a sum of improper integrals, for $t \in (-\infty, 0)$ and $t \in (0, \infty)$. We shall see in the proof of Theorem 2.3 that each integral converges, if ϕ belongs to L^q for large enough q . In particular, using Fubini, we recover the formulas (6) and (7) stated in the introduction.

We are now ready to state our main result. Recall $\Lambda > 1$ from Definition 2.1. For a function $g(x, t)$ of two real variables, we denote by $M^\eta g|_{t=t_0}$ the Marchaud derivative of g in the t variable, at $t = t_0$.

Theorem 2.3 (Fractional susceptibility function). *Let $(f_t)_{|t| \leq \varepsilon}$ be a C^2 perturbation of a mixing piecewise C^2 expanding unimodal map f . Assume that either f is good or that all the f_t are topologically conjugated to f . Then there exist $\kappa < 1$ (depending only on f_0) and $\varepsilon_1 \in (0, \varepsilon)$ such that for any $0 < \eta < 1$, and for any $\phi \in L^q(I)$ with $q > (1 - \eta)^{-1}$, the following holds for the fractional susceptibility functions of $(f_t^{(\varepsilon_1)})$ and ϕ :*

- (a) $\Psi_\phi(\eta, z)$ and $\Psi_\phi^{\text{fr}}(\eta, z)$ are holomorphic in the open disc of radius $\min(\Lambda^{1-\eta}, \kappa^{-1})$.
- (b) $\Psi_\phi(\eta, 1) = M^\eta \left(\int_I \phi(x) \rho_t(x) dx \right) |_{t=0}$.
- (c) $\Psi_\phi^{\text{fr}}(\eta, z) = \int_I \phi(x) \cdot ((I - z\mathcal{L}_0)^{-1} M^\eta(\mathcal{L}_t \rho_0)|_{t=0})(x) dx$ for all $|z| \leq 1$.

As explained in §1.2, formula (b) in the above theorem can be viewed as the key result of this work. The proof of Theorem 2.3 will be given in Section 4, after we introduce some necessary tools in Section 3. We next make a few remarks about the statement:

Clearly, if $\phi \in L^\infty$, then we can consider all values of $\eta \in (0, 1)$.

The proof of Theorem 2.3 actually shows that for $\phi \in L^q(I)$, the response function is Hölder continuous with exponent η provided that $q > (1 - \eta)^{-1}$.

We do not claim that the holomorphy radius given in claim a) is optimal. However, we expect that, generically, the maximal holomorphic extension radius of $\Psi_\phi(\eta, z)$ tends to one as $\eta \rightarrow 1$. It is unclear whether the frozen fractional susceptibility function is holomorphic in a disc of radius larger than one, uniformly in $\eta \rightarrow 1$.

Claim c) implies that, for any $\eta \in (0, 1)$, and all $|z| \leq 1$

$$(14) \quad \Psi_\phi^{\text{fr}}(\eta, z) = \sum_{k=0}^{\infty} z^k \int_I \phi(f_0^k(x)) \cdot (M^\eta(\mathcal{L}_t \rho_0)|_{t=0})(x) dx.$$

For $z = 1$ this is reminiscent of the fluctuation–dissipation formula for linear response (see e.g. [7]). Note, however, that we cannot integrate by parts (in spite of [35, (6.27)])

because the Marchaud derivative is with respect to the parameter t . See Corollary 2.4 for more information on the frozen susceptibility function in the “horizontal” case.

To state an interesting corollary of our Theorem 2.3, letting H_u denote the Heaviside jump at $u \in \mathbb{R}$, i.e., $H_u(x) = -1$ if $x < u$, while $H_u(x) = 0$ if $x > u$, and $H_u(u) = -1/2$, and setting $c_k = f^k(c)$, we recall that [3, Proposition 3.3] we may decompose ρ_0 as $\rho_0 = \rho_0^{\text{reg}} + \rho_0^{\text{sal}}$, where the regular part ρ_0^{reg} is differentiable and supported in I , with derivative in BV , and the saltus (or singular) part $\rho_0^{\text{sal}} = \sum_{k=1}^{N_f} s_k H_{c_k}$ where $N_f = \#\{c_k \mid k \geq 1\} \in [2, \infty]$. In addition, if the critical point c is not periodic, we have

$$(15) \quad \rho_0^{\text{sal}} = \sum_{k=1}^{\infty} \bar{s}_k H_{c_k}, \quad \text{where } \bar{s}_1 = -\lim_{x \uparrow c_1} \rho_0(x) < 0, \quad \bar{s}_k = \frac{s_1}{(f^{k-1})'(c_1)}.$$

Note that if N_f is finite but c is not periodic (it is then preperiodic) then the jump s_j at c_j is given by the sum of all \bar{s}_k for k such that $c_k = c_j$.

We say that a bounded function v is *horizontal* for f if, setting $P_f = N_f$ if c is periodic with minimal period $P_f \geq 2$, and $P_f = \infty$ otherwise, we have

$$(16) \quad \sum_{k=0}^{P_f-1} \frac{v(c_k)}{(f^k)'(c_1)} = 0.$$

If v is not horizontal, we say that v is *transversal*. In the horizontal case, we have (Corollary 2.4 is proved in Section 4.4):

Corollary 2.4 (Fractional susceptibility of horizontal perturbations). *Let $(f_t)_{|t| \leq \varepsilon}$ be a C^3 perturbation of a mixing piecewise C^3 expanding unimodal map f . Assume that either f is good or that all the f_t are topologically conjugated to f . Assume in addition⁹ that $f_t(-1) = f_t(1) = -1$ for all t , that $v_0 = \partial_t f_t|_{t=0} = X_0 \circ f_0$ is horizontal for f_0 , where X_0 is C^2 on $f(I)$, and, finally that $(x, t) \mapsto f_t(x)$ extends as a C^2 map to $(\tilde{I}_+ \cup \tilde{I}_-) \times [-\varepsilon, \varepsilon]$. Then, there exists $\varepsilon_2 \in (0, \varepsilon)$ such that for any $\phi \in C^0$, the following holds for the fractional susceptibility function of $(f_t^{(\varepsilon_2)})$ and ϕ :*

$$(17) \quad \begin{aligned} \lim_{\eta \uparrow 1} \Psi_\phi(\eta, 1) &= \partial_t \left(\int \phi(x) \rho_t(x) dx \right) \Big|_{t=0} \\ &= - \sum_{j=1}^{P_f} \phi(c_j) \sum_{k=1}^j \bar{s}_k X_0(c_k) - \int \phi \cdot (I - \mathcal{L}_0)^{-1} (X_0' \rho_0^{\text{sal}} + (X_0 \rho_0^{\text{reg}})') dx. \end{aligned}$$

Assume furthermore that c is not periodic for f . Then, we have

$$(18) \quad \lim_{\eta \uparrow 1} ((\Psi_\phi(\eta, 1) - \Psi_\phi^{\text{ff}}(\eta, 1))) = 0.$$

Remark 2.5. *The first term of (17) can be rewritten as*

$$- \sum_{j=1}^{P_f} \phi(c_j) \sum_{k=1}^j \bar{s}_k X_0(c_k) = - \int \phi \alpha(\rho_0^{\text{sal}})', \quad \text{where } \alpha(x) := - \sum_{j=0}^{P_f-1} \frac{v(f^j(x))}{(f^{j+1})'(x)}.$$

⁹This implies $v_0(-1) = v_0(1) = 0$ and if $f(c) = 1$ then $v_0(c) = 0$.

Indeed, α is the solution (which [9, Lemma 2.2, Remark 2.3, Proposition 2.4] is unique and continuous under the assumptions of the corollary) of the twisted cohomological equation

$$(19) \quad X_0(f(x)) = v_0(x) = \alpha(f(x)) - f'(x)\alpha(x), \quad x \neq c, \quad X_0(c_1) = v(c) = \alpha(c_1),$$

so that, for $j \leq P_f$,

$$(20) \quad \sum_{k=1}^j \bar{s}_k X_0(c_k) = \bar{s}_1 \sum_{k=1}^j \frac{X_0(c_k)}{(f^{k-1})'(c_1)} = \bar{s}_1 \left(X_0(c_1) - \alpha(c_1) + \frac{\alpha(c_j)}{(f^{j-1})'(c_1)} \right) = \bar{s}_j \alpha(c_j).$$

2.2. Fractional moduli of continuity and Keller's $x \log x$ bound. In the definition of the Marchaud derivatives $M_+^\eta g$ and $M_-^\eta g$ of a function g , we may replace $t^{1+\eta}$ with other weights. Suppose for instance that $\ell: [0, \infty) \rightarrow \mathbb{C}$ is a continuous function and that $\gamma \in [0, 1)$ is a constant such that

$$(21) \quad \ell(0) = 0, \quad \int_0^1 \frac{t^\gamma}{|\ell(t)|} dt < \infty \quad \text{and} \quad \int_1^\infty \frac{1}{|\ell(t)|} dt < \infty.$$

We then define the right and left-sided ℓ -Marchaud derivatives of a bounded γ -Hölder function g by

$$M_+^{(\ell)} g(t_0) = \int_0^\infty \frac{g(t_0) - g(t_0 - t)}{\ell(t)} dt, \quad M_-^{(\ell)} g(t_0) = \int_0^\infty \frac{g(t_0) - g(t_0 + t)}{\ell(t)} dt.$$

We define the two-sided ℓ -Marchaud derivative by $M^{(\ell)} g = \frac{1}{2}(M_+^{(\ell)} g - M_-^{(\ell)} g)$. Finally, we define the ℓ -susceptibility function to be the formal power series

$$\Psi_\phi((\ell), z) = \sum_{k=0}^\infty z^k \int_I \phi \int_{-\infty}^\infty \left(\mathcal{L}_t^k \frac{(\mathcal{L}_t - \mathcal{L}_0)\rho_0}{\ell(|t|)} \right) \text{sign}(t) dt dx.$$

Similarly, we define the frozen ℓ -susceptibility function by the formal power series

$$\Psi_\phi^{\text{fr}}((\ell), z) = \sum_{k=0}^\infty z^k \int_I \phi \int_{-\infty}^\infty \left(\mathcal{L}_0^k \frac{(\mathcal{L}_t - \mathcal{L}_0)\rho_0}{\ell(|t|)} \right) \text{sign}(t) dt dx.$$

The following theorem is proved in almost the same way as Theorem 2.3 (see Section 4.2).

Theorem 2.6 (Generalized fractional susceptibility function). *Let $(f_t)_{|t| \leq \varepsilon}$ be a C^2 perturbation of a mixing piecewise C^2 expanding unimodal map f . Assume that either f is good or all the f_t are topologically conjugated to f . Let ℓ and $\gamma \geq 0$ be such that (21) holds. Then, for any $q \geq (1 - \gamma)^{-1}$ and any $\phi \in L^q$, the following holds:*

- (a) *the susceptibility functions $\Psi_\phi((\ell), z)$ and $\Psi_\phi^{\text{fr}}((\ell), z)$ are well-defined and holomorphic in a disc of radius strictly larger than one.*
- (b) *$\Psi_\phi((\ell), 1) = M^{(\ell)}(\int_I \phi(x)\rho_t(x) dx)|_{t=0}$.*
- (c) *$\Psi_\phi^{\text{fr}}((\ell), z) = \int_I \phi(x) \cdot ((I - z\mathcal{L}_0)^{-1} M^{(\ell)}(\mathcal{L}_t \rho_0)|_{t=0})(x) dx$ for all $|z| \leq 1$.*

For instance, (cf. the power-logarithmic kernel operators discussed in [35, §21]) we may fix $0 < \eta < 1$ and $\beta \geq 0$ and set

$$\ell(t) = \ell_{\eta, \beta}(t) = t^{1+\eta} |\min\{-1, \log t\}|^\beta.$$

Then, if $\phi \in L^q$, $\eta = 1 - \frac{1}{q}$ and $\beta > 1$, Theorem 2.6 for $\gamma < \eta$ implies

$$M^{(\ell_{\eta,\beta})} \left(\int_I \phi(x) \rho_t(x) dx \right) \Big|_{t=0} = \Psi_\phi((\ell_{\eta,\beta}), 1).$$

In particular, if $q = \infty$, we may take $\eta = 1$ and $\beta > 1$ arbitrarily close to 1 which is reminiscent of Keller's [24] $t |\log |t||$ modulus of continuity.

Another consequence of Theorem 2.6 is that we may consider the Marchaud derivative $M^\eta \left(\int_I \phi(x) \rho_t(x) dt \right) \Big|_{t=0}$ and the susceptibility functions $\Psi_\phi(\eta, z)$ and $\Psi_\phi^{\text{fr}}(\eta, z)$ for non-real values of η . A corollary of Theorem 2.6 is the following result, which we prove in Section 4.3.

Corollary 2.7 (Holomorphic extension in (η, z)). *Let $(f_t)_{|t| \leq \varepsilon}$ be a C^2 perturbation of a mixing piecewise C^2 expanding unimodal map f . Assume that either f is good or all the f_t are topologically conjugated to f , and let $\kappa < 1$ and ε_1 be from Theorem 2.3. Let $\phi \in L^q$, for some $q > 1$. Then the function $\eta \mapsto M^\eta \left(\int_I \phi(x) \rho_t(x) dt \right) \Big|_{t=0}$ is holomorphic in the strip $0 < \Re \eta < 1 - \frac{1}{q}$, and the functions $(\eta, \zeta) \mapsto \Psi_\phi(\eta, z)$ and $(\eta, \zeta) \mapsto \Psi_\phi^{\text{fr}}(\eta, z)$ are holomorphic in the domain $\{(\eta, z) \mid 0 < \Re \eta < 1 - \frac{1}{q}, |z| < \min(\kappa, \Lambda^{-\frac{1}{q}})\}$.*

2.3. Two linear examples. We illustrate our definitions with two families of linear tent maps. One simplifying feature is that ρ_t^{reg} vanishes identically for all t in both examples. To study ρ_0^{sal} , we shall use that for any $x \neq u$, we have

$$(22) \quad M_t^\eta H_t(x) \Big|_{t=u} = -\frac{1}{2\Gamma(1-\eta)} \frac{1}{|x-u|^\eta}.$$

To prove (22), observe that if $x - u > 0$, then

$$H_{t+u}(x) - H_u(x) = H_t(x-u) - H_0(x-u) = \begin{cases} 0 & \text{if } t < x-u \\ -1 & \text{if } t > x-u. \end{cases}$$

Similarly, if $x - u < 0$, then

$$H_{t+u}(x) - H_u(x) = H_t(x-u) - H_0(x-u) = \begin{cases} 1 & \text{if } t < x-u \\ 0 & \text{if } t > x-u. \end{cases}$$

It follows that

$$\begin{aligned} M_t^\eta H_t(x) \Big|_{t=u} &= \frac{\Gamma_\eta}{2} \int_{-\infty}^{\infty} \frac{H_{t+u}(x) - H_u(x)}{|t|^{1+\eta}} \text{sign}(t) dt \\ &= -\frac{\Gamma_\eta}{2} \int_{|x-u|}^{\infty} \frac{1}{|t|^{1+\eta}} dt = -\frac{1}{2\Gamma(1-\eta)} \frac{1}{|x-u|^\eta}, \end{aligned}$$

as claimed. Note also that since $H_{u+t}(x) = H_u(x-t)$ we have

$$(23) \quad M_y^\eta H_u(y) \Big|_{y=x} = -M_t^\eta H_t(x) \Big|_{t=u}, \quad \forall x \neq u.$$

The first example is the family of tent maps with fixed slopes $\lambda_0 \in (1, 2)$ given by $\tilde{f}_t(x) = \lambda_0 x + \lambda_0 - 1 + t_0 + t$ for $-1 \leq x \leq 0$, and $\tilde{f}_t(x) = -\lambda_0 x + \lambda_0 - 1 + t_0 + t$ for $0 \leq x \leq 1$, where $|t| < \varepsilon_1$, for small enough $\varepsilon_1 \in (0, \min\{|t_0|, 2 - \lambda_0 - t_0\})$. (For $|t| > \varepsilon_1$ we set $\tilde{f}_t(x) = \tilde{f}_{\pm\varepsilon_1}(x)$.) This family is horizontal because each \tilde{f}_t is topologically conjugated to \tilde{f}_0 . (Indeed, setting, $h_t(x) = (\lambda_0 - 1 + t_0 + t)x$, we have $h_t \circ \tilde{f}_0(x) = \tilde{f}_t \circ h_t(x)$. We can also use that the topological entropy of \tilde{f}_t is $\log \lambda_0$ for all t .)

We assume that the orbit of $c = 0$ is infinite for the sake of simplicity. We have $\tilde{X}_0 \equiv 1$ on the support of $\tilde{\rho}_0$, so that (22) and (23) imply

$$M_x^\eta(\tilde{X}_0\tilde{\rho}_0)(x) = \sum_{n=1}^{\infty} \bar{s}_n M_x^\eta H_{c_n}(x) = \frac{1}{2\Gamma(1-\eta)} \sum_{n=1}^{\infty} \frac{s_1}{(\tilde{f}_0^{n-1})'(c_1)} \frac{1}{|x - c_n|^\eta},$$

where $(\tilde{f}_0^{n-1})'(c_1) = \sigma_n \lambda_0^{n-1}$, for $\sigma_n = \text{sgn}(\tilde{f}_0^{n-1})'(c_1)$. It follows that

$$\Psi_\phi^{\text{rsp}}(\eta, z) = -\frac{1}{2\Gamma(1-\eta)} \sum_{k=0}^{\infty} z^k \int_I (\phi \circ \tilde{f}_0^k) \sum_{n=1}^{\infty} \frac{s_1 \sigma_n}{\lambda_0^{n-1}} \frac{1}{|x - c_n|^\eta} dx.$$

Next, observe that since $\tilde{f}_t(y) = \tilde{f}_0(y) + t$ for $|t| < \varepsilon_1$, we have $(\tilde{\mathcal{L}}_t \varphi)(x) = (\tilde{\mathcal{L}}_0 \varphi)(x - t)$ for such t , so that, using $\tilde{\mathcal{L}}_0 \rho_0^{\text{sal}} = \rho_0^{\text{sal}}$, we find, for any $|t| < \varepsilon_1$,

$$(24) \quad (\tilde{\mathcal{L}}_t \tilde{\rho}_0)(x) = \sum_{n=1}^{\infty} \frac{s_1}{(\tilde{f}_0^{n-1})'(c_1)} H_{c_n}(x - t) = \sum_{n=1}^{\infty} \frac{s_1 \sigma_n}{\lambda_0^{n-1}} H_t(x - c_n).$$

Set $H_t^{(\varepsilon_1)}(y) = H_t(y)$ if $|t| < \varepsilon_1$, and $H_t^{(\varepsilon_1)}(y) = H_{\pm \varepsilon_1}(y)$ if $\pm t > \varepsilon_1$. Then, we have,

$$(25) \quad M_t^\eta H_t^{(\varepsilon_1)}(y)|_{t=u} = \begin{cases} -\frac{1}{2\Gamma(1-\eta)} \frac{1}{|y-u|^\eta} & \text{if } 0 < |y-u| < \varepsilon_1, \\ 0 & \text{if } |y-u| \geq \varepsilon_1. \end{cases}$$

(The claims above are proved in the same way as (22).) Using (24) and (25), we get,

$$M_t^\eta (\tilde{\mathcal{L}}_t \tilde{\rho}_0)(x)|_{t=0} = -\frac{1}{2\Gamma(1-\eta)} \sum_{n=1}^{\infty} \frac{s_1 \sigma_n}{\lambda_0^{n-1}} \left(\frac{1_{[c_n-\varepsilon_1, c_n+\varepsilon_1]}(x)}{|x - c_n|^\eta} \right).$$

Finally,

$$\Psi_\phi^{\text{fr}}(\eta, z) = -\frac{1}{2\Gamma(1-\eta)} \sum_{k=0}^{\infty} z^k \int_I (\phi \circ \tilde{f}_0^k) \sum_{n=1}^{\infty} \frac{s_1 \sigma_n}{\lambda_0^{n-1}} \frac{1_{[c_n-\varepsilon_1, c_n+\varepsilon_1]}(x)}{|x - c_n|^\eta} dx.$$

So we see that, in the horizontal linear case given by the family (\tilde{f}_t) , the frozen and response susceptibility function coincide if $\varepsilon_1 \geq \sup_n(c_1 - c_n, c_n - c_2)$. This does not seem possible, since $\varepsilon_1 \leq \varepsilon_0$, where ε_0 is given by the uniform Lasota–Yorke bound Lemma 3.1. However, the two susceptibility functions are qualitatively similar. (In fact, the relation $\tilde{f}_t = \tilde{f}_0 + t$ for $|t| < \varepsilon_1$ implies that the response and frozen susceptibilities differ by a function holomorphic in a disc of radius larger than one; this can be shown as in [11, Proposition 2.5]). In addition, if we replaced¹⁰ in the definitions of all fractional susceptibility functions the Marchaud derivative by the truncated Marchaud derivative

$$(26) \quad M^{\eta, (\varepsilon_1)} g(t_0) := \frac{\Gamma_\eta}{2} \int_{-\varepsilon_1}^{\varepsilon_1} \frac{g(t_0 + t) - g(t_0)}{|t|^{1+\eta}} \text{sign}(t) dt,$$

then the frozen and response susceptibility functions would coincide for \tilde{f}_t . (Lemma 2.2 and the integration by parts formula mentioned in footnote 5 both hold for $M^{\eta, (\varepsilon_1)}$.)

The second example is the family of tent maps with varying slopes $\bar{f}_t(x) = \lambda_t x + \lambda_t - 1$ for $-1 \leq x \leq 0$ and $\bar{f}_t(x) = -\lambda_t x + \lambda_t - 1$ for $0 \leq x \leq 1$, where $\lambda_t = \lambda_0 + t$, for $\lambda_0 \in (1, 2)$ and $|t| < \varepsilon_1$, with small enough $\varepsilon_1 < \min(2 - \lambda_0, \lambda_0 - 1)$. This family is not horizontal

¹⁰With this modification, the use of $f_t^{(\varepsilon_1)}$ as defined in (11) would not be needed.

(see e.g. [41, §4]). We assume again that the orbit of $c = 0$ for \bar{f}_0 is infinite for the sake of simplicity.

Setting $\bar{v}_t = \partial_s \bar{f}_s|_{s=t}$, and defining \bar{X}_t by $\bar{v}_t = \bar{X}_t \circ \bar{f}_t$, we have $\bar{v}_t(x) = x + 1$ if $-1 < x < 0$ and $\bar{v}_t(x) = -x + 1$ if $0 < x < 1$, so that $\bar{X}_t(y) = (y + 1)/\lambda_t$ and $\bar{X}_0(y) = (y + 1)/\lambda_0$, for $y \in [-1, 1]$. We find,

$$(27) \quad M_x^\eta(\bar{X}_0 \bar{\rho}_0)(x) = M_x^\eta \left(\sum_{n=1}^{\infty} \frac{s_1 \sigma_n}{\lambda_0^{n-1}} \frac{x+1}{\lambda_0} H_{c_n}(x) \right).$$

(The computation of the Marchaud derivative above is straightforward, but cumbersome, using that the sum over n is supported in $[c_2, c_1]$, and we do not carry it out here. We just point out that, if we use $M^{\eta, (\varepsilon_1)}$ from (26) instead of M^η , we have

$$\begin{aligned} M^{\eta, (\varepsilon_1)}(x H_{c_n}(x)) &= \frac{\Gamma_\eta}{2} \int_{-\varepsilon_1}^{\varepsilon_1} \text{sign}(y) \frac{(x+y) \cdot H_{c_n}(x+y) - x \cdot H_{c_n}(x)}{|y|^{1+\eta}} dy \\ &= \frac{\Gamma_\eta}{2} \int_{-\varepsilon_1}^{\varepsilon_1} \frac{H_{c_n-x}(y)}{|y|^\eta} dy + x \cdot M^{\eta, (\varepsilon_1)}(H_{c_n}(x)), \end{aligned}$$

so that the singularity type of $M^{\eta, (\varepsilon_1)}((x+1)H_{c_n}(x))$ is $|x - c_n|^{-\eta}$.)

We move to the frozen susceptibility function. We have $\bar{f}_t(x) = (\lambda_t/\lambda_0)(\bar{f}_0(x)+1) - 1$, so that

$$(28) \quad \bar{\mathcal{L}}_t \varphi(x) = \frac{\lambda_0}{\lambda_t} \cdot \bar{\mathcal{L}}_0 \varphi \left(\frac{\lambda_0 x - t}{\lambda_t} \right),$$

for all $|t| < \varepsilon_1$. Combining (28) with $\bar{\mathcal{L}}_0 \bar{\rho}_0^{\text{sal}} = \bar{\rho}_0^{\text{sal}}$, we find

$$\begin{aligned} \bar{\mathcal{L}}_t \bar{\rho}_0^{\text{sal}}(x) &= \frac{\lambda_0}{\lambda_t} \bar{\rho}_0^{\text{sal}} \left(\frac{\lambda_0 x - t}{\lambda_t} \right) = \frac{\lambda_0}{\lambda_0 + t} \sum_{n=1}^{\infty} \frac{s_1 \sigma_n}{\lambda_0^{n-1}} H_{c_n} \left(\frac{\lambda_0 x - t}{\lambda_0 + t} \right) \\ &= \frac{1}{1 + t/\lambda_0} \sum_{n=1}^{\infty} \frac{s_1 \sigma_n}{\lambda_0^{n-1}} H_{c_n + (c_n+1)t/\lambda_0}(x) \\ &= \left(\sum_{k=0}^{\infty} \left(-\frac{t}{\lambda_0} \right)^k \right) \cdot \sum_{n=1}^{\infty} \frac{s_1 \sigma_n}{\lambda_0^{n-1}} H_t \left(\lambda_0 \frac{x - c_n}{c_n + 1} \right), \end{aligned}$$

for all $|t| < \varepsilon_1$, so that

$$(29) \quad M_t^\eta(\bar{\mathcal{L}}_t(\bar{\rho}_0)(x))|_{t=0} = \sum_{k=0}^{\infty} \frac{(-1)^k}{\lambda_0^k} M_t^\eta \left(t^k \cdot \sum_{n=1}^{\infty} \frac{s_1 \sigma_n}{\lambda_0^{n-1}} H_t \left(\lambda_0 \frac{x - c_n}{c_n + 1} \right) \right) \Big|_{t=0}.$$

Comparing (27) with (29), we see that the discrepancy between $M_x^\eta(\bar{X}_0 \bar{\rho}_0)(x)$ and $-M_t^\eta(\bar{\mathcal{L}}_t(\bar{\rho}_0)(x))|_{t=0}$, and thus between $\Psi_\phi^{\text{fr}}(\eta, z)$ and $\Psi_\phi^{\text{rsp}}(\eta, z)$, is more marked for the family \bar{f}_t than for the family \tilde{f}_t .

3. TRANSFER OPERATORS AND SOBOLEV SPACES

3.1. Basic definition and properties. We will use the Sobolev spaces $\mathcal{H}^{\tau, p} = \mathcal{H}^{\tau, p}(I)$ of functions $\varphi \in L^p(\mathbb{R})$ supported in I such that

$$\|\varphi\|_{\mathcal{H}^{\tau, p}} = \|\mathcal{F}^{-1}((1 + |\xi|^2)^{\tau/2}(\mathcal{F}\varphi))\|_{L^p} < \infty,$$

where $p > 1$ and $\tau \in [0, 1/p)$ are real numbers, and \mathcal{F} denotes the Fourier transform. (Note that $\mathcal{H}^{\tau,p}(I) \subset \mathcal{H}^{\tilde{\tau},p}(I)$ if $\tilde{\tau} < \tau$, and that the embedding is compact.)

Let f_t be a piecewise C^2 expanding unimodal map. Thomine [38] showed that the transfer operator \mathcal{L}_t is bounded on $\mathcal{H}^{\tau,p}$, for any $1 < p < \infty$ and $0 \leq \tau < \frac{1}{p}$. More precisely, there exists a constant $C = C(\sup |f_t''|, \inf |f_t'|)$ such that

$$(30) \quad \|\mathcal{L}_t \varphi\|_{\mathcal{H}^{\tau,p}} \leq C \|\varphi\|_{\mathcal{H}^{\tau,p}}, \quad \forall \varphi, \forall 0 \leq \tau < 1/p.$$

Recall that the essential spectral radius $r_{\text{ess}}(\mathcal{L}|_{\mathcal{B}})$ of a bounded operator $\mathcal{L}: \mathcal{B} \rightarrow \mathcal{B}$ on a Banach space \mathcal{B} is the smallest $r \geq 0$ such that the spectrum of \mathcal{L} on \mathcal{B} in the complement of the disc of radius r consists of isolated eigenvalues of finite multiplicity. Thomine [38, Theorem 1.3] proved that, for any $1 < p < \infty$ and $0 < \tau < \frac{1}{p}$, we have¹¹

$$\begin{aligned} r_{\text{ess}}(\mathcal{L}_t|_{\mathcal{H}^{\tau,p}(I)}) &\leq \lim_{n \rightarrow \infty} \left\| 2^{\frac{1}{p}} \cdot 2^{n(1-\frac{1}{p})} |(f_t^n)'|^{-1+\frac{1}{p}} (\lambda_n(f_t))^{-\tau} \right\|_{\infty}^{\frac{1}{n}} \\ &\leq \lim_{n \rightarrow \infty} 2^{1-\frac{1}{p}} (\lambda_n(f_t))^{(-\tau-1+\frac{1}{p})/n}. \end{aligned}$$

In particular, if $\tilde{\Lambda} \in (1, \Lambda(f_t))$, we find $p(f_t) = p(f_t, \tilde{\Lambda}) > 1$ such that

$$(31) \quad r_{\text{ess}}(\mathcal{L}_t|_{\mathcal{H}^{\tau,p}}) < \tilde{\Lambda}^{-\tau-1+\frac{1}{p}} < 1, \quad \forall 1 < p < p(f), \forall 0 < \tau < \frac{1}{p}.$$

By standard arguments (see [38, Theorem 1.6]), using that f_t is unimodal and thus topologically transitive on $[f_t^2(c), f_t(c)]$, and that the dual of \mathcal{L}_t preserves Lebesgue measure dx on I , it follows that for such τ and p the spectral radius of \mathcal{L}_t on $\mathcal{H}^{\tau,p}$ is equal to one, that the invariant density ρ_t belongs to $\mathcal{H}^{\tau,p}$, (extending ρ_t by zero outside of its domain), that ρ_t is the unique fixed point of \mathcal{L}_t in $\mathcal{H}^{\tau,p}$, and that the algebraic multiplicity of the eigenvalue 1 is equal to one.

If $0 < \tilde{\tau} < \tau$ then, outside of the closed disc of radius $\tilde{\Lambda}^{-\tilde{\tau}-1+\frac{1}{p}}$, the spectrum of \mathcal{L}_t on $\mathcal{H}^{\tilde{\tau},p}(I)$ and $\mathcal{H}^{\tau,p}(I)$ coincide, including multiplicity and generalised eigenspaces (this follows from the fact that both spaces are continuously embedded in L^p while C^1 functions are dense in both spaces [34, 2.1.3, Proposition 1], applying the result in [12, App. A], see also [5, App. A.2]).

3.2. Stability of mixing and mixing rates for good families. To prove stability of the spectrum of the perturbation (f_t) , the following uniform Lasota–Yorke lemma is crucial. (A version of Lemma 3.1 is established for the BV and L^1 norm, see¹² e.g. [25, comment above Remark 2]. The proof for our Sobolev spaces is given in Appendix B.)

Lemma 3.1 (Uniform Lasota–Yorke bound). *Let $(f_t)_{|t| \leq \varepsilon}$ be a C^2 perturbation of a piecewise C^2 expanding unimodal map $f = f_0$. Assume that either f is good or all the f_t are topologically conjugated to f , and recall $\Lambda > 1$ from Definition 2.1. Then for*

¹¹Indeed, in our one-dimensional unimodal setting Thomine’s “ n -complexity at the beginning” is bounded by 2, and his “ n -complexity at the end” is bounded by 2^n . Cf. the proof of our uniform Lasota–Yorke estimate Lemma 3.1.

¹²Above equation (26) in [9] there is a mistaken reference to Remark 5 in [25] instead.

any $\tilde{\Lambda} < \Lambda$ there exist $p_0 \in (1, p(f))$ and $\varepsilon_0 \leq \varepsilon$ such that for any $p \in (1, p_0)$ and $0 \leq \tilde{\tau} < \tau < \frac{1}{p}$ there exist finite constants C_0 and C such that

$$(32) \quad \|\mathcal{L}_t^n \varphi\|_{\mathcal{H}^{\tau,p}} \leq C \tilde{\Lambda}^{(-\tau-1+\frac{1}{p})n} \|\varphi\|_{\mathcal{H}^{\tau,p}} + C C_0^n \|\varphi\|_{\mathcal{H}^{\tilde{\tau},p}}, \quad \forall |t| < \varepsilon_0, \forall n \geq 0, \forall \varphi.$$

We shall also use the following perturbation estimate, proved in §3.3:

Lemma 3.2 (Perturbation bound). *Let $(f_t)_{|t| \leq \varepsilon}$ be a C^2 perturbation of a piecewise C^2 expanding unimodal map f_0 . For any $p > 1$ and $0 < \tilde{\tau} < \frac{1}{p}$ there exists $C < \infty$ such that*

$$(33) \quad \|(\mathcal{L}_t - \mathcal{L}_0)\varphi\|_{\mathcal{H}^{\tilde{\tau},p}} \leq C |t|^{\tau-\tilde{\tau}} \|\varphi\|_{\mathcal{H}^{\tau,p}}, \quad \forall |t| < \varepsilon, \forall \tau \in (\tilde{\tau}, \frac{1}{p}), \forall \varphi.$$

Assume now that $f_t = f$ is mixing. Then for any $\mathcal{H}^{\tau,p}$ with $p \in (p(f), 1)$ and $\tau < 1/p$, we have that 1 is the only eigenvalue of the transfer operator \mathcal{L}_0 of f on the unit circle (adapting e.g. the proof [2, Theorem 3.5] for \mathcal{L}_t acting on BV). In other words, the operator \mathcal{L}_0 has a spectral gap on $\mathcal{H}^{\tau,p}$. The following notation will be useful:

Definition 3.3 (Maximal eigenvalue $\kappa < 1$ of a mixing piecewise C^2 expanding unimodal map f). *Let f be a mixing piecewise C^2 expanding unimodal map. If there exist $p > 1$ and $\tau < 1/p$ such that \mathcal{L}_0 on $\mathcal{H}^{\tau,p}$ has an eigenvalue $\zeta \neq 1$ with $|\zeta| > \Lambda(f)^{-\tau-1+\frac{1}{p}}$, then we set $\kappa(f) < 1$ to be the maximal such modulus $|\zeta|$. Otherwise we set $\kappa(f) = 0$.*

Recalling the notation from (31), fix $\tau < 1/p$ for $p < p(f, \tilde{\Lambda})$. Then for any $\kappa_0 > \max(\kappa(f), \tilde{\Lambda}^{-\tau-1+\frac{1}{p}})$ there exists C such that

$$(34) \quad \|\mathcal{L}_0^n \varphi\|_{\mathcal{H}^{\tau,p}} \leq C \kappa_0^n \|\varphi\|_{\mathcal{H}^{\tau,p}}, \quad \forall n \geq 0, \forall \varphi \in \mathcal{H}_0^{\tau,p} := \{\varphi \in \mathcal{H}^{\tau,p}(I) \mid \int \varphi dx = 0\}.$$

Lemmas 3.1 and 3.2 put us in a position to apply the results of Keller–Liverani and show stability of (34): Since $\mathcal{H}_0^{\tau,p}$ is invariant under each \mathcal{L}_t , [25, Corollary 2 (2)] gives for any $\kappa_1 > \max(\kappa(f), \Lambda^{-\tau-1+\frac{1}{p}})$ constants¹³ $C = C(\tau, p) < \infty$, $\varepsilon_1 \in (0, \varepsilon_0)$ such that

$$(35) \quad \|\mathcal{L}_t^n \varphi\|_{\mathcal{H}^{\tau,p}} \leq C \kappa_1^n \|\varphi\|_{\mathcal{H}^{\tau,p}}, \quad \forall n \geq 1, \forall |t| \leq \varepsilon_1, \forall \varphi \in \mathcal{H}_0^{\tau,p}.$$

The uniform mixing rate given by the above bound will be crucial to establish our main theorem.

3.3. Four basic lemmas and the proof of the perturbation Lemma 3.2. We end this section by showing the perturbation Lemma 3.2. The proof will use the following four standard lemmas (the first three lemmas are also instrumental in the proof of Lemma 3.1):

Lemma 3.4 (Triebel [40, Section 4.2.2]). *Suppose that $g \in C^\gamma$ where $\gamma > \tau$, and let $p > 1$. Then there exists $C = C(\tau, p, \gamma) > 0$ such that $\|g\varphi\|_{\mathcal{H}^{\tau,p}} \leq C \|g\|_{C^\gamma} \|\varphi\|_{\mathcal{H}^{\tau,p}}$.*

Lemma 3.5 (See e.g. [38, Lemma 3.3]). *Let T be a C^1 diffeomorphism of \mathbb{R} such that $\frac{A}{2} \leq |T'(x)| \leq 2A$ for some $A > 0$ and all $x \in \mathbb{R}$. Then, for all $0 \leq \tau \leq 1$ and $p > 1$ there exists $C = C(\tau, p) > 0$ such that $\|\varphi \circ T\|_{\mathcal{H}^{\tau,p}} \leq C A^{\tau-\frac{1}{p}} \|\varphi\|_{\mathcal{H}^{\tau,p}} + C A^{-\frac{1}{p}} \|\varphi\|_{L^p}$.*

¹³ $C(\tau, p)$ tends to infinity as η tends to 1.

Lemma 3.6 (Strichartz [37, Corollary I.4.2]). *Suppose that $0 \leq \tau < \frac{1}{p} < 1$. Then there is a constant $C = C(\tau, p)$ such that $\|1_J \varphi\|_{\mathcal{H}^{\tau, p}} \leq C \|\varphi\|_{\mathcal{H}^{\tau, p}}$ for any interval J .*

Lemma 3.7 (See e.g. [5, Lemma 2.39]). *For any $p > 1$ and any C^1 map $T: \mathbb{R} \rightarrow \mathbb{R}$ there exists $C = C(p, \|T\|_{C^1})$ such that $\|\varphi - \varphi \circ T\|_{\mathcal{H}^{\tilde{\tau}, p}} \leq C \|I - T\|_{C^1}^{\tau - \tilde{\tau}} \|\varphi\|_{\mathcal{H}^{\tau, p}}$ for all $0 < \tilde{\tau} < \tau < 1$ and all φ .*

Proof of Lemma 3.2. Let $I_t = f_t(I)$, and put $J_t = I_t \setminus (I_0 \cap I_t)$ and $K_t = I_0 \setminus (I_0 \cap I_t)$. (It is possible that J_t or K_t is empty.) We have

$$\mathcal{L}_t \varphi - \mathcal{L}_0 \varphi = 1_{I_0 \cap I_t} (\mathcal{L}_t \varphi - \mathcal{L}_0 \varphi) + 1_{J_t} \mathcal{L}_t \varphi - 1_{K_t} \mathcal{L}_0 \varphi,$$

since $\mathcal{L}_t \varphi$ is supported in $I_t = J_t \cup (I_0 \cap I_t)$ and $\mathcal{L}_0 \varphi$ is supported in $I_0 = K_t \cup (I_0 \cap I_t)$. Hence

$$(36) \quad \|\mathcal{L}_t \varphi - \mathcal{L}_0 \varphi\|_{\mathcal{H}^{\tilde{\tau}, p}} \leq \|1_{I_0 \cap I_t} (\mathcal{L}_t \varphi - \mathcal{L}_0 \varphi)\|_{\mathcal{H}^{\tilde{\tau}, p}} + \|1_{J_t} \mathcal{L}_t \varphi\|_{\mathcal{H}^{\tilde{\tau}, p}} + \|1_{K_t} \mathcal{L}_0 \varphi\|_{\mathcal{H}^{\tilde{\tau}, p}}.$$

We consider first the term $\|1_{I_0 \cap I_t} (\mathcal{L}_t \varphi - \mathcal{L}_0 \varphi)\|_{\mathcal{H}^{\tilde{\tau}, p}}$. If $x \in I_0 \cap I_t$, both $f_{t, \pm}^{-1}(x)$ and $f_{0, \pm}^{-1}(x)$ are defined, and we may write

$$\begin{aligned} & \|1_{I_0 \cap I_t} (\mathcal{L}_t \varphi - \mathcal{L}_0 \varphi)\|_{\mathcal{H}^{\tilde{\tau}, p}} \\ & \leq \|1_{I_0 \cap I_t} \left(\frac{\varphi}{|f'_t|} \circ f_{t, -}^{-1} - \frac{\varphi}{|f'_0|} \circ f_{0, -}^{-1} \right)\|_{\mathcal{H}^{\tilde{\tau}, p}} + \|1_{I_0 \cap I_t} \left(\frac{\varphi}{|f'_t|} \circ f_{t, +}^{-1} - \frac{\varphi}{|f'_0|} \circ f_{0, +}^{-1} \right)\|_{\mathcal{H}^{\tilde{\tau}, p}}. \end{aligned}$$

We consider the first term on the right-hand side (the second one is treated in the same fashion). Since (f_t) is a C^2 perturbation, we may extend $f_{t, -}: [-1, c] \rightarrow I_t$ to a C^2 -diffeomorphism of \mathbb{R} , still denoted $f_{t, -}: \mathbb{R} \rightarrow \mathbb{R}$, such that $\inf_{|t| \leq \varepsilon} |f'_{t, -}| > 1$, while $f''_{t, -}$ has compact support and is bounded uniformly in t , and $\|f_{t, -} - f_{0, -}\|_{C^1} = O(|t|)$ as $|t| \rightarrow 0$. Since $0 < \tilde{\tau} < 1/p$, by the Strichartz Lemma 3.6,

$$\|1_{I_0 \cap I_t} \left(\frac{\varphi}{|f'_{t, -}|} \circ f_{t, -}^{-1} - \frac{\varphi}{|f'_{0, -}|} \circ f_{0, -}^{-1} \right)\|_{\mathcal{H}^{\tilde{\tau}, p}} \leq C \left\| \frac{\varphi}{|f'_{t, -}|} \circ f_{t, -}^{-1} - \frac{\varphi}{|f'_{0, -}|} \circ f_{0, -}^{-1} \right\|_{\mathcal{H}^{\tilde{\tau}, p}},$$

where φ is extended by zero outside of its domain I . We then split

$$(37) \quad \begin{aligned} \left\| \frac{\varphi}{|f'_{t, -}|} \circ f_{t, -}^{-1} - \frac{\varphi}{|f'_{0, -}|} \circ f_{0, -}^{-1} \right\|_{\mathcal{H}^{\tilde{\tau}, p}} & \leq \left\| \frac{\varphi}{|f'_{t, -}|} \circ f_{t, -}^{-1} - \frac{\varphi}{|f'_{t, -}|} \circ f_{0, -}^{-1} \right\|_{\mathcal{H}^{\tilde{\tau}, p}} \\ & \quad + \left\| \left(\frac{1}{|f'_{t, -}|} \circ f_{0, -}^{-1} - \frac{1}{|f'_{0, -}|} \circ f_{0, -}^{-1} \right) \cdot (\varphi \circ f_{0, -}^{-1}) \right\|_{\mathcal{H}^{\tilde{\tau}, p}}. \end{aligned}$$

The first term in (37) is estimated using Lemmas 3.4, 3.5, and 3.7:

$$\begin{aligned} & \left\| \frac{\varphi}{|f'_{t, -}|} \circ f_{t, -}^{-1} - \frac{\varphi}{|f'_{t, -}|} \circ f_{0, -}^{-1} \right\|_{\mathcal{H}^{\tilde{\tau}, p}} = \left\| \left(\frac{\varphi}{|f'_{t, -}|} \circ f_{t, -}^{-1} \circ f_{0, -} - \frac{\varphi}{|f'_{t, -}|} \right) \circ f_{0, -}^{-1} \right\|_{\mathcal{H}^{\tilde{\tau}, p}} \\ & \leq C \left\| \frac{\varphi}{|f'_{t, -}|} \circ f_{t, -}^{-1} \circ f_{0, -} - \frac{\varphi}{|f'_{t, -}|} \right\|_{\mathcal{H}^{\tilde{\tau}, p}} \leq C \|I - f_{t, -}^{-1} \circ f_{0, -}\|_{C^1}^{\tau - \tilde{\tau}} \left\| \frac{\varphi}{|f'_{t, -}|} \right\|_{\mathcal{H}^{\tau, p}} \\ & \leq C \|I - f_{t, -}^{-1} \circ f_{0, -}\|_{C^1}^{\tau - \tilde{\tau}} \|\varphi\|_{\mathcal{H}^{\tau, p}}, \end{aligned}$$

if $0 < \tilde{\tau} < \tau < 1$. It is easy to check that $\|f_{t, -} - f_{0, -}\|_{C^1} = O(|t|)$ implies that $\|I - f_{t, -}^{-1} \circ f_{0, -}\|_{C^1} = O(|t|)$.

Note that $\|\frac{1}{|f'_{t,-}|} \circ f_{0,-}^{-1} - \frac{1}{|f'_{0,-}|} \circ f_{0,-}^{-1}\|_{C^1} = O(|t|)$ as $t \rightarrow 0$. Since $0 \leq \tilde{\tau} < \tau < 1$, for the second term in (37) we then have by Lemmas 3.4 and 3.5 the upper bound

$$\left\| \left(\frac{1}{|f'_{t,-}|} \circ f_{0,-}^{-1} - \frac{1}{|f'_{0,-}|} \circ f_{0,-}^{-1} \right) \cdot (\varphi \circ f_{0,-}^{-1}) \right\|_{\mathcal{H}^{\tilde{\tau},p}} \leq C|t| \|\varphi\|_{\mathcal{H}^{\tilde{\tau},p}} \leq C|t| \|\varphi\|_{\mathcal{H}^{\tau,p}}.$$

We have shown that if $0 < \tilde{\tau} < \tau < \frac{1}{p}$ then

$$\|1_{I_0 \cap I_t}(\mathcal{L}_t \varphi - \mathcal{L}_0 \varphi)\|_{\mathcal{H}^{\tilde{\tau},p}} \leq C|t|^{\tau - \tilde{\tau}} \|\varphi\|_{\mathcal{H}^{\tau,p}}.$$

We return to (36) and consider $\|1_{J_t} \mathcal{L}_t \varphi\|_{\mathcal{H}^{\tilde{\tau},p}}$.

If $q \in (p, \infty)$, letting r be the conjugate of q/p , then Hölder's inequality gives for any interval $J \subset I$ and any $u \in L^q(I)$ that

$$\|1_J u\|_{L^p(I)} = \|1_J u^p\|_{L^1(I)}^{\frac{1}{p}} \leq (\|1_J\|_{L^r(I)} \|u^p\|_{L^{q/p}(I)})^{\frac{1}{p}} = |J|^{\frac{1}{r p}} \|u\|_{L^q(I)} = |J|^{\frac{1}{p} - \frac{1}{q}} \|u\|_{L^q(I)}.$$

Since $|J_t| \leq C|t|$, it follows by the Sobolev embedding theorem that, taking $q > p$ such that $\tau = \frac{1}{p} - \frac{1}{q}$,

$$(38) \quad \|1_{J_t} \mathcal{L}_t \varphi\|_{L^p} \leq |t|^{\frac{1}{p} - \frac{1}{q}} \|\mathcal{L}_t \varphi\|_{L^q} \leq C|t|^\tau \|\mathcal{L}_t \varphi\|_{\mathcal{H}^{\tau,p}} \leq C|t|^\tau \|\varphi\|_{\mathcal{H}^{\tau,p}}.$$

Also, (30) and Lemma 3.6 give, for $\tau < 1/p$,

$$(39) \quad \|1_{J_t} \mathcal{L}_t \varphi\|_{\mathcal{H}^{\tau,p}} \leq C \|\varphi\|_{\mathcal{H}^{\tau,p}}.$$

Finally, since $\mathcal{H}^{\tilde{\tau},p} = [L^p, \mathcal{H}^{\tau,p}]_{\tilde{\tau}/\tau}$ (where $[\mathcal{B}_0, \mathcal{B}_1]_\theta$ denotes complex interpolation) interpolating at $\theta = \tilde{\tau}/\tau$ (see [34, §2.5.2] and [39, §1.9]) between (38) and (39), we get

$$\|1_{J_t} \mathcal{L}_t \varphi\|_{\mathcal{H}^{\tilde{\tau},p}} \leq C|t|^{\tau(1-\theta)} \|\varphi\|_{\mathcal{H}^{\tau,p}} = C|t|^{\tau - \tilde{\tau}} \|\varphi\|_{\mathcal{H}^{\tau,p}}.$$

In the same way, since $|K_t| \leq C|t|$, we get $\|1_{K_t} \mathcal{L}_0 \varphi\|_{\mathcal{H}^{\tilde{\tau},p}} \leq C|t|^{\tau - \tilde{\tau}} \|\varphi\|_{\mathcal{H}^{\tau,p}}$. Recalling (36), we have proved $\|\mathcal{L}_t \varphi - \mathcal{L}_0 \varphi\|_{\mathcal{H}^{\tilde{\tau},p}} \leq C|t|^{\tau - \tilde{\tau}} \|\varphi\|_{\mathcal{H}^{\tau,p}}$ for $0 < \tilde{\tau} < \tau < \frac{1}{p}$. \square

4. PROOFS OF THEOREMS 2.3, 2.6 AND COROLLARIES 2.4, 2.7

4.1. Proof of Theorem 2.3 on the fractional susceptibility function. Clearly,

$$(40) \quad \int_I (\mathcal{L}_t - \mathcal{L}_0) \varphi(x) dx = 0, \quad \forall \varphi \in L^1.$$

Since I is compact, we may assume that $q \neq \infty$ (otherwise replace q by any finite number larger than $(1 - \eta)^{-1}$). It suffices to consider the contribution of the improper integral \int_0^∞ in $\Psi_\phi(\eta, z)$, the computation for the integral $\int_{-\infty}^0$ is exactly the same. We will prove claim (a) for $\Psi_\phi(\eta, z)$, the proof for $\Psi_\phi^{\text{fr}}(\eta, z)$ is obtained by a slight simplification. Let $q' > 1$ be the conjugate of q , that is $\frac{1}{q} + \frac{1}{q'} = 1$. Then, by Hölder's inequality,

$$\int_I \left| \phi \cdot \mathcal{L}_t^k (\mathcal{L}_t - \mathcal{L}_0) \rho_0 \right| dx \leq \|\mathcal{L}_t^k (\mathcal{L}_t - \mathcal{L}_0) \rho_0\|_{L^{q'}} \|\phi\|_{L^q}.$$

Next, fixing $p \in (1, q')$ (we shall need to take p close to 1 soon) and setting

$$(41) \quad \tilde{\tau} := \frac{1}{p} - \frac{1}{q'} = \frac{1}{p} - 1 + \frac{1}{q} \in \left(0, \frac{1}{p} - \eta\right),$$

the Sobolev embedding theorem [20, Theorem 1.3.5], gives a constant $C_{p,q'}$ such that

$$\int_I \left| \phi \cdot \mathcal{L}_t^k (\mathcal{L}_t - \mathcal{L}_0) \rho_0 \right| dx \leq C_{p,q'} \|\mathcal{L}_t^k (\mathcal{L}_t - \mathcal{L}_0) \rho_0\|_{\mathcal{H}^{\tilde{\tau},p}} \|\phi\|_{L^q}.$$

Thus, for each fixed $k \geq 0$, by Lemma 3.2, the improper integral defining the coefficient of z^k in $\Psi(\eta, z)$ or $\Psi^{\text{fr}}(\eta, z)$ is a well-defined complex number.

Assume from now on that $p \in (1, \min(p_0, q'))$ where p_0 is from Lemma 3.1, and let κ be as in Definition 3.3.

Since $\Lambda^{-\tilde{\tau}-1+\frac{1}{p}} = \Lambda^{-\frac{1}{q}}$, using (40) for $\varphi = \rho_0$, we have for any $\kappa_1 > \max(\kappa, \Lambda^{-\frac{1}{q}})$, by Keller and Liverani's consequence (35) of the uniform Lasota–Yorke and perturbation bounds, constants $\varepsilon_1 > 0$ and¹⁴ $C_\eta < \infty$ such that

$$(42) \quad \|\mathcal{L}_t^k (\mathcal{L}_t - \mathcal{L}_0) \rho_0\|_{\mathcal{H}^{\tilde{\tau},p}} \leq C_\eta \kappa_1^k \|(\mathcal{L}_t - \mathcal{L}_0) \rho_0\|_{\mathcal{H}^{\tilde{\tau},p}}, \forall |t| \leq \varepsilon_1.$$

For any $\tau \in (\tilde{\tau}, \frac{1}{p})$, the bound (33) implies

$$(43) \quad \|(\mathcal{L}_t - \mathcal{L}_0) \rho_0\|_{\mathcal{H}^{\tilde{\tau},p}} \leq C |t|^{\tau-\tilde{\tau}} \|\rho_0\|_{\mathcal{H}^{\tau,p}}, \forall |t| \leq \varepsilon_1.$$

We conclude that for any $0 \leq |t| < \varepsilon_1$

$$(44) \quad \int_I \left| \phi \cdot \mathcal{L}_t^k \frac{(\mathcal{L}_t - \mathcal{L}_0) \rho_0}{|t|^{1+\eta}} \right| dx \leq C_\eta \kappa_1^k |t|^{-1-\eta+\tau-\tilde{\tau}} \|\rho_0\|_{\mathcal{H}^{\tau,p}} \|\phi\|_{L^q}.$$

Since $q > (1-\eta)^{-1}$, we may choose $\tau < 1/p$ and $\tilde{\tau} = 1/p - 1 + 1/q$ such that $\eta < \tau - \tilde{\tau}$. For such τ and $\tilde{\tau}$, we may integrate (44) over $t \in (0, \varepsilon_1)$, and we get

$$(45) \quad \int_0^{\varepsilon_1} \int_I \left| \phi(x) \mathcal{L}_t^k \frac{(\mathcal{L}_t - \mathcal{L}_0) \rho_0}{t^{1+\eta}} \right| dx dt \leq C \kappa_1^k \|\rho_0\|_{\mathcal{H}^{\tau,p}} \|\phi\|_{L^q}.$$

For $t > \varepsilon_1$, we have $\mathcal{L}_t = \mathcal{L}_{\pm \varepsilon_1}$, so that, using again the arguments above,

$$(46) \quad \begin{aligned} \int_{t>\varepsilon_1} \int_I \left| \phi \cdot \mathcal{L}_t^k \frac{(\mathcal{L}_t - \mathcal{L}_0) \rho_0}{t^{1+\eta}} \right| dx dt &\leq \int_{t>\varepsilon_1} \frac{1}{t^{1+\eta}} dt \cdot \int_I \left| \phi \cdot \mathcal{L}_{\pm \varepsilon_1}^k (\mathcal{L}_{\varepsilon_1} - \mathcal{L}_0) \rho_0 \right| dx \\ &\leq C_\eta \kappa_1^k \varepsilon_1^{\tau-\tilde{\tau}-\eta} \|\rho_0\|_{\mathcal{H}^{\tau,p}} \|\phi\|_{L^q}. \end{aligned}$$

By Fubini, the bounds (45) and (46) imply that $\Psi_\phi(\eta, z)$ and $\Psi_\phi^{\text{fr}}(\eta, z)$ are holomorphic in the disc of radius κ_1^{-1} , if $\phi \in L^q$ for $q > (1-\eta)^{-1}$. Since we may take q arbitrarily close to $(1-\eta)^{-1}$ and κ_1 arbitrarily close to $\min(\kappa, \Lambda^{-\frac{1}{q}})$, this shows (a).

We next show claim (b) of the theorem. The bounds (40), (42), (45), and (46) together with Fubini and the dominated convergence theorem imply that, as holomorphic functions in the disc of radius $\kappa_1^{-1} > 1$,

$$\begin{aligned} \Psi_\phi(\eta, z) &= \frac{\Gamma_\eta}{2} \sum_{k=0}^{\infty} z^k \int_{-\infty}^{\infty} \int_I \phi \left(\mathcal{L}_t^k \frac{(\mathcal{L}_t - \mathcal{L}_0) \rho_0}{|t|^{1+\eta}} \right) \text{sign}(t) dx dt \\ &= \frac{\Gamma_\eta}{2} \int_{-\infty}^{\infty} \int_I \phi \left((I - z \mathcal{L}_t)^{-1} \frac{(\mathcal{L}_t - \mathcal{L}_0) \rho_0}{|t|^{1+\eta}} \right) \text{sign}(t) dx dt. \end{aligned}$$

¹⁴As $\eta \rightarrow 1$, we have $C_\eta \leq C(1-\eta)^{-1}$.

In view of (40), the trivial identity $(\mathcal{L}_t - \mathcal{L}_0)\rho_0 = (I - \mathcal{L}_t)(\rho_t - \rho_0)$ implies the key formula (1). In particular,

$$\begin{aligned}\Psi_\phi(\eta, 1) &= \frac{\Gamma_\eta}{2} \int_{-\infty}^{\infty} \frac{1}{|t|^{1+\eta}} \int_I \phi(x)(\rho_0(x) - \rho_t(x)) dx \operatorname{sign}(t) dt \\ &= M^\eta \left(\int_I \phi(x)\rho_t(x) dx \right) \Big|_{t=0}.\end{aligned}$$

This proves claim (b).

To conclude the proof of Theorem 2.3, it only remains to prove (c). Note that \mathcal{L}_0^k does not depend on t and acts on functions depending on x . Since we have shown that every improper integral defining $\Psi_\phi^{\text{fr}}(\eta, z)$ is convergent, and since the sum over k in (13) converges absolutely for $|z| \leq 1$, we may write

$$\begin{aligned}\Psi_\phi^{\text{fr}}(\eta, z) &= \sum_{k=0}^{\infty} \int_I \phi z^k \mathcal{L}_0^k \left(\frac{\Gamma_\eta}{2} \int_{-\infty}^{\infty} \left(\frac{(\mathcal{L}_0 - \mathcal{L}_t)\rho_0}{|t|^{1+\eta}} \right) \operatorname{sign}(t) dt \right) dx \\ &= \int_I \phi (I - z\mathcal{L}_0)^{-1} \left(\frac{\Gamma_\eta}{2} \int_{-\infty}^{\infty} \left(\frac{(\mathcal{L}_0 - \mathcal{L}_t)\rho_0}{|t|^{1+\eta}} \right) \operatorname{sign}(t) dt \right) dx.\end{aligned}$$

Thus, (c) just follows from the definition of the Marchaud derivative M^η . \square

4.2. Proof of Theorem 2.6 on the generalized fractional susceptibility function. The proof of Theorem 2.6 is the same as the proof of Theorem 2.3 up to equation (44). Instead of (44) we have

$$\int_I \left| \phi \cdot \mathcal{L}_t^k \frac{(\mathcal{L}_t - \mathcal{L}_0)\rho_0}{\ell(|t|)} \right| dx \leq C\kappa_1^k \frac{|t|^{\tau-\tilde{\tau}}}{\ell(|t|)} \|\rho_0\|_{\mathcal{H}^{\tau,p}} \|\phi\|_{L^q}.$$

We choose $\tau < 1/p$ and $\tilde{\tau} = 1/p - 1 + 1/q$. Then $\tau - \tilde{\tau} = 1 - 1/q \geq \gamma$ and

$$\int_0^{\varepsilon_1} \frac{t^{\tau-\tilde{\tau}}}{\ell(t)} dt \leq \int_0^{\varepsilon_1} \frac{t^\gamma}{\ell(t)} dt < \infty,$$

by the assumption on γ . Hence we get, similar to (45), that

$$\int_0^{\varepsilon_1} \int_I \left| \phi(x) \mathcal{L}_t^k \frac{(\mathcal{L}_t - \mathcal{L}_0)\rho_0}{\ell(t)} \right| dx dt \leq C\kappa_1^k \|\rho_0\|_{\mathcal{H}^{\tau,p}} \|\phi\|_{L^q}.$$

Similar to (46), we have

$$\begin{aligned}\int_{t>\varepsilon_1} \int_I \left| \phi \cdot \mathcal{L}_t^k \frac{(\mathcal{L}_t - \mathcal{L}_0)\rho_0}{\ell(t)} \right| dx dt &\leq \int_{t>\varepsilon_1} \frac{1}{\ell(t)} dt \cdot \int_I \left| \phi \cdot \mathcal{L}_{\varepsilon_1}^k (\mathcal{L}_{\varepsilon_1} - \mathcal{L}_0)\rho_0 \right| dx \\ &\leq C\kappa_1^k \|\rho_0\|_{\mathcal{H}^{\tau,p}} \|\phi\|_{L^q},\end{aligned}$$

where we used the assumption $\int_0^1 1/\ell(t) dt < \infty$. We may choose κ_1 arbitrary close to $\min(\kappa, \Lambda^{-\frac{1}{q}})$, which proves (a).

Claims (b) and (c) are proved just as the corresponding claims in Theorem 2.3, replacing $|t|^{1+\eta}$ by $\ell(t)$ in the proofs.

4.3. Proof of Corollary 2.7 on holomorphic extensions.

Remark 4.1 (Sketch of an alternative proof using Morera's theorem). *The proof below is by estimating the growth of derivatives. We sketch here a more conceptual proof: First extend the definition of M^η and Theorem 2.3 to complex η with $0 < \Re \eta < 1 - q^{-1}$. This implies in particular that the double integrals appearing in $\Psi_\phi(\eta, z)$ are well-defined for such η , and Fubini is justified. Then, since Euler's Gamma function and $t^{-1-\eta}$ are holomorphic in the domain considered, Morera's theorem gives the desired holomorphic extension.*

We first show that the function

$$\eta \mapsto M^\eta \left(\int_I \phi(x) \rho_t(x) dt \right) \Big|_{t=0}$$

is holomorphic in the strip $0 < \Re \eta < 1 - \frac{1}{q}$. We give the argument for the right-sided derivative, the left-sided case is the same up to introducing signs. Let $0 < \gamma < 1 - \frac{1}{q}$ and take η_0 such that $0 < \Re \eta_0 < \gamma$. The right-sided fractional derivative with the parameter $\eta = \eta_0 + \zeta$ is given by

$$\begin{aligned} F_{\eta_0}(\zeta) &:= M_-^{\eta_0+\zeta} \left(\int_I \phi(x) \rho_t(x) dt \right) \Big|_{t=0} = \Psi_\phi^-(\eta_0 + \zeta, 1) \\ &= \sum_{k=0}^{\infty} \frac{\Gamma_{\eta_0+\zeta}}{2} \int_I \phi \int_0^\infty \left(\mathcal{L}_t^k \frac{(\mathcal{L}_t - \mathcal{L}_0) \rho_0}{t^{1+\eta_0+\zeta}} \right) dt dx. \end{aligned}$$

Let $\ell_n(t) = |t|^{1+\eta_0} (\log |t|)^{-n}$. Using Theorem 2.6 and $|t|^{-\zeta} = \sum_{n=0}^{\infty} \frac{1}{n!} (\log |t|)^n \zeta^n$, we then have

$$\begin{aligned} F_{\eta_0}(\zeta) &= \sum_{k=0}^{\infty} \frac{\Gamma_{\eta_0+\zeta}}{2} \int_I \phi \int_0^\infty \left(\mathcal{L}_t^k (\mathcal{L}_t - \mathcal{L}_0) \rho_0 \sum_{n=0}^{\infty} \frac{(\log t)^n \zeta^n}{t^{1+\eta_0} n!} \right) dt dx \\ &= \frac{\Gamma_{\eta_0+\zeta}}{2} \sum_{k=0}^{\infty} \sum_{n=0}^{\infty} \int_I \phi \int_0^\infty \left(\mathcal{L}_t^k (\mathcal{L}_t - \mathcal{L}_0) \rho_0 \frac{(\log t)^n \zeta^n}{t^{1+\eta_0} n!} \right) dt dx \\ &= \frac{\Gamma_{\eta_0+\zeta}}{2} \sum_{n=0}^{\infty} \frac{\Psi_\phi^-((\ell_n), 1)}{n!} \zeta^n. \end{aligned}$$

To justify exchanging the order of sums and integrations above, we will estimate below

$$(47) \quad \frac{\zeta^n}{n!} \int_I \phi \int_{-\infty}^{\infty} \left(\mathcal{L}_t^k (\mathcal{L}_t - \mathcal{L}_0) \rho_0 \frac{(\log |t|)^n}{|t|^{1+\eta_0}} \right) \text{sign}(t) dt dx.$$

Since $F_{\eta_0}(\zeta)$ is a product of $\frac{\Gamma_{\eta_0+\zeta}}{2}$ and $\sum_{n=0}^{\infty} \frac{\Psi_\phi^-((\ell_n), 1)}{n!} \zeta^n$, which are both holomorphic in ζ , provided $|\zeta|$ is sufficiently small, this will show that $F_{\eta_0}(\zeta)$ is holomorphic.

To estimate (47), we first use the proof of Theorem 2.6, to get

$$\left| \frac{\zeta^n}{n!} \int_I \phi \int_{-\infty}^{\infty} \left(\mathcal{L}_t^k (\mathcal{L}_t - \mathcal{L}_0) \rho_0 \frac{(\log |t|)^n}{|t|^{1+\eta_0}} \right) \text{sign}(t) dt dx \right| \leq C_n \kappa_1^k \zeta^n \|\rho_0\|_{\mathcal{H}^{\tau,p}} \|\phi\|_{L^q},$$

where

$$C_n = \frac{C}{n!} \left(\int_0^{\varepsilon_1} \frac{t^\gamma}{|\ell_n(t)|} dt + \int_{t>\varepsilon_1} \frac{1}{|\ell_n(t)|} dt \right),$$

and C is a constant that does not depend on n . Next, since $\gamma > \Re\eta_0$ we have

$$\left| \int_{-\varepsilon_1}^{\varepsilon_1} \frac{|t|^\gamma}{\ell_n(|t|)} dt \right| = 2 \int_0^{\varepsilon_1} \frac{|\log t|^n}{t^{1-(\gamma-\Re\eta_0)}} dt \leq 2 \int_0^1 \frac{|\log t|^n}{t^{1-(\gamma-\Re\eta_0)}} dt = 2 \frac{n!}{(\gamma - \Re\eta_0)^{n+1}}$$

and

$$\begin{aligned} \left| \int_{|t|>\varepsilon_1} \frac{1}{\ell_n(|t|)} dt \right| &\leq 2|\varepsilon_1|^{1+\Re\eta_0} |\log \varepsilon_1|^n + 2 \int_1^\infty \frac{(\log t)^n}{t^{1+\Re\eta_0}} dt \\ &= 2|\varepsilon_1|^{1+\Re\eta_0} |\log \varepsilon_1|^n + 2 \frac{n!}{(\Re\eta_0)^{n+1}}. \end{aligned}$$

In conclusion, C_n grows at most with an exponential speed. Hence, we may change order of integration if ζ is in a sufficiently small disc. In particular, our estimates show that $\frac{\Psi_\phi((\ell_n), 1)}{n!}$ grows at most exponentially in n , so that

$$F_{\eta_0}(\zeta) = \frac{\Gamma_{\eta_0+\zeta}}{2} \sum_{n=0}^{\infty} \frac{\Psi_\phi((\ell_n), 1)}{n!} \zeta^n,$$

holds when ζ is in a disc around the origin.

The proof that the function $(\eta, \zeta) \mapsto \Psi_\phi(\eta, z)$ is holomorphic in the domain

$$\left\{ (\eta, z) \mid 0 < \Re\eta < 1 - \frac{1}{q}, \quad |z| < \min(\kappa, \Lambda^{-\frac{1}{q}}) \right\}$$

uses the same estimates, writing instead

$$\begin{aligned} \Psi_\phi(\eta_0 + \zeta, z) &= \sum_{k=0}^{\infty} \frac{\Gamma_{\eta_0+\zeta}}{2} z^k \int_I \phi \int_{-\infty}^{\infty} \left(\mathcal{L}_t^k \frac{(\mathcal{L}_t - \mathcal{L}_0)\rho_0}{|t|^{1+\eta_0+\zeta}} \right) \text{sign}(t) dt dx \\ &= \sum_{k=0}^{\infty} \frac{\Gamma_{\eta_0+\zeta}}{2} z^k \int_I \phi \int_{-\infty}^{\infty} \left(\mathcal{L}_t^k (\mathcal{L}_t - \mathcal{L}_0)\rho_0 \sum_{n=0}^{\infty} \frac{(\log |t|)^n \zeta^n}{|t|^{1+\eta_0} n!} \right) \text{sign}(t) dt dx. \end{aligned}$$

4.4. Proof of Corollary 2.4 on horizontal perturbations. By [9, Theorem 2.8] (see [10]), the horizontality assumption implies that f_t is tangential to the topological class of f_0 that is, there exist a C^2 perturbation (\tilde{f}_t) of f_0 with $|\tilde{f}_t - f_t| = O(t^2)$ (as $|t| \rightarrow 0$) and homeomorphisms h_t of I with $h_t(c) = c$ and $\tilde{f}_t = h_t \circ f \circ h_t^{-1}$. Note that, letting $\tilde{\mathcal{L}}_t$ be the operator associated to \tilde{f}_t , the proof of Lemma 3.2 gives ε_2 such that (cf. [9, (27) proof of Proposition 3.3]) that

$$\|(\mathcal{L}_t - \tilde{\mathcal{L}}_t)\varphi\|_{\mathcal{H}^{\tilde{\tau}, p}} \leq C|t|^{2(\tau-\tilde{\tau})} \|\varphi\|_{\mathcal{H}^{\tau, p}}, \quad \forall |t| < \varepsilon_2,$$

(this will be used to show (18)). Since we may choose $0 < \tilde{\tau} < \tau < 1/p$ such that $2(\tau-\tilde{\tau}) > 1$, it is not hard to see that we may replace f_t by \tilde{f}_t in the proof of Corollary 2.4.

The two first equalities claimed in Corollary 2.4 follow from statement (b) of Theorem 2.3, combined with Lemma 2.2 and [9, Proposition 4.3, Theorem. 5.1], respectively.

The last statement in the corollary, (18), can be deduced¹⁵ from statement (c) of Theorem 2.3 together with the following claim (the last term in the right-hand side is

¹⁵It would be interesting to have a more direct proof of (18).

understood in the sense of distributions, integrating against a continuous function)

$$(48) \quad \lim_{\eta \uparrow 1} M^\eta(\mathcal{L}_t \rho_0)|_{t=0} = -X'_0 \rho_0 - X_0(\rho_0^{\text{reg}})' - \sum_{k=1}^{\infty} \bar{s}_k X_0(c_k) \delta_{c_k}.$$

Indeed, it suffices to note that (recall that c is not periodic for f)

$$\int \phi \sum_{k=0}^{\infty} \sum_{j=1}^{\infty} \mathcal{L}_0^k(\bar{s}_j X_0(c_j) \delta_{c_j}) = \sum_{k=0}^{\infty} \sum_{j=1}^{\infty} \bar{s}_j X_0(c_j) \phi(c_{j+k}) = \sum_{\ell=0}^{\infty} \phi_{c_\ell} \sum_{j=1}^{\ell} \bar{s}_j X_0(c_j).$$

The desired identity (48) will be an immediate consequence of the decomposition $\rho = \rho^{\text{reg}} + \rho^{\text{sal}}$, with (15), and Lemma 2.2 after we establish that $t \mapsto \int_I \phi(x)(\mathcal{L}_t \rho_0)(x) dx$ is differentiable at $t = 0$ for any $\phi \in C^0$, with derivative

$$(49) \quad \partial_t \left(\int_I \phi(x)(\mathcal{L}_t \rho_0)(x) dx \right) \Big|_{t=0} = - \int_I \phi(X_0 \rho_0)' dx.$$

To show (49), we first recall Step 2 in the proof of [9, Theorem 5.1]. For this, letting $c_{k,t} = f_t^k(c)$ and denoting by \mathcal{B}_t the vector space of functions $\varphi = \varphi^{\text{reg}} + \sum_{k=1}^{\infty} a_k H_{c_{k,t}}$ where $\varphi^{\text{reg}} \in BV \cap C^0$ is supported in I and is differentiable with derivative in BV , and the a_k are complex numbers, we recall the invertible map $G_t: \mathcal{B}_t \rightarrow \mathcal{B}_0$ from [9], defined by

$$G_t(\varphi) = \varphi^{\text{reg}} + \sum_{k=1}^{\infty} a_k H_{c_k}.$$

In this notation, Step 2 in the proof of [9, Theorem 5.1] (the proof uses $f_t = \tilde{f}_t$) says

$$(50) \quad \partial_t(G_t(\mathcal{L}_t(G_t^{-1} \rho_0)))|_{t=0} = -X'_0(\rho_0) - X_0(\rho_0^{\text{reg}})' \in BV.$$

To use the above identity, we decompose $\mathcal{L}_t \rho_0 - \rho_0$ into

$$(51) \quad (G_t(\mathcal{L}_t(G_t^{-1} \rho_0)) - \rho_0) + (\mathcal{L}_t(G_t^{-1}(\rho_0)) - G_t(\mathcal{L}_t(G_t^{-1} \rho_0))) + (\mathcal{L}_t \rho_0 - \mathcal{L}_t(G_t^{-1}(\rho_0))).$$

The first term above is handled by (50). For the other two terms, recall that c is not periodic, so $P_f = \infty$. For the last term in (51), note that, since $c_{k,t} = h_t(c_k)$, we have (see [9, Lemma 2.2, Remark 2.5])

$$(52) \quad \lim_{t \rightarrow 0} \frac{c_{k,t} - c_k}{t} = - \sum_{j=0}^{\infty} \frac{X_0(c_{j+k})}{(f^{j+1})'(c_k)} = \alpha(c_k), \quad \forall k \geq 1.$$

Therefore, since $\mathcal{L}_0(\delta_{c_k}) = \delta_{c_{k+1}}$, using bounded distortion

$$(53) \quad \begin{aligned} \lim_{t \rightarrow 0} \frac{\int \phi(\mathcal{L}_t \rho_0 - \mathcal{L}_t(G_t^{-1}(\rho_0))) dx}{t} &= \lim_{t \rightarrow 0} \frac{\int \phi \mathcal{L}_t(\sum_{k=1}^{\infty} \bar{s}_k H_{c_k} - \sum_{k=1}^{\infty} \bar{s}_k H_{c_{k,t}}) dx}{t} \\ &= \sum_{k=1}^{\infty} \bar{s}_k \alpha(c_k) \phi(c_{k+1}) = \sum_{k=1}^{\infty} \bar{s}_{k+1} f'(c_k) \alpha(c_k) \phi(c_{k+1}). \end{aligned}$$

Finally, for the second term in (51), we have

$$\begin{aligned}
 & (\mathcal{L}_t(G_t^{-1}(\rho_0)) - G_t(\mathcal{L}_t(G_t^{-1}\rho_0))) \\
 &= [\mathcal{L}_t(\rho_0^{\text{reg}} + \sum_{k=1}^{\infty} \bar{s}_k H_{c_{k,t}})]^{\text{sal}} - G_t[\mathcal{L}_t(\rho_0^{\text{reg}} + \sum_{k=1}^{\infty} \bar{s}_k H_{c_{k,t}})]^{\text{sal}} \\
 &= \left(\frac{1}{|f'_{t,-}(c)|} + \frac{1}{|f'_{t,+}(c)|} \right) \rho_0^{\text{reg}}(c)(H_{c_{1,t}} - H_{c_1}) + \sum_{k=1}^{\infty} \frac{\bar{s}_k}{|f'(c_{k,t})|} (H_{c_{k+1,t}} - H_{c_{k+1}}).
 \end{aligned}$$

Therefore, using $\mathcal{L}_0\rho_0(c_1) = \rho_0(c_1) = -\bar{s}_1$ and $\rho_0^{\text{sal}}(c) = 0$,

$$(54) \quad \lim_{t \rightarrow 0} \int \phi \frac{(\mathcal{L}_t(G_t^{-1}(\rho_0)) - G_t(\mathcal{L}_t(G_t^{-1}\rho_0)))}{t} dx$$

$$(55) \quad = -\bar{s}_1 X_0(c_1) \phi(c_1) - \sum_{k=1}^{\infty} \bar{s}_{k+1} \alpha(c_{k+1}) \phi(c_{k+1}).$$

The twisted cohomological equation (19) implies that the sum of (53) and (54) is

$$\begin{aligned}
 & -\bar{s}_1 X_0(c_1) \phi(c_1) + \sum_{k=1}^{\infty} \bar{s}_{k+1} (f'(c_k) \alpha(c_k) - \alpha(f(c_k))) \phi(c_{k+1}) \\
 &= - \sum_{k=0}^{\infty} \bar{s}_{k+1} X_0(c_{k+1}) \phi(c_{k+1}) = - \int \phi (X_0 \rho_0^{\text{sal}})'.
 \end{aligned}$$

This concludes the proof of (49), and of Corollary 2.4. \square

APPENDIX A. PROOF OF LEMMA 2.2

Without loss of generality, we set $t_0 = 0$. Since g is bounded, there exists A such that

$$\left| \int_1^{\infty} \frac{g(0) - g(-t)}{t^{1+\eta}} dt \right| \leq \int_1^{\infty} \frac{2A}{t^{1+\eta}} dt < \infty.$$

Since g is differentiable at 0, there exists B such that

$$\left| \int_0^1 \frac{g(0) - g(-t)}{t^{1+\eta}} dt \right| \leq \int_0^1 \frac{B}{t^\eta} dt < \infty.$$

This shows that $M_+^\eta g(0)$ exists for all $\eta < 1$. Exchanging g with $t \mapsto g(-t)$, we get that $M_-^\eta g(0)$ exists for all $\eta < 1$.

Let $\varepsilon > 0$. We will prove $g'(0) - \varepsilon \leq \liminf_{\eta \uparrow 1} M_+^\eta g(0) \leq \limsup_{\eta \uparrow 1} M_+^\eta g(0) \leq g'(0) + \varepsilon$. Since ε is arbitrary, this implies that $\lim_{\eta \uparrow 1} M_+^\eta g(0) = g'(0)$.

Since the limit $g'(0) = \lim_{h \rightarrow 0} \frac{g(h) - g(0)}{h}$ exists, we can find $\delta > 0$ such that

$$-\varepsilon \leq \frac{g(h) - g(0)}{h} - g'(0) \leq \varepsilon, \quad \forall |h| \leq \delta.$$

Next, since $0 < \eta < 1$, we can write

$$\begin{aligned} M_+^\eta g(0) &= \frac{\eta}{\Gamma(1-\eta)} \left(\int_0^\delta \frac{g(0) - g(-t)}{t} \frac{1}{t^\eta} dt + \int_\delta^\infty \frac{g(0) - g(-t)}{t^{1+\eta}} dt \right) \\ &\leq \frac{\eta}{\Gamma(1-\eta)} \left(\int_0^\delta (g'(0) + \varepsilon) \frac{1}{t^\eta} dt + \int_\delta^\infty \frac{g(0) - g(-t)}{t^{1+\eta}} dt \right) \\ &\leq \frac{\eta}{(1-\eta)\Gamma(1-\eta)} \delta^{1-\eta} (g'(0) + \varepsilon) + \frac{\eta}{\Gamma(1-\eta)} \int_\delta^\infty \frac{2A}{t^{1+\eta}} dt \\ &= \frac{\eta}{\Gamma(2-\eta)} \delta^{1-\eta} (g'(0) + \varepsilon) + \frac{1}{\Gamma(1-\eta)} 2A\delta^{-\eta}. \end{aligned}$$

Using that $\Gamma(1) = 1$, $\lim_{\eta \uparrow 1} \Gamma(1-\eta) = \infty$ and $\lim_{\eta \uparrow 1} \delta^{1-\eta} = 1$, this shows that

$$\limsup_{\eta \uparrow 1} M_+^\eta g(0) \leq g'(0) + \varepsilon.$$

In the same way, we get

$$M_+^\eta g(0) \geq \frac{\eta}{\Gamma(2-\eta)} \delta^{1-\eta} (g'(0) - \varepsilon) - \frac{1}{\Gamma(1-\eta)} 2A\delta^{-\eta},$$

and thus $\liminf_{\eta \uparrow 1} M_+^\eta g(0) \geq g'(0) - \varepsilon$. Using $t \mapsto g(-t)$, we get $\lim_{\eta \uparrow 1} M_-^\eta g(0) = -g'(0)$.

APPENDIX B. PROOF OF THE UNIFORM LASOTA–YORKE ESTIMATE (LEMMA 3.1)

Note that in view of (30) it is enough to show that for any $\tilde{\Lambda} < \Lambda$, there exist $p_0 \in (1, p(f))$ and $\varepsilon_0 \leq \varepsilon$ such that for any $p \in (1, p_0)$ and $0 \leq \tilde{\tau} < \tau < \frac{1}{p}$ there exist $N \geq 1$ and a constant $C(N) < \infty$ such that

$$(56) \quad \|\mathcal{L}_t^N \varphi\|_{\mathcal{H}^{\tau,p}} \leq \tilde{\Lambda}^{(-\tau-1+\frac{1}{p})N} \|\varphi\|_{\mathcal{H}^{\tau,p}} + C(N) \|\varphi\|_{\mathcal{H}^{\tilde{\tau},p}}, \quad \forall |t| < \varepsilon_0, \forall \varphi.$$

(To conclude for general $n = jN + k$, use a geometric series for jN and (30) for $k \leq N-1$.)

Our strategy to prove (56) is to follow Thomine's argument [38], using a ‘‘zoom’’:¹⁶ For this, note that Lemma 3.5 gives $C_{\text{zoom}} = C(\tau, p) > 0$ such that for any positive number $Z > 0$, setting $\mathcal{Z}(x) = Z \cdot x$,

$$(57) \quad \|\varphi \circ \mathcal{Z}\|_{\mathcal{H}^{\tau,p}} \leq C_{\text{zoom}} Z^{\tau-\frac{1}{p}} \|\varphi\|_{\mathcal{H}^{\tau,p}} + C_{\text{zoom}} Z^{-\frac{1}{p}} \|\varphi\|_{L^p}, \forall n.$$

Define $\mathcal{Z}_n: \mathbb{R} \rightarrow \mathbb{R}$ by $\mathcal{Z}_n(x) = Z_n \cdot x$, where¹⁷ the sequence of zooms $Z_n > 1$, for $n \geq 1$, will be specified later. We define an auxiliary zoomed norm by $\|\varphi\|_n = \|\varphi \circ \mathcal{Z}_n^{-1}\|_{\mathcal{H}^{\tau,p}}$, which is equivalent with $\|\cdot\|_{\mathcal{H}^{\tau,p}}$, for each fixed n .

Let $\psi: \mathbb{R} \rightarrow [0, 1]$ be a C^∞ -function supported in $(-1, 1)$ with $\sum_{m \in \mathbb{Z}} \psi_m = 1$, where $\psi_m(x) = \psi(x + m)$. We define

$$\psi_{m,n}(x) = \psi_m \circ \mathcal{Z}_n(x) = \psi_m(Z_n \cdot x), \quad m \in \mathbb{Z}, n \in \mathbb{Z}_+.$$

¹⁶The zoom and localisation can perhaps be replaced by the fragmentation and reconstitution lemmas in [5, Chapter 2].

¹⁷It is essential that Z_n will be chosen uniformly in t for each fixed n .

Note that the support of $\psi_{m,n}$ becomes small if Z_n is large. For any $x \in I$ and any $n \geq 1$, there are at most three integers m such that $\psi_{m,n}(x) \neq 0$, and we may decompose

$$(58) \quad \mathcal{L}_t^n \varphi(x) = \sum_{m \in \mathbb{Z}} \mathcal{L}_t^n(\psi_{m,n} \varphi)(x) = \sum_{m \in \mathbb{Z}} \sum_{J \in \mathcal{A}(t,n)} \left(1_J \psi_{m,n} \frac{\varphi}{|(f_t^n)'|} \right) \circ f_{t,J}^{-n}(x),$$

where $\mathcal{A}(t,n)$ is the partition of monotonicity associated to f_t^n , while $f_{t,J}^{-n}$ is the extension to I of the inverse of $f_t^n|_J$.

For any t , the intersection multiplicity

$$\sup_x \#\{ (m, J) \mid m \in \mathbb{Z}, J \in \mathcal{A}(t,n), x \in f_t^n(J \cap \text{supp } \psi_{m,n}) \}$$

is bounded by $3 \cdot 2^n$ (this is called the ‘‘complexity at the end’’ in Thomine’s paper). Therefore, by Lemma 3.5 in [38] there is $C = C(\tau, p) > 0$ and for each n there exists $C_n = C(\tau, p, \psi, n)$ such that

$$(59) \quad \begin{aligned} \|\mathcal{L}_t^n \varphi\|_n^p &\leq C \sum_{m \in \mathbb{Z}} \sum_{J \in \mathcal{A}(t,n)} 2^{n(p-1)} \left\| \left(1_J \psi_{m,n} \frac{\varphi}{|(f_t^n)'|} \right) \circ f_{t,J}^{-n} \right\|_n^p \\ &\quad + C_n \sum_{m \in \mathbb{Z}} \sum_{J \in \mathcal{A}(t,n)} \left\| \left(1_J \psi_{m,n} \frac{\varphi}{|(f_t^n)'|} \right) \circ f_{t,J}^{-n} \circ \mathcal{Z}_n^{-1} \right\|_{L^p}^p \\ &\leq C \sum_{m \in \mathbb{Z}} \sum_{J \in \mathcal{A}(t,n)} 2^{n(p-1)} \left\| \left(1_J \psi_{m,n} \frac{\varphi}{|(f_t^n)'|} \right) \circ f_{t,J}^{-n} \right\|_n^p + C_n Z_n \|\varphi\|_{L^p}^p. \end{aligned}$$

To control the double sum in (59), we write

$$(1_J \psi_{m,n} \varphi) \circ f_{t,J}^{-n} \circ \mathcal{Z}_n^{-1} = \tilde{\varphi}_{m,n,J} \circ \mathcal{Z}_n \circ f_{t,J}^{-n} \circ \mathcal{Z}_n^{-1},$$

where $\tilde{\varphi}_{m,n,J} = (1_J \psi_{m,n} \varphi) \circ \mathcal{Z}_n^{-1}$. Since $\sup_t |f_t''| < \infty$ and $\inf_t \inf |f_t'| > 1$, a standard bounded distortion¹⁸ estimate gives for any n a large enough Z_n (uniformly in t, m , and J) such that, for each t and $J \in \mathcal{A}(t,n)$, and for any $y \in \text{supp}(1_J \psi_{m,n})$, we have,

$$\frac{1}{2|(f_t^n)'(y)|} \leq \left| \frac{1}{(f_t^n)' \circ f_{t,J}^{-n} \circ \mathcal{Z}_n^{-1}(x)} \right| \leq \frac{3}{2|(f_t^n)'(y)|}, \quad \forall x \in \text{supp}((1_J \psi_{m,n}) \circ f_{t,J}^{-n} \circ \mathcal{Z}_n^{-1}).$$

Applying Lemma 3.5 to $\mathcal{Z}_n \circ f_{t,J}^{-n} \circ \mathcal{Z}_n^{-1}$, we obtain $C = C(\tau, p)$ independent of n and t , and $C_n = C_n(p)$ independent of t such that

$$(60) \quad \begin{aligned} \|(1_J \psi_{m,n} \varphi) \circ f_{t,J}^{-n} \circ \mathcal{Z}_n^{-1}\|_{\mathcal{H}^{\tau,p}} \\ \leq C |(f_t^n)'(y)|^{\frac{1}{p}} (\lambda_n(f_t))^{-\tau} \|\tilde{\varphi}_{m,n,J}\|_{\mathcal{H}^{\tau,p}} + C_n \|\tilde{\varphi}_{m,n,J}\|_{L^p}, \quad \forall |t| \leq \varepsilon. \end{aligned}$$

¹⁸We mean $|1 - (f_t^n)'(\bar{x})/(f_t^n)'(y)| \leq K(f_t'', f_t') \cdot d(f_t^n(\bar{x}), f_t^n(y))$, and the zoom \mathcal{Z}_n ensures $K(f_t'', f_t') \cdot d(f_t^n(\bar{x}), f_t^n(y)) < 1/2$.

Since $1/p < 1$, the bound (60) together with Lemma 3.4 for¹⁹ $\gamma = 1$ imply

$$\begin{aligned} \left\| \left(1_J \psi_{m,n} \frac{\varphi}{|(f_t^n)'|} \right) \circ f_{t,J}^{-n} \right\|_n &\leq C |(f_t^n)'(y)|^{\frac{1}{p}} (\lambda_n(f_t))^{-\tau-1} \|\tilde{\varphi}_{m,n,J}\|_{\mathcal{H}^{\tau,p}} + C_n \|\tilde{\varphi}_{m,n,J}\|_{L^p} \\ &\leq C (\lambda_n(f_t))^{-(\tau+1-\frac{1}{p})} \|\tilde{\varphi}_{m,n,J}\|_{\mathcal{H}^{\tau,p}} + C_n \|\tilde{\varphi}_{m,n,J}\|_{L^p}. \end{aligned}$$

Therefore, by the convexity inequality $(|a| + |b|)^p \leq 2^{p-1}(|a|^p + |b|^p)$, we have

$$(61) \quad \left\| \left(1_J \psi_{m,n} \frac{\varphi}{|(f_t^n)'|} \right) \circ f_{t,J}^{-n} \right\|_n^p \leq C(p) (\lambda_n(f_t))^{-p(\tau+1-1)} \|\tilde{\varphi}_{m,n,J}\|_{\mathcal{H}^{\tau,p}}^p + C_n \|\tilde{\varphi}_{m,n,J}\|_{L^p}^p,$$

where $C(p)$ and C_n are independent of t . By the Strichartz Lemma 3.6 (we use here $0 \leq \tau < 1/p$) and the (classical) localisation Theorem 2.4.7 in [40] (or Lemma 3.4 in [38]) there exists $K = K(\psi, \tau, p) > 0$ such that for all m, n and φ

$$(62) \quad \begin{aligned} \sum_{m \in \mathbb{Z}} \|\tilde{\varphi}_{m,n,J}\|_{\mathcal{H}^{\tau,p}}^p &= \sum_{m \in \mathbb{Z}} \|(1_J \cdot (\psi_m \circ \mathcal{Z}_n) \cdot \varphi) \circ \mathcal{Z}_n^{-1}\|_{\mathcal{H}^{\tau,p}}^p \\ &\leq K \sum_{m \in \mathbb{Z}} \|((\psi_m \circ \mathcal{Z}_n) \cdot \varphi) \circ \mathcal{Z}_n^{-1}\|_{\mathcal{H}^{\tau,p}}^p \leq K^2 \|\varphi \circ \mathcal{Z}_n^{-1}\|_{\mathcal{H}^{\tau,p}}^p = K^2 \|\varphi\|_n^p. \end{aligned}$$

For a fixed map f_0 and for any n , if Z_n is large enough, then for any m there are at most two intervals $J \in \mathcal{A}(0, n)$ such that $(1_J \psi_{m,n}) \circ f_J^{-n}$ is not identically zero. (This is the ‘‘complexity at the beginning’’ in Thomine’s argument.) For a perturbation (f_t) , caution is needed, especially (but not only) if c is periodic for f (see footnote 7).

Assume first that c is not periodic. Fix $\tilde{\Lambda} < \bar{\Lambda} < \Lambda$, choose $p_0 \in (1, p(f))$ and $n_0 \geq 1$ such that

$$(63) \quad 2^{p_0-1} \cdot (\lambda_{n_0}(f_t))^{-p_0(\tau+1)-1} \leq \bar{\Lambda}^{n_0(-p_0(\tau+1)-1)}, \quad \forall |t| < \varepsilon_0.$$

Next, for $p \in (1, p_0)$, let $C(p)$ be the constant from (61), let $K(p)$ be the constant from (62). Then, there exists $N \geq n_0$ such that

$$(64) \quad 2 \cdot C(p) \cdot K(p)^2 \cdot 2^{N(p-1)} \cdot (\lambda_N(f_t))^{-(p(\tau+1)-1)} < \tilde{\Lambda}^{-N(p(\tau+1)-1)}.$$

Now, since c is not periodic, (see e.g. [13, p. 376]) there exists $\varepsilon' = \varepsilon'(N) < \varepsilon_0$ such that for any $|t| \leq \varepsilon'(N)$ there is a natural bijection $\mathcal{J}_t = \mathcal{J}_{t,N}$ between $\mathcal{A}(0, N)$ and $\mathcal{A}(t, N)$. In addition, we may ensure that $|f_t^N(\mathcal{J}_t(J))| \geq |f^N(J)|/2$ for all $J \in \mathcal{A}(0, N)$ and all $|t| \leq \varepsilon'(N)$. By bounded distortion, there exists Z_N such that for all $|t| \leq \varepsilon'(N)$ and any m there are at most two intervals $J \in \mathcal{A}(t, N)$ such that $(1_J \psi_{m,N} \varphi) \circ f_{t,J}^{-N}$ is not

¹⁹Since the C^1 norm of $1/(f_t^n)'$ is bounded uniformly in n and t by some D , choosing a suitable zoom Z_n allows to bound the C^1 norm of $(1_J \psi_{m,n}/(f_t^n)') \circ Z_n^{-1}$ by $\lambda_n(f_t)^{-1} + D/Z_n \leq C \lambda_n(f_t)^{-1}$.

identically zero. Therefore, it follows from (61) and (62) that

$$\begin{aligned}
 & \sum_{m \in \mathbb{Z}} \sum_{J \in \mathcal{A}(t, N)} \left\| \frac{(1_J \psi_{m, N} \varphi) \circ f_{t, J}^{-N}}{(f_t^N)' \circ f_{t, J}^{-N}} \right\|_N^p \\
 (65) \quad & \leq \sum_{m \in \mathbb{Z}} \sum_{J \in \mathcal{A}(t, N)} \left(C(\lambda_N(f_t))^{-(p(\tau+1)-1)} \|\tilde{\varphi}_{m, N, J}\|_{\mathcal{H}^{\tau, p}}^p + C_N \|\tilde{\varphi}_{m, N, J}\|_{L^p}^p \right) \\
 (66) \quad & \leq CK^2 (\lambda_N(f_t))^{-(p(\tau+1)-1)} \|\varphi\|_N^p + 2C_N \|\varphi\|_{L^p}^p, \quad \forall |t| \leq \varepsilon'(N).
 \end{aligned}$$

Recalling (59), we have shown that for all $|t| \leq \varepsilon'(N)$,

$$\|\mathcal{L}_t^N \varphi\|_N^p \leq CK^2 2^{N(p-1)} (\lambda_N(f_t))^{-(p(\tau+1)-1)} \|\varphi\|_N^p + 2C_N \|\varphi\|_{L^p}^p.$$

Finally, we use (57) twice to obtain

$$(67) \quad \|\mathcal{L}_t^N \varphi\|_{\mathcal{H}^{\tau, p}} \leq 2CK^2 2^{N(1-\frac{1}{p})} (\lambda_N(f_t))^{-(\tau+1-\frac{1}{p})} \|\varphi\|_{\mathcal{H}^{\tau, p}} + 2(C_N + C_{\text{zoom}}) \|\varphi\|_{L^p},$$

for all $|t| \leq \varepsilon'(N)$, which together with our choice (64) of N and the Sobolev embedding theorem completes the proof of (56) if c is not periodic, for $\varepsilon_0 := \varepsilon'(N)$.

If c is periodic but all the f_t are topologically conjugated to f_0 , the argument in the non periodic case can be applied.

Otherwise, if c is periodic of minimal period $P_f \geq 2$, we first take $\tilde{\Lambda} < \bar{\Lambda} < \Lambda$ and choose $p_0 \in (1, p(f))$ and $n_0 = k_0 P_f$ so that

$$(68) \quad 2^{k_0} 2^{p_0-1} \cdot (\lambda_{n_0}(f_t))^{-p_0(\tau+1)-1} \leq \bar{\Lambda}^{n_0(-p_0(\tau+1)-1)}, \quad \forall |t| < \varepsilon_0.$$

Then we take $p \in (1, p_0)$, and we fix $N = k_1 P_f \geq n_0$ large enough such that

$$(69) \quad 2^{k_1} \cdot C(p) \cdot K(p)^2 \cdot 2^{N(p-1)} \cdot (\lambda_N(f_t))^{-p(\tau+1)-1} < \tilde{\Lambda}^{-N(p(\tau+1)-1)}, \quad \forall |t| < \varepsilon_0.$$

Now, ([13, p. 376]), for $N = k_1 P_f$, there exists $\varepsilon'' = \varepsilon''(N) < \varepsilon_0$ such that for any $|t| \leq \varepsilon''(N)$, there is a natural injection $\mathcal{J}_t = \mathcal{J}_{t, N}$ from $\mathcal{A}(0, N)$ to $\mathcal{A}(t, N)$ such that $|f_t^N(\mathcal{J}_t(J))| \geq |f^N(J)|/2$ for all $J \in \mathcal{A}(0, N)$ and all $|t| \leq \varepsilon'(N)$, and, in addition, between any two intervals of $\mathcal{A}(t, N)$ in the image of $\mathcal{J}_{t, N}$, there are at most $2^{k_1} - 2$ intervals of $\mathcal{A}(t, N)$ which are *not* in the image of $\mathcal{J}_{t, N}$. Therefore, there exists Z_N (uniform in t) such that for any $|t| \leq \varepsilon''(N)$ and any $m \in \mathbb{Z}$, there are at most 2^{k_1} intervals $J \in \mathcal{A}(t, N)$ such that $(1_J \psi_{m, N} \varphi) \circ f_{t, J}^{-N}$ is not identically zero. Therefore, (67) is replaced by

$$\|\mathcal{L}_t^N \varphi\|_{\mathcal{H}^{\tau, p}} \leq 2^{k_1} [CK^2 2^{N(1-\frac{1}{p})} (\lambda_N(f_t))^{-(\tau+1-\frac{1}{p})}] \|\varphi\|_{\mathcal{H}^{\tau, p}} + 2(C_N + C_{\text{zoom}}) \|\varphi\|_{L^p},$$

for $N = k_1 P_f$ and all $|t| < \varepsilon''(N)$. Recalling (10) and (69), we may conclude for $\varepsilon_0 := \varepsilon''(N)$.

REFERENCES

- [1] A. Avila, *Infinitesimal perturbations of rational maps*, Nonlinearity **15** (2002) 695–704.
- [2] V. Baladi, *Positive Transfer Operators and Decay of Correlations*, World Scientific Publishing (2000).
- [3] V. Baladi, *On the susceptibility function of piecewise expanding interval maps*, Comm. Math. Phys. **275** (2007) 839–859.

- [4] V. Baladi, *Linear Response – Or Else*, Proceedings of the International Congress of Mathematicians-Seoul 2014. Vol. III. Invited lectures. 525–545. Papers from the congress (ICM 2014) held August 13–21, 2014. Seoul, 2014. vii+1250 pp. ISBN: 978-89-6105-806-3; 978-89-6105-803-2
- [5] V. Baladi, *Dynamical Zeta Functions and Dynamical Determinants for Hyperbolic Maps, A Functional Approach*, Ergebnisse der Mathematik und ihrer Grenzgebiete, Springer, Cham (2018).
- [6] V. Baladi, M. Benedicks, and D. Schnellmann, *Whitney Hölder continuity of the SRB measure for transversal families of smooth unimodal maps*, Invent. Math. **201** (2015) 773–844.
- [7] V. Baladi, T. Kuna, and V. Lucarini, *Linear and fractional response for the SRB measure of smooth hyperbolic attractors and discontinuous observables*, Nonlinearity **30** (2017) 1204–1220. (Corrigendum: Nonlinearity **30** (2017) C4–C6.)
- [8] V. Baladi, S. Marmi, and D. Sauzin, *Natural boundary for the susceptibility function of generic piecewise expanding unimodal maps*, Ergodic Theory Dynam. Systems **34** (2014) 777–800.
- [9] V. Baladi and D. Smania, *Linear response formula for piecewise expanding unimodal maps*, Nonlinearity **21** (2008) 677–711. (Corrigendum: Nonlinearity **25** (2012) 2203–2205.)
- [10] V. Baladi and D. Smania, *Smooth deformations of piecewise expanding unimodal maps*, Discrete Contin. Dyn. Syst. **23** (2009) 685–703.
- [11] V. Baladi and D. Smania, *Fractional susceptibility function for the quadratic family: Misiurewicz–Thurston parameters* Comm. Math. Phys. (2021) doi.org/10.1007/s00220-021-04015-z
- [12] V. Baladi and M. Tsujii, *Anisotropic Hölder and Sobolev spaces for hyperbolic diffeomorphisms*, Ann. Inst. Fourier **57** (2007) 127–154.
- [13] V. Baladi and L.-S. Young, *On the spectra of randomly perturbed expanding maps*, Comm. Math. Phys. **156** (1993) 355–385. (Erratum: Comm. Math. Phys. **166** (1994) 219–220.)
- [14] G. Contreras, *Regularity of topological and metric entropy of hyperbolic flows*, Math. Z. **210** (1992) 97–111.
- [15] F. Contreras, *Modulus of continuity of averages of SRB measures for a transversal family of piecewise expanding unimodal maps* (2016) arXiv:1604.03365.
- [16] A. de Lima and D. Smania, *Central limit theorem for the modulus of continuity of averages of observables on transversal families of piecewise expanding unimodal maps*, J. Inst. Math. Jussieu **17** (2018) 673–733.
- [17] S. V. Ershov, *Is a perturbation theory for dynamical chaos possible?* Physics Letters A **177** (1993) 180–185.
- [18] P. Eslami and P. Góra, *Stronger Lasota-Yorke inequality for one-dimensional piecewise expanding transformations*, Proc. Amer. Math. Soc. **141** (2013) 4249–4260.
- [19] S. Gouëzel and C. Liverani, *Banach spaces adapted to Anosov systems*, Ergodic Theory Dynam. Systems **26** (2006) 189–218.
- [20] L. Grafakos, *Modern Fourier Analysis*, Graduate Texts in Mathematics **250**, Third edition. Springer (2014).
- [21] R. Kubo, *The fluctuation-dissipation theorem*, Rep. Prog. Phys. **29** (1966) 255–284.
- [22] Y. Jiang and D. Ruelle, *Analyticity of the susceptibility function for unimodal Markovian maps of the interval*, Nonlinearity **18** (2005) 2447–2453.
- [23] A. Katok, G. Knieper, M. Pollicott, and H. Weiss, *Differentiability and analyticity of topological entropy for Anosov and geodesic flows*, Invent. Math. **98** (1989) 581–597.
- [24] G. Keller, *Stochastic stability in some chaotic dynamical systems*, Mon. Math. **94** (1982) 313–333.
- [25] G. Keller and C. Liverani, *Stability of the spectrum for transfer operators*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. **28** (1999) 141–152.
- [26] G. Levin, *On an analytic approach to the Fatou conjecture*, Fund. Math. **171** (2002) 177–196.
- [27] M. Mazzoleni, *Dinamiche espansive unidimensionali: dipendenza della misura invariante da un parametro*, Master’s Thesis, Roma 2 (2007).
- [28] K.S. Miller and B. Ross, *An Introduction to the Fractional Calculus and Fractional Differential Equations*, John Wiley & Sons (1993).
- [29] M. Porte, *Linear response for Dirac observables of Anosov diffeomorphisms*, Discrete Contin. Dyn. Syst. **39** (2019) 1799–1819.
- [30] D. Ruelle, *Differentiation of SRB states*, Comm. Math. Phys. **187** (1997) 227–241.

- [31] D. Ruelle, *General linear response formula in statistical mechanics, and the fluctuation-dissipation theorem far from equilibrium*, Phys. Lett. A **245** (1998) 220–224.
- [32] D. Ruelle, *Differentiating the absolutely continuous invariant measure of an interval map f with respect to f* , Comm. Math. Phys. **258** (2005) 445–453.
- [33] D. Ruelle, *Linear response theory for diffeomorphisms with tangencies of stable and unstable manifolds — a contribution to the Gallavotti-Cohen chaotic hypothesis*, Nonlinearity **31** (2018) 5683–5691.
- [34] T. Runst and W. Sickel, *Sobolev spaces of Fractional Order, Nemytskij Operators, and Nonlinear Partial Differential Equations*, Walter de Gruyter & Co., Berlin (1996).
- [35] S. G. Samko, A. A. Kilbas, and O. I. Marichev, *Fractional Integrals and Derivatives. Theory and Applications*, Gordon and Breach Science Publishers (1993).
- [36] B. Saussol, *Absolutely continuous invariant measures for multidimensional expanding maps*, Israel J. Math. **116** (2000) 223–248.
- [37] R. S. Strichartz, *Multipliers on fractional Sobolev spaces*, J. Math. Mech. **16** (1967) 1031–1060.
- [38] D. Thomine, *A spectral gap for transfer operators of piecewise expanding maps*, Discrete Contin. Dyn. Syst. **30** (2011) 917–944.
- [39] H. Triebel, *Interpolation Theory, Function Spaces, Differential Operators*, North Holland, Amsterdam (1978).
- [40] H. Triebel, *Theory of function spaces II*, Birkhäuser, Basel (1992).
- [41] M. Tsujii, *A simple proof for monotonicity of entropy in the quadratic family*, Ergodic Theory Dynam. Systems **20** (2000) 925–933.
- [42] L.-S. Young, *What are SRB measures, and which dynamical systems have them?*, J. Stat. Phys. **108** (2002) 733–754.

†CENTRE FOR MATHEMATICAL SCIENCES, LUND UNIVERSITY, BOX 118, 221 00 LUND, SWEDEN

Email address: `magnusa@maths.lth.se`

Email address: `tomas@maths.lth.se`

††LABORATOIRE DE PROBABILITÉS, STATISTIQUE ET MODÉLISATION (LPSM), CNRS,
SORBONNE UNIVERSITÉ, UNIVERSITÉ DE PARIS, 4 PLACE JUSSIEU, 75005 PARIS, FRANCE

Email address: `baladi@lpsm.paris`

Email address: `lepp.juho@gmail.com`