

# Stability Conditions and Liapunov Functions for Quasipolynomial Systems

BENITO HERNÁNDEZ–BERMEJO<sup>1</sup>

Departamento de Física Matemática y Fluidos, Facultad de Ciencias  
Universidad Nacional de Educación a Distancia  
Senda del Rey S/N, 28040 Madrid, Spain.

**Abstract**— The stability of equilibrium points of quasipolynomial systems of ODEs is considered. The criteria and Liapunov functions found generalize those traditionally known for Lotka-Volterra equations, that now appear as a particular case.

**Keywords**— Stability, Liapunov functions, quasipolynomial systems, Lotka-Volterra systems.

<sup>1</sup> E-mail: bhernand@apphys.uned.es

Telephone: (+ 34 91) 398 72 19. Fax: (+ 34 91) 398 66 97.

Consider the well-known Lotka-Volterra (LV) system

$$\dot{x}_i = x_i \left( \lambda_i + \sum_{j=1}^n A_{ij} x_j \right), \quad i = 1, \dots, n \quad (1)$$

which is assumed to have a unique positive equilibrium point:  $x^* = (x_1^*, \dots, x_n^*)^T \in \text{int}(\mathbb{R}_+^n)$ . A question of fundamental importance in the analysis of these equations concerns the stability of the steady state (see [1, pp. 3-5] for the main definitions of stability, which are the ones adopted in this work). In this sense, many of the most relevant results about stability of LV systems are those based on the Liapunov function [2-5]:

$$V(x) = \sum_{i=1}^n c_i \left( x_i - x_i^* - x_i^* \ln \frac{x_i}{x_i^*} \right), \quad c_i > 0 \quad \forall i \quad (2)$$

The time derivative of (2) along the trajectories of (1) is

$$\dot{V}(x) = \frac{1}{2} (x - x^*)^T (C \cdot A + A^T \cdot C) (x - x^*) \quad (3)$$

where  $C = \text{diag}(c_1, \dots, c_n)$ . Thus, it can be stated that if there exists a positive definite diagonal matrix  $C$  such that  $C \cdot A + A^T \cdot C$  is negative semi-definite (i.e.  $A \in \bar{S}_w$ ) then  $x^*$  is stable. Moreover if, instead,  $C \cdot A + A^T \cdot C$  is negative definite (that is,  $A \in S_w$ ) then  $x^*$  is globally asymptotically stable in  $\text{int}(\mathbb{R}_+^n)$ .

A natural and extensive generalization of LV systems is provided by quasipolynomial (QP) systems of ODEs (see [6] and references therein):

$$\dot{x}_i = x_i \left( \lambda_i + \sum_{j=1}^m A_{ij} \prod_{k=1}^n x_k^{B_{jk}} \right), \quad i = 1, \dots, n, \quad m \geq n, \quad \text{Rank}(B) = n \quad (4)$$

Conditions  $m \geq n$  and  $\text{Rank}(B) = n$  can be assumed without loss of generality [6] because systems not verifying them are reducible to the form (4) in  $\text{int}(\mathbb{R}_+^n)$ . Therefore, (4) is a completely generic starting point for what is to follow.

The aim of this letter is to demonstrate that the aforementioned properties regarding the stability of LV equations (1) can be generalized to a great extent for QP systems (4). This is interesting from an applied perspective, because most systems of ODEs arising in practice are QP or can be algorithmically recasted in such form [6,7].

For convenience let us first define the  $m \times m$  matrix  $Q = B \cdot A$ , as well as the quasimonomial functions:

$$\varphi_i(x) = \prod_{j=1}^n x_j^{B_{ij}}, \quad i = 1, \dots, m \quad (5)$$

The main result is the following one:

**THEOREM 1.** For every fixed point  $x^* \in \text{int}(\mathbb{R}_+^n)$  of a QP system (4):

a) If  $Q \in \bar{S}_w$  then  $x^*$  is stable.

b) If  $Q \in S_w$  then  $x^*$  is globally asymptotically stable in  $\text{int}(\mathbb{R}_+^n)$ .

c) Let  $C = \text{diag}(c_1, \dots, c_m)$ , with  $c_i > 0 \forall i$ , be a matrix such that  $C \cdot Q + Q^T \cdot C$  is negative semi-definite in case (a) or negative definite in case (b). Then, in either (a) or (b),

$$W(x) = \sum_{i=1}^m c_i \left\{ \prod_{j=1}^n x_j^{B_{ij}} - \prod_{j=1}^n (x_j^*)^{B_{ij}} - \left( \prod_{j=1}^n (x_j^*)^{B_{ij}} \right) \ln \left[ \prod_{j=1}^n \left( \frac{x_j}{x_j^*} \right)^{B_{ij}} \right] \right\} \quad (6)$$

is a Liapunov function for  $x^*$  in  $\text{int}(\mathbb{R}_+^n)$ .

d) The time variation of the Liapunov function (6) along the trajectories of (4) is:

$$\dot{W} = \frac{1}{2} (\varphi(x) - \varphi(x^*))^T (C \cdot Q + Q^T \cdot C) (\varphi(x) - \varphi(x^*)) \quad (7)$$

where  $\varphi(x) = (\varphi_1(x), \dots, \varphi_m(x))^T$  and the  $\varphi_i(x)$  are defined in (5).

**PROOF.** Notice that  $W(x^*) = 0$ . In addition, we shall first demonstrate that function  $W(x)$  in (6) verifies  $W(x) > 0$  for all  $x \neq x^*$ ,  $x \in \text{int}(\mathbb{R}_+^n)$ . For this, consider the  $m$ -dimensional function:

$$V(y_1, \dots, y_m) = \sum_{i=1}^m c_i \left( y_i - y_i^* - y_i^* \ln \frac{y_i}{y_i^*} \right), \quad c_i > 0 \forall i, \quad y_i^* > 0 \forall i \quad (8)$$

Obviously,  $V(y^*) = 0$  and  $V(y) > 0$  for every  $y \neq y^*$ ,  $y \in \text{int}(\mathbb{R}_+^m)$ . Let us now perform the following change of variables:

$$y_i = \prod_{j=1}^m z_j^{\tilde{B}_{ij}}, \quad i = 1, \dots, m \quad (9)$$

In (9) we define  $\tilde{B} = (B \mid B_{m \times (m-n)}^*)$ , where  $B^*$  is an  $m \times (m-n)$  submatrix of arbitrary entries to be chosen in such a way that  $\tilde{B}$  is invertible (note that this is always possible since  $\text{Rank}(B) = n$ ). Consequently, transformation (9) is bijective in  $\text{int}(\mathbb{R}_+^m)$ . Let  $z^* = (z_1^*, \dots, z_m^*)^T \in \text{int}(\mathbb{R}_+^m)$  be the unique point whose image is  $y^*$  after transformation (9). Then, function  $V(y)$  in (8) is mapped to the  $m$ -dimensional function  $\tilde{W}(z) = V(y(z))$ , defined in  $\text{int}(\mathbb{R}_+^m)$ :

$$\tilde{W}(z) = \sum_{i=1}^m c_i \left\{ \prod_{j=1}^m z_j^{\tilde{B}_{ij}} - \prod_{j=1}^m (z_j^*)^{\tilde{B}_{ij}} - \left( \prod_{j=1}^m (z_j^*)^{\tilde{B}_{ij}} \right) \ln \left[ \prod_{j=1}^m \left( \frac{z_j}{z_j^*} \right)^{\tilde{B}_{ij}} \right] \right\} \quad (10)$$

Thus we have that  $\tilde{W}(z^*) = 0$  and  $\tilde{W}(z) > 0$  for every  $z \neq z^*$ ,  $z \in \text{int}(\mathbb{R}_+^m)$ .

If  $m = n$ , then  $\tilde{B}$  reduces to  $B$  and if we replace in (10)  $z$  and  $z^*$  by  $x$  and  $x^*$ , respectively, it is demonstrated that  $W(x)$  in (6) verifies  $W(x) > 0$  for all  $x \neq x^*$ ,  $x \in \text{int}(\mathbb{R}_+^n)$ .

To prove the same in the complementary  $m > n$  case, let us choose a point  $z^*$  of the form  $z^* = (x_1^*, \dots, x_n^*, 1, \overset{(m-n)}{\dots}, 1)^T$ . We know that  $\tilde{W}(z) > 0$  for every  $z \neq z^*$ ,  $z \in \text{int}(\mathbb{R}_+^m)$ . Then this will be the case, in particular, for the points of the hyperplane  $\{z_i = 1, i = n+1, \dots, m\}$  belonging to  $\text{int}(\mathbb{R}_+^m)$ . Let us parametrize those points as  $z = (x_1, \dots, x_n, 1, \overset{(m-n)}{\dots}, 1)^T$ , with  $x_i > 0$  for all  $i = 1, \dots, n$ . Thus if we substitute  $z^*$  and the parametrized form of  $z$  in  $\tilde{W}$ , it is immediately obtained that  $\tilde{W}(x_1, \dots, x_n, 1, \overset{(m-n)}{\dots}, 1) = W(x_1, \dots, x_n)$  in (6) because  $\tilde{B}_{ij} = B_{ij}$  for  $j = 1, \dots, n$ . This proves that  $W(x) > 0$  for all  $x \neq x^*$ ,  $x \in \text{int}(\mathbb{R}_+^n)$ , in the  $m > n$  case.

Finally, let us look at the time evolution of  $W(x)$ . For this, note that we can write the QP system (4) as:

$$\dot{x}_i = x_i \sum_{j=1}^m A_{ij}(\varphi_j(x) - \varphi_j(x^*)), \quad i = 1, \dots, n \quad (11)$$

where  $\varphi_i(x)$  is defined in (5). Note also that

$$\frac{\partial \varphi_i(x)}{\partial x_j} = \frac{B_{ij} \varphi_i(x)}{x_j} \quad (12)$$

We then have:

$$\dot{W} = \sum_{i=1}^m c_i \dot{\varphi}_i(x) \left(1 - \frac{\varphi_i(x^*)}{\varphi_i(x)}\right) = \sum_{i=1}^m c_i \left(1 - \frac{\varphi_i(x^*)}{\varphi_i(x)}\right) \sum_{k=1}^n \frac{\partial \varphi_i(x)}{\partial x_k} \dot{x}_k \quad (13)$$

Taking (11) and (12) into account, (7) is found after some simple algebra.

Consequently [1, pp. 3-5], we have that if  $Q \in \bar{S}_w$  ( $Q \in S_w$ ) then  $W(x)$  is a Liapunov function for the system in  $\text{int}(\mathbb{R}_+^n)$  and  $x^*$  is stable (globally asymptotically stable in  $\text{int}(\mathbb{R}_+^n)$ ).

This completes the proof of the theorem.

REMARK 1. Notice that the aforementioned results for LV systems appear now as a particular case of Theorem 1 when  $m = n$ ,  $B$  is the identity matrix and  $\text{Rank}(A) = n$ .

REMARK 2. The previously known criteria for the belonging of a matrix to  $S_w$  or  $\bar{S}_w$ , widely investigated in the context of the stability of LV equations (see [5] and references therein) can now be extended straightforwardly to QP systems.

COROLLARY 1. Under the same hypotheses of Theorem 1, if  $\dot{W}(x) = 0$  for all  $x \in \text{int}(\mathbb{R}_+^n)$ , then the trajectories of system (4) lie on the surfaces:

$$\sum_{i=1}^m c_i \left\{ \prod_{j=1}^n x_j^{B_{ij}} - \prod_{j=1}^n (x_j^*)^{B_{ij}} - \left( \prod_{j=1}^n (x_j^*)^{B_{ij}} \right) \ln \left[ \prod_{j=1}^n \left( \frac{x_j}{x_j^*} \right)^{B_{ij}} \right] \right\} = \text{constant} \quad (14)$$

REMARK 3. Corollary 1 takes place, for instance, in QP systems described in terms of a Poisson structure, which are closely related [8] to conservative LV systems (see [4] for a review of LV conservativeness). In such cases the Liapunov function (14) is a first integral playing the role of Hamiltonian.

## REFERENCES

1. J. Guckenheimer and P. Holmes, *Nonlinear Oscillations, Dynamical Systems, and Bifurcations of Vector Fields*, Springer-Verlag, New York (1983).
2. B. S. Goh, *Management and Analysis of Biological Populations*, Elsevier, Oxford-Amsterdam-New York (1980).
3. J. Hofbauer and K. Sigmund, *The Theory of Evolution and Dynamical Systems*, Cambridge University Press, Cambridge (1988).
4. D. O. Logofet, *Matrices and Graphs. Stability Problems in Mathematical Ecology*, CRC Press, Boca Raton, Florida (1993).
5. Y. Takeuchi, *Global Dynamical Properties of Lotka-Volterra Systems*, World Scientific, Singapore (1996).
6. B. Hernández-Bermejo, V. Fairén and L. Brenig, Algebraic recasting of nonlinear systems of ODEs into universal formats, *J. Phys. A, Math. Gen.* **31**, 2415-2430 (1998).
7. B. Hernández-Bermejo and V. Fairén, Lotka-Volterra representation of general nonlinear systems, *Math. Biosci.* **140**, 1-32 (1997).
8. B. Hernández-Bermejo and V. Fairén, Hamiltonian structure and Darboux theorem for families of generalized Lotka-Volterra systems, *J. Math. Phys.* **39**, 6162-6174 (1998).