

Tau-functions and monodromy symplectomorphisms

M. Bertola^{†‡1}, D. Korotkin^{† 2},

[†] *Department of Mathematics and Statistics, Concordia University
1455 de Maisonneuve W., Montréal, Québec, Canada H3G 1M8*

[‡] *SISSA/ISAS, Area of Mathematics
via Bonomea 265, 34136 Trieste, Italy*

Abstract

We derive a new Hamiltonian formulation of Schlesinger equations in terms of the dynamical r -matrix structure. The corresponding symplectic form is shown to be the pullback, under the monodromy map, of a natural symplectic form on the extended monodromy manifold. We show that Fock-Goncharov coordinates are log-canonical for the symplectic form. Using these coordinates we define the symplectic potential on the monodromy manifold and interpret the Jimbo-Miwa-Ueno tau-function as the generating function of the monodromy map. This, in particular, solves a recent conjecture by A. Its, O. Lisovyy and A. Prokhorov.

Contents

1	Introduction	1
2	Dynamical r-matrix formulation of the Schlesinger system	8
2.1	Quadratic Poisson bracket via dynamical r -matrix	8
2.1.1	Reduction to $SL(n)$	10
2.1.2	Relation to the Kirillov-Kostant bracket	11
2.2	Hamiltonian formulation of the Schlesinger system in G -variables	12
2.3	Symplectic form and potential	12
3	Monodromy symplectomorphism via Malgrange's form	13
4	Standard monodromy map and weak version of Its-Lisovyy-Prokhorov conjecture	18
5	Log-canonical coordinates and symplectic potential	22
5.1	Fock-Goncharov coordinates	22
5.2	Parametrization of the space \mathcal{M}	26
5.3	Symplectic form	26
5.4	Symplectic potential	28
5.5	$SL(2)$ case	28
5.5.1	Change of triangulation.	28
6	Tau-function as generating function of monodromy symplectomorphism	29
6.1	$SL(2)$ tau-function	31
6.2	Equations with respect to Fock-Goncharov coordinates	31

1 Introduction

Symplectic aspects of the monodromy map for the Fuchsian systems were studied starting from [33, 3, 42]; in these papers it was proved that the monodromy map is a symplectomorphism from a symplectic leaf in the space of coefficients of the system to a symplectic leaf in the monodromy manifold. The non-Fuchsian case was considered in [19, 13, 51, 11, 15]. Remarkably, the simplest non-Fuchsian case of the Painlevé II hierarchy was treated in the paper [19] in 1981, about 15 years before the Fuchsian

¹Marco.Bertola@{concordia.ca,sissa.it}

²Dmitry.Korotkin@concordia.ca

case was studied in detail in [33, 3, 42]. The key object associated to any Fuchsian or non-Fuchsian system of linear ODE's is the tau-function introduced by Jimbo, Miwa and Ueno [48]; until now its significance in the framework of the monodromy symplectomorphism remained unclear; the main goal of this paper is to fill this gap and prove a recent conjecture by Its-Lisovsky-Prokhorov [36].

Our interest in this subject stems from the study of monodromy map of a second order equation on a Riemann surface; such a map was also proven to be a symplectomorphism [41, 9, 43, 10]. Remarkably, the generating function of the monodromy symplectomorphism (the "Yang-Yang" function) plays an important role in the theory of supersymmetric Yang-Mills equations [46]; several steps towards understanding of this generating function were made in [9].

In this paper we address the question about the role of the generating function of the monodromy symplectomorphism in the context of Fuchsian equations on the Riemann sphere. The conclusion we arrive to is somewhat unexpected: such generating function can be naturally identified with the isomonodromic tau-function; moreover, this interpretation allows to define the dependence of the tau-function on monodromy data.

The version of the monodromy map for Fuchsian systems used in the current paper is slightly different from the monodromy map considered in [33, 3, 42]. This version is standard in the theory of isomonodromy deformations [48] and it was also considered in [37, 11] from the symplectic point of view.

To describe the monodromy map we remind the basics of the theory of solutions of Fuchsian systems of differential equations on \mathbb{P}^1 , following [48]. Consider the equation

$$\frac{\partial \Psi}{\partial z} = \sum_{i=1}^N \frac{A_i}{z - t_i} \Psi, \quad \Psi(z = \infty) = \mathbf{1}, \quad (1.1)$$

where $A_i \in sl(n)$ and $t_j \neq t_k$ such that $\sum_{i=1}^N A_i = 0$. Assume also that eigenvalues of each A_j are simple and furthermore do not differ by an integer. Choose a system of cuts $\gamma_1, \dots, \gamma_N$ connecting ∞ with t_1, \dots, t_N respectively, and assume that the ends of these cuts emanating from ∞ are ordered as $(1, \dots, N)$ counter-clockwise (Fig.1). The normalization condition $\Psi(\infty) = \mathbf{1}$ is then understood in the sense that $\lim_{z \rightarrow \infty} \Psi(z) = \mathbf{1}$ where the limit is taken in the sector between γ_1 and γ_N .

The set of generators $\sigma_1, \dots, \sigma_N$ of the fundamental group $\pi_1(\mathbb{P}^1 \setminus \{t_j\}_{j=1}^N, \infty)$ is chosen such that the loop representing σ_j crosses only the cut γ_j , and its orientation is chosen so that the relation between σ_j takes the form $\sigma_N \cdots \sigma_1 = \text{Id}$ (Fig.1).




Figure 1: Choice of cuts and generators of the fundamental group.

The solution Ψ of (1.1) is single-valued in the simply connected domain $\mathbb{P}^1 \setminus \{\gamma_j\}_{j=1}^N$. Denote the diagonal form of the matrix A_j by L_j , $j = 1, \dots, N$ (the matrices A_j are diagonalizable due to our assumption about their eigenvalues). Then the asymptotics of Ψ near t_j has the standard form [48]:

$$\Psi(z) = (G_j + O(z - t_j))(z - t_j)^{L_j} C_j^{-1}. \quad (1.2)$$

The matrix G_j is a diagonalizing matrix for A_j :

$$A_j = G_j L_j G_j^{-1} . \quad (1.3)$$

The matrices C_j are called the *connection matrices*. Notice that the matrices G_j and C_j are not uniquely defined by equation (1.1) since a simultaneous transformation $G_j \rightarrow G_j D_j$ and $C_j \rightarrow C_j D_j$ with diagonal D_j 's changes neither the asymptotics (1.2) nor the equation (1.1).

Analytic continuation of $\Psi(z)$ along σ_j yields $\Psi(z)M_j^{-1}$, where the *monodromy matrix* $M_j \in SL(n)$ is related to the connection matrix C_j and the exponent of monodromy L_j by:

$$M_j = C_j \Lambda_j C_j^{-1} , \quad \Lambda_j := e^{2i\pi L_j} . \quad (1.4)$$

Alternatively, the matrix M_j can be viewed as the jump matrix on γ_j : orienting γ_j from ∞ towards t_j , the boundary values of Ψ on the right ("+") and left ("-") sides of γ_j are related by $\Psi_+ = \Psi_- M_j$. Our assumption about the ordering of the branch cuts γ_j and generators σ_j implies the relation

$$M_1 \cdots M_N = \mathbf{1} . \quad (1.5)$$

The *monodromy map* introduced in [48] sends the set of pairs (G_j, L_j) to the set of pairs (C_j, Λ_j) for a given set of poles t_j .

The map between the set of coefficients A_j and the set of monodromy matrices M_j is a different version of monodromy map associated to equation (1.1); the symplectic aspects of this version of the monodromy map were studied in [33, 3, 42].

To describe our framework in more details we introduce the following two spaces. The first space is the quotient

$$\mathcal{A} = \left\{ (G_j, L_j)_{j=1}^N, \quad G_j \in SL(n), \quad L_j \in \mathfrak{h}_{ss}^{nr}, \quad \forall j = 1, \dots, N : \quad \sum_{j=1}^N G_j L_j G_j^{-1} = 0 \right\} / \sim \quad (1.6)$$

where \mathfrak{h}_{ss}^{nr} denotes the set of diagonal matrices with simple eigenvalues not differing by integers (*non-resonant*). The equivalence relation is given by the $SL(n)$ action $G_j \mapsto S G_j$ with S independent of j .

The second space is the quotient

$$\mathcal{M} = \left\{ \{C_j, L_j\}_{j=1}^N, \quad C_j \in SL(n), \quad L_j \in \mathfrak{h}_{ss}^{nr} : \quad \prod_{j=1}^N C_j e^{2\pi i L_j} C_j^{-1} = \mathbf{1} \right\} / \sim . \quad (1.7)$$

Similarly to (1.6), the equivalence is given by the $SL(n)$ action $C_j \mapsto S C_j$ (with the same S for all j 's).

For a fixed set of poles $\{t_j\}_{j=1}^N$ we denote the "monodromy map" from the (G, L) -space to (C, L) -space by

$$\mathcal{F}^t : \mathcal{A} \rightarrow \mathcal{M} . \quad (1.8)$$

Poisson and symplectic structures on \mathcal{A} and dynamical r -matrix. Let $\mathcal{H} := SL(n, \mathbb{C}) \times \mathfrak{h}_{ss} = \{(G, L)\}$. Here $G \in SL(N)$ and $L = \text{diag}(\lambda_1, \dots, \lambda_n)$ is a diagonal traceless matrix with $\lambda_j \neq \lambda_k$ and $\lambda_1 + \dots + \lambda_n = 0$. Consider the following 1-form on \mathcal{H} :

$$\theta = \text{tr}(L G^{-1} dG) . \quad (1.9)$$

We prove in Prop. 2.1 that the form $\omega = d\theta$ is non-degenerate, and therefore, is a symplectic form on \mathcal{H} . For any matrix M we use the following notation for the Kronecker products

$$\overset{1}{M} = M \otimes \mathbf{1} , \quad \overset{2}{M} = \mathbf{1} \otimes M .$$

Then the Poisson structure on \mathcal{H} associated to the symplectic form ω is (see Prop. 2.2):

$$\left\{ \overset{1}{G}, \overset{2}{G} \right\} = -\overset{1}{G} \overset{2}{G} r(L) , \quad \left\{ \overset{1}{G}, \overset{2}{L} \right\} = -\overset{1}{G} \Omega , \quad (1.10)$$

where

$$r(L) = \sum_{i < j} \frac{E_{ij} \otimes E_{ji} - E_{ji} \otimes E_{ij}}{\lambda_i - \lambda_j} \quad (1.11)$$

and

$$\Omega = \sum_{i=1}^n E_{ii} \otimes E_{ii} - \frac{1}{n} \mathbf{1} \otimes \mathbf{1} ;$$

we use the standard notation E_{ij} for the matrix with only one non-vanishing element equal to 1 in the (i, j) entry. The matrix $r(L)$ is a simplest example of dynamical r -matrix [17]. Theorem 2.1 shows that the bracket (1.10) induces the Kirillov-Kostant Poisson bracket for $A = GLG^{-1}$.

The bracket (1.10) can be used to define the Poisson structure on the space \mathcal{A} as follows. Denote first by \mathcal{A}_0 the space of pairs $\{(G_j, L_j)\}_{j=1}^N$ with the product symplectic structure, or, equivalently, with the following Poisson bracket:

$$\{\overset{1}{G}_j, \overset{2}{G}_k\} = -\overset{1}{G}_j \overset{2}{G}_k r(L_k) \delta_{jk} , \quad \{\overset{1}{G}_j, \overset{2}{L}_k\} = -\overset{1}{G}_k \Omega \delta_{jk} . \quad (1.12)$$

The moment map corresponding to the group action $G_j \rightarrow SG_j$ ($S \in SL(n)$) on \mathcal{A}_0 is given by $\sum_{j=1}^N G_j L_j G_j^{-1}$. The space \mathcal{A} (1.6) inherits a symplectic form from \mathcal{A}_0 via the standard symplectic reduction [4]:

Theorem 1 (see Thm. 2.3). *The Poisson structure induced on \mathcal{A} from the Poisson structure (1.12) on \mathcal{A}_0 via the reduction on the level set $\sum_{j=1}^N G_j L_j G_j^{-1} = 0$ of the moment map, corresponding to the group action $G_j \rightarrow SG_j$, is non-degenerate and the corresponding symplectic form is $\omega_{\mathcal{A}} = d\theta_{\mathcal{A}}$, where the symplectic potential $\theta_{\mathcal{A}}$ for $\omega_{\mathcal{A}}$ is given by*

$$\theta_{\mathcal{A}} = \sum_{k=1}^N \text{tr}(L_k G_k^{-1} dG_k) .$$

This symplectic structure appeared in [11] but the connection to dynamical r -matrix and associated Poisson structure was not known until now.

Symplectic structure on \mathcal{M} . Define the following 2-form on the space \mathcal{M} :

$$\omega_{\mathcal{M}} = \frac{1}{4\pi i} (\omega_1 + \omega_2)$$

where

$$\begin{aligned} \omega_1 &= \text{tr} \sum_{\ell=1}^N (M_{\ell}^{-1} dM_{\ell} \wedge K_{\ell}^{-1} dK_{\ell}) + \text{tr} \sum_{\ell=1}^N (\Lambda_{\ell}^{-1} C_{\ell}^{-1} dC_{\ell} \wedge \Lambda_{\ell} C_{\ell}^{-1} dC_{\ell}) \\ \omega_2 &= 2 \sum_{\ell=1}^N \text{tr} (\Lambda_{\ell}^{-1} d\Lambda_{\ell} \wedge C_{\ell}^{-1} dC_{\ell}) \end{aligned}$$

with $K_{\ell} = M_1 \cdots M_{\ell}$ and $\Lambda_j = e^{2\pi i L_j}$.

The form $-\omega_1/2$ coincides with the symplectic form on the symplectic leaves $\Lambda_j = \text{const}$ of the $SL(n)$ Goldman bracket (see (3.14) of [2] in the case $g = 0$). The first result of this paper (see Thm.3.2 and its proof in Section 3) is that given a set of poles $\{t_j\}_{j=1}^N$ and a point $p_0 \in \mathcal{M}$ in a neighbourhood of which the monodromy map is invertible, the pullback of the form $\omega_{\mathcal{M}}$ under the map $\mathcal{F}^t : \mathcal{A} \rightarrow \mathcal{M}$ coincides with $\omega_{\mathcal{A}}$:

$$(\mathcal{F}^t)^* \omega_{\mathcal{M}} = \omega_{\mathcal{A}} . \quad (1.13)$$

This statement implies that (see Corollary 3.1 and its proof) the form $\omega_{\mathcal{M}}$ is closed and non-degenerate, and, therefore, defines a symplectic structure on \mathcal{M} .

For a given set of monodromy data the monodromy map is invertible outside of a locus of codimension 1 in the space of poles [12]. Since the form $\omega_{\mathcal{M}}$ is independent of $\{t_j\}$, this form is always non-degenerate on the monodromy manifold.

The equality (1.13) generalizes the results of [33, 3, 42], where it was proved that the monodromy map between the ‘‘smaller’’ spaces - the space of coefficients A_j with fixed eigenvalues and the symplectic leaf of the $GL(n)$ character variety of N -punctured sphere - is a symplectomorphism; the formula (1.13) was proved in a different way in [11].

Time dependence. To describe the dependence on the t_j 's (the ‘‘times’’) we extend the spaces \mathcal{A} and \mathcal{M} to include also the coordinates $\{t_j\}$:

$$\tilde{\mathcal{A}} = \left\{ (p, \{t_j\}_{j=1}^N), p \in \mathcal{A}, t_j \in \mathbb{C}, t_j \neq t_k \right\}, \quad (1.14)$$

$$\tilde{\mathcal{M}} = \left\{ (p, \{t_j\}_{j=1}^N), p \in \mathcal{M}, t_j \in \mathbb{C}, t_j \neq t_k \right\}. \quad (1.15)$$

The monodromy map \mathcal{F}^t then extends to the map $\mathcal{F} : \tilde{\mathcal{A}} \rightarrow \tilde{\mathcal{M}}$. The locus in $\tilde{\mathcal{M}}$ where the map is not invertible is usually referred to as the *Malgrange divisor*. Denote the pullback of the form $\omega_{\mathcal{A}}$ from \mathcal{A} to $\tilde{\mathcal{A}}$ by $\tilde{\omega}_{\mathcal{A}}$ and the pullback of the form $\omega_{\mathcal{M}}$ from \mathcal{M} to $\tilde{\mathcal{M}}$ by $\tilde{\omega}_{\mathcal{M}}$ (notice that the forms $\tilde{\omega}_{\mathcal{A}}$ and $\tilde{\omega}_{\mathcal{M}}$ are closed but degenerate). Now we are in a position to formulate the next theorem (see Thm.3.1)

Theorem 2 (see Thm.3.1 together with Thm.3.2). *The following identity holds between two-forms on $\tilde{\mathcal{A}}$*

$$\mathcal{F}^* \tilde{\omega}_{\mathcal{M}} = \tilde{\omega}_{\mathcal{A}} - \sum_{k=1}^N dH_k \wedge dt_k \quad (1.16)$$

where

$$H_k = \sum_{j \neq k} \frac{\text{tr} A_j A_k}{t_k - t_j}, \quad k = 1, \dots, N \quad (1.17)$$

are the canonical Hamiltonians of the Schlesinger system.

We remind that the Schlesinger equations [12] consist of the following system of PDEs for the coefficients of $A(z)$

$$\frac{\partial A_k}{\partial t_j} = \frac{[A_k, A_j]}{t_k - t_j}, \quad j \neq k; \quad \frac{\partial A_j}{\partial t_j} = - \sum_{k \neq j} \frac{[A_k, A_j]}{t_k - t_j} \quad (1.18)$$

and they define the deformations of the connection $A(z)$ which preserve the monodromy representation. They are Hamiltonian equations with respect to the standard Kirillov-Kostant Poisson bracket with time-dependent Hamiltonians H_k as in (1.17).

Tau function and generating function of the monodromy map. The above theorem allows to establish the relationship between the isomonodromic tau-function and the generating function of the monodromy map. Namely, consider some local symplectic potential $\theta_{\mathcal{M}}$ for the form $\omega_{\mathcal{M}}$ such that

$$d\theta_{\mathcal{M}} = \omega_{\mathcal{M}}$$

on the space \mathcal{M} (globally $\theta_{\mathcal{M}}$ can be defined on a covering of \mathcal{M}) and denote its pullback to $\tilde{\mathcal{M}}$ by $\tilde{\theta}_{\mathcal{M}}$. Denote by $\tilde{\theta}_{\mathcal{A}}$ the pullback of $\theta_{\mathcal{A}}$ under the natural projection $\tilde{\mathcal{A}} \rightarrow \mathcal{A}$. Then $\tilde{\theta}_{\mathcal{A}}$ is the potential of the symplectic form $\tilde{\omega}_{\mathcal{A}}$ on $\tilde{\mathcal{A}}$, and (1.16) implies existence of a locally defined generating function \mathcal{G} on $\tilde{\mathcal{A}}$.

Definition 1.1. *The generating function (corresponding to a given choice of the symplectic potential $\theta_{\mathcal{M}}$) of the monodromy map between spaces $\tilde{\mathcal{A}}$ and $\tilde{\mathcal{M}}$ is defined by*

$$d\mathcal{G} = \tilde{\theta}_{\mathcal{A}} - \sum_{j=1}^N H_j dt_j - \mathcal{F}^* \tilde{\theta}_{\mathcal{M}}. \quad (1.19)$$

A different choice of $\theta_{\mathcal{M}}$ (and hence of its pullback $\tilde{\theta}_{\mathcal{M}}$) adds a $\{t_k\}$ -independent term to \mathcal{G} i.e. it corresponds to a transformation $\mathcal{G} \rightarrow \mathcal{G} + f(\{C, L\})$ for some local function f on \mathcal{M} .

The dependence of \mathcal{G} on $\{t_j\}$ is, however, completely fixed by (1.19). Namely, locally one can write (1.19) in the coordinate system where $\{t_j\}_{j=1}^N$ and $\{C_j, L_j\}_{j=1}^N$ are considered as independent variables.

Then derivatives of G_j on $\{t_k\}$ for constant $\{C_j, L_j\}_{j=1}^N$ i.e. for constant monodromy data, are given by Schlesinger equations of isomonodromic deformations in G -variables:

$$\frac{\partial G_k}{\partial t_j} = \frac{A_j G_k}{t_j - t_k}, \quad j \neq k; \quad \frac{\partial G_k}{\partial t_k} = - \sum_{k \neq j} \frac{A_j G_k}{t_j - t_k}. \quad (1.20)$$

The equations (1.20) imply (1.18) *but not viceversa*. In this paper we obtain the following Hamiltonian formulation of equations (1.20):

Theorem 3 (see Theorem 2.2). *The equations (1.20) are Hamiltonian,*

$$\frac{\partial G_k}{\partial t_j} = \{H_j, G_k\} \quad (1.21)$$

where $\{.,.\}$ is the quadratic Poisson bracket (1.10) and the Hamiltonians are given by (1.17).

A direct computation shows that in (t_j, C_j, L_j) coordinates the part of $\tilde{\theta}_A$ containing dt_j 's is given by $2 \sum_{j=1}^N H_j dt_j$; together with (1.19) this implies

$$\frac{\partial \mathcal{G}}{\partial t_j} = H_j. \quad (1.22)$$

Therefore, we get the following theorem:

Theorem 4 (see Thm.3.3). *For any choice of symplectic potential $\theta_{\mathcal{M}}$ on \mathcal{M} the dependence of the generating function \mathcal{G} (1.19) on $\{t_j\}_{j=1}^N$ coincides with the t_j -dependence of the isomonodromic Jimbo-Miwa tau-function. In other words, $e^{-\mathcal{G}} \tau_{JM}$ depends only on monodromy data $\{C_j, L_j\}_{j=1}^N$.*

The above theorem shows that the generating function \mathcal{G} can be used to define the tau-function not only as a function of positions of singularities of the Fuchsian differential equation but also as a function of monodromy matrices. The ambiguity built into this definition corresponds to the freedom to choose different symplectic potentials on different open sets of the monodromy manifold.

The symplectic potential we use in this paper was found in [8] using the coordinates introduced by Fock and Goncharov in [21] (for $SL(2, \mathbb{R})$ case these coordinates called *shear coordinates* are attributed to Thurston, see [20, 16]; see also [14] where the complex analogs of the shear coordinates were used for the explicit parametrization of the open subset of full dimension of the $SL(2, \mathbb{C})$ character variety of four-punctured sphere).

Definition 1.2. *The $SL(n)$ tau-function τ on $\tilde{\mathcal{M}}$ is locally defined by the following set of compatible equations. The equations with respect to t_j are given by the formulas*

$$\frac{\partial \log \tau}{\partial t_j} = \frac{1}{2} \operatorname{res}_{z=t_j} \operatorname{tr} A^2(z), \quad (1.23)$$

where

$$A(z) := \sum_{i=1}^N \frac{A_i}{z - t_i}. \quad (1.24)$$

The equations with respect to coordinates on monodromy manifold \mathcal{M} are given by

$$d_{\mathcal{M}} \log \tau = \sum_{j=1}^N \operatorname{tr}(L_j G_j^{-1} d_{\mathcal{M}} G_j) - \theta_{\mathcal{M}}[\Sigma_0] \quad (1.25)$$

where $\theta_{\mathcal{M}}[\Sigma_0]$ is a symplectic potential (5.34) for the form $\omega_{\mathcal{M}}$ defined using the Fock-Goncharov coordinates corresponding to a ciliated triangulation Σ_0 (see Section 5.4).

Explicit formulas for derivatives of τ with respect to Fock-Goncharov coordinates will be given in Section 6.

Conjecture by A.Its, O.Lisovyy and A.Prokhorov. Theorem 2 emphasizes a close relationship with the recent work [36] where the issue of dependence of the Jimbo-Miwa tau-function on monodromy matrices was also addressed. In particular, the relevance of the Goldman bracket and the corresponding symplectic form on its symplectic leaves was observed in [36] in the case of 2×2 system with four simple poles (the associate isomonodromic deformations give Painlevé 6 equation).

Moreover, the authors of [36] introduced a form which we denote by Θ_{ILP} (this form is denoted by ω in (2.7) of [36]). This form appeared in [36] as a result of computation involving the 1-form introduced by Malgrange in [44], similarly to this work, which in our notations is given by

$$\Theta_{ILP} = \sum_{j < k}^N \text{tr} A_j A_k d \log(t_j - t_k) + \sum_{j=1}^N \text{tr}(L_j G_j^{-1} d_{\mathcal{M}} G_j) \quad (1.26)$$

where $d_{\mathcal{M}}$ denotes the differential with respect to monodromy data. Proposition 2.3 of [36] shows that the external derivative of the form (1.26) is a closed 2-form independent of $\{t_j\}_{j=1}^N$. Furthermore, in Section 1.6 the authors of [36] formulate the following

Conjecture 1. [Its-Lisovyy-Prokhorov] *The form $d\Theta_{ILP}$ coincides with the natural symplectic form on the monodromy manifold.*

There are two natural versions of this conjecture:

- *The "weak" ILP conjecture.* In this version $d_{\mathcal{M}}$ means the differential on a symplectic leaf $\{\Lambda_j = \text{const}\}_{j=1}^N$ of the $SL(n)$ character variety of $\pi_1(\mathbb{P}^1 \setminus \{t_j\}_{j=1}^N)$ (we denote this symplectic leaf by \mathcal{M}_{Λ}). The canonical symplectic form on \mathcal{M}_{Λ} is given by the inversion of the $SL(n)$ Goldman's bracket [30] and can be written explicitly in terms of monodromy data as shown in ([2], formula (3.14) for $g = 0$ and $k = 2\pi$).

By "weak" ILP conjecture we understand the coincidence of $d\Theta_{ILP}$ (1.26) with the Goldman's symplectic form on the symplectic leaves.

The problem with this formulation is that the choice of matrices G_j should be such that they satisfy the Schlesinger equations (1.20); this requirement is not natural from the symplectic point of view.

- *The "strong" ILP conjecture.* In this version the differential $d_{\mathcal{M}}$ in (1.26) means the differential on the full space \mathcal{M} (1.6) which contains both the eigenvalues of the monodromy matrices and the connection matrices. Then (omitting the pullbacks) the *strong ILP conjecture* states that

$$d\Theta_{ILP} = \omega_{\mathcal{M}}. \quad (1.27)$$

The weak version of the ILP conjecture can be derived from known results of [33, 3] or [42], as shown in Section 4.

The strong version of the ILP conjecture is equivalent to our Theorem 2. To see this equivalence it is sufficient to write (1.19) in coordinates which are split into "times" $\{t_j\}$ and some coordinates on the monodromy manifold \mathcal{M} . Then the "t-part" of the form $\tilde{\theta}_{\mathcal{A}}$ is given by $2 \sum_{k=1}^N H_k dt_k$ (this follows from the isomonodromic equations (1.20) for $\{G_j\}$) and the monodromy part coincides with the second term of the form (1.26) where the differential $d_{\mathcal{M}}$ is understood as the differential on \mathcal{M} . Now, taking the external derivative of (1.19) we come to (1.16) where the right-hand side coincides with the form $d\Theta_{ILP}$ of [36]. Finally, we notice that the formula (1.19) allows to interpret the generating function \mathcal{G} as the action of the multi-time hamiltonian system, according to Conjecture 2 of [34] (see also [35]).

Summarizing, the main results of this paper are the following:

1. We give a new hamiltonian formulation of Schlesinger system written in terms of (G, L) -variables; this formulation involves a quadratic Poisson structure defined by the dynamical r -matrix (Sec.2).
2. We prove that the monodromy map for a Fuchsian system is a symplectomorphism between (G, L) and (C, Λ) spaces (Sec.3).
3. We prove the "weak" (Sec.4) and "strong" (Thm.3.2) versions of the Its-Lisovyy-Prokhorov conjecture about coincidence of the external derivative of the Malgrange form with the natural symplectic form on the monodromy manifold
4. We introduce defining equations for the Jimbo-Miwa-Ueno tau-function with respect to Fock-Goncharov coordinates on the monodromy manifold (Def.6.1 and formula (6.5)).
5. In the $SL(2)$ case we derive equations which define the monodromy dependence of Ψ , G_j and tau-functions (Thm.6.1, Cor.6.1 and Prop.6.3).

2 Dynamical r -matrix formulation of the Schlesinger system

In this section we describe the Hamiltonian formulation of Schlesinger equations. We start from considering the $GL(n)$ case and then indicate the modifications required in the $SL(n)$ case.

2.1 Quadratic Poisson bracket via dynamical r -matrix

Let us introduce the space

$$\mathcal{H} := GL(n, \mathbb{C}) \times \mathfrak{h}_{ss} \quad (2.1)$$

where \mathfrak{h}_{ss} is the space of diagonal matrices with distinct eigenvalues. We denote an element of \mathcal{H} by (G, L) where $G \in GL(n)$ and $L \in \mathfrak{h}_{ss}$.

Proposition 2.1. *Consider the following one-form on \mathcal{H} :*

$$\theta := \text{tr}(LG^{-1} dG). \quad (2.2)$$

Then the 2-form $\omega = d\theta$ given by

$$\omega = \text{tr}(dL \wedge G^{-1} dG) - \text{tr}(LG^{-1} dG \wedge G^{-1} dG) \quad (2.3)$$

is symplectic on \mathcal{H} .

Proof. The form ω is obviously closed; to verify its non-degeneracy we consider two tangent vectors in $T_{(G,L)}\mathcal{H}$ and represent them as $(X_i, D_i) \in \mathfrak{gl}(n) \oplus \mathfrak{h}$ ($i = 1, 2$) where \mathfrak{h} denotes the Cartan subalgebra of $\mathfrak{gl}(n)$. Then

$$\omega((X_1, D_1), (X_2, D_2)) = \text{tr}\left(D_1 X_2 - D_2 X_1 - L[X_1, X_2]\right) = \text{tr}\left(D_1 X_2 - (D_2 + [X_2, L])X_1\right). \quad (2.4)$$

Suppose that ω is degenerate i.e. the vector (X_2, D_2) can be chosen so that (2.4) vanishes identically for all (X_1, D_1) . Then, choosing $D_1 = 0$, we have $\text{tr}((D_2 + [X_2, L])X_1) = 0$. Then, since X_1 is arbitrary, we have $D_2 + [X_2, L] = 0$; since L is diagonal, the commutator is diagonal-free and hence $D_2 = 0$; since L is semisimple (the eigenvalues are distinct), it follows that X_2 must be diagonal.

Then, choosing $X_1 = 0$ and D_1 arbitrary we see that the diagonal part of X_2 must vanish as well. Thus the pairing is nondegenerate and the form ω (2.3) is symplectic. \blacksquare

The corresponding Poisson structure is given by the following proposition.

Proposition 2.2. *The nonzero Poisson brackets corresponding to the symplectic form ω are*

$$\{G_{bj}, G_{c\ell}\} = \frac{G_{bt}G_{cj}}{\lambda_j - \lambda_\ell}, \quad j \neq \ell, \quad \{G_{bk}, \lambda_\ell\} = -G_{bk}\delta_{\ell k}. \quad (2.5)$$

Proof. The form (2.4) defines a map $\Phi_{(G,L)} : T_{(G,L)}\mathcal{H} \rightarrow T_{(G,L)}^*\mathcal{H}$ given by

$$\langle \Phi_{(G,L)}(X_1, D_1), (X_2, D_2) \rangle := \omega((X_1, D_1), (X_2, D_2)) \quad (2.6)$$

for all (X_2, D_2) . Then (2.4) implies

$$\Phi_{(G,L)}(X, D) = \left(-D - [X, L], X^D \right) \in T_{(G,L)}^*\mathcal{H} \quad (2.7)$$

where X^D and X^{OD} denote the diagonal and off-diagonal parts of the matrix X , respectively and the identification between a matrix and its dual is defined by the trace pairing. We denote

$$Q = -D - [X, L], \quad \delta = X^D. \quad (2.8)$$

Given now $(Q, \delta) \in T_{(G,L)}^*\mathcal{H}$ we observe from (2.8) that $D = -Q^D$ and $X = \delta + \text{ad}_L^{-1}(Q^{OD})$. The inverse of $\text{ad}_L(\cdot) = [L, \cdot]$ is given explicitly by

$$\text{ad}_L^{-1}(M)_{ab} = \frac{M_{ab}}{L_{aa} - L_{bb}}, \quad a \neq b \quad (2.9)$$

as a linear invertible map on the space of off-diagonal matrices.

Thus $\Phi_{(G,L)}^{-1} : T_{(G,L)}^* \mathcal{H} \rightarrow T_{(G,L)} \mathcal{H}$ is given by

$$\Phi_{(G,L)}^{-1}(Q, \delta) = \left(\delta + \text{ad}_L^{-1}(Q^{OD}), -Q^D \right) \quad (2.10)$$

where Q^{OD} and Q^D denote the off-diagonal and diagonal parts, respectively. The Poisson tensor $\mathbb{P} \in \wedge^2 T_{(G,L)} \mathcal{H} \simeq (\wedge^2 T_{(G,L)}^* \mathcal{H})^\vee$ is defined by

$$\mathbb{P}_{(G,L)} \left((Q_1, \delta_1), (Q_2, \delta_2) \right) := \omega \left(\Phi_{(G,L)}^{-1}(Q_1, \delta_1), \Phi_{(G,L)}^{-1}(Q_2, \delta_2) \right). \quad (2.11)$$

Using the definition (2.4), (2.6) we get

$$\begin{aligned} & \mathbb{P}_{(G,L)} \left((Q_1, \delta_1), (Q_2, \delta_2) \right) \\ = & \text{tr} \left(-Q_1^D \left(\delta_2 + \text{ad}_L^{-1}(Q_2^{OD}) \right) + Q_2^D \left(\delta_1 + \text{ad}_L^{-1}(Q_1^{OD}) \right) + \text{ad}_L \left(\delta_1 + \text{ad}_L^{-1}(Q_1^{OD}) \right) \left(\delta_2 + \text{ad}_L^{-1}(Q_2^{OD}) \right) \right) \end{aligned}$$

which is equal to

$$\text{tr} \left(Q_2^D \delta_1 - Q_1^D \delta_2 + Q_1^{OD} \text{ad}_L^{-1}(Q_2^{OD}) \right).$$

To obtain the Poisson bracket between the matrix entries of G and L we now write $Q = G^{-1} dG$ and $\delta = dL = \text{diag}(d\lambda_1, \dots, d\lambda_n)$.

Choosing $Q_1 = \mathbb{E}_{jk}, \delta_1 = 0$ and $Q_2 = 0, \delta_2 = \mathbb{E}_{\ell\ell}$ we have

$$(G^{-1})_{jb} \{G_{bk}, \lambda_\ell\} = \mathbb{P}((G^{-1} dG)_{jk}, d\lambda_\ell) = -\delta_{jk} \delta_{\ell k} \Rightarrow \{G_{bk}, \lambda_\ell\} = -G_{bk} \delta_{\ell k}.$$

Choosing $Q_1 = \mathbb{E}_{ij}, Q_2 = \mathbb{E}_{k\ell}, \delta_1 = \delta_2 = 0$ we have

$$\mathbb{P}_{(G,L)}((G^{-1} dG)_{ij}, (G^{-1} dG)_{k\ell}) = (G^{-1})_{ib} (G^{-1})_{kc} \{G_{bj}, G_{c\ell}\} = \frac{\delta_{jk} \delta_{i\ell}}{\lambda_j - \lambda_\ell}.$$

■

Proposition 2.3. *Introduce the $GL(n)$ dynamical r -matrix ([17], p.4):*

$$r(L) = \sum_{i < j} \frac{E_{ij} \otimes E_{ji} - E_{ji} \otimes E_{ij}}{\lambda_i - \lambda_j}.$$

where E_{ij} is an $n \times n$ matrix whose (ij) entry equals 1 while all other entries vanish. Introduce also the matrix

$$\Omega = \Omega_{\mathfrak{gl}(n)} := \sum_{i=1}^n E_{ii} \otimes E_{ii}.$$

Then the bracket (2.5) can be written as follows:

$$\{G^1, G^2\} = -G^2 G^1 r(L), \quad (2.12)$$

$$\{G^1, L\} = -G^1 \Omega. \quad (2.13)$$

The proof is a straightforward computation. Notice that the formula (2.13) can alternatively be written as follows:

$$\{G, \lambda_j\} = -G E_{jj}. \quad (2.14)$$

The Jacobi identity involving the brackets $\{\{G^1, G^2\}, G^3\}$ implies (taking into account that $\overset{ij}{r} = -\overset{ji}{r}$) the classical dynamical Yang-Baxter equation: (see (3) of [17]).

$$\left[\overset{12}{r}, \overset{13}{r} \right] + \left[\overset{12}{r}, \overset{23}{r} \right] + \left[\overset{23}{r}, \overset{31}{r} \right] + \sum_{i=1}^n \frac{\partial \overset{12}{r}(L)}{\partial \lambda_i} E_{ii}^3 + \frac{\partial \overset{23}{r}(L)}{\partial \lambda_i} E_{ii}^1 + \frac{\partial \overset{31}{r}(L)}{\partial \lambda_i} E_{ii}^2 = 0. \quad (2.15)$$

Remark 2.1. We did not find the construction of this section in the existing literature. In the special case of the $SL(2)$ group, the Poisson algebra (2.12), (2.13) appeared in the work [1] in the context of classical Poisson geometry of $T^*SL(2)$, see formulas (2),(3) in loc.cit.

As it was mentioned to us by L.Feher, the Poisson structure (2.12), (2.13) can be obtained from the canonical Poisson structure on $T^*SL(n)$ as follows. Consider an element $(G, A) \in T^*SL(n)$ and denote by L the diagonal form of the matrix $A \in sl(n)$ (on an open part of the space where the matrix A is diagonalizable). The condition that A is diagonal i.e $A = L$ is then a constraint of the second kind, according to Dirac's classification. The computation of the Dirac bracket for the pair (G, L) starting from the canonical Poisson structure on $T^*SL(n)$ leads to the Poisson structure (2.12), (2.13), similarly to a computation given in [18].

2.1.1 Reduction to $SL(n)$

To reduce to $SL(n)$ we observe that the proof of Prop. 2.1 holds also if we assume $\text{tr}L = 0$ and $\det G = 1$. To compute the corresponding Poisson bracket we recall that inverting the restriction of a symplectic form to a symplectic submanifold is equivalent to the computation of the Dirac bracket.

Let $h_1 := \log \det G$ and $h_2 := \text{tr}L$; the Dirac bracket is then

$$\{F, H\}_D = \{F, H\} - \sum_{j=1}^2 \{F, h_j\} A_{jk} \{h_k, H\}$$

where A_{jk} is the inverse matrix to $\{h_j, h_k\}$: in our case we have

$$\{\log \det G, \text{tr}L\} = -n \quad \Rightarrow \quad A = \frac{1}{n} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}.$$

Moreover a simple computation using (2.5) shows that

$$\{G_{jk}, \det G\} = 0, \quad \{G_{jk}, \text{tr}L\} = -G_{jk}.$$

Then (we denote by $\{\}_{SL(n)}$ the Dirac bracket restricted to $\det G = 1, \text{tr}L = 0$)

$$\begin{aligned} \{G_{bj}, G_{cl}\}_{SL(n)} &= \{G_{bj}, G_{cl}\}, \\ \{\lambda_j, \lambda_k\}_{SL(n)} &= \{\lambda_j, \lambda_k\} = 0, \\ \{G_{bk}, \lambda_\ell\}_{SL(n)} &= \{G_{bk}, \lambda_\ell\} + \frac{1}{n} \{G_{bk}, \text{tr}L\} \{\log \det G, \lambda_\ell\} \\ &= -G_{bk} \delta_{\ell k} + \frac{1}{n} G_{bk} = G_{bk} \left(\frac{1}{n} - \delta_{\ell k} \right). \end{aligned} \tag{2.16}$$

Equivalently the $SL(n)$ bracket is written as

$$\{\overset{1}{G}, \overset{2}{G}\}_{SL(n)} = -\overset{1}{G} \overset{2}{G} r(L), \quad \{\overset{1}{G}, \overset{2}{L}\}_{SL(n)} = -\overset{1}{G} \Omega \tag{2.17}$$

where now the matrix Ω is given by

$$\Omega := \Omega_{sl(n)} = \sum_{j=1}^n E_{jj} \otimes E_{jj} - \frac{1}{n} \mathbf{1} \otimes \mathbf{1} = \sum_{j,k=1}^{n-1} (\mathbb{A}^{-1})_{jk} \alpha_j \otimes \alpha_k$$

and $\alpha_j = \text{diag}(0, \dots, 1, -1, 0, \dots)$ are the simple roots of $SL(n)$ and \mathbb{A} is the Cartan matrix of $SL(n)$;

$$\mathbb{A} = \begin{pmatrix} 2 & -1 & 0 & \dots & & \\ -1 & 2 & -1 & 0 & \dots & \\ 0 & -1 & 2 & -1 & \dots & \\ 0 & 0 & \ddots & & & \\ \dots & & & -1 & 2 & \end{pmatrix}. \tag{2.18}$$

2.1.2 Relation to the Kirillov-Kostant bracket

The Kirillov-Kostant bracket on $GL(n)$, in tensor notation, takes the form

$$\{A, A\} = [A, P] = P(A - A). \quad (2.19)$$

Here P is the permutation matrix of size $n^2 \times n^2$ given by

$$P = \sum_{i,j=1}^n E_{ij} \otimes E_{ij}. \quad (2.20)$$

The regular symplectic leaves are the (co)adjoint orbits of diagonal matrices L with distinct eigenvalues, and on the orbit passing through L the symplectic form of the Kirillov-Kostant bracket (2.19) is equal to (see [5], pp. 44, 45):

$$\omega_{KK} = -\text{tr}(LG^{-1}dG \wedge G^{-1}dG) \quad (2.21)$$

where G is any matrix diagonalizing A i.e. $A = GLG^{-1}$. The form ω_{KK} is invariant under the transformation $G \rightarrow GD$ where D is a diagonal matrix which may depend on G ; such transformation leaves A invariant.

Theorem 2.1. *The map $(G, L) \mapsto A = GLG^{-1}$ is a Poisson morphism between the Poisson structure (2.5) and the Kirillov-Kostant Poisson structure on A ;*

$$\{\text{tr}(AF), \text{tr}(AH)\}_{KK} = \text{tr}\left(A[H, F]\right), \quad \forall F, H \in \mathfrak{gl}_n \simeq \mathfrak{gl}_n^\vee \quad (2.22)$$

or, equivalently,

$$\{A, A\} = [A, P]. \quad (2.23)$$

Proof. We have

$$dA = [dGG^{-1}, A] + G dLG^{-1} = G\left([X, L] + \Lambda\right)G^{-1}.$$

Then

$$\begin{aligned} \{(GLG^{-1})_{ab}, (GLG^{-1})_{cd}\} &= \mathbb{P}_{(G,L)}\left(d(GLG^{-1})_{ab}, d(GLG^{-1})_{cd}\right) \\ &= \sum_{i,j,k,\ell=1}^n G_{ai}(G^{-1})_{jb}G_{ck}(G^{-1})_{\ell d}\mathbb{P}_{(G,L)}\left(\left([G^{-1}dG, L] + dL\right)_{ij}, \left([G^{-1}dG, L] + dL\right)_{k\ell}\right). \end{aligned} \quad (2.24)$$

From the Poisson bracket (2.14) we have

$$\begin{aligned} \mathbb{P}\left((G^{-1}dG)_{jk}, d\lambda_\ell\right) &= -\delta_{jk}\delta_{\ell k}\mathbb{P}\left((G^{-1}dG)_{ij}, (G^{-1}dG)_{k\ell}\right) = \frac{\delta_{jk}\delta_{\ell i}}{\lambda_j - \lambda_\ell}, \quad j \neq \ell, \\ \mathbb{P}(d\lambda_j, d\lambda_\ell) &= 0. \end{aligned} \quad (2.25)$$

Plugging (2.25) in (2.24) we see that the only terms giving non-trivial contributions are the following:

$$\begin{aligned} &\sum_{i,j,k,\ell=1}^n G_{ai}(G^{-1})_{jb}G_{ck}(G^{-1})_{\ell d}\mathbb{P}_{(G,L)}\left(\left([G^{-1}dG, L] + dL\right)_{ij}, \left([G^{-1}dG, L] + dL\right)_{k\ell}\right) \\ &= \sum_{i,j,k,\ell=1}^n G_{ai}(G^{-1})_{jb}G_{ck}(G^{-1})_{\ell d}(\lambda_i - \lambda_j)(\lambda_k - \lambda_\ell)\mathbb{P}\left((G^{-1}dG)_{ij}, (G^{-1}dG)_{k\ell}\right) \\ &= \sum_{j,\ell=1}^n G_{a\ell}(G^{-1})_{jb}G_{cj}(G^{-1})_{\ell d}(\lambda_\ell - \lambda_j) = A_{ad}\delta_{bc} - A_{cb}\delta_{ad}. \end{aligned}$$

This expression coincides with the Kirillov-Kostant Poisson bracket (2.22). ■

A slight modification of this computation shows that the quadratic Poisson bracket (2.5) implies the Kirillov-Kostant bracket in $SL(n)$ case.

2.2 Hamiltonian formulation of the Schlesinger system in G -variables

Consider the Schlesinger system written in terms of the matrices $G_j \in SL(n)$:

$$\frac{\partial G_k}{\partial t_j} = \frac{A_j G_k}{t_j - t_k}, \quad j \neq k; \quad \frac{\partial G_k}{\partial t_k} = - \sum_{k \neq j} \frac{A_j G_k}{t_j - t_k}, \quad \frac{\partial L_k}{\partial t_j} = 0 \quad (2.26)$$

where

$$A_j = G_j L_j G_j^{-1}. \quad (2.27)$$

Matrices $L_j \in \mathfrak{sl}(n)$ are diagonal and the eigenvalues of L_j are assumed to be distinct.

The Poisson structure of the Schlesinger system (1.18) written in terms of A_j is known to be linear: it is based on the Kirillov-Kostant bracket. On the other hand, the hamiltonian formulation of the system (2.26) involves the quadratic bracket defined by the dynamical r -matrix.

The following theorem can be checked by direct calculation:

Theorem 2.2. *Denote by \mathcal{A}_0 the space of matrices $\{(G_j, L_j)\}_{j=1}^N$ where L_j are diagonal matrices with distinct eigenvalues. Then the system (2.26) is a multi-time hamiltonian system with respect to the Poisson structure on \mathcal{A}_0*

$$\{G_j^1, G_k^2\} = -G_j^1 G_k^2 r(L_k) \delta_{jk}, \quad \{G_j^1, L_k^2\} = -G_k^1 \Omega \delta_{jk}. \quad (2.28)$$

where δ_{jk} is the Kronecker delta. The Hamiltonian defining the evolution with respect to "time" t_k is given by

$$H_k = \sum_{j \neq k} \frac{\text{tr} A_j A_k}{t_k - t_j}, \quad k = 1, \dots, N.$$

We notice that for the Schlesinger system for matrices A_j (1.18) the Hamiltonians H_j are the same as for the system (2.26).

2.3 Symplectic form and potential

In the sequel we shall use the symplectic form associated to the bracket (2.28). A direct computation using the Poisson bracket (2.16) shows that the matrix $A = GLG^{-1}$ has the following Poisson brackets with G and L :

$$\{A_{ab}, G_{jk}\} = G_{ak} \delta_{bj} - G_{jk} \frac{\delta_{ab}}{n}, \quad \{A_{ab}, \lambda_k\} = 0. \quad (2.29)$$

Thus $\{\text{tr}(XA), G\} = XG$ for any fixed matrix $X \in \mathfrak{sl}(n)$ and, therefore, the matrix $A = GLG^{-1}$ is the moment map for the group action $G \mapsto SG$ on the space \mathcal{H} . A similar statement, of course, holds for $GL(n)$ using the Poisson bracket (2.5) instead.

Consider now the diagonal group action on \mathcal{A}_0 given by

$$\{G_j, L_j\}_{j=1}^N \rightarrow \{SG_j, L_j\}_{j=1}^N \quad (2.30)$$

where S is an $SL(n)$ matrix. The previous computation shows immediately that the moment map corresponding to the group action $G_j \rightarrow SG_j$ on \mathcal{A}_0 is given by

$$\{G_j, L_j\}_{j=1}^N \rightarrow \mathfrak{m} = \sum_{j=1}^N G_j L_j G_j^{-1}. \quad (2.31)$$

The space \mathcal{A} is defined by (1.6) as the space of the orbits of the action (2.30) of in the zero level set of the moment map (2.31). This implies the following theorem proven via the standard symplectic reduction [4]:

Theorem 2.3. *The Poisson structure induced on \mathcal{A} from the Poisson structure (1.12) on \mathcal{A}_0 via the reduction on the level set $\sum_{j=1}^N G_j L_j G_j^{-1} = 0$ of the moment map, corresponding to the group action $G_j \rightarrow SG_j$, is non-degenerate and the corresponding symplectic form is given by*

$$\omega_{\mathcal{A}} = - \sum_{k=1}^N \text{tr}(L_k G_k^{-1} dG_k \wedge G_k^{-1} dG_k) + \text{tr}(dL_k \wedge G_k^{-1} dG_k). \quad (2.32)$$

A symplectic potential $\theta_{\mathcal{A}}$ for $\omega_{\mathcal{A}}$ is given by

$$\theta_{\mathcal{A}} = \sum_{k=1}^N \text{tr}(L_k G_k^{-1} dG_k) . \quad (2.33)$$

3 Monodromy symplectomorphism via Malgrange's form

We start from introducing the Malgrange form associated to a Riemann-Hilbert problem on an oriented graph and discussing some of its properties, following [44, 6, 7]. From now on we work with the $SL(n)$ case.

Let Σ be an embedded graph on \mathbb{CP}^1 whose edges are smooth oriented arcs meeting transversally at the vertices. We denote by \mathbf{V} the set of vertices of Σ . Consider a "jump matrix" i.e. a function $J(z) : \Sigma \setminus \mathbf{V} \rightarrow SL(n)$ that satisfies the following properties

- Assumption 3.1.** 1. In a small neighbourhood of each point $z_0 \in \Sigma \setminus \mathbf{V}$ the matrix $J(z)$ is given by a germ of analytic function;
2. for each $v \in \mathbf{V}$, denote by $\gamma_1, \dots, \gamma_{n_v}$ the edges incident at v in a small disk centered thereof. Suppose first that all these edges are oriented away from v and enumerated in counter-clockwise order. Denote by $J_j^{(v)}(z)$ the analytic restrictions of J to γ_j . Assume that each $J_j^{(v)}(z)$ admits an analytic extension to a full neighbourhood of v and that these extensions satisfy the local non-monodromy condition

$$J_1^{(v)}(z) \cdots J_{n_v}^{(v)}(z) = \mathbf{1} . \quad (3.1)$$

If the edge γ_j is oriented towards v then $J_j^{(v)}(z)$ is taken to be the inverse of $J(z)$.

Suppose now that the jump matrices form an analytic family depending on some deformation parameters and satisfying Assumption 3.1, and consider a family of Riemann-Hilbert problems on Σ .

Malgrange form for an arbitrary Riemann Hilbert Problem. Let $\Phi(z) : \mathbb{CP}^1 \setminus \Sigma \rightarrow SL(n)$ be a matrix-valued function, bounded everywhere and analytic on each face of Σ . We also assume that the boundary values on the two sides of each edge of Σ are related by

$$\Phi_+(z) = \Phi_-(z)J(z) , \quad \forall z \in \Sigma \setminus \mathbf{V} , \quad \Phi(\infty) = \mathbf{1} . \quad (3.2)$$

where the $+/-$ boundary value is from the left/right, respectively, of the oriented edge.

Definition 3.1. [44] The Malgrange 1-form on the deformation space of Riemann-Hilbert problems with given graph Σ and jump matrices J is defined by

$$\Theta[\Sigma, J] = \frac{1}{2i\pi} \int_{\Sigma} \text{tr} \left(\Phi_-^{-1} \frac{d\Phi_-}{dz} dJ(z) J^{-1}(z) \right) dz \quad (3.3)$$

where dJ denotes the total differential of J in the space of deformation parameters for fixed z .

In ([7], Thm. 2.1) it was proved the following formula for the exterior derivative of (3.10):

$$d\Theta[\Sigma, J] = -\frac{1}{2} \int_{\Sigma} \frac{dz}{2i\pi} \text{tr} \left(\frac{d}{dz} (dJ J^{-1}) \wedge (dJ J^{-1}) \right) + \eta_{\mathbf{V}} \quad (3.4)$$

where

$$\eta_{\mathbf{V}} = -\frac{1}{4i\pi} \sum_{v \in \mathbf{V}} \sum_{\ell=1}^{n_v-1} \text{tr} \left((J_{\ell}^{(v)})^{-1} dJ_{\ell}^{(v)} \wedge dJ_{[\ell+1:n_v]}^{(v)} (J_{[\ell+1:n_v]}^{(v)})^{-1} \right) . \quad (3.5)$$

Here the notation $J_{[a:b]}^{(v)}$ stands for the product $J_a^{(v)} \cdots J_b^{(v)}$ for any two indices $a < b$.

In [7] the formula for $\eta_{\mathbf{V}}$ is written in a slightly different form and can be recast as the above expression by using the conditions (3.1).

Malgrange form and Schlesinger systems. Let us now discuss how the form (3.3) can be used in the context of the Fuchsian equation (1.1) and the associated Riemann-Hilbert problem.

Let \mathbb{D}_j be small, pairwise non-intersecting disks centered at t_j , $j = 1, \dots, N$.

In order to define the inverse monodromy map unambiguously, we need to fix the determination of the power in (1.2). To this end, fix a point β_j on the boundary of each of the disks \mathbb{D}_j and declare that, within the disk \mathbb{D}_j , the power $(z - t_j)^{L_j}$ stands for $|z - t_j|^{L_j} e^{i \arg(z - t_j) L_j}$, where the argument is chosen between $\arg(\beta_j - t_j)$ and $\arg(\beta_j - t_j) + 2\pi$. In particular the logarithm $\ln(z - t_j)$ is assumed to have the branch cut connecting t_j with β_j and the determination implied by the above.

Choose now a collection of non-intersecting edges l_1, \dots, l_N connecting ∞ with each of the β_j 's (l_j is assumed to be transversal to the boundary $\partial\mathbb{D}_j$ at β_j). Denote by Σ the union of all the circles $\partial\mathbb{D}_j$ and the edges l_j . Denote by \mathcal{D}_∞ the "exterior" domain which is the complement of the union of the disks \mathbb{D}_j and the graph Σ .

The solution $\Psi(z)$ of (1.1) is a single-valued matrix function in \mathcal{D}_∞ normalized by $\lim_{z \rightarrow \infty} \Psi(z) = \mathbf{1}$ where the direction lies within a sector lying between edges l_1 and l_N . Within each disk, with the above choice of determination of the logarithm, the analytic continuation of Ψ has the local expression (1.2). The "connection matrices" C_j are uniquely determined by a choice of G_j and the determination of the logarithm.

We have therefore defined the (extended) *monodromy map*

$$\left\{ (G_j, L_j) : \sum_{j=1}^N G_j L_j G_j^{-1} = 0 \right\} \rightarrow \left\{ (C_j, L_j) : \prod_{j=1}^N C_j e^{2i\pi L_j} C_j^{-1} = \mathbf{1} \right\}. \quad (3.6)$$

Although this monodromy map depends on Σ and the determinations of the logarithms, we are not going to indicate it explicitly.




Figure 2: Graph Σ and jump matrices on its edges used in the calculation of the form Θ .

An example of the graph Σ is shown in Fig.2; the graph looks like N "cherries" whose "stems" are attached to the point $z = \infty$. Introduce the piecewise analytic matrix on its faces as follows

$$\Phi(z) = \begin{cases} \Psi(z), & z \in \mathbb{D} = \mathbb{CP}^1 \setminus \Sigma \setminus \bigcup_{j=1}^N \mathbb{D}_j; \\ \Phi_j(z) := \Psi(z) C_j (z - t_j)^{-L_j}, & z \in \mathbb{D}_j. \end{cases} \quad (3.7)$$

The function Φ solves a Riemann-Hilbert Problem on Σ with the jump matrices on its edges indicated in Fig.2:

$$J = \begin{cases} M_j = C_j e^{2i\pi L_j} C_j^{-1}, & z \in l_j; \\ C_j (z - t_j)^{-L_j}, & z \in \partial\mathbb{D}_j \end{cases} \quad (3.8)$$

where l_j is the "stem" of the j th cherry.

The matrix function Φ given by (3.7) is the unique solution of the Riemann–Hilbert problem with jump matrices (3.8):

$$\Phi_+(z) = \Phi_-(z)J(z), \quad \Phi(\infty) = \mathbf{1}, \quad (3.9)$$

The solution of the Riemann–Hilbert problem exists for generic set of data $\{C_j, L_j, t_j\}$; this solution provides the inverse of the map in (3.6). We emphasize that the inverse monodromy map depends on the isotopy class of Σ and on the fixing of the branches of the logarithms.

In the context of Fuchsian systems the general Malgrange form in Def. 3.1 specializes to the following definition:

Definition 3.2. *The Malgrange one form $\Theta \in T_p^* \widetilde{\mathcal{M}}$ is the form defined by the expression (3.3), where Φ is the solution of the Riemann–Hilbert problem (3.8), (3.9).*

It is known [44] that the form Θ is a meromorphic form on $\widetilde{\mathcal{M}}$; the set of poles of Θ is called the ”Malgrange divisor”; on this divisor the Riemann–Hilbert problem fails to have a solution. Moreover, the residue along this divisor is a *positive* integer [44].

The deformation parameters involved in the expression (3.3) for Θ are C_j, L_j subject to the monodromy relation $\prod_{j=1}^N C_j e^{2i\pi L_j} C_j^{-1} = \mathbf{1}$, and the locations of the poles t_1, \dots, t_N .

Theorem 3.1. *The form $\Theta \in T^* \widetilde{\mathcal{M}}$ (3.3) and the potential $\tilde{\theta}_{\mathcal{A}} \in T^* \widetilde{\mathcal{A}}$ are related by*

$$\Theta = (\tilde{\mathcal{F}}^{-1})^* \left(\theta_{\mathcal{A}} - \sum_{j=1}^N H_j dt_j \right) \quad (3.10)$$

where H_j are the Hamiltonians (1.17). Denote now by $\partial_{t_j} \in T\widetilde{\mathcal{M}}$ the vector field of differentiation w.r.t. t_j keeping the monodromy data constant. Then the contraction of Θ with ∂_{t_j} is given by

$$\Theta(\partial_{t_j}) = H_j. \quad (3.11)$$

Equivalently, the contraction of (the pullback via the inverse monodromy map of) $\tilde{\theta}_{\mathcal{A}}$ with ∂_{t_j} equals $2H_j$.

Proof. The simplest way to prove (3.10) is via the *localization formula* [36] using the Riemann–Hilbert problem defined on the graph Σ shown in Fig.2. To simplify the notation we will not indicate explicitly the pullbacks, but simply consider the matrices G_j as functions of times and monodromy data via the inverse monodromy map.

In the formula (3.3) the function Φ_- coincides with the boundary value of the solution, Ψ , of the ODE (1.1) in the domain \mathbb{D} . Therefore, denoting $d\Phi/dz$ by Φ' we have:

$$\text{tr}(\Phi_-^{-1} \Phi'_- dJJ^{-1}) = \text{tr}(A(z) \Phi_- dJJ^{-1} \Phi_-^{-1}). \quad (3.12)$$

Here we have used the fact that Φ_- coincides with Ψ and therefore $\Phi'_- \Phi_-^{-1} = A(z)$. Moreover we have

$$\Phi_- dJJ^{-1} \Phi_-^{-1} = d(\Phi_- J) J^{-1} \Phi_-^{-1} - d\Phi_- \Phi_-^{-1} = d\Phi_+ \Phi_+^{-1} - d\Phi_- \Phi_-^{-1}$$

since $\Phi_+ = \Phi_- J$. Thus (3.3) can be equivalently written as follows

$$\Theta = \frac{1}{2\pi i} \int_{\Sigma} \text{tr}(A(z)(d\Phi_+ \Phi_+^{-1} - d\Phi_- \Phi_-^{-1})) dz \quad (3.13)$$

and further represented as

$$\Theta = \frac{1}{2\pi i} \int_{\partial\mathbb{D}} \text{tr}(A(z) d\Psi \Psi^{-1}) dz + \frac{1}{2\pi i} \sum_{j=1}^N \int_{\partial\mathbb{D}_j} \text{tr}(A(z) d\Phi_+ \Phi_+^{-1}) dz. \quad (3.14)$$

The first integral in the r.h.s. of (3.14) vanishes since the integrand is holomorphic in \mathbb{D} . Thus (3.14) reduces to (this is the expression that also appears in [36], formula (1.11)):

$$\Theta = \sum_j \text{res}_{z=t_j} \text{tr}(A(z) d\Phi_j(z) \Phi_j^{-1}(z)) dz. \quad (3.15)$$

The expression (3.15) can be further evaluated in the coordinate system given by (C_j, L_j, t_j) . Namely, the contribution of derivatives with respect to monodromy data (C_j, L_j) into (3.15) is obtained by evaluation of $d\Phi_j(z)\Phi_j^{-1}(z)$ at the poles t_j which gives the monodromy part of $\tilde{\theta}_{\mathcal{A}}$ in (3.10).

A straightforward local analysis using (3.7) shows that:

$$\left. \partial_{t_k} \Phi_j(z)\Phi_j^{-1}(z) \right|_{z=t_j} = \partial_{t_k} G_j G_j^{-1} - \delta_{kj} \partial_{t_k} G_k G_k^{-1} - \delta_{kj} [A_k, \Phi_j'(t_j)\Phi_j(t_j)^{-1}].$$

Thus

$$\Theta = \sum_j \operatorname{res}_{z=t_j} \operatorname{tr} \left(A(z) d\Phi_j(z)\Phi_j^{-1}(z) \right) dz = \sum_j \operatorname{tr} \left(A_j dG_j G_j^{-1} \right) - \sum_j dt_j \operatorname{tr} \left(A_j \partial_{t_j} G_j G_j^{-1} \right).$$

Finally, due to the Schlesinger equations for G_j (1.20) we get

$$\Theta = \sum_j \operatorname{tr} \left(A_j dG_j G_j^{-1} \right) - \sum_j dt_j \sum_{k \neq j} \frac{\operatorname{tr} A_j A_k}{t_j - t_k}.$$

Recalling that the Jimbo-Miwa Hamiltonians are given by $H_j = \sum_{k \neq j} \frac{\operatorname{tr} A_j A_k}{t_j - t_k}$ and that the first term equals the potential $\tilde{\theta}_{\mathcal{A}}$ on $\tilde{\mathcal{A}}$, we arrive at (3.10).

As a corollary of the Schlesinger equations (1.20) the contraction of $\tilde{\theta}_{\mathcal{A}}$ with a vector field ∂_{t_j} (for fixed monodromy data) is

$$\tilde{\theta}_{\mathcal{A}}(\partial_{t_j}) = 2H_j.$$

Therefore, the total dt_j - part of the form Θ for fixed monodromies equals to $\sum_{j=1}^N H_j dt_j$. ■

Symplectic form on the monodromy manifold. We start from defining the two-form on the monodromy manifold which is one of central objects of this paper.

Definition 3.3. Define the following 2-form on \mathcal{M} (1.7):

$$\omega_{\mathcal{M}} = \frac{1}{4\pi i} (\omega_1 + \omega_2) \quad (3.16)$$

where

$$\omega_1 = \sum_{\ell=1}^N \operatorname{tr} (M_{\ell}^{-1} dM_{\ell} \wedge K_{\ell}^{-1} dK_{\ell}) + \sum_{\ell=1}^N \operatorname{tr} (\Lambda_{\ell}^{-1} C_{\ell}^{-1} dC_{\ell} \wedge \Lambda_{\ell} C_{\ell}^{-1} dC_{\ell}), \quad (3.17)$$

$$\omega_2 = 2 \sum_{\ell=1}^N \operatorname{tr} (\Lambda_{\ell}^{-1} d\Lambda_{\ell} \wedge C_{\ell}^{-1} dC_{\ell}) \quad (3.18)$$

and $K_{\ell} = M_1 \dots M_{\ell}$.

On the monodromy manifold $M_1 \dots M_N = \mathbf{1}$ the form $\omega_{\mathcal{M}}$ is invariant under simultaneous transformation $C_j \rightarrow SC_j$ with S is an arbitrary $SL(n)$ -valued function on \mathcal{M} .

Remark 3.1. The restriction of the form $-2i\pi\omega_{\mathcal{M}}$ on the leaves $\Lambda_j = \text{constant}$ (under such restriction $\omega_2 = 0$ and hence $-2i\pi\omega_{\mathcal{M}} = -\omega_1/2$) coincides with the symplectic form on the symplectic leaves of the $GL(n)$ Goldman bracket found in [2] (formula (3.14); the case of this formula relevant for us corresponds to $k = 2\pi$ and $g = 0$ in the notation of [2]).

As we prove below in Corollary 3.1, the form $\omega_{\mathcal{M}}$ is non-degenerate on the space \mathcal{M} , which is a torus fibration (with fiber the product of N copies of the $SL(n)$ torus of diagonal matrices) over the union of all the symplectic leaves of the Goldman bracket. The fact that \mathcal{M} is a torus fibration is simply due to the fact that the fibers of the map $(C_j, \Lambda_j) \rightarrow M_j = C_j \Lambda_j C_j^{-1}$ are obtained by multiplication of the C_j 's on the right by diagonal matrices.

Let us trivially extend the form $\omega_{\mathcal{M}}$ to the space $\widetilde{\mathcal{M}}$ (1.15) which includes also the variables t_j . This extension is denoted by $\widetilde{\omega}_{\mathcal{M}}$.

Relation between forms Θ and $\omega_{\mathcal{M}}$. The following theorem was stated in [6] in slightly different notations without direct proof. The proof is given below.

Theorem 3.2. *The exterior derivative of the form Θ is given by the pullback of the form $\widetilde{\omega}_{\mathcal{M}}$ (3.16) under the monodromy map:*

$$d\Theta = [\widetilde{\mathcal{F}}]^* \widetilde{\omega}_{\mathcal{M}}. \quad (3.19)$$

Proof. Let us apply the formulas (3.4), (3.5) to the graph Σ depicted in Fig.2 with indicated jump matrices. The integral over Σ in the formula (3.4) then reduces to a sum of integrals over $\partial\mathbb{D}_\ell$'s because the jump matrix $J(z)$ on the cuts is constant with respect to z . We denote by β_ℓ the three-valent vertices where the circles around t_ℓ meet with the edges going towards z_0 . Let us consider the contribution of one of the integrals over $\partial\mathbb{D}_\ell$ to (3.4).

We will drop the index ℓ for brevity in the formulas below. Notice also that $dL \wedge dL = 0$ because the matrix L is diagonal. Letting $J(z) = C(z-t)^{-L}$ we get

$$-\frac{1}{2} \oint \frac{dz}{2i\pi} \operatorname{tr} \left(\frac{d}{dz} \left(dJ(z)J(z)^{-1} \right) \wedge dJ(z)J(z)^{-1} \right) = \frac{1}{2} \left(\frac{dt \wedge L dL}{(\beta-t)} + dL \wedge C^{-1} dC \right). \quad (3.20)$$

In the course of the computation we have used that

$$\int_{\beta}^{\beta} \frac{dz \log(z-t)}{2i\pi (z-t)^2} = -\frac{1}{\beta-t}$$

where the integration goes along the circle $|z-t| = |\beta-t|$ starting at $z = \beta$. We now turn to the evaluation of the term $\eta_{\mathbf{v}}$ (3.5). The set of vertices \mathbf{V} consists of $\mathbf{V} = \{z_0, \beta_1, \dots, \beta_N\}$. The contribution coming from the vertex z_0 is precisely the first term in ω_1 (3.17) (in (3.17) this term is simplified using the local no-monodromy condition (3.1)).

To evaluate the contribution of the vertex $\beta = \beta_\ell \in \mathbf{V}$ we observe that this vertex is tri-valent and the jump matrices on the three incident arcs are

$$J_1 = C\Lambda^{-1}C^{-1}, \quad J_2 = C(\beta-t)^{-L}, \quad J_3 = (\beta-t)^L e^{2i\pi L} C^{-1}$$

where $\Lambda := e^{2i\pi L}$. In the definition it is assumed that $(z-t)^L$ is defined with a branch cut extending from t to β . Since $J_1 J_2 J_3 = \mathbf{1}$ the contribution of the vertex to (3.5) reduces to the term

$$\frac{-1}{4i\pi} \operatorname{tr} (J_1 dJ_2 \wedge dJ_3) = \frac{-1}{4i\pi} \operatorname{tr} (J_2^{-1} dJ_2 \wedge dJ_3 J_3^{-1}).$$

Recall that L, Λ are diagonal; we have then

$$\begin{aligned} J_2^{-1} dJ_2 &= (\beta-t)^L C^{-1} dC (\beta-t)^{-L} + (\beta-t)^L \frac{L dt}{\beta-t} (\beta-t)^{-L} - dL \log(\beta-t), \\ dJ_3 J_3^{-1} &= \frac{-dtL}{(\beta-t)} + (\log(\beta-t) + 2i\pi) dL - (\beta-t)^L \Lambda C^{-1} dC \Lambda^{-1} (\beta-t)^{-L}. \end{aligned} \quad (3.21)$$

Then a straightforward computation gives

$$\begin{aligned} &\frac{-1}{4i\pi} \operatorname{tr} (J_2^{-1} dJ_2 \wedge dJ_3 J_3^{-1}) \\ &= \frac{-1}{4i\pi} \operatorname{tr} \left(C^{-1} dC \wedge \Lambda^{-1} d\Lambda - C^{-1} dC \wedge \Lambda C^{-1} dC \Lambda^{-1} + 2i\pi \frac{L dt}{\beta-t} \wedge dL \right). \end{aligned} \quad (3.22)$$

Summing up (3.20) (the contribution of the integral) with (3.22) (the contribution coming from the vertex $\beta = \beta_\ell$) we get

$$(3.20) + (3.22) = \frac{1}{4i\pi} \operatorname{tr} \left(-2C^{-1} dC \wedge \Lambda^{-1} d\Lambda + C^{-1} dC \wedge \Lambda C^{-1} dC \Lambda^{-1} \right).$$

Then summing over all contributions from vertices β_ℓ leads to (3.16).

Summarizing, the first term in (3.17) corresponds to the N -valent vertex. The second term in (3.17) together with the term (3.18) arise from the contributions of cherries and the three-valent vertices formed by cherries and their stems. \blacksquare

This theorem immediately implies the following corollary, which can also be deduced from previous results of [11].

Corollary 3.1. *The form $\omega_{\mathcal{M}}$ (3.16) is closed and non-degenerate on the monodromy manifold \mathcal{M} .*

Strong version of Its-Lisovyy-Prokhorov conjecture. The theorem 3.2 proves the "strong" version of the ILP conjecture (1.26). To state this conjecture in the present setting we consider the form (1.11) or (2.7) of [36] which we denote by Θ_{ILP} to avoid confusion with the notations of this paper (see also the identity (4.20) below):

$$\Theta_{ILP} = \sum_{j < k}^N \text{tr} A_j A_k d \log(t_j - t_k) + \sum_{j=1}^N \text{tr}(L_j G_j^{-1} d_{\mathcal{M}} G_j). \quad (3.23)$$

The Conjecture from section 1.6 of [36] refers to the restriction of the form to the symplectic leaves $L_j = \text{constants}$. We refer to this as the *weak Its-Lisovyy-Prokhorov conjecture*; in this formulation $d_{\mathcal{M}}$ refers to the differential only with respect to the connection matrices C_j . This "weak" version of the conjecture is proved on the basis of known results [33, 3, 42] in the next section.

The statement of Theorem 3.2 is the *strong* version of the above conjecture: in this version the differential $d_{\mathcal{M}}$ is with respect to all monodromy data including the L_j 's.

Generating function of the monodromy map. The closure of $\omega_{\mathcal{M}}$ guarantees the local existence of a *symplectic potential*. Denoting any such local potential by $\theta_{\mathcal{M}}$ (such that $d\theta_{\mathcal{M}} = \omega_{\mathcal{M}}$) we define the (local on $\widetilde{\mathcal{M}}$) generating function \mathcal{G} as follows

$$d\mathcal{G} = \sum_{k=1}^N \text{tr}(L_k G_k^{-1} dG_k) - \sum_{j=1}^N H_k dt_k - \widetilde{\theta}_{\mathcal{M}} \quad (3.24)$$

where G_k and H_k are considered as functions on $\widetilde{\mathcal{M}}$ under the inverse monodromy map.

The equation (3.24) can be used to extend the definition of Jimbo-Miwa tau-function to include its dependence on monodromies. Irrespectively of the choice of $\theta_{\mathcal{M}}$, the formula (3.11) implies the following theorem

Theorem 3.3. *For any choice of symplectic potential $\theta_{\mathcal{M}}$ on \mathcal{M} the dependence of the generating function \mathcal{G} (1.19) on $\{t_j\}_{j=1}^N$ coincides with t_j -dependence of the isomonodromic Jimbo-Miwa tau-function. In other words, $e^{-\mathcal{G}}_{\tau_{JM}}$ depends only on monodromy data $\{C_j, L_j\}_{j=1}^N$.*

In Section 6 we are going to use this theorem to *define* the isomonodromic tau function as exponent of the generating function \mathcal{G} under a special choice of the symplectic potential $\theta_{\mathcal{M}}$ based on the use of Fock-Goncharov coordinates.

Remark 3.2. "Extended" character varieties with non-degenerate symplectic form were considered in the '94 paper [37] and later in the paper [11]. In ([11] Corollary 1) it was proven that the pullback of a symplectic form from the extended monodromy manifold coincides with a symplectic form on (L_j, G_j) side. The description of the corresponding Poisson bracket, construction of symplectic potentials, Malgrange form, the tau-function and coordinatization in term of Fock-Goncharov parameters were not considered before, to the best of our knowledge.

4 Standard monodromy map and weak version of Its-Lisovyy-Prokhorov conjecture

Here we show that a weak version of Its-Lisovyy-Prokhorov conjecture can be derived in a simple way from previous results of [33, 3] or [42] where a symplectomorphism between the space of coefficients

$\{A_j\}$ with given set of eigenvalues of the Fuchsian equation (1.1) and a symplectic leaf of Goldman bracket was proved.

First, consider the submanifold \mathcal{A}_L of \mathcal{A} such that the diagonal form of each of the matrices A_j is fixed:

$$\mathcal{A}_L = \left\{ \{A_i\}_{i=1}^N, A_i \in \mathcal{O}(L_i), \sum_{i=1}^N A_i = 0 \right\} / \sim \quad (4.1)$$

where \sim is the equivalence over simultaneous adjoint transformation $A_i \rightarrow SA_iS^{-1}$ of all A_i for $S \in SL(n)$; $L = (L_1, \dots, L_N)$ where L_j is the diagonal form of A_j and $\mathcal{O}(L)$ is the (co)-adjoint orbit of the diagonal matrix L . We assume that diagonal entries of each L_j do not differ by an integer.

Consider similarly also the space \mathcal{M}_L which is the subspace of the $SL(n)$ character variety of $\pi_1(\mathbb{P}^1 \setminus \{t_j\}_{j=1}^N)$ such that the diagonal form of the matrix M_j equals to $\Lambda_j = e^{2\pi i L_j}$.

The Kirillov-Kostant brackets (2.19) for each A_j :

$$\{A_j^1, A_k^2\} = [A_j^1, P] \delta_{jk} \quad (4.2)$$

can be equivalently rewritten in the r -matrix form

$$\{A(z)^1, A(w)^2\} = \frac{1}{z-w} [P, A(z)^1 + A(w)^2]. \quad (4.3)$$

The Schlesinger equations for $A_j = G_j L_j G_j^{-1}$ which follow from the system (1.20) for G_j take the form:

$$\frac{\partial A_k}{\partial t_j} = \frac{[A_k, A_j]}{t_k - t_j}, \quad j \neq k; \quad \frac{\partial A_j}{\partial t_j} = - \sum_{k \neq j} \frac{[A_k, A_j]}{t_k - t_j}. \quad (4.4)$$

These equations are Hamiltonian,

$$\frac{\partial A_k}{\partial t_j} = \{H_j, A_k\},$$

with the Poisson structure given by (4.3) and the (time dependent) Hamiltonians H_j defined by (1.17). Notice that these Hamiltonians commute $\{H_k, H_j\} = 0$ and satisfy the equations $\partial_{t_k} H_j = \partial_{t_j} H_k$.

After the symplectic reduction to the space of orbits of the global $Ad_{GL(N)}$ action and restriction to the level set $\sum_{j=1}^N A_j = 0$ of the corresponding moment map one gets a degenerate Poisson structure; its symplectic leaves coincide with \mathcal{A}_L [33]. The symplectic form on \mathcal{A}_L can be written as

$$\omega_{\mathcal{A}}^L = - \sum_{k=1}^N \text{tr}(L_k G_k^{-1} dG_k \wedge G_k^{-1} dG_k). \quad (4.5)$$

The form (4.5) is independent of the choice of matrices G_j which diagonalize A_j ; moreover, it is invariant under simultaneous transformation $A_j \rightarrow SA_j S^{-1}$ and thus it is indeed defined on the space \mathcal{A}_L .

The $SL(n)$ character variety is equipped with the Poisson structure given by the Goldman bracket defined as follows (see p.266 of [31]): for any two loops $\sigma, \tilde{\sigma} \in \pi_1(\mathbb{P}^1 \setminus \{t_i\}_{i=1}^N)$ the Poisson bracket between the traces of the corresponding monodromies is given by

$$\left\{ \text{tr} M_\sigma, \text{tr} M_{\tilde{\sigma}} \right\}_G = \sum_{p \in \sigma \cap \tilde{\sigma}} \nu(p) \left(\text{tr}(M_{\sigma_p \tilde{\sigma}}) - \frac{1}{n} \text{tr} M_\sigma \text{tr} M_{\tilde{\sigma}} \right). \quad (4.6)$$

where $\nu(p) = \pm 1$ is the contribution of point p to the intersection index of σ and $\tilde{\sigma}$.

The space \mathcal{M}_L is a symplectic leaf of the $SL(n)$ Goldman bracket; the Goldman's symplectic form on \mathcal{M}_L coincides with $-\frac{1}{2}\omega_1$ [2] where ω_1 is defined in (3.17). We define

$$\omega_{\mathcal{M}}^L = \frac{1}{4\pi i} \omega_1. \quad (4.7)$$

The study of the symplectic properties of the map (1.8) was initiated in [33, 3, 42]. In [33, 3] two different proofs were given of the fact that the monodromy map \mathcal{F}^t is a symplectomorphism i.e.

$$(\mathcal{F}^t)^* \omega_{\mathcal{M}}^L = \omega_{\mathcal{A}}^L. \quad (4.8)$$

In [42] the brackets between the monodromy matrices themselves were obtained starting from (4.3); the result is given by

$$\{M_i^1, M_i^2\}^* = \pi i P(M_i^1 M_i^1 - M_i^2 M_i^2), \quad (4.9)$$

$$\{M_i^1, M_j^2\}^* = \pi i P\left(M_j^1 M_i^1 + M_i^2 M_j^2 - M_i^1 M_j^2 - M_j^1 M_i^2\right), \quad i < j \quad (4.10)$$

where P is the matrix of permutation of two spaces. The brackets (4.9), (4.10) were computed for the basepoint $z_0 = \infty$ on the level set $\sum_{j=1}^N A_j = 0$ of the moment map; thus the algebra (4.9), (4.10) does not satisfy the Jacobi identity. However, the Jacobi identity is restored for the algebra of Ad -invariant objects i.e. for traces of monodromies; moreover, for any two loops σ and $\tilde{\sigma}$ we have ([50]; see also Thm. 5.2 of [14] where this statement was proved for $n = 4, N = 2$ case):

$$\{\mathrm{tr} M_\sigma, \mathrm{tr} M_{\tilde{\sigma}}\}^* = -2\pi i \{\mathrm{tr} M_\sigma, \mathrm{tr} M_{\tilde{\sigma}}\}_G \quad (4.11)$$

which gives an alternative proof of (4.8).

Let us now show that (4.8) implies the weak version of the Its-Lisovyy-Prokhorov conjecture. Similarly to (1.14) and (1.15) we introduce the two spaces

$$\tilde{\mathcal{A}}_L = \left\{ (p, \{t_j\}_{j=1}^N), p \in \mathcal{A}_L, t_j \in \mathbb{C}, t_j \neq t_k \right\}, \quad (4.12)$$

$$\tilde{\mathcal{M}}_L = \left\{ (p, \{t_j\}_{j=1}^N), p \in \mathcal{M}_L, t_j \in \mathbb{C}, t_j \neq t_k \right\}. \quad (4.13)$$

Denote the pullback of the form $\omega_{\mathcal{A}}^L$ with respect to the natural projection of $\tilde{\mathcal{A}}_L$ to \mathcal{A}_L by $\tilde{\omega}_{\mathcal{A}}^L$ and the pullback of the form $\omega_{\mathcal{M}}$ with respect to the natural projection of $\tilde{\mathcal{M}}_L$ to \mathcal{M}_L by $\tilde{\omega}_{\mathcal{M}}^L$.

Proposition 4.1. *The following identity holds between two-forms on $\tilde{\mathcal{A}}_L$:*

$$\tilde{\mathcal{F}}^*[\tilde{\omega}_{\mathcal{M}}^L] = \tilde{\omega}_{\mathcal{A}}^L - \sum_{k=1}^N dH_k \wedge dt_k \quad (4.14)$$

where H_k are the Hamiltonians (1.17).

Proof. Denote by $2d$ the dimension of the spaces \mathcal{A}_L and \mathcal{M}_L . Introduce some local Darboux coordinates (p_i, q_i) on \mathcal{A}^L for the form $\omega_{\mathcal{A}}^L$ (4.5) and also some Darboux coordinates (P_j, Q_j) on \mathcal{M}^L for the form $\omega_{\mathcal{M}}^L$ given by (4.7).

We are going to verify (4.14) using coordinates $\{t_j\}_{j=1}^N$ and $\{P_j, Q_j\}_{j=1}^d$. Let us split the operator d into two parts:

$$d = d_t + d_{\mathcal{M}}$$

where $d_{\mathcal{M}}$ is the differential with respect to $\{P_j, Q_j\}_{j=1}^d$. Then relation (4.8) can be written as

$$\sum_{j=1}^d dP_j \wedge dQ_j = \sum_{j=1}^d d_{\mathcal{M}} p_i \wedge d_{\mathcal{M}} q_i. \quad (4.15)$$

The right-hand side can be further rewritten using the Hamilton equations $\frac{\partial p_i}{\partial t_k} = -\frac{\partial H_k}{\partial q_i}$, $\frac{\partial q_i}{\partial t_k} = \frac{\partial H_k}{\partial p_i}$ (where the Hamiltonians H_k are given by (1.17)). Using

$$d_{\mathcal{M}} p_i = dp_i + \sum_{k=1}^N \frac{\partial H_k}{\partial q_i} dt_k \quad d_{\mathcal{M}} q_i = dq_i - \sum_{k=1}^N \frac{\partial H_k}{\partial p_i} dt_k$$

one gets

$$\sum_{i=1}^d d_{\mathcal{M}} p_i \wedge d_{\mathcal{M}} q_i = \sum_{i=1}^d dp_i \wedge dq_i + \sum_{k=1}^N dt_k \wedge \sum_{i=1}^d \left(\frac{\partial H_k}{\partial q_i} dq_i + \frac{\partial H_k}{\partial p_i} dp_i \right)$$

$$- \sum_{\ell < k=1}^N \sum_{i=1}^d \left(\frac{\partial H_\ell}{\partial q_i} \frac{\partial H_k}{\partial p_i} - \frac{\partial H_\ell}{\partial p_i} \frac{\partial H_k}{\partial q_i} \right) dt_\ell \wedge dt_k. \quad (4.16)$$

To simplify the second sum in (4.16) we recall that

$$dH_k = \sum_{i=1}^d \left(\frac{\partial H_k}{\partial q_i} dq_i + \frac{\partial H_k}{\partial p_i} dp_i \right) + \sum_{\ell=1}^N \frac{\partial H_k}{\partial t_\ell} \Big|_{p_i, q_i = \text{const}} dt_\ell;$$

thus the second sum can be written as

$$\sum_{k=1}^H dt_k \wedge dH_k + \sum_{l, k, l < k} \left(\frac{\partial H_k}{\partial t_l} \Big|_{p, q} - \frac{\partial H_l}{\partial t_k} \Big|_{p, q} \right) dt_l \wedge dt_k.$$

Adding all the terms in (4.16) we obtain

$$\sum_{j=1}^d dP_j \wedge dQ_j = \sum_{i=1}^d dp_i \wedge dq_i + \sum_{k=1}^N dt_k \wedge dH_k - \sum_{\ell < k} \left(\frac{\partial H_\ell}{\partial t_k} \Big|_{p, q} - \frac{\partial H_k}{\partial t_\ell} \Big|_{p, q} + \{H_k, H_\ell\} \right) dt_\ell \wedge dt_k.$$

The coefficient of $dt_\ell \wedge dt_k$ vanishes because the Hamiltonians satisfy the zero-curvature equations implied by commutativity of the flows with respect to t_j and t_ℓ ; in fact in this particular case they satisfy a stronger compatibility: $\{H_k, H_\ell\} = 0$ and $\partial_{t_\ell} H_k = \partial_{t_k} H_\ell$. Therefore we arrive at (4.14). \blacksquare

Let us show that (4.14) implies

Proposition 4.2 (Weak ILP conjecture). *The following identity holds on the space $\widetilde{\mathcal{M}}_L$:*

$$d\Theta_{ILP}^L = \widetilde{\omega}_{\mathcal{M}}^L \quad (4.17)$$

where

$$\Theta_{ILP}^L = \sum_{j < k}^N \text{tr} A_j A_k d \log(t_j - t_k) + \sum_{j=1}^N \text{tr}(L_j G_j^{-1} d_{\mathcal{M}} G_j) \quad (4.18)$$

and matrices G_j diagonalizing A_j are chosen to satisfy the Schlesinger equations (1.20); $d_{\mathcal{M}}$ denotes the differential with respect to monodromy coordinates. The form Θ_{ILP}^L is the "weak" version of the form (1.26). The form $\widetilde{\omega}_{\mathcal{M}}^L$ is the pullback of Alekseev-Malkin form (4.7) from \mathcal{M}_L to $\widetilde{\mathcal{M}}_L$.

Proof. The symplectic potential for the form $\widetilde{\omega}_{\mathcal{A}}^L$ can be written as

$$\widetilde{\theta}_{\mathcal{A}}^L = \sum_{j=1}^n \text{tr}[L_j G_j^{-1} (d_t + d_{\mathcal{M}}) G_j]. \quad (4.19)$$

We notice that the potential $\widetilde{\theta}_{\mathcal{A}}^L$, in contrast to the form $\widetilde{\omega}_{\mathcal{A}}^L$ itself, is not well-defined on the space $\widetilde{\mathcal{A}}_L$ due to ambiguity $G_j \rightarrow G_j D_j$ for diagonal D_j in the definition of G_j . Under such transformation $\widetilde{\theta}_{\mathcal{A}}^L$ changes by an exact form. Therefore for the purpose of proving (4.17) one can pick any concrete representative for each G_j . The most natural choice is to assume that $\{G_j\}$ satisfy the system (1.20). Then the "t"-part of potential (4.19) can be computed using (1.20) and the definition of the Hamiltonians (1.17) to give

$$\sum_{j=1}^n \text{tr}(L_j G_j^{-1} d_t G_j) = 2 \sum_{j=1}^N H_j dt_j. \quad (4.20)$$

Therefore, the relation (4.14) can be rewritten as

$$\widetilde{\mathcal{F}}^*[\widetilde{\omega}_{\mathcal{M}}^L] = d \left(\sum_{k=1}^N dH_k \wedge dt_k + \sum_{j=1}^n \text{tr}(L_j G_j^{-1} d_{\mathcal{M}} G_j) \right) \quad (4.21)$$

which coincides with (4.17). \blacksquare

Comparison of weak and strong ILP conjectures. In spite of the formal similarity, there is a significant difference between the statements of the weak and strong ILP conjectures. In the strong version the form $\sum \text{tr}(L_j dG_j G_j^{-1})$ is a well-defined form on the phase space \mathcal{A} as well as on its extension $\tilde{\mathcal{A}}$.

In the weak version the same form is not defined on the space \mathcal{A}^L since to get the equality (4.17) one needs to take the residues A_j (which are given by a point of \mathcal{A}^L up to a conjugation) and then diagonalize each A_j into $G_j L_j G_j^{-1}$ in a way which is non-local in times t_j : the matrices G_j 's themselves must satisfy the Schlesinger system (1.20). This requirement can not be satisfied staying entirely within the space \mathcal{A}^L and thus G_j 's can not be chosen as functionals of A_j 's only; their choice encodes a highly non-trivial t_j -dependence which fixes the freedom in the right multiplication of each G_j by a diagonal matrix which also can be time-dependent.

The strong version of the ILP conjecture (Theorem 3.2) is a stronger statement since the form $\theta_{\mathcal{A}}$ is a 1-form defined on the underlying phase space.

5 Log-canonical coordinates and symplectic potential

Here we summarize results of [8] where the form $\omega_{\mathcal{M}}$ was expressed in log-canonical form on an open subspace of highest dimension of \mathcal{M} using the (extended) system of Fock-Goncharov coordinates [21]. This allows to find the corresponding symplectic potential and use it in the definition of the tau-function.

5.1 Fock-Goncharov coordinates

To define the Fock-Goncharov coordinates we introduce the following auxiliary graphs (see Fig. 3):

1. The graph Σ_0 with N vertices v_1, \dots, v_N which defines a triangulation of the N -punctured sphere; we assume that each vertex v_j lies in a small neighbourhood of the corresponding pole t_j . Since Σ_0 is a triangulation there are $2N - 4$ faces $\{f_k\}_{k=1}^{2N-4}$ and $3N - 6$ edges $\{e_k\}_{k=1}^{3N-6}$; the edges are assumed to be oriented.
2. Consider a small loop around each t_k (the *cherry*) and attach it to the vertex v_k by an edge (the *stem of the cherry*). The cherries are assumed to not intersect the edges of Σ_0 . The union of Σ_0 , the stems and the cherries is denoted by Σ_1 .
The graph Σ_1 is fixed by Σ_0 if one chooses the ciliation at each vertex of the graph Σ_0 ; the ciliation determines the position of the stem of the corresponding cherry.
3. Choose a point p_f inside each face f_k of Σ_0 and connect it by edges $\mathcal{E}_f^{(i)}$, $i = 1, 2, 3$ to the vertices of the face, oriented towards the point p_f . We will denote by Σ the graph obtained by the augmentation of Σ_1 and these new edges. It is the graph Σ which will be used to compute the form $\omega_{\mathcal{M}}$.

We will make use of the following notations: by α_i , $i = 1, \dots, n - 1$ we denote the simple positive roots of $SL(n)$; by \mathfrak{h}_i we denote the dual roots:

$$\alpha_i := \text{diag}(0, \dots, \overset{i\text{-pos}}{1}, -1, 0, \dots), \quad \mathfrak{h}_i := \begin{pmatrix} (n-i)\mathbf{1}_i & 0 \\ 0 & -i\mathbf{1}_{n-i} \end{pmatrix}, \quad \text{tr}(\alpha_i \mathfrak{h}_k) = n\delta_{ik}. \quad (5.1)$$

For any matrix M we define $M^* := \mathbf{P}M\mathbf{P}$ where \mathbf{P} is the ‘‘long permutation’’ in the Weyl group,

$$\mathbf{P}_{ab} = \delta_{a, n+1-b}.$$

In particular $\alpha_i^* = -\alpha_{n-i}$, $\mathfrak{h}_i^* = -\mathfrak{h}_{n-i}$. Let

$$\sigma = \text{diag}(1, -1, 1, -1, \dots)$$

be the signature matrix.

Introduce the $(n-1) \times (n-1)$ matrix \mathbb{G} given by

$$\mathbb{G}_{jk} = \text{tr}(\mathfrak{h}_j \mathfrak{h}_k) = n^2 \left(\min(j, k) - \frac{jk}{n} \right). \quad (5.2)$$

The matrix \mathbb{G} coincides with $n^2 A_{n-1}^{-1}$ with A_{n-1} being the Cartan matrix of $SL(n)$.

The full set of coordinates on \mathcal{M} consists of three groups: the coordinates assigned to vertices of the graph Σ_0 , to its edges and faces. Below we describe these three groups separately and use them to parametrize the jump matrices of the Riemann-Hilbert problem on the graph Σ .




Figure 3: The support of the jump matrices J . The graph Σ_0 is in black (the triangulation).

Edge coordinates and jump matrices on e_j . To each edge $e \in E(\Sigma_0)$ we associate $n-1$ non-vanishing variables

$$\mathbf{z} = \mathbf{z}_e = (z_1, \dots, z_{n-1}) \in (\mathbb{C}^\times)^{n-1} \quad (5.3)$$

and introduce their exponential counterparts:

$$\zeta = \zeta_e = (\zeta_1, \dots, \zeta_{n-1}) \in \mathbb{C}^{n-1}, \quad \zeta_j = \frac{1}{n} \log z_j^n. \quad (5.4)$$

The jump matrix on the oriented edge $e \in \mathbf{E}(\Sigma_0)$ is given by

$$S(\mathbf{z}) = \mathbf{z}^{-\mathbf{h}} \mathbf{P} \sigma := \prod_{j=1}^{n-1} z_j^{-h_j} \mathbf{P} \sigma = \prod_{\ell=1}^{n-1} z_\ell^\ell \begin{pmatrix} 0 & \dots & & (-1)^{n-1} \prod_{j=1}^{n-1} z_j^{-n} \\ & & \ddots & 0 \\ \vdots & & & \\ 0 & -z_{n-2}^{-n} z_{n-1}^{-n} & 0 & \dots \\ 1 & 0 \dots & & \end{pmatrix} \quad (5.5)$$

where h_i are the dual roots (5.1). For the inverse matrix we have

$$S^{-1}(\mathbf{z}) = \sigma \mathbf{P} \mathbf{z}^{\mathbf{h}} = (-1)^{n-1} \mathbf{z}^{\mathbf{h}^*} \mathbf{P} \sigma.$$

The notation $\mathbf{z}^{\mathbf{h}}$ stands for

$$\mathbf{z}^{\mathbf{h}} = z_1^{h_1} \dots z_{n-1}^{h_{n-1}}. \quad (5.6)$$

The sets of variables (5.3), (5.4) corresponding to an oriented edge e of Σ_0 and the opposite edge $-e$ are related as follows:

$$\zeta_{-e} = (\zeta_{e,n-1}, \dots, \zeta_{e,1}); \quad \mathbf{z}_{-e} := (-1)^{n-1} (z_{e,n-1}, \dots, z_{e,1}). \quad (5.7)$$

Face coordinates and jump matrices on $\mathcal{E}_f^{(i)}$. To each face $f \in F(\Sigma_0)$ (i.e. a triangle of the original triangulation) we associate $\frac{(n-1)(n-2)}{2}$ variables $\xi_f = \{\xi_{f;abc} : a, b, c \in \mathbb{N}, a + b + c = n\}$ and their exponential counterparts $x_{f;abc} := e^{\xi_{f;abc}}$ as follows.

The variables $\xi_{f;abc}$ define the jump matrices $A_i(\xi_f)$ on three edges $\{\mathcal{E}_f^{(i)}\}_{i=1}^3$, which connect a chosen point p_f in each face f of the graph Σ_0 with its three vertices (these edges are shown in red in

Fig. 3). The enumeration of vertices v_1, v_2 and v_3 is chosen arbitrarily for each face f . Namely, for a given vertex v and the face f of Σ_0 such that $v \in \partial f$ we define the index $f(v) \in \{1, 2, 3\}$ depending on the enumeration that we have chosen for the three edges $\{\mathcal{E}_f^{(i)}\}$ lying in the face f . For example in Fig 3 for the face f containing point p_i we define $f(v_\ell) = 1, f(v_k) = 3$ and $f(v_s) = 2$.

The matrices $A_{1,2,3}(\boldsymbol{\xi}_f)$ are defined following [21]. First, the matrix A_1 is defined by the formula

$$A_1(\mathbf{x}) = \sigma \left(\prod_{k=n-1}^1 N_k \right) \mathbf{P}, \quad (5.8)$$

where E_{ik} are the elementary matrices and

$$F_i = \mathbf{1} + E_{i+1,i}, \quad H_i(x) := x^{h_i} = \text{diag}(\overbrace{x^{i-n}, \dots, x^{i-n}}^{i \text{ times}}, x^i, \dots, x^i), \quad i = 1, \dots, n-1; \quad (5.9)$$

$$N_k = \left(\prod_{k \leq i \leq n-2} H_{i+1}(x_{n-i-1, i-k+1, k}) F_i \right) F_{n-1}. \quad (5.10)$$

The matrices A_2 and A_3 are obtained from A_1 by cyclically permuting the indices of the variables:

$$A_2(\mathbf{x}) = A_1(\{x_{bca}\}), \quad A_3(\mathbf{x}) = A_1(\{x_{cab}\}); \quad (5.11)$$

the important property of the matrices A_i is the equality

$$A_1 A_2 A_3 = \mathbf{1} \quad (5.12)$$

which guarantees the triviality of total monodromy around the point p_f on each face f . In the first two non-trivial cases the matrices A_i have the following forms:

$SL(2)$: there are no face variables and all matrices $A_i = A$ are given by

$$A = \begin{pmatrix} 0 & 1 \\ -1 & -1 \end{pmatrix}. \quad (5.13)$$

$SL(3)$: there is one parameter $x = x_{111}$ for each face. The matrices A_1, A_2 and A_3 coincide in this case, too; they are given by

$$A(x) = \frac{1}{x} \begin{pmatrix} 0 & 0 & 1 \\ 0 & -1 & -1 \\ x^3 & x^3 + 1 & 1 \end{pmatrix}. \quad (5.14)$$

Jump matrices on stems. The jump matrix on the stem of the cherry connected to a vertex v is defined from the triviality of total monodromy around v .

For each vertex v of Σ_0 of valence n_v the jump matrix on the stem of the cherry attached to v is given by

$$M_v^0 = \left(\prod_{i=1}^{n_v} A_{f_i} S_{e_i} \right)^{-1} \quad (5.15)$$

where f_1, \dots, f_{n_v} and e_1, \dots, e_{n_v} are the faces/edges ordered counterclockwise starting from the stem of the cherry, with the edges oriented away from the vertex (using if necessary the formula (5.7)). Since each product $A_{f_i} S_{e_i}$ is a lower triangular matrix, the matrices M_v^0 are also lower-triangular. The diagonal part of M_v^0 will be denoted by Λ_v and parametrized as follows:

$$\Lambda_v = \text{diag} \left(m_{v;1}, \frac{m_{v;2}}{m_{v;1}}, \dots, \frac{m_{v;n-1}}{m_{v;n-2}}, \frac{1}{m_{v;n-1}} \right). \quad (5.16)$$

Notice that the relation (5.16) can also be written as $\Lambda_v = \mathbf{m}_v^\alpha = \prod_{j=1}^{n-1} m_{v;j}^{\alpha_j}$ where α_j are the roots (5.1).

In order to express Λ_v in terms of ζ and ξ -coordinates, we enumerate the faces and edges incident at the vertex v by f_1, \dots, f_{n_v} and e_1, \dots, e_{n_v} , respectively. We assume the edges to be oriented away from v using (5.7). We also assume without loss of generality that the arc $\mathcal{E}_{f_j}^{(1)}$ is the one connected to the vertex v for all $j = 1, \dots, n_v$. Then (see (4.20) of [8]) we have

$$\Lambda_v = e^{2i\pi L_v} = \mathbf{P} \left(\prod_{f \perp v} \mathbf{x}_f^{\mathbf{h}_f} \right) \left(\prod_{e \perp v} \mathbf{z}_e^{\mathbf{h}_e} \right) \mathbf{P} \quad (5.17)$$

Introduce now the variables $\mu_{v;\ell}$ via

$$\mu_{v;n-\ell} = \sum_{f \perp v} \sum_{\substack{a+b+c=n \\ a,b,c \geq 1}} \xi_{f;abc} \mathbb{G}_{a\ell} + \sum_{e \perp v} \sum_{j=1}^{n-1} \zeta_{e;j} \mathbb{G}_{j\ell} \quad (5.18)$$

where the matrix \mathbb{G} equals to n^2 times the inverse Cartan matrix (see (5.2)).

The relationship between μ 's and variables m_v is

$$m_{v;\ell}^n = e^{n\mu_{v;\ell}}$$

i.e. $\mu_{v;\ell}$ defines $m_{v;\ell}$ up to an n th root of unity. Therefore, the entries $\lambda_{v;j}$ of the diagonal matrices L_v are related to $\mu_{v;j}$ as follows:

$$\lambda_{v;j} \equiv \frac{1}{2\pi i} (\mu_{v;j} - \mu_{v;j-1}) \pmod{\mathbb{Z}}, \quad j = 1, \dots, n. \quad (5.19)$$

Vertex coordinates and jump matrices on cherries To each vertex v of the graph Σ_0 we associate a set of $n-1$ non-vanishing complex numbers $r_{v;i}$, $i = 1, \dots, n-1$ in the following way.

Since the matrix M_v^0 is lower-triangular it can be diagonalized by a lower-triangular matrix C_v^0 such that all diagonal entries of C_v^0 equal to 1:

$$M_v^0 = C_v^0 \Lambda_v (C_v^0)^{-1}. \quad (5.20)$$

Any other lower-triangular matrix C_v diagonalizing M_v^0 can be written as

$$C_v = C_v^0 R_v \quad (5.21)$$

where the matrix R_v (which equals to the diagonal part of C_v , $R_v = (C_v)^D$), is parametrized by $n-1$ variables r_1, \dots, r_{n-1} and their logarithmic counterparts

$$\rho_i = \log r_i, \quad i = 1, \dots, n-1,$$

as follows (we omit the index v below):

$$R = \prod_{i=1}^{n-1} r_i^{\mathbf{h}_i} = \mathbf{r}^{\mathbf{h}} = \left(\prod_{i=1}^{n-1} r_i^i \right)^{-1} \text{diag} \left(\prod_{i=1}^{n-1} r_i^n, \prod_{i=2}^{n-1} r_i^n, \dots, r_{n-2}^n r_{n-1}^n, r_{n-1}^n, 1 \right). \quad (5.22)$$

The jump on the boundary of the cherry is defined to be

$$J_v = C_v(z - t_v)^{-L_v}. \quad (5.23)$$

The point of discontinuity of the function J_v on the boundary of the cherry is assumed to coincide with the point where the stem is connected to the cherry (this point is denoted by β in Fig.4).




Figure 4: On contribution of one of the loops to the form $d\Theta$

5.2 Parametrization of the space \mathcal{M}

The set of jump matrices on the graph Σ constructed in the previous section can be used to parametrize the space \mathcal{M} . Recall that the vertices of the graph Σ_0 are in one-to-one correspondence with points t_j ; thus the vertex connected to the cherry around t_j will be denoted by v_j . To construct the monodromy map as $SL(n)$ representation of $\pi_1(\mathbf{CP}^1 \setminus \{t_1, \dots, t_N\}, \infty)$ we topologically identify the punctured sphere with the complement of connected and simply connected neighbourhoods of the t_j 's that contain also the distal vertex of the stem. The fundamental group of the punctured sphere and of this sphere with deleted neighbourhood is the same. Equivalently, for an element in the fundamental group we choose a representative that does not intersect the cherry and stem.

Then the map is then defined as follows; for $\sigma \in \pi_1(\mathbf{CP}^1 \setminus \{U_1, \dots, U_N\}, \infty)$ the corresponding monodromy is given by

$$M_\sigma := \prod_{e \in \sigma \cap \mathbf{E}(\Sigma)} J_e^{\nu(e, \sigma)} \quad (5.24)$$

where the product is taken in the same order as the order of the edges being crossed by σ and $\nu(e, \sigma) \in \{\pm 1\}$ is the orientation of the intersection of the (oriented) edge e and σ at the point of intersection. With this definition the analytic continuation of Ψ satisfies $\Psi(z^\sigma) = \Psi(z)M_\sigma^{-1}$. This allows us to relate the normalization of the eigenvector matrices C_j with that of the matrices C_j^0 (5.20). To this end, choose z_0^j in the connected region of $\mathbb{P}^1 \setminus \Sigma$ that contains the j -th cherry (see Fig. 5).

Then the monodromy matrix M_j equals to the ordered product of jump matrices at the edges of Σ crossed by σ_j and it has the form

$$M_j = T_j M_{v_j}^0 T_j^{-1} \quad (5.25)$$

where the matrix T_j equals to the product of jump matrices on the edges of Σ crossed by σ_j as it is traversed from $z_0 = \infty$ to z_0^j . Therefore, the diagonal form of the monodromy matrix M_j is:

$$M_j = C_j \Lambda_j C_j^{-1}, \quad C_j = T_j C_{v_j}^0 R_{v_j}. \quad (5.26)$$

This determines, unambiguously, the normalization of the matrix C_j in terms of the Fock–Goncharov coordinates, thus providing a complete parametrization of $\widehat{\mathcal{M}}$.

5.3 Symplectic form

The computation of the symplectic form $d\Theta = \omega_{\mathcal{M}}$ is given in [8].

Theorem 5.1 (Thm. 4.1 of [8]). *In the coordinate chart parametrized by coordinates*

$$\left\{ \mathbf{z}_e, \mathbf{x}_f, \mathbf{r}_v : e \in E(\Sigma_0), f \in F(\Sigma_0), v \in V(\Sigma_0) \right\} \quad (5.27)$$

the symplectic form $\omega_{\mathcal{M}}$ (3.16) is given by

$$-2\pi i \omega_{\mathcal{M}} = \frac{1}{2} \sum_{v \in V(\Sigma_0)} \omega_v + \frac{1}{2} \sum_{f \in F(\Sigma_0)} \omega_f + n \sum_{v \in V(\Sigma_0)} \sum_{i=1}^{n-1} d\rho_{v;i} \wedge d\mu_{v;i}. \quad (5.28)$$

The variables $\mu_{v;j}$ are defined by (5.18).




Figure 5: Local triangular monodromy $M_{v_j}^0$ and global monodromy $M_{v_j}^0 = T_j M_{v_j}^0 T_j^{-1}$.

The form ω_v in (5.28) is defined as follows: for each vertex $v \in V(\Sigma_0)$ of valence n_v let $\{e_1, \dots, e_{n_v}\}$ be the incident edges ordered counterclockwise starting from the one on the left of the stem and oriented away from v . Let $\{f_1, \dots, f_{n_v}\} \in F(T)$ be the faces incident to v and counted in counterclockwise order from the one containing the cherry. We denote the order relation by \prec . Then

$$\begin{aligned} \omega_v = & \sum_{e' \prec e \perp v} \mathbb{G}_{ij} d\zeta_{e';i} \wedge d\zeta_{e;j} + \sum_{f \prec e \perp v} \sum_{a+b+c=n} \sum_{\ell=1}^{n-1} \mathbb{G}_{f(v),\ell} d\xi_{f;abc} \wedge d\zeta_{e;\ell} \\ & + \sum_{e \prec f \perp v} \sum_{a+b+c=n} \sum_{\ell=1}^{n-1} \mathbb{G}_{f(v),\ell} d\zeta_{e;\ell} \wedge d\xi_{f;abc} + \sum_{f' \prec f \perp v} \sum_{\substack{a+b+c=n \\ a^4 b^4 c^4 = n}} \mathbb{G}_{f'(v),f(v)} d\xi_{f';a'b'c'} \wedge d\xi_{f;abc} \end{aligned} \quad (5.29)$$

where the subscript $f(v)$ indicates the index a, b or c depending on the value $f(v) \in \{1, 2, 3\}$, respectively. The form ω_f for face f is given by

$$\omega_f = \sum_{\substack{i+j+k=n \\ i'+j'+k'=n}} F_{ijk;i'j'k'} d\xi_{f;ijk} \wedge d\xi_{f;i'j'k'} \quad (5.30)$$

where $F_{ijk;i'j'k'}$ are the following constants

$$\frac{1}{n} F_{ijk;i'j'k'} = (k\Delta i - i\Delta k)H(\Delta i \Delta k) + (j\Delta k - k\Delta j)H(\Delta j \Delta k) + (i\Delta j - j\Delta i)H(\Delta i \Delta j) \quad (5.31)$$

and

$$\Delta i = i' - i; \quad \Delta j = j' - j; \quad \Delta k = k' - k;$$

$H(x)$ is the Heaviside function:

$$H(x) = \begin{cases} 1 & x > 0 \\ \frac{1}{2} & x = 0 \\ 0 & x < 0 \end{cases} . \quad (5.32)$$

We point out that while the coordinates ξ, ζ, ρ are defined on a covering space of the character variety (with the deck transformations being shifts by integer multiples of $2i\pi$), the symplectic form (5.29) is defined on the character variety itself. Notice also that for $SL(2)$ and $SL(3)$ the form ω_f vanishes.

5.4 Symplectic potential

We are going to choose a symplectic potential $\theta_{\mathcal{M}}$ satisfying the equation $d\theta_{\mathcal{M}} = \omega_{\mathcal{M}}$ for the symplectic form $\omega_{\mathcal{M}}$ using the representation (5.28). For convenience we introduce a uniform notation for coordinates ζ_e and $\xi_{f;ijk}$; the number of these coordinates equals $\dim \mathcal{M} - (n-1)N$ (we subtract the number of coordinates ρ_j from the total dimension of \mathcal{M}). These coordinates we denote collectively by

$$\{\kappa_j\}_{j=1}^{\dim \mathcal{M} - (n-1)N}$$

Then the formula (5.28) can be written as

$$-2\pi i \omega_{\mathcal{M}} = \frac{1}{2} \sum_{j < \ell} n_{j\ell} d\kappa_j \wedge d\kappa_\ell + n \sum_{v \in V(\Sigma_0)} \sum_{j=1}^{n-1} d\rho_{v;j} \wedge d\mu_{v;j} \quad (5.33)$$

where all $n_{j\ell}$ are integer numbers and $\mu_{v;j}$ are linear functions of κ_j 's.

Definition 5.1. The symplectic potential $\theta_{\mathcal{M}}$ is defined by the following relation:

$$2\pi i \theta_{\mathcal{M}} = \frac{1}{4} \sum_{j < \ell} n_{j\ell} (\kappa_\ell d\kappa_j - \kappa_j d\kappa_\ell) + \frac{n}{2} \sum_{v \in V(\Sigma_0)} \sum_{j=1}^{n-1} (\mu_{v;j} d\rho_{v;j} - \rho_{v;j} d\mu_{v;j}). \quad (5.34)$$

Obviously, there exist infinitely many choices of the potential for the form $\omega_{\mathcal{M}}$. Our choice (5.34) is due to Theorem 7.1 of [8] which states that the potential $\theta_{\mathcal{M}}$ (5.34) remains invariant if any of the cherries is moved to the neighbouring face.

5.5 $SL(2)$ case

For $n = 2$ the general formula in Thm. 5.1 simplifies considerably to the following (for details see (7.5) of [8])

$$-2\pi i \omega_{\mathcal{M}} = \sum_{k=1}^N \left(\sum_{\substack{e, e' \perp v_k \\ e' \prec e}} d\zeta_{e'} \wedge d\zeta_e + 2 \sum_{e \perp v_k} d\rho_k \wedge d\zeta_e \right); \quad (5.35)$$

the symplectic potential (5.34) $\theta_{\mathcal{M}}$ is given by:

$$2\pi i \theta_{\mathcal{M}} = \sum_{k=1}^N \left(\frac{1}{2} \sum_{\substack{e, e' \perp v_k \\ e' \prec e}} (\zeta_e d\zeta_{e'} - \zeta_{e'} d\zeta_e) + \sum_{e \perp v_k} (\zeta_e d\rho_k - \rho_k d\zeta_e) \right). \quad (5.36)$$

Notice that the expression (5.36) “forgets” about the orientation of vertices since the coordinate ζ_e remains invariant if the orientation of the edge e is changed i.e. when z_e transforms to $-z_e$. Unlike the form $\omega_{\mathcal{M}}$, the potential $\theta_{\mathcal{M}}$ depends on the choice of triangulation Σ_0 ; the change of triangulation implies a non-trivial change of $\theta_{\mathcal{M}}$.

5.5.1 Change of triangulation.

One triangulation Σ_0 can be transformed to any other by a sequence of “flips” of the diagonal in the quadrilateral formed by two triangles with a common edge, see Fig.6. Let us assume that the four cherries attached to the vertices are placed as shown in Fig.6. Then, the assumption that all the monodromies around the four vertices of these triangles are preserved, implies the following equations [8]:

$$\tilde{\kappa}_1 = \frac{\kappa}{\kappa+1} \kappa_1, \quad \tilde{\kappa}_2 = (\kappa+1) \kappa_2, \quad \tilde{\kappa}_3 = \frac{\kappa}{\kappa+1} \kappa_3, \quad \tilde{\kappa}_4 = (\kappa+1) \kappa_4, \quad \tilde{\kappa} = \frac{1}{\kappa} \quad (5.37)$$

where $\kappa_j = z_j^2$, $\tilde{\kappa}_j = \tilde{z}_j^2$, $j = 1, \dots, 4$; $\tilde{\kappa} = \tilde{z}^2$ is the variable on the “flipped” edge. The variables r_j remain invariant under the change of triangulation due to the choice of cherries positions in Fig.6.




Figure 6: Transformation of edges and jump matrices under an elementary flip of an edge of Σ_0 .

Denote the symplectic potential corresponding to the new triangulation by $\tilde{\theta}_{\mathcal{M}}$ and introduce the *Rogers dilogarithm* L which for $x \geq 0$ is defined by (see (1.9) of [45]):

$$L\left(\frac{x}{x+1}\right) := \frac{1}{2} \int_0^x \left\{ \frac{\log(1+y)}{y} - \frac{\log y}{1+y} \right\}. \quad (5.38)$$

As it was shown in Prop.7.1 of [8], the symplectic potentials $\tilde{\theta}_{\mathcal{M}}$ and $\theta_{\mathcal{M}}$ are related as follows:

$$2\pi i(\tilde{\theta}_{\mathcal{M}} - \theta_{\mathcal{M}}) = d \left[L\left(\frac{\kappa}{1+\kappa}\right) \right]. \quad (5.39)$$

Therefore, the function $L\left(\frac{\kappa}{1+\kappa}\right)$ is the generating function of the symplectomorphism corresponding to the elementary flip of the edge of Σ_0 .

6 Tau-function as generating function of monodromy symplectomorphism

Here we extend the definition of the Jimbo-Miwa-Ueno tau-function by including an explicit description of its dependence on the monodromy data.

Definition 6.1. *The tau function is defined by the following set of compatible equations. The equations with respect to t_j are given by the Jimbo-Miwa-Ueno formulæ*

$$\frac{\partial \log \tau}{\partial t_j} = \frac{1}{2} \operatorname{res}_{z=t_j} \operatorname{tr} A^2(z); \quad (6.1)$$

the equations with respect to coordinates on monodromy manifold \mathcal{M} are given by

$$d_{\mathcal{M}} \log \tau = \sum_{j=1}^N \operatorname{tr}(L_j G_j^{-1} d_{\mathcal{M}} G_j) - \theta_{\mathcal{M}}[\Sigma_0] \quad (6.2)$$

where $\theta_{\mathcal{M}}[\Sigma_0]$ is the symplectic potential (5.34) for the form $\omega_{\mathcal{M}}$; we consider the matrices G_j as (meromorphic) functions on $\tilde{\mathcal{M}}$ defined by the formula

$$G_j = \Phi(t_j) \quad (6.3)$$

with Φ the solution of the Riemann-Hilbert problem (3.9).

Using Thm.3.1 and in particular (3.11), we can rewrite this definition in an alternative form, which encodes the complete system (6.1), (6.2):

Definition 6.1'. *The tau-function on $\widetilde{\mathcal{M}}$ is locally defined by equations*

$$d \log \tau = \Theta - \widetilde{\theta}_{\mathcal{M}}[\Sigma_0] \quad (6.4)$$

where Θ is the Malgrange form (3.3) corresponding to solution Φ (3.9) and $\widetilde{\theta}_{\mathcal{M}}[\Sigma_0]$ is the pullback to $\widetilde{\mathcal{M}}$ of $\theta_{\mathcal{M}}[\Sigma_0]$.

The formula (6.2) means that $\log \tau$ is nothing but the generating function of the monodromy symplectomorphism: $d \log \tau$ equals to the difference of symplectic potentials defined in terms of the (extended) Kirillov–Kostant symplectic potential $\theta_{\mathcal{A}}$ and the symplectic potential on the monodromy manifold. It was proven in [44] that the residue of Θ along the points of the Malgrange divisor is a positive integer; thus τ is actually locally analytic on $\widetilde{\mathcal{M}}$; multiplicity of zero of τ equals to the residue of Θ .

We now analyze in more detail the dependence of τ on the Fock–Goncharov coordinates. The tau-function τ defined by (6.2) depends on the full set of variables (z, x, r) on \mathcal{M} . The right-hand sides of equations (6.2) depend on the choice of the triangulation Σ_0 defining the symplectic potential $\theta_{\mathcal{M}}$. However, according to Thm.6.1 of [8], the potential (5.34) is independent of the choice of ciliation of the graph Σ_0 .

The next proposition shows how the tau-function defined in Def.6.1 depends on variables $\rho_{j,i}$: namely, define the second tau-function τ_1 by

$$\tau_1 = \tau \exp \left\{ -\frac{n}{2\pi i} \sum_{j=1}^N \sum_{i=1}^{n-1} \rho_{j,i} \mu_{j,i} \right\}. \quad (6.5)$$

Proposition 6.1. *The tau-function τ_1 (6.5) is independent of the variables $\{r_{j,i}\}$, $j = 1 \dots N$, $i = 1 \dots n - 1$ i.e.*

$$\frac{\partial \log \tau_1}{\partial r_{j,i}} = 0. \quad (6.6)$$

Proof. Denote by G_j^0 the set of matrices G_j which correspond to all variables $r_{j,i} = 1$. Then matrices G_j can be expressed in terms of G_j^0 and $r_{j,i}$ as follows:

$$G_j = G_j^0 R_j \quad (6.7)$$

where the diagonal matrix R_j is given by (5.22). Then,

$$G_j^{-1} dG_j - (G_j^0)^{-1} dG_j^0 = R_j^{-1} dR_j.$$

Therefore, the first sum in (6.2) gets an additive term equal to

$$\sum_{j=1}^N \operatorname{tr} L_j R_j^{-1} dR_j. \quad (6.8)$$

On the other hand, matrices C_j transform under (6.7) in the same way:

$$C_j = C_j^0 R_j \quad (6.9)$$

where the matrices C_j^0 are assumed to be triangular with all 1's on the diagonal.

To get the variation of $\theta_{\mathcal{M}}$ under the transformation (6.9) we observe that the form $\omega_{\mathcal{M}}$ (3.16) transforms under (6.9) as follows:

$$\omega_{\mathcal{M}} \rightarrow \omega_{\mathcal{M}} + \sum_{j=1}^N \operatorname{tr} dL_j \wedge R_j^{-1} dR_j.$$

Therefore, according to our definition of $\theta_{\mathcal{M}}$, the last sum in this expression should be integrated to give

$$\theta_{\mathcal{M}} \rightarrow \theta_{\mathcal{M}} + \sum_{j=1}^N \text{tr} L_j R_j^{-1} dR_j \quad (6.10)$$

which cancels against (6.8) (alternatively, one can derive (6.10) using the definition of $\mu_{v;i}$ and $\rho_{v;i}$ and (5.34)). ■

The equations for the tau-function with respect to variables \mathbf{z} and \mathbf{x} (or, equivalently, $\boldsymbol{\zeta}$ and $\boldsymbol{\xi}$) implied by Def.6.1 can be obtained from expression (5.34) for the potential $\theta_{\mathcal{M}}$. Below we write these equations explicitly in the $SL(2)$ case.

6.1 $SL(2)$ tau-function

In the $SL(2)$ case the coordinates on $\mathcal{M}_N^{SL(2)}$ are given by edge coordinates $\{\zeta_e\}$ and vertex coordinates $\{\rho_k\}_{k=1}^N$; the potential $\theta_{\mathcal{M}}$ is given by (5.36). Then

$$L_j = \begin{pmatrix} \lambda_j & 0 \\ 0 & -\lambda_j \end{pmatrix} = \frac{1}{2\pi i} \begin{pmatrix} \mu_j & 0 \\ 0 & -\mu_j \end{pmatrix}.$$

and the relationship (6.5) becomes:

$$\tau(\{\zeta_e, \rho_j\}, \{t_j\}) = \tau_1(\{\zeta_e\}, \{t_j\}) \exp \left\{ \frac{1}{\pi i} \sum_{j=1}^N \rho_j \mu_j \right\}. \quad (6.11)$$

where μ_j is the sum of the ζ_e for all edges incident to the j -th vertex. The equations for τ_1 with respect to the edge coordinates take the following form:

Definition 6.2. For a given triangulation Σ_0 the tau-function τ_1 of an $SL(2)$ Fuchsian system is defined by the system (6.1) with respect to poles $\{t_j\}_{j=1}^N$ and the following equations with respect to coordinates $\{\zeta_{e_j}\}_{j=1}^{3N-6}$:

$$\frac{\partial}{\partial \zeta_e} \log \tau_1 = \sum_{j=1}^N \text{tr} \left(L_j G_j^{-1} \frac{\partial G_j}{\partial \zeta_e} \right) - \frac{1}{4\pi i} \left(\sum_{\substack{e' \perp v_1 \\ e < e'}} \zeta_{e'} - \sum_{\substack{e' \perp v_1 \\ e' < e}} \zeta_{e'} + \sum_{\substack{e' \perp v_2 \\ e < e'}} \zeta_{e'} - \sum_{\substack{e' \perp v_2 \\ e' < e}} \zeta_{e'} \right) \quad (6.12)$$

where v_1 and v_2 are vertices of Σ_0 connected by the edge e .

This definition depends on the choice of triangulation Σ_0 . The change of the tau-function τ under an elementary flip of an edge of the triangulation Σ_0 acting on the underlying triangulation follows from (5.39):

Proposition 6.2. Let τ and $\tilde{\tau}$ be tau-functions corresponding to triangulations related by the flip of the edge e shown in Fig.6 . Then

$$\frac{\tilde{\tau}}{\tau} = \exp \left[-\frac{1}{2\pi i} L \left(\frac{e^{2\zeta_e}}{e^{2\zeta_e} + 1} \right) \right] \quad (6.13)$$

under an appropriate choice of branch of the Rogers' dilogarithm L (5.38).

6.2 Equations with respect to Fock-Goncharov coordinates

Here we derive equations for Ψ , G_j and τ with respect to an edge coordinate ζ .

First we notice that for any Riemann-Hilbert problem on an oriented contour C with jump matrix J the variation of the solution of the Riemann-Hilbert problem takes the form :

$$\delta \Psi \Psi^{-1}(w) = \frac{1}{2\pi i} \int_C \frac{\Psi_- \delta J J^{-1} \Psi_-^{-1}}{z - w} dw. \quad (6.14)$$

The formula (6.14) can be easily derived by applying the variation δ to the equation $\Psi_+ = \Psi_- J$ on C which gives $\delta\Psi_+ = \delta\Psi_- J + \Psi_- \delta J$ and then solving the resulting non-homogeneous Riemann-Hilbert problem via Cauchy kernel.

We apply (6.14) to variations of the solution Ψ of the Riemann-Hilbert problem on the Fock-Goncharov graph with respect to the variable ζ corresponding to the edge e . Without loss of generality we assume that the positions of cherries are chosen as in Fig. 7.




Figure 7: Positioning of the cherries for the computation of the derivative in ζ_e .

For simplicity we assume that both vertices v_1 and v_2 connected by e are three-valent but it is not a significant restriction.

The jump matrices depending on ζ are the following: the jump matrix on the edge e (and the one on the reverse edge) according to the general rules in (5.7) is

$$S(\zeta) = \begin{pmatrix} 0 & -e^{-\zeta} \\ e^\zeta & 0 \end{pmatrix}; \quad \zeta_{-e} = \zeta + i\pi. \quad (6.15)$$

Furthermore, using the expression (5.13) for the matrix A and denoting $S_j = S(\zeta_j)$ the jump matrices Q_1 and Q_2 on the stems s_1 and s_2 depend on ζ as follows:

$$Q_1 = (SAS_1AS_2A)^{-1} = \begin{pmatrix} e^{\zeta+\zeta_1+\zeta_2} & -e^{-(\zeta+\zeta_1+\zeta_2)} - e^{-\zeta-\zeta_1+\zeta_2} - e^{-\zeta+\zeta_1+\zeta_2} \\ 0 & e^{-(\zeta+\zeta_1+\zeta_2)} \end{pmatrix},$$

$$Q_2 = (AS_4AS_3AS^{-1})^{-1} = \begin{pmatrix} e^{-(\zeta+\zeta_3+\zeta_4+i\pi)} & 0 \\ e^{\zeta+\zeta_3+\zeta_4+i\pi} + e^{\zeta+\zeta_3-\zeta_4+i\pi} + e^{\zeta-\zeta_3-\zeta_4+i\pi} & e^{\zeta+\zeta_3+\zeta_4+i\pi} \end{pmatrix}.$$

Notice that logarithmic derivatives of the matrices S and Q_1^{-1} and Q_2 with respect to ζ are the same and are given by

$$J_\zeta J^{-1} = -\sigma_3 = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}. \quad (6.16)$$

Then the exponents of monodromy are

$$\lambda_1 = \frac{1}{2i\pi}(\zeta + \zeta_1 + \zeta_2), \quad \lambda_2 = \frac{-1}{2i\pi}(\zeta + \zeta_3 + \zeta_4 + i\pi).$$

Thus

$$L_1 = \frac{1}{2\pi i} \begin{pmatrix} \zeta + \zeta_1 + \zeta_2 & 0 \\ 0 & -(\zeta + \zeta_1 + \zeta_2) \end{pmatrix}, \quad C_1 = \begin{pmatrix} 1 & f_1(\zeta) \\ 0 & 1 \end{pmatrix} \quad (6.17)$$

where

$$f_1(\zeta) = \frac{e^{-\zeta}(e^{-\zeta_1-\zeta_2} + e^{-\zeta_1+\zeta_2} + e^{\zeta_1+\zeta_2})}{e^{\zeta+\zeta_1+\zeta_2} - e^{-(\zeta+\zeta_1+\zeta_2)}}$$

and

$$L_2 = \frac{1}{2\pi i} \begin{pmatrix} -(\zeta + \zeta_3 + \zeta_4 + i\pi) & 0 \\ 0 & \zeta + \zeta_3 + \zeta_4 + i\pi \end{pmatrix}, \quad C_2 = \begin{pmatrix} 1 & 0 \\ f_2(\zeta) & 1 \end{pmatrix} \quad (6.18)$$

where

$$f_2(\zeta) = \frac{e^\zeta(e^{\zeta_3+\zeta_4} + e^{\zeta_3-\zeta_4} + e^{-\zeta_3-\zeta_4})}{e^{-(\zeta+\zeta_3+\zeta_4)} - e^{\zeta+\zeta_3+\zeta_4}}.$$

Introduce the graph Σ'_0 by identifying the vertex v_j with the corresponding pole t_j . The variational formula takes the simplest form if we make an explicit assumption on the growth (1.2) of the solution Ψ near the poles, that is on the real part of the eigenvalues of L_j ; indeed it is known [40] that for a given monodromy representation there is a lattice of solutions to the inverse monodromy representation. The reason is simply that the eigenvalues matrices L_j are defined up to addition of diagonal matrices in $sl_n(\mathbb{Z})$. For $n = 2$ there is therefore a \mathbb{Z}^N (N being the number of poles) lattice of inverse solutions. The transformations between different solutions in this lattice are referred to as ‘‘discrete Schlesinger transformations’’. We use this observation to shift the real part of λ_j 's to a value within the interval $\Re\lambda_j \in [-\frac{1}{2}, \frac{1}{2}]$. Excluding the non-generic cases $\Re\lambda_j = -\frac{1}{2}$ we have

Theorem 6.1. *Denote by ζ_{jk} the FG coordinate corresponding to the edge e_{jk} . Suppose all eigenvalues of matrices L_j satisfy the conditions*

$$|\Re\lambda_j| < \frac{1}{2}. \quad (6.19)$$

Then the function Ψ satisfies the following system of equations with respect to coordinates ζ_{jk} :

$$\frac{d\Psi(z)}{d\zeta_{jk}} = \frac{1}{2\pi i} \left[\int_{t_j}^{t_k} \frac{\Psi_-(w)\sigma_3\Psi_-^{-1}(w)}{z-w} dw \right] \Psi(z), \quad z \neq t_j, t_k \quad (6.20)$$

where the integral along the oriented edge e_{jk} of Σ'_0 in the right hand side is convergent at the endpoints due to condition (6.19). The integral in (6.20) is discontinuous across the edge e_{jk} .

Proof. We denote $\zeta = \zeta_{jk}$ for simplicity, and $j = 1, k = 2$. The expression $\partial_\zeta J J^{-1}$ is nonzero only on the edge, the two stems and the boundaries of the two cherries. A direct computation shows (with the edges oriented as shown in Fig.7), that the expression of $\partial_\zeta J J^{-1}$ on the edge and on the two stems is equal to σ_3 . The jump matrix on the cherry 1 equals to $J_1 = C_1(z - t_1)^{-L_1}$ and on c_2 it equals to $J_2 = C_2(z - t_2)^{-L_2}$. Then a direct computation gives (since $L_{1\zeta} = \frac{\sigma_3}{2i\pi}$ and $L_{2\zeta} = \frac{-\sigma_3}{2i\pi}$):

$$J_{1\zeta} J_1^{-1} = C_{1\zeta} C_1^{-1} - \frac{\log(z - t_1)}{2i\pi} C_1 L_{1\zeta} C_1^{-1} = \begin{pmatrix} 0 & -\frac{2f_1}{1 - e^{-4\pi i \lambda_1}} \\ 0 & 0 \end{pmatrix} - \frac{\log(z - t_1)}{2i\pi} \begin{pmatrix} 1 & -2f_1 \\ 0 & -1 \end{pmatrix},$$

$$J_{2\zeta} J_2^{-1} = C_{2\zeta} C_2^{-1} - \frac{\log(z - t_2)}{2i\pi} C_2 L_{2\zeta} C_2^{-1} = \begin{pmatrix} 0 & 0 \\ \frac{2f_2}{1 - e^{-4\pi i \lambda_2}} & 0 \end{pmatrix} + \frac{\log(z - t_2)}{2i\pi} \begin{pmatrix} -1 & 0 \\ -2f_2(\zeta) & 1 \end{pmatrix}.$$

Consider the first cherry (the second cherry can be treated in parallel); we shall call β the point of intersection of the stem and the cherry. Within a neighbourhood containing the cherry, we have $\Psi(z) = \Phi_1(z)(z - t_1)^{L_1} C_1^{-1}$, with $\Phi_1(t_1) = G_1$. In the integral (6.14) the contribution coming from the first cherry is then the integral

$$\int_{\beta_+}^{\beta_-} \frac{\Psi_- \partial_\zeta J_1 J_1^{-1} \Psi_-^{-1} dw}{w - z} \frac{dw}{2i\pi} = \int_{\beta_+}^{\beta_-} \Phi_1 \left(f_1 \sigma_+ \left(\frac{ie^{2i\pi\lambda_1}}{\sin(2\pi\lambda_1)} (w - t_1)^{2\lambda_1} - \frac{\log(w - t_1)}{i\pi} \right) - \frac{\log(w - t_1)}{2i\pi} C_1^{-1} \sigma_3 C_1 \right) \Phi_1^{-1} \frac{dw}{2i\pi(w - z)} \quad (6.21)$$

where the contour of integration is the circle $|w - t_1| = \text{const}$ starting at β_+ and ending at β_- and the branch-cut of the power is the segment from t_1 to β . We also have used that $[C_1, \sigma_+] = 0$. Under the condition $-1 < 2\Re\lambda_1$ the contribution of the integration on the cherry tends to zero in the limit of zero radius of the cherry. ■

Remark 6.1. While the general formula (6.21) is valid without any restriction on the real parts of λ_j 's, the integral on the boundary of the cherries provide some sort of ‘‘regularization’’ to the integral along the edge. However, if the conditions (6.19) in fulfilled, the regularization are not needed and we arrive at the simplified formula (6.20).

Introducing the notation

$$F(z) = -\Psi_-(z)\sigma_3\Psi_-^{-1}(z) \quad (6.22)$$

we can formulate the following

Corollary 6.1. *The derivatives of G_ℓ on coordinates ζ_{jk} take the form:*

$$\frac{dG_\ell}{d\zeta_{jk}} = \frac{1}{2\pi i} \left[\int_{t_j}^{t_k} \frac{F(w)}{w - t_\ell} dw \right] G_\ell, \quad \ell \neq j, k, \quad (6.23)$$

$$\frac{dG_j}{d\zeta_{jk}} G_j^{-1} = \lim_{z \rightarrow t_j} \left[\int_{t_j}^{t_k} \frac{F(w)}{w - z} \frac{dw}{2i\pi} + G_j \left(\frac{if_j(z - t_j)^{2\lambda_j} \sigma_+}{\sin(2\pi\lambda_j)} - \frac{\log(z - t_j)}{2i\pi} \sigma_3 \right) G_j^{-1} \right], \quad (6.24)$$

$$\frac{dG_k}{d\zeta_{jk}} G_k^{-1} = \lim_{z \rightarrow t_k} \left[\int_{t_j}^{t_k} \frac{F(w)}{w - z} \frac{dw}{2i\pi} + G_k \left(\frac{-if_k(z - t_k)^{-2\lambda_k} \sigma_-}{\sin(2\pi\lambda_k)} + \frac{\log(z - t_k)}{2i\pi} \sigma_3 \right) G_k^{-1} \right] \quad (6.25)$$

where $f_j = (C_j)_{12}$ and $f_k = (C_k)_{21}$ as defined in (6.17) and (6.18).

Proof. The first formula follows from the fact that the connection matrices and exponents at the vertices not connected to the edge are constant under the variation. For definiteness assume $j = 1$, $k = 2$. To find G_1 (G_2 can be treated in the same way) we need to evaluate $\Phi_1(z) = \Psi(z)C_1(z - t_1)^{L_1}$ at $z = t_1$. Differentiating the identity $\Phi_1(z) = \Psi(z)C_1(z - t_1)^{-L_1}$ with respect to ζ and taking the limit $z \rightarrow t_1$ gives the formula, recalling that $G_1 = \Phi_1(t_1)$. Indeed

$$\partial\Phi_1(z)\Phi_1(z)^{-1} = \partial\Psi(z)\Psi(z)^{-1} + \Phi_1(z) \left(\frac{if_1}{\sin(2\pi\lambda_1)} \sigma_+(z - t_1)^{2\lambda_1} - \frac{\log(z - t_1)}{2i\pi} \sigma_3 \right) \Phi_1(z)^{-1}. \quad (6.26)$$

If $\Re 2\lambda_1 > -1$ (which is our standing assumption), in the limit as $z \rightarrow t_1$ we can substitute $\Phi_1(z)$ by G_1 in the above formula, and we obtain the statement.

On a related note; the same result is obtained by looking at the singular behaviour of the integral (6.20) (using results on Cauchy-type integrals in [27], for example) and simply removing the singular part. \blacksquare

Now we come to the following

Proposition 6.3. *The equations (6.12) for the tau-function τ_1 with respect to FG coordinates can be equivalently written as follows:*

$$2\pi i \frac{\partial}{\partial \zeta_{jk}} \log \tau_1 = \sum_{\ell=1}^N \text{reg} \int_{t_j}^{t_k} \frac{\text{tr} A_\ell F(w)}{w - t_\ell} dw - \frac{1}{2} \left(\sum_{\substack{e' \perp v_j \\ e < e'}} \zeta_{e'} - \sum_{\substack{e' \perp v_j \\ e' < e}} \zeta_{e'} + \sum_{\substack{e' \perp v_k \\ e < e'}} \zeta_{e'} - \sum_{\substack{e' \perp v_k \\ e' < e}} \zeta_{e'} \right)$$

where F is given by (6.22); the terms of the sum which need the regularization correspond to $\ell = j$ and $\ell = k$; they are given by

$$\text{reg} \int_{t_j}^{t_k} \frac{\text{tr} A_j F(w)}{w - t_j} dw = \lim_{z \rightarrow t_j} \left[\int_{t_j}^{t_k} \frac{\text{tr} A_j F(w)}{w - z} dw - 2\lambda_j \log(z - t_j) \right] \quad (6.27)$$

and

$$\text{reg} \int_{t_j}^{t_k} \frac{\text{tr} A_k F(w)}{w - t_k} dw = \lim_{z \rightarrow t_k} \left[\int_{t_j}^{t_k} \frac{\text{tr} A_k F(w)}{w - z} dw + 2\lambda_k \log(z - t_k) \right]. \quad (6.28)$$

The proof follows from equations (6.23), (6.24) and (6.25). ■

Acknowledgements. We thank T. Bridgeland, L. Chekhov, L. Feher, V. Fock, S. Fomin, A. Goncharov, J. Harnad, M. Gekhtman, R. Kashaev, A. Nietzke and M. Shapiro for illuminating discussions. We thank also A. Its, O. Lisovyy and A. Prokhorov for clarifying comments. We thank P. Boalch for bringing the references [37, 11] to our attention. Finally, we thank the anonymous referees for helpful commentary. The work of M.B. was supported in part by the Natural Sciences and Engineering Research Council of Canada (NSERC) grant RGPIN-2016-06660. The work of D.K. was supported in part by the NSERC grant RGPIN/3827-2015. The completion of this work was supported by the National Science Foundation under Grant No. DMS-1440140 while the authors were in residence at the Mathematical Sciences Research Institute in Berkeley, California, during the Fall 2019 semester *Holomorphic Differentials in Mathematics and Physics*.

References

- [1] Alekseev, A. Yu.; Faddeev, L. D. $(T^*G)_t$: a toy model for conformal field theory, *Comm. Math. Phys.* **141** No.2 413-422 (1991)
- [2] Alekseev, A., Malkin, A., *Symplectic structure of the moduli space of flat connection on a Riemann surface*, *Comm.Math.Phys.* **169** 99-120 (1995)
- [3] Alekseev, A., Malkin, A., *The hyperbolic moduli space of flat connections and the isomorphism of symplectic multiplicity spaces* *Duke Math. J.*, **93**, No. 3 575-595 (1998)
- [4] Audin, M., *Torus actions on symplectic manifolds*, *Progress in Mathematics*, **93** Birkhäuser (2004)
- [5] Babelon, O., Bernard, D., Talon, M., *Introduction to Classical Integrable Systems*, Cambridge University Press, 2003
- [6] Bertola, M., *The dependence on the monodromy data of the isomonodromic tau function*, *Comm. Math. Phys.* **294**, (2010), 539-579;
- [7] Bertola, M. *Correction to: The Dependence on the Monodromy Data of the Isomonodromic Tau Function*, *Comm. Math. Phys.*, 381(3):1445–1461, 2021.
- [8] Bertola, M., Korotkin, D., *Extended Goldman symplectic structure in Fock-Goncharov coordinates*, [arXiv.org: 1910.06744](https://arxiv.org/abs/1910.06744), to appear in *J. Diff. Geom.* (2021)
- [9] Bertola, M., Korotkin, D., Norton, C., *Symplectic geometry of the moduli space of projective structures in homological coordinates*, *Invent. Math.* **210**, Issue 3, p. 759-814 (2017)
- [10] Bertola, M., Korotkin, D., *WKB expansion for a Yang-Yang generating function and the Berman tau function*, *Theor. Math. Phys.* **206**(3): 259–295 (2021)
- [11] Boalch, P. , *Quasi-Hamiltonian geometry of meromorphic connections*, *Duke Math. J.*, (2007), **139**, Issue 2, 369-405.
- [12] Bolibruch, A.A. , *Inverse monodromy problems of the analytic theory of differential equations*, *Mathematical events of the twentieth century*, Springer, Berlin (2006), 49–74.
- [13] Bondal, A., *A symplectic groupoid of triangular bilinear forms and the braid group*, *Izv. RAN. Ser. Mat.* **68** Issue 4, 19-74 (2004)
- [14] Chekhov, L., Mazzocco, M., *Shear coordinate description of the quantised versal unfolding of D_4 singularity*, *J. Phys. A* **43** no. 44, 442002, 13 pp. (2010)
- [15] Chekhov, L., Mazzocco, M., Roubtsov, V., *Painlevé monodromy manifolds, decorated character varieties, and cluster algebras*, *IMRN*, **2017** (24), 7639-7691
- [16] Chekhov, L., Fock, V., *Quantum Teichmüller space*, *Theor.Math.Phys.* **120** 1245-1259 (1999)
- [17] Etingof, P., *On the dynamical Yang-Baxter equation*, *Proceedings of the International Congress of Mathematicians*, Vol. II (Beijing, 2002), 555–570, Higher Ed. Press, Beijing, 2002. See also Erratum in *Proceedings of the International Congress of Mathematicians*, Vol. I (Beijing, 2002), 649–650, Higher Ed. Press, Beijing, 2002
- [18] Fehér, L., Gábor, A., Pusztai, B.G., *On dynamical r -matrices obtained from Dirac reduction and their generalizations to affine Lie algebras*, *J. Phys. A* **34** (2001), no. 36, 7335–7348.

- [19] Flaschka, H., Newell, A.C. , *The inverse monodromy transform is a canonical transformation*, Nonlinear problems: present and future (Los Alamos, N. M. , 1981), **61**, 65-89
- [20] Fock, V.V., *Description of moduli space of projective structures via fat graphs*, arXiv:hep-th/9312193
- [21] Fock, V., Goncharov, A., *Moduli spaces of local systems and higher Teichmüller theory*, Publications Mathématiques de l'Institut des Hautes Études Scientifiques **103**, Issue 1, pp 1-211 (2006)
- [22] Fock, V., Goncharov, A., *The quantum dilogarithm and of representations quantum cluster varieties* , Invent. Math., **175**, Issue 2, 223-286 (2009)
- [23] Fock, V., Rosly, A., *Poisson structure on moduli of flat connections on Riemann surfaces and r -matrix*, Moscow Seminar in Mathematical Physics, 67–86, Amer. Math. Soc. Transl. Ser. 2, 191, Adv. Math. Sci., 43, Amer. Math. Soc., Providence, RI, 1999.
- [24] Fomin, S., Zelevinsky, A., *Cluster algebras. I. Foundations*. J. Amer. Math. Soc. **15** no. 2, 497-529 (2002)
- [25] Gaiotto, D., Moore, G., Nietzke, A., *Wall-crossing, Hitchin systems and WKB approximation*, hep-th 0907.3987
- [26] Gamayun, O., Iorgov, N., Lisovyy, O., *Conformal field theory of Painlevé VI*, J. High Energy Phys., **10** 038 pages, (2012).
- [27] Gakhov, F. D., *Boundary value problems*, Dover Publications, (1990).
- [28] M.Gekhtman, M.Shapiro, A.Veinstein, *Cluster algebras and Poisson geometry*, Math.Surveys and Monographs, **167**, AMS (2010)
- [29] Gekhtman, M, Nakanishi, T., Rupel, D., *Hamiltonian and Lagrangian formalisms of mutations in cluster algebras and application to dilogarithm identities*, J. Integrable Syst. 2 (2017), no. 1, xyx005, 35 pp.
- [30] Goldman, W. *The symplectic nature of fundamental groups of surfaces*, Adv. in Math. **54**, 200-225 (1984)
- [31] Goldman, W. *Invariant functions on Lie groups and Hamiltonian flows of surface group representations*, Invent. Math. **85**, 263-302 (1986)
- [32] Hikami, K., *Note on Character Varieties and Cluster Algebras*, SIGMA, **15** , 003, 32 p. (2019)
- [33] Hitchin, N., *Frobenius manifolds*, in "Gauge Theory and Symplectic Geometry", J.Hurtubise and F.Lalonde (eds), NATO ASI Series C 488, Kluwer, Dordrecht (1997)
- [34] Its, A., Prokhorov, A., *On some Hamiltonian properties of the isomonodromic tau functions*, Rev. Math. Phys. **30** (2018), no. 7, 1840008, 38 pp.
- [35] Its, A., Prokhorov, A., *Connection problem for the tau-function of the sine-Gordon reduction of Painlevé-III equation via the Riemann-Hilbert approach*, IMRN **22** , 6856-6883 (2016)
- [36] Its, A., Lisovyy, O., Prokhorov, A., *Monodromy dependence and connection formulae for isomonodromic tau functions*, Duke Math. J. **167**, no. 7 (2018), 1347-1432
- [37] Jeffrey, L., *Extended moduli spaces of flat connections on Riemann surfaces*, Math. Ann. **298** (1994), 667–692.
- [38] Jeffrey, L., Weitsman J., *Bohr-Sommerfeld orbits in the moduli space of flat connections and the Verlinde dimension formula*, Comm. Math. Phys., **150**, No.3, 593-630 (1992)
- [39] Jimbo, M., Miwa, T., and Ueno, K., *Monodromy preserving deformation of linear ordinary differential equations with rational coefficients. I. General theory and τ -function*, Phys. D, **2** (2): 306-352 (1981)
- [40] Jimbo, M., Miwa, T., *Monodromy preserving deformation of linear ordinary differential equations with rational coefficients II*, Phys. D, **2** (3): 407-448 (1981)
- [41] Kawai, S., *The symplectic nature of the space of projective connections on Riemann surfaces*, Math Ann **305** 161-182 (1996)
- [42] Korotkin, D.; Samtleben, H. *Quantization of Coset Space σ -Models Coupled to Two-Dimensional Gravity*, Comm.Math.Phys., **190**, Issue 2, pp. 411-457 (1997).

- [43] Korotkin, D., *Periods of meromorphic quadratic differentials and Goldman bracket*, in Proceedings of Symposia in Pure Mathematics, **100** ed. by Chiu-Chu, M. and Mulase, M., (2018)
- [44] Malgrange, B., *Sur les Déformation Isomonodromiques*, in *Mathématique et Physique (E.N.S. Séminaire 1979-1982)*, p.401-426, Birkhäuser, Boston (1983)
- [45] Nakanishi, T., *Rogers dilogarithms of higher degree and generalized cluster algebras*, J. Math. Soc. Japan **70** (2018), no. 4, 1269–1304.
- [46] Nekrasov, N., Rosly, A., Shatashvili, S., *Darboux coordinates, Yang-Yang functional, and gauge theory*, Nuclear Phys. B Proc. Suppl. **216**, 69-93 (2011)
- [47] Palesi, F., *Introduction to positive representations and Fock-Goncharov coordinates*, <https://hal.archives-ouvertes.fr/hal-01218570>
- [48] Sato, M., Miwa, T., Jimbo, M., *Holonomic quantum fields II. The Riemann-Hilbert problem*, Publ. RIMS, Kyoto Univ., **15**, 201-278 (1979)
- [49] Soudères, I., *Functional equations for Rogers dilogarithm*, Ann. Inst. Fourier (Grenoble) **68** no. 1, 151–169 (2018)
- [50] Schemmel, M., *Stationary Axisymmetric Gravity in the Isomonodromic Canonical Formulation*, Diplomarbeit, 86 p., Hamburg preprint: Schemmel, M., January 1997.
- [51] Ugaglia, M., *On a Poisson structure on the space of Stokes matrices*, IMRN, issue 9, 473-493 (1999)