

# On Generalized $QQ$ -systems

Zoltán Bajnok<sup>1</sup>, Etienne Granet<sup>2,3</sup>, Jesper Lykke Jacobsen<sup>2,3,4</sup>  
and Rafael I. Nepomechie<sup>5</sup>

## Abstract

We formulate  $QQ$ -systems for the closed XXZ, open XXX and open quantum-group-invariant XXZ quantum spin chains. Polynomial solutions of these  $QQ$ -systems can be found efficiently, which in turn lead directly to the admissible solutions of the corresponding Bethe ansatz equations.

---

<sup>1</sup> Wigner Research Centre for Physics, Konkoly-Thege Miklós u. 29-33, 1121 Budapest, Hungary, bajnok.zoltan@wigner.mta.hu

<sup>2</sup> Institut de Physique Théorique, Paris Saclay, CEA, CNRS, 91191 Gif-sur-Yvette, France

<sup>3</sup> Laboratoire de Physique de l'Ecole Normale Supérieure, ENS, Université PSL, CNRS, Sorbonne Université, Université Paris-Diderot, Sorbonne Paris Cité, Paris, France, etienne.granet@gmail.com

<sup>4</sup> Sorbonne Université, École Normale Supérieure, CNRS, Laboratoire de Physique (LPENS), 75005 Paris, France, jesper.jacobsen@ens.fr

<sup>5</sup> Physics Department, P.O. Box 248046, University of Miami, Coral Gables, FL 33124 USA, nepomechie@miami.edu

# 1 Introduction

An inconvenient truth about quantum integrable models – well-known to experts but seldom acknowledged – is that the corresponding Bethe ansatz (BA) equations (to which exact solutions of such models invariably reduce) are very difficult to solve. Various approaches to solving BA equations have been investigated, see e.g. [1, 2] and references therein. Significant further progress on this problem was recently achieved in [3], which formulated so-called  $QQ$ -systems, whose polynomial solutions can be found efficiently; the zeros of the fundamental  $Q$ -function are the sought-after Bethe roots. The  $SU(2)$ -invariant  $QQ$ -system was an essential ingredient in the recent computation of torus partition functions [4, 5], which exploited also techniques from algebraic geometry.

The  $QQ$ -systems in [3] were restricted to *rational* BA equations for *closed* spin chains with periodic boundary conditions. The purpose of this paper is to generalize the  $SU(2)$ -invariant  $QQ$ -system [3] in two different directions: from rational to trigonometric, and from closed to open. These new  $QQ$ -systems will be used to compute partition functions for trigonometric vertex models and for vertex models with boundaries [6].

The outline of this paper is as follows. In Sec. 2, we review the  $QQ$ -system from [3] for the closed XXX spin chain with periodic boundary conditions. However, we provide an alternative derivation based on [7], which is convenient for deriving generalizations. In Sec. 3, we formulate a  $QQ$ -system for the closed XXZ spin chain with periodic boundary conditions. We then turn to open spin chains. In Sec. 4, we formulate a  $QQ$ -system for the  $SU(2)$ -invariant open XXX spin chain. A  $QQ$ -system for the quantum-group-invariant open XXZ spin chain [8] is formulated in Sec. 5. We conclude in Sec. 6 with a brief summary and a list of some interesting open problems.

## 2 Closed XXX $QQ$ -system

In this section we review the  $QQ$ -system [3] whose polynomial solutions provide the full spectrum of the closed XXX spin chain of length  $N$  with periodic boundary conditions, whose Hamiltonian is given by

$$\mathbb{H} = \sum_{k=1}^N \vec{\sigma}_k \cdot \vec{\sigma}_{k+1}, \quad \vec{\sigma}_{N+1} \equiv \vec{\sigma}_1. \quad (2.1)$$

First the model is introduced and its solution by the algebraic BA method is recalled, together with the physicality conditions for the Bethe roots. We then describe the relevant  $QQ$ -system and construct its solution explicitly. We show that polynomial solutions are in one-to-one correspondence with the physical solutions.

### 2.1 Review of the algebraic BA solution

The closed spin-1/2 XXX spin chain and its solution can be succinctly formulated with the help of an  $SU(2)$ -invariant solution of the Yang-Baxter equation given by the  $4 \times 4$   $R$ -matrix

(see e.g. [9])

$$\mathbb{R}(u) = (u - \frac{i}{2})\mathbb{I} + i\mathbb{P}, \quad (2.2)$$

where  $\mathbb{P}$  is the permutation matrix,  $\mathbb{I}$  is the identity matrix, and  $u$  is the spectral parameter. For  $N$  sites with periodic boundary conditions, one can introduce the monodromy matrix  $\mathbb{M}$  and the transfer matrix  $\mathbb{T}$  as

$$\mathbb{T}(u) = \text{tr}_0(\mathbb{M}_0(u)), \quad \mathbb{M}_0(u) = \mathbb{R}_{01}(u) \mathbb{R}_{02}(u) \dots \mathbb{R}_{0N}(u). \quad (2.3)$$

An auxiliary space denoted by index 0 has been introduced, and lower indices help indicate the spaces in which operators act. The transfer matrix is obtained by tracing over the auxiliary space, and thus acts on the quantum space, which is the  $N$ -fold tensor product of  $\mathbb{C}^2$  accommodating all possible states of spin-up and spin-down. As a consequence of the Yang-Baxter equation, the transfer matrix forms a one-parameter family of commuting operators

$$[\mathbb{T}(u), \mathbb{T}(v)] = 0, \quad (2.4)$$

and generates conserved charges in involution, including the Hamiltonian of the system (2.1).

We are interested in the eigenvectors and eigenvalues of the transfer matrix. The former can be generated from the all spin-up reference state

$$|0\rangle = \left(\begin{smallmatrix} 1 \\ 0 \end{smallmatrix}\right)^{\otimes N} \quad (2.5)$$

by acting with a matrix element of the monodromy matrix as

$$\mathbb{B}(u_1) \dots \mathbb{B}(u_M)|0\rangle \equiv |u_1, \dots, u_M\rangle, \quad \mathbb{M}_0(u) = \begin{pmatrix} \mathbb{A}(u) & \mathbb{B}(u) \\ \mathbb{C}(u) & \mathbb{D}(u) \end{pmatrix}. \quad (2.6)$$

The eigenvalues  $T(u)$  of the transfer matrix

$$\mathbb{T}(u)|u_1, \dots, u_M\rangle = T(u)|u_1, \dots, u_M\rangle \quad (2.7)$$

satisfy the  $TQ$ -relation

$$T(u) Q(u) = (u + \frac{i}{2})^N Q(u - i) + (u - \frac{i}{2})^N Q(u + i), \quad (2.8)$$

where  $Q$  encodes the Bethe roots  $\{u_i\}$ :

$$Q(u) = \prod_{j=1}^M (u - u_j). \quad (2.9)$$

As follows from the definition of the transfer matrix and its commutativity property (2.4),  $T(u)$  is a polynomial in  $u$ , and is thus regular at  $u_j$ . The  $TQ$ -relation (2.8) then leads to the BA equations for the roots:

$$\left(\frac{u_j + \frac{i}{2}}{u_j - \frac{i}{2}}\right)^N = - \prod_{k=1}^M \frac{u_j - u_k + i}{u_j - u_k - i}, \quad j = 1, \dots, M. \quad (2.10)$$

For roots with multiplicities, we have further equations [10, 11]. Since repeated roots do not seem to appear in this model (see e.g. [2]), we assume that roots never coincide.

We call a solution of the BA equations *physical* if the corresponding Bethe vector (2.6) is an eigenvector of the transfer matrix. Unfortunately, not all solutions of the BA equations are physical. Solutions that contain the roots  $\pm \frac{i}{2}$  can be unphysical, i.e. they might solve the BA equations, but there is no related eigenvector of the transfer matrix. We define a solution  $\{u_1, \dots, u_M\}$  of the BA equations to be *admissible*, if all roots are finite and pairwise distinct; and, if they are of the form  $\{\frac{i}{2}, -\frac{i}{2}, u_1, \dots, u_{M-2}\}$  (which we call a *singular* solution), then the further constraint

$$\prod_{j=1}^{M-2} \frac{(u_j + \frac{i}{2})(u_j + \frac{3i}{2})}{(u_j - \frac{i}{2})(u_j - \frac{3i}{2})} = (-1)^N \quad (2.11)$$

is satisfied. It was shown in [12] that admissibility implies physicality, and the converse follows from Lemmata 2 and 4 of [7]. Hence, admissibility and physicality are equivalent.

Alternatively, it was observed in [3] that the polynomial solutions of a  $QQ$ -system on an appropriately chosen diagram can be computed efficiently, and correctly account for the physical solutions. In the remainder of this section, we provide an alternative derivation of these results based on [7], which we will subsequently use to generalize this  $QQ$ -system.

## 2.2 $QQ$ -system

For given values of  $N$  and  $M$ , the  $Q$ -functions  $Q_{a,s}$  are defined on a Young tableau with the indices referring to the vertex  $(a, s)$ , where the  $a$ -axis is vertical and the  $s$ -axis is horizontal, see Fig. 1. These  $Q$ -functions satisfy the  $QQ$ -equations, which are formulated around a face as <sup>1</sup>

$$Q_{a+1,s}(u) Q_{a,s+1}(u) \propto Q_{a+1,s+1}^+(u) Q_{a,s}^-(u) - Q_{a+1,s+1}^-(u) Q_{a,s}^+(u), \quad (2.12)$$

where  $f^\pm(u) = f(u \pm \frac{i}{2})$ . The relevant diagram for the closed XXX spin chain with the boundary conditions,  $Q_{2,s} = 1$ ,  $Q_{1,s>M} = 1$ , is displayed in Fig. 1. The initial condition

$$Q_{0,0}(u) = u^N, \quad Q_{1,0}(u) = Q(u) = \prod_{j=1}^M (u - u_j), \quad (2.13)$$

leads to a unique solution of the  $QQ$ -system. The degree of the polynomial  $Q_{a,s}(u)$  is given by the number of boxes in the Young tableau to the right and top of the vertex  $(a, s)$ . Let us see how we can proceed column-by-column and express all  $Q$ -functions in terms of  $Q_{0,0}$  and  $Q_{1,0}$ .

The  $QQ$ -equation for  $(a, s) = (1, 0)$  can be solved easily

$$Q_{1,1}(u) = Q_{1,0}^+(u) - Q_{1,0}^-(u) \equiv Q'_{1,0}(u) = Q'(u), \quad (2.14)$$

---

<sup>1</sup>A  $Q$ -function is defined up to a multiplicative constant. For definiteness, we generally treat  $\propto$  as equality, or (as in (2.14)) with an extra minus sign.




Figure 1: Non-trivial  $Q$ -functions for the closed XXX spin chain.

where we have introduced the discrete derivative defined by

$$f'(u) = f^+(u) - f^-(u). \quad (2.15)$$

The function  $Q_{1,1}$  is automatically a polynomial of degree  $M - 1$ . The equation for  $(a, s) = (0, 0)$  gives

$$Q_{0,1}Q_{1,0} = Q_{0,0}^-Q_{1,1}^+ - Q_{0,0}^+Q_{1,1}^-. \quad (2.16)$$

Making use of (2.14) and (2.16), it follows that

$$Q_{0,1}Q = Q_{0,0}^-Q^{++} + Q_{0,0}^+Q^{--} - Q(Q_{0,0}^- + Q_{0,0}^+), \quad (2.17)$$

or

$$(Q_{0,1} + Q_{0,0}^- + Q_{0,0}^+)Q = Q_{0,0}^-Q^{++} + Q_{0,0}^+Q^{--}. \quad (2.18)$$

Recognizing the RHS of the above equation as the RHS of the  $TQ$ -relation (2.8), one obtains

$$T = Q_{0,1} + Q_{0,0}^- + Q_{0,0}^+. \quad (2.19)$$

Polynomiality of  $Q_{0,1}$  is equivalent to the polynomiality of  $T$ , which leads to the BA equations (2.10).

### 2.2.1 $Q_{a,s}$ in terms of $Q$ and $P$

We now show that the polynomiality of the remaining  $Q$ -functions is equivalent to the admissibility of  $\{u_1, \dots, u_M\}$ . To this end, we define a function  $P(u)$ , such that

$$Q_{0,0} = P^+Q^- - P^-Q^+. \quad (2.20)$$

Using this parametrization for  $Q_{0,0}$ , one can easily show that

$$Q_{0,1} \propto P'^+Q'^- - P'^-Q'^+, \quad (2.21)$$

where prime denotes discrete derivative (2.15). Repeating the calculations starting from  $(a, s) = (1, 1)$  and  $(a, s) = (0, 1)$ , we arrive at

$$Q_{1,2} = Q'', \quad Q_{0,2} \propto P''^+Q''^- - P''^-Q''^+. \quad (2.22)$$

This can be iterated further

$$Q_{1,n} = Q^{(n)}, \quad Q_{0,n} \propto P^{(n)+} Q^{(n)-} - P^{(n)-} Q^{(n)+}, \quad (2.23)$$

where the superscript  $(n)$  denotes the  $n^{\text{th}}$  discrete derivative. In short, all  $Q$ -functions can be expressed in terms of  $P$  and  $Q$ . Clearly, if  $P$  is a polynomial, then all  $Q$ -functions are polynomial. In the following we show that the polynomiality of  $P$  is in fact equivalent to the polynomiality of  $Q_{0,2}$ . We also derive that polynomiality of  $P$  is equivalent to the admissibility of the roots  $\{u_1, \dots, u_M\}$ .

### 2.2.2 Construction of $P$

We construct  $P$  as in [7] by generalizing the approach in [13] (which implicitly assumes that all Bethe roots are regular) to the case of a singular solution. String configurations have roots that differ by  $i$ :  $u_{i_1} - u_{i_2} = i$ , and it is well known (see e.g. [7]) that the only exact string solution consists of a pair of singular roots  $u_1 = \frac{i}{2}$  and  $u_2 = -\frac{i}{2}$ . In the presence of such singular roots, the  $Q$ -function takes the following form

$$Q(u) = u^+ u^- \bar{Q}(u), \quad \bar{Q}(u) = \prod_{u_j \neq \pm \frac{i}{2}} (u - u_j). \quad (2.24)$$

We start by dividing (2.20) by  $Q^+ Q^-$ . We need to write

$$R(u) = \frac{u^N}{Q^+ Q^-} = \frac{u^{N-2}}{u^{++} u^{--} \bar{Q}^+ \bar{Q}^-} \quad (2.25)$$

in the form

$$R = \frac{P^+}{Q^+} - \frac{P^-}{Q^-} = \left( \frac{P}{Q} \right)', \quad (2.26)$$

i.e. we need to “integrate”  $R$  in the discrete sense. To this end, we perform a partial fraction decomposition of (2.25)

$$R = \pi + \frac{q_+}{\bar{Q}^+} + \frac{q_-}{\bar{Q}^-} + \frac{a_+}{u^{++}} + \frac{a_-}{u^{--}}, \quad (2.27)$$

where  $\pi$  is a polynomial of order  $N - 2M$ , the polynomials  $q_{\pm}$  have degree less than  $\bar{Q}$ , while  $a_{\pm}$  are constants. Using the relation

$$R^+ + R^- = \frac{T}{Q^{++} Q^{--}} = \frac{T}{u^+ u^- u^{+++} u^{---} \bar{Q}^{++} \bar{Q}^{--}} \quad (2.28)$$

which follows from the  $TQ$ -relation (2.8), one can investigate all the singularities explicitly. In particular, the RHS of (2.28) has no singularities at the zeros of  $\bar{Q}$ , implying

$$q_+ = q^+, \quad q_- = -q^-, \quad (2.29)$$

for some polynomial  $q(u)$ . The coefficients  $a_{\pm}$  can be determined from the residues of (2.28) at  $u = \mp \frac{i}{2}$ :

$$a_{\pm} = \mp \frac{T(\mp \frac{i}{2})}{2i \bar{Q}(\pm \frac{i}{2}) \bar{Q}(\mp \frac{3i}{2})}. \quad (2.30)$$

The polynomial  $\pi$  can always be written as

$$\pi = \rho' = \rho^+ - \rho^-, \quad (2.31)$$

where  $\rho$  is a polynomial. Clearly,  $\rho$  is defined up to a constant. Changing this constant by  $a$  modifies  $P$  as  $P + aQ$ . This additional term, however, disappears from  $R$  and  $Q_{0,0}$ , thus is irrelevant for us.

In the absence of singular roots, we have  $\bar{Q} = Q$  and  $a_{\pm} = 0$ ; hence, the polynomial  $P = \rho Q + q$  satisfies (2.26), which implies the required Eq. (2.20), see also [13]. In the presence of singular roots, the “integration” of  $R$  in (2.26) requires to “integrate”  $u^{-1}$ , appearing in (2.27). To this end, we define the function  $p(u)$  by

$$p'(u) = \frac{1}{u}, \quad p(u) = -i\psi(-iu + \frac{1}{2}), \quad (2.32)$$

where  $\psi(u)$  is the digamma function

$$\psi(u) = -\gamma + \sum_{n=0}^{\infty} \left( \frac{1}{n+1} - \frac{1}{n+u} \right). \quad (2.33)$$

In view of the fact

$$\begin{aligned} \frac{a_+}{u^{++}} + \frac{a_-}{u^{--}} &= (a_+ p^{++} + a_- p^{--})' = \left[ \frac{1}{2}(a_+ - a_-)(p^{++} - p^{--}) + \frac{1}{2}(a_+ + a_-)(p^{++} + p^{--}) \right]' \\ &= \left[ \frac{1}{2}(a_+ - a_-)\left(\frac{1}{u^+} + \frac{1}{u^-}\right) + \frac{1}{2}(a_+ + a_-)(p^{++} + p^{--}) \right]', \end{aligned} \quad (2.34)$$

we see that the function  $P$  satisfying (2.26) takes the form

$$P = \rho Q + u^+ u^- q + (a_+ - a_-)u\bar{Q} + \frac{1}{2}(a_+ + a_-)(p^{++} + p^{--})Q. \quad (2.35)$$

It is a polynomial if and only if  $a_+ = -a_-$ , i.e. when

$$(-1)^M \frac{\bar{Q}(+\frac{i}{2})\bar{Q}(+\frac{3i}{2})}{\bar{Q}(-\frac{i}{2})\bar{Q}(-\frac{3i}{2})} = 1 \quad (2.36)$$

is satisfied. Here we used the  $TQ$ -relation (2.8) to eliminate  $T(\pm\frac{i}{2})$  in (2.30). Clearly this is the admissibility condition for singular solutions (2.11). Thus, we have just proven that polynomiality of  $P$  is equivalent to the admissibility of the roots.

Even if  $P$  is not a polynomial, the relation (2.26) implies (2.20), which leads to

$$T = P^{++}Q^{--} - P^{--}Q^{++}. \quad (2.37)$$

This implies that  $P$  also satisfies the  $TQ$ -relation

$$TP = (u^+)^N P^{--} + (u^-)^N P^{++}. \quad (2.38)$$

Thus  $P$  and  $Q$  are the two independent solutions of this second order difference equation, and (2.20) is the corresponding Wronskian relation. It has been known (see e.g. [14, 15]) that the two independent solutions of the  $TQ$ -relation are both polynomial iff the Bethe state (2.6) is an eigenstate of the transfer matrix.

Finally, let us investigate the polynomiality of

$$Q_{0,2} = Q_{0,1}^+ + Q_{0,1}^- + P'^{++}Q'^{-} - P'^{-}Q'^{++}. \quad (2.39)$$

Since  $Q_{0,1}$  is a polynomial<sup>2</sup>, we investigate the regularity of the remaining part at  $u = 0$ . Since  $Q(\pm \frac{i}{2}) = 0$  we can see that

$$Q'^{-}(0) = -Q(-\frac{3i}{2}), \quad Q'^{++}(0) = Q(\frac{3i}{2}), \quad (2.40)$$

which are not zero. We now focus on the pole contributions at  $u = 0$ . They can only come from the terms proportional to  $p(u)$ , which have poles at  $u = -i(n + \frac{1}{2})$  for any integer  $n \geq 0$ , with residues  $-1$ . Thus the singular parts can arise as

$$\begin{aligned} P'^{++}(\epsilon) &= P(\frac{3i}{2} + \epsilon) + \dots = \frac{1}{2}(a_+ + a_-)Q(\frac{3i}{2})(p^{++++}(\epsilon) + p^+(\epsilon)) = 0 + \dots \\ P'^{-}(\epsilon) &= -P(-\frac{3i}{2} + \epsilon) + \dots = -\frac{1}{2}(a_+ + a_-)Q(-\frac{3i}{2})(p^{-----}(\epsilon) + p^-(\epsilon)) \\ &= \frac{1}{\epsilon}(a_+ + a_-)Q(-\frac{3i}{2}) + \dots, \end{aligned} \quad (2.41)$$

where we have omitted regular terms in  $\epsilon$ . The singular part of  $Q_{0,2}(\epsilon)$  is then  $-\frac{1}{\epsilon}(a_+ + a_-)Q(\frac{3i}{2})Q(-\frac{3i}{2})$  whose vanishing implies  $a_+ = -a_-$ , i.e. the polynomiality of  $P$ .

We can thus conclude that the following four properties are equivalent:

- (i)  $P$  is a polynomial
- (ii) all  $Q_{a,s}$ -functions are polynomial
- (iii) the roots  $\{u_1, \dots, u_M\}$  are admissible solutions of the BA equations
- (iv) the Bethe vector is an eigenvector of the transfer matrix

---

<sup>2</sup>Recall from (2.19) that polynomiality of  $Q_{0,1}$  is equivalent to polynomiality of  $T$ ; and the latter is evident from

$$T = (u^+)^{N-1}u^{---}\frac{\bar{Q}^{--}}{\bar{Q}} + (u^-)^{N-1}u^{+++}\frac{\bar{Q}^{++}}{\bar{Q}},$$

which follows from the  $TQ$ -relation (2.8) and (2.24), and which has vanishing residues at the zeros of  $\bar{Q}$  by virtue of the BA equations.



### 3 Closed XXZ $QQ$ -system

In this section we present a generalization of the  $QQ$ -system for the closed XXZ spin chain of length  $N$  with periodic boundary conditions, whose Hamiltonian is given by

$$\mathbb{H} = \sum_{k=1}^N \left[ \sigma_k^x \sigma_{k+1}^x + \sigma_k^y \sigma_{k+1}^y + \frac{1}{2}(q + q^{-1}) \sigma_k^z \sigma_{k+1}^z \right], \quad \vec{\sigma}_{N+1} \equiv \vec{\sigma}_1. \quad (3.1)$$

We show that there is a notion of polynomial solutions of the  $QQ$ -equations, which determine the spectrum of the closed XXZ model.

#### 3.1 Review of the algebraic BA solution

The XXZ spin chain is related to the trigonometric generalization of the rational  $R$ -matrix (2.2):<sup>3</sup>

$$\mathbb{R}(u) = \begin{pmatrix} \sinh(u + \frac{\eta}{2}) & 0 & 0 & 0 \\ 0 & \sinh(u - \frac{\eta}{2}) & \sinh(\eta) & 0 \\ 0 & \sinh(\eta) & \sinh(u - \frac{\eta}{2}) & 0 \\ 0 & 0 & 0 & \sinh(u + \frac{\eta}{2}) \end{pmatrix} \quad (3.2)$$

The monodromy and transfer matrices can be introduced by the analogous formulae to the XXX case (2.3). The off-diagonal elements of the monodromy matrix (2.6) can be used as creation and annihilation operators. The  $\mathbb{B}$  operators, by acting on the all spin-up reference state (2.5), create eigenstates of the transfer matrix. The eigenvalue  $T(u)$  of the transfer matrix  $\mathbb{T}(u)$  satisfies the  $TQ$ -relation

$$T(u) Q(u) = \sinh^N(u + \frac{\eta}{2}) Q(u - \eta) + \sinh^N(u - \frac{\eta}{2}) Q(u + \eta), \quad (3.3)$$

where now

$$Q(u) = \prod_{j=1}^M \sinh(u - u_j). \quad (3.4)$$

Alternatively, we can switch to the variable  $t = e^u$ . By construction,  $T(u)$  is a polynomial of  $t$  and  $t^{-1}$ , regular at  $t_j = e^{u_j}$ , which implies the BA equations

$$\left( \frac{\sinh(u_k + \frac{\eta}{2})}{\sinh(u_k - \frac{\eta}{2})} \right)^N = - \prod_{j=1}^M \frac{\sinh(u_k - u_j + \eta)}{\sinh(u_k - u_j - \eta)}, \quad k = 1, \dots, M. \quad (3.5)$$

Singular BA solutions appear also for the XXZ spin chain, and the admissibility of the solution  $\{-\frac{\eta}{2}, \frac{\eta}{2}, u_1, \dots, u_{M-2}\}$  with pairwise distinct and finite roots can be formulated similarly to the XXX case as [16]

$$\frac{\bar{Q}(+\frac{\eta}{2})\bar{Q}(+\frac{3\eta}{2})}{\bar{Q}(-\frac{\eta}{2})\bar{Q}(-\frac{3\eta}{2})} = (-1)^N, \quad (3.6)$$

---

<sup>3</sup>The XXX limit can be recovered by the rescalings  $u \rightarrow \epsilon u$ ,  $\eta \rightarrow i\epsilon$  and by then taking the  $\epsilon \rightarrow 0$  limit.

where  $\bar{Q}(u) = \prod_{j=1}^{M-2} \sinh(u - u_j)$ .

In the following we introduce a  $QQ$ -system whose polynomial (in  $t$  and  $t^{-1}$ ) solutions account for the physical solutions. We work for generic  $\eta$ , i.e. when  $q = e^\eta$  is not a root of unity.

### 3.2 $QQ$ -system and its solution

For given  $N, M$ , the nontrivial  $Q$ -functions are defined on the same Young tableau as in the XXX case, and with the same boundary conditions  $Q_{2,s} = 1$ ,  $Q_{1,s>M} = 1$ , see Fig. 1. However, we now regard the  $Q$ -functions as functions of the argument  $t = e^u$ . Moreover, shifts now denote  $f^\pm(t) = f(tq^{\pm\frac{1}{2}})$ , and  $QQ$ -equations are formulated around each face as

$$Q_{a+1,s}(t) Q_{a,s+1}(t) \propto Q_{a+1,s+1}^+(t) Q_{a,s}^-(t) - Q_{a+1,s+1}^-(t) Q_{a,s}^+(t). \quad (3.7)$$

The initial conditions are

$$Q_{0,0}(t) = (t - t^{-1})^N, \quad Q_{1,0}(t) = Q(t) = \prod_{j=1}^M (tt_j^{-1} - t^{-1}t_j). \quad (3.8)$$

Both of these  $Q$ -functions are polynomial in the variables  $t$  and  $t^{-1}$ . As in the XXX case, we introduce the analogue of  $P$ , and then proceed to express all  $Q$ -functions in terms of  $P$  and  $Q$ .

The  $QQ$ -equation for  $(a, s) = (1, 0)$  leads again to the discrete derivative of  $Q$ :

$$Q_{1,1}(t) = Q_{1,0}^+(t) - Q_{1,0}^-(t) \equiv Q'_{1,0}(t) = Q'(t). \quad (3.9)$$

However, contrary to the XXX case, the order of  $Q_{1,1}$  is the same as that of  $Q$ . Since the  $QQ$ -equations for the XXZ case (3.7) are the same as for the XXX case (2.12), the solutions are the same, too. In particular, formulas such as (2.17)-(2.19) are exactly the same, and polynomiality of  $Q_{0,1}$  is equivalent to the polynomiality of  $T$ , which gives the BA equations (3.5). In proceeding as before, we search for a function  $P$  that satisfies

$$Q_{0,0} = P^+ Q^- - P^- Q^+, \quad (3.10)$$

that is,

$$(t - t^{-1})^N = P(tq^{\frac{1}{2}}) Q(tq^{-\frac{1}{2}}) - P(tq^{-\frac{1}{2}}) Q(tq^{\frac{1}{2}}). \quad (3.11)$$

With this  $P$  and  $Q$ , all  $Q$ -functions can be written as in the XXX case (2.23)

$$Q_{1,n} = Q^{(n)}, \quad Q_{0,n} \propto P^{(n)+} Q^{(n)-} - P^{(n)-} Q^{(n)+}, \quad (3.12)$$

except that the superscript  $(n)$  denotes the  $n^{\text{th}}$  discrete derivative obtained from multiplicative shifts in  $t$ , with  $f^\pm(t) = f(tq^{\pm\frac{1}{2}})$  and  $f'(t) = f^+(t) - f^-(t)$ .

The construction of the function  $P$ , once written in terms of the shifts, literally repeats the steps in the XXX case. One first shows that the only singular solutions are  $t = q^{\pm\frac{1}{2}}$ .

One then separates the singular solutions as  $Q(t) = (t - t^{-1})^+(t - t^{-1})^-\bar{Q}(t)$ , and performs a partial fraction decomposition of  $R(t)$  as

$$R(t) = \frac{(t - t^{-1})^N}{Q^+Q^-} = \pi + \frac{r_+}{\bar{Q}^+} + \frac{r_-}{\bar{Q}^-} + \frac{a_+}{(t - t^{-1})^{++}} + \frac{a_-}{(t - t^{-1})^{--}}, \quad (3.13)$$

where  $\pi(t)$  is polynomial. From the singularity structure, one can obtain  $r_+ = r^+$  and  $r_- = -r^-$  for some polynomial function  $r(t)$ , together with

$$a_{\pm} = \pm \frac{2^N T(q^{\mp \frac{1}{2}})}{(q - q^{-1})^2 (q^2 - q^{-2})} \frac{1}{\bar{Q}(q^{\pm \frac{1}{2}}) \bar{Q}(q^{\mp \frac{3}{2}})}. \quad (3.14)$$

The important new step now is the discrete integration of  $(t - t^{-1})^{-1} = \frac{1}{2}(\frac{1}{t-1} + \frac{1}{t+1})$ . To this end, we define the function  $p_q(t)$  by

$$p'_q(t) = \frac{1}{t - t^{-1}}, \quad p_q(t) = \frac{1}{2 \log q} \left\{ \psi_{q^{-1}} \left( \frac{\log t}{\log q} + \frac{1}{2} \right) - \psi_{q^{-1}} \left( \frac{\log(-t)}{\log q} + \frac{1}{2} \right) \right\}, \quad (3.15)$$

where  $\psi_q(x)$  denotes the  $q$ -deformed digamma function [17, 18], which satisfies

$$\psi_q(x+1) - \psi_q(x) = \frac{\log q}{1 - q^{-x}}. \quad (3.16)$$

The function  $P$  finally takes the form

$$P = \rho Q + (t - t^{-1})^+(t - t^{-1})^- r + \frac{1}{2}(a_+ - a_-)(t - t^{-1})(q^{\frac{1}{2}} - q^{-\frac{1}{2}})\bar{Q} + \frac{1}{2}(a_+ + a_-)(p_q^{++} + p_q^{--})Q. \quad (3.17)$$

Polynomiality of  $P$  (in  $t, t^{-1}$ ) requires  $a_+ = -a_-$ , which is equivalent to the admissibility of the Bethe roots (3.6). Investigating the polynomiality of the function  $Q_{0,2}$ , one discovers that it is also equivalent to the  $a_+ = -a_-$  condition. Thus, similarly to the XXX case, the following statements are equivalent: (i)  $P$  is a polynomial, (ii) all  $Q_{a,s}$ -functions are polynomial, (iii) the roots  $\{u_1, \dots, u_M\}$  are admissible solutions of the BA equations, (iv) the Bethe vector is an eigenvector of the transfer matrix.

Let us briefly comment on the root of unity case  $q = e^{i\pi/p}$  with integer  $p \geq 2$ . In this case, another exact string besides  $\pm\eta/2$  becomes possible, namely, a complete string of length  $p$  [19, 20, 21, 22, 23], due to the periodicity of  $\sinh$  in the imaginary direction. Thus the construction of the function  $P$  would involve more  $q$ -deformed digamma functions located at the center of these new exact strings, and the polynomiality of  $P$  would be equivalent to the cancellation of multiple constants. It would a priori require more work to show that their cancellation are equivalent to the  $QQ$  relations, as we should expect.

## 4 Open XXX $QQ$ -system

We turn now to the open XXX spin chain of length  $N$ , with Hamiltonian

$$\mathbb{H} = \sum_{k=1}^{N-1} \vec{\sigma}_k \cdot \vec{\sigma}_{k+1}, \quad (4.1)$$

which is  $SU(2)$  invariant. After reviewing its BA solution, we propose a corresponding  $QQ$ -system, and argue that all the  $Q$ 's are polynomial if and only if the Bethe state is an eigenstate of the transfer matrix.

## 4.1 Review of the algebraic BA solution

The transfer matrix  $\mathbb{T}(u)$  is given by [24]

$$\mathbb{T}(u) = \text{tr}_0 \mathbb{U}_0(u), \quad \mathbb{U}_0(u) = \mathbb{M}_0(u) \widehat{\mathbb{M}}_0(u), \quad (4.2)$$

where  $\mathbb{M}_0(u)$  is the monodromy matrix in (2.3), and  $\widehat{\mathbb{M}}_0(u)$  is given by

$$\widehat{\mathbb{M}}(u) = \mathbb{R}_{0N}(u) \cdots \mathbb{R}_{02}(u) \mathbb{R}_{01}(u). \quad (4.3)$$

The  $R$ -matrix is again given by (2.2). Its boundary equivalent, the  $K$ -matrix, is the identity in the case considered here. By construction, the open-chain transfer matrix (4.2) has the commutativity property

$$[\mathbb{T}(u), \mathbb{T}(v)] = 0, \quad (4.4)$$

and it also has the crossing symmetry

$$\mathbb{T}(-u) = \mathbb{T}(u). \quad (4.5)$$

The Hamiltonian (4.1) is proportional to  $\left. \frac{d\mathbb{T}(u)}{du} \right|_{u=i/2}$ , up to an additive constant.

We denote the matrix elements of  $\mathbb{U}_0(u)$  (4.2) as follows

$$\mathbb{U}_0(u) = \begin{pmatrix} \mathbb{A}(u) & \mathbb{B}(u) \\ \mathbb{C}(u) & \frac{u^-}{u} \mathbb{D}(u) + \frac{i}{2u} \mathbb{A}(u) \end{pmatrix}. \quad (4.6)$$

The reference state (2.5) is annihilated by  $\mathbb{C}(u)$ , and is an eigenstate of  $\mathbb{A}(u)$  and  $\mathbb{D}(u)$ , with

$$\mathbb{A}(u)|0\rangle = (u^+)^{2N}|0\rangle, \quad \mathbb{D}(u)|0\rangle = (u^-)^{2N}|0\rangle. \quad (4.7)$$

The Bethe states are defined by

$$|u_1 \dots u_M\rangle = \prod_{k=1}^M \mathbb{B}(u_k)|0\rangle. \quad (4.8)$$

The Bethe states satisfy (for any  $\{u_1, \dots, u_M\}$ ) the off-shell relation

$$\mathbb{T}(u)|u_1 \dots u_M\rangle = T(u)|u_1 \dots u_M\rangle + \sum_{j=1}^M F_j |u, u_1 \dots \hat{u}_j \dots u_M\rangle, \quad (4.9)$$

where  $\hat{u}_j$  is omitted. Moreover,  $T(u)$  is given by the  $TQ$ -relation

$$u T(u) Q(u) = (u^+)^{2N+1} Q^{--}(u) + (u^-)^{2N+1} Q^{++}(u), \quad Q(u) = \prod_{k=1}^M (u - u_k)(u + u_k), \quad (4.10)$$

and the coefficients  $F_j$  of the “unwanted” terms are given by

$$F_j = \frac{2iu^+(u_j - \frac{i}{2})}{u_j(u - u_j)(u + u_j)} \left[ (u_j + \frac{i}{2})^{2N} \prod_{\substack{k \neq j \\ k=1}}^M \frac{(u_j - u_k - i)(u_j + u_k - i)}{(u_j - u_k)(u_j + u_k)} \right. \\ \left. - (u_j - \frac{i}{2})^{2N} \prod_{\substack{k \neq j \\ k=1}}^M \frac{(u_j - u_k + i)(u_j + u_k + i)}{(u_j - u_k)(u_j + u_k)} \right]. \quad (4.11)$$

We again write  $f^\pm(u) = f(u \pm \frac{i}{2})$ , as in the closed XXX case. Note that both  $Q$  and  $T$  are even functions of  $u$

$$Q(-u) = Q(u), \quad T(-u) = T(u). \quad (4.12)$$

Substituting  $u = u_j$  in the  $TQ$ -relation (4.10), we see that the LHS vanishes, and we obtain

$$(u_j + \frac{i}{2})^{2N+1} \prod_{k=1}^M (u_j - u_k - i)(u_j + u_k - i) + (u_j - \frac{i}{2})^{2N+1} \prod_{k=1}^M (u_j - u_k + i)(u_j + u_k + i) = 0, \quad j = 1, \dots, M. \quad (4.13)$$

If  $u_j \neq \pm \frac{i}{2}$ , then these equations are equivalent to the BA equations

$$\left( \frac{u_j + \frac{i}{2}}{u_j - \frac{i}{2}} \right)^{2N} = \prod_{\substack{k \neq j \\ k=1}}^M \frac{(u_j - u_k + i)(u_j + u_k + i)}{(u_j - u_k - i)(u_j + u_k - i)}, \quad j = 1, \dots, M. \quad (4.14)$$

The BA equations have the reflection symmetry  $u_j \mapsto -u_j$ , while keeping the other  $u$ 's (i.e.  $u_k$  with  $k \neq j$ ) unchanged. Hence, without loss of generality, we henceforth assume that  $\Re(u_j) > 0$ , or  $\Re(u_j) = 0$  and  $\Im(u_j) \geq 0$ .

Roughly speaking, if  $\{u_1, \dots, u_M\}$  satisfy the BA equations (4.14), then all  $F_j = 0$ ; i.e., the “unwanted” terms in the off-shell relation (4.9) vanish, hence the Bethe state (4.8) is an eigenstate of the transfer matrix, with corresponding eigenvalue  $T(u)$ . However, there are some important caveats. We argue in Appendix A that certain “exceptional” solutions of the BA equations (namely 0 and  $\pm \frac{i}{2}$ ) do *not* lead to eigenstates of the transfer matrix. Moreover, we make the standard assumption (supported by numerical evidence, see e.g. [23]) that the Bethe roots are pairwise distinct, i.e.  $u_j \neq u_k$  if  $j \neq k$ .

We therefore define an *admissible* solution  $\{u_1, \dots, u_M\}$  of the BA equations (4.14), such that all the  $u_j$ 's are finite, not equal to  $\pm \frac{i}{2}$  or 0, and pairwise distinct (no two are equal), and each  $u_j$  satisfies either

$$\Re(u_j) > 0 \quad (4.15)$$

or

$$\Re(u_j) = 0 \quad \text{and} \quad \Im(u_j) > 0. \quad (4.16)$$

The set  $\{u_1, \dots, u_M\}$  is an admissible solution of the BA equations if and only if the Bethe state  $|u_1 \dots u_M\rangle$  is an eigenstate of the transfer matrix  $\mathbb{T}(u)$ . We emphasize that, for the open XXX chain, there are no physical singular solutions of the Bethe equations – all the singular solutions are unphysical.

## 4.2 $QQ$ -system

We propose the following  $QQ$ -system

$$u Q_{a+1,s}(u) Q_{a,s+1}(u) \propto Q_{a+1,s+1}^+(u) Q_{a,s}^-(u) - Q_{a+1,s+1}^-(u) Q_{a,s}^+(u), \quad (4.17)$$

where the nontrivial  $Q$ -functions for given values of  $N$  and  $M$  are again defined on the Young tableau in Fig. 1, with the boundary conditions  $Q_{2,s} = 1$ ,  $Q_{1,s>M} = 1$ , and with the initial condition

$$Q_{0,0}(u) = u^{2N} \quad \text{with} \quad Q_{1,0}(u) = Q(u) = \prod_{k=1}^M (u - u_k)(u + u_k). \quad (4.18)$$

In contrast to the  $QQ$ -system for periodic XXX (2.12), there is an extra factor of  $u$  on the LHS of (4.17), and the  $Q$ 's are *even* functions of  $u$ . The degree of the polynomial  $Q_{a,s}(u)$  is *doubled* with respect to the periodic XXX case (namely, *twice* the number of boxes in the Young tableau to the right and top of the vertex  $(a, s)$ ).<sup>4</sup> We claim that all the  $Q$ 's are polynomial if and only if the Bethe state  $|u_1 \dots u_M\rangle$  (4.8) is an eigenstate of the transfer matrix  $\mathbb{T}(u)$  (4.2).

Before entering into the proof, let us quickly check that this  $QQ$ -system indeed leads to the correct BA equations for  $\{u_1, \dots, u_M\}$ . We write the  $QQ$ -equations for  $(a, s) = (0, 0)$ :

$$u Q_{1,0} Q_{0,1} \propto Q_{1,1}^+ Q_{0,0}^- - Q_{1,1}^- Q_{0,0}^+, \quad (4.19)$$

and for  $(a, s) = (1, 0)$ :

$$u Q_{2,0} Q_{1,1} \propto Q_{2,1}^+ Q_{1,0}^- - Q_{2,1}^- Q_{1,0}^+. \quad (4.20)$$

Since  $Q_{2,0} = Q_{2,1} = 1$ , the latter reduces to

$$u Q_{1,1} \propto Q^- - Q^+. \quad (4.21)$$

Performing the shifts  $u \mapsto u \pm \frac{i}{2}$  in (4.21) and evaluating at  $u = u_j$ , we obtain

$$(u_j + \frac{i}{2}) Q_{1,1}^+(u_j) \propto -Q^{++}(u_j), \quad (u_j - \frac{i}{2}) Q_{1,1}^-(u_j) \propto Q^{--}(u_j), \quad (4.22)$$

since  $Q(u_j) = 0$ . Moreover, evaluating (4.19) at  $u = u_j$  gives

$$Q_{1,1}^+(u_j) Q_{0,0}^-(u_j) = Q_{1,1}^-(u_j) Q_{0,0}^+(u_j), \quad (4.23)$$

Substituting (4.22) into the above relation gives

$$-(u_j - \frac{i}{2}) Q^{++}(u_j) Q_{0,0}^-(u_j) = (u_j + \frac{i}{2}) Q^{--}(u_j) Q_{0,0}^+(u_j), \quad (4.24)$$

which coincides with the BA equations (4.13).

---

<sup>4</sup>Equivalently, the  $Q$ -functions are polynomials in the variable  $u^2$  of the same degree as in the periodic XXX case.

### 4.2.1 $Q_{a,s}$ in terms of $Q$ and $P$

We now solve the  $QQ$ -system (4.17) in terms of  $Q(u)$  and a function  $P(u)$ , such that polynomiality of  $P(u)$  implies polynomiality of all the  $Q$ 's. We define  $P(u)$  by <sup>5</sup>

$$P^+Q^- - P^-Q^+ = u Q_{0,0}, \quad (4.25)$$

where  $Q_{0,0}$  is given by (4.18). It follows from (4.19) and (4.21) that

$$Q_{1,1} \propto \frac{Q'}{u} = DQ, \quad u Q_{0,1} \propto (DP)^+(DQ)^- - (DP)^-(DQ)^+, \quad (4.26)$$

where we have used the following compact notation for discrete derivatives with certain  $1/u$  factors

$$\begin{aligned} Df &= \frac{1}{u}(f^+ - f^-) = \frac{f'}{u}, \\ D^2f &= \frac{1}{u}[(Df)^+ - (Df)^-], \dots \\ D^n f &= \frac{1}{u}[(D^{n-1}f)^+ - (D^{n-1}f)^-]. \end{aligned} \quad (4.27)$$

Similarly, we obtain

$$Q_{1,2} \propto D^2Q, \quad u Q_{0,2} \propto (D^2P)^+(D^2Q)^- - (D^2P)^-(D^2Q)^+, \quad (4.28)$$

and in general

$$Q_{1,n} \propto D^n Q, \quad u Q_{0,n} \propto (D^n P)^+(D^n Q)^- - (D^n P)^-(D^n Q)^+. \quad (4.29)$$

Since both  $Q(u)$  and  $P(u)$  are even functions of  $u$ , it follows that  $DQ$  and  $DP$  are also even functions of  $u$ . Hence, if  $P(u)$  is a polynomial function of  $u$ , then the RHS of the second equation in (4.26) is divisible by  $u$ , thus  $Q_{0,1}$  is polynomial; and, from (4.29), we similarly conclude that all the  $Q$ 's are polynomial.

We observe, similarly to the closed-chain case, that the  $TQ$ -equation (4.10) together with the definition of  $P$  (4.25) imply

$$uT = P^{++}Q^{--} - P^{--}Q^{++}. \quad (4.30)$$

It follows that  $P$  is also a solution of the  $TQ$ -equation

$$uTP = (u^+)^{2N+1}P^{--} + (u^-)^{2N+1}P^{++}. \quad (4.31)$$

Hence, (4.25) can be regarded as the Wronskian relation obeyed by the two solutions  $Q$  and  $P$  of the  $TQ$ -equation (4.10).

---

<sup>5</sup>As in the closed-chain case, for given  $Q$ , this equation does not uniquely define  $P$ : if  $P(u)$  is a solution of (4.25), then so is  $P(u) + \alpha Q(u)$ , for any constant value of  $\alpha$ .

### 4.2.2 Construction of $P$

We now construct the function  $P(u)$  for a set  $\{u_1, \dots, u_M\}$ , and argue that  $P(u)$  is polynomial if and only if  $\{u_1, \dots, u_M\}$  is an admissible solution of the BA equations.

The construction of the  $P$ -function for the open chain is similar to that for the closed chain, but with some significant differences. In the presence of one singular root  $\frac{i}{2}$  and one zero root 0, the  $Q$ -function takes the form <sup>6</sup>

$$Q(u) = u^+ u^- u^2 \bar{Q}(u), \quad \bar{Q}(u) = \prod_{\substack{u_k \neq \frac{i}{2}, 0 \\ k=1}}^M (u - u_k)(u + u_k). \quad (4.32)$$

We define the function  $R(u)$

$$R = \frac{u^{2N+1}}{Q^+ Q^-} = \frac{u^{2N-1}}{u^{++} u^{--} (u^+ u^-)^2 \bar{Q}^+ \bar{Q}^-}, \quad (4.33)$$

which is related to  $P(u)$  defined in (4.25) by

$$R = \left( \frac{P}{Q} \right)' = \frac{P^+}{Q^+} - \frac{P^-}{Q^-}. \quad (4.34)$$

Decomposing (4.33) in partial fractions, we obtain

$$R = \pi + \frac{q_+}{\bar{Q}^+} + \frac{q_-}{\bar{Q}^-} + \frac{a_+}{u^{++}} + \frac{b_+}{u^+} + \frac{c_+}{(u^+)^2} + \frac{a_-}{u^{--}} + \frac{b_-}{u^-} + \frac{c_-}{(u^-)^2}, \quad (4.35)$$

where  $\pi$  is a polynomial of order  $2N - 4M + 1$ ,  $q_{\pm}$  are polynomials of degree less than that of  $\bar{Q}$ , and  $a_{\pm}, b_{\pm}, c_{\pm}$  are constants. Note that  $a_{\pm}$  arise from the presence of the singular root, while  $b_{\pm}, c_{\pm}$  are due to the presence of the zero root. From the  $TQ$ -relation (4.10), we obtain

$$R^+ + R^- = \frac{uT}{Q^{++} Q^{--}} = \frac{uT}{u^+ u^- u^{+++} u^{---} (u^{++} u^{--})^2 \bar{Q}^{++} \bar{Q}^{--}}. \quad (4.36)$$

We now evaluate the LHS of (4.36) using (4.35), and consider the values of  $u$  where singularities could arise. The RHS of (4.36) has no singularities at the zeros of  $\bar{Q}$  (recall that  $T(u)$  is regular for values of  $u$  corresponding to admissible Bethe roots), hence

$$q_+ = q^+, \quad q_- = -q^-, \quad (4.37)$$

for some polynomial  $q(u)$ . From the residues of (4.36) at  $u = \mp \frac{i}{2}$ , we obtain

$$a_{\pm} = \frac{4T(\mp \frac{i}{2})}{9\bar{Q}(\pm \frac{i}{2})\bar{Q}(\mp \frac{3i}{2})} = \frac{8(-1)^{N+1}}{9\bar{Q}(\pm \frac{i}{2})\bar{Q}(\mp \frac{3i}{2})}. \quad (4.38)$$

---

<sup>6</sup>The cases of either one singular root or one zero root are essentially special cases, for which  $b_{\pm} = c_{\pm} = 0$  or  $a_{\pm} = 0$  in (4.35) below, respectively.



Since  $Q(u)$  is an even function of  $u$ , we conclude – in significant contrast from the closed-chain case – that  $a_+ = a_- \equiv a$ . From the residues of (4.36) at  $u = \mp i$ , we can obtain expressions for  $c_{\pm}$  and  $b_{\pm}$ , and we find that

$$c \equiv c_+ = -c_-, \quad b \equiv b_+ = b_-, \quad (4.39)$$

which is consistent with the constraints coming from the residues of (4.36) at  $u = 0$  (note that the presence of a zero root implies that  $T(u)$  has a double pole at  $u = 0$ ).

We write the polynomial  $\pi$  in (4.35) as

$$\pi = \rho' = \rho^+ - \rho^-, \quad (4.40)$$

where  $\rho$  is a polynomial. Recalling the definition of the function  $p(u)$  (2.32), we see that

$$R = \left( \rho + \frac{q}{Q} + a(p^{++} + p^{--}) + b(p^+ + p^-) + \frac{c}{u^2} \right)'. \quad (4.41)$$

It immediately follows from (4.34) that the  $P$ -function is given by

$$P = \rho Q + u^+ u^- u^2 q + c u^+ u^- \bar{Q} + a(p^{++} + p^{--})Q + b(p^+ + p^-)Q, \quad (4.42)$$

which is a polynomial iff  $a = b = 0$ . That is, the  $P$ -function is polynomial iff there is no singular root and no zero root, in which case  $\{u_1, \dots, u_M\}$  is admissible. Moreover,  $Q_{0,2}$  is polynomial iff  $a = b = 0$ .

The proof of the  $QQ$ -system (4.17) is now complete, since we have argued as before that the following statements are equivalent: (i)  $P$  is a polynomial, (ii) all  $Q_{a,s}$ -functions are polynomial, (iii) the roots  $\{u_1, \dots, u_M\}$  are admissible solutions of the BA equations, (iv) the Bethe vector is an eigenvector of the transfer matrix.

## 5 Open quantum-group-invariant XXZ $QQ$ -system

We now generalize the preceding results to the open quantum-group-invariant XXZ spin chain, whose Hamiltonian is given by [8]

$$\mathbb{H} = \sum_{k=1}^{N-1} \left[ \sigma_k^x \sigma_{k+1}^x + \sigma_k^y \sigma_{k+1}^y + \frac{1}{2}(q + q^{-1})\sigma_k^z \sigma_{k+1}^z \right] - \frac{1}{2}(q - q^{-1})(\sigma_1^z - \sigma_N^z). \quad (5.1)$$

For simplicity, we restrict to generic values of  $q = e^\eta$ .

### 5.1 Review of the algebraic BA solution

The transfer matrix is now given by [24]

$$\mathbb{T}(u) = \text{tr}_0 \mathbb{K}_0^L(u) \mathbb{U}_0(u), \quad \mathbb{U}_0(u) = \mathbb{M}_0(u) \mathbb{K}_0^R(u) \widehat{\mathbb{M}}_0(u), \quad (5.2)$$

where the  $R$ -matrix is again given by (3.2), and the left and right  $K$ -matrices (solutions of boundary Yang-Baxter equations) are given by the diagonal matrices

$$\mathbb{K}^L(u) = \text{diag}(e^{-u-\frac{\eta}{2}}, e^{u+\frac{\eta}{2}}), \quad \mathbb{K}^R(u) = \text{diag}(e^{u-\frac{\eta}{2}}, e^{-u+\frac{\eta}{2}}). \quad (5.3)$$

The transfer matrix (5.2) has the commutativity property (4.4) as well as the crossing symmetry (4.5). The Hamiltonian (5.1) is proportional to  $d\mathbb{T}(u)/du|_{u=\eta/2}$ , up to an additive constant.

We define the elements of  $\mathbb{U}_0(u)$  (5.2) as follows

$$\mathbb{U}_0(u) = \begin{pmatrix} e^{u-\frac{\eta}{2}}\mathbb{A}(u) & \mathbb{B}(u) \\ \mathbb{C}(u) & \frac{e^{-u-\frac{\eta}{2}} \sinh(2u-\eta)}{\sinh(2u)}\mathbb{D}(u) + \frac{e^{u-\frac{\eta}{2}} \sinh(\eta)}{\sinh(2u)}\mathbb{A}(u) \end{pmatrix}. \quad (5.4)$$

The reference state (2.5) is annihilated by  $\mathbb{C}(u)$ , and is an eigenstate of  $\mathbb{A}(u)$  and  $\mathbb{D}(u)$

$$\mathbb{A}(u)|0\rangle = \sinh^{2N}(u + \frac{\eta}{2})|0\rangle, \quad \mathbb{D}(u)|0\rangle = \sinh^{2N}(u - \frac{\eta}{2})|0\rangle. \quad (5.5)$$

The Bethe states are again defined by

$$|u_1 \dots u_M\rangle = \prod_{k=1}^M \mathbb{B}(u_k)|0\rangle. \quad (5.6)$$

The off-shell equation is

$$\mathbb{T}(u)|u_1 \dots u_M\rangle = T(u)|u_1 \dots u_M\rangle + \sum_{j=1}^M F_j|u, u_1 \dots \hat{u}_j \dots u_M\rangle, \quad (5.7)$$

where  $T(u)$  is given by the  $TQ$ -relation

$$\sinh(2u) T(u) Q(u) = \sinh(2u+\eta) \sinh^{2N}(u+\frac{\eta}{2}) Q(u-\eta) + \sinh(2u-\eta) \sinh^{2N}(u-\frac{\eta}{2}) Q(u+\eta), \quad (5.8)$$

with

$$Q(u) = \prod_{k=1}^M \sinh(u - u_k) \sinh(u + u_k), \quad (5.9)$$

and  $F_j$  is given by

$$F_j = \frac{\sinh(2u+\eta) \sinh(2u_j-\eta) \sinh(\eta)}{\sinh(2u_j) \sinh(u-u_j) \sinh(u+u_j)} \left[ \sinh^{2N}(u_j + \frac{\eta}{2}) \prod_{\substack{k \neq j \\ k=1}}^M \frac{\sinh(u_j - u_k - \eta) \sinh(u_j + u_k - \eta)}{\sinh(u_j - u_k) \sinh(u_j + u_k)} \right. \\ \left. - \sinh^{2N}(u_j - \frac{\eta}{2}) \prod_{\substack{k \neq j \\ k=1}}^M \frac{\sinh(u_j - u_k + \eta) \sinh(u_j + u_k + \eta)}{\sinh(u_j - u_k) \sinh(u_j + u_k)} \right]. \quad (5.10)$$

Substituting  $u = u_j$  in the  $TQ$ -equation (5.8), we see that the LHS vanishes, and we obtain

$$\begin{aligned} & \sinh(2u_j + \eta) \sinh^{2N}(u_j + \frac{\eta}{2}) \prod_{k=1}^M \sinh(u_j - u_k - \eta) \sinh(u_j + u_k - \eta) \\ & + \sinh(2u_j - \eta) \sinh^{2N}(u_j - \frac{\eta}{2}) \prod_{k=1}^M \sinh(u_j - u_k + \eta) \sinh(u_j + u_k + \eta) = 0, \quad j = 1, \dots, M. \end{aligned} \quad (5.11)$$

If  $u_j \neq \pm \frac{\eta}{2}$ , then these equations are equivalent to the BA equations

$$\left( \frac{\sinh(u_j + \frac{\eta}{2})}{\sinh(u_j - \frac{\eta}{2})} \right)^{2N} = \prod_{\substack{k=1 \\ k \neq j}}^M \frac{\sinh(u_j - u_k + \eta) \sinh(u_j + u_k + \eta)}{\sinh(u_j - u_k - \eta) \sinh(u_j + u_k - \eta)}, \quad j = 1, \dots, M. \quad (5.12)$$

The BA equations have the reflection symmetry  $u_j \mapsto -u_j$  (while keeping the other  $u$ 's unchanged), as well as the periodicity  $u_j \mapsto u_j + i\pi$ .

We must exclude solutions with roots  $0, \pm \frac{i\pi}{2}$  and  $\pm \frac{\eta}{2}$ , see Sec. A.2. We therefore define an *admissible* solution  $\{u_1, \dots, u_M\}$  of the BA equations (5.12), such that all the  $u_j$ 's are finite, not equal to  $0, \pm \frac{i\pi}{2}$  or  $\pm \frac{\eta}{2}$ , pairwise distinct, and each  $u_j$  satisfies either

$$\Re(u_j) > 0 \quad \text{and} \quad -\frac{\pi}{2} < \Im(u_j) \leq \frac{\pi}{2} \quad (5.13)$$

or

$$\Re(u_j) = 0 \quad \text{and} \quad 0 < \Im(u_j) < \frac{\pi}{2}, \quad (5.14)$$

The set  $\{u_1, \dots, u_M\}$  is an admissible solution of the BA equations if and only if the Bethe state  $|u_1 \dots u_M\rangle$  is an eigenstate of the transfer matrix  $\mathbb{T}(u)$ .

## 5.2 $QQ$ -system

We propose the following  $QQ$ -system<sup>7</sup>

$$(t^2 - t^{-2}) Q_{a+1,s}(t) Q_{a,s+1}(t) \propto Q_{a+1,s+1}^+(t) Q_{a,s}^-(t) - Q_{a+1,s+1}^-(t) Q_{a,s}^+(t), \quad (5.15)$$

where the nontrivial  $Q$ -functions for given values of  $N$  and  $M$  are again defined on the Young tableau in Fig. 1, with the boundary conditions  $Q_{2,s} = 1$ ,  $Q_{1,s>M} = 1$ , and with the initial condition

$$Q_{0,0}(t) = (t - t^{-1})^{2N} \quad \text{with} \quad Q_{1,0}(t) = Q(t) = \prod_{k=1}^M (tt_k^{-1} - t^{-1}t_k) (tt_k - t^{-1}t_k^{-1}). \quad (5.16)$$

In contrast to the periodic XXZ case (3.7), there is an extra factor  $(t^2 - t^{-2}) \propto \sinh(2u)$  on the LHS of (5.15). This  $QQ$ -system indeed leads to the BA equations (5.11), as can be seen by following the same logic (4.19) - (4.24) of the rational case. The degree of the polynomials is the same as in the open XXX case.

<sup>7</sup>As for the periodic XXZ case, here again  $t = e^u$ , and  $f^\pm(t) = f(tq^{\pm \frac{1}{2}})$ .

### 5.2.1 $Q_{a,s}$ in terms of $Q$ and $P$

We now define  $P$  by

$$P^+ Q^- - P^- Q^+ = (t^2 - t^{-2}) Q_{0,0}, \quad (5.17)$$

where  $Q_{0,0}$  is given by (5.16). Similarly to the rational case (4.29), we find that the  $QQ$ -system (5.15) implies that

$$Q_{1,n} \propto D^n Q, \quad (t^2 - t^{-2}) Q_{0,n} \propto (D^n P)^+ (D^n Q)^- - (D^n P)^- (D^n Q)^+, \quad (5.18)$$

where  $D$  is now defined by

$$\begin{aligned} Df &= \frac{1}{(t^2 - t^{-2})} (f^+ - f^-) = \frac{f'}{(t^2 - t^{-2})}, \\ D^2 f &= \frac{1}{(t^2 - t^{-2})} [(Df)^+ - (Df)^-], \dots \\ D^n f &= \frac{1}{(t^2 - t^{-2})} [(D^{n-1} f)^+ - (D^{n-1} f)^-], \end{aligned} \quad (5.19)$$

cf. (4.27). Note that  $P$  is also a solution of the  $TQ$ -relation (5.8)

$$\sinh(2u) T P = \sinh(2u + \eta) Q_{0,0}^+ P^{--} + \sinh(2u - \eta) Q_{0,0}^- P^{++}, \quad (5.20)$$

cf. (4.31).

### 5.2.2 Construction of $P$

The construction of  $P$  parallels the rational case. In the presence of one singular root ( $u = \frac{\eta}{2}$ ,  $t = q^{\frac{1}{2}}$ ) and one zero root ( $u = 0$ ,  $t = 1$ ; the case  $u = \frac{i\pi}{2}$ ,  $t = -1$  is similar), the  $Q$ -function becomes

$$Q(t) = (t - t^{-1})^+ (t - t^{-1})^- (t - t^{-1})^2 \bar{Q}(t). \quad (5.21)$$

We now define  $R(t)$  as

$$R(t) = \frac{(t^2 - t^{-2})(t - t^{-1})^{2N}}{Q^+ Q^-} = \frac{(t^2 - t^{-2})(t - t^{-1})^{2N-2}}{(t - t^{-1})^{++} (t - t^{-1})^{--} [(t - t^{-1})^+ (t - t^{-1})^-]^2 \bar{Q}^+ \bar{Q}^-}, \quad (5.22)$$

which is related to  $P$  (5.17) by  $R = (\frac{P}{Q})'$ . We decompose  $R$  as follows

$$\begin{aligned} R &= \pi + \frac{r_+}{Q^+} + \frac{r_-}{\bar{Q}^-} + \frac{a_+}{(t - t^{-1})^{++}} + \frac{b_+}{(t - t^{-1})^+} + \frac{c_+}{[(t - t^{-1})^+]^2} \\ &\quad + \frac{a_-}{(t - t^{-1})^{--}} + \frac{b_-}{(t - t^{-1})^-} + \frac{c_-}{[(t - t^{-1})^-]^2}, \end{aligned} \quad (5.23)$$

where  $\pi$  is polynomial. From the  $TQ$ -relation (5.8), we obtain

$$\begin{aligned} R^+ + R^- &= \frac{2^{2N}(t^2 - t^{-2})T}{Q^{++}Q^{--}} \\ &= \frac{2^{2N}(t^2 - t^{-2})T}{(t - t^{-1})^+(t - t^{-1})^-(t - t^{-1})^{++}(t - t^{-1})^{--}[(t - t^{-1})^{++}(t - t^{-1})^{--}]^2 \bar{Q}^{++}\bar{Q}^{--}}. \end{aligned} \quad (5.24)$$

From the singularity structure of this equation and the fact that  $Q(t)$  and  $T(t)$  are invariant under  $t \mapsto t^{-1}$ , we again obtain

$$r_+ = r^+, \quad r_- = -r^-, \quad (5.25)$$

where  $r(t)$  is a polynomial in  $t$  and  $t^{-1}$ , and

$$a_+ = a_- \equiv a, \quad c \equiv c_+ = -c_-, \quad b \equiv b_+ = b_-. \quad (5.26)$$

The expression (5.23) for  $R$  can therefore be rewritten in the form

$$R = \left( \rho + \frac{r}{\bar{Q}} + a(p_q^{++} + p_q^{--}) + b(p_q^+ + p_q^-) + \frac{c}{(t - t^{-1})^2} \right)', \quad (5.27)$$

where  $p_q(t)$  is defined in (3.15), cf. (4.41). Since  $R = (\frac{P}{Q})'$ , we conclude that  $P$  is given by

$$\begin{aligned} P &= \rho Q + (t - t^{-1})^+(t - t^{-1})^-(t - t^{-1})^2 r + c(t - t^{-1})^+(t - t^{-1})^-\bar{Q} \\ &\quad + a(p_q^{++} + p_q^{--})Q + b(p_q^+ + p_q^-)Q, \end{aligned} \quad (5.28)$$

which is a polynomial in  $t$  and  $t^{-1}$  iff  $a = b = 0$ . That is, as in the rational case,  $P$  is polynomial iff there is no singular root and no zero root, in which case  $\{u_1, \dots, u_M\}$  is admissible. Moreover,  $Q_{0,2}$  is polynomial iff  $a = b = 0$ .

## 6 Conclusions

Our main results are  $QQ$ -systems for the closed XXZ (3.7)-(3.8), open XXX (4.17)-(4.18) and open quantum-group-invariant XXZ (5.15)-(5.16) quantum spin chains. Polynomial solutions of these  $QQ$ -systems can be found efficiently, which in turn lead directly to the admissible solutions of the corresponding BA equations.

Numerous applications of these results are possible. In conjunction with techniques from algebraic geometry, these  $QQ$ -systems allow the exact computation of partition functions for trigonometric vertex models and for vertex models with boundaries [6].

We restricted here to open spin chains with  $SU(2)$  or  $U_q(SU(2))$  symmetry. It would be both interesting and useful to formulate  $QQ$ -systems for open spin chains with other integrable boundary conditions, as well as for integrable models based on  $R$ -matrices for higher-rank algebras and/or higher-dimensional representations.

# Acknowledgments

ZB and RN are grateful for the warm hospitality extended to them at the University of Miami and the Wigner Research Center, respectively. ZB was supported in part by NKFIH grant K116505. The work of EG and JLJ was supported by the European Research Council under the Advanced Grant NuQFT. RN was supported in part by a Cooper fellowship.

## A Exceptional solutions for open spin chains

### A.1 XXX

#### A.1.1 $u_1 = 0$

Let us consider a solution  $u_1, \dots, u_M$  of the BA equations (4.14) with one zero Bethe root, say  $u_1 = 0$  (and  $u_2, \dots, u_M$  pairwise distinct and not equal to 0). Since  $F_1$  in (4.11) is not well-defined, we set  $u_1 = \epsilon$  and consider the limit  $\lim_{\epsilon \rightarrow 0} F_1$ . It is straightforward to see that this limit exists and is nonzero. Hence, the corresponding Bethe state in the off-shell equation (4.9) is *not* an eigenstate of the transfer matrix, see also [25].

#### A.1.2 $u_1 = \frac{i}{2}$

The equations (4.13) evidently have solutions  $u_1, \dots, u_M$  with one “singular” root, say  $u_1 = \frac{i}{2}$  (and  $u_2, \dots, u_M$  pairwise distinct and not equal to  $\pm \frac{i}{2}$ ). However, the BA equations (4.14) do not have such solutions (recall that the latter equations are not equivalent to (4.13) for this case). Hence, it is not surprising that the corresponding Bethe state is not an eigenstate of the transfer matrix. Indeed, let us define a renormalized B-operator

$$\tilde{\mathbb{B}}(u) = \frac{1}{u^-} \mathbb{B}(u), \quad (\text{A.1})$$

such that  $\lim_{\epsilon \rightarrow 0} \tilde{\mathbb{B}}(\frac{i}{2} + \epsilon)$  is finite and non-singular.<sup>8</sup> Bethe states created with this renormalized operator satisfy an off-shell relation similar to (4.9), except with  $F_j$  replaced by

$$\tilde{F}_j = \frac{u^-}{u_j - \frac{i}{2}} F_j. \quad (\text{A.2})$$

We find that  $\lim_{\epsilon \rightarrow 0} \tilde{F}_1$  exists and is nonzero for  $u_1 = \pm \frac{i}{2} + \epsilon$ . Hence, the corresponding Bethe state is *not* an eigenstate of the transfer matrix.

The BA equations (4.14) do have solutions with a pair of singular Bethe roots, e.g.  $u_1 = \frac{i}{2}$  and  $u_2 = \pm \frac{i}{2}$ , which must be discarded since  $|u_1|$  and  $|u_2|$  are not distinct.

---

<sup>8</sup>We have checked this explicitly for small values of  $N$ , and we expect that it can be proved by induction in  $N$ , similarly to the case of the closed chain [12].

## A.2 XXZ

For the XXZ case, the coefficients  $F_j$  of the “unwanted” terms are given by (5.10). We exclude both  $u_j = 0$  and  $u_j = \frac{i\pi}{2}$ , since (similarly to Sec. A.1.1) the limit  $\lim_{\epsilon \rightarrow 0} F_1$  exists and is nonzero for both  $u_1 = \epsilon$  and  $u_1 = \frac{i\pi}{2} + \epsilon$ . We must also exclude  $u_j = \pm \frac{\eta}{2}$ : similarly to Sec. A.1.2, we renormalize the B-operator

$$\tilde{\mathbb{B}}(u) = \frac{1}{\sinh(2u - \eta)} \mathbb{B}(u), \quad (\text{A.3})$$

so that  $\lim_{\epsilon \rightarrow 0} \tilde{\mathbb{B}}(\frac{\eta}{2} + \epsilon)$  is finite and non-singular. The coefficients of the “unwanted” terms become

$$\tilde{F}_j = \frac{\sinh(2u - \eta)}{\sinh(2u_j - \eta)} F_j. \quad (\text{A.4})$$

Then  $\lim_{\epsilon \rightarrow 0} \tilde{F}_1$  exists and is nonzero for  $u_1 = \pm \frac{\eta}{2} + \epsilon$ .

## References

- [1] R. Hagemans and J.-S. Caux, “Deformed strings in the Heisenberg model,” *J.Phys.A* **40** (2007) 14605, [arXiv:0707.2803](#) [cond-mat].
- [2] W. Hao, R. I. Nepomechie, and A. J. Sommes, “Completeness of solutions of Bethe’s equations,” *Phys. Rev.* **E88** no. 5, (2013) 052113, [arXiv:1308.4645](#) [math-ph].
- [3] C. Marboe and D. Volin, “Fast analytic solver of rational Bethe equations,” *J. Phys.* **A50** no. 20, (2017) 204002, [arXiv:1608.06504](#) [math-ph].
- [4] Y. Jiang and Y. Zhang, “Algebraic geometry and Bethe ansatz. Part I. The quotient ring for BAE,” *JHEP* **03** (2018) 087, [arXiv:1710.04693](#) [hep-th].
- [5] J. L. Jacobsen, Y. Jiang, and Y. Zhang, “Torus partition function of the six-vertex model from algebraic geometry,” [arXiv:1812.00447](#) [hep-th].
- [6] Z. Bajnok, J. L. Jacobsen, Y. Jiang, R. I. Nepomechie, and Y. Zhang, “in progress,”.
- [7] E. Granet and J. L. Jacobsen, “On zero-remainder conditions in the Bethe ansatz”.
- [8] V. Pasquier and H. Saleur, “Common Structures Between Finite Systems and Conformal Field Theories Through Quantum Groups,” *Nucl. Phys.* **B330** (1990) 523–556.
- [9] L. D. Faddeev, “How algebraic Bethe ansatz works for integrable models,” in *Symétries Quantiques (Les Houches Summer School Proceedings vol 64)*, A. Connes, K. Gawedzki, and J. Zinn-Justin, eds., pp. 149–219. North Holland, 1998. [arXiv:hep-th/9605187](#) [hep-th].

- [10] A. Izergin and V. Korepin, “Pauli principle for one-dimensional bosons and the algebraic Bethe ansatz,” *Lett.Math.Phys.* **6** (1982) 283–288.
- [11] L. Avdeev and A. Vladimirov, “Exceptional solutions of the Bethe ansatz equations,” *Theor. Math. Phys.* **69** (1987) 1071.
- [12] R. I. Nepomechie and C. Wang, “Algebraic Bethe ansatz for singular solutions,” *J. Phys.* **A46** (2013) 325002, [arXiv:1304.7978 \[hep-th\]](#).
- [13] G. P. Pronko and Yu. G. Stroganov, “Bethe equations ‘on the wrong side of equator’,” *J. Phys.* **A32** (1999) 2333–2340, [arXiv:hep-th/9808153 \[hep-th\]](#).
- [14] E. Mukhin, V. Tarasov, and A. Varchenko, “Bethe algebra of homogeneous XXX Heisenberg model has simple spectrum,” *Commun. Math. Phys.* **288** (2009) 1–42, [arXiv:0706.0688 \[math\]](#).
- [15] V. Tarasov, “Completeness of the Bethe ansatz for the periodic isotropic Heisenberg model,” *Rev. Math. Phys.* **30** (2018) 1840018.
- [16] R. I. Nepomechie and C. Wang, “Twisting singular solutions of Bethe’s equations,” *J. Phys.* **A47** no. 50, (2014) 505004, [arXiv:1409.7382 \[math-ph\]](#).
- [17] J. Thomae, “Beiträge zur Theorie der durch die Heinesche Reihe: . . . darstellbaren Functionen,” *J. für die reine und angewandte Mathematik* **70** (1869) 258–281.
- [18] F. H. Jackson, “The basic gamma-function and the elliptic functions,” *Proc. Roy. Soc. London* **76** (1905) 127–144.
- [19] R. J. Baxter, “Eight vertex model in lattice statistics and one-dimensional anisotropic Heisenberg chain. I. Some fundamental eigenvectors,” *Annals Phys.* **76** (1973) 1–24.
- [20] K. Fabricius and B. M. McCoy, “Bethe’s equation is incomplete for the XXZ model at roots of unity,” *J.Statist.Phys.* **103** (2001) 647–678, [arXiv:cond-mat/0009279 \[cond-mat.stat-mech\]](#).
- [21] R. J. Baxter, “Completeness of the Bethe ansatz for the six and eight vertex models,” *J.Statist.Phys.* **108** (2002) 1–48, [arXiv:cond-mat/0111188 \[cond-mat\]](#).
- [22] V. O. Tarasov, “On Bethe vectors for the XXZ model at roots of unity,” *J. Math. Sciences* **125** (2005) 242–248, [arXiv:math/0306032 \[math.QA\]](#).
- [23] A. M. Gainutdinov, W. Hao, R. I. Nepomechie, and A. J. Sommes, “Counting solutions of the Bethe equations of the quantum group invariant open XXZ chain at roots of unity,” *J. Phys.* **A48** no. 49, (2015) 494003, [arXiv:1505.02104 \[math-ph\]](#).
- [24] E. K. Sklyanin, “Boundary Conditions for Integrable Quantum Systems,” *J. Phys.* **A21** (1988) 2375.
- [25] P. Fendley and H. Saleur, “Deriving boundary S matrices,” *Nucl.Phys.* **B428** (1994) 681–693, [arXiv:hep-th/9402045 \[hep-th\]](#).