

Universal Bounds for Fluctuations in Quasistatic Small Heat Engines

Kosuke Ito,^{1,2} Chao Jiang,¹ and Gentaro Watanabe^{1,3,*}

¹Department of Physics and Zhejiang Institute of Modern Physics,
Zhejiang University, Hangzhou, Zhejiang 310027, China

²Graduate School of Engineering Science, Osaka University,
1-3 Machikaneyama, Toyonaka, Osaka 560-8531, Japan

³Zhejiang Province Key Laboratory of Quantum Technology and Device,
Zhejiang University, Hangzhou, Zhejiang 310027, China
(Dated: June 20, 2022)

We study higher order statistical properties of the Carnot cycle whose working substance consists of a small (classical) system. We show that the ratio between the fluctuations of work and heat is given by a universal form depending solely on the temperature ratio between the hot and cold heat baths. We also show that the Carnot cycle provides an upper bound on the former fluctuation ratio among cycles consisting of quasistatic strokes.

Introduction.—Advances in technology so far have enabled us to downsize the heat engines to the single colloidal particle or macromolecule level [1–7] or even smaller [8–14]. Such microscopic heat engines are expected to deepen our understanding of the thermodynamics for small systems [15–21] and have broad applications to nanomachines and biological machines [6, 18, 22, 23].

For conventional macroscopic heat engines, their performance is well characterized by the efficiency and power, which are based on the mean values of work and heat. However, for microscopic heat engines whose working substance consists of a system with a small number of degrees of freedom, fluctuations in thermodynamic quantities relative to their mean values are non-negligible [2, 4, 5, 24–31], and thus it is essentially important to characterize their performance not only by mean values, but also by higher order statistical quantities such as variance. One strategy to approach this issue is to identify the efficiency as a random variable [32–42]. In the present work, we take an alternative route by focusing on the quasistatic limit. The quasistatic case is practically important for the microscopic engines: Such engines can effectively reach this limit within a time scale much shorter than the macroscopic one [30] because the phase space structure of small systems are generally simple, and all the processes there are involved with only a small number of degrees of freedom, so that they are governed by microscopic time scales.

We study higher order statistical properties of the Carnot cycle whose working substance is a small (classical) system. We find that the higher order moment of the performance of the Carnot cycle shows a universal behavior similar to its efficiency depending solely on the temperature ratio T_c/T_h between the hot (T_h) and cold (T_c) heat baths. Furthermore, we show that the ratio between the fluctuations of work and heat for the Carnot cycle is maximum among cycles operating between the temperature T_c and T_h consisting of quasistatic strokes.

Working substance.—Let us first discuss the property of the working substance with which Carnot cycle can be realized. Since systems with a small number of degrees of freedom do not necessarily relax into an equilibrium state during adiabatic processes, the final state of a quasistatic adiabatic stroke does

not follow the canonical distribution in general. In order to make the cycle reversible, the final state of the quasistatic adiabatic strokes and the canonical state for the initial point of the subsequent quasistatic isothermal stroke should be statistically the same [16, 43]. Otherwise, irreversible heat exchange between the working substance and heat bath occurs when we make a thermal contact with a heat bath after the quasistatic adiabatic stroke.

Suppose the system is initially in thermal equilibrium at temperature T_1 when the external control parameter λ (such as the position of the piston in a setup of a gas in a cylinder with a piston) is at λ_1 , and a quasistatic adiabatic process is performed by slowly changing λ from λ_1 to λ_2 . After the quasistatic adiabatic process, we make a thermal contact between the system and a heat bath at some temperature T_2 . The internal energy of the initial and the final state of the adiabatic process is denoted by E_1 and E_2 , respectively. At the beginning of the quasistatic adiabatic process, the system is in a microstate Γ_1 in the phase space with energy $E_1 = H_{\lambda_1}(\Gamma_1)$ whose distribution is given by the canonical ensemble for H_{λ_1} at T_1 . During the quasistatic adiabatic process, the internal energy of the system changes in accordance with the adiabatic theorem such that the phase space volume $I_\lambda(E)$ enclosed by the iso-energy surface at E (so-called adiabatic invariant) is invariant under the change of λ . Here, $I_\lambda(E)$ is defined as $I_\lambda(E) \equiv \int \theta(E - H_\lambda(\Gamma)) d\Gamma$. A necessary and sufficient condition in order that the canonical ensemble for H_{λ_2} at some temperature T_2 is consistent with the final state of the quasistatic adiabatic process is [43]:

$$E_1/T_1 = E_2/T_2 \quad (1)$$

up to a constant.

In the present work, we consider working substance in which such T_2 can always be found for any values of λ_1 , λ_2 , and T_1 . Namely, we take working substance which allows us to form a Carnot cycle for an arbitrary choice of the bath temperatures T_h and T_c , and arbitrary values λ at the starting points of the quasistatic adiabatic strokes (points 1 and 3 in Fig. 1). As a consequence, throughout quasistatic adiabatic processes, the energy distribution function of the working substance is always given by the canonical one for some temper-




FIG. 1. The temperature versus external control parameter (T - λ) diagram of the Carnot cycle working with a hot heat bath at temperature T_h and a cold one at T_c .

ature. For example, working substance whose adiabatic invariant is in the form of $I_\lambda(E) = f(\lambda)E^\alpha$ with an arbitrary function f and a real constant α is a special case of the above class of working substance. Typical examples of small systems such as a single particle trapped in a harmonic oscillator potential and that in a box potential fall into this case.

Higher order statistics of the performance of the Carnot cycle.— Next, we discuss the central moment of work and heat for the Carnot cycle, and show the universality of the ratio between them. The n th order central moment $\langle(\Delta X)^n\rangle$ of a random variable X is defined as $\langle(\Delta X)^n\rangle \equiv \langle(X - \langle X \rangle)^n\rangle$, where $\langle A \rangle$ is the statistical average of “ A ”. Especially, the case of $n = 2$ is the variance $\langle\Delta X^2\rangle = \langle X^2 \rangle - \langle X \rangle^2$.

The four strokes of the Carnot cycle (see Fig. 1) are performed as follows. (0): *Initial state of the cycle.*— First, we set the external parameter at λ_0 and start with a randomly chosen microstate from the canonical ensemble for H_{λ_0} at temperature T_h . (0 \rightarrow 1): *Quasistatic isothermal expansion.*— We make a thermal contact between the engine and the hot heat bath with temperature T_h , then increase the parameter from λ_0 to λ_1 quasistatically. (1 \rightarrow 2[−]): *Quasistatic adiabatic expansion.*— We remove the thermal contact between the engine and the bath, and increase the parameter from λ_1 to λ_2 quasistatically. (2⁺ \rightarrow 3): *Quasistatic isothermal compression.*— At point 2⁺, we make a thermal contact between the engine and the cold heat bath with temperature T_c , then decrease the parameter from λ_2 to λ_3 . (3 \rightarrow 4[−]): *Quasistatic adiabatic compression.*— We remove the thermal contact between the engine and the bath, and decrease the parameter from λ_3 to λ_4 quasistatically. Here, point 4[−] is equivalent to point 0, and the parameter returns to the initial value, i.e. $\lambda_4 = \lambda_0$, to close the cycle.

In general, work and heat through each stroke, and the internal energy of the initial and the final states of each stroke are random variables. However, fluctuation ΔW_{isoth} of the work output W_{isoth} through the quasistatic isothermal process becomes negligible if the duration τ of the process is sufficiently long, and it vanishes no slower than $\sim \tau^{-1/2}$ [16, 44], i.e. $\Delta W_{\text{isoth}} \equiv W_{\text{isoth}} - \langle W_{\text{isoth}} \rangle = O(\tau^{-1/2})$. This is because there is no long-time correlation in the variation of the force exerted by the working substance. Let us now focus on the quasistatic isothermal expansion stroke 0 \rightarrow 1. From the first law of thermodynamics, work output $W_{0 \rightarrow 1}$ by the engine, heat input Q_h from the hot heat bath to the working substance, and the in-

ternal energy of the working substance E_0 and E_1 at the initial and the final state of the stroke should satisfy

$$Q_h = E_1 - E_0 + W_{0 \rightarrow 1}. \quad (2)$$

Since the fluctuation of $W_{0 \rightarrow 1}$ is negligible (i.e., $W_{0 \rightarrow 1} = \langle W_{0 \rightarrow 1} \rangle$) while Q_h , E_0 , and E_1 are not the case, we get

$$\Delta Q_h \equiv Q_h - \langle Q_h \rangle = (E_1 - \langle E_1 \rangle) - (E_0 - \langle E_0 \rangle). \quad (3)$$

Here, $\langle Q_h \rangle$ is the ensemble average of Q_h over possible sample paths [44], and $\langle E_i \rangle$ is the average of E_i over the canonical ensemble at point i .

Next, we consider the total work output W through the whole cycle:

$$W = W_{0 \rightarrow 1} + W_{1 \rightarrow 2} + W_{2 \rightarrow 3} + W_{3 \rightarrow 4}, \quad (4)$$

where $W_{i \rightarrow i+1}$ is work output through the stroke from point i to $i+1$ as $W_{0 \rightarrow 1}$ defined previously. Since the fluctuations of $W_{0 \rightarrow 1}$ and $W_{2 \rightarrow 3}$ by the quasistatic isothermal strokes are negligible, we obtain

$$\Delta W \equiv W - \langle W \rangle = (W_{1 \rightarrow 2} - \langle W_{1 \rightarrow 2} \rangle) + (W_{3 \rightarrow 4} - \langle W_{3 \rightarrow 4} \rangle). \quad (5)$$

Here, the strokes 1 \rightarrow 2[−] and 3 \rightarrow 4[−] are quasistatic adiabatic processes. Since there is no heat exchange between the working substance and the heat bath during these strokes, $W_{1 \rightarrow 2}$, for example, reads

$$W_{1 \rightarrow 2} = E_1 - E_2^- \quad (6)$$

with E_1 and E_2^- being the internal energy of the working substance at points 1 and 2[−], respectively. In addition, for our working substance, since the initial and the final state of the quasistatic adiabatic stroke should satisfy the relation (1), we have $E_1/T_h = E_2^-/T_c$. From this relation and Eq. (6), we get

$$W_{1 \rightarrow 2} = [1 - (T_c/T_h)] E_1. \quad (7)$$

Similarly, for the quasistatic adiabatic compression stroke 3 \rightarrow 4[−], we get

$$W_{3 \rightarrow 4} = E_3 - E_4^- = [(T_c/T_h) - 1] E_4^-. \quad (8)$$

Substituting Eqs. (7) and (8) into Eq. (5), we obtain

$$\Delta W = [1 - (T_c/T_h)] [(E_1 - \langle E_1 \rangle) - (E_4^- - \langle E_4^- \rangle)]. \quad (9)$$

Since E_0 and E_4^- are equivalent random variables, from Eqs. (3) and (9) we finally get

$$\eta^{(n)} \equiv \frac{\langle(\Delta W)^n\rangle}{\langle(\Delta Q_h)^n\rangle} = \left(1 - \frac{T_c}{T_h}\right)^n. \quad (10)$$

This is one of the main results of this work. It is noted that the ratio $\eta^{(n)}$ for the Carnot cycle given by Eq. (10) shows the universal form which depends only on the ratio T_c/T_h .




FIG. 2. Four kinds of processes considered [(a) quasistatic isothermal, (b) quasistatic adiabatic, (c) isochoric, and (d) quasistatic isobaric strokes] on the T - λ plane.

For the quasistatic isothermal compression stroke $2^+ \rightarrow 3$, heat output Q_c from the working substance to the cold heat bath through this stroke is given by

$$Q_c = E_2^+ - E_3 - W_{2 \rightarrow 3}. \quad (11)$$

Since the fluctuation of $W_{2 \rightarrow 3}$ is negligible, we get

$$\Delta Q_c \equiv Q_c - \langle Q_c \rangle = (E_2^+ - \langle E_2^+ \rangle) - (E_3 - \langle E_3 \rangle). \quad (12)$$

The condition (1) for the quasistatic adiabatic strokes $1 \rightarrow 2^-$ and $3 \rightarrow 4^-$ reads $E_1/T_h = E_2^-/T_c$ and $E_3/T_c = E_4^-/T_h$, respectively. Since E_2^+ and E_3 are independent and the distributions of E_2^- and E_2^+ are the same, Eq. (12) with the above conditions leads to

$$\begin{aligned} \langle (\Delta Q_c)^n \rangle &= (T_c/T_h)^n \langle [(E_1 - \langle E_1 \rangle) - (E_0 - \langle E_0 \rangle)]^n \rangle \\ &= (T_c/T_h)^n \langle (\Delta Q_h)^n \rangle. \end{aligned} \quad (13)$$

Because Q_h and Q_c are independent, the variance $\langle \Delta Q^2 \rangle$ of the total heat input $Q \equiv Q_h - Q_c$ can be written as

$$\langle \Delta Q^2 \rangle = \langle \Delta Q_h^2 \rangle + \langle \Delta Q_c^2 \rangle = [1 + (T_c/T_h)^2] \langle \Delta Q_h^2 \rangle. \quad (14)$$

From Eqs. (10) and (14), we finally obtain

$$\xi^{(2)} \equiv \frac{\langle \Delta W^2 \rangle}{\langle \Delta Q^2 \rangle} = \frac{[1 - (T_c/T_h)]^2}{1 + (T_c/T_h)^2}. \quad (15)$$

This is another main result showing that the variance of work W relative to that of heat Q of the Carnot cycle also has a universal form depending only on the temperature ratio T_c/T_h .

Bounds on $\eta^{(2)}$ and $\xi^{(2)}$.—Finally, we show that work fluctuations $\eta^{(2)}$ and $\xi^{(2)}$ relative to heat fluctuation for the Carnot cycle given by Eqs. (10) and (15) provide the upper bound for cycles using the working substance considered which allows us to form the Carnot cycle [45]. We consider an arbitrary cycle consisting of any of the quasistatic isothermal, quasistatic adiabatic, isochoric (equal “volume”), or quasistatic isobaric (equal “pressure”) strokes operating between the temperature

TABLE I. Work fluctuation $\langle \Delta W_{i \rightarrow i+1}^2 \rangle$ and heat fluctuation $\langle \Delta Q_{i \rightarrow i+1}^2 \rangle$ for various thermodynamic processes from point i to $i+1$. $\langle \Delta E_i^2 \rangle$ is the variance of the internal energy E_i of the working substance for the canonical distribution at point i .

Process	$\langle \Delta W_{i \rightarrow i+1}^2 \rangle$	$\langle \Delta Q_{i \rightarrow i+1}^2 \rangle$
(1) Isothermal	0	$\langle \Delta E_i^2 \rangle + \langle \Delta E_{i+1}^2 \rangle$
(2) Adiabatic	$[1 - (T_{i+1}/T_i)]^2 \langle \Delta E_i^2 \rangle$ $= [(T_i/T_{i+1}) - 1]^2 \langle \Delta E_{i+1}^2 \rangle$	0
(3) Isochoric	0	$\langle \Delta E_i^2 \rangle + \langle \Delta E_{i+1}^2 \rangle$
(4) Isobaric	0	$\langle \Delta E_i^2 \rangle + \langle \Delta E_{i+1}^2 \rangle$

T_c and T_h [46]. On the temperature versus external parameter (T - λ) plane, a quasistatic isothermal stroke is shown by a horizontal line at T_h or T_c [Fig. 2(a)], and a quasistatic adiabatic stroke is by a downward-sloping curve [Fig. 2(b)].

Isobaric processes are ones during which the generalized pressure $P \equiv -\langle \partial H_\lambda / \partial \lambda \rangle$ is constant. Here, change of P due to the variation of λ should be compensated by that due to the heat exchange between the working substance and the heat bath. Thus a quasistatic adiabatic stroke is shown by an upward-sloping curve on the T - λ plane [Fig. 2(d)]. To perform quasistatic isobaric processes, we introduce “quasi-adiabatic wall”, which is made of an imperfect heat insulator. When we make a thermal contact between the working substance and the heat bath, heat conduction between them and thermalization in the working substance occur simultaneously. By surrounding the working substance by the quasi-adiabatic wall and making a thermal contact with a heat bath through this wall, we can make the time scale of heat conduction much larger than that of thermalization. In this situation, the working substance is always in thermal equilibrium while there is continuous heat exchange between the working substance and the heat bath until the temperature of the former reaches that of the latter. Then, quasistatic isobaric processes can be performed by changing λ with keeping P constant, which is given by the canonical average of $-\partial H_\lambda / \partial \lambda$ for instantaneous values of the parameter λ and the temperature T of the working substance. For example, for working substance with $I_\lambda(E) = f(\lambda)E^\alpha$, the generalized pressure is given by $P = k_B T f(\lambda)^{-1} \partial_\lambda f(\lambda)$, where k_B is the Boltzmann constant, and the instantaneous values of λ and T are related to keep P fixed at the initial value.

Isochoric processes are ones during which λ is fixed and heat exchange between the working substance and the heat bath is performed. On the T - λ plane, an isochoric stroke with the hot (cold) heat bath is shown by an upward (downward) vertical line [Fig. 2(c)]. If we make a direct thermal contact between the working substance and the heat bath, the final state of the isochoric process is a canonical state at the temperature of the bath; however, if we make a thermal contact through the quasi-adiabatic wall, the temperature of the final canonical state can be any value between the initial state and the temperature of the bath.

Now we consider fluctuations of work output and heat input through the above four kinds of processes. Suppose the initial and the final points of the stroke are points i and $i+1$, and the variances of work output and heat input through the stroke $i \rightarrow i+1$ are denoted by $\langle \Delta W_{i \rightarrow i+1}^2 \rangle$ and $\langle \Delta Q_{i \rightarrow i+1}^2 \rangle$, respectively. (1) For quasistatic isothermal strokes, $\langle \Delta W_{i \rightarrow i+1}^2 \rangle = 0$ as mentioned before. Since $\Delta Q_{i \rightarrow i+1} = (E_{i+1} - \langle E_{i+1} \rangle) - (E_i - \langle E_i \rangle)$ and the internal energies E_i and E_{i+1} at points i and $i+1$ are independent, heat fluctuation is given by $\langle \Delta Q_{i \rightarrow i+1}^2 \rangle = \langle \Delta E_{i+1}^2 \rangle + \langle \Delta E_i^2 \rangle$. (2) For quasistatic adiabatic strokes, $\langle \Delta Q_{i \rightarrow i+1}^2 \rangle = 0$ by definition. Since $W_{i \rightarrow i+1} = [1 - (T_{i+1}/T_i)]E_i = [(T_i/T_{i+1}) - 1]E_{i+1}$ obtained in the same way as Eqs. (7) and (8), work fluctuation is given by $\langle \Delta W_{i \rightarrow i+1}^2 \rangle = [1 - (T_{i+1}/T_i)]^2 \langle \Delta E_i^2 \rangle = [(T_i/T_{i+1}) - 1]^2 \langle \Delta E_{i+1}^2 \rangle$. (3) For isochoric strokes, $\langle \Delta W_{i \rightarrow i+1}^2 \rangle = 0$ by definition, and $\langle \Delta Q_{i \rightarrow i+1}^2 \rangle = \langle \Delta E_{i+1}^2 \rangle + \langle \Delta E_i^2 \rangle$ according to the similar reason as in the case of quasistatic isothermal strokes. (4) For quasistatic isobaric strokes, $\langle \Delta W_{i \rightarrow i+1}^2 \rangle = 0$ and $\langle \Delta Q_{i \rightarrow i+1}^2 \rangle = \langle \Delta E_{i+1}^2 \rangle + \langle \Delta E_i^2 \rangle$ due to the similar reason for quasistatic isothermal strokes. The results of $\langle \Delta W_{i \rightarrow i+1}^2 \rangle$ and $\langle \Delta Q_{i \rightarrow i+1}^2 \rangle$ for each kind of processes are summarized in Table I.

Suppose we have an arbitrary cycle consisting of the above four kinds of processes operating between the bath temperatures T_c and T_h . We choose the starting point such that the final stroke is an adiabatic one [48]; for cycles with no adiabatic strokes, trivially $\langle \Delta W^2 \rangle = 0$ and thus $\xi^{(2)} = \eta^{(2)} = 0$. Among all the N nodes of the cycle $k = 0, 1, 2, \dots, N$ with points 0 and N being identical, we consider their subsets of the both ends of quasistatic adiabatic expansion and compression strokes. For the i th adiabatic expansion (compression) stroke, the initial and the final points are denoted by J_i and $J_i + 1$ (K_i and $K_i + 1$), and the energy and the temperature of these points satisfy the condition (1) reading $E_{J_i+1} = (T_{J_i+1}/T_{J_i})E_{J_i}$ [$E_{K_i} = (T_{K_i}/T_{K_i+1})E_{K_i+1}$]. Since only the adiabatic strokes yield work fluctuation, fluctuation of total work output through the cycle is given by

$$\langle \Delta W^2 \rangle = \sum_i [1 - (T_{J_i+1}/T_{J_i})]^2 \langle \Delta E_{J_i}^2 \rangle + \sum_j [1 - (T_{K_j}/T_{K_j+1})]^2 \langle \Delta E_{K_j+1}^2 \rangle. \quad (16)$$

Regarding the fluctuation $\langle \Delta Q^2 \rangle$ of total heat input, the end points of adiabatic strokes must be connected to the other kinds of strokes, which have heat fluctuation. On the other hand, if there are heat exchanging strokes (those other than quasistatic adiabatic stroke) consecutively, the node connecting them does not contribute to $\langle \Delta Q^2 \rangle$ because the internal energies at the final point of the preceding stroke and at the initial point of the following one cancel with each other. Thus,

$$\begin{aligned} \langle \Delta Q^2 \rangle &= \sum_i (\langle \Delta E_{J_i}^2 \rangle + \langle \Delta E_{J_i+1}^2 \rangle) + \sum_j (\langle \Delta E_{K_j}^2 \rangle + \langle \Delta E_{K_j+1}^2 \rangle) \\ &= \sum_i [1 + (T_{J_i+1}/T_{J_i})^2] \langle \Delta E_{J_i}^2 \rangle \\ &\quad + \sum_j [1 + (T_{K_j}/T_{K_j+1})^2] \langle \Delta E_{K_j+1}^2 \rangle. \end{aligned} \quad (17)$$

Since the temperatures T_{J_i} and T_{J_i+1} (T_{K_j} and T_{K_j+1}) are in the region of $[T_c, T_h]$ and $T_{J_i} \geq T_{J_i+1}$ ($T_{K_j+1} \geq T_{K_j}$), we get $1 \geq T_{J_i+1}/T_{J_i} \geq T_c/T_h \geq T_c/T_h$ ($1 \geq T_{K_j}/T_{K_j+1} \geq T_c/T_h$). Therefore, $[1 - (T_{J_i+1}/T_{J_i})]^2, [1 - (T_{K_j}/T_{K_j+1})]^2 \leq [1 - (T_c/T_h)]^2$ and $1 + (T_{J_i+1}/T_{J_i})^2, 1 + (T_{K_j}/T_{K_j+1})^2 \geq 1 + (T_c/T_h)^2$. Applying these relations to Eqs. (16) and (17), we obtain

$$\langle \Delta W^2 \rangle \leq [1 - (T_c/T_h)]^2 \left(\sum_i \langle \Delta E_{J_i}^2 \rangle + \sum_j \langle \Delta E_{K_j+1}^2 \rangle \right), \quad (18)$$

$$\langle \Delta Q^2 \rangle \geq [1 + (T_c/T_h)^2] \left(\sum_i \langle \Delta E_{J_i}^2 \rangle + \sum_j \langle \Delta E_{K_j+1}^2 \rangle \right). \quad (19)$$

Thus we finally get

$$\xi^{(2)} \leq \xi_C^{(2)}, \quad (20)$$

where $\xi_C^{(2)} \equiv [1 - (T_c/T_h)]^2 / [1 + (T_c/T_h)^2]$ is for the Carnot cycle given by Eq. (15). Since a sum of the fluctuations of heat inputs for each stroke trivially satisfies $\sum_{i=0}^{N-1} \langle \Delta Q_{i \rightarrow i+1}^2 \rangle \geq \langle \Delta Q^2 \rangle$, another ratio $\tilde{\xi}^{(2)} \equiv \langle \Delta W^2 \rangle / \sum_{i=0}^{N-1} \langle \Delta Q_{i \rightarrow i+1}^2 \rangle$ defined with this quantity is also bounded by ξ_C : $\tilde{\xi}^{(2)} \leq \xi^{(2)} \leq \xi_C^{(2)}$. Note that $\tilde{\xi}^{(2)} = \xi^{(2)}$ for the Carnot cycle.

Regarding $\eta^{(2)}$, the one $\eta_C^{(2)}$ for the Carnot cycle gives the maximum value among any cycle all of whose quasistatic adiabatic expansion (compression) strokes are preceded (followed) by a stroke with the hot bath [47]. Typical cycles such as the Otto, Brayton, Stirling, Ericsson cycles, etc. satisfy this condition. In such cycles, the energy fluctuation at the initial point J_i (final point $K_j + 1$) of all the quasistatic adiabatic expansion (compression) strokes contribute to $\langle \Delta Q_h^2 \rangle$. Thus,

$$\langle \Delta Q_h^2 \rangle \geq \sum_i \langle \Delta E_{J_i}^2 \rangle + \sum_j \langle \Delta E_{K_j+1}^2 \rangle. \quad (21)$$

From Eqs. (18) and (21), we get

$$\eta^{(2)} \leq \eta_C^{(2)}, \quad (22)$$

with $\eta_C^{(2)} \equiv [1 - (T_c/T_h)]^2$ given by Eq. (10) for $n = 2$. For the Stirling and Ericsson cycles, $\langle \Delta W^2 \rangle = 0$ and thus $\eta^{(2)} = 0$. For the Otto and the Brayton cycles, taking the working substance with $I_\lambda(E) = f(\lambda)E^\alpha$ as an example, we get $\eta^{(2)} = [1 - (T_2/T_1)]^2$, where T_1 and T_2 are the initial and the final temperatures of the quasistatic adiabatic expansion stroke. Since $1 \geq T_2/T_1 \geq T_c/T_h$, this $\eta^{(2)}$ indeed satisfies $\eta^{(2)} \leq \eta_C^{(2)}$.

Concluding remarks.— We have derived the arbitrary order of the central moment of work and heat of the Carnot cycle, and have shown that the ratio between them has universal relations (10) and (15). We have also shown that the Carnot cycle provides the upper bound for the ratio between the fluctuations of work and heat [Eqs. (22) and (20)]. Our results would provide a new guiding principle to design microscopic heat engines.

We thank Yuki Izumida and Peter Talkner for helpful discussions and comments. This work is supported by the

Zhejiang Provincial Natural Science Foundation Key Project (Grant No. LZ19A050001), by NSF of China (Grant No. 11674283), by the Fundamental Research Funds for the Central Universities (2017QNA3005, 2018QNA3004), by the Zhejiang University 100 Plan, and by the Junior 1000 Talents Plan of China. K. I. is also supported by MEXT Quantum Leap Flagship Program (MEXT Q-LEAP) (Grant No. JPMXS0118067394).

* gentaro@zju.edu.cn

- [1] T. Hugel, N. B. Holland, A. Cattani, L. Moroder, M. Seitz, H. E. Gaub, *Science* **296**, 1103 (2002).
- [2] V. Blickle and C. Bechinger, *Nat. Phys.* **8**, 143 (2012).
- [3] P. A. Quinto-Su, *Nat. Commun.* **5**, 5889 (2014).
- [4] I. A. Martínez, É. Roldán, L. Dinis, D. Petrov, J. M. R. Parrondo, and R. A. Rica, *Nat. Phys.* **12**, 67 (2016).
- [5] S. Krishnamurthy, S. Ghosh, D. Chatterji, R. Ganapathy, and A. K. Sood, *Nat. Phys.* **12**, 1134 (2016).
- [6] I. A. Martínez, É. Roldán, L. Dinis, and R. A. Rica, *Soft Matter*, **13**, 22 (2017).
- [7] A. Argun, J. Soni, L. Dabelow, S. Bo, G. Pesce, R. Eichhorn, and G. Volpe, *Phys. Rev. E* **96**, 052106 (2017).
- [8] P. G. Steeneken, K. Le Phan, M. J. Goossens, G. E. J. Koops, G. J. A. M. Brom, C. van der Avoort, and J. T. M. van Beek, *Nat. Phys.* **7**, 354 (2011).
- [9] J. V. Koski, V. F. Maisi, J. P. Pekola, and D. V. Averin, *Proc. Natl. Acad. Sci. U.S.A.* **111**, 13786 (2014).
- [10] J. Roßnagel, S. T. Dawkins, K. N. Tolazzi, O. Abah, E. Lutz, F. Schmidt-Kaler, and K. Singer, *Science* **352**, 325 (2016).
- [11] G. Maslennikov, S. Ding, R. Hablützel, J. Gan, A. Roulet, S. Nimmrichter, J. Dai, V. Scarani, and D. Matsukevich, *Nat. Commun.* **10**, 202 (2019).
- [12] J. Klatzow, J. N. Becker, P. M. Ledingham, C. Weinzetl, K. T. Kaczmarek, D. J. Saunders, J. Nunn, I. A. Walmsley, R. Uzdin, and E. Poem, *Phys. Rev. Lett.* **122**, 110601 (2019).
- [13] D. von Lindenfels, O. Gräß, C. T. Schmiegelow, V. Kaushal, J. Schulz, M. T. Mitchison, J. Goold, F. Schmidt-Kaler, and U. G. Poschinger, *Phys. Rev. Lett.* **123**, 080602 (2019).
- [14] J. P. S. Peterson, T. B. Batalhão, M. Herrera, A. M. Souza, R. S. Sarthour, I. S. Oliveira, and R. M. Serra, *arXiv:1803.06021*.
- [15] K. Sekimoto, *Prog. Theor. Phys. Suppl.* **130**, 17 (1998).
- [16] K. Sekimoto, *Stochastic energetics*, Lecture Notes in Physics Vol. 799 (Springer, Berlin, 2010).
- [17] C. Jarzynski, *Ann. Rev. Cond. Mat. Phys.* **2**, 329 (2011).
- [18] U. Seifert, *Rep. Prog. Phys.* **75**, 126001 (2012).
- [19] C. Van den Broeck and M. Esposito, *Physica A* **418**, 6 (2015).
- [20] S. Ciliberto, *Phys. Rev. X* **7**, 021051 (2017).
- [21] G. Nicolis and Y. De Decker, *Entropy* **19**, 434 (2017).
- [22] C. Bustamante, J. Liphardt, and F. Ritort, *Phys. Today* **58**, 43 (2005).
- [23] S. Toyabe and M. Sato, *J. Phys. Soc. Jpn.* **84**, 102001 (2015).
- [24] K. Sekimoto, F. Takagi, and T. Hondou, *Phys. Rev. E* **62**, 7759 (2000).
- [25] P. Jop, A. Petrosyan, and S. Ciliberto, *Europhys. Lett.* **81**, 50005 (2008).
- [26] J. Hoppenau, M. Niemann, and A. Engel, *Phys. Rev. E* **87**, 062127 (2013).
- [27] V. Holubec, *J. Stat. Mech.* **2014**, P05022 (2014).
- [28] S. Rana, P. S. Pal, A. Saha, and A. M. Jayannavar, *Phys. Rev. E* **90**, 042146 (2014).
- [29] L. Cerino, A. Puglisi, and A. Vulpiani, *Phys. Rev. E* **91**, 032128 (2015).
- [30] I. A. Martínez, É. Roldán, L. Dinis, D. Petrov, and R. A. Rica, *Phys. Rev. Lett.* **114**, 120601 (2015).
- [31] V. Holubec and A. Ryabov, *Phys. Rev. E* **96**, 030102(R) (2017).
- [32] G. Verley, M. Esposito, T. Willaert, and C. Van den Broeck, *Nat. Commun.* **5**, 4721 (2014).
- [33] G. Verley, T. Willaert, C. Van den Broeck, and M. Esposito, *Phys. Rev. E* **90**, 052145 (2014).
- [34] M. Polettini, G. Verley, and M. Esposito, *Phys. Rev. Lett.* **114**, 050601 (2015).
- [35] K. Proesmans and C. Van den Broeck, *New. J. Phys.* **17**, 065004 (2015).
- [36] J.-H. Jiang, B. K. Agarwalla, and D. Segal, *Phys. Rev. Lett.* **115**, 040601 (2015).
- [37] K. Proesmans, C. Driesen, B. Cleuren, and C. Van den Broeck, *Phys. Rev. E* **92**, 032105 (2015).
- [38] H. Vroylandt, A. Bonfils, and G. Verley, *Phys. Rev. E* **93**, 052123 (2016).
- [39] J.-M. Park, H.-M. Chun, and J. D. Noh, *Phys. Rev. E* **94**, 012127 (2016).
- [40] D. Gupta and S. Sabhapandit, *Phys. Rev. E* **96**, 042130 (2017).
- [41] M. Suñé and A. Imparato, *J. Phys. A: Math. Theor.* **52**, 045003 (2019).
- [42] S. K. Manikandan, L. Dabelow, R. Eichhorn, and S. Krishnamurthy, *Phys. Rev. Lett.* **122**, 140601 (2019).
- [43] K. Sato, K. Sekimoto, T. Hondou, and F. Takagi, *Phys. Rev. E* **66**, 016119 (2002).
- [44] See Supplemental Material.
- [45] However, there exist other cycles which give the same value as $\eta_C^{(2)}$ and $\xi_C^{(2)}$ [44].
- [46] Although it is possible to go outside the region of $T_c \leq T \leq T_h$ by a quasistatic adiabatic stroke, getting out of this region is necessarily accompanied by a subcycle of opposite direction giving negative work output to close the whole cycle. Therefore, such cases are less important and we shall constrain ourselves within the region of $T_c \leq T \leq T_h$.
- [47] If it is not the case, it is possible to have $\eta^{(2)} > \eta_C^{(2)}$. See [44] for such an example.
- [48] This choice is based on our definition of heat input through one cycle, where we avoid the situation in which both of the equivalent points 0 and N contribute to the heat fluctuation.

Supplemental Material

Fluctuations of work and heat in the quasistatic isothermal process

It has been discussed that the fluctuation of work output throughout the quasistatic isothermal process vanishes as $\sim o(\tau^{-1/2})$ for long duration τ of the process [S1, S2]. Namely, in the limit of large τ , each sample path gives the same value of the work for a given protocol in the quasistatic isothermal process. This is because there is no long-time correlation in the variation of the force exerted by the working substance: this force varies due to thermal fluctuation caused by the contact with a heat bath, and thus the variation does not have a long-time correlation. In the following, we shall show this vanishing the fluctuation using the path integral representation.

Let us consider a trajectory γ in the phase space between the initial time t_{init} and the final time t_{fin} under the driving of the external control parameter $\lambda(t)$ (whose initial and the final values are λ_{init} and λ_{fin} , respectively). The work output W from the system along this trajectory is given by

$$W[\gamma] \equiv - \int_{t_{\text{init}}}^{t_{\text{fin}}} dt \frac{d\lambda}{dt} \frac{\partial H_\lambda[\gamma]}{\partial \lambda}. \quad (\text{S1})$$

The first and the second moments of the work averaged over sample paths are

$$\langle W \rangle = \int \mathcal{D}\Gamma \mathcal{P}[\gamma] W[\gamma], \quad (\text{S2})$$

$$\langle W^2 \rangle = \int \mathcal{D}\Gamma \mathcal{P}[\gamma] (W[\gamma])^2, \quad (\text{S3})$$

where $\mathcal{P}[\gamma]$ is the probability (density) of the trajectory γ , and $\int \mathcal{D}\Gamma$ denotes the functional integral with respect to the trajectories.

We set the duration of the process as $\tau \equiv t_{\text{fin}} - t_{\text{init}} = M \tau_{\text{unit}}$ with M being an integer and τ_{unit} being a sufficiently long time so that the variation of the parameter λ is slow enough. (As a prerequisite, τ_{unit} is taken to be much larger than the correlation time τ_{corr} of the thermal fluctuation of the force.) Then we shall see the variance $\langle \Delta W^2 \rangle$ of work W vanishes for large M . To evaluate the right-hand side of Eqs. (S2) and (S3), we discretize the time τ_{unit} into a sufficiently large number of $N + 1$ points, $\{t_0 \equiv t_{\text{init}}, t_1, t_2, \dots, t_N \equiv t_{\text{init}} + \tau_{\text{unit}}\}$, by the time step $\Delta t \equiv t_{n+1} - t_n = \tau_{\text{unit}}/N$, which is taken to be much larger than the correlation time τ_{corr} so that the force exerted by the working substance at different time slices t_n is uncorrelated. Therefore, the probability density $\mathcal{P}[\gamma]$ of the trajectory can be written as a product of the phase space distribution function $P_{\lambda_n}(\Gamma_n; t_n)$ at each time slice t_n . The phase space point and the external parameter at time t_n ($0 \leq n \leq MN$) are denoted by $\Gamma_n \equiv (\mathbf{q}_n, \mathbf{p}_n)$ and λ_n , respectively. Here, \mathbf{q} and \mathbf{p} represent the D generalized coordinates and momenta, respectively, in the $2D$ -dimensional phase space for the system with D degrees of freedom. Then $\mathcal{D}\Gamma$ reduces to $\mathcal{D}\Gamma \rightarrow \prod_{n=0}^{MN} d\Gamma_n$, where $d\Gamma_n \equiv C d\mathbf{q}_n d\mathbf{p}_n$ is the phase space volume element at time t_n including the numerical factor C coming from the phase space volume of a microstate.

Thus, the average of work $\langle W \rangle$ given by Eq. (S2) reads

$$\begin{aligned} \langle W \rangle &\simeq \int \prod_{n=0}^{MN} d\Gamma_n P_{\lambda_n}(\Gamma_n; t_n) \left[- \sum_{m=0}^{MN-1} \frac{\Delta \lambda_{m/M}}{M} \frac{\partial H_{\lambda_m}(\Gamma_m)}{\partial \lambda} \right] \\ &= - \sum_{m=0}^{MN-1} \frac{\Delta \lambda_{m/M}}{M} \int d\Gamma_m P_{\lambda_m}(\Gamma_m; t_m) \frac{\partial H_{\lambda_m}(\Gamma_m)}{\partial \lambda}, \end{aligned} \quad (\text{S4})$$

where $\Delta \lambda_i$ with $0 \leq i \leq N$ sets the protocol of the parameter change in the case of $\tau = \tau_{\text{unit}}$ with $M = 1$: $\Delta \lambda_i \equiv \lambda_{i+1} - \lambda_i$ if $i = m/M$ is an integer; otherwise, $\Delta \lambda_{m/M}$ is between $\Delta \lambda_{[m/M]}$ and $\Delta \lambda_{[m/M]+1}$, where $[m/M]$ is the integer part of m/M . Note that the factor of $1/M$ appears because the speed of the parameter change is slowed down by increasing the total duration τ of the process by a factor of M . Similarly, the second moment (S3) becomes

$$\begin{aligned} \langle W^2 \rangle &\simeq \int \prod_{n=0}^{MN} d\Gamma_n P_{\lambda_n}(\Gamma_n; t_n) \left[- \sum_{m=0}^{MN-1} \frac{\Delta \lambda_{m/M}}{M} \frac{\partial H_{\lambda_m}(\Gamma_m)}{\partial \lambda} \right]^2 \\ &= \left[\sum_{m=0}^{MN-1} \frac{\Delta \lambda_{m/M}}{M} \int d\Gamma_m P_{\lambda_m}(\Gamma_m; t_m) \frac{\partial H_{\lambda_m}(\Gamma_m)}{\partial \lambda} \right]^2 \\ &\quad - \sum_{m=0}^{MN-1} \left(\frac{\Delta \lambda_{m/M}}{M} \right)^2 \left[\int d\Gamma_m P_{\lambda_m}(\Gamma_m; t_m) \frac{\partial H_{\lambda_m}(\Gamma_m)}{\partial \lambda} \right]^2 + \sum_{m=0}^{MN-1} \left(\frac{\Delta \lambda_{m/M}}{M} \right)^2 \int d\Gamma_m P_{\lambda_m}(\Gamma_m; t_m) \left[\frac{\partial H_{\lambda_m}(\Gamma_m)}{\partial \lambda} \right]^2. \end{aligned} \quad (\text{S5})$$

Here, the second and the third terms of the right-hand side scales as $\sim 1/M$ since the summation $\sum_{m=0}^{MN-1}$ yields a contribution of a factor of M , which is multiplied by the factor of $1/M^2$.

From Eqs. (S4) and (S5), we readily see that the variance of work $\langle \Delta W^2 \rangle$ in the quasistatic isothermal process is given by the second and the third terms of Eq. (S5) both of which scale as $\sim 1/M$ and have an opposite sign with each other. Therefore, the variance $\langle \Delta W^2 \rangle$ vanishes as $\sim \tau^{-1}$ or faster:

$$\langle \Delta W^2 \rangle \equiv \langle W^2 \rangle - \langle W \rangle^2 = O(\tau^{-1}), \quad (\text{S6})$$

or the fluctuation ΔW vanishes as $\sqrt{\langle \Delta W^2 \rangle} = O(\tau^{-1/2})$. As a consequence, fluctuation of W through the quasistatic isothermal process becomes negligible provided the duration τ of the process is sufficiently long, and $\langle \Delta W^2 \rangle \rightarrow 0$ in the limit of $\tau \rightarrow \infty$.

Next, we shall also discuss the fluctuation of heat during the quasistatic isothermal process. From the first law of thermodynamics for an individual trajectory, the heat absorbed by the system along the trajectory γ is given by

$$Q[\gamma] \equiv H_{\lambda_{\text{fin}}}(\Gamma_{\text{fin}}) - H_{\lambda_{\text{init}}}(\Gamma_{\text{init}}) + W[\gamma], \quad (\text{S7})$$

where Γ_{init} and Γ_{fin} are the initial and the final phase space points of the trajectory γ . The first and the second moments of the heat averaged over sample paths are

$$\langle Q \rangle = \int \mathcal{D}\Gamma \mathcal{P}[\gamma] Q[\gamma], \quad (\text{S8})$$

$$\langle Q^2 \rangle = \int \mathcal{D}\Gamma \mathcal{P}[\gamma] (Q[\gamma])^2. \quad (\text{S9})$$

The average of heat (S8) can be written as

$$\langle Q \rangle = \langle H_{\lambda_{\text{fin}}} \rangle_{\lambda_{\text{fin}}, t_{\text{fin}}} - \langle H_{\lambda_{\text{init}}} \rangle_{\lambda_{\text{init}}, t_{\text{init}}} + \langle W \rangle, \quad (\text{S10})$$

where $\langle W \rangle$ is given by Eq. (S4), and $\langle A \rangle_{\lambda, t} \equiv \int d\Gamma P_{\lambda}(\Gamma; t) A(\Gamma)$ is the average of “ A ” at time t .

Similarly, the second moment (S9) reads

$$\begin{aligned} \langle Q^2 \rangle &= \langle H_{\lambda_{\text{fin}}}^2 \rangle_{\lambda_{\text{fin}}, t_{\text{fin}}} - 2 \langle H_{\lambda_{\text{fin}}} \rangle_{\lambda_{\text{fin}}, t_{\text{fin}}} \langle H_{\lambda_{\text{init}}} \rangle_{\lambda_{\text{init}}, t_{\text{init}}} + \langle H_{\lambda_{\text{init}}}^2 \rangle_{\lambda_{\text{init}}, t_{\text{init}}} + \langle W^2 \rangle \\ &\quad - 2 \int \mathcal{D}\Gamma \mathcal{P}[\gamma] [H_{\lambda_{\text{fin}}}(\Gamma_{\text{fin}}) - H_{\lambda_{\text{init}}}(\Gamma_{\text{init}})] \int_{\lambda_{\text{init}}}^{\lambda_{\text{fin}}} dt \frac{d\lambda}{dt} \frac{\partial H_{\lambda}[\gamma]}{\partial \lambda}. \end{aligned} \quad (\text{S11})$$

Here, the fifth term in the right-hand side can be written as

$$\begin{aligned} &\int \mathcal{D}\Gamma \mathcal{P}[\gamma] [H_{\lambda_{\text{fin}}}(\Gamma_{\text{fin}}) - H_{\lambda_{\text{init}}}(\Gamma_{\text{init}})] \int_{\lambda_{\text{init}}}^{\lambda_{\text{fin}}} dt \frac{d\lambda}{dt} \frac{\partial H_{\lambda}[\gamma]}{\partial \lambda} \\ &\simeq \int \prod_{n=0}^{MN} d\Gamma_n P_{\lambda_n}(\Gamma_n; t_n) [H_{\lambda_{\text{fin}}}(\Gamma_{\text{fin}}) - H_{\lambda_{\text{init}}}(\Gamma_{\text{init}})] \sum_{m=0}^{MN-1} \frac{\Delta \lambda_{m/M}}{M} \frac{\partial H_{\lambda_m}(\Gamma_m)}{\partial \lambda} \\ &= (\langle H_{\lambda_{\text{fin}}} \rangle_{\lambda_{\text{fin}}, t_{\text{fin}}} - \langle H_{\lambda_{\text{init}}} \rangle_{\lambda_{\text{init}}, t_{\text{init}}}) \sum_{m=0}^{MN-1} \frac{\Delta \lambda_{m/M}}{M} \int d\Gamma_m P_{\lambda_m}(\Gamma_m; t_m) \frac{\partial H_{\lambda_m}(\Gamma_m)}{\partial \lambda} \\ &\quad - \frac{\Delta \lambda_0}{M} (\langle H_{\lambda_{\text{fin}}} \rangle_{\lambda_{\text{fin}}, t_{\text{fin}}} - \langle H_{\lambda_{\text{init}}} \rangle_{\lambda_{\text{init}}, t_{\text{init}}}) \int d\Gamma_{\text{init}} P_{\lambda_{\text{init}}}(\Gamma_{\text{init}}; t_{\text{init}}) \frac{\partial H_{\lambda_{\text{init}}}(\Gamma_{\text{init}})}{\partial \lambda} \\ &\quad - \frac{\Delta \lambda_0}{M} \int d\Gamma_{\text{init}} P_{\lambda_{\text{init}}}(\Gamma_{\text{init}}; t_{\text{init}}) \frac{\partial H_{\lambda_{\text{init}}}(\Gamma_{\text{init}})}{\partial \lambda} \\ &= -(\langle H_{\lambda_{\text{fin}}} \rangle_{\lambda_{\text{fin}}, t_{\text{fin}}} - \langle H_{\lambda_{\text{init}}} \rangle_{\lambda_{\text{init}}, t_{\text{init}}}) \langle W \rangle \\ &\quad - \frac{\Delta \lambda_0}{M} (\langle H_{\lambda_{\text{fin}}} \rangle_{\lambda_{\text{fin}}, t_{\text{fin}}} - \langle H_{\lambda_{\text{init}}} \rangle_{\lambda_{\text{init}}, t_{\text{init}}}) \int d\Gamma_{\text{init}} P_{\lambda_{\text{init}}}(\Gamma_{\text{init}}; t_{\text{init}}) \frac{\partial H_{\lambda_{\text{init}}}(\Gamma_{\text{init}})}{\partial \lambda} \\ &\quad - \frac{\Delta \lambda_0}{M} \int d\Gamma_{\text{init}} P_{\lambda_{\text{init}}}(\Gamma_{\text{init}}; t_{\text{init}}) \frac{\partial H_{\lambda_{\text{init}}}(\Gamma_{\text{init}})}{\partial \lambda}. \end{aligned} \quad (\text{S12})$$

Note that, for the variance $\langle \Delta Q^2 \rangle \equiv \langle Q^2 \rangle - \langle Q \rangle^2$ of heat Q , the contribution from the first term in the right-hand side of Eq. (S12) cancels and only those from the second and the third terms remain, which scale as $\sim 1/M$. Therefore, from Eqs. (S10), (S11),




FIG. S1. An example of a cycle whose $\eta^{(2)}$ and $\xi^{(2)}$ is the same as those of the Carnot cycle. The temperature versus external control parameter (T - λ) diagram of this cycle working with a hot heat bath at temperature T_h and a cold one at T_c .

and (S12), the variance of heat in the quasistatic isothermal process reads

$$\begin{aligned}
 \langle \Delta Q^2 \rangle &= \left[\langle H_{\lambda_{\text{fin}}}^2 \rangle_{\lambda_{\text{fin}}, t_{\text{fin}}} - (\langle H_{\lambda_{\text{fin}}} \rangle_{\lambda_{\text{fin}}, t_{\text{fin}}})^2 \right] + \left[\langle H_{\lambda_{\text{init}}}^2 \rangle_{\lambda_{\text{init}}, t_{\text{init}}} - (\langle H_{\lambda_{\text{init}}} \rangle_{\lambda_{\text{init}}, t_{\text{init}}})^2 \right] + \langle \Delta W^2 \rangle \\
 &+ 2 \frac{\Delta \lambda_0}{M} (\langle H_{\lambda_{\text{fin}}} \rangle_{\lambda_{\text{fin}}, t_{\text{fin}}} - \langle H_{\lambda_{\text{init}}} \rangle_{\lambda_{\text{init}}, t_{\text{init}}}) \int d\Gamma_{\text{init}} P_{\lambda_{\text{init}}}(\Gamma_{\text{init}}; t_{\text{init}}) \frac{\partial H_{\lambda_{\text{init}}}(\Gamma_{\text{init}})}{\partial \lambda} \\
 &+ 2 \frac{\Delta \lambda_0}{M} \int d\Gamma_{\text{init}} P_{\lambda_{\text{init}}}(\Gamma_{\text{init}}; t_{\text{init}}) \frac{\partial H_{\lambda_{\text{init}}}(\Gamma_{\text{init}})}{\partial \lambda} \\
 &= \langle \Delta H_{\lambda_{\text{fin}}}^2 \rangle_{\lambda_{\text{fin}}, t_{\text{fin}}} + \langle \Delta H_{\lambda_{\text{init}}}^2 \rangle_{\lambda_{\text{init}}, t_{\text{init}}} + O(\tau^{-1}), \tag{S13}
 \end{aligned}$$

with $\langle \Delta H_{\lambda}^2 \rangle_{\lambda, t} \equiv \langle H_{\lambda}^2 \rangle_{\lambda, t} - (\langle H_{\lambda} \rangle_{\lambda, t})^2$. Thus, $\langle \Delta Q^2 \rangle \rightarrow \langle \Delta H_{\lambda_{\text{fin}}}^2 \rangle_{\lambda_{\text{fin}}, t_{\text{fin}}} + \langle \Delta H_{\lambda_{\text{init}}}^2 \rangle_{\lambda_{\text{init}}, t_{\text{init}}}$ in the limit of $\tau \rightarrow \infty$.

Another example giving $\eta_C^{(2)}$ and $\xi_C^{(2)}$

In the main text, we have shown that the Carnot cycle gives upper bounds $\eta_C^{(2)}$ and $\xi_C^{(2)}$ for $\eta^{(2)} \equiv \langle \Delta W^2 \rangle / \langle \Delta Q_h^2 \rangle$ and $\xi^{(2)} \equiv \langle \Delta W^2 \rangle / \langle \Delta Q^2 \rangle$, respectively. However, the Carnot cycle is not the unique case which gives these bounds. Here we provide an example whose $\eta^{(2)}$ and $\xi^{(2)}$ are the same as those for the Carnot cycle.

The example on the temperature versus external parameter (T - λ) plane is shown in Fig. S1. As can be seen from this figure, this cycle can be regarded as a combination of two Carnot cycles. At point 0, we set the external parameter at λ_0 and take a randomly chosen microstate from the canonical ensemble for H_{λ_0} at temperature T_h . Starting with this initial state, we perform the following eight strokes in numerical order of the points: two quasistatic isothermal strokes at T_h [(0 \rightarrow 1) and (4⁺ \rightarrow 5)], those at T_c [(2⁺ \rightarrow 3) and (6⁺ \rightarrow 7)], two quasistatic adiabatic expansion strokes [(1 \rightarrow 2⁻) and (5 \rightarrow 6⁻)], and two quasistatic adiabatic compression strokes [(3 \rightarrow 4⁻) and (7 \rightarrow 8⁻)]. The final point 8⁻ is equivalent to the initial point 0, and the cycle is closed. Here, the superscripts “-” and “+” respectively represent just before or after making a thermal contact with a heat bath when finishing the quasistatic adiabatic stroke.

Heat input Q_h from the hot heat bath throughout this cycle is given by

$$Q_h = Q_{0 \rightarrow 1} + Q_{4 \rightarrow 5}, \tag{S14}$$

where $Q_{0 \rightarrow 1}$ and $Q_{4 \rightarrow 5}$ are heat input through the quasistatic isothermal strokes (0 \rightarrow 1) and (4⁺ \rightarrow 5) at temperature T_h , respectively:

$$Q_{0 \rightarrow 1} = E_1 - E_0 + W_{0 \rightarrow 1}, \tag{S15}$$

$$Q_{4 \rightarrow 5} = E_5 - E_4^+ + W_{4 \rightarrow 5}. \tag{S16}$$

Here, E_i represents the internal energy of the working substance at point i , and $W_{0 \rightarrow 1}$ and $W_{4 \rightarrow 5}$ are work output through the

strokes $(0 \rightarrow 1)$ and $(4^+ \rightarrow 5)$, respectively. Since the fluctuations of $W_{0 \rightarrow 1}$ and $W_{4 \rightarrow 5}$ are negligible, we get

$$\begin{aligned}\Delta Q_h &\equiv Q_h - \langle Q_h \rangle \\ &= (\Delta E_1 - \Delta E_0) + (\Delta E_5 - \Delta E_4^+).\end{aligned}\quad (\text{S17})$$

with $\Delta E_i \equiv E_i - \langle E_i \rangle$.

Similarly, heat output Q_c to the cold heat bath throughout the cycle is

$$Q_c = -(Q_{2 \rightarrow 3} + Q_{6 \rightarrow 7}), \quad (\text{S18})$$

where $Q_{2 \rightarrow 3}$ and $Q_{6 \rightarrow 7}$ are heat input through the quasistatic isothermal strokes $(2^+ \rightarrow 3)$ and $(6^+ \rightarrow 7)$ at T_c , respectively:

$$Q_{2 \rightarrow 3} = E_3 - E_2^+ + W_{2 \rightarrow 3}, \quad (\text{S19})$$

$$Q_{6 \rightarrow 7} = E_7 - E_6^+ + W_{6 \rightarrow 7}. \quad (\text{S20})$$

Since the fluctuations of $W_{2 \rightarrow 3}$ and $W_{6 \rightarrow 7}$ are negligible, we get

$$\Delta Q_c \equiv Q_c - \langle Q_c \rangle = -[(\Delta E_3 - \Delta E_2^+) + (\Delta E_7 - \Delta E_6^+)]. \quad (\text{S21})$$

Thus the variance $\langle \Delta Q_c^2 \rangle$ of Q_c can be written as

$$\begin{aligned}\langle \Delta Q_c^2 \rangle &= \langle [(\Delta E_3 - \Delta E_2^+) + (\Delta E_7 - \Delta E_6^+)]^2 \rangle \\ &= \langle [(\Delta E_3 - \Delta E_2^-) + (\Delta E_7 - \Delta E_6^-)]^2 \rangle \\ &= \left(\frac{T_c}{T_h}\right)^2 \langle [(\Delta E_4^- - \Delta E_1) + (\Delta E_8^- - \Delta E_5)]^2 \rangle \\ &= \left(\frac{T_c}{T_h}\right)^2 \langle [(\Delta E_4^+ - \Delta E_1) + (\Delta E_0 - \Delta E_5)]^2 \rangle \\ &= \left(\frac{T_c}{T_h}\right)^2 \langle \Delta Q_h^2 \rangle.\end{aligned}\quad (\text{S22})$$

From the first to the second line, we have used the facts that E_2^\pm , E_3 , E_6^\pm , and E_7 are independent, and the distributions of E_i^+ and E_i^- ($i = 2$ and 6) are the same, which allows us to replace ΔE_i^+ by ΔE_i^- inside $\langle \dots \rangle$. From the second to the third line, we have used the relation given by Eq. (1) of the main text for the quasistatic adiabatic strokes $(1 \rightarrow 2^-)$, $(3 \rightarrow 4^-)$, $(5 \rightarrow 6^-)$, and $(7 \rightarrow 8^-)$ reading $E_2^- = (T_c/T_h)E_1$, $E_3 = (T_c/T_h)E_4^-$, $E_6^- = (T_c/T_h)E_5$, and $E_7 = (T_c/T_h)E_8^-$, respectively. Then, from the third to the fourth line, we have used the facts that E_1 , E_4^\pm , E_5 , and E_8^- are independent, E_8^- and E_0 are equivalent random variables, and the distributions of E_4^+ and E_4^- are the same.

From Eq. (S22) and the fact that Q_h and Q_c are independent, the variance $\langle \Delta Q^2 \rangle$ of the total heat input $Q \equiv Q_h - Q_c$ can be written as

$$\langle \Delta Q^2 \rangle = \langle \Delta Q_h^2 \rangle + \langle \Delta Q_c^2 \rangle = \left[1 + \left(\frac{T_c}{T_h}\right)^2\right] \langle \Delta Q_h^2 \rangle. \quad (\text{S23})$$

Next, we consider the total work output W through the whole cycle:

$$W = W_{0 \rightarrow 1} + W_{1 \rightarrow 2} + W_{2 \rightarrow 3} + W_{3 \rightarrow 4} + W_{4 \rightarrow 5} + W_{5 \rightarrow 6} + W_{6 \rightarrow 7} + W_{7 \rightarrow 8}. \quad (\text{S24})$$

Since the fluctuations of the work outputs $W_{0 \rightarrow 1}$, $W_{2 \rightarrow 3}$, $W_{4 \rightarrow 5}$, and $W_{6 \rightarrow 7}$ through the quasistatic isothermal strokes are negligible, we obtain

$$\Delta W \equiv W - \langle W \rangle = \Delta W_{1 \rightarrow 2} + \Delta W_{3 \rightarrow 4} + \Delta W_{5 \rightarrow 6} + \Delta W_{7 \rightarrow 8}. \quad (\text{S25})$$

Here, work through an adiabatic stroke is given by the difference between the internal energies of the initial and the final states of the stroke, which satisfy the relation (1) in the main text. For example, regarding the stroke $1 \rightarrow 2^-$, the work output $W_{1 \rightarrow 2}$ through this stroke can be written as

$$W_{1 \rightarrow 2} = E_1 - E_2^- = \left(1 - \frac{T_c}{T_h}\right) E_1, \quad (\text{S26})$$




FIG. S2. An example of a cycle on the T - λ plane whose $\eta^{(2)}$ can be larger than that of the Carnot cycle $\eta_C^{(2)}$. In this cycle, a quasistatic adiabatic compression stroke $1 \rightarrow 2^-$ is followed by a cold isochoric stroke $2^+ \rightarrow 3$ instead of a hot one.

and thus

$$\Delta W_{1 \rightarrow 2} = \left(1 - \frac{T_c}{T_h}\right) \Delta E_1. \quad (\text{S27})$$

Similarly, for the other quasistatic adiabatic strokes, we get $\Delta W_{3 \rightarrow 4} = [(T_c/T_h) - 1] \Delta E_4^-$, $\Delta W_{5 \rightarrow 6} [1 - (T_c/T_h)] \Delta E_5$, and $\Delta W_{7 \rightarrow 8} = [(T_c/T_h) - 1] \Delta E_8^-$. Substituting them into Eq. (S25), we obtain

$$\Delta W = \left(1 - \frac{T_c}{T_h}\right) [(\Delta E_1 - \Delta E_8^-) + (\Delta E_5 - \Delta E_4^-)]. \quad (\text{S28})$$

Since E_1 , E_4^- , E_5 , and E_8^- are independent, E_8^- and E_0 are equivalent random variables, and the distributions of E_4^+ and E_4^- are the same, we get

$$\begin{aligned} \langle \Delta W^2 \rangle &= \left(1 - \frac{T_c}{T_h}\right)^2 \langle [(\Delta E_1 - \Delta E_8^-) + (\Delta E_5 - \Delta E_4^-)]^2 \rangle \\ &= \left(1 - \frac{T_c}{T_h}\right)^2 \langle \Delta Q_h^2 \rangle. \end{aligned} \quad (\text{S29})$$

From Eqs. (S23) and (S29), we finally obtain

$$\eta^{(2)} \equiv \frac{\langle \Delta W^2 \rangle}{\langle \Delta Q_h^2 \rangle} = \left(1 - \frac{T_c}{T_h}\right)^2, \quad (\text{S30})$$

$$\xi^{(2)} \equiv \frac{\langle \Delta W^2 \rangle}{\langle \Delta Q^2 \rangle} = \frac{[1 - (T_c/T_h)]^2}{1 + (T_c/T_h)^2}, \quad (\text{S31})$$

which are the same as those for the Carnot cycle $\eta_C^{(2)}$ and $\xi_C^{(2)}$, respectively.

An example giving $\eta^{(2)} > \eta_C^{(2)}$

The relation $\eta^{(2)} \leq \eta_C^{(2)}$ holds provided all the quasistatic adiabatic expansion strokes are preceded by a stroke with a hot bath and all the quasistatic adiabatic compression strokes are followed by a stroke with a hot bath. However, if it is not the case, this relation is not guaranteed. Here we provide such an example.

Figure S2 shows the T - λ diagram of the cycle which we shall consider. At point 0, we set the external parameter at λ_0 and take a randomly chosen microstate from the canonical ensemble for H_{λ_0} at temperature T_c . Starting with this initial state, we perform the following six strokes. ($0 \rightarrow 1$): Quasistatic isothermal compression at T_c . ($1 \rightarrow 2^-$): Quasistatic adiabatic compression.

($2^+ \rightarrow 3$): Cold isochore. ($3 \rightarrow 4$): Quasistatic isothermal compression at T_c . ($4 \rightarrow 5$): Hot isochore. ($5 \rightarrow 6^-$): Quasistatic adiabatic expansion. The final point 6^- is equivalent to the initial point 0, and the cycle is closed.

We take working substance whose adiabatic invariant is in the form of

$$I_\lambda(E) \equiv \int d\Gamma \theta(E - H_\lambda(\Gamma)) = f(\lambda) E^\alpha, \quad (\text{S32})$$

with an arbitrary function f and a real constant α . In this case, the density of state $g_\lambda(E)$ reads

$$g_\lambda(E) \equiv \frac{\partial I_\lambda(E)}{\partial E} = \int d\Gamma \delta(E - H_\lambda(\Gamma)) = \alpha f(\lambda) E^{\alpha-1}. \quad (\text{S33})$$

Then, the canonical distribution $P_{\beta,\lambda}^{\text{eq}}$ at the inverse temperature $\beta \equiv 1/k_B T$ and the external parameter λ is

$$P_{\beta,\lambda}^{\text{eq}}(E) = \frac{g_\lambda(E)}{Z_{\beta,\lambda}} e^{-\beta E} = \frac{\beta^\alpha E^{\alpha-1}}{\Gamma(\alpha)} e^{-\beta E}, \quad (\text{S34})$$

with $Z_{\beta,\lambda}$ being the partition function given by

$$Z_{\beta,\lambda} = \int_0^\infty dE g_\lambda(E) e^{-\beta E} = \alpha \Gamma(\alpha) f(\lambda) \beta^{-\alpha}, \quad (\text{S35})$$

where $\Gamma(\alpha) \equiv \int_0^\infty dx x^{\alpha-1} e^{-x}$ is the gamma function.

For the canonical distribution given by Eq. (S34), the average of the internal energy E_i and of its square E_i^2 at point i with temperature T_i and parameter λ_i can be calculated as

$$\langle E_i \rangle = \int_0^\infty dE E P_{\beta_i,\lambda_i}^{\text{eq}}(E) = k_B T_i \alpha, \quad (\text{S36})$$

$$\langle E_i^2 \rangle = \int_0^\infty dE E^2 P_{\beta_i,\lambda_i}^{\text{eq}}(E) = (k_B T_i)^2 (\alpha + 1) \alpha. \quad (\text{S37})$$

Therefore, the variance $\langle \Delta E_i^2 \rangle$ of E_i reads

$$\langle \Delta E_i^2 \rangle = \langle E_i^2 \rangle - \langle E_i \rangle^2 = (k_B T_i)^2 \alpha. \quad (\text{S38})$$

Heat input Q_h from the hot heat bath is done only in the stroke $4 \rightarrow 5$ throughout the cycle: $Q_h = Q_{4 \rightarrow 5}$ and $\Delta Q_h \equiv Q_h - \langle Q_h \rangle = \Delta Q_{4 \rightarrow 5}$. Therefore, according to Table I in the main text and Eq. (S38), the variance of Q_h can be written as

$$\langle \Delta Q_h^2 \rangle = \langle \Delta Q_{4 \rightarrow 5}^2 \rangle = \langle \Delta E_4^2 \rangle + \langle \Delta E_5^2 \rangle = k_B^2 (T_c^2 + T_h^2) \alpha. \quad (\text{S39})$$

Next, work output W through the whole cycle is given by

$$W = W_{0 \rightarrow 1} + W_{1 \rightarrow 2} + W_{2 \rightarrow 3} + W_{3 \rightarrow 4} + W_{4 \rightarrow 5} + W_{5 \rightarrow 6} = W_{0 \rightarrow 1} + W_{1 \rightarrow 2} + W_{3 \rightarrow 4} + W_{5 \rightarrow 6} \quad (\text{S40})$$

because no work is done in the isochoric strokes: $W_{2 \rightarrow 3} = W_{4 \rightarrow 5} = 0$. In addition, since the fluctuations of work outputs $W_{0 \rightarrow 1}$ and $W_{3 \rightarrow 4}$ through the quasistatic isothermal strokes are negligible, we get

$$\Delta W \equiv W - \langle W \rangle = \Delta W_{1 \rightarrow 2} + \Delta W_{5 \rightarrow 6}. \quad (\text{S41})$$

Using Eq. (S38) and the expression of the work fluctuation through a quasistatic adiabatic stroke given in Table I of the main text, we get $\langle \Delta W_{1 \rightarrow 2}^2 \rangle = k_B^2 (T_c - T_2)^2 \alpha$ and $\langle \Delta W_{5 \rightarrow 6}^2 \rangle = k_B^2 (T_h - T_c) \alpha$. Since $W_{1 \rightarrow 2}$ and $W_{5 \rightarrow 6}$ are independent random variables, the variance $\langle \Delta W^2 \rangle$ of W reads

$$\langle \Delta W^2 \rangle = \langle \Delta W_{1 \rightarrow 2}^2 \rangle + \langle \Delta W_{5 \rightarrow 6}^2 \rangle = k_B^2 (T_c - T_2)^2 \alpha + k_B^2 (T_h - T_c)^2 \alpha. \quad (\text{S42})$$

From Eqs. (S39) and (S42), we finally obtain

$$\eta^{(2)} \equiv \frac{\langle \Delta W^2 \rangle}{\langle \Delta Q_h^2 \rangle} = \frac{(T_h - T_c)^2 + (T_2 - T_c)^2}{T_c^2 + T_h^2}. \quad (\text{S43})$$

This $\eta^{(2)}$ is apparently an increasing function of T_2 , and it becomes $\eta^{(2)} > \eta_C^{(2)}$ when

$$T_2 > T_c \left(2 - \frac{T_c}{T_h} \right) \equiv T_* . \quad (\text{S44})$$

Note that T_* is always smaller than T_h provided $T_c/T_h < 1$.

* gentaro@zju.edu.cn

[S1] K. Sekimoto, *Stochastic energetics*, Lecture Notes in Physics Vol. 799 (Springer, Berlin, 2010).

[S2] K. Sekimoto, *Stochastic energetics*, Japanese edition (Iwanami, Tokyo, 2004).