

# Strong Cosmic Censorship and the Universal Relaxation Bound

Shahar Hod

*The Ruppin Academic Center, Emeq Hefer 40250, Israel*

*and*

*The Hadassah Institute, Jerusalem 91010, Israel*

(Dated: October 23, 2019)

## Abstract

The strong cosmic censorship conjecture, introduced by Penrose five decades ago, asserts that, in self-consistent theories of gravity, Cauchy horizons inside dynamically formed black holes should be unstable to remnant perturbation fields that fall into the newly born black holes. The question of the (in)validity of this intriguing conjecture in non-asymptotically flat charged black-hole spacetimes has recently attracted much attention from physicists and mathematicians. We here provide a general proof, which is based on Bekenstein's generalized second law of thermodynamics, for the validity of this fundamental conjecture in non-asymptotically flat de Sitter black-hole spacetimes.

The influential singularity theorems of Hawking and Penrose [1–3] have dramatically changed our understanding of gravity and curved black-hole spacetimes. In particular, these mathematically elegant theorems have revealed the physically important fact that spacetime singularities, regions in which classical general relativity loses its predictive power, may naturally be formed by self-gravitating matter configurations whose dynamical collapse is governed by the Einstein field equations.

Following this intriguing (and highly disturbing) observation, Penrose [2, 3] has suggested that the deterministic nature of classical theories of gravity is nevertheless preserved by a mysterious “cosmic censor” which prevents spacetime singularities from being observed by distant observers. In particular, the strong version of the Penrose cosmic censorship conjecture asserts that Cauchy horizons inside black holes, which mark the deterministic boundaries of spacetime for dynamical evolutions of matter fields which are governed by the classical Einstein field equations, are singular and cannot be crossed by physical observers [2, 3].

It should be emphasized that, at present, there is no general mathematically rigorous proof for the validity of the Penrose strong cosmic censorship conjecture (SCCC) in curved spacetimes. In particular, the evidence in favor (or against) the existence of a mysterious cosmic censor who protects the deterministic power of classical general relativity is based on our (rather limited) experience in solving the non-linearly coupled Einstein-matter field equations [4–8].

In order to disprove the Penrose SCCC, many researches have tried during the last five decades to identify curved black-hole spacetimes which possess inner stable Cauchy horizons which are regular enough to allow a continuation of the spacetime metric beyond that horizons in a non-unique (that is, in a non-deterministic) way. If exist, such black-hole spacetimes would imply the catastrophic failure of the SCCC in some curved spacetimes and the corresponding breakdown of determinism in classical general relativity.

The question of the (in)validity of the Penrose SCCC in non-asymptotically flat de Sitter black-hole spacetimes has attracted much attention from physicists and mathematicians during the last couple of years [8]. In particular, it has recently been shown [8] that, in de Sitter black-hole spacetimes, the fate (singular/non-singular) of the potentially pathological inner Cauchy horizons is determined by a delicate competition between two opposite physical mechanisms:

(1) The exponential late-time *decay*  $\psi^{\text{external}}(t \rightarrow \infty) \sim e^{-\Im\omega_0 t}$  of remnant matter fields which propagate in the external spacetime regions of the dynamically formed black holes. The characteristic rate of this decay mechanism is determined by the fundamental (least damped) quasinormal resonant frequency  $\omega_0$  of the composed black-hole-field system.

(2) The exponential *amplification* (blue-shift) mechanism  $\psi^{\text{internal}}(v \rightarrow \infty) \sim e^{\kappa_- v}$  [9] which characterizes the dynamics of the remnant fields as they fall into the newly born black holes and accumulate along their inner (Cauchy) horizons. The characteristic rate of this amplification mechanism is determined by the surface gravity  $\kappa_-$  of the inner black-hole horizon.

Specifically, the (in)stability properties of the inner Cauchy horizons inside dynamically formed black holes in non-asymptotically flat de Sitter spacetimes are determined by the dimensionless ratio [8]

$$\Gamma \equiv \frac{\Im\omega_0}{\kappa_-} . \quad (1)$$

In particular, dynamically formed black holes in asymptotically de Sitter spacetimes which are characterized by the dimensionless inequality  $\Gamma > 1/2$  are pathological from the point of view of the Penrose SCCC in the sense that they contain inner Cauchy horizons which are stable and regular enough to allow a continuation of the spacetime metric in a *non-unique* (non-deterministic) way [8].

Thus, a necessary condition for classical general relativity to preserve its predictive power in non-asymptotically flat de Sitter black-hole spacetimes is given by the simple inequality

$$\Gamma \leq \frac{1}{2} . \quad (2)$$

For the advocates of the Penrose SCCC, the task remains to prove that the resonant relaxation spectra of *all* dynamically formed black holes in non-asymptotically flat curved spacetimes possess (at least) one quasinormal resonant frequency with the property [see Eqs. (1) and (2)]

$$\Im\omega_0 \leq \frac{1}{2}\kappa_- . \quad (3)$$

Interestingly, using analytical techniques, it has been explicitly proved in [10] that Kerr black holes in asymptotically de Sitter spacetimes conform to the inequality (3). These spinning curved spacetime solutions of the Einstein field equations therefore respect determinism.

As for the case of charged black holes in asymptotically de Sitter spacetimes, the situation is more involved [4–8]. In particular, it has been proved [5] that the resonant relaxation spectra of composed charged-Reissner-Nordström-de-Sitter-black-holes-charged-massive-fields systems in the dimensionless physical regime  $\mu r_+ \ll qQ \ll (\mu r_+)^2$  [here  $\{Q, r_+\}$  are respectively the electric charge and the outer horizon radius of the black hole and  $\{\mu, q\}$  are respectively the proper mass and charge coupling constant of the matter field] are characterized by the fundamental inequality (3). These charged black-hole-field systems therefore respect the Penrose SCCC.

It is clear, however, that the proof presented in [5] for the validity of the inequality (3) [11] in the regime  $\mu r_+ \ll qQ \ll (\mu r_+)^2$  is not enough. In particular, one would like to have a generic (that is, parameter-*independent*) proof that the resonant relaxation spectra of *all* dynamically formed Reissner-Nordström-de Sitter (RNdS) black holes are characterized by the fundamental inequality (3) which acts as a necessary condition for the validity of the SCCC in these non-asymptotically flat curved spacetimes.

Due to the complexity of the non-linearly coupled Einstein-charged-matter field equations, any attempt to obtain a general proof for the validity of the Penrose SCCC in charged curved spacetimes seems, at least at first glance, hopeless. In particular, it should be realized that a direct brute force test of the (in)validity of the Penrose SCCC in charged de Sitter spacetimes would require one to study (probably numerically) the non-linear dynamics of self-gravitating charged matter fields that collapse to form charged black holes. Specifically, one should scan the entire (infinitely large) phase space of the physical parameters  $\{r_-, r_+, r_c, q, \mu\}$  [12] which characterize the newly born black-hole spacetimes in order to determine whether or not there exist dynamically formed RNdS black holes that violate the fundamental inequality (3) [11, 13]. This direct approach to the problem seems to be a truly Sisyphean task!

But we need not lose heart – one should always remember that the basic physical laws of nature can provide, in a relatively straightforward way, general insights about the physical properties of relatively complex systems. One such law is Bekenstein’s generalized second law of thermodynamics [14] which asserts that the total entropy of a black-hole spacetime (that is, the entropy  $S_{\text{BH}} = A/4\hbar$  [15] of the black hole itself plus the entropy of the external matter fields) is an irreducible quantity in any physically self-consistent quantum theory of gravity.

Interestingly, and most importantly for the question of the (in)validity of the Penrose

SCCC in black-hole spacetimes, it has been explicitly shown in [16] that Bekenstein’s generalized second law of thermodynamics yields, in a rather straightforward way, a fundamental upper bound on the characteristic relaxation rates of dynamical physical systems with a well defined thermodynamic description. Specifically, for a thermodynamic system which is characterized by a temperature  $T$ , the universal relaxation bound can be expressed by the remarkably compact time-times-temperature (TTT) quantum relation [16]

$$\tau \times T \geq \frac{\hbar}{\pi} , \quad (4)$$

where  $\tau$  is the characteristic relaxation time of the thermodynamic system.

Remarkably, curved black-hole spacetimes are known to be characterized by a well defined temperature, the Bekenstein-Hawking temperature [14, 17]

$$T_{\text{BH}} = \frac{\kappa_+}{2\pi} \cdot \hbar . \quad (5)$$

Taking cognizance of the universal relaxation bound (4) and the quantum Bekenstein-Hawking relation (5) one immediately deduces that the resonant relaxation spectra of *all* dynamically formed black holes are characterized by the simple upper bound [18]

$$\Im\omega_0 \leq \frac{1}{2}\kappa_+ , \quad (6)$$

where  $\omega_0$  is the fundamental (least damped) resonant frequency of the composed dynamically-formed-black-hole-remnant-field system.

What we find most interesting is the fact that, in black-hole physics where  $T_{\text{BH}} \propto \hbar$  [see Eq. (5)], the *quantum* lower bound (4) becomes a *classical* (that is,  $\hbar$ -independent) relation between the characteristic relaxation rates of the dynamically formed black-hole spacetimes and the corresponding black-hole surface gravities.

*Summary.*— The conjectured strong cosmic censorship principle [1–3] has attracted much attention from physicists and mathematicians since its introduction by Penrose five decades ago. In particular, the question of the (in)validity of this highly influential conjecture in charged de Sitter black-hole spacetimes has been the focus of an intense debate in the last couple of years (see [4–7] and references therein).

The SCCC asserts that, in self-consistent theories of gravity, spacetime metrics inside physically realistic (*dynamically* formed) black holes cannot be extended in a non-unique (non-deterministic) way beyond the inner Cauchy horizons. For this to happen, the inner

Cauchy horizons of dynamically formed black-hole spacetimes should be unstable to remnant perturbation fields that fall into the newly born black holes. In particular, a necessary condition for the validity of the SCCC in asymptotically de Sitter black-hole spacetimes is provided by the simple inequality (3).

A direct way to prove the validity of the fundamental inequality (3) would be to scan numerically the infinitely large phase space of the resonant spectra which characterize the relaxation of dynamically formed black-hole spacetimes. Instead of following this truly Sisyphean (and numerically time consuming) task, we have pointed out that the Bekenstein generalized second law of thermodynamics [14], a physical law which is widely believed to reflect a fundamental aspect of a physically self-consistent quantum theory of gravity, yields the upper relaxation bound (6) [16, 19, 20]. Using the characteristic inequality  $\kappa_+ \leq \kappa_-$  for the surface gravities of the black-hole spacetime [4], one immediately realizes that the relaxation spectra of all dynamically formed black holes are indeed characterized by the desired inequality (3).

We therefore conclude that physically realistic [19, 20] black holes in asymptotically de Sitter spacetimes *respect* determinism and conform to the Penrose strong cosmic censorship conjecture.

## ACKNOWLEDGMENTS

This research is supported by the Carmel Science Foundation. I would like to thank Yael Oren, Arbel M. Ongo, Ayelet B. Lata, and Alona B. Tea for helpful discussions.

- 
- [1] S. W. Hawking and R. Penrose, Proc. R. Soc. Lond. **A314**, 529 (1970).
- [2] R. Penrose, Riv. Nuovo Cimento I **1**, 252 (1969).
- [3] R. Penrose in *General Relativity, an Einstein Centenary Survey*, eds. S.W. Hawking and W. Israel (Cambridge University Press, 1979).
- [4] C. Chambers, arXiv:gr-qc/9709025.
- [5] S. Hod, Nucl. Phys. B **941**, 636 (2019) [arXiv:1801.07261].
- [6] S. Hod, arXiv:1810.04853.
- [7] B. Ge, J. Jiang, B. Wang, H. Zhang, and Z. Zhong, JHEP **01**, 123 (2019) [arXiv:1810.12128].
- [8] See details in [4–7] and references therein.
- [9] We use gravitational units in which  $G = c = 1$ . Here  $v$  is the standard advanced null coordinate.
- [10] See S. Hod, Phys. Lett. B **780**, 221 (2018) [arXiv:1803.05443] and references therein.
- [11] It is worth emphasizing again that the inequality (3) serves as a necessary condition for the validity of the Penrose SCCC in de Sitter black-hole spacetimes.
- [12] Here  $\{r_-, r_+, r_c\}$  are respectively the inner (Cauchy), outer (event), and cosmological horizons of the non-asymptotically flat RNdS black-hole spacetime.
- [13] It is worth emphasizing that we consider here only physically realistic black holes that can be formed *dynamically* from spatially regular self-gravitating matter fields that evolve (collapse) according to the Einstein field equations. It is a rather trivial task to find sets of parameters  $\{q, \mu\}$  which correspond to matter fields whose resonant spectra in the background of pre-existing black holes (that is, eternal black-hole spacetimes that were just postulated by hand and were not formed dynamically during a physically realistic collapse scenario) violate the relation (3). For instance, the resonant spectra of neutral fields in the background of highly charged (near-extremal,  $r_- \simeq r_+$ ) RNdS black holes are characterized by the relation  $\Im\omega_0 > \kappa_-/2$ . However, it should be realized that charged black holes cannot be formed dynamically from the collapse of neutral fields, and thus the composed charged-RNdS-black-hole-neutral-field system with the corresponding set of parameters  $\{r_- \simeq r_+, r_+, r_c, q = 0, \mu\}$  *cannot* provide a physically realistic counter-example to the Penrose SCCC.
- [14] J. D. Bekenstein, Phys. Rev. D **7**, 2333 (1973).
- [15] Here  $A$  is the black-hole surface area.

- [16] S. Hod, Phys. Rev. D **75**, 064013 (2007) [arXiv:gr-qc/0611004].
- [17] S. W. Hawking, Commun. Math. Phys. **43**, 199 (1975).
- [18] Here we have used the relation  $\tau_{\text{relax}} \equiv 1/\Im\omega_0$  for the characteristic relaxation time of the dynamically formed black-hole spacetime, where  $\omega_0$  is the fundamental (least damped) resonant frequency of the composed black-hole-remnant-field system.
- [19] It is important to stress that the compact relaxation bound (6) should be regarded as a fundamental property of physically realistic black holes. That is, the relation (6) characterizes black-hole spacetimes which *can* be formed *dynamically* during a non-linear gravitational collapse which starts from physically realistic (spatially regular, horizonless) initial conditions. It should be pointed out that, as shown in the physically interesting work [7], matter fields with small enough values of the charge coupling parameter  $q$  can violate the relation (6) in near-extremal (highly charged) black-hole spacetimes. Does it mean that these weakly charged fields violate the fundamental Penrose SCCC? The answer to this important question is ‘no’. In particular, the fundamental bound (6), which is a direct consequence of Bekenstein’s generalized second law of thermodynamics [14], a law whose very meaning stems from a *quantum* theory of gravity, asserts that these highly charged (near-extremal) black-hole spacetimes cannot be formed dynamically during a non-linear gravitational collapse (which starts from spatially regular horizonless initial conditions) of these weakly charged fields. In addition, within the framework of a self-consistent *quantum* theory of gravity, one should take into account the well-known phenomenon of the quantum Schwinger discharge of charged black holes [20], a physical mechanism which tends to prevent the dynamical formation of highly charged black-hole spacetimes that could risk the validity of both the Bekenstein generalized second law of thermodynamics [14] and the Penrose SCCC [2, 3].
- [20] W. Heisenberg and H. Euler, Z. Phys. **98**, 714 (1936); J. Schwinger, Phys. Rev. **82**, 664 (1951); M. A. Markov and V. P. Frolov, Teor. Mat. Fiz. **3**, 3 (1970); W. T. Zaumen, Nature **247**, 531 (1974); B. Carter, Phys. Rev. Lett. **33**, 558 (1974); G. W. Gibbons, Comm. Math. Phys. **44**, 245 (1975); T. Damour and R. Ruffini, Phys. Rev. Lett. **35**, 463 (1975).