

# Geometric Flows of Curves, Two-Component Camassa-Holm Equation and Generalized Heisenberg Ferromagnet Equation

Aigul Taishiyeva\*, Tolkynay Myrzakul†, Gulgassyl Nugmanova‡, Shynaray Myrzakul§, Kuralay Yesmakhanova¶ and Ratbay Myrzakulov||

*Eurasian International Center for Theoretical Physics and  
Department of General & Theoretical Physics,  
Eurasian National University, Nur-Sultan, 010008, Kazakhstan*

## Abstract

In this paper, we study the generalized Heisenberg ferromagnet equation, namely, the M-CVI equation. This equation is integrable. The integrable motion of the space curves induced by the M-CVI equation is presented. Using this result, the Lakshmanan (geometrical) equivalence between the M-CVI equation and the two-component Camassa-Holm equation is established.

**Mathematics Subject Classifications(2000).** 35Q53, 37K35

**Key words:** Camassa-Holm equation, AKNS hierarchy, reciprocal transformation, Heisenberg ferromagnet equation, spin systems, motion of space curves, differential geometry of curves and surfaces.

## 1 Introduction

The celebrated Camassa-Holm equation (CHE) has the form

$$u_t + \kappa u_x - u_{xxt} + 3uu_x = 2u_x u_{xx} + uu_{xxx}, \quad (1.1)$$

where  $u = u(x, t)$  is the fluid velocity in the  $x$  direction and  $\kappa = const$  is related to the critical shallow water wave speed. The CHE shares most of the important properties of integrable equations like the  $N$ -soliton solutions, the bi-Hamiltonian structure, the Lax representation (LR) and so on. In the case, when  $\kappa = 0$ , the CHE (1.1) has the so-called peakon solutions. Several important generalizations of the CHE including integrable cases but many other (non-integrable or whose integrability has not been determined) have been discovered [1]-[29]. In particular, the two-component CHE (2-CHE) was constructed. Our main interest in this paper is to go further with the investigation initiated in our previous papers (see, e.g. [65]-[70]). In this paper, we study the 2-CHE, its relation with the geometry of space curves and the equivalent spin system.

The paper is organized as follows. In Section 2 we present main fact for the M-CVI equation. Basic information on the 2-CHE we give in Section 3. The integrable motion of space curves induced by the M-CVI equation and the 2-CHE is studied in Section 4. In Section 4, we consider the gauge

---

\*Email: ataishiyeva@gmail.com

†Email: trmyrzakul@gmail.com

‡Email: nugmanovagn@gmail.com

§Email: srmyrzakul@gmail.com

¶Email: kryesmakhanova@gmail.com

||Email: rmyrzakulov@gmail.com

equivalence between the M-CVI equation and the 2-CHE. Finally, in Section 5 we present a discussion of our achievements and how they impact some recent results found in the recent literature.

## 2 M-CVI equation

There are several integrable and non-integrable generalized Heisenberg ferromagnet equations (gHFE) (see, e.g. [30]-[70]). In this paper, we consider one of the gHFE, namely, the M-CVI equation. The M-CVI equation is integrable. It shares most of the important properties of integrable systems like the Lax representation (LR), the bi-hamiltonian structure, the  $N$ -soliton solutions, infinite hierarchy of symmetries and conservation laws and so on. The M-CVI equation also can admits the so-called peakon solutions.

### 2.1 Equation

Consider the M-CVI equation

$$[A, A_{xt} + (uA_x)_x] - \frac{1}{\beta^2}A_x - 4\beta\rho\rho_x Z = 0. \quad (2.1)$$

Here  $m = \frac{\det(A_x^2)}{4\beta^2} = u - u_{xx}$ ,  $\rho^2 = -\frac{\text{tr}(A_x^2) + 2\det(A_x)}{8\beta^4}$ ,  $u = 0.25\beta^{-2}(1 - \partial_x^2)^{-1} \det(A_x^2)$  are some real functions,  $\beta = \text{const}$  and

$$Z = \frac{0.5\beta}{u_x + u_{xx}}[A, A_t + (u - 0.5\beta^{-2})A_x], \quad \mathbf{A} = (A_1, A_2, A_3), \quad (2.2)$$

$$A = \begin{pmatrix} A_3 & A^- \\ A^+ & -A_3 \end{pmatrix}, \quad A^\pm = A_1 \pm iA_2, \quad A^2 = I, \quad \mathbf{A}^2 = 1. \quad (2.3)$$

### 2.2 Lax representation

The LR of the M-CVI equation reads as

$$\Psi_x = U_1 \Psi, \quad (2.4)$$

$$\Psi_s = V_1 \Psi. \quad (2.5)$$

Here

$$U_1 = \left( \frac{\lambda}{4\beta} - \frac{1}{4} \right) [A, A_x] + (\lambda^3 - \beta^2\lambda)\rho^2 Z, \quad (2.6)$$

$$V_1 = \left( \frac{1}{4\beta^2} - \frac{1}{4\lambda^2} \right) A + \frac{u}{4} \left( \frac{\beta}{\lambda} - \frac{\lambda}{\beta} \right) [A, A_x] + \left( \frac{\beta}{4\lambda} - \frac{1}{4} \right) [A, A_t] + v\rho^2 Z \quad (2.7)$$

where  $v = \lambda(0.5 + \beta^2 u) - \lambda^3 u - 0.5\beta^2 \lambda^{-1}$ . The compatibility condition  $U_{1t} - V_{1x} + [U_1, V_1] = 0$  is equivalent to the M-CVI equation (2.1).

### 2.3 Reductions

One of the reductions of the M-CVI equation is the so-called M-CIV equation. Let  $\rho = 0$ . Then the M-CVI equation takes the form

$$[A, A_{xt} + (uA_x)_x] - \frac{1}{\beta^2}A_x = 0. \quad (2.8)$$

It is nothing but the M-CIV equation (see, e.g. [65]-[70]). The LR of the M-CIV equation follows from the LR of the M-CVI equation (2.4)-(2.5) as  $\rho = 0$ . So that the LR of the M-CIV equation is given by [65]

$$\Psi_x = U_3 \Psi, \quad (2.9)$$

$$\Psi_s = V_3 \Psi, \quad (2.10)$$

where

$$U_3 = \left( \frac{\lambda}{4\beta} - \frac{1}{4} \right) [A, A_x], \quad (2.11)$$

$$V_3 = \left( \frac{1}{4\beta^2} - \frac{1}{4\lambda^2} \right) A + \frac{u}{4} \left( \frac{\beta}{\lambda} - \frac{\lambda}{\beta} \right) [A, A_x] + \left( \frac{\beta}{4\lambda} - \frac{1}{4} \right) [A, A_t]. \quad (2.12)$$

### 3 2-CHE

#### 3.1 Equation

The two-component CHE (2-CHE) is given by [1]

$$m_t + um_x + 2mu_x - \rho\rho_x = 0, \quad (3.1)$$

$$\rho_t + (\rho u)_x = 0, \quad (3.2)$$

where  $m = u - u_{xx} + 0.5\kappa$ . If  $\rho = 0$ , the 2-CHE reduces to the CHE (1.1).

#### 3.2 Lax representation

The LR of the 2-CHE is given by (see, e.g. [1])

$$\phi_{xx} = \left( \frac{1}{4} - m\zeta + \rho^2\zeta^2 \right) \phi, \quad (3.3)$$

$$\phi_t = -\left( \frac{1}{2\zeta} + u \right) \phi_x + \frac{u_x}{2} \phi, \quad (3.4)$$

where  $\zeta$  is a spectral parameter and  $m = u - u_{xx} + 0.5\kappa$  ( $\kappa = \text{const}$ ).

#### 3.3 Reciprocal transformation

From the equation (3.2) follows that the 1-form

$$\omega = \rho dx - \rho u dt \quad (3.5)$$

is closed. This means that we can define a reciprocal transformation  $(x, t) \mapsto (y, s)$  by the relation [1]

$$dy = \rho dx - \rho u dt, \quad ds = dt. \quad (3.6)$$

So we obtain

$$\frac{\partial}{\partial x} = \rho \frac{\partial}{\partial y}, \quad \frac{\partial}{\partial t} = \frac{\partial}{\partial s} - \rho u \frac{\partial}{\partial y}. \quad (3.7)$$

Then the spectral problem (3.3)-(3.4) takes the form [1]

$$\varphi_{yy} = (\lambda^2 - P\lambda - Q) \varphi, \quad (3.8)$$

$$\varphi_s = -\frac{\rho}{2\lambda} \varphi_y + \frac{\rho_y}{4\lambda} \varphi, \quad (3.9)$$

where

$$\varphi = \sqrt{\rho} \phi, \quad P = \frac{m}{\rho^2}, \quad Q = -\frac{1}{4\rho^2} - \frac{\rho_{yy}}{2\rho} + \frac{\rho_y^2}{4\rho^2}. \quad (3.10)$$

The compatibility condition of the equations (3.8)-(3.9) gives

$$P_s = \rho_y, \quad (3.11)$$

$$Q_s + \frac{1}{2}\rho P_y + P\rho_y = 0, \quad (3.12)$$

$$\frac{1}{2}\rho Q_y + Q\rho_y + \frac{1}{4}\rho_{yyy} = 0. \quad (3.13)$$

Hence we get the equation [1]

$$\rho^2 Q + \frac{1}{2}\rho \rho_{yy} - \frac{1}{4}\rho_y^2 = C = -\frac{1}{4}. \quad (3.14)$$

From the equation (3.11) follows

$$P = \frac{\partial f(y, s)}{\partial y}, \quad \rho = \frac{\partial f(y, s)}{\partial s}, \quad (3.15)$$

where  $f(y, s)$  is some function. This function satisfies the equation [1]

$$\frac{f_{ss}}{2f_s^3} + f_y f_{ys} - \frac{f_{ss} f_{ys}^2}{2f_s^3} + \frac{f_{ys} f_{yss}}{2f_s^2} + \frac{1}{2} f_s f_{yy} + \frac{f_{ss} f_{yys}}{2f_s^2} - \frac{f_{yyss}}{2f_s} = 0. \quad (3.16)$$

Finally we come to the following theorems [1]

**Theorem 3.1** *Let  $f$  be a solution of the equation (3.16), and*

$$u = f_y f_s^2 + \frac{f_{ss} f_{ys}}{f_s} - f_{yss}, \quad \rho = f_s. \quad (3.17)$$

*If  $x(y, s)$  is a solution of the following system of ODEs:*

$$\frac{\partial x}{\partial y} = \frac{1}{\rho}, \quad \frac{\partial x}{\partial s} = u, \quad (3.18)$$

*then  $(u(y, t), \rho(y, t), x(y, t))$  is a parametric solution of the 2-CHE (3.1)-(3.2).*

**Theorem 3.2** *Let  $f(y, s)$  be a solution of the equation (3.16). Define the functions  $x = x(y, s), u = u(y, s), \rho = \rho(y, s)$  by*

$$x = f(s, y), \quad u = \frac{\partial x}{\partial s}, \quad \frac{1}{\rho} = \frac{\partial x}{\partial y}. \quad (3.19)$$

*Then  $(u(y, t), \rho(y, t), x(y, t))$  is a parametric solution of the 2-CHE (3.1)-(3.2).*

### 3.4 Relations to the first negative flow of the AKNS hierarchy

Let us briefly present the well-known result about relation between the 2-CHE and the AKNS spectral problem following the paper [1]. The AKNS spectral problem reads as

$$\begin{pmatrix} \phi_1 \\ \phi_2 \end{pmatrix}_y = \begin{pmatrix} \lambda & -q \\ r & -\lambda \end{pmatrix} \begin{pmatrix} \phi_1 \\ \phi_2 \end{pmatrix}. \quad (3.20)$$

The first negative flow of the AKNS problem is given by

$$\begin{pmatrix} \phi_1 \\ \phi_2 \end{pmatrix}_s = \frac{1}{4\lambda} \begin{pmatrix} a & b \\ c & -a \end{pmatrix} \begin{pmatrix} \phi_1 \\ \phi_2 \end{pmatrix}. \quad (3.21)$$

The compatibility condition of the equations (3.20)-(3.21) gives

$$q_s = \frac{1}{2} b, \quad (3.22)$$

$$r_s = \frac{1}{2} c, \quad (3.23)$$

$$b_y = 2 a q, \quad (3.24)$$

$$c_y = 2 a r, \quad (3.25)$$

$$a_y + b r + c q = 0. \quad (3.26)$$

Hence we get the condition [1]

$$a^2 + b c = \varepsilon^2, \quad (3.27)$$

where  $\varepsilon = \text{const}$ . We have the following theorems [1]:

**Theorem 3.3** *Let  $(a, b, c, q, r)$  be a solution of the equations (3.23)–(3.26) with  $\varepsilon^2 = 1$ , then any function  $f(y, s)$  satisfying*

$$2a = b e^{-f} - c e^f \quad (3.28)$$

*gives a primary solution of the 2-CH system.*

**Theorem 3.4** *If  $f$  is a primary solution of the 2-CHE system (3.1)–(3.2), then we can construct a solution of the first negative flow of the AKNS hierarchy by the following formulae*

$$q = \frac{e^f}{2} \left( f_y + \frac{\varepsilon - f_{ys}}{f_s} \right), \quad r = \frac{e^{-f}}{2} \left( f_y - \frac{\varepsilon - f_{ys}}{f_s} \right), \quad b = 2 q_s, \quad c = 2 r_s, \quad a = \frac{b e^{-f} - c e^f}{2}. \quad (3.29)$$

where  $\varepsilon = 1$  or  $\varepsilon = -1$ .

### 3.5 Bi-Hamiltonian structure

Let us again briefly present some basic facts about the bi-Hamiltonian structure of the 2-CHE following the paper [1]. Note that both bi-Hamiltonian structures of the CHE and the KdV hierarchies are deformations of the following bi-Hamiltonian structure of hydrodynamic type [1]

$$\begin{aligned} \{u(x), u(y)\}_1 &= \delta'(x - y), \\ \{u(x), u(y)\}_2 &= u(x)\delta'(x - y) + \frac{1}{2} u(x)'\delta(x - y). \end{aligned} \quad (3.30)$$

This means that the dispersionless limits of the CHE and KdV hierarchies have the same forms. One of the main features of the integrable hierarchies that correspond to bi-Hamiltonian structures with constant central invariants is the existence of  $\tau$ -functions. It is well known that integrable hierarchies of nonlinear partial differential equations with one spatial variable possess bi-Hamiltonian structures that are deformations of bi-Hamiltonian structure of hydrodynamic type with constant central invariants, and the existence of  $\tau$ -functions plays an important role in the study of these integrable systems. The CHE hierarchy is an exceptional example of integrable systems which does not possess  $\tau$ -functions.

## 4 Motion of space curves induced by the M-CIV equation. Lakshmanan (geometrical) equivalence

In this section, we would like to find the integrable motion of the space curves induced by the M-CVI equation. To do that, let us consider a smooth space curve in  $R^3$  given by

$$\gamma(x, t) : [0, X] \times [0, T] \rightarrow R^3, \quad (4.1)$$

where  $x$  is the arc length of the curve at each time  $t$ . Then the following three vectors

$$\mathbf{e}_1 = \gamma_x, \quad \mathbf{e}_2 = \frac{\gamma_{xx}}{|\gamma_{xx}|}, \quad \mathbf{e}_3 = \mathbf{e}_1 \wedge \mathbf{e}_2, \quad (4.2)$$

are the unit tangent vector, the principal normal vector and the binormal vector of the curve, respectively. The corresponding Frenet-Serret equation is given by

$$\begin{pmatrix} \mathbf{e}_1 \\ \mathbf{e}_2 \\ \mathbf{e}_3 \end{pmatrix}_x = C \begin{pmatrix} \mathbf{e}_1 \\ \mathbf{e}_2 \\ \mathbf{e}_3 \end{pmatrix} = \begin{pmatrix} 0 & \kappa_1 & \kappa_2 \\ -\kappa_1 & 0 & \tau \\ -\kappa_2 & -\tau & 0 \end{pmatrix} \begin{pmatrix} \mathbf{e}_1 \\ \mathbf{e}_2 \\ \mathbf{e}_3 \end{pmatrix}, \quad (4.3)$$

where  $\tau$ ,  $\kappa_1$  and  $\kappa_2$  are torsion, geodesic curvature and normal curvature of the curve, respectively. Let the deformation of the curves are given by

$$\begin{pmatrix} \mathbf{e}_1 \\ \mathbf{e}_2 \\ \mathbf{e}_3 \end{pmatrix}_x = C \begin{pmatrix} \mathbf{e}_1 \\ \mathbf{e}_2 \\ \mathbf{e}_3 \end{pmatrix}, \quad \begin{pmatrix} \mathbf{e}_1 \\ \mathbf{e}_2 \\ \mathbf{e}_3 \end{pmatrix}_t = G \begin{pmatrix} \mathbf{e}_1 \\ \mathbf{e}_2 \\ \mathbf{e}_3 \end{pmatrix}. \quad (4.4)$$

Here

$$C = -\tau L_1 + \kappa_2 L_2 - \kappa_1 L_3 = \begin{pmatrix} 0 & \kappa_1 & \kappa_2 \\ -\kappa_1 & 0 & \tau \\ -\kappa_2 & -\tau & 0 \end{pmatrix}, \quad (4.5)$$

$$G = -\omega_1 L_1 + \omega_2 L_2 - \omega_3 L_3 = \begin{pmatrix} 0 & \omega_3 & \omega_2 \\ -\omega_3 & 0 & \omega_1 \\ -\omega_2 & -\omega_1 & 0 \end{pmatrix}, \quad (4.6)$$

where

$$L_1 = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{bmatrix}, \quad L_2 = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{bmatrix}, \quad L_3 = \begin{bmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \quad (4.7)$$

are basis elements of  $so(3)$  algebra. The compatibility condition of the equations (4.4) has the form

$$C_t - G_x + [C, G] = 0 \quad (4.8)$$

or in components

$$\kappa_{1t} - \omega_{3x} - \kappa_2 \omega_1 + \tau \omega_2 = 0, \quad (4.9)$$

$$\kappa_{2t} - \omega_{2x} + \kappa_1 \omega_1 - \tau \omega_3 = 0, \quad (4.10)$$

$$\tau_t - \omega_{1x} - \kappa_1 \omega_2 + \kappa_2 \omega_3 = 0. \quad (4.11)$$

As usual, we consider the following identification  $\mathbf{A} \equiv \mathbf{e}_1$ . We have

$$\kappa_1 = i, \quad \kappa_2 = \lambda(m-1) + \lambda^3 \rho^2, \quad \tau = -i[\lambda(m+1) + \lambda^3 \rho^2], \quad (4.12)$$

where  $\kappa_2 + i\tau = 2m\lambda + 2\rho^2\lambda^3$  and  $\kappa_2 - i\tau = -2\lambda$ ,  $\lambda = \text{const}$ . Finally we get the following expressions for the functions  $\omega_j$

$$\omega_1 = i[(u\lambda - 0.5\lambda^{-1})(m+1) - 0.5\lambda^{-1}(u_x + u_{xx}) - 0.5\lambda\rho^2 + \lambda^3u\rho^2], \quad (4.13)$$

$$\omega_2 = [(0.5\lambda^{-1} - \lambda u)(m-1) + 0.5\lambda^{-1}(u_x + u_{xx}) + 0.5\lambda\rho^2 - \lambda^3u\rho^2], \quad (4.14)$$

$$\omega_3 = i[0.5\lambda^{-2} - u - u_x]. \quad (4.15)$$

Substituting these expressions to Eqs.(4.9)-(4.11) we get the following equations for  $m, \rho$ :

$$m_t + 2u_x m + um_x - \rho\rho_x = 0, \quad (4.16)$$

$$\rho_t + (u\rho)_x = 0, \quad (4.17)$$

$$m - u + u_{xx} - 0.5\kappa = 0, \quad (4.18)$$

which is the 2-CHE. So, we have proved the Lakshmanan (geometrical) equivalence between the M-CVI equation (2.1) and the 2-CHE (3.1)-(3.2).

## 5 Gauge equivalence

In the previous section, we have shown that the M-CIV equation (2.1) and the 2-CHE (3.1)-(3.2) are the geometrical equivalent each to other. As it was established in [67], between these equations takes place also the gauge equivalence. In fact, consider the gauge transformation  $\Phi = g\Psi$ , where  $g = \Phi|_{\lambda=\beta}$ . Then we have

$$U_1 = g^{-1}U_2g - g^{-1}g_x, \quad V_1 = g^{-1}V_2g - g^{-1}g_t. \quad (5.1)$$

As result, we get the following LR for the 2-CHE

$$\Phi_x = U_2\Phi, \quad (5.2)$$

$$\Phi_t = V_2\Phi, \quad (5.3)$$

where

$$U_2 = \begin{pmatrix} -0.5 & \lambda \\ m\lambda + \rho^2\lambda^3 & 0.5 \end{pmatrix}, \quad (5.4)$$

$$V_2 = \begin{pmatrix} 0.5(u + u_x) - 0.25\lambda^{-2} & 0.5\lambda^{-1} - u\lambda \\ 0.5(m + u_x + u_{xx})\lambda^{-1} - um + 0.5\rho^2\lambda - u\rho^2\lambda^3 & 0.25\lambda^{-2} - 0.5(u + u_x) \end{pmatrix}. \quad (5.5)$$

The compatibility condition

$$U_{2t} - V_{2x} + [U_2, V_2] = 0 \quad (5.6)$$

gives the 2-CHE.

## 6 Conclusion

In this paper, we have considered the M-CVI equation and the 2-CHE. First, we have presented some well-known main facts on these equations. In particular, we briefly present the reciprocal transformation between the 2-CHE and the first negative flow of the AKNS hierarchy which includes in particular the well known sine-Gordon and the sinh-Gordon equations. This transformation gives the correspondence between solutions of the first negative flow of the AKNS hierarchy and the 2-CHE. Then we have studied the motion of the space curves induced by these equations. Using this result, we have

proved that the M-CVI equation and the 2-CHE is the Lakshmanan (geometrical) equivalent each to other. Last but not least, we would like to note and believe that the "spinalization" of integrable systems gives some new informations on their nature.

**Acknowledgments.** This work was supported by the Ministry of Education and Science of Kazakhstan under grants 011800935 and 011800693.

## References

- [1] Ming Chen, Si-Qi Liu, Youjin Zhang. *A 2-Component Generalization of the Camassa-Holm Equation and Its Solutions*, [arXiv:nlin/0501028]
- [2] Z. J. Qiao. *The Camassa-Holm Hierarchy, N-Dimensional Integrable Systems, and Algebro-Geometric Solution on a Symplectic Submanifold*, Commun. Math. Phys., **239**, 309341 (2003)
- [3] Z. J. Qiao, G. P. Zhang. *On peaked and smooth solitons for the Camassa-Holm equation*, Europhys. Lett., **73**, N5, 657663 (2006)
- [4] . Zhijun Qiao, Xia B., Li J. . An integrable system with peakon, complex peakon, weak kink, and kink-peakon interactional solutions. Communications in Nonlinear Science and Numerical Simulation, **63**, 292 - 306 (2018)
- [5] Zhijun Qiao, Wang F., Li F. *On the Cauchy problem for a higher-order mu-Camassa-Holm equation*, Discrete & Continuous Dynamical Systems - A, **38**, N8, 4163-4187 (2018)
- [6] Abenta S., Grava T., Modulation of Camassa-Holm equation and reciprocal transformations, preprint 2004.
- [7] Alber M. S., Camassa R., Fedorov Yu. N., Holm D. D., Marsden J. E., The complex geometry of weak piecewise smooth solutions of integrable nonlinear PDEs of shallow water and Dym type, Comm. Math. Phys. **221** (2001) 197-227.
- [8] Antonowicz M., Fordy A. P., Coupled KdV equations with multi-Hamiltonian structures. Phys. D **28** (1987) 345-357.
- [9] Antonowicz M., Fordy A. P., Coupled Harry Dym equations with multi-Hamiltonian structures, J. Phys. A **21** (1988) L269-L275.
- [10] Antonowicz M., Fordy A. P., Factorisation of energy dependent Schrödinger operators: Miura maps and modified systems. Comm. Math. Phys. **124** (1989) 465-486.
- [11] Camassa R. and Holm D. D., An integrable shallow water equation with peaked solitons, Phys. Rev. Lett. **71** (1993) 1661-1664.
- [12] Camassa R., Holm D. D. and Hyman J. M., A new integrable shallow water equation, Adv. Appl. Mech. **31** (1994), 1-33.
- [13] Constantin A., On the scattering problem for the Camassa-Holm equation. R. Soc. Lond. Proc. Ser. A Math. Phys. Eng. Sci. **457** (2001), 953-970.
- [14] Degasperis A., Holm D. D., Hone A. N. W., Integrable and non-integrable equations with peakons. Nonlinear physics: theory and experiment, II (Gallipoli, 2002), 37-43, World Sci. Publishing, River Edge, NJ, 2003.
- [15] Dubrovin B., Zhang Y., Normal forms of integrable PDEs, Frobenius manifolds and Gromov-Witten invariants, math.DG/0108160.



- [16] Dubrovin B., Liu S. Q., Zhang Y., On Hamiltonian perturbations of hyperbolic systems of conservation laws, math.DG/0410027.
- [17] Falqui G., On a two-component generalization of the CH equation, talk given at the conference “Analytic and geometric theory of the Camassa-Holm equation and Integrable systems”, Bologna, September 22- 25, 2004.
- [18] Fokas A. S., On a class of physically important integrable equations, *Physica* **D 87**(1995) 145-150.
- [19] Fuchssteiner B., Some tricks from the symmetry-toolbox for nonlinear equations: Generalizations of the Camassa-Holm equation, *Physica* **D 95** (1996) 229-243.
- [20] Hone A. N. W., The associated Camassa-Holm equation and the KdV equation. *J. Phys. A* **32** (1999), L307–L314.
- [21] Johnson R. S., On solutions of the Camassa-Holm equation. *R. Soc. Lond. Proc. Ser. A Math. Phys. Eng. Sci.* **459** (2003), 1687–1708.
- [22] Li Y. S., Zhang J. E., The multiple-soliton solution of the Camassa-Holm equation, *R. Soc. Lond. Proc. Ser. A Math. Phys. Eng. Sci.* **460** (2004),
- [23] Li Y. S., Zhang J. E., Analytical multiple-soliton solution of the Camassa-Holm equation, preprint 2004, to appear in *J. Nonlinear Math. Phys.*
- [24] McKean H., The Liouville correspondence between the Korteweg-de Vries and the Camassa-Holm hierarchies. Dedicated to the memory of Jrgen K. Moser. *Comm. Pure Appl. Math.* **56** (2003), 998–1015.
- [25] McKean H., Breakdown of the Camassa-Holm equation. *Comm. Pure Appl. Math.* **57** (2004), 416–418.
- [26] Schiff J., The Camassa-Holm equation: a loop group approach. *Phys. D* **121** (1998), no. 1-2, 24–43.
- [27] Chang X-K., Hu X-B., Szmigiels J. *Multipeakons of a two-component modified Camassa-Holm equation and the relation with the finite Kac-Van Moerbeke lattice*, [arXiv:1512.08300]
- [28] Ivanov R.I. *Two component integrable systems modelling shallow water waves: the constant vorticity case*, [arXiv:0906.0780]
- [29] Hone A.N.W., Wang J.P. *Integrable peakon equations with cubic nonlinearity*, [arXiv:0805.4310]
- [30] G. Nugmanova, Z. Zhunussova, K. Yesmakhanova, G. Mamyrbekova, R. Myrzakulov. *International Journal of Mathematical, Computational, Statistical, Natural and Physical Engineering*, **9**, N8, 328-331 (2015).
- [31] U. Saleem, M. Hasan. *J. Phys. A: Math. Theor.*, **43**, 045204 (2010).
- [32] R. Myrzakulov, S. Vijayalakshmi, G. Nugmanova , M. Lakshmanan *Physics Letters A*, **233** , 14-6, 391-396 (1997).
- [33] R. Myrzakulov, S. Vijayalakshmi, R. Syzdykova, M. Lakshmanan, *J. Math. Phys.*, **39**, 2122-2139 (1998).
- [34] R. Myrzakulov, M. Lakshmanan, S. Vijayalakshmi, A. Danlybaeva , *J. Math. Phys.*, **39**, 3765-3771 (1998).
- [35] Myrzakulov R, Danlybaeva A.K, Nugmanova G.N. *Theoretical and Mathematical Physics*, V.118, 13, P. 441-451 (1999).

- [36] Myrzakulov R., Nugmanova G., Syzdykova R. Journal of Physics A: Mathematical & Theoretical, V.31, 147, P.9535-9545 (1998).
- [37] Myrzakulov R., Daniel M., Amuda R. Physica A., V.234, 13-4, P.715-724 (1997).
- [38] Myrzakulov R., Makhankov V.G., Pashaev O.?. Letters in Mathematical Physics, V.16, N1, P.83-92 (1989)
- [39] Myrzakulov R., Makhankov V.G., Makhankov A. Physica Scripta, V.35, N3, P. 233-237 (1987)
- [40] Myrzakulov R., Pashaev O.?. Kholmurodov Kh. Physica Scripta, V.33, N4, P. 378-384 (1986)
- [41] Anco S.C., Myrzakulov R. Journal of Geometry and Physics, v.60, 1576-1603 (2010)
- [42] Myrzakulov R., Rahimov F.K., Myrzakul K., Serikbaev N.S. *On the geometry of stationary Heisenberg ferromagnets*. In: "Non-linear waves: Classical and Quantum Aspects", Kluwer Academic Publishers, Dordrecht, Netherlands, P. 543-549 (2004)
- [43] Myrzakulov R., Serikbaev N.S., Myrzakul Kur., Rahimov F.K. *On continuous limits of some generalized compressible Heisenberg spin chains*. Journal of NATO Science Series II. Mathematics, Physics and Chemistry, V 153, P. 535-542 (2004)
- [44] R.Myrzakulov, G. K. Mamyrbekova, G. N. Nugmanova, M. Lakshmanan. Symmetry, **7**(3), 1352-1375 (2015). [arXiv:1305.0098]
- [45] R.Myrzakulov, G. K. Mamyrbekova, G. N. Nugmanova, K. Yesmakhanova, M. Lakshmanan. Physics Letters A, **378**, N30-31, 2118-2123 (2014). [arXiv:1404.2088]
- [46] Myrzakulov R., Martina L., Kozhamkulov T.A., Myrzakul Kur. *Integrable Heisenberg ferromagnets and soliton geometry of curves and surfaces*. In book: "Nonlinear Physics: Theory and Experiment. II". World Scientific, London, P. 248-253 (2003)
- [47] Myrzakulov R. *Integrability of the Gauss-Codazzi-Mainardi equation in 2+1 dimensions*. In "Mathematical Problems of Nonlinear Dynamics", Proc. of the Int. Conf. "Progress in Nonlinear sciences", Nizhny Novgorod, Russia, July 2-6, 2001, V.1, P.314-319 (2001)
- [48] Chen Chi, Zhou Zi-Xiang. *Darboux Transformation and Exact Solutions of the Myrzakulov-I Equations*. Chin. Phys. Lett., **26**, N8, 080504 (2009)
- [49] Chen Hai, Zhou Zi-Xiang. *Darboux Transformation with a Double Spectral Parameter for the Myrzakulov-I Equation*. Chin. Phys. Lett., **31**, N12, 120504 (2014)
- [50] Zhao-Wen Yan, Min-Ru Chen, Ke Wu, Wei-Zhong Zhao. J. Phys. Soc. Jpn., **81**, 094006 (2012)
- [51] Yan Zhao-Wen, Chen Min-Ru, Wu Ke, Zhao Wei-Zhong. Commun. Theor. Phys., **58**, 463-468 (2012)
- [52] K.R. Ysmakhanova, G.N. Nugmanova, Wei-Zhong Zhao, Ke Wu. *Integrable inhomogeneous Lakshmanan-Myrzakulov equation*, [nlin/0604034]
- [53] Zhen-Huan Zhang, Ming Deng, Wei-Zhong Zhao, Ke Wu. *On the integrable inhomogeneous Myrzakulov-I equation*, [arXiv: nlin/0603069]
- [54] Martina L, Myrzakul Kur., Myrzakulov R, Soliani G. Journal of Mathematical Physics, V.42, 13, P.1397-1417 (2001).
- [55] Xiao-Yu Wu, Bo Tian, Hui-Ling Zhen, Wen-Rong Sun and Ya Sun. Journal of Modern Optics, 2015.

- [56] Z.S. Yersultanova, M. Zhassybayeva, K. Yesmakhanova, G. Nugmanova, R. Myrzakulov. International Journal of Geometric Methods in Modern Physics, **13**, N1, 1550134 (2016).. [arXiv:1404.2270]
- [57] Myrzakul Akbota and Myrzakulov Ratbay. *Integrable Motion of Two Interacting Curves and Heisenberg Ferromagnetic Equations*, Abstracts of XVIII-th Intern. Conference "Geometry, Integrability and Quantization", June 3-8, 2016, Bulgaria.
- [58] Myrzakul Akbota and Myrzakulov Ratbay. *Integrable motion of two interacting curves, spin systems and the Manakov system*. International Journal of Geometric Methods in Modern Physics, **13**, N1, 1550134 (2016). [arXiv:1606.06598]
- [59] Myrzakul Akbota and Myrzakulov Ratbay. *Darboux transformations and exact soliton solutions of integrable coupled spin systems related with the Manakov system*, [arXiv:1607.08151]
- [60] Myrzakul Akbota and Myrzakulov Ratbay. *Integrable geometric flows of interacting curves/surfaces, multilayer spin systems and the vector nonlinear Schrodinger equation*. International Journal of Geometric Methods in Modern Physics, **13**, N1, 1550134 (2016). [arXiv:1608.08553]
- [61] Myrzakulova Zh., Myrzakul A., Nugmanova G., Myrzakulov R. *Notes on Integrable Motion of Two Interacting Curves and Two-layer Generalized Heisenberg Ferromagnet Equations*, [arXiv:1811:12216]
- [62] Hussien R.A., Mohamed S.G. *Generated Surfaces via Inextensible Flows of Curves in  $R^3$* . Journal of Applied Mathematics, v.2016, Article ID 6178961 (2016).
- [63] Bekova G., Yesmakhanova K., Shaikhova G., Nugmanova G., Myrzakulov R. *Geometric formulation and soliton solutions of the Myrzakulov-LXXIII equation and the complex short pulse equation*,
- [64] C. Qu, J. Song and R. Yao. *Multi-Component Integrable Systems and Invariant Curve Flows in Certain Geometries*, Symmetry, Integrability and Geometry: Methods and Applications SIGMA 9 (2013), 001, 19 pages [arXiv:1301.0180]
- [65] Mussatayeva A., Myrzakul T., Nugmanova G., Yesmakhanova K., Myrzakulov R. *Integrable Motion of Curves, Spin Equation and Camassa-Holm Equation*, [arXiv:1907.10910]
- [66] Kutum B. *Gauge equivalence between the M-CV equation and the two-component Camassa-Holm equation*
- [67] Taishiyeva A., Myrzakul T., Nugmanova G., Yesmakhanova K. *On the Gauge Equivalence between the 2-component Camassa-Holm and Myrzakulov-CVI equations*
- [68] Mussatayeva A., Myrzakul T., Nugmanova G., Yesmakhanova K. *Gauge equivalence between the Myrzakulov-CIV equation and the Camassa-Holm Equation*
- [69] Taishiyeva A., Myrzakul T., Nugmanova G., Yesmakhanova K., Myrzakulov R. *Geometric Flows of Curves, Two-Component Camassa-Holm Equation and Generalized Heisenberg Ferromagnet Equation*
- [70] Kutum B., Nugmanova G., Myrzakul T., Yesmakhanova K., Myrzakulov R. *Integrable Deformation of Space Curves, Generalized Heisenberg Ferromagnet Equation and Two-Component Modified Camassa-Holm Equation*, [arXiv:1908.01371]