

# Interaction phenomena between lump and solitary wave of a generalized $(3 + 1)$ -dimensional variable-coefficient nonlinear-wave equation in liquid with gas bubbles

Jian-Guo Liu\* · Wen-Hui Zhu\* · Yan He\* · Ya-Kui Wu\*

Received: date / Accepted: date

**Abstract** In this paper, a generalized  $(3 + 1)$ -dimensional variable-coefficient nonlinear-wave equation is studied in liquid with gas bubbles. Based on the Hirota's bilinear form and symbolic computation, lump and interaction solutions between lump and solitary wave are obtained. Their interaction phenomena is shown in some 3d graphs and contour plots, which include a periodic-shape lump solution, a parabolic-shape lump solution, a cubic-shape lump solution, interaction solutions between lump and one solitary wave, and between lump and two solitary waves. The spatial structures called the bright lump wave and the bright-dark lump wave are discussed. Interaction behaviors of two bright-dark lump waves and a periodic-shape bright lump wave are also presented.

**Keywords** Solitary wave, lump wave, variable-coefficient nonlinear-wave equation, interaction behaviors

**2010 Mathematics Subject Classification:** 35Q55, 35C08, 78A60

---

Project supported by National Natural Science Foundation of China (Grant No 81960715) and Science and Technology Project of Education Department of Jiangxi Province(GJJ151079).

---

Jian-Guo Liu(\*Corresponding author), Yan He(\*Corresponding author)  
College of Computer, Jiangxi University of Traditional Chinese Medicine, Jiangxi 330004, China  
Tel.: +8613970042436  
E-mail: 20101059@jxutcm.edu.cn(J.G. Liu)  
E-mail: 274667818@qq.com(Y. He)

Wen-Hui Zhu(\*Corresponding author)  
Institute of artificial intelligence, Nanchang Institute of Science and Technology, Jiangxi 330108, China  
E-mail: 415422402@qq.com

Ya-Kui Wu(\*Corresponding author)  
School of science, Jiujiang University, Jiangxi 332005, China  
E-mail: 2180501008@cnu.edu.cn

## 1 Introduction

In some branches of science and engineering such as fluid mechanics, quantum mechanics, particle physics, mass transfer, plasma physics, nanoliquids and biological mathematics [1-5], Nonlinear partial differential equations (NPDES) are used to describe many nonlinear phenomena and wave propagation characteristics. As the lump solutions of the NPDES are the special, powerful destructive ocean wave in the real world, it is important to search for the lump solutions of the NPDES, especially the constant coefficient NPDES have attracted the attention of many scholars [6-14].

Recently, a generalized  $(3 + 1)$ -dimensional nonlinear-wave equation has been presented as [15]

$$[4u_t + 4uu_x + u_{xxx} - 4u_x]_x + 3(u_{yy} + u_{zz}) = 0, \quad (1)$$

which describes a liquid with gas bubbles in the three dimensional case.

However, the variable-coefficient NPDES provide us more real phenomena in the inhomogeneities of media and non-uniformities of boundaries than corresponding constant-coefficient counterparts in some physical cases. In this paper, a generalized  $(3 + 1)$ -dimensional variable-coefficient nonlinear-wave equation is investigated [16,17]

$$\alpha(t)u_x^2 + \alpha(t)uu_{xx} + \beta(t)u_{xxxx} + \gamma(t)u_{xx} + \delta(t)u_{yy} + \varrho(t)u_{zz} + u_{xt} = 0, \quad (2)$$

where  $u = u(x, y, z, t)$  is the wave-amplitude function. The bilinear form, Bäcklund transformation, Lax pair, infinitely-many conservation laws, multi-soliton solutions, travelling-wave solutions and one-periodic wave solutions are presented by virtue of the binary Bell polynomials, the Hirota method, the polynomial expansion method and the Hirota-Riemann method [18]. However, lump and interaction solutions between lump and solitary wave of Eq. (2) have not been obtained yet, which will make the main work of our paper.

This paper will be organised as follows. Section 2 obtains the lump solutions of Eq. (2) with the aid of the Hirota's bilinear form [19-24] and Mathematical software [25-35] and demonstrates their physical structures by some 3d graphs and contour plots; Section 3 presents the interaction solutions between lump and one solitary wave; Section 4 derives the interaction solutions between lump and two solitary waves; Section 5 gives the conclusion.

## 2 Lump solutions of Eq. (2)

Setting  $u = 12 [\ln \xi(x, y, z, t)]_{xx}$  and  $\alpha(t) = \beta(t)$ , the bilinear form of Eq. (2) can be introduced as (see Ref. [18]) by using the multi-dimensional Bell polynomials

$$[D_x D_t + \beta(t)D_x^4 + \gamma(t)D_x^2 + \delta(t)D_y^2 + \varrho(t)D_z^2] \xi \cdot \xi = 0. \quad (3)$$

This is equivalent to

$$\begin{aligned} & \xi[\beta(t)\xi_{xxxx} + \gamma(t)\xi_{xx} + \delta(t)\xi_{yy} + \varrho(t)\xi_{zz} + \xi_{xt}] + 3\beta(t)\xi_{xx}^2 \\ & - 4\beta(t)\xi_x\xi_{xxx} - \gamma(t)\xi_x^2 - \delta(t)\xi_y^2 - \varrho(t)\xi_z^2 - \xi_t\xi_x = 0. \end{aligned} \quad (4)$$

In order to seek the lump solutions of Eq. (2), we suppose

$$\begin{aligned} \zeta &= x\alpha_1 + y\alpha_2 + z\alpha_3 + \alpha_4(t), \\ \varsigma &= x\alpha_5 + y\alpha_6 + z\alpha_7 + \alpha_8(t), \\ \xi &= \zeta^2 + \varsigma^2 + \alpha_9(t), \end{aligned} \quad (5)$$

where  $\alpha_1, \alpha_2, \alpha_3, \alpha_5, \alpha_6$  and  $\alpha_7$  are unknown constants.  $\alpha_4(t), \alpha_8(t), \alpha_9(t)$  are undefined real functions. Substituting Eq. (5) into Eq. (4) through Mathematical symbolic computations, we get

$$\begin{aligned} (I) : \alpha_8(t) &= \eta_1 + \int_1^t -\frac{(\alpha_1^2 + \alpha_5^2) [\alpha_1^2\gamma(t) + \alpha_2^2\delta(t) + \alpha_3^2\varrho(t)] + \alpha_1^3\alpha_4'(t)}{\alpha_1^2\alpha_5} dt, \\ \alpha_9(t) &= \eta_2 + \int_1^t [2[\alpha_1(\alpha_1\gamma(t) + \alpha_4'(t)) + \alpha_2^2\delta(t) + \alpha_3^2\varrho(t)][\alpha_1[\eta_3 \\ &+ \int_1^t -\frac{(\alpha_1^2 + \alpha_5^2) [\alpha_1^2\gamma(t) + \alpha_2^2\delta(t) + \alpha_3^2\varrho(t)] + \alpha_1^3\alpha_4'(t)}{\alpha_1^2\alpha_5} dt] - \alpha_5 \\ &* \alpha_4(t)]/(\alpha_1\alpha_5) dt, \alpha(t) = \beta(t) = 0, \alpha_6 = \frac{\alpha_2\alpha_5}{\alpha_1}, \alpha_7 = \frac{\alpha_3\alpha_5}{\alpha_1}, \end{aligned} \quad (6)$$


with  $\alpha_1 \neq 0, \alpha_5 \neq 0$ . Substituting Eq. (5) and Eq. (6) into the transformation  $u = 12 [\ln \xi(x, y, z, t)]_{xx}$ , we have the following lump solution of Eq. (2)

$$\begin{aligned} u^{(I)} &= [12[2(\alpha_1^2 + \alpha_5^2) [\eta_2 + \int_1^t [2[\alpha_1[\alpha_1\gamma(t) + \alpha_4'(t)] + \alpha_2^2\delta(t) + \alpha_3^2\varrho(t)][\alpha_1[\eta_1 \\ &+ \int_1^t -\frac{(\alpha_1^2 + \alpha_5^2) [\alpha_1^2\gamma(t) + \alpha_2^2\delta(t) + \alpha_3^2\varrho(t)] + \alpha_1^3\alpha_4'(t)}{\alpha_1^2\alpha_5} dt] - \alpha_5\alpha_4(t)]] \\ &/ (\alpha_1\alpha_5) dt + [\eta_1 - \int_1^t \frac{(\alpha_1^2 + \alpha_5^2) [\alpha_1^2\gamma(t) + \alpha_2^2\delta(t) + \alpha_3^2\varrho(t)] + \alpha_1^3\alpha_4'(t)}{\alpha_1^2\alpha_5} dt \\ &+ \frac{\alpha_5(\alpha_1x + \alpha_2y + \alpha_3z)}{\alpha_1}]^2 + [\alpha_4(t) + \alpha_1x + \alpha_2y + \alpha_3z]^2] - 4[\alpha_5[\eta_1 \\ &+ \int_1^t -\frac{(\alpha_1^2 + \alpha_5^2) [\alpha_1^2\gamma(t) + \alpha_2^2\delta(t) + \alpha_3^2\varrho(t)] + \alpha_1^3\alpha_4'(t)}{\alpha_1^2\alpha_5} dt + \alpha_5x] \\ &+ \alpha_1(\alpha_4(t) + \alpha_2y + \alpha_3z) + \alpha_1^2x + \frac{\alpha_5^2(\alpha_2y + \alpha_3z)}{\alpha_1}]^2]/[\eta_2 \\ &+ \int_1^t [2[\alpha_1(\alpha_1\gamma(t) + \alpha_4'(t)) + \alpha_2^2\delta(t) + \alpha_3^2\varrho(t)][\alpha_1[\eta_1 \\ &- \int_1^t \frac{(\alpha_1^2 + \alpha_5^2) [\alpha_1^2\gamma(t) + \alpha_2^2\delta(t) + \alpha_3^2\varrho(t)] + \alpha_1^3\alpha_4'(t)}{\alpha_1^2\alpha_5} dt] - \alpha_5\alpha_4(t)]] \end{aligned}$$

$$\begin{aligned}
& / (\alpha_1 \alpha_5) dt + [\eta_1 - \int_1^t \frac{(\alpha_1^2 + \alpha_5^2) [\alpha_1^2 \gamma(t) + \alpha_2^2 \delta(t) + \alpha_3^2 \varrho(t)] + \alpha_1^3 \alpha_4'(t)}{\alpha_1^2 \alpha_5} dt \\
& + \frac{\alpha_5 (\alpha_1 x + \alpha_2 y + \alpha_3 z)}{\alpha_1}]^2 + (\alpha_4(t) + \alpha_1 x + \alpha_2 y + \alpha_3 z)^2] dt, \quad (7)
\end{aligned}$$

where  $\alpha_4(t)$  is arbitrary function,  $\eta_1$  and  $\eta_2$  are integral constants.

The physical structures for  $(u^{(I)})$  are described in Fig. 1 by the 3d graphs and contour plots. Fig. 1 shows the propagation of solution  $(u^{(I)})$  when  $\gamma(t)$ ,  $\delta(t)$ ,  $\varrho(t)$  and  $\alpha_4(t)$  select different functions. When  $\gamma(t) = \delta(t) = \varrho(t) = \cos t$  and  $\alpha_4(t) = \sin t$ , a periodic-shape rational solution is listed in Fig. 1(a) and Fig. 1(d). When  $\gamma(t) = \delta(t) = \varrho(t) = t$  and  $\alpha_4(t) = \sin t$ , a parabolic-shape rational solution is presented in Fig. 1(b) and Fig. 1(e). When  $\gamma(t) = \cosh t$ ,  $\varrho(t) = \exp t$  and  $\alpha_4(t) = \delta(t) = t$ , a cubic-shape rational solution is shown in Fig. 1(c) and Fig. 1(f).




**Fig. 1.** Lump solution  $u^{(I)}$  with  $\alpha_1 = 1$ ,  $\alpha_2 = 2$ ,  $\alpha_3 = -1$ ,  $\alpha_5 = -3$ ,  $x = -5$ ,  $\eta_1 = \eta_2 = y = 0$ , when  $\gamma(t) = \delta(t) = \varrho(t) = \cos t$ ,  $\alpha_4(t) = \sin t$  in (a) (d),  $\gamma(t) = \delta(t) = \varrho(t) = t$ ,  $\alpha_4(t) = \sin t$  in (b) (e) and  $\gamma(t) = \cosh t$ ,  $\varrho(t) = \exp t$  and  $\alpha_4(t) = \delta(t) = t$  in (c) (f).

$$\begin{aligned}
(II) : \alpha_8(t) &= \eta_3 + \int_1^t \alpha_5 \left[ -\frac{3(\alpha_1^2 + \alpha_5^2) \beta(t)}{\alpha_9} - \frac{\alpha_3^2 \varrho(t)}{\alpha_1^2} - \gamma(t) \right] dt, \\
\alpha_9(t) &= \alpha_9, \alpha_6 = -\frac{\alpha_1 \alpha_2}{\alpha_5}, \alpha_7 = \frac{\alpha_3 \alpha_5}{\alpha_1}, \delta(t) = -\frac{3\alpha_5^2 (\alpha_1^2 + \alpha_5^2) \beta(t)}{\alpha_2^2 \alpha_9}, \\
\alpha_4(t) &= \eta_4 + \int_1^t \left[ -\frac{\alpha_1 [3(\alpha_1^2 + \alpha_5^2) \beta(t) + \alpha_9 \gamma(t)]}{\alpha_9} - \frac{\alpha_3^2 \varrho(t)}{\alpha_1} \right] dt, \quad (8)
\end{aligned}$$


with  $\alpha_1 \neq 0$ ,  $\alpha_5 \neq 0$ ,  $\alpha_9 \neq 0$  and  $\alpha_2\alpha_9 \neq 0$ . Substituting Eq. (5) and Eq. (8) into the transformation  $u = 12 [\ln \xi(x, y, z, t)]_{xx}$ , we have the following lump solution of Eq. (2)

$$\begin{aligned}
u^{(II)} = & [12[2(\alpha_1^2 + \alpha_5^2) [\alpha_9 + [\int_1^t \alpha_5 [-\frac{3(\alpha_1^2 + \alpha_5^2)\beta(t)}{\alpha_9} - \frac{\alpha_3^2 \varrho(t)}{\alpha_1^2} - \gamma(t)] dt \\
& + \eta_3 + \alpha_5 x - \frac{\alpha_1 \alpha_2 y}{\alpha_5} + \frac{\alpha_3 \alpha_5 z}{\alpha_1}]^2 + [\int_1^t [-\frac{\alpha_1 [3(\alpha_1^2 + \alpha_5^2)\beta(t) + \alpha_9 \gamma(t)]}{\alpha_9} \\
& - \frac{\alpha_3^2 \varrho(t)}{\alpha_1}]] dt + \eta_4 + \alpha_1 x + \alpha_2 y + \alpha_3 z]^2] \\
& - [4[\alpha_1^2 [\int_1^t [-\frac{\alpha_1 [3(\alpha_1^2 + \alpha_5^2)\beta(t) + \alpha_9 \gamma(t)]}{\alpha_9} - \frac{\alpha_3^2 \varrho(t)}{\alpha_1}]] dt] + \alpha_5 \alpha_1 [\eta_3 \\
& + \int_1^t \alpha_5 [-\frac{3(\alpha_1^2 + \alpha_5^2)\beta(t)}{\alpha_9} - \frac{\alpha_3^2 \varrho(t)}{\alpha_1^2} - \gamma(t)] dt + \alpha_5 x] + \alpha_1^3 x \\
& + \alpha_1^2 (\eta_4 + \alpha_3 z) + \alpha_3 \alpha_5^2 z^2] / (\alpha_1^2)] / [\alpha_9 + [\eta_3 + \alpha_5 x - \frac{\alpha_1 \alpha_2 y}{\alpha_5} + \frac{\alpha_3 \alpha_5 z}{\alpha_1} \\
& + \int_1^t \alpha_5 [-\frac{3(\alpha_1^2 + \alpha_5^2)\beta(t)}{\alpha_9} - \frac{\alpha_3^2 \varrho(t)}{\alpha_1^2} - \gamma(t)] dt]^2 + [\eta_4 + \alpha_1 x + \alpha_2 y \\
& + \int_1^t [-\frac{\alpha_1 [3(\alpha_1^2 + \alpha_5^2)\beta(t) + \alpha_9 \gamma(t)]}{\alpha_9} - \frac{\alpha_3^2 \varrho(t)}{\alpha_1}]] dt + \alpha_3 z]^2]^2], \quad (9)
\end{aligned}$$


where  $\eta_3$  and  $\eta_4$  are integral constants.




**Fig. 2.** Lump solution (9) with  $\alpha_1 = 1$ ,  $\alpha_2 = 2$ ,  $\alpha_3 = -1$ ,  $\alpha_5 = \alpha_9 = -3$ ,  $\eta_3 = \eta_4 = 0$ ,  $z = -10$ , when  $t = -1$  in (a) (d),  $t = 0$  in (b) (e) and  $t = 1$  in (c) (f).



**Fig. 3.** Lump solution (9) with  $\alpha_1 = -1$ ,  $\alpha_2 = 2$ ,  $\alpha_3 = -1$ ,  $\alpha_5 = 3$ ,  $\alpha_9 = -3$ ,  $\eta_3 = \eta_4 = z = 0$ , when  $x = -30$  in (a) (d),  $x = 0$  in (b) (e) and  $x = 30$  in (c) (f).



**Fig. 4.** Lump solution (9) with  $\alpha_1 = -1$ ,  $\alpha_2 = 2$ ,  $\alpha_3 = -1$ ,  $\alpha_5 = 3$ ,  $\alpha_9 = -3$ ,  $\eta_3 = \eta_4 = z = 0$ , when  $x = -8$  in (a) (d),  $x = 0$  in (b) (e) and  $x = 8$  in (c) (f).



**Fig. 5.** Lump solution (9) with  $\alpha_1 = -1$ ,  $\alpha_2 = 2$ ,  $\alpha_3 = -1$ ,  $\alpha_5 = 3$ ,  $\alpha_9 = -3$ ,  $\eta_3 = \eta_4 = z = 0$ , when  $x = -3$  in (a) (d),  $x = 0$  in (b) (e) and  $x = 3$  in (c) (f).

Then, the physical structures for  $(u^{(II)})$  are shown in Figs. 2-5 with some 3d graphs and contour plots. When  $\gamma(t) = -1$ ,  $\varrho(t) = \beta(t) = 1$ , the spatial structure called the bright lump wave is seen in Fig. 2 at  $t = -1; 0; 1$ , the spatial structure called the bright-dark lump wave is shown in Fig. 3 at  $x = -30; 0; 30$ . When  $\gamma(t) = -t$ ,  $\varrho(t) = \beta(t) = t$ , interaction behaviors of two bright-dark lump waves are presented in Fig. 4 at  $x = -8; 0; 8$ . As the value of  $x$  changes, the two bright-dark lump waves move towards each other, and finally merge together. When  $\gamma(t) = \varrho(t) = \beta(t) = \cos t$ , a periodic-shape bright lump wave is found in Fig. 5 at  $x = -3; 0; 3$ .

$$(III) : \varrho(t) = -\frac{3(\alpha_1^2 + \alpha_5^2)^3 \beta(t) + (\alpha_2 \alpha_5 - \alpha_1 \alpha_6)^2 \alpha_9 \delta(t)}{(\alpha_3 \alpha_5 - \alpha_1 \alpha_7)^2 \alpha_9}, \alpha_9(t) = \alpha_9,$$


$$\alpha_4(t) = \eta_5 - \int_1^t [\alpha_1 [\alpha_5^2 \gamma(t) + (\alpha_2^2 - \alpha_6^2) \delta(t) + (\alpha_3^2 - \alpha_7^2) \varrho(t)]$$

$$+ \alpha_1^3 \gamma(t) + 2\alpha_5 [\alpha_2 \alpha_6 \delta(t) + \alpha_3 \alpha_7 \varrho(t)]] / (\alpha_1^2 + \alpha_5^2) dt,$$


$$\alpha_8(t) = \eta_6 - \int_1^t [\alpha_5 [\alpha_5^2 \gamma(t) + (\alpha_6^2 - \alpha_2^2) \delta(t) + (\alpha_7^2 - \alpha_3^2) \varrho(t)]$$

$$+ \alpha_5 \alpha_1^2 \gamma(t) + 2\alpha_1 [\alpha_2 \alpha_6 \delta(t) + \alpha_3 \alpha_7 \varrho(t)]] / (\alpha_1^2 + \alpha_5^2) dt \quad (10)$$

with  $(\alpha_3 \alpha_5 - \alpha_1 \alpha_7)^2 \alpha_9 \neq 0$ ,  $\alpha_1^2 + \alpha_5^2 \neq 0$ ,  $\eta_5$  and  $\eta_6$  are integral constants. Substituting Eq. (5) and Eq. (10) into the transformation  $u = 12 [\ln \xi(x, y, z, t)]_{xx}$ , we derive another lump solution of Eq. (2)



**Fig. 6.** Lump solution (11) with  $\alpha_1 = \alpha_2 = \alpha_3 = \alpha_5 = -1$ ,  $\alpha_6 = \alpha_7 = \eta_5 = 3$ ,  $\eta_6 = -2$ ,  $\alpha_9 = 2$ ,  $z = 0$ , when  $x = -10$  in (a) (d),  $x = 0$  in (b) (e) and  $x = 10$  in (c) (f).




**Fig. 7.** Lump solution (11) with  $\alpha_1 = \alpha_3 = \alpha_5 = -1$ ,  $\alpha_2 = 1$ ,  $\alpha_6 = -3$ ,  $\alpha_7 = 3$ ,  $\eta_6 = \eta_5 = z = 0$ ,  $\alpha_9 = 2$ , when  $y = -3$  in (a) (d),  $y = 0$  in (b) (e) and  $y = 3$  in (c) (f).



$$\begin{aligned}
u^{(III)} = & [12[2(\alpha_1^2 + \alpha_5^2)[\alpha_9 + [\alpha_4(t) + \alpha_1x + \alpha_2y + \alpha_3z]^2 + [\alpha_8(t) + \alpha_5x \\
& + \alpha_6y + \alpha_7z]^2] - [2\alpha_1[\alpha_4(t) + \alpha_1x + \alpha_2y + \alpha_3z] + 2\alpha_5[\alpha_8(t) + \alpha_5x \\
& + \alpha_6y + \alpha_7z]^2]] / [[\alpha_9 + [\alpha_4(t) + \alpha_1x + \alpha_2y + \alpha_3z]^2 \\
& + [\alpha_8(t) + \alpha_5x + \alpha_6y + \alpha_7z]^2]^2], \tag{11}
\end{aligned}$$

where  $\alpha_4(t)$ ,  $\alpha_4(t)$  and  $\varrho(t)$  satisfy constraint (10).

The physical structures for  $(u^{(III)})$  are shown in Figs. 6-8 with some 3d graphs and contour plots. When  $\gamma(t) = -1, \delta(t) = \beta(t) = 1$ , the spatial structure called the bright lump wave is seen in Fig. 6 at  $x = -10; 0; 10$ . When  $\gamma(t) = -t, \delta(t) = \beta(t) = t$ , interaction behaviors of two bright lump waves are presented in Fig. 7 at  $y = -3; 0; 3$ . As the value of  $y$  changes, the two bright lump waves move towards each other, and finally merge together. When  $\gamma(t) = \delta(t) = \beta(t) = \cos t$ , a periodic-shape bright lump wave is found in Fig. 8 at  $z = -10; 0; 10$  and  $x = 0$ .



**Fig. 8.** Lump solution (11) with  $\alpha_1 = \alpha_2 = \alpha_3 = \alpha_5 = -1$ ,  $\alpha_6 = \alpha_7 = \eta_5 = 3$ ,  $\eta_6 = -2$ ,  $\alpha_9 = 2$ ,  $x = 0$ , when  $z = -10$  in (a) (d),  $z = 0$  in (b) (e) and  $z = 10$  in (c) (f).

### 3 Interaction solutions between lump and one solitary wave

In order to find the interaction solutions between lump and one solitary wave, we add an exponential function in Eq. (5) as follows


$$\zeta = x\alpha_1 + y\alpha_2 + z\alpha_3 + \alpha_4(t),$$

$$\begin{aligned}\zeta &= x\alpha_5 + y\alpha_6 + z\alpha_7 + \alpha_8(t), \\ \xi &= \zeta^2 + \zeta^2 + \alpha_9(t) + \alpha_{14}(t)e^{\alpha_{13}(t) + \alpha_{10}x + \alpha_{11}y + \alpha_{12}z},\end{aligned}\quad (12)$$

where  $\alpha_{10}$ ,  $\alpha_{11}$  and  $\alpha_{12}$  are unknown constants.  $\alpha_{13}(t)$  and  $\alpha_{14}(t)$  are unknown real functions. Substituting Eq. (12) into Eq. (4) through Mathematical software, we have

$$\begin{aligned}\alpha_6 &= -\frac{\alpha_1\alpha_2}{\alpha_5}, \alpha_7 = -\frac{\alpha_1\alpha_3}{\alpha_5}, \alpha_{11} = \alpha_{12} = 0, \alpha_9(t) = \frac{\alpha_1^2 + \alpha_5^2}{\alpha_{10}^2}, \\ \delta(t) &= -\frac{3\alpha_5^2\alpha_{10}^2\beta(t) + \alpha_3^2\varrho(t)}{\alpha_2^2}, \alpha_8(t) = \eta_8 - \alpha_5 \int [3\alpha_{10}^2\beta(t) + \gamma(t)] dt, \\ \alpha_4(t) &= \eta_9 - \alpha_1 \int [3\alpha_{10}^2\beta(t) + \gamma(t)] dt, \\ \alpha_{13}(t) &= \eta_{10} - \alpha_{10} \int [\alpha_{10}^2\beta(t) + \gamma(t)] dt - \ln \alpha_{14}(t),\end{aligned}\quad (13)$$

with  $\alpha_2 \neq 0$ ,  $\alpha_5 \neq 0$  and  $\alpha_{10} \neq 0$ . Substituting Eq. (5) and Eq. (13) into the transformation  $u = 12 [\ln \xi(x, y, z, t)]_{xx}$ , we get




**Fig. 9.** Interaction solution (14) with  $\alpha_1 = \eta_8 = \eta_9 = \eta_{10} = \beta(t) = 1$ ,  $\alpha_3 = \gamma(t) = -1$ ,  $\alpha_2 = \alpha_{10} = 2$ ,  $\alpha_5 = -3$ ,  $z = 0$ , when  $t = -1$  in (a) (d),  $t = 0$  in (b) (e) and  $t = 1$  in (c) (f).

$$\begin{aligned}u^{(IV)} &= [12[[2(\alpha_1^2 + \alpha_5^2) + \alpha_{10}^2 \exp[\eta_{10} + \alpha_{10}(x - \int (\alpha_{10}^2\beta(t) + \gamma(t)) dt]]] \\ &\quad * [\frac{\alpha_1^2 + \alpha_5^2}{\alpha_{10}^2} + \exp[\eta_{10} + \alpha_{10} \left( x - \int (\alpha_{10}^2\beta(t) + \gamma(t)) dt \right)]]\end{aligned}$$

$$\begin{aligned}
& + \frac{[\alpha_1(\alpha_2 y + \alpha_3 z) - \alpha_5[\eta_8 + \alpha_5(x - \int (3\alpha_{10}^2 \beta(t) + \gamma(t)) dt)]]^2}{\alpha_5^2} \\
& + \left( \eta_9 + \alpha_1 \left( x - \int (3\alpha_{10}^2 \beta(t) + \gamma(t)) dt \right) + \alpha_2 y + \alpha_3 z \right)^2 \\
& - [2\alpha_1 \eta_9 + \alpha_{10} \exp \left( \eta_{10} + \alpha_{10} \left( x - \int (\alpha_{10}^2 \beta(t) + \gamma(t)) dt \right) \right) \\
& + 2\alpha_5 [\eta_8 + \alpha_5 \left( x - \int (3\alpha_{10}^2 \beta(t) + \gamma(t)) dt \right) \\
& + 2\alpha_1^2 [x - \int (3\alpha_{10}^2 \beta(t) + \gamma(t)) dt]^2] / \left[ \frac{\alpha_1^2 + \alpha_5^2}{\alpha_{10}^2} \right. \\
& + \exp \left( \eta_{10} + \alpha_{10} \left( x - \int (\alpha_{10}^2 \beta(t) + \gamma(t)) dt \right) \right) + [\alpha_1(\alpha_2 y + \alpha_3 z) \\
& - \alpha_5 \left( \eta_8 + \alpha_5 \left( x - \int (3\alpha_{10}^2 \beta(t) + \gamma(t)) dt \right) \right)]^2 / \alpha_5^2 \\
& \left. + [\eta_9 + \alpha_1 [x - \int (3\alpha_{10}^2 \beta(t) + \gamma(t)) dt] + \alpha_2 y + \alpha_3 z]^2 \right]^2, \quad (14)
\end{aligned}$$

where  $\eta_8$ ,  $\eta_9$  and  $\eta_{10}$  are integral constants. Interaction phenomena between lump and one solitary wave in Eq. (14) is shown in Fig. 9 and Fig. 10. Obviously, it can be seen a solitary wave and a lump wave in Fig. 9(a) and Fig. 9(d). In Fig. 9(b) and Fig. 9(e), the solitary and lump wave are slowly approaching at  $t = 0$ . In Fig. 9(c) and Fig. 9(f), the solitary and lump wave merge together to propagate forward at  $t = 1$ . Fig. 10 displays the effect of variable coefficient  $\gamma(t)$  on the interaction phenomena between lump and one solitary wave.



**Fig. 10.** Interaction solution (14) with  $\alpha_1 = \eta_8 = \eta_9 = \eta_{10} = \alpha_{10} = \beta(t) = 1$ ,

$\alpha_3 = \alpha_5 = -1$ ,  $\alpha_2 = 2$ ,  $y = z = 0$ , when  $\gamma(t) = 1$  in (a) (d),  $\gamma(t) = t$  in (b) (e) and  $\gamma(t) = \cos t$  in (c) (f).

#### 4 Interaction solutions between lump and two solitary waves


In order to derive the interaction solutions between lump and two solitary waves, we add two exponential functions in Eq. (5) as follows

$$\begin{aligned}\zeta &= x\alpha_1 + y\alpha_2 + z\alpha_3 + \alpha_4(t), \\ \varsigma &= x\alpha_5 + y\alpha_6 + z\alpha_7 + \alpha_8(t), \\ \xi &= \zeta^2 + \varsigma^2 + \alpha_9(t) + \alpha_{14}(t)e^{\alpha_{13}(t) + \alpha_{10}x + \alpha_{11}y + \alpha_{12}z} \\ &\quad + \alpha_{15}(t)e^{-\alpha_{13}(t) - \alpha_{10}x - \alpha_{11}y - \alpha_{12}z},\end{aligned}\quad (15)$$

where  $\alpha_{15}(t)$  are unknown real functions. Substituting Eq. (15) into Eq. (4) through Mathematical software, we obtain

$$\begin{aligned}\alpha_6 &= -\frac{\alpha_1\alpha_2}{\alpha_5}, \alpha_7 = -\frac{\alpha_1\alpha_3}{\alpha_5}, \alpha_9(t) = \frac{\alpha_{10}^4\eta_{12} + \alpha_1^4 + 2\alpha_5^2\alpha_1^2 + \alpha_5^4}{(\alpha_1^2 + \alpha_5^2)\alpha_{10}^2}, \\ \varrho(t) &= \frac{\alpha_2^2(-\delta(t)) - 3\alpha_5^2\alpha_{10}^2\beta(t)}{\alpha_3^2}, \alpha_8(t) = \eta_{13} - \alpha_5 \int [3\alpha_{10}^2\beta(t) + \gamma(t)] dt, \\ \alpha_4(t) &= \eta_{14} - \alpha_1 \int [3\alpha_{10}^2\beta(t) + \gamma(t)] dt, \alpha_{11} = \alpha_{12} = 0, \\ \alpha_{13}(t) &= \eta_{15} - \alpha_{10} \int [\alpha_{10}^2\beta(t) + \gamma(t)] dt - \ln \alpha_{14}(t), \alpha_{15}(t) = \frac{\eta_{12}}{\alpha_{14}(t)},\end{aligned}\quad (16)$$


with  $\alpha_3 \neq 0$ ,  $\alpha_5 \neq 0$ ,  $\alpha_{14}(t) \neq 0$ ,  $\alpha_1^2 + \alpha_5^2 \neq 0$  and  $\alpha_{10} \neq 0$ . Substituting Eq. (5) and Eq. (16) into the transformation  $u = 12 [\ln \xi(x, y, z, t)]_{xx}$ , we get



**Fig. 11.** Interaction solution (16) with  $\alpha_3 = \gamma(t) = -1$ ,  $\alpha_2 = \alpha_{10} = 2$ ,  $\alpha_5 = -3$ ,  $\alpha_1 = \eta_{12} = \eta_{13} = \eta_{14} = \eta_{15} = \beta(t) = 1$ ,  $z = 0$ , when  $t = -1$  in (a),  $t = -0.3$  in (b),  $t = 0$  in (c),  $t = 0.3$  in (d),  $t = 1$  in (e).

$$\begin{aligned}
u^{(V)} = & 12[[2\alpha_1^2 + 2\alpha_5^2 + \alpha_{10}^2\alpha_{14}(t)e^{\alpha_{13}(t)+\alpha_{10}x} + \alpha_{10}^2\alpha_{15}(t)e^{-\alpha_{10}x-\alpha_{13}(t)}] \\
& / [\alpha_9(t) + \alpha_{14}(t)e^{\alpha_{13}(t)+\alpha_{10}x} + \alpha_{15}(t)e^{-\alpha_{10}x-\alpha_{13}(t)} + [\alpha_4(t) + \alpha_1x \\
& + \alpha_2y + \alpha_3z]^2 + (\alpha_8(t) + \alpha_5x + \alpha_6y + \alpha_7z)^2] \\
& - [[\alpha_{10}\alpha_{14}(t)e^{\alpha_{13}(t)+\alpha_{10}x} - \alpha_{10}\alpha_{15}(t)e^{-\alpha_{10}x-\alpha_{13}(t)} + 2\alpha_1[\alpha_4(t) \\
& + \alpha_1x + \alpha_2y + \alpha_3z] + 2\alpha_5(\alpha_8(t) + \alpha_5x + \alpha_6y + \alpha_7z)]^2 / [[\alpha_9(t) \\
& + \alpha_{14}(t)e^{\alpha_{13}(t)+\alpha_{10}x} + \alpha_{15}(t)e^{-\alpha_{10}x-\alpha_{13}(t)} + [\alpha_4(t) + \alpha_1x \\
& + \alpha_2y + \alpha_3z]^2 + [\alpha_8(t) + \alpha_5x + \alpha_6y + \alpha_7z]^2]^2]], \tag{17}
\end{aligned}$$

where  $\eta_{12}$ ,  $\eta_{13}$ ,  $\eta_{14}$  and  $\eta_{15}$  are integral constants. Interaction phenomena between lump and two solitary waves in Eq. (16) is shown in Fig. 11 and Fig. 12. Two solitary waves can be found in Fig. 11(a). A lump wave appears in one of two solitary waves in Fig. 11(b). In Fig. 11(c) and Fig. 11(d), the lump wave slowly shifts to another solitary wave, until vanishes in Fig. 11(e). Fig. 12 displays the corresponding contour plots of Fig. 11.



**Fig. 12.** The corresponding contour plots of Fig. 11..

## 5 Conclusion

In this paper, based on the Hirota's bilinear form and Mathematical software, the lump and interaction solutions between lump and solitary wave of a

generalized  $(3 + 1)$ -dimensional variable-coefficient nonlinear-wave equation in liquid with gas bubbles are studied. Their physical structures are described in some 3d graphs and contour plots. A periodic-shape rational solution is listed in Fig. 1(a) and Fig. 1(d). A parabolic-shape rational solution is presented in Fig. 1(b) and Fig. 1(e). A cubic-shape rational solution is shown in Fig. 1(c) and Fig. 1(f). In lump solutions ( $u^{(II)}$ ), The spatial structure called the bright lump wave is seen in Fig. 2, the spatial structure called the bright-dark lump wave is shown in Fig. 3. Interaction behaviors of two bright-dark lump waves are presented in Fig. 4. A periodic-shape bright lump wave is found in Fig. 5. In lump solutions ( $u^{(III)}$ ), the spatial structure called the bright lump wave is seen in Fig. 6. Interaction behaviors of two bright lump waves are presented in Fig. 7. A periodic-shape bright lump wave is found in Fig. 8. Fig. 9 and Fig. 10 display the interaction phenomena between lump and one solitary wave. Fig. 11 and Fig. 12 discuss the interaction phenomena between lump and two solitary waves.

### Compliance with ethical standards

**Conflict of interests** The authors declare that there is no conflict of interests regarding the publication of this article.

**Ethical standard** The authors state that this research complies with ethical standards. This research does not involve either human participants or animals.

### References

1. Deng, G.F., Gao, Y.T.: Integrability, solitons, periodic and travelling waves of a generalized  $(3+1)$ -dimensional variable-coefficient nonlinear-wave equation in liquid with gas bubbles. *Eur. Phys. J. Plus.*, 132(6), 255-271 (2017)
2. Wazwaz, A.M.: Painlevé analysis for a new integrable equation combining the modified Calogero-Bogoyavlenskii-Schiff (mcbs) equation with its negative-order form. *Nonlinear Dyn.*, 91(2), 877-883 (2018)
3. Osman, M.S.: Nonlinear interaction of solitary waves described by multi-rational wave solutions of the  $(2+1)$ -dimensional Kadomtsev-Petviashvili equation with variable coefficients. *Nonlinear Dyn.*, 87(2), 1209-1216 (2017)
4. Wazwaz, A.M., El-Tantawy, S.A.: Optical Gaussons for nonlinear logarithmic Schrödinger equations via the variational iteration method, *Optik*, 180, 414-418 (2019)
5. Tan, W., Dai, Z.D.: Spatiotemporal dynamics of lump solution to the  $(1 + 1)$ -dimensional benjamin-ono equation. *Nonlinear Dyn.*, 89, 2723-2728 (2017)
6. Wazwaz, A.M.: Multiple-soliton solutions for extended-dimensional jimbo-miwa equations. *Appl. Math. Lett.*, 64, 21-26 (2017)
7. Xie, X.Y., Meng, G.Q.: Multi-dark soliton solutions for a coupled AB system in the geophysical flows. *Appl. Math. Lett.*, 92, 201-207 (2019)
8. Yang, J.Y., Ma, W.X., Qin, Z.: Lump and lump-soliton solutions to the  $(2+1)$ -dimensional ito equation. *Anal. Math. Phys.*, 8(3), 427-436 (2017)
9. Li, Y.Z., Liu, J.G.: Multiple periodic-soliton solutions of the  $(3+1)$ -dimensional generalised shallow water equation. *Pramana*, 90(6), 71 (2018)
10. Alquran, M., Jaradat, H.M., Syam, M.I.: Analytical solution of the time-fractional phi-4 equation by using modified residual power series method. *Nonlinear Dyn.*, 90(4), 2525-2529 (2017)

11. Hu, C.C., Tian, B., Wu, X.Y., Yuan, Y.Q., Du, Z.: Mixed lump-kink and rogue wave-kink solutions for a  $(3 + 1)$ -dimensional B-type Kadomtsev-Petviashvili equation in fluid mechanics. *Eur. Phys. J. Plus.*, 133, 40 (2018)
12. Sun, H.Q., Chen, A.H.: Lump and lump-kink solutions of the  $(3+1)$ -dimensional Jimbo-Miwa and two extended jimbo-miwa equations. *Appl. Math. Lett.*, 68, 55-61 (2016)
13. Zhang, J.B., Ma, W.X.: Mixed lump-kink solutions to the bkp equation. *Comput. Math. Appl.*, 74, 591-596 (2017)
14. Osman, M.S., Machado, J.A.T., Baleanu, D.: On nonautonomous complex wave solutions described by the coupled schrödinger-Boussinesq equation with variable-coefficients. *Opt. Quant. Electron.*, 50(73), 1-11 (2018)
15. Kudryashov, N.A., Sinelshchikov, D.I.: Equation for the three-dimensional nonlinear waves in liquid with gas bubbles. *Phys Scr.*, 85, 025402 (2012)
16. Tu, J.M., Tian, S.F., Xu, M.J., Song, X.Q., Zhang, T.T.: Bäcklund transformation, infinite conservation laws and periodic wave solutions of a generalized  $(3+1)$ -dimensional nonlinear wave in liquid with gas bubbles. *Nonlinear Dyn.*, 83(3), 1199-1215 (2016)
17. Huang, W.H.: A polynomial expansion method and its application in the coupled Zakharov-Kuznetsov equations. *Chaos. Soliton. Fract.*, 29(2), 365-371 (2006)
18. Rao J.G., Mihalache, D., Cheng, Y., He, J.S.: Lump-soliton solutions to the Fokas system. *Phys. Lett. A.*, 383(11), 1138-1142 (2019)
19. Zhou, Q., Zhu, Q., Liu, Y., Yao, P. Bhrawy, A.H., Moraru, L., Biswas, A.: Bright-dark combo optical solitons with non-local nonlinearity in parabolic law medium. *Optoelectron Adv. Mat.* 8(9-10), 837-839 (2014)
20. Ma, W.X., Zhou, Y.: Lump solutions to nonlinear partial differential equations via Hirota bilinear forms. *J. Diff. Eqs.*, 264, 2633-2659 (2018)
21. Gao, X.Y.: Looking at a nonlinear inhomogeneous optical fiber through the generalized higher-order variable-coefficient Hirota equation. *Appl. Math. Lett.*, 73, 143-149 (2017)
22. Gao, X.Y.: Mathematical view with observational/experimental consideration on certain  $(2+1)$ -dimensional waves in the cosmic/laboratory dusty plasmas. *Appl. Math. Lett.* 91, 165-172 (2019)
23. L, W.H., Zhang, Y.F., Shi, D.D.: Lump waves, solitary waves and interaction phenomena to the  $(2+1)$ -dimensional Konopelchenko-Dubrovsky equation. *Phys. Lett. A.*, 383, 97-102 (2019)
24. Qawasmeh, A., Alquran, M.: Reliable Study of Some New Fifth-Order Nonlinear Equations by Means of  $(G'/G)$ -Expansion Method and Rational Sine-Cosine Method. *Appl. Math. Sci.*, 8(120), 5985-5994 (2014)
25. Biswas, A., Sonmezoglu, A., Ekici, M., Mirzazadeh, M., Zhou, Q., Alshomrani, A.S., Moshokoa, S.P., Belic, M.: Optical soliton perturbation with fractional temporal evolution by extended  $(G'/G)$ -expansion method. *Optik* 161, 301-320 (2018)
26. Huang, L.L., Chen, Y.: Lump solutions and interaction phenomenon for  $(2+1)$ -dimensional sawada-kotera equation. *Commun. Theor. Phys.*, 67(5), 473-478 (2017)
27. Xie, X.Y., Meng, G.Q.: Dark solitons for the  $(2+1)$ -dimensional Davey-Stewartson-like equations in the electrostatic wave packets. *Nonlinear Dyn.*, 93, 779-783 (2018)
28. Ma, W.X., Li, J., Khalique, C.M.: A Study on Lump Solutions to a Generalized Hirota-Satsuma-Ito Equation in  $(2+1)$ -Dimensions. *Complexity*, 11, 9059858 (2018)
29. Chen, R.P., Dai, C.Q.: Vortex solitons of the  $(3+1)$ -dimensional spatially modulated cubic-quintic nonlinear schrödinger equation with the transverse modulation. *Nonlinear Dyn.*, 90(3), 1563-1570 (2017)
30. Chen, M.D., Li, X., Wang, Y., Li, B.: A pair of resonance stripe solitons and lump solutions to a reduced  $(3+1)$ -dimensional nonlinear evolution equation. *Commun. Theor. Phys.*, 67(6), 595-600 (2017)
31. Huang, Q.M., Gao, Y.T.: Wronskian, Pfaffian and periodic wave solutions for a  $(2 + 1)$ -dimensional extended shallow water wave equation. *Nonlinear Dyn.*, 89(4), 2855-2866 (2017)
32. Ma, W.X.: Riemann-Hilbert problems of a six-component fourth-order AKNS system and its soliton solutions. *Comput. Appl. Math.*, 37(5), 6359-6375 (2018)
33. Ma, W.X.: lumps and their interaction solutions of  $(3+1)$ -dimensional linear PDEs. *J. Geom. Phys.*, 133, 10-16 (2018)
34. Li, Y.Z., Liu, J.G.: New periodic solitary wave solutions for the new  $(2+1)$ -dimensional Korteweg-de Vries equation. *Nonlinear Dyn.*, 91(1), 497-504 (2018)

- 
35. Lan, Z.Z., Hu, W.Q., Guo, B.L.: General propagation lattice Boltzmann model for a variable-coefficient compound KdV-Burgers equation. *Appl. Math. Model.*, 73, 695-714 (2019)